Trichoderma asperelloides enhances local (LAR) and systemic acquired resistance (SAR) response under low nitrate nutrition in *Arabidopsis*

Aakanksha Wany, Pradeep K. Pathak, Kapuganti Jagadis Gupta *

National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi, 110067, India

Correspondence

*Kapuganti Jagadis Gupta

National Institute of Plant Genome Research

Aruna Asaf Ali Marg

110067

New Delhi

Email: jgk@nipgr.ac.in

Phone +91-11-26735111

E-mail: jgk@nipgr.ac.in

Summary

- Nitrogen (N) is essential for growth, development and defense but how low N effects defense response is not known.

- Low nitrate fed Arabidopsis plants displayed reduced growth and compromised local and systemic acquired resistance (LAR & SAR) response when infected with avirulent and virulent *Pseudomonas syringae* DC3000. Growth and defense responses were enhanced in the presence of *Trichoderma asperelloides*.

- The mechanism of increased LAR and SAR involve increased N uptake and enhanced protein levels via modulation of various low and high affinity nitrate transporter genes (*CLC-A*, *NPF1.2*, *NRT2.1*, *NRT2.2* and *NRT2.4*). *nrt2.1* mutant is compromised in LAR and SAR response suggesting a link between enhanced N transport and defense.

- Enhanced N uptake was mediated by Trichoderma elicited nitric oxide (NO). Low NO producing mutants *nia1,2*, *nsHb+* were unable to induce nitrate transporters. These mutants displayed reduced protein levels and *PR1* gene expression in the presence of Trichoderma under low N suggesting a signaling role of NO elicited by *Trichoderma*.

- *Trichoderma* also induced SA levels and defense gene expression under low N. Salicylate hydroxylase mutant (*NahG*) and Non-expresser of Pathogenesis Related gene1 mutant (*npr1*) were compromised in LAR and SAR response in the presence of Trichoderma.

- The mechanism of enhanced plant defense under low N mediated by Trichoderma involve NO, ROS production, *NRT* gene induction, production of SA and induction of defense, SAR marker genes.

Key words: Nitric oxide, nitrate transporters, *Pseudomonas syringae,* SAR, Trichoderma

Introduction

Nitrogen (N) is essential for normal growth and development of plants. It is a crucial component in the chlorophyll structure, nucleic acids and amino acids. N plays an important role in operation of primary and secondary metabolism and also involved in protection of plants from biotic and abiotic stresses (O'Brien *et al.* 2016; Mur *et al.* 2017). N deficiency occurs in soil due to slow mineralization, lack of sufficient organic matter, leaching due to heavy rainfalls and increased activities of denitrifying bacteria. Low N can retard the growth of plants and cause severe physiological and morphological defects (Walker *et al.* 2001; Landrein *et al.* 2018). To cope with this, plants have evolved N uptake systems to support their survival under N deficiency (Li *et al.* 2017). These uptake systems are based on their affinity with nitrates (NO_3^-); mediated by the family of nitrate transporter proteins in plants. First is low affinity nitrate transporters (LATs), which operate under optimum N conditions (>50 mM), second is high affinity nitrate transporters (HATs), which can operate under very low N concentrations (<1 mM) (Tsay *et al.* 2007; Li *et al.* 2017).

N plays a very significant role in plant defense (Gupta *et al.* 2013; Mur *et al.* 2017). Hence, operation of efficient N transport system help plants to defend against pathogens. This defense response involve activation of innate immune responses comprising of two unified branches termed PTI (PAMP Triggered Immunity) and ETI (Effector-Triggered Immunity) that initiate extensive transcriptional reprogramming (Alves *et al.* 2014; Jones and Dangl 2006). These processes require N. Moreover, the host mobilizes salicylic acid (SA) for the plant defense (Oliva and Quibod 2017, Mur *et al.* 2017). This pathway itself requires N for synthesis of various intermediates leading to SA production. The defense mechanism for invading pathogen importantly includes rapid programmed cell death known as the hypersensitive response (HR), which develops during incompatible plant-pathogen (*R-avr*) interactions (Delledonne *et al.* 1998). An early characteristic of HR is the rapid generation of superoxide (O_2^{--}), nitric oxide (NO) and accumulation of H_2O_2 following perception of avirulent pathogen signals (Lamb and Dixon, 1997). Moreover, SA produced during HR, plays an important role in plant defense (Mur *et al.* 2000, 2008; Gupta *et al.* 2013). Thus, N not only improves the nutritional status of the plant, but, also its concentration plays a role in defense. Moreover, the form of N nutrition can greatly influence the HR-mediated resistance in plants (Gupta *et al.* 2013).

Nitrate (NO_3^-) nutrition greatly influences programmed cell death (HR) via the production of NO which is a regulatory signal in plant defense (Delledonne *et al.* 1998) and is involved in the development of cell death (Gupta *et al.* 2011; Delledonne *et al.* 1998). NO production depends on NO_3^-. Therefore, N deficiency can also lead to reduced levels of NO. SA is known to be induced by NO, hence, low N leads to low NO and reduced SA levels. Since NO production also requires NO_3^-, it may also play a role in systemic acquired resistance (SAR) which results in broad-spectrum disease resistance against secondary infections after primary infection (Cameron *et al.* 1994, 1999). SAR develops either as a consequence of HR where NO has its proven role (Delledonne *et al.* 1998), or in the course of a compatible interaction resulting in disease development (Gupta *et al.* 2014). SAR is dependent on SA or its derivatives (Park *et al.* 2007, Metraux *et al.* 1990) and other metabolic pathways such as increased phenylalanine ammonia-lyase (PAL) activity (Smith-Becker *et al.*. 1998; Kim and Hwang 2014) and *PR* gene expression (Ryals *et al.* 1996). Previously, it was shown that NO plays a role in nitrate assimilation/uptake during stress by modulating nitrate transporters (Frungillo *et al.* 2013). Thus nitrate, and defense pathways those dependent on N, play multifaceted roles in plant defense.

Therefore, any mechanism that can increase N uptake may assist in plant defense under low NO_3^-. Many plant symbiotic microbes such as mycorrhiza, *Trichoderma*, PGPR are known to increase nutrient uptake. However, the operation of plant defense under low N and the effect of these microbes in increasing plant defense under N deficiency is not known.

Several species of *Trichoderma* play an important role in plant growth promotion and resistance against various biotic and abiotic stresses. They confer resistance to plants via various mechanisms such as mycoparasitism, activation of plant defense responses via induced systemic resistance response (ISR) and increasing basal resistance. Previously, it was demonstrated that roots colonized by *Trichoderma* species can activate defense responses very rapidly and intensively via priming in *Arabidopsis* (Brotman *et al.* 2012). This occurs via modulation of transcription factors and key metabolites of plant defense (Shoresh 2010). Upon pathogen attack, *Trichoderma* treated plants show accumulated transcript levels of

PAL (Yedidia *et al.* 2003) and increased levels of defense-related plant enzymes
(peroxidases, chitinases, and β-1, 3-glucanases; Shoresh *et al.* 2005).

If *Trichoderma* enhances N uptake, it can have positive effect on plant
defense via LAR and SAR responses altogether. So far is it not known whether
Trichoderma can increase plant defense under low N. Therefore, in the present
study, we have assessed, the systemic defense response of low N fed *Arabidopsis*
thaliana plants to the phytopathogen; *Pseudomonas syringae* p.v. tomato DC3000
induced by the beneficial fungus *Trichoderma asperelloides* (T203) under low and
optimum nitrate concentrations. Here, we describe that T203 enhances SAR
response in Arabidopsis grown under low N via modulating nitrate transporters and
this mechanism involves both NO and SA.

Material and Methods

Plant material and growth conditions

Seeds of *Arabidopsis thaliana* ecotype Columbia 0 (*Col0*; WT) was sown in
plastic pots (10 cm diameter) containing soilrite: agropeat (1:1) mix (mix contains
zero N) and stratified at 4°C in dark for 48 h. Then, the pots were kept in growth
room under short day conditions (8h-light, 16h-dark), 22/18°C (day/ night)
temperatures, relative humidity of 60% and 200 µE m^{-2} s^{-1} light intensity. Initially,
plants were bottom irrigated for a week, once with half strength Hoagland solution
and once with water. Then, 0.1 mM and 3 mM N concentrations were given to the
growing plants weekly. The nitrate nutrient solution in Hoagland's media contained
either 0.1 mM or 3 mM KNO_3, 1 mM $CaCl_2$, 1 mM $MgSO_4$, 1 mM K_2HPO_4, 1 mM
KH_2PO_4, 15 µM H_3BO_3, 3 µM $MnCl_2.4H_2O$, 0.25 µM $ZnSO_4.7H_2O$, 0.1 µM
$CuSO_4.5H_2O$, 0.04 µM Na_2MoO_4 and 25 µM NaFe-EDTA. 30-36 day old plants with
fully developed rosette were used for the experiments. The seeds of *nrt2.1*
(SALK_035429C) were procured from ABRC, and only homozygous T-DNA
insertion lines were used for the experiments.

***Trichoderma* supplementation**

152   *T. asperelloides* (T203 strain) was grown on potato dextrose agar (PDA;
153 Himedia) plates for 15 days under low light conditions until sporulation. Conidia were
154 harvested by gently scraping the petridish and poured with 10 ml of sterile water.
155 The spores were evaluated up to 1x10^9 spores/ml. It was thoroughly mixed into the

soilrite mix and distributed into the individual plastic pots.

Bacterial strain and growth conditions

Pseudomonas syringae pv. tomato DC3000 (*Pst*DC3000; *avrRpm1,* avirulent) was used as the non-host pathogen strain on *A. thaliana*. Bacteria were grown in King's B (KB) medium containing 50 µg ml^{-1} rifampicin. The culture was revived in 5 ml of King's B medium (Himedia, India) and incubated at 28°C with continuous shaking at 180 rpm for 12h. Upon reaching OD600 of 0.2, bacterial cells were harvested by centrifugation at 5000 rpm for 15 min. The pellet containing bacterial cells was washed thrice with water and re-suspended in 10 mM MgCl$_2$. The OD600 of the bacterial suspension was measured and diluted to OD600 of 0.005 (1×10^6 CFU ml^{-1}). The OD600 was equated to OD 0.1= 2×10^7 CFU ml^{-1}. Similarly, virulent *P. syringae* was cultured and used at OD of 0.005 (2×10^6 CFU ml^{-1}) for challenge inoculation.

Bacterial infiltration in leaves

Mock infiltration (control) was performed with 10 mM MgCl$_2$. *Pst* inoculations (1×10^6 CFU ml^{-1}) were made by syringe infiltration on the abaxial side of the leaves manually. There were total four groups of plants; Group 1: 0.1 mM + MgCl$_2$ plants, Group 2: 3 mM + MgCl$_2$ plants, Group 3: 0.1 mM + *Pst* plants and Group 4: 3 mM + *Pst* plants. Each group consisted of 20-25 plants. SAR experiment initiated with primary inoculation (a*vrRpm1 Pst* DC3000) on one leaf per plant, and the secondary challenge inoculation with (virulent *P. syringae*) on 4 other (distant) leaves per plant, leaving 5-6 healthy un-inoculated leaves per plant (Cameron *et al*. 1999) after 2 days of primary inoculation. The Inoculated (I) and un-inoculated (U) leaves from each treatment were harvested (Fig. S1) for RNA extraction and other experiments post challenge inoculation. Leaves were analyzed for 0, 6, 24 and 48 hours post inoculation (hpi) or challenge (hpc) and the symptoms was determined periodically.

Electrolyte leakage

Leaf discs were taken and electrolyte leakage was monitored by conductivity meter (Lab Man Scientific Instruments, India) at different time points from inoculated and un-inoculated leaves and electrolyte leakage was calculated as described in (Gupta *et al.* 2013).

In planta bacterial number quantification assay

Bacterial multiplication in leaves from *Pst* treated plants was assessed after primary as well as challenge inoculation. Each infected leaf disc (5 mm in diameter) was surface sterilized with 0.02% H_2O_2 for 8 minutes, and was homogenized in 1 ml of sterile 10 mM $MgCl_2$. Upon further serial dilution in 10 mM $MgCl_2$, it was plated on KB agar medium supplemented with rifampicin (50 µg ml^{-1}). For all the bacterial treatments, *in planta* bacterial number was assayed up to 2 days at 24h interval. Bacterial number was calculated as per Gupta *et al.* (2016).

Expression profiling by qRT-PCR

The inoculated and uninoculated leaves were collected at specific time points and were immediately frozen in liquid nitrogen and stored at -80°C. RNA extraction, cDNA synthesis and qPCR was performed according to Wany *et al.* (2017; 2018). The synthesized cDNAs were used as templates in qRT-PCRs performed using primers (Table S1). Fold change in the target genes was normalized to *Arabidopsis* ubiquitin, *18sRNA* (GQ380689) and *YSL8* gene (X69885.1). Fold expression relative to control treatment was determined by ΔΔCT values. Three biological experiments (with three independent replicates for each experiment) were performed for each treatment. The comparisons of the gene expression between the different treatments to control was performed by student's t-test. Differences between the treatments with $p<0.05$, $p<0.01$ and $p<0.001$ were considered significant.

Ex- vivo NO estimation

For this experiment, NO was measured from the following five different combinations in roots; 1. WT; 2. WT + T203 (*Trichoderma*); 3. *nia1,2* double mutants +T203; 4. WT + cPTIO; 5. WT + cPTIO + T203. WT and *nia1, 2* plants were grown for one week in plates containing 0.1 mM and 3 mM nitrogen concentrations. The spore suspension of *Trichoderma* was poured over these one week old plants and incubated for 2 minutes, 10 minutes and 24 hours. Then, the roots were incubated in 10 µM DAF-FM DA (4-amino-5-methylamino- 2′,7′-difluorofluorescein diacetate) in 100 mM HEPES buffer (pH 7.2) placed in a 1.5 ml tube and incubated for 15 minutes in dark. Then, the roots were washed three to four times using same buffer and transferred onto the slide and mounted with a drop of 10% glycerol with a cover-slip.

Each sample was photographed using fluorescence microscope (Nikon*80i*, Japan) with a CCD camera (DXM1200C) equipped with Image software (NIS elements ver3.2) at 495 nm excitation and 515 nm emission wavelength.

Determination of ROS levels

Production of hydrogen peroxide (H_2O_2) in inoculated and un-inoculated leaves was detected by Diaminobenzidine tetrahydrochloride (DAB) staining as per Daudi *et al*. (2012) with slight modifications. The superoxide levels were measured by *in vivo* staining with Nitroblue tetrazolium chloride (NBT, SA, USA) (Jambunathan 2010).

Histochemical detection of HR

Hypersensitive cell death in inoculated leaves was visualized by trypan blue staining method according to Fernández-Bautista *et al.* (2002).

SA levels

The SA levels were measured by HPLC according to the protocol described in Singh *et al.* (2013). The inoculated and uninoculated leaves of both WT and WT + T203 plants (0.1 mM and 3 mM) were harvested at 0, 6, 24 and 48 hpc for SA estimation.

Results

Optimal and low nitrogen effects LAR and SAR differently

Nitrogen plays important role in plant defense, hence, we first tested the effect of low N on plant defense mediated by LAR and SAR response. For this purpose, WT plants were grown under optimum (3 mM) and low (0.1 mM) NO_3^- conditions and challenged with *P. syringae* DC3000 (*avrRpm1*) and observed local HR at different time points. As shown in Fig.1a (lower panel) within 24 hpi, the 3 mM NO_3^--fed plants showed HR (lesion formation) at the inoculation sites and it was further progressed to higher levels till 48 hpi (Fig.1a) whereas 0.1 mM NO_3^- plants showed chlorotic lesions at the site of inoculation at 24 hpi and these spots were enhanced in 48 hpi. Electrolyte leakage (EL) (Fig.1b) was more rapid in 3 mM *Pst* infiltrated plants than

0.1 mM plants which is in accordance with HR response. In order to link HR
formation with plant resistance, *in planta* bacterial number were determined (Fig. 1c).
A significant increase in CFU count was observed in 0.1 mM grown plants at 48 hpi
in comparison to bacterial numbers in 3 mM grown plants. We further found that, the
PR1 (a marker for HR) transcript levels have increased in 3 mM *Pst* treated plants
(up to 25 fold till 48hpi), but in 0.1 mM WT plants, this gene was not at all induced at
48 hpi (Fig1d). Taken together these results suggest that low NO_3^- grown plants are
compromised with HR linked LAR response.

After observing LAR, we studied SAR (Fig. S1b; Cameron *et al.* 1994 and
1999). Three days post-secondary challenge with virulent *Pst* DC3000, EL, CFU and
PR1 gene expression were assessed. During the primary inoculation, symptoms
were visible after 24 and 48 hpi in 3 mM *Pst*-treated leaves, than 0.1 mM *Pst*-treated
leaves (Fig. 2a- first panel), which showed enhanced chlorotic and necrotic
symptoms. Three days after secondary challenge, disease symptoms were observed
from 0 to 48 hpc. 0.1 mM *Pst*-challenged plants show more prominent necrotic and
dark yellow lesions spread to both sides of the leaf (Fig. 2a-second panel) confirming
susceptible symptoms (yellow specks) where as 3Mm grown plants showed reduced
yellowing and necrotic symptoms. This clearly suggests that, 3 mM NO_3^- plants
defended better during secondary challenge than 0.1 mM NO_3^- plants suggesting
that optimal N is required for SAR development.

Trypan blue images of inoculated and uninoculated leaves are shown in Fig.
2b. It was observed that, after 3 days post challenge, the cell death observed in
infiltrated tissues of 0.1 mM NO_3^- plants is profoundly less in comparison to 3 mM
NO_3^- plants till 24 hpc. At 48 hpc, there was increased cell death in both 0.1 and 3
mM inoculated leaves but cell death phenotype is different. Interestingly, the
uninoculated leaves of 0.1 and 3 mM WT plants, showed small and uniform spread
of macrobursts throughout the leaf blades unanimously over the leaf, in all the time
points, in exception that, 3 mM leaves displayed slightly larger macrobursts (Fig. 2b).
This suggests that during SAR establishment, the mobile signal perceived by the
uninoculated leaves results in the occurrence of low frequency microscopic HRs
(Cameron *et al.* 1999; Alvarez *et al.* 1998).

Then, we assessed EL (Fig. 2c), we found leakage was significantly higher in
3 mM plants than 0.1 mM plants in both the cases (1° and 2°). But, much higher EL
was observed during 2°challenge (Fig. 2c, right panel). Mock inoculated plants

showed moderate EL in both 0.1 mM and 3 mM plants (Fig. S2a). Higher bacterial growth was observed in 0.1 mM *Pst* treated plants as expected till 72 hpc, while there was a significant reduction in the bacterial numbers in 3 mM *Pst* inoculated plants (Fig2d).

Low LAR under low nitrogen is reversed by *Trichoderma*

Several strains of Trichoderma are known to play important role in plant growth, development and stress resistance (Brotman *et al.* 2010). Hence, we investigated whether *Trichoderma asperelloides* plays a role in improving LAR and SAR response under low NO_3^- availability. Interestingly, we found that under 0.1 mM NO_3^-, *Trichoderma* treated plants displayed healthy phenotype than untreated plants (absence of Trichoderma) (Fig. S1a), this enhanced growth was linked to increased leaf number, fresh weight and total chlorophyll content (Fig. S2b, c and d). During LAR, Trichoderma grown 0.1 mM NO_3^- plants did not show any severe disease symptoms, rather they mimicked the HR phenotype of 3 mM plants and showed resistance (Fig. 3a). This suggests that Trichoderma provides resistance under low N stress (Compare Fig. 3a with Fig. 1a). EL was significantly higher in both 3 mM and 0.1 mM Trichoderma treated plants (Fig. 3b) in comparison to untreated plants (Compare Fig. 3b with Fig. 1b). As shown in Fig. 3c, bacteria multiplied to a high level in both 0.1 mM and 3 mM *Pst* inoculated plants till 8 hpi, but strikingly, there was a significant reduction in the bacterial count in 0.1 mM *Pst* plants at 24 hpi, suggesting that Trichoderma enhances LAR response under low N. The higher accumulation of *PR1* transcripts in both 0.1 mM (~9 folds at 48 hpi) and 3 mM (~28 folds at 48 hpi) in Trichoderma grown *Pst* treated plants (Compare Fig 3d with Fig1d) again providing evidence that Trichoderma can enhance LAR under low N.

***Trichoderma* can induce SAR under low N**

During SAR assay, the T203 treated 0.1 mM plants have shown a drastic change in resistance where clear resistant symptoms were evident at 24 and 48 hpc (Compare Fig. 4a left panel with Fig. 2a left panel). During secondary challenge also, there was a clear reduction in yellowing in T203 grown 0.1 mM plants in comparison to untreated plants (Compare Fig. 4a right panel with Fig. 2a right panel). In response to T203, 0.1 mM NO_3^- plants showed more Trypan blue spots during secondary challenge. Upon infection, cell death spots were more widely spread in

these leaves (Compare Fig. 4b with Fig. 2b and Fig. S3 a&b). EL was significantly higher (~4 fold) in both 0.1 and 3 mM T203 treated plants during 2° challenge, in comparison to 1° inoculation at 24 and 48 hpc (Fig. 4c). This suggests that T203 treatment primes the defense responses in the uninoculated leaves immediately at 0 hpi and after 2° challenge. Bacterial count was in accordance with EL that, the *Pst* population decreased several folds in T203 treated low N-fed plants, suggesting that T203 imparts SAR response to the NO_3^- stressed plants (compare Fig. 4d with Fig 2d).

Trichoderma activates nitrate transporters, facilitates N uptake and promotes SAR in low N-fed plants

Increased LAR and SAR response under low N in the presence of *Trichoderma* is probably due to increased nutrient uptake facilitated by *Trichoderma* as these responses are absent in *Trichoderma* untreated plants. Increased nitrate transport is probably the reason for resistance hence, we checked the expression of both LATs (*CLC-A*, *NPF1.2*) and HATs (*NRT2.1*, *NRT2.2* and *NRT2.4*).

CLCa (the chloride channel family) is a tonoplast located antiporter channel system which drives nitrate accumulation in the vacuoles (Krapp *et al.* 2014). It was observed that, *CLCa* is significantly upregulated 4 to 6 folds in *Pst* treated leaves of 3 mM plants in 6h-48h; than *Pst* treated 0.1 mM plants (Fig. 5a), which displayed slight induction at 6 hpc (2 folds) and decreased at later time points. This suggest that under low NO_3^-, C*LCa* is less inducible. But, in response to *Pst* treatment T203 grown 3 mM WT plants (Fig. 5b), significantly induced to 4 to 5 folds at 24 and 48 hpc, whereas it was greatly induced to 6 folds in *Pst* treated 0.1 mM plants at 6 hpc, which was stabilized at later time points. The uninoculated leaves of 0.1 mM T203 grown WT plants showed slight induction of *CLCa* in comparision to non-T203 grown plants (Compare Fig. 5a & 5b). We further checked the expression of *NPF1.2;* which belongs to NRT1 (peptide transporter) family of nitrate transporters (Fig. 5c) and is involved in the transfer of xylem-borne nitrate to the phloem in the petiole (Krapp *et al.* 2014). It was found that, in *Pst* treated 3 mM WT plants, *NPF1.2* levels were highly induced till 48 hpc, in comparison to 0.1 mM *Pst* inoculated leaves (Fig. 5c). This might be the reason behind resistance of 3 mM WT plants and susceptibility of 0.1 mM WT plants after pathogen challenge. But, upon T203 pre-treatment, *NPF1.2* transcript levels in *Pst*-inoculated 0.1 mM plants (Fig. 5d), showed significant

induction in comparison to same treatment without T203 (Compare Fig. 5c & 5d). This suggests that, T203 may cause the regulation of *NPF1.2*.

NRT2 transporters (*NRT2.1*, *NRT2.2* and *NRT2.4*) are HATS and gets activated at low N concentrations (<1 mM). Under optimum N conditions, *NRT2.1* expression levels in *Pst* treated 3 mM plants were highly induced from 2.5 to 6 fold post challenge. Previously, it was shown that *NRT2.1* is active only under N starving conditions (Dechorgnat *et al.* 2012). We found similar results that *NRT2.1* transcript was induced under 0.1 mM *Pst* treated plants (Fig. 6a). Likewise, the uninoculated leaves of 3 mM WT plants showed slight induction at 6 hpc, which significantly down-regulated at later time points. Whereas, in 0.1 mM uninoculated leaves of WT plants, *NRT2.1* showed sharp induction of 4.2 folds at 0 hpc, 6.3 folds at 6 hpc, which drastically reduced at later time points (Fig. 6a). This suggests that during the early initiation phase of SAR, i.e., upto 6 hpc, low N-fed WT plants tried to initiate the immunization potential by maintaining the N supply by displaying higher *NRT2.1* transcripts. The T203 pre-treatment displayed reprogrammed transcript levels of *NRT2.1* in *Pst* treated 3 mM WT plants, which showed extremely rapid induction up to 10 folds at 0 hpc, 5 folds at 6 and 24 hpc which got stabilized at >11 folds at 48 hpc. In line with this, *Pst* treated 0.1 mM T203 grown WT plants showed 14 fold up regulation at 24 hpc, which stabilized at 10 folds at 48 hpc (Fig. 6a). The SAR establishment stage represented by uninoculated leaves showed dynamic and hiked up-regulation of *NRT2.1* transcripts up to 26 folds in 24 hpc and 28 folds in 48 hpc, whereas, there was <5 folds induced level of *NRT2.1* expression in all the time points in 3 mM plants. This revealed that, T203 colonization benefits the plant by facilitating the N supply even in low NO_3^- conditions, after 48 hpc and more, thereby showing improved plant defense mediating SAR.

Further, we checked *NRT2.2* expression levels (Fig. 6b), it was observed that, in *Pst* treated 3 mM plants, there was an induction of this transcript from 1.6 to 3.2 fold at various time points. In comparison with 3 mM plants, 0.1 mM *Pst* treated plants showed very strong induction from 4 to 6 fold at 6 to 24hpc. This suggests that, the *NRT2.2* expression is highly induced in low N-fed plants facilitating their NO_3^- uptake and is also highly correlated to *Pst* treatment. The systemic (U) leaves of 3 mM plants, showed a stable and uniform *NRT2.2* transcript accumulation with at 2-4 fold at 24 and 48 hpc respectively, whereas, 0.1 mM systemic leaves showed

highest of 5.3 fold change at 6 hpc. In Fig. 6b, T203 treated *Pst* infiltrated 3 mM plants showed constant induction from 2.5 fold at 0 hpc, 10 fold at 6 hpc, and 5- 6 fold at 24 and 48 hpc respectively. In 0.1 mM *Pst* treated plants, there was very strong induction of 6 fold at 0 hpc, which hiked to 25 fold at 6 hpc, later came down to 9 fold and 7 fold in 24 and 48 hpc respectively (Fig. 6b). The uninoculated distal leaves showed remarkable 12 fold induction at 6 hpc, 6 and 3.2 fold change at 24 and 48 hpc in 3 mM plants. In 0.1 mM uninoculated WT + T203 leaves, there was rapid accumulation upto 23 folds of *NRT2.2* transcripts, 11 folds in 24 hpc and >12 folds at 48 hpc. This again revealed that faster N uptake by T203 treated low NO_3^- plants can help to defend better.

Consequently, we checked *NRT2.4* expression levels in both set of treatments. It was found that in *Pst* treated 3 mM WT plants, there was consistent induction of *NRT2.4* expression from 0 hpc (1.5 fold) to 48 hpc (<4 fold), whereas in *Pst* treated 0.1 mM plants, there was induction of 6 fold at 6 hpc, which drastically reduced to <0.5 fold in 24 and 48 hpc (Fig. 6c) was observed. The uninoculated leaves of 3 mM plants showed induced expression only at 0 hpc, but later got declined in consecutive time points to <0.1 fold, conversely, 0.1 mM plants showed higher induction of 2.5 fold at 0 hpc till 5.5 fold hike at 6 hpc. Upon T203 treatment (Fig. 6c), 3 mM *Pst* treated plants showed consistent and induced expression pattern from 2.9, 2.8, 6.5 and 4.5 fold at 0, 6, 24 and 48 hpc, whereas, 0.1 mM plants showed initial induction till 6 hpc (6 fold), but, later declined. There was only slight difference in the expression levels of 3 mM and 0.1 mM uninoculated leaves of T203 treated and untreated WT plants.

Taken together, these results suggests that Trichoderma induces HATs and LATs to facilitate N uptake under low NO_3^- conditions. To confirm the role of HATs in increasing plant defense via N uptake, we checked the *PR1* expression in WT and *nrt2.1* mutants (Fig 6d). The expression of *PR1* gene in *Trichoderma* inoculated plants increased 12-13 folds in 24 h but in the case of *nrt2.1* mutant (Fig. 6d), *PR1* gene was not at all induced suggesting that *NRT2.1* plays an important role in increasing plant defense under low nitrogen mediated by Trichoderma. Then, we further found that in *nrt2.1* mutant, the expression of *NRT2.2* gene also suppressed in both control and Trichoderma treatment (Fig. 6e) suggesting that *NRT2.1* is mainly responsible for increasing uptake of N under low NO_3^- facilitated by *Trichoderma*.

Then, we checked the protein levels in WT and *nrt2.1* mutant in the presence or absence of Trichoderma under low N. In response to Trichoderma, protein levels have increased in WT, whereas in *nrt2.1* mutant protein levels were less than WT but upon T203 treatment only slight increase (Fig. 6f) was observed suggesting T203 application leads to increase nitrate transport hence increase protein levels.

Trichoderma elicits NO production during early stages of inoculation which is required for induction of HATs and PR gene expression

Previously, it was shown that Trichoderma elicits NO at early stages (Gupta *et al.* 2014). This NO could have a role in induction of transporter genes. Hence, we checked NO production in WT, *nia1,2* mutants and WT seedlings grown on NO scavenger cPTIO inoculated with Trichoderma. Control roots in 0.1 and 3 mM NO_3^- produced reduced levels of NO. Within 2 min of T203 application, WT plants showed much increased levels of NO in both 0.1 and 3 mM nitrate (Fig. 7) but the increase was slightly higher in 0.1 mM than 3 mM. But, in 10 minutes and 24 h of T203 incubation, the WT plants showed extremely low fluorescence (Fig. 7). This suggests T203 greatly induces NO in low NO_3^- grown plants within 2 min of Trichoderma application. Thus, Trichoderma induced NO probably plays a role in priming and induction of LATs and HATs. Assessment of NO in *nia1,2* mutant revealed that NR is responsible for NO production. C PTIO grown plants showed reduced levels of NO.

Then, we further checked the importance of Trichoderma elicited early stages of NO in the induction of HATs, total protein levels and expression of *PR1* gene in low NO producing phytoglobin over-expressing *nsHb+* line, *nia1,2* and *nrt2.1* mutants. Expression of *NRT2.1* and *2.2* levels have increased in WT in response to Trichoderma, whereas in Hb+, *nia1,2* and *nrt2.1* mutants these two HATs were not induced (Fig 8a & 8b) suggesting that Trichoderma elicited NO is responsible induction of HATs. Further reduced protein levels were observed in in *nia1,2*, Hb+ plants suggesting that induction of transporters mediated by *Trichoderma* elicited NO plays a role in increased N uptake under low N (Fig. 8c), further, we found that *nia1,2* and Hb+ were unable to induce *PR1* expression under these conditions. Taken together, these results suggests that *Trichoderma* elicited NO plays a role in overall increase in defense response under low N.

Further, we investigated the role of NO in LAR and SAR development in the

presence of Trichoderma. Under low NO_3^-, WT plants performed better LAR

response than Hb^+ and *nia1,2* mutant. Reduced bacterial growth and increased *PR1*

gene expression was observed in WT than Hb^+ and *nia1,2* mutant (Fig. 9 a,b&c).

Similarly, SAR response was also compromised in than Hb+ and *nia1,2* like LAR, a

similar reduced bacterial number and increased *PR1* gene expression was observed

in WT in comparison to Hb^+ and *nia1,2* mutant (Fig. 9 d,e&f).

ROS is a component of *Trichoderma* induced increased resistance via SAR

Both NO and ROS are involved in plant resistance response hence, we

investigated the role of ROS. The distal un-inoculated leaves of 0.1 mM NO_3^- WT+T

plants showed H_2O_2 production as concentrated patches in comparison to 0.1 mM

un-inoculated leaves of WT plants (Fig. 10a-upper panel). Also, the level of H_2O_2

was intense and distributed in 3 mM un-inoculated leaves of WT+T plants in

comparison to untreated plants (Fig. 10a-lower panel). This is probably due to

suppression of catalase activity by SA. This suggests that T203 inoculation can

enhance H_2O_2 pattern in *Pst* uninoculated leaves during SAR, whereas, both the 3

mM *Pst* inoculated leaves from WT and WT+T203 plants showed higher H_2O_2

production in the infiltrated area which is visible as a dark brown patch at 24 and 48

hpc (Fig. 10a-lower panel). This clearly suggests that T203 plays a role in inducing

H_2O_2 in low NO_3^- fed WT plants during SAR.

In order to investigate that, $O_2^{-.}$ are key players in cell death, we measured

their levels during SAR response (Fig. 10b). It was observed that, there was

increased formazan formation (dark blue spots) in inoculated leaves of both 0.1 and

3 mM fed WT and T203 treated WT plants, but increase was much higher in T203

treated WT plants than Trichoderma untreated plants (compare Fig. 10a-lower left

and 10b-right panel). This suggests that there was a rapid oxidative burst (HR)

which offers resistance to the plants in response to T203 treatment. These results

also suggests that, T203 treatment can enhance $O_2^{-.}$ levels of 0.1 mM NO_3^- grown

plants to contribute towards H_2O_2 production to enhance resistance. Taken together,

the coordinated interplay of H_2O_2, $O_2^{-.}$ and NO leads to the HR associated cell death

that generously improvises the LAR and SAR responses in low N-stress plants.

Trichoderma induces defense genes during SAR in low N-fed plants

486 Next, we checked the expression profiles of defense related genes (*PAL1,*
487 *PR1, PR2* and *PR5*) during SAR. In both 3 mM and 0.1 mM *Pst* inoculated WT
488 plants, *PAL1* transcripts were highly induced in all time points, but, the uninoculated
489 leaves showed *PAL1* induction only till 6 hpc, but drastically declined at later time
490 points (Fig. 11a-I). Trichoderma grown WT plants under 3 mM and 0.1 mM, showed
491 even more enhanced levels of *PAL1* transcripts (Fig. 11a-II) in inoculated as well as
492 uninoculated leaves till 48 hpc. Similar trend was observed in *PR1* (Fig. 11b), *PR2*
493 (Fig. 11c) and *PR5* (Fig. 11d) expression profiles, which got elevated after
494 Trichoderma application. Out of all the four defense genes examined, *PR1* gene (SA
495 marker) displayed the extremely higher expression levels in Trichoderma grown
496 plants suggesting that T203 might induce SA levels to a greater extent during SAR.

Trichoderma enhances expression of SAR mediated regulatory genes

498 Further, we checked whether Trichoderma can induce SAR response via
499 induction of regulatory genes such as *DIR1, NPR1, SARD1 and TGA3*. The lipid
500 transfer protein, DEFECTIVE IN INDUCED RESISTANCE1 (*DIR1*) is a key mobile
501 component of SAR response (Maldonaldo *et al.* 2002) involved in long-distance
502 translocation from local to distant leaves (Carella *et al.* 2015, Champigny *et al.*
503 2013). It was found that, DIR1 induction took place only in the initial time points i.e. 0
504 and 6 hpc in both 0.1 and 3 mM inoculated and uninoculated WT plants (Fig. 12a-I).
505 But, in T203 grown plants, irrespective of early time points, a slight increased *DIR1*
506 expression was observed at 24 and 48 hpc (Fig. 12a-II). Upon perception of SAR
507 mobile signals, SAR master regulator, Non-Expresser of Pathogenesis-Related
508 Genes1 (*NPR1*) is required to activate defense in challenged plants (Cao *et al.*
509 1997). We found a similar trend in the *NPR1* expression profile like *DIR1* in WT
510 plants (Fig. 12b-I). Moreover, upon the *Trichoderma* treatment, the levels of *NPR1*
511 gradually increased in 24 and 48 hpc (~10 fold) in both 0.1 and 3 mM inoculated and
512 uninoculated challenged leaves (Fig. 12b-II).

513 Next, we checked the expression of SAR DEFICIENT 1 (*SARD1*) a pathogen-
514 induced transcription factor (Zhang *et al.* 2010) and a key regulator for Isochorismate
515 Synthase 1 (*ICS1*) and SA synthesis (Wang *et al.* 2011). A remarkably stronger
516 induction of *SARD1* expression levels in T203 treated 0.1 mM uninoculated leaves at

6 hpc (~82 fold) revealed that Trichoderma is a potential inducer of SA biosynthesis (Fig. 12c-II), whereas the other treatments at different time points remained less than 15 fold in both the nitrate conditions (Fig.12c). Another important regulatory gene, TGA3, is an *NPR1*-interacting protein (NIPs) and is a critical component in the SA signaling mechanism regulating the expression of SA-induced genes such as *PR1* (Zhou *et al.* 2000). Upon T203 treatment, it was found that, the TGA3 levels were also re-programmed in all the treatments (Fig. 12d) suggesting that Trichoderma might act as channel in SA signaling pathways.

nrt2.1 and npr1 mutants are compromised in LAR and SAR

Further we checked LAR in *nrt2.1* (nitrate transporter mutant) and *npr1* (SA master regulator mutant) in response to virulent *Pst*DC3000. In 0.1 mM N-fed *nrt2.1* and *npr1* mutants developed more severe symptoms than 0.1 mM WT plants (Fig. 13a- 0.1 mM panel). Whereas the 3 mM NO_3^--fed WT plants showed less disease symptoms as compared to 0.1 mM NO_3^- plants (Compare Fig 1a and Fig. 13a- 3 mM panel). In order to check plant resistance, *in planta* bacterial number were determined (Fig 13b). A continuous increase in CFU count was observed in 0.1 mM grown WT, *nrt2.1* and *npr1* at 24 and 48 hpi in comparison to bacterial numbers in 3 mM NO_3^- grown plants. This suggests that optimum N concentration plays an important role in resistance response towards virulent *Pst* DC3000.

Similarly, we studied the SAR response in the WT, *nrt2.1* and *npr1* plants. 0.1 mM Pst-challenged WT plants show more prominent necrotic and discolored lesions spread to half of the leaf (Fig. 2a) confirming susceptible symptoms (yellow specks). On the other hand, the mutant's *nrt2.1* and *npr1* showed even more disease symptoms (extensive chlorosis and necrosis) (Fig. 13c) under low NO_3^- conditions. Results suggests (Fig. 13c) that, 3 mM NO_3^- plants defended better during secondary challenge than 0.1 mM NO_3^- plants. Bacterial populations also increased in *nrt2.1* and *npr1* mutants (Fig. 13d) suggesting that *NRT2.1* and *NPR1* plays role in increasing defense mediated by Trichoderma.

Salicylic acid pathway is part of enhanced plant resistance under low N mediated by *Trichoderma*

Examination of SA levels revealed that Trichoderma presence accelerated
total SA levels in 0.1 mM uninoculated distal leaves, in comparison to 3 mM
uninoculated leaves, whereas there was no significant increase in SA levels
observed in inoculated leaves of 0.1 mM Trichoderma grown plants, while in 3 mM
Trichoderma grown plants a significant increase in SA levels were observed (Fig.
14a). We sought to further confirm role of SA in Trichoderma increased SAR, hence
NahG plants were challenged in the presence or absence of Trichoderma. An
intense chlorotic lesion was evident in both 0.1 and 3 mM grown *NahG* in response
to challenge inoculation while *Trichoderma* grown plants when challenged they
defended much better which can be seen in the terms of decreased chlorotic lesions
and decreased bacterial numbers (Fig. 14b& c). Surprisingly, in *Trichoderma* grown
NahG plants showed slightly enhanced *PR1* transcript levels were observed (Fig.
14d).

Discussion

Nitrogen availability and supply can severely impact growth and development
of plants (Walker *et al.* 2001; Landrein *et al.* 2018). N deficiency can cause chlorosis,
which can impact photosynthesis and overall energy demand for growth and
defense. Since N is also important for synthesis of various secondary metabolites,
severe depletion of N can impact defense related pathways. Hence, plants may not
be able to activate defense pathways for tolerance or resistance (Snoeijers *et al.*
2000). Plant take up N in the form of NH_4^+ or NO_3^- or combination of both. Different N
forms differentially effect various free radicals such as NO and ROS (Wany *et al.*
2018; Gupta *et al.* 2013). Ammonium uptake and assimilation is less costly to the
plants in comparison to NO_3^- but excess of NH_4^+ can cause toxic effects to the plants
(Boudsocq *et al.* 2012; Liu *et al.* 2017), hence, many plants use NO_3^- as the
preferable N source. Moreover, NO_3^- can help in better charge balance (Boudsocq *et*
al. 2012). Nitrate nutrition can also enhance plant defence via increased generation
of NO, polyamines and SA (Gupta *et al.* 2013; Fagard *et al.* 2014). Hence, we
preferred to check the plant defense response under NO_3^- nutrition rather than under
NH_4^+.

Plants which are grown under 0.1 mM NO_3^- showed reduced growth (Fig. S1)
and reduced leaf number, chlorophyll (Fig. S2), suggesting the supplied 0.1 mM NO_3^-

is not sufficient for the plants for growth, under these conditions plants become more
susceptible. In our experiments, 0.1 mM NO_3^- grown plants were compromised in
both LAR and SAR (Fig. 1&2) response, suggesting that N is required for better
defense. Plants grown on low N produced less SA (Fig. 14a), further supporting that
N is needed for SA biosynthesis. Since plants need N for growth and disease
resistance, increasing N use efficiency can help them to defend. However, very less
information is available whether increasing N uptake can enhance plant resistance or
not. Some group of Trichoderma help in nutrient absorption leading to increased
growth and enhanced plant defense (Brotman *et al.* 2010). Previously, it was shown
that the supplementation of *Trichoderma asperelloides* enhances plant growth
(Brotman et *al.* 2012), and protects against abiotic and biotic stressors, moreover, it
induces systemic resistance responses (Contreras-Cornejo *et al.* 2016; Brotman *et*
al. 2012). Trichoderma induced increased growth attributed to auxin and ethylene
(Garnica-Vergara *et al.* 2016) and increased expression of genes involved in carbon
and nitrogen metabolism (Domínguez *et al.* 2016). However, there are hardly any
reports on investigating the mechanism of improved plant growth and defense under
N starvation mediated by Trichoderma. This study we unraveled the mechanism of
Trichoderma induced plant growth and defense under low N. Hence, in this current
work, we studied the impact of Trichoderma on enhancing N uptake and support in
LAR & SAR response under low NO_3^- stress.

The low NO_3^- fed plants which were grown in the presence of Trichoderma,
showed increased growth, increased leaf number, fresh weight and enhanced
chlorophyll content (Fig. S2) suggesting that Trichoderma can enhance growth under
low NO_3^- (Fig. S1a). We found that this enhanced growth is partly due to increase
nitrogen uptake which was evidenced by increased expression of *NRT2.1, NRT2.2,*
NRT 2.4 nitrate transporters (Fig. 6a, b, c, d, e &f) and increased protein levels (Fig.
6i). Since low NO_3^- fed plants becomes susceptible (Snoeijers *et al.* 2000), it is
logical to assume that Trichoderma can increase N uptake and enhance resistance.
Hence, in further experiments, we focused on LAR and SAR response under low and
high NO_3^- in the presence or absence of Trichoderma. During local *Pst* infection, the
plants display LAR response, and the systemic/distant uninoculated leaves induces
SAR response. Both are important for plants to defend against pathogens.

During SAR, after primary inoculation, within 4–6 hours, in the inoculated site, where, localized cell death (HR) occurs, plant displays hyponastic response. 3 mM N-fed WT plants showed clear hyponasty after primary inoculation, whereas 0.1 mM did not display such response (Fig. S1c). Hyponastic response is accompanied by SA release, ethylene burst and other auxin transporters present in the leaf lamina and leaf blade to let the petiole elongate (Shah and Zeier 2013). Examination of symptoms in response to primary and secondary challenge (Fig. 2a; Fig. S1b) in 3 mM and 0.1 mM N-fed plants revealed that in 3 mM WT plants the defense response after secondary challenge was more rapid, robust and even long-lasting till 5 days post challenge (data not shown) whereas, 0.1 mM WT plants showed

chlorotic and necrotic lesions, disease symptoms accompanied with reduced electrolyte leakage and increased CFU suggesting that NO_3^- concentration plays a role in LAR and SAR development. Mock infiltrated WT plants (primary and secondary inoculation-both 10 mM $MgCl_2$) does not show any symptoms of wound induced cell death in the treatments (Fig. S2a) during SAR.

In our experiments, roots were treated with *Trichoderma* hence, we examined high and low affinity transporters to see whether *Trichoderma* modulates these transporters to enhance N uptake under low nitrate. HATs operates at <1 mM N concentrations hence, the expression of these transporters were examined in the presence or absence of *Trichoderma*. *NRT2.1* is the main HAT, localized at the plasma membrane. Previously, it was shown that these transporters becomes active during N starvation and is severely inhibited when reduced nitrate sources such as glutamine or ammonium are provided (Dechorgnat *et al.* 2012). Similarly, *NRT2* also induced under low N (Dechorgnat *et al.* (2012). In response to *Trichoderma* treatment, a rapid re-programmed *NRT2.1, NRT2.2* transcripts in both 0.1 and 3 mM was observed during *Pst* inoculation. There was extremely high levels of *NRT2.1* expression in 0.1 mM NO_3^- in both inoculated and uninoculated leaves (Fig. 6b), suggesting that *Trichoderma* can induce HATs at very high levels under low NO_3^-. This feature is very important since plants are NO_3^- starved. *NRT2.1* involvement was further evidenced by the fact that *nrt2.1* mutant produced less protein under low N and *Trichoderma* was unable to increase protein content in this mutant (Fig. 6f). This mutant become very much susceptible to LAR and SAR (Fig. 13).

LATs operates relatively under high N concentrations >50 mM. In our study
we examined *CLCa* and *NPF1.2*. In both cases, we found slightly enhanced
expression of these genes in response to Trichoderma growth. The expression
levels have increased further in inoculated and uninoculated leaves suggesting that
Trichoderma can induce expression of LATs even when grown under low NO_3^-
suggesting active participation of this opportunistic fungus in activating nitrate
transporters proteins irrespective of their affinity (Fig. 5) to nitrate.

We further unraveled the mechanism behind increased expression of LATs
and HATs mediated by *Trichoderma*. One of the features of *Trichoderma* is the
induction of short-term NO and ROS. These molecules play important role in
induction of plant defense responses (Gupta *et al.* 2014). We suspected the role of
NO in activation genes of these transporters. *Trichoderma* induced increase
expression of *NRTs* are most likely mediated by short term increase in NO upon
Trichoderma inoculation. An increased NO was observed in WT but not in *nia1,2*
mutant suggesting that the NO production is mediated by nitrate reductase. The NO
elicited by *Trichoderma* probably responsible for the increased expression of HATs
as our experiments revealed that in *nia1,2* and Hb^+, despite of growing in the
presence of *Trichoderma* were unable to induce HATs they showed decline in
protein levels (Fig. 8c). *Trichoderma* triggers SA dependent SAR pathway (Pieterse
et al. 2014). *Trichoderma* spp. induced ROS is the plant's resistance response
against many biotic stressors (Asmawati *et al.* 2017). It also plays an important role
in hypersensitive cell death together with NO (Durner and Klessig 1999, Dorey *et al.*
1999). Among ROS, H_2O_2 is the most stable form of ROIs that play an important role
as signal transducer in the plant cell death process (Pieterse *et al.* 2014), and acts
as key modulators of NO in triggering cell death. As shown in Fig. 10a, it is observed
that, T203 treated low N fed plants displayed increased H_2O_2 levels (Fig. S4 a&b),
thus enabling the SAR initiation and the establishment stage. Superoxides (O_2^{-}) are
mainly produced via mitochondrial electron transport and NADPH oxidase chain
during stress. Pathogens usually employ this mitochondrial disturbance as a strategy
to suppress host immunity and thus, increased ROS may contribute some resistance
to plants against pathogen attack (Torres *et al.* 2002, 2005). Earlier, it was shown
that the extracellular elicitors isolated from *Trichoderma viridae* also induces the O_2^{-}
levels (Calderon *et al.* 1994). Here, we also found that T203 grown, 0.1 mM NO_3^-,

Pst inoculated plants show higher $O_2^{-\cdot}$ levels than untreated ones (Fig. 10b, Fig.
S4c), suggesting that Trichoderma can enhance $O_2^{-\cdot}$ production during infection
which can aid in defense.

Consequently, the expression of defense marker genes such as *PR1*, *PR2*,
PR5 and *PAL1* (Fig. 11) were also highly induced in the presence of *Trichoderma* in
low N-fed plants suggesting that Trichoderma mediated ROS, NO along with
increased N probably responsible for higher induction of these genes. Martinez-
Medina *et al.* (2013), reported several Trichoderma strains are known to induce
systemic responses by acting as a "short circuit" in plant defense signaling. *PAL1* is
an important marker gene in SA mediated defense (Kim and Hwang 2014) and
accumulates in cells undergoing HR (Dorey *et al.* 1997) and to be essential for local
and systemic resistance (Delaney *et al.* 1994). Apart from showing defense, *PAL1* is
also an important precursor in phenylpropanoid pathway, which is induced under
nutrient stress (Lea *et al.* 2007). Upon T203 pre-treatment to low N fed plants, *PAL1*
expression levels were maintained in the inoculated as well as uninoculated
systemic leaves which revealed that T203 induces systemic responses in
uninoculated organs by mobilizing *PAL1* transcripts (Brotman *et al.* 2012; Shoresh
and Harman 2008; Yedidia *et al.* 2003). SAR response is associated with a specific
set of SAR genes, including genes encoding pathogenesis related (PR) proteins
(Pieterse *et al.* 1996). *PR* gene activates and accumulates during SAR (Fig. 11b,c,
and d) which correlated with findings of Brotman *et al.* (2012) and Pieterse *et al.*
(1996).

In our study, SAR regulatory genes *DIR1*, *NPR1* and *TGA3* are induced in the
presence of Trichoderma under low N stress. The *npr1* mutant compromised in LAR
and SAR in the presence of *Trichoderma* suggesting the role of this gene in
enhancing defense. In *nahg* mutant, defense responses are slightly enhanced in the
presence of Trichoderma despite of reduced SA levels, suggesting that apart from
SA other factors such as increased NO, ROS probably responsible for slight
enhancement of defense in this mutant. Overall this study proved optimum N is
required for LAR and SAR. Trichoderma can enhance N uptake via modulating N
transporters via eliciting short term NO under nitrate nutrition. The enhanced N
uptake plays a role in enhancing SA levels and defense gene expression in local and
distal levels to increase overall plant defense. These defense responses are not

activated in *npr1, nahg, nrt2.1, Hb+* and *nia1,2* mutant providing strong evidence that Trichoderma mediated enhanced resistance under low nitrate involves, NO, ROS, SA.

Acknowledgements

Authors acknowledge Ms. Aanchal Chaudhary for providing virulent strain of *Pst* DC3000. We thank Dr. Yariv Brotman for providing the *T203* strain of Trichoderma. Seeds of *nahG* and *npr1* mutants were kindly provided by Prof. A.K. Nandi, School of Life Sciences, JNU, New Delhi, seeds of non-symbiotic hemoglobin over-expressing lines (Hb+) was kindly provided by Prof. Kim Hebelstrup, *nia1,2* (NASC-6936) and *nrt2.1* mutants were procured from ABRC. The protocol of SA estimation was also provided by Prof. A.K. Nandi. This research was funded by Science and Engineering Research Board (SERB), DST, GOI, New Delhi (National Post Doctoral Fellowship to AW), UGC, New Delhi (SRF to PKP) and SERB-ECR (Early Career Research Award) and DBT-IYBA award to KJG. Authors greatly acknowledge Central Instrumentation Facility (CIF) of the Institute (NIPGR, New Delhi).

Figure legends

Fig. 1: **LAR response elicited by *Pst*DC3000/*avrRpm1* is compromised in low nitrate grown WT plants**

(a) *Pst*DC3000-*avrRpm1* mediated HR observed in 3mM and 0.1 mM NO_3^- grown WT plants at different time points. An early HR response observed in 3 mM *Pst* treated WT plants immediately after 4 hpi whereas 0.1 mM *Pst* treated WT plants showed extensive leaf discoloration and necrotic lesions shown by red arrows at 24 and 48 hpi. **(b)** Electrolyte leakage from leaf areas of 0.1 mM and 3 mM NO_3^- plants infiltrated with *Pst* and 10 mM $MgCl_2$. Data are mean values ± SE. A significant difference between all treatments is analyzed by t-test at $p < 0.001$ (***), $p < 0.01$ (**) and $p < 0.05$ (*) with 3 mM $MgCl_2$ as control. **(c)** Bacterial number in terms of colony forming units (CFU) in *Pst* infiltrated leaves of 0.1 mM and 3 mM NO_3^- plants

immediately after infiltration (0 hpi). Data are mean values ± SE. A significant difference between all treatments is analyzed by t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*) with 3 mM nitrate as control. **(d)** Relative *PR1* transcript levels from 0.1 and 3 mM *Pst* treated leaves at different time points. Data are mean values ± SE. Asterisks indicate statistical significance between 0.1 mM and 3 mM NO_3^- plants is analyzed by t-test at p<0.001 (***), p<0.01 (**). All the results obtained are representative of three independent experiments. Representative images of leaves after *Pst* infiltration taken at respective time points is shown here.

Fig. 2: **Low nitrate grown WT plants show susceptible systemic responses during SAR**

(a) Lesion development post primary inoculation (first panel) and secondary challenge (second panel) in inoculated and uninoculated leaves of 0.1 mM and 3 mM NO_3^- plants. A characteristic lesion pattern in inoculated leaves of both 0.1 and 3 mM NO_3^- plants is noted **(b)** Histochemical staining for the detection of HR- mediated cell death post secondary challenge in inoculated and uninoculated leaves of 0.1 mM and 3 mM NO_3^- plants. Scale bar- 1mm **(c)** Electrolyte leakage observed post primary (first panel) inoculation and secondary challenge (second panel) in inoculated and uninoculated leaves of 0.1 mM and 3 mM NO_3^- plants. A significant difference between all treatments is analyzed by t-test at p<0.05 (*) with 3 mM WT $MgCl_2$ as control **(d)** Bacterial number in log CFU observed post secondary challenge in inoculated leaves of 0.1 mM and 3 mM NO_3^- plants. A significant difference between all treatments is analyzed by t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*).

Fig. 3: **Trichoderma supplementation enhances LAR response in low nitrate grown WT plants**

(a) *Pst*DC3000-*avrRpm1* induced HR in 0.1 and 3 mM WT plants at different time points after Trichoderma (T203) supplementation. The images are representative of three independent experiments. First panel shows the mock infiltrated leaves i.e. control leaves treated with 10 mM $MgCl_2$ and second panel shows *Pst* infiltrated leaves of 0.1 and 3 mM NO_3^- concentrations **(b)** Electrolyte leakage from 0.1 mM and 3 mM WT + T203 treated plants infiltrated with *Pst* and 10 mM $MgCl_2$. Data are mean values ± SE. A significant difference between all treatments is analyzed by t-

test at p<0.001 (***) and p<0.01 (**) with 3 mM WT-MgCl$_2$ as control **(c)** Bacterial

number in terms of log CFU in *Pst* infiltrated leaves of T203 treated 0.1 mM and 3

mM NO$_3^-$ WT plants. Data are mean values ± SE. A significant difference between

all treatments is analyzed by t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*) **(d)**

Relative *PR1* transcript levels from 0.1 and 3 mM *Pst* treated leaves of T203 treated

WT plants at different time points (hpi). Data are mean values ± SE. Asterisks

indicate statistical significance between 0.1 mM and 3 mM plants is analyzed by t-

test p<0.001 (***) and p<0.01 (**), by taking 3 mM plants as control.

Fig. 4: **Trichoderma enhances SAR response in low nitrate plants**

(a) HR development post primary inoculation (first panel and secondary challenge

(second panel) in inoculated and uninoculated leaves of 0.1 mM and 3 mM T203

treated WT plants. Note a characteristic lesion pattern in inoculated leaves of both

0.1 and 3 mM plants

(b) Histochemical staining for the detection of HR-mediated cell death post

secondary challenge in inoculated and uninoculated leaves of 0.1 mM and 3 mM

NO$_3^-$, T203 treated WT plants. Scale bar- 1 mm **(c)** Electrolyte leakage observed

post primary (first panel) inoculation and secondary challenge (second panel) in

inoculated and uninoculated leaves of 0.1 mM and 3 mM T203 treated WT plants. A

significant difference between all treatments is analyzed by t-test at p<0.001 (***),

p<0.01 (**) and p<0.05 (*) with 3 mM WT MgCl$_2$ as control **(d)** Bacterial number in

log CFU observed post secondary challenge in inoculated leaves of 0.1 mM and 3

mM, T203 treated WT plants. A significant difference between 0.1 and 3 mM NO$_3^-$ is

analyzed by t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*).

Fig. 5: **Expression profile of low affinity nitrate transporters genes (LATs)**

during SAR

(a) Relative expression of *CLCa* gene in WT shown in (I panel) and WT+T203 shown

in (II panel) grown under 0.1 mM and 3 mM NO$_3^-$ concentration, post secondary

challenge in both inoculated and un-inoculated leaves **(b)** Relative expression of

NPF1.2 gene in WT shown in (I panel) and WT+T203 shown in (II panel) grown

under 0.1 mM and 3 mM NO$_3^-$ concentration post secondary challenge in both

inoculated and un-inoculated leaves. For all the target genes, fold expression values

are means (n=3) ± SE. A significant difference between all treatments is analyzed by

t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*) with 3 mM un-inoculated leaves as control in each gene.

Fig. 6: Expression profile of high affinity nitrate transporters genes (HATs) during SAR and response of genotypes (WT and *nrt2.1* mutant) on priming effect of 24 h-Trichoderma pre-treatment w.r.t *PR1* and *NRT2.2* expression on low nitrate WT plants

(a) Relative expression of *NRT2.1* in WT shown in (I panel) and WT+T203 shown in (II panel) grown under 0.1 mM and 3 mM NO_3^- concentration post secondary challenge in both inoculated and uninoculated leaves **(b)** Relative expression of *NRT2.2* in WT shown in (I panel) and WT+T203 shown in (II panel) grown under 0.1 mM and 3 mM NO_3^- concentration post secondary challenge in both inoculated and uninoculated leaves. **(c)** Relative expression of *NRT2.4* in WT shown in (I panel) and WT+T203 shown in (II panel) grown under 0.1 mM and 3 mM NO_3^- concentration post secondary challenge in both inoculated and un-inoculated leaves. For all the target genes, fold expression values are means (n=3) ± SE. A significant difference between all treatments is analyzed by t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*) with 3 mM un-inoculated leaves as control in each gene.

 (d) Relative *PR1* expression in roots of WT and *nrt2.1* plants grown under low (0.1 mM) nitrate concentration (with and without T203 treatment; 24 h). A significant difference between with and without T203 treatment is analyzed by t-test at p<0.01 (**) and p<0.05 (*) **(e)** Relative *NRT2.2* expression in roots of WT and *nrt2.1* plants grown under 0.1 mM NO_3^- concentration (with and without T203 treatment; 24 h). **(f)** Protein levels in WT and *nrt2.1* plants grown under 0.1 mM NO_3^- concentration in the presence or absence of T-203. A significant difference between with and without T203 treatment is analyzed by t-test at p<0.01 (**) and p<0.05 (*).

Fig. 7: Visualization of nitric oxide by diaminofluorescein (DAF) fluorescence

Nitric oxide estimation by diaminofluorescein (DAF-FM) fluorescence under 0.1 and 3 mM NO_3^- concentrations in I. WT, II. *nia1,2* and III. cPTIO (100 µM) grown WT plants during Trichoderma supplementation The experiment was performed three times independently with similar results.

Fig. 8: Response of the genotypes during early stages of Trichoderma inoculation on *NRT2.1, 2.2* expression, protein levels and *PR1* expression

(a) Relative *NRT2.1* expression in roots of WT, Hb+, *nia1,2* and *nrt2.1* plants grown under 0.1 mM and 3 mM NO_3^- concentrations, with and without T203 treatment, given for 5 minutes and 24 h to the plants. A significant difference between with and without T203 treatment is analyzed by t-test at $p<0.01$ (**) and $p<0.05$ (*) **(b)** Relative *NRT2.2* expression in roots of WT, Hb+, *nia1,2* and *nrt2.1* plants grown under 0.1 mM and 3 mM NO_3^- concentrations, with and without T203 treatment, given for 5 minutes and 24 h to the plants. A significant difference between with and without T203 treatment is analyzed by t-test at $p<0.01$ (**) and $p<0.05$ (*) **(c)** Protein levels measured in WT, Hb+ and *nia1,2* seedlings grown under 0.1 mM NO_3^- for 15 days in vertical plates using Bradford's assay. T203 treatment was given for 24 h to the plants. A significant difference between with and without T203 treatment is analyzed by t-test at $p<0.01$ (**) and $p<0.05$ (*) **(d)** *PR1* gene expression in WT, Hb+, *nia1,2* and *nrt2.1* under 0.1 mM NO_3^- plants, with and without *Trichoderma* treatment, given for 5 minutes and 24 h to the plants. A significant difference between with and without T203 treatment is analyzed by t-test at $p<0.01$ (**) and $p<0.05$ (*).

Fig. 9: Response of WT and NO mutants (Hb+ and *nia1,2*) grown under 0.1 mM NO_3^- during LAR and SAR

In LAR response **(a)** HR phenotype in inoculated leaves of WT, Hb+ and *nia1,2* plants grown under 0.1 mM NO_3^- concentration at 0, 6, 24 and 48 hpi **(b)** *In planta* bacterial growth (Log CFU) at 6, 24 and 48 hpi, asterisks indicate a significant difference ($p < 0.01$; Student's t test) taking WT as control **(c)** Relative *PR1* expression in infiltrated leaves during localized infiltration. Values are means (n=3) ± SE. Asterisks indicate a significant difference ($p < 0.01$; student's t-test) taking 0 hpi as control. In SAR response **(d)** HR phenotype in inoculated and uninoculated leaves of WT, Hb+ and *nia1,2* plants grown under 0.1 mM NO_3^- concentration post secondary challenge **(e)** Bacterial population represented by CFU per leaf disc at 6, 24 and 48 hpc, asterisks indicate a significant difference ($p < 0.01$; Student's t test) taking 3 mM WT as control **(f)** Relative *PR1* gene expression in infiltrated leaves

post secondary challenge. Values are means (n=3) ± SE. Asterisks indicate a
significant difference (p < 0.01; Student's t test) taking 0 hpc as control.

Fig. 10: Detection of ROS by measuring H_2O_2 and O_2^{-} in 0.1 mM and 3 mM NO_3^{-}
WT and Trichoderma grown WT plants during SAR response

(a) DAB staining used to measure H_2O_2 levels in the inoculated leaves and
uninoculated leaves after 3 days of challenge inoculation at 0, 6, 24 and 48 hpc.
Representative leaves are shown, and the experiment was repeated three times with
10 leaves each. Scale bar =1 mm.

(b) NBT staining used to measure O_2^{-} levels in the inoculated leaves and
uninoculated leaves post challenge inoculation at 0, 6, 24 and 48 hpc.
Representative leaves are shown, and the experiment was repeated three times with
10 leaves each. Scale bar =1 mm.

Fig. 11: **Expression profiles of defense related genes during SAR**

Relative expression of defense related genes in wild-type (WT) shown in (I panel)
and Trichoderma treated wild-type plants (WT+T203) shown in (II panel) grown
under low (0.1 mM) and optimum (3 mM) NO_3^{-} concentration post secondary
challenge of *Pst*DC3000 in both inoculated and uninoculated leaves **(a)** Relative
expression of *PAL1* **(b)** Relative expression of *PR1* **(c)** Relative expression of *PR2*
(d) Relative expression of *PR5*. For all the target genes, fold expression values are
means (n=3) ± SE. A significant difference between all treatments is analyzed by t-
test at p<0.001 (***), p<0.01 (**) and p<0.05 (*) with 3mM uninoculated leaves as
control in each gene.

Fig. 12: **Expression profiles of regulatory SAR genes**
Relative expression of defense genes in WT shown in (I panel) and WT+T203 shown
in (II panel) grown under low (0.1 mM) and optimum (3 mM) NO_3^{-} concentration post
secondary challenge of *Pst*DC3000 in both inoculated and uninoculated leaves **(a)**
Relative expression of *DIR1* **(b)** Relative expression of *NPR1* **(c)** Relative expression
of *SARD1* **(d)** Relative expression of *TGA3*. For all the target genes, fold expression
values are means (n=3) ± SE. A significant difference between all treatments is
analyzed by t-test at p<0.001 (***), p<0.01 (**) and p<0.05 (*) with 3mM
uninoculated leaves as control in each gene.

Fig 13: (a) LAR disease symptoms in *nrt2.1* and *npr1* mutant in 0.1 and 3 mM nitrate

(b) Bacterial growth in WT, *nrt2.1* and *npr1* mutant (c) SAR response in *nrt2.1* and

npr1 mutant in 0.1 and 3 mM nitrate (d) Bacterial growth in WT, *nrt2.1* and *npr1*

mutant during SAR. A significant difference between with and without T203

treatment is analyzed by t-test at p<0.01 (**) and p<0.05 (*).

Fig. 14: **SA Accumulation during SAR, and response of *nahg* mutant in the**
presence and absence of Trichoderma during SAR

(a) Total (free glucose conjugate) SA levels ($\mu g\ g^{-1}$ fresh weight) were determined

post challenge inoculation (24 hpc) in inoculated and uninoculated leaves of 0.1 mM

and 3 mM NO_3^- fed WT plants, with and without Trichoderma. Two samples for each

treatment were collected and the range of the mean is represented by the error bars.

Asterisks indicate mean values that are significantly different according to ($p < 0.01$;

Student's t test) with 3 mM uninoculated leaves as control (b) HR phenotype in

inoculated and un-inoculated distal leaves of *nahg* mutants grown under 0.1 and 3

mM NO_3^- concentration post secondary challenge at 6, 24 and 48 hpc **(c)** Bacterial

number represented by Log CFU per leaf disc at 6, 24 and 48 hpc from the

inoculated leaves post secondary challenge **(f)** *PR1* gene expression in inoculated

and uninoculated leaves of *nahg* mutants grown under 0.1 and 3 mM NO_3^- ,with and

without *Trichoderma* treatment, post secondary challenge. Values are means (n=3)

± SE. A significant difference between with and without T203 treatment is analyzed

by t-test at p<0.01 (**) and p<0.05 (*).

Supplementary Information

Supplementary Fig 1: (a) The growth pattern and phenotype of WT and T203

919 treated WT plants in 3 mM and 0.1 mM NO_3^- nutrition; (b). Effect of post secondary

920 challenge on plant phenotype, 1°- primary inoculation, 2°-secondary challenge and

921 U- uninoculated leaves; (c) Hyponastic response (petiole elongation) shown by 0.1

922 mM and 3 mM NO_3^- plants after primary inoculation.

Supplementary Fig 2: (a) Electrolyte leakage from mock infiltrated plants in SAR;

924 Morphological growth parameters observed in WT and T203 treated WT plants; (b)

925 Leaf number; (c) Biomass measured as fresh weight (FW); (d) Total chlorophyll

926 content in *Pst* treated and untreated plants.

Supplementary Fig 3: **(a)** Enlarged microscopic view of the *Pst* inoculated leaves

showing macrobursts of H_2O_2 shown by red arrows after DAB staining. Scale bar =

500 µm. **(b)** Quantification of DAB staining as intensity per stained area from ten
leaves per treatment measured using ImageJ in arbitrary units with the mean values
± SE. **(c)** Quantification of NBT staining as intensity per stained area from ten leaves
per treatment measured using ImageJ in arbitrary units with the mean values ± SE.
Supplementary Fig 4: **(a)** Enlarged microscopic view of the necrotic areas of only
Pst treated 0.1 and 3 mM WT and T203 treated WT plants after trypan blue staining.
Scale bar- 500µm **(b)** Asterisks indicate statistically significant differences between
treatments and the time points using Tukey's post-hoc test for pairwise comparisons
(α = 0.05) in only *Pst* treated leaves. Results are presented as necrotized leaf area
compared with the total surface of leaf analyzed by ImageJ (ver 3.2) and represent
means ± SE of 6 leaves per treatment.
Supplementary Table 1: List of primers

References:

Alvarez ME, Pennell RI, Meijer PJ, Ishikawa A, Dixon RA, Lamb C. 1998.
Reactive oxygen intermediates mediate a systemic signal network in the
establishment of plant immunity. *Cell* **92:** 773-784.

Alves M, Dadalto S, Gonçalves A, de Souza G, Barros V, Fietto L. 2014.
Transcription factor functional protein-protein interactions in plant defense
responses. *Proteomes* **2:** 85-106.

Asmawati L, Widiastuti A, Sumardiyono C. 2017. Induction of Reactive Oxygen
Species by Trichoderma spp. Against Downy Mildew in Maize. In *Proceeding of the*
1st International Conference on Tropical Agriculture. Springer, Cham **:**139-146.

Boudsocq S, Niboyet A, Lata JC, Raynaud X, Loeuille N, Mathieu J, Blouin M,
Abbadie L, Barot S. 2012. Plant preference for ammonium versus nitrate: a
neglected determinant of ecosystem functioning?. *The American Naturalist* **180:** 60-
69.

Brotman Y, Lisec J, Méret M, Chet I, Willmitzer L, Viterbo A. 2012. Transcript
and metabolite analysis of the Trichoderma-induced systemic resistance response to
Pseudomonas syringae in Arabidopsis thaliana. *Microbiology* **158:** 139-146.

Brotman Y, Gupta KJ, Vetribo A. (2010) *Trichoderma:* Quick Guide. *Current*
Biology **20:** 390-391.

Calderon AA, Zapata JM, Barcelo AR. 1994. Peroxidase-mediated formation of
resveratrol oxidation products during the hypersensitive-like reaction of grapevine
cells to an elicitor from Trichoderma viride. *Physiological and Molecular Plant*
Pathology **44:** 289-299.

Cameron RK, Dixon RA, Lamb CJ. 1994. Biologically induced systemic acquired
resistance in Arabidopsis thaliana. *The Plant Journal* **5:** 715-725.

Cameron RK, Paiva NL, Lamb CJ, Dixon RA. 1999. Accumulation of salicylic acid
and PR-1 gene transcripts in relation to the systemic acquired resistance (SAR)
response induced by Pseudomonas syringae pv. tomato in
Arabidopsis. *Physiological and Molecular Plant Pathology* **55:** 121-130.

Cao H, Glazebrook J, Clarke JD, Volko S, Dong X. 1997. The Arabidopsis NPR1
gene that controls systemic acquired resistance encodes a novel protein containing
ankyrin repeats. *Cell* **88:** 57-63.

Carella P, Isaacs M, Cameron RK. 2015. Plasmodesmata-located protein
overexpression negatively impacts the manifestation of systemic acquired resistance
and the long-distance movement of Defective in Induced Resistance1 in A
rabidopsis. *Plant Biology* **17:** 395-401.

Champigny MJ, Isaacs M, Carella P, Faubert J, Fobert PR, Cameron RK. 2013.
Long distance movement of DIR1 and investigation of the role of DIR1-like during
systemic acquired resistance in Arabidopsis. *Frontiers in Plant Science* **4:** 230.

Contreras-Cornejo HA, Macías-Rodríguez L, del-Val E, Larsen J. 2016.
Ecological functions of Trichoderma spp. and their secondary metabolites in the
rhizosphere: interactions with plants. *FEMS microbiology ecology* **92:** fiw036.

Daudi A, O'Brien JA. 2012. Detection of hydrogen peroxide by DAB staining in
Arabidopsis leaves. *Bio-Protocol* **2:** 1-4.

Dechorgnat J, Patrit O, Krapp A, Fagard M, Daniel-Vedele F. 2012.
Characterization of the Nrt2. 6 gene in Arabidopsis thaliana: a link with plant
response to biotic and abiotic stress. *PloS one* **7:** e42491.

Delaney TP, Uknes S, Vernooij B, Friedrich L, Weymann K, Negrotto D, Gaffney
T, Gut-Rella M, Kessmann H, Ward E, Ryals J. 1994. A central role of salicylic acid
in plant disease resistance. *Science* **266:** 1247-1250.

Delledonne M, Xia Y, Dixon R A, Lamb C. 1998. Nitric oxide functions as a signal
in plant disease resistance. *Nature* **394:** 585.

Domínguez S, Rubio MB, Cardoza RE, Gutiérrez S, Nicolás C, Bettiol W,
Hermosa R, Monte E. 2016. Nitrogen metabolism and growth enhancement in
tomato plants challenged with Trichoderma harzianum expressing the Aspergillus
nidulans acetamidase amdS gene. *Frontiers in Microbiology* **7:** 1182.

Dorey S, Baillieul F, Pierrel MA, Saindrenan P, Fritig B, Kauffmann S. 1997.
Spatial and temporal induction of cell death, defense genes, and accumulation of
salicylic acid in tobacco leaves reacting hypersensitively to a fungal glycoprotein
elicitor. *Molecular Plant-Microbe Interactions* **10:** 646–655.

Dorey S, Kopp M, Geoffroy P, Fritig B, Kauffmann S. 1999. Hydrogen peroxide
from the oxidative burst is neither necessary nor sufficient for hypersensitive cell
death induction, phenylalanine ammonia lyase stimulation, salicylic acid
accumulation, or scopoletin consumption in cultured tobacco cells treated with
elicitin. *Plant Physiology* **121:** 163-172.

Durner J, Klessig DF. 1999. Nitric oxide as a signal in plants. *Current Opinion in*
Plant Biology **2:** 369-374.

Fagard M, Launay A, Clément G, Courtial J, Dellagi A, Farjad M, Krapp A,
Soulié MC, Masclaux-Daubresse C. 2014. Nitrogen metabolism meets
phytopathology. *Journal of Experimental Botany* **65:** 5643-5656.

Fernández-Bautista N, Domínguez-Núñez JA, Moreno MC, Berrocal-Lobo M.
2002. Plant tissue trypan blue staining during phytopathogen infection. *Bio-*
Protocol **6**: e2078.

Frungillo L, de Oliveira JFP, Saviani EE, Oliveira HC, Martínez MC, Salgado I.
2013. Modulation of mitochondrial activity by S-nitrosoglutathione reductase in
Arabidopsis thaliana transgenic cell lines. Biochimica et Biophysica Acta -
Bioenergetics **1827:** 239-247.

Garnica-Vergara A, Barrera-Ortiz S, Muñoz-Parra E, Raya-González J, Méndez-Bravo A, Macías-Rodríguez L, Ruiz-Herrera LF, López-Bucio J. 2016. The volatile 6-pentyl-2H-pyran-2-one from Trichoderma atroviride regulates Arabidopsis thaliana root morphogenesis via auxin signaling and ETHYLENE INSENSITIVE 2 functioning. *New Phytologist* **209:** 1496-1512.

Gupta A, Dixit SK, Senthil-Kumar M. 2016. Drought stress predominantly endures Arabidopsis thaliana to Pseudomonas syringae infection. *Frontiers in Plant Science* **7:** 808.

Gupta KJ, Mur LA, Brotman Y. 2014. Trichoderma asperelloides suppresses nitric oxide generation elicited by Fusarium oxysporum in Arabidopsis roots. *Molecular Plant-Microbe Interactions* **27:** 307-314.

Gupta KJ, Brotman Y, Segu S, Zeier T, Zeier J, Persijn ST, Cristescu SM, Harren FJM, Bauwe H, Fernie A.R. 2013. The form of nitrogen nutrition affects resistance against *Pseudomonas syringae pv. phaseolicola* in tobacco. *Journal of Experimental Botany* **64:** 553-568.

Jambunathan N. 2010. Determination and detection of reactive oxygen species (ROS), lipid peroxidation, and electrolyte leakage in plants. *Methods in Molecular Biology* **639:** 292–298.

Jones JD, Dangl JL. 2006. The plant immune system. *Nature* **444:** 323.

Kim DS, Hwang BK. 2014. An important role of the pepper phenylalanine ammonia-lyase gene (PAL1) in salicylic acid-dependent signalling of the defence response to microbial pathogens. *Journal of Experimental Botany* **65:** 2295-2306.

Krapp A, David LC, Chardin C, Girin T, Marmagne A, Leprince AS, Chaillou S, Ferrario-Méry S, Meyer C, Daniel-Vedele F. 2014. Nitrate transport and signalling in Arabidopsis. *Journal of Experimental Botany* **65:** 789-798.

Lamb C, Dixon RA. 1997. The oxidative burst in plant disease resistance. *Annual Review of Plant Biology* **48:** 251-275.

Landrein B, Formosa-Jordan P, Malivert A, Schuster C, Melnyk CW, Yang W, Turnbull C, Meyerowitz EM, Locke JC, Jönsson H. 2018. Nitrate modulates stem

cell dynamics in Arabidopsis shoot meristems through cytokinins. *Proceedings of the National Academy of Sciences* **115:** 1382-1387.

Lea US, Slimestad R, Smedvig P, Lillo C. 2007. Nitrogen deficiency enhances expression of specific MYB and bHLH transcription factors and accumulation of end products in the flavonoid pathway. *Planta* **225:** 1245–1253.

Li H, Hu B, Chu C. 2017. Nitrogen use efficiency in crops: lessons from Arabidopsis and rice. *Journal of Experimental Botany* **68:** 2477-2488.

Liu Y, von Wirén N. 2017. Ammonium as a signal for physiological and morphological responses in plants. *Journal of Experimental Botany* **68:** 2581-2592.

Maldonado AM, Doerner P, Dixon RA, Lamb CJ, Cameron RK. 2002. A putative lipid transfer protein involved in systemic resistance signalling in Arabidopsis. *Nature* **419:** 399.

Martínez-Medina A, Fernández I, Sánchez-Guzmán MJ, Jung SC, Pascual JA, Pozo MJ. 2013. Deciphering the hormonal signalling network behind the systemic resistance induced by Trichoderma harzianum in tomato. *Frontiers in Plant Science* **4:** 206.

Métraux JP, Signer H, Ryals J, Ward E, Wyss-Benz M, Gaudin J, Raschdorf K, Schmid E, Blum W, Inverardi B. 1990. Increase in salicylic acid at the onset of systemic acquired resistance in cucumber. *Science* **250:** 1004-1006.

Mur LA, Brown IR, Darby RM, Bestwick CS, Bi YM, Mansfield JW, Draper J. 2000. A loss of resistance to avirulent bacterial pathogens in tobacco is associated with the attenuation of a salicylic acid-potentiated oxidative burst. *The Plant Journal* **23:** 609-621.

Mur LA, Kenton P, Lloyd AJ, Ougham H, Prats E. 2008. The hypersensitive response; the centenary is upon us but how much do we know?. *Journal of Experimental Botany* **59:** 501-520.

Mur LA, Simpson C, Kumari A, Gupta AK, Gupta KJ. 2017. Moving nitrogen to the centre of plant defence against pathogens. *Annals of botany* **119:** 703-709.

O'Brien JA, Vega A, Bouguyon E, Krouk G, Gojon A, Coruzzi G, Gutiérrez RA. 2016. Nitrate transport, sensing, and responses in plants. *Molecular Plant* **9:** 837-856.

Oliva R, Quibod IL. 2017. Immunity and starvation: new opportunities to elevate disease resistance in crops. *Current Opinion in Plant Biology* **38:** 84-91.

Park SW, Kaimoyo E, Kumar D, Mosher S, Klessig DF. 2007. Methyl salicylate is a critical mobile signal for plant systemic acquired resistance. *Science* **318:** 113-116.

Pieterse CM, Van Wees SC, Hoffland E, Van Pelt JA, Van Loon LC. 1996. Systemic resistance in Arabidopsis induced by biocontrol bacteria is independent of salicylic acid accumulation and pathogenesis-related gene expression. *The Plant Cell* **8:** 1225-1237.

Pieterse CM, Zamioudis C, Berendsen RL, Weller DM, Van Wees SC, Bakker PA. 2014. Induced systemic resistance by beneficial microbes. *Annual Review of Phytopathology* **52:** 347-375

Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner HY, Hunt MD. 1996. Systemic acquired resistance. *The Plant Cell* **8:** 1809.

Shah J, Zeier J. 2013. Long-distance communication and signal amplification in systemic acquired resistance. *Frontiers in Plant Science* ***4:*** 30.

Shoresh M, Harman GE, Mastouri F. 2010. Induced systemic resistance and plant responses to fungal biocontrol agents. *Annual Review of Phytopathology* **48:** 21-43.

Shoresh M, Harman GE. 2008. The molecular basis of shoot responses of maize seedlings to Trichoderma harzianum T22 inoculation of the root: a proteomic approach. *Plant Physiology* **147:** 2147-2163.

Shoresh M, Yedidia I, Chet I. 2005. Involvement of jasmonic acid/ethylene signaling pathway in the systemic resistance induced in cucumber by Trichoderma asperellum T203. *Phytopathology* **95:** 76-84.

Singh V, Roy S, Giri MK, Chaturvedi R, Chowdhury Z, Shah J, Nandi AK. 2013. Arabidopsis thaliana FLOWERING LOCUS D is required for systemic acquired resistance. *Molecular Plant-Microbe Interactions* **26:** 1079-1088.

Smith-Becker J, Marois E, Huguet EJ, Midland SL, Sims JJ, Keen NT. 1998.
Accumulation of salicylic acid and 4-hydroxybenzoic acid in phloem fluids of
cucumber during systemic acquired resistance is preceded by a transient increase in
phenylalanine ammonia-lyase activity in petioles and stems. *Plant Physiology* **116:**
231-238.

Snoeijers SS, Pérez-García A, Joosten MH, De Wit PJ. 2000. The effect of
nitrogen on disease development and gene expression in bacterial and fungal plant
pathogens. *European Journal of Plant Pathology* **106:** 493-506.

Torres MA, Dangl JL, Jones JD. 2002. Arabidopsis gp91phox homologues
AtrbohD and AtrbohF are required for accumulation of reactive oxygen intermediates
in the plant defense response. *Proceedings of the National Academy of Sciences of*
the United States of America **99:** 517–522.

Torres MA, Jones JD, Dangl JL. 2005. Pathogen-induced, NADPH oxidasederived
reactive oxygen intermediates suppress spread of cell death in Arabidopsis thaliana.
Nature Genetics **37:** 1130–1134.

Tsay YF, Chiu CC, Tsai CB, Ho CH, Hsu PK. 2007. Nitrate transporters and
peptide transporters. *FEBS letters* **581:** 2290-2300.

Walker RL, Burns IG, Moorby J. 2001. Responses of plant growth rate to nitrogen
supply: a comparison of relative addition and N interruption treatments. *Journal of*
Experimental Botany **52:** 309-317.

Wang L, Tsuda K, Truman W, Sato M, Nguyen LV, Katagiri F, Glazebrook J.
2011. CBP60g and SARD1 play partially redundant critical roles in salicylic acid
signaling. *The Plant Journal* **67:** 1029-1041.

Wany A, Gupta AK, Kumari, Mishra S, Singh N, Pandey S, Vanvari R,
Igamberdiev AU, Fernie AR, Gupta KJ. 2018. Nitrate nutrition influences multiple
factors in order to increase energy efficiency under hypoxia in Arabidopsis. *Annals of*
Botany (In Press).

Wany A, Kumari A, Gupta KJ. 2017. Nitric oxide is essential for the development of
aerenchyma in wheat roots under hypoxic stress. *Plant, Cell & Environment* **40:**
3002-3017.

Yedidia I, Shoresh M, Kerem Z, Benhamou N, Kapulnik Y, Chet I. 2003. Concomitant induction of systemic resistance to Pseudomonas syringae pv. lachrymans in cucumber by Trichoderma asperellum (T-203) and accumulation of phytoalexins. *Applied and Environmental Microbiology* **69:** 7343-7353.

Zhang Y, Xu S, Ding P, Wang D, Cheng YT, He J, Gao M, Xu F, Li Y, Zhu Z, Li X. 2010. Control of salicylic acid synthesis and systemic acquired resistance by two members of a plant-specific family of transcription factors. *Proceedings of the National Academy of Sciences* USA 107 (42) 18220-18225

Zhou JM, Trifa Y, Silva H, Pontier D, Lam E, Shah J, Klessig DF. 2000. NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the PR-1 gene required for induction by salicylic acid. *Molecular Plant-Microbe Interactions* **13:** 191-202.

Fig 1

(a)

(b)

Fig. 2

Fig. 3

Fig. 4

Fig 5

(a)

(b)

(c)

(d)

Fig 6

Fig7

Fig. 8

(a)

(b)

(c)

(d)

Fig. 9

(a)

(b)

(c)

(d)

(e)

(f)

Fig. 10

(a)

(b)

Fig. 11

I. WT   **II. WT+*T203***

Fig. 12

Fig. 13

(a)

(b)

(c)

(d)

Fig. 14