p62-mediated Selective Autophagy Endows Virus-transformed Cells with Insusceptibility to DNA Damage under Oxidative Stress

Ling Wang[1, 2,*], Mary E. A. Howell[1], Ayrianna Sparks-Wallace[1], Caroline Hawkins[1], Camri Nicksic[1],

Carissa Kohne[1], Kenton H. Hall[1], Jonathan P. Moorman[1, 2, 3,] Zhi Q. Yao[1, 2, 3], Shunbin Ning[1, 2,*]

1. Department of Internal Medicine, Quillen College of Medicine, East Tennessee State University, Johnson City, TN 37614.
2. Center of Excellence for Inflammation, Infectious Diseases and Immunity, Quillen College of Medicine, East Tennessee State University, Johnson City, TN 37614.
3. The HCV/HIV Program, James H Quillen VA Medical Center, Johnson City, TN 37614.

Running title: p62-mediated autophagy promotes DNA damage response in virus-

transformed cells

Key Words: p62, Autophagy, EBV, DDR

 * **Corresponding Authors:**

Ling Wang, Ph.D. Email: wangl3@etsu.edu

Shunbin Ning, Ph.D. Email: nings1@etsu.edu

Center of Excellence for Inflammation, Infectious Diseases and Immunity; Department of Internal Medicine, Quillen College of Medicine, East Tennessee State University, Johnson City, TN 37614

Tel: 423-439-8063; Fax: 423-439-7010

Abstract

DNA damage response (DDR) and selective autophagy both can be activated by reactive oxygen/nitrogen species (ROS/RNS), and both are of paramount importance in cancer development. The selective autophagy receptor and ubiquitin (Ub) sensor p62 plays a key role in their crosstalk. ROS production has been well documented in latent infection of oncogenic viruses including Epstein-Barr Virus (EBV). However, p62-mediated selective autophagy and its interplay with DDR have not been investigated in these settings. In this study, we provide evidence that p62 is upregulated through the ROS-Keap1-NRF2 pathway, and considerable levels of p62-mediated selective autophagy are constitutively induced, in virus-transformed cells. Inhibition of autophagy results in p62 accumulation, and promotes ROS-induced DNA damage and cell death, as well as downregulates the DNA repair proteins CHK1 and RAD51. In contrast, MG132-mediated proteasome inhibition, which induces rigorous autophagy, promotes p62 degradation but accumulation of the DNA repair proteins CHK1 and RAD51. However, small hairpin RNA (shRNA)-mediated p62 depletion or transient expression of p62 in B cell lines had no apparent effects on the protein levels of CHK1 and RAD51. These results indicate that p62 accumulation resulting from autophagy inhibition is prerequisite for p62-mediated proteasomal degradation of CHK1 and RAD51. Furthermore, shRNA-mediated p62 depletion in EBV-transformed lymphoblastic cell lines results in significant increases of endogenous RNF168-γH2AX damage foci and chromatin ubiquitination in the nucleus, indicative of activation of RNF168-mediated DNA repair mechanisms. Finally, we have evaluated endogenous p62 and autophagy levels and their correlation with endogenous ROS and DNA damage levels in B cell population of AIDS lymphoma patients versus age-matched subjects. Our results have unveiled a pivotal role for p62-mediated selective autophagy that governs DDR in the setting of oncogenic virus latent infection, and provide a novel insight into virus-mediated oncogenesis.

Author Summary

Reactive oxygen/nitrogen species (ROS/RNS) can induce both DNA damage response (DDR) and selective autophagy, which play crucial roles in cancer development. The selective autophagy receptor and ubiquitin (Ub) sensor p62 links their crosstalk. However, p62-mediated selective autophagy and its interplay with DDR have not been investigated in latent infection of oncogenic viruses including Epstein-Barr Virus (EBV). In this study, we provide evidence that p62-mediated selective autophagy is constitutively induced in virus-transformed cells, and that its inhibition exacerbates ROS-induced DNA damage, and promotes proteasomal degradation of CHK1 and RAD51 in a p62-dependent manner. However, rigorous autophagy induction results in accumulation of DNA repair proteins CHK1 and RAD51, and p62 degradation. Furthermore, small hairpin RNA (shRNA)-mediated p62 depletion or transient expression of p62 did not affect CHK1 and RAD51 protein levels; rather, shRNA-mediated p62 depletion activates RNF168-dependent DNA repair mechanisms. Our results have unveiled a pivotal role for p62-mediated selective autophagy in regulation of DDR by overriding traditional DDR mechanisms in the setting of oncogenic virus latent infection, and provide a novel insight into the etiology of viral cancers.

Introduction

p62 (also named EBIAP, ZIP3, SQSTM1/Sequestosome-1), a human homolog of mouse ZIPs

(Zeta PKC-interacting proteins), is well known as a selective autophagy receptor and a ubiquitn

sensor, which controls myraid cellular processes, including redox homeostasis, DNA damage

response (DDR), cancer development, aging, inflammation and immunity, osteoclastogenesis, and

obesity, with or without the involvement of autophagy [1-3].

Autophagy, with either non-selective (random) or selective fashion, is a unique intracellular

process that engulfs damaged and even functional cellular constituents and delivers them to

lysosomes for digestion and recycling in the cytosol under diverse stresses, such as nutrient

deprivation, viral replication, cancer hypoxia, and genotoxic stress. Autophagy is thereby a crucial

cellular machinery conserved from yeast to higher eukaryotes that maintains organ metabolism

and cell survival, and functions as either tumor suppressor at early stage or promotor at late stage

[4-6]. Distinct from non-selective autophagy, selective autophagy sort specific substrates to

lysosomes, and is mediated by an increasing pool of receptors, including p62, NBR1, TAX1BP1,

NDP52, OPTN, TRIMs and TOLLIP [7-9].

Reactive oxygen/nitrogen species (ROS and RNS), the major cause of endogenous DNA

damage, can be produced in chronic viral infections, in which viral replication is generally absent

[10]. They can directly modify DNA and generate different levels of lesions, including double strand

breaks (DSBs) [11, 12]. Eukaryotic organisms have developed sophisticated strategies to repair

DNA damage to ensure genomic integrity, with homologous recombination (HR) and

nonhomologous end joining (NHEJ) being two non-redundant repair mechanisms for DSBs [13].

Unrepaired DSBs, however, incite chronic inflammation, resulting in genomic instability that

promotes malignant transformation under certain conditions [14].

ROS/RNS also induce p62 expression through the Keap1-NRF2 pathway, licensing the induction of p62-mediated selective autophagy. Mounting evidence indicates that DDR and selective autophagy closely crosstalk in response to oxidative stress, in which p62 plays a key role [15]. While p62 inhibits DNA damage repair, p62-mediated selective autophagy promotes DNA repair by targeting ubiquitinated substrates including p62 itself for degradation in cancer cells [16, 17], which usually harbor deficient traditional DNA repair mechanisms and heavily rely on autophagy as an alternative repair mechanism for survival [18, 19]. In this sense, p62-mediated selective autophagy, which is activated upon DNA damage caused by various stresses such as conventional chemotherapeutic agents, allows these cancer cells to escape DNA damage-induced cell death [20, 21].

ROS/RNS overproduction, deregulation of host DDR machinery, and chronic inflammation, are the most common features of viral persistent infections, and together with non-selective autophagy, have also been documented in latency of herpesviruses including Epstein-Barr Virus (EBV) [22-29]. Moreover, we and others have provided overwhelming evidence supporting that EBV latent infection reprograms the host ubiquitin machinery for its own benefits [30-32], including the employment of linear ubiquitin chain assembly complex (LUBAC)-mediated ubiquitination to modulate LMP1 signal transduction [33]. However, as the major selective autophagy receptor and a ubiquitin sensor, p62 and its relationship with EBV latency and oncogenesis have never been investigated. In our recent publication, our findings imply a role for the p62-autophagy interplay in ROS-elicited DDR in EBV latency [34].

In this study, we aimed to investigate the potential role of p62-mediated selective autophagy in regulating DDR in EBV latent infection. Our results show that p62 is transcriptionally induced by the ROS-Keap1-NRF2 pathway, and that a well-balanced basal level of p62-mediated selective autophagy is essential for maintaining genomic stability in EBV latency.

Results

p62-mediated selective autophagy is constitutively induced in viral latency and correlates with ROS-Keap1-NRF2 pathway activity

Our recent findings have shown that treatment of EBV$^+$ cells with the calcium ionophore ionomycin, which raises the intracellular level of calcium (Ca^{2+}) essential for ROS production, elevates the protein levels of both p62 and LC3b-II (the smaller cleavage product of LC3b; Generally represents an autophagosomal marker in mammalian cells), and induces DNA damage; autophagy deficiency also elevates p62 protein levels [34]. Since both p62 and LC3b are targeted by autophagy for degradation, their turnover represents the autophagic flux (autophagic degradation activity) [35, 36]. These results imply that the p62-autophagy interplay may be involved in oxidative stress in EBV latent infection.

We thus sought to evaluate the correlation between intracellular ROS, and p62 and autophagy levels in EBV latency programs. Results show that, although nearly 100% cells of each tested virus-associated cancer cell line produce ROS, their levels, as indicated by mean of fluorescence intensity (MFI), are higher in SavIII, JiJoye, and MT4, compared to SavI, P3HR1, and CEM, respectively (Fig 1, A). Correspondingly, the cell lines with higher ROS production remarkably express higher p62 at both protein and mRNA levels (Fig 1, B and C). Furthermore, the basal levels of p62-mediated selective autophagy activity, as indicated by both the cleaved LC3b product LC3b-II and phosphorylation of p62(Ser403), correspond to the endogenous ROS levels, and are readily detectable by immunoblotting in EBV type III latency and human T-cell leukemia virus-1 (HTLV1)-transformed MT4 cell line (Fig 1, B). Phosphorylation of p62(Ser403) promotes p62-Ub binding for activation of p62-selective autophagy [37]. These results indicate that a basal level of p62-mediated selective autophagy is constitutively induced and correlates with the endogenous ROS level in viral latency.

The antioxidant transcription factor NRF2 is spontaneously degraded by the ubiquitin (Ub) E3 ligase complex Keap1/Cul3/RBX1 under normoxia. ROS/oxidative stress triggers autophagic degradation of Keap1, resulting in the accumulation and activation of NRF2, which then induces p62 expression [38]. Thus, we evaluated the Keap1-NRF2 pathway activities, indicated by Keap1 and NRF2 expression levels, in these cell lines. As indicated in Fig 1, B, the endogenous p62 levels positively correlate with the Keap-NRF2 pathway activities, strongly suggesting that the Keap1-NRF2 pathway induces p62 expression at least in part in viral latency.

Together, these results indicate that p62-mediated selective autophagy is constitutively induced in oncovirus latency, and correlates with the endogenous ROS-Keap1-NRF2 pathway activity.

The ROS-Keap1-NRF2 pathway contributes to p62 expression and activation of p62-mediated autophagy in viral latency

We next aimed to verify the induction of p62 by the ROS-Keap1-NRF2 pathway in viral latency. To this end, we first treated SavI and SavIII cells with the clinical topoisomerase II inhibitor doxorubicin (Doxo), which generates the highest level of mitochondrial ROS causing DSBs [39]. Results show that Doxo augments the activities of the Keap1-NRF2 pathway in both cell lines in a time-dependent manner. Interestingly, p62 protein levels and p-p62(S403) are increased by Doxo treatment in SavI cells, but p62 protein levels are decreased in SavIII cells. Moreover, LC3b-II is increased in both cell lines but decreased at late stage in SavIII cells (Fig 2, A). These results are consistent with the notion that considerable levels of ROS are required for induction of p62 and mediated autophagy, but excess ROS and selective autophagy result in autophagic degradation of p62 and LC3b in that both are targets of p62-mediated selective autophagy [35, 36].

We then used 3-amino-1,2,4-triazole (3-AT) to inhibit the endogenous activities of catalase, an enzyme converting H_2O_2 to H_2O+O_2, in cell lines with lower ROS levels to elevate their

endogenous ROS levels. Then, we evaluated the Keap1-NRF2 pathway activity, and p62,

autophagy, and DNA damage levels. Results show that 3-AT treatment substantially elevates

endogenous levels of the Keap1-NRF2 pathway activities and p62 expression at both protein and

mRNA levels, and also induces p62(S403) phosphorylation, autophagy and the DNA damage

hallmark γH2AX that are readily detectable (Fig 2, B). Accumulation of LC3-II does not necessarily

reflect an increased autophagic activity; instead it may represent its decreased clearance due to

the blockage of autophagic degradation. Thus, we further measured autophagy flux, as indicated

by MFI, by flow cytometry (Fig 2, C). In contrast, quenching endogenous ROS with the ROS

scavenger N-acetylcysteine amide (NACA) in indicated cell lines substantially dampens p62 levels

and autophagy activities due to blockage of their endogenous Keap1-NRF2 pathway activities, as

well as attenuates endogenous DNA damage (Fig 2, D). Furthermore, confocal microscopy and

flow cytometry results show that treatment of IB4 cells with 3-AT or with the traditional oxidative

DNA damage inducer H_2O_2 remarkably increases p62 expression, autophagosomes and

autophagy flux, and that most p62 foci co-localize with autophagosomal bodies in the cytoplasm

(Fig 2, E and F).

   Taken together, these results indicate that endogenous ROS are responsible for p62

expression by activating the Keap1-NRF2 pathway, and for induction of p62-mediated selective

autophagy, in viral latency.

Autophagy inhibition sensitizes EBV[+] cells to ROS-induced DNA damage that is associated

with p62 accumulation

Since we have previously shown that mild ionomycin treatments induce p62 expression and

profound autophagy in lymphoblastic cell lines (LCLs), but stringent treatments promote p62

degradation due to induction of massive autophagy [34], we used ionomycin treatment here to

study the p62-autohagy interplay in regulating DDR in LCLs, which serve as a system crucial for

genetic and functional study of carcinogen sensitivity and DNA repair [40].

We first used the lysosome-specific inhibitor bafilomycin A1 (BafA1), which inhibits lysosomal

activity that occurs after LC3 processing, to inhibit autophagy activity induced by ionomycin in

LCLs. Results show that both p62 and LC3b, which are both selectively degraded by p62-mediated

autophagy [35, 36], are accumulated in ionomycin-treated cells due to impaired autophagy

activities. As a consequence, the levels of γH2AX are remarkably augmented (Fig 3, A and B). To

minimize the interference of potential "off-target" effects of BafA1, we performed this experiment

using another lysosome-specific inhibitor chloroquine, and obtained similar results (Fig 3, C).

These results indicate that the autophagy-p62 interplay plays a role in DDR in EBV latency.

We further show that ionomycin triggers profound ROS production and DNA damage in LCLs in

a time-dependent manner (Fig 3, D and E), and the ROS scavenger NACA offsets the effects of

ionomycin (Fig 3, E). Thus, these results indicate that ROS are responsible for ionomycin-induced

autophagy and DNA damage in EBV latency.

Autophagy inhibition promotes cell death

To confirm the requirement of ROS for induction of autophagy and DNA damage in EBV latency,

we further used H_2O_2 to treat IB4 cells. Results show that H_2O_2 treatment induces profound DNA

damage and reduces the endogenous p62 protein level in a dose-dependent manner, and

blockage of autophagy activity with BafA1 potentiates the DNA damage that correlates with

elevated p62 protein levels (Fig 4, A).

Furthermore, autophagy inhibition by BafA1 promotes cell death (as indicated by 7-AAD

expression and Annexin-V binding, or caspase-3 activity) induced by H_2O_2 (Fig 4, B and C) or

ionomycin (Fig 3, B), respectively.

Taken together, these results (Figs 3 and 4) indicate that autophagy inhibition exacerbates ROS-induced DNA damage by stabilizing p62 protein, further supporting that p62-mediated autophagy promotes DDR in EBV latency.

p62 accumulation upon autophagy inhibition destabilizes HR DNA repair proteins CHK1 and RAD51 in viral latency

It has been reported that autophagy inhibition or nuclear p62 accumulation accruing from autophagy deficiency promotes proteasomal degradation of HR DNA repair proteins such as RAD51, CHK1 and FLNA [15, 41]. Our results show that SavIII, JiJoye, MT4, and IB4, which have higher endogenous p62 and autophagy levels (Fig 1, B), have lower CHK1 and RAD51 protein levels, compared to SavI, P3HR1, CEM, and BJAB, respectively (Fig 5, A). These results suggest that the p62-autophagy interplay may also regulate proteasome-dependent stability of CHK1 and RAD51 proteins in viral latency. It has also been reported that p62 promotes NHEJ by activating the Keap1-NRF2 pathway in a feedback loop, which consequently induces expression of NHEJ-specific repair proteins such as 53BP1 [42]. However, our results show that 53BP1 reversely correlates with p62 at the protein level in viral latency (Fig 5, A), indicating that NHEJ-mediated DNA repair activity is also compromised in virus-transformed cells. In addition, endogenous DNA damage is consistently lower in viral latency with higher p62-mediated autophagy levels, in which both HR and NHEJ pathways are deficient (Fig 5, A), supporting our hypothesis that p62-mediated autophagy enables these cells to resist to DNA damage.

To validate whether proteasome also mediates degradation of CHK1 and RAD51 in virus-transformed cells, we used the proteasome inhibitor MG132 to treat these cells that express high levels of endogenous p62. Then, CHK1 and RAD51 were analyzed by immunoblotting. As

expected, our results show that MG132 treatment remarkably increases the protein levels of CHK1 and RAD51 (Fig 5, B), confirming that CHK1 and RAD51 protein stability is controlled in a proteasome-dependent manner in virus-transformed cells.

In addition to its ability to inhibit proteasomal activity, MG132 is capable of inducing autophagy in various cancer cells [43]. Our results show that MG132 also increases the LC3b cleavage product LC3b-II in virus-transformed cells, and dramatically reduces p62 protein levels (Fig 5, B). Together, these results indicate that excess induction of autophagy in viral latency resulting in elevation of CHK1 and RAD51 protein levels is associated with p62 degradation. However, the accumulation of CHK1 and RAD51 and decrease of p62 after MG132 treatment did not mitigate DNA damage; instead, DNA damage is strikingly increased (Fig 5, B). This observation can be explained by the fact that proteasome function is required for DNA damage repair [44].

We then inhibited endogenous autophagy activities in virus-transformed cells with BafA1. Results show that autophagy inhibition did not affect 53BP1, but clearly decreases CHK1 and RAD51 protein levels that are associated with elevated endogenous p62 protein levels (Fig 5, C).

Next, we sought to evaluate whether transient expression of p62 regulates DNA repair protein stability. To this end, we transfected the expression plasmids harboring HA-p62 or pcDNA4 vector control into BJAB, P3HR1, and CEM cells, which express low levels of endogenous p62. Then, we analyzed CHK1, RAD51, and 53BP1. Surprisingly, results show that exogenic expression of p62 did not affect the protein levels of CHK1, RAD51 and 53BP1 (Fig 5, D). By checking autophagy, results show that exogenic p62 expression did not induce LC3b cleavage either, implying that p62 alone is not sufficient for autophagy induction (Fig 5, D). Together with Fig 5, B and C, these results suggest that p62-mediated destabilization of DNA repair proteins is dependent on regulation of p62 by autophagy; autophagy inhibition may result in p62 nuclear translocation that is

required for its destabilization of DNA repair proteins [16].

To further confirm the conditional role of p62 in downregulation of the DNA repair proteins, we depleted p62 expression in IB4 cells using shRNA-mediated gene knockdown. Results show that p62 depletion reduces autophagy activity, as shown by decreased LC3b cleavage and autophagy flux (Fig 5, E and F), consistent with the notion that p62 is required for endogenous autophagy induction. However, the protein levels of CHK1, RAD51, and 53BP1 have no apparent changes in p62-depleted cells (Fig 5, E and G). Elevating endogenous ROS levels with 3-AT, which induces autophagy (Fig 2, B), in p62-depleted cells did not elevate the DNA repair protein levels, but consistently results in slight decreases of CHK1, in addition to inducing DNA damage (Fig 5, E). Additional inhibition of the residual autophagy activities in p62-depleted cells slightly decreases CHK1 and RAD51 levels (Fig 5, G). These results further indicate that p62 is required for destabilization of DNA repair proteins in response to autophagy inhibition.

Taken together, these results demonstrate that p62 accumulation in response to autophagy inhibition promotes proteasomal degradation of RAD51 and CHK1 in virus-transformed cells. The conditional role of p62 in DDR is in line with a recent report showing that the ability of p62 to inhibit DNA repair requires its localization in the nucleus [16]. Further study is required to determine whether and how autophagy inhibition causes p62 accumulation in the nucleus. These results also indicate that a fine balance of p62 and autophagy levels is required to confine endogenous p62 in the cytoplasm of virus-transformed cells.

p62 depletion by RNA interference promotes RNF168-mediated chromatin ubiquitination and DNA repair in viral latency

It has been shown that p62 inhibits both HR and NHEJ independently of the autophagy machinery [16], through its physical interaction with RNF168, which mediates histone ubiquitination that is

prelude to activation of both HR and NHEJ DSB repair mechanisms [17].

Consistently, our confocal microscopy results show that RNF168 co-localizes with endogenous γH2AX DNA damage foci in IB4 cells. More importantly, shRNA-mediated p62 depletion significantly increases RNF168-γH2AX foci (Fig 6, A and B). Further, using the anti-ubiquitinated proteins antibody FK2, we show that shRNA-mediated p62 depletion significantly increases histone H3 ubiquitination and FK2-γH2AX foci (Fig 6, C and D). We further validated the interaction of endogenous p62 with RNF168 by immunoprecipitation (IP) in virus-transformed cells treated with BafA1 (Fig 6, E), and the increased histone H3 ubiquitination due to p62 depletion in cells treated with ionomycin (Fig 6, F). Our IP assays failed to detect definite p62-RNF168 interaction and H3 ubiquitination in these cells without treatments (not shown). In conclusion, these results indicate that p62 depletion promotes chromatin ubiquitination and RNF168-mediated DNA repair mechanisms.

p62 and autophagy are upregulated and correlate with ROS and DNA damage levels in B cell population of AIDS lymphoma patients

Finally, we evaluated the biological relevance of our findings in a disease setting. To this end, we assessed the correlation of ROS production, p62 expression, autophagy activity, and DNA damage in B cell populations of AIDS lymphoma patients, and compared these parameters with age-matched health controls (HS). Our results show that, CD19$^+$ B cell frequency in AIDS patients on ART has no significant difference compared to age-matched HS (Fig 7, A). p62-positive CD19$^+$ B cell population and the MFI of p62 in CD19$^+$ B cell population are significantly greater in AIDS patients, compared with age-matched controls (Fig 7, B and C). Correspondingly, endogenous ROS and autophagy levels are higher (Fig 7, D and E), concomitant with significantly lower DNA damage (Fig 7, F), in B cell population of AIDS patients, compared to HS.

Both EBV and HIV, as well as other co-morbidities, have the potential to manipulate DDR and produce ROS in AIDS patients; unfortunately, we could not rule out the separate contribution of EBV to their deregulation in the setting. However, these preliminary *ex vivo* observations are of importance in that they provide biological relevance to our findings from mechanistic studies using cell line *in vitro* system.

Discussion

In this study, we provide several lines of evidence that support a crucial role for p62-mediated autophagy in regulation of DDR in oncogenic virus latent infection. First, p62 is upregulated by ROS through activation of the Keap1-NRF2 pathway, and considerable levels of p62-mediated selective autophagy are constitutively induced in this setting. Second, inhibition of autophagy in virus-transformed cells exacerbates ROS-induced DNA damage, and destabilizes the DNA repair proteins RAD51 and CHK1 in a p62-dependent manner; in contrast, excess autophagy induction promotes accumulation of the DNA repair proteins CHK1 and RAD51 that is associated with p62 degradation. Third, shRNA-mediated p62 depletion promotes RNF168-mediated chromatin ubiquitination and DNA repair in EBV latency. These findings have defined a crucial role for p62-mediated autophagy in regulation of DDR in viral latency (Fig 8).

The p62-autophagy interplay is well balanced and controlled in diverse contexts, with cancer and aging being two representative systems [4-6, 45]. Loss of this balance by exogenic stresses may result in different impacts on DDR. Pharmaceutical inhibition of autophagy, or spontaneous autophagy deficiency during aging, chronic inflammation, or neurodegeneration, leads to p62 accumulation, consequently attenuating DNA repair that accounts for the etiology of age-related disorders [46, 47]. In contrast, substantial enhancement of basal levels of autophagy in cancer cells by anticancer chemotherapeutic drugs or by radiation therapy promotes p62 degradation, and

consequently confers these cells resistance to DNA damage-induced cell death [21, 48, 49].

Moreover, our results indicate that the basal levels of p62-mediated autophagy are distinctly regulated in different EBV latency programs. ROS are produced separately by the EBV products LMP1, EBNA1/2, and EBERs, amongst which LMP1 induces predominant ROS [50-54]. In consistent, our results show that EBV type III latency produces a greater level of ROS compared with type I latency (Fig 1, A). Thus, cells with type III latency express higher endogenous levels of p62-mediated autophagy (Fig 1, B), which are required to overcome the higher risk of DNA damage in response to endogenous higher oxidative stress and replication stress to support their more aggressive proliferation, and higher growth and survival demands. As such, a higher level of p62-mediated autophagy also confers these cells greater resistance to DNA damage in response to drug treatments. The type III latency cell line P3HR1, which was derived from the parental JiJoye but lacks LMP1 expression, resembles type I latency cell lines in ROS production, expression of p62 and autophagy (Fig 1), and DNA repair proteins (Fig 5), further supporting that LMP1 contributes the majority to these events.

Surprisingly, our results show that, in contrast to p62 elimination by massive autophagy, p62 depletion by shRNA potentiates DNA damage (Fig 5, D). There are a few possibilities to explain this paradox. First, p62 depletion impairs p62-mediated selective autophagy that is resident in the cytoplasm (Fig 2, E) and required for maintaining the ability of the cell to resist to DNA damage (Fig 5, E and F); second, other p62-mediated, autophagy-independent, DNA damage-protecting functions are impaired after shRNA-mediated p62 depletion and consequently DNA damage is accelerated, given the fact that p62 is a multifunctional protein [3]. Investigation of these potential p62-mediated but autophagy-independent functions in viral latency and oncogenesis is underway. In fact, p62 depletion resulting in worsen DNA damage is coincident with its role as a tumor promoter, which is induced by Ras that accounts for at least 25% of human cancers [55]. p62

overexpression in hepatocellular carcinoma (HCC) predicts poor prognosis [56].

Based on the observations from us and other relevant studies, we propose that p62 plays a dichotomous role in DDR, depending on the presence or absence of autophagy. Higher levels of p62 resulting from defective autophagy inhibit DNA repair and therefore perturb genomic instability that facilitates tumor initiation [57]. In mouse models with defective autophagy, p62 ablation decreases tumorigenesis [55]. However, considerable levels of p62 in cancer cells promote DNA repair by mediating selective autophagy activation, and consequently confer these cancer cells resistance to DNA damage. p62 is upregulated at considerable levels in different cancer cells, including breast and prostate cancers, where it is required for induction of selective autophagy to support cancer cell metabolism and survival [56, 58, 59].

Regarding the mechanisms underneath deregulation of DDR by the p62-autophagy interplay, our results indicate that autophagy inhibition promotes proteasomal degradation of RAD51 and CHK1 in a p62-depdendent manner (Fig 5), and that p62 depletion promotes RNF168-mediated DDR (Fig 6). Our results show that the majority of endogenous p62 is tethered to autophagosomes in the cytoplasm (Fig 2, E), but inhibition of DNA repair requires p62 in the nucleus [16]. Thus, our results define a new role for p62-mediated autophagy in preventing DNA damage by confining p62 in the cytoplasm. In conclusion, p62-mediated selective autophagy not only confers invulnerability to DNA damage, but also at least partially contributes to the deficiency of traditional DDR mechanisms, in virus-transformed cells. In this regard, it is to our understanding that an oncogenic virus gains a dual benefit by invoking p62-mediated autophagy: one facet of p62-mediated autophagy endows its host cell with ability to resist DNA damage to support cell survival; the other facet hijacks the traditional DDR mechanisms in the host cell to facilitate genomic instability.

Although p62 was reported to promote NHEJ by inducing 53BP1 expression through the Keap1-NRF2 pathway that requests p62 S349 phosphorylation [42, 60], our results show that p62

is reversely correlated with 53BP1 in viral latency (Fig 5, A), and that p62 accumulation by autophagy inhibition, transient expression of p62, or p62 depletion by shRNA failed to regulate 53BP1 levels (Fig 5). Thus, p62 does not regulate 53BP1 in our system, and how 53BP1 is downregulated in virus-transformed cells is worthy of further examination (Fig 5, A). Rather, p62 interacts with RNF168 in response to autophagy inhibition, and consequently inhibits RNF168-mediated chromatin ubiquitination in viral latency (Fig 6). In this regard, our findings are consistent with a recent study [16], which has shown that p62 can impede both HR and NHEJ through its interaction with RNF168, given that RNF168-mediated histone ubiquitination is prerequisite for activation of all DSB repair mechanisms [17].

Viruses have evolved diverse strategies to hijack host traditional DDR machinery during their chronic infections to perturb genomic integrity, including their ability to deregulate the p62-autophagy balance, which we believe only makes partial contribution. In fact, our group has recently shown that traditional DDR mechanisms are also deficient in aging T cells in chronic HCV infection [61, 62], at least partially attributable to endogenous p62 accumulation accruing from deficient autophagy in these cells. p62 also inhibits DDR through other mechanisms that have not been fully elucidated. For example, nuclear p62 interacts with and inhibits PML nuclear bodies, which are involved in DNA repair [63, 64]. Moreover, other autophagy mechanisms, such as chaperone-mediated autophagy [65], also participate in DDR, by regulating stability of DDR-related proteins such as HP1α and CHK1 [66, 67], and by regulating p62-dependent or -independent cellular functions [68]. It is of great interest to investigate these potential mechanisms and their related cellular mechanisms in virus-mediated oncogenesis.

Endogenous ROS/RNS trigger signal cascades that activate both DDR and autophagy programs. Unrepaired damaged DNA can serve as a major source of genomic instability particularly in cancer cells where traditional DDR and cell death pathways are compromised.

Thus, cancer cells heavily rely on autophagy, not only to replenish their deficient DNA repair

mechanisms, but also to corroborate their higher metabolic demand than normal cells do.

Therefore, cancer cells are more vulnerable to autophagy inhibition, providing viable opportunities

for therapeutic strategy by targeting autophagy, in particular in combination with another cellular

mechanism that is specifically coupled with autophagy in a given cancer context for improving

clinical efficacy and specificity [3, 18, 69].

Materials and Methods

Cell Lines

SavI, SavIII, P3HR1 and JiJoye are human B cell lines derived from EBV-positive Burkitt's

lymphoma (BL) patients. P3HR1 was derived from JiJoye but does not express LMP1 due to

lacking the entire EBNA2 ORF in the viral genome [70]. BJAB is an EBV-negative BL line. The

lymphoblastic cell line (LCL) IB4 was derived from umbilical cord B-lymphocytes latently infected

with EBV *in vitro*. KR4 is a LCL with gamma irradiation resistance [71]. LCL45 is a newly

established LCL by *in vitro* transforming primary B cells of a healthy adult peripheral blood with the

EBV strain B95.8. CEM is a HTLV1-negative, EBV-negative T cell line derived from acute

leukemia, and MT4 is a HTLV1-transformed CD4$^+$ T cell line derived from umbilical cord blood

lymphocytes. B and T cell lines and primary PBMCs from patients are cultured with RPMI1640

medium plus 10% FBS and antibiotics. All cell culture supplies were purchased from Life

Technologies.

Ethics Statement

AIDS lymphoma blood samples were from our established East Tennessee State University James

H. Quillen VA Medical Center (ETSU/VA) clinical repository [61, 72, 73], which has been collected

from local AIDS subjects on ART with undetectable viremia for at least 1 year and a reasonable CD4 count (>200 cells/mm^3). All subjects were anonymized. Subjects co-infected with HBV, HCV or HTLV1 were excluded. Blood from healthy subjects (Physicians Plasma Alliance, Gray, TN) serve as controls. The mean age of the two populations was comparable (p>0.05). The study protocol was approved by the joint institutional review board (IRB) at ETSU/VA (IRB, #01-105s). Written informed consent was obtained from all participants.

Antibodies and Reagents

p62 (D-3), LIMD1 (H-4), and histone H3 (1G1) mouse monoclonal antibodies were from Santa Cruz for immunoprecipitation or immunoblotting. p62-Alexa Fluor 488 mouse antibody for flow cytometry was from Millipore or R&D Systems. Phospho-p62(S403), K63-linked ubiquitin rabbit monoclonal antibody, NRF2 (D1Z9C) mouse monoclonal antibody, and HRP-coupled secondary antibodies were from Cell Signaling Technologies. RNF168 rabbit polyclonal antibody and the FK2 mouse monoclonal antibody that recognizes K29-, K48-, and K63-linked polyubiquitin chains and monoubiquitin conjugation but not free ubiquitin for immunofluorescence were from Millipore. The RNF168 rabbit polyclonal antibody for IP was from Proteintech Group Inc. RNF168 sheep polyclonal, and LC3b and histone H3 rabbit polyclonal antibodies were from Invitrogen. RAD51 rabbit polyclonal and CHK1 (G-4) mouse monoclonal antibodies were from Abcam and Santa Cruz, respectively. The γH2AX(S139) mouse and rabbit antibodies were from BioLegend and Cell Signaling Technol., respectively. Secondary antibodies coupled with FITC, Alexa Fluor, APC, PE, Cy5, or PerCP and human anti-CD19-PE were from BioLegend, BD Biosciences, Invitrogen, or eBioscience.

HA-p62 cloned in pcDNA4 was a gift from Qing Zhong (Addgene #28027) [74]. CellRox Green, MG132, chloroquine, and doxorubicin HCl, were purchased from Invitrogen, EMD Millipore, MP

Biomedicals, and UBPBio, respectively. Ionomycin calcium salt, N-acetylcysteine amide,

bafilomycin A1, doxycycline, and mouse and rabbit IgG were purchased from Sigma. H_2O_2 was

from Santa Cruz.

shRNA Transfection and Selection of Stable Transfectants

A set of p62 shRNA cloned in pTRIPz/Puro comprising three individual p62 shRNA and a

scramble control plasmids were purchased from Dharmacon. We selected two of them for this

study and the targeting sequences on the human p62/SQSTM1 gene are: shRNA#1: 5'-

TCTCTTTAATGTAGATTCG-3' and shRNA#2: 5'- TCAGGAAATTCACACTCGG-3'. Lentiviral

packing, preparation, infection, and selection of stable cells by puromycin (0.5 µg/ml) were

performed as detailed in our previous publication [34]. shRNA expression was induced by 1

µg/ml doxycycline for 3 days, and then cells (expressing red fluorescence) were subjected to a

second selection by FACS on a FACS/Aria Fusion Cell Sorter (BD Biosciences). Stable

transfectants were maintained in complete medium plus 1.0 µg/ml puromycin.

Confocal Microscopy

Cells were fixed in 2% paraformaldehyde (PFA) for 20 min, permeabilized with 0.3% Triton X-100

in phosphate-buffered saline (PBS) for 10 min, blocked with 5% bovine serum albumin (BSA) in

PBS for 1 h, and then incubated with indicated primary antibodies at 4 °C overnight. Cells were

washed with PBS with 0.1% Tween-20 for three times, and then incubated with corresponding

secondary antibodies coupled with FITC, Alexa Fluor, APC, PE, Cy5, or PerCP, at room

temperature for 1 h. Cells were then washed and mounted with DAPI Fluoromount-G

(SouthernBiotech, Birmingham, AL). Images were acquired with a confocal laser-scanning inverted

microscope (Leica Confocal, Model TCS sp8, Germany).

Immunoprecipitation and Immunoblotting

To assess endogenous p62-RNF168 interaction, 1×10^7 cells for each sample were lysed with NP40 lysis buffer (150 mM NaCl, 1% NP-40, 50 mM Tris-pH 8.0, plus protease inhibitors), and cell lysates were subjected to immunoprecipitation (IP) with 1.5 µg anti-RNF168 for overnight, and then incubated with 40 µl Protein A/G beads (Santa Cruz) for 1 h. After three washes, proteins on beads were denatured in 1% SDS before subjected to immunoblotting (IB). IB was carried out with indicated antibodies and signals were detected with an enhanced chemiluminescence (ECL) kit following the manufacturer's protocol (Amersham Pharmacia Biotech). For H3 ubiquitination assay, endogenous H3 was pulled down with an H3 antibody in denatured IP, as detailed in our publication [75].

RNA Extraction and Real-time Quantitative PCR

Total RNA was isolated from tested cells using an RNeasy Mini kit according to the manufacturer's protocols (Qiagen). The eluted RNA was subjected to reverse transcriptase reactions, which were performed with the use of GoScript RT kit following the manufacturer's instructions (Promega).

Quantitative real-time PCR (qPCR) was performed with the use of SYBR Green (Applied Biosystems), on a CFX96^{TM} Real-time PCR Detection System (Bio-Rad). All reactions were run in triplicates. Mean cycle threshold (C_t) values were normalized to 18s rRNA, yielding a normalized C_t (ΔC_t). $\Delta\Delta C_t$ value was calculated by subtracting respective control from the ΔC_t, and expression level was then calculated by 2 raised to the power of respective -$\Delta\Delta C_t$ value. The averages of 2^(-$\Delta\Delta C_t$) in the control samples were set to 1 or 100%. Results are the average $\pm$ standard error (SE) of triplicates for each sample. Primers for real-time qPCR are as follows: p62: F: 5'-CAGGCGCACTACCGCGATG-3' and R: 5'-ACACAAGTCGTAGTCTGGGCAGAC-3'. Keap1: F: 5'-CCATGGGCGAGAAGTGTGTCC-3'; R: 5'-ACAGGTTGAAGAACTCCTCTTGCTTG-3'. 18s rRNA:

F: 5'-GGCCCTGTAATTGGAATGAGTC-3' and R: 5'-CCAAGATCCAACTACGAGCTT-3'.

Flow Cytometry

Samples were fixed with 2% PFA for 20 min at RT, then wash with flow buffer (eBioscience). Samples were then incubated with PE-conjugated anti-human CD19 antibody (eBioscience) or isotype controls for 20 min at RT, then wash with flow buffer, followed by incubation with p62-Alexa Fluor 488 antibody for 60 min at RT. Samples were then washed with flow buffer, and analyzed with BD C6 plus flow cytometer.

For intracellular ROS measurement, $1X10^6$ cells in 500 µl medium per well were seeded in 24-well plates, and cultured overnight. 1 µl CellROX Green Reagent (Invitrogen) was added to each well and incubated for 30 min. Cells were then washed 3 times with PBS, and fixed with 2% PFA for 20min at RT, followed by extensive washes and then incubated with PE-conjugated anti-human CD19 antibody (eBioscience) for 20min at RT, before subjected to flow cytometry.

Apoptosis and Proliferation Assays

Apoptosis was quantified using flow cytometry as detailed in our previous publication [34], for Annex V binding (BD Biosciences) and 7-Aminoactinomycin D (7-AAD) expression (eBioscience). Caspase 3 activity was evaluated by Western blotting. Cell proliferation was quantified using a cell proliferation kit (Promega).

Statistical Analysis

Unpaired, two-tailed student t tests were executed using GraphPad Prism (version 6) to determine the differences between two data sets obtained from three independent experiments. $p<0.05$ (*) and $p<0.01$ (**), and $p<0.001$ (***) were considered significant. Data are expressed

as mean ± standard error (SE) of duplicate or triplicate samples, and representative results from

at least three independent repeats with similar results are shown.

Acknowledgements

This study was conducted with resources and the use of facilities at the James H. Quillen Veterans

Affairs Medical Center. The contents in this publication are solely the responsibility of the authors

and does not necessarily represent the official views of the NIH, the Department of Veterans

Affairs, or the United States Government. The authors declare no competing interests.

References

1. Moscat J, Karin M, Diaz-Meco MT. p62 in Cancer: Signaling Adaptor Beyond Autophagy. Cell. 2016;167(3):606-9. doi: http://dx.doi.org/10.1016/j.cell.2016.09.030.
2. Bitto A, Lerner CA, Nacarelli T, Crowe E, Torres C, Sell C. p62/SQSTM1 at the interface of aging, autophagy, and disease. AGE. 2014;36(3):9626. doi: 10.1007/s11357-014-9626-3.
3. Ning S, Wang L. The Multifunctional Protein p62 and Its Mechanistic Roles in Cancers. Current Cancer Drug Targets. 2019;19:1-11. doi: http://dx.doi.org/10.2174/1568009618666181016164920.
4. Wilde L, Tanson K, Curry J, Martinez-Outschoorn U. Autophagy in cancer: a complex relationship. Biochemical Journal. 2018;475(11):1939-54. doi: 10.1042/bcj20170847.
5. Singh SS, Vats S, Chia AY-Q, Tan TZ, Deng S, Ong MS, et al. Dual role of autophagy in hallmarks of cancer. Oncogene. 2018;37(9):1142-58. doi: 10.1038/s41388-017-0046-6.
6. Rybstein MD, Bravo-San Pedro JM, Kroemer G, Galluzzi L. The autophagic network and cancer. Nature Cell Biology. 2018;20(3):243-51. doi: 10.1038/s41556-018-0042-2.
7. Farre J-C, Subramani S. Mechanistic insights into selective autophagy pathways: lessons from yeast. Nat Rev Mol Cell Biol. 2016;17(9):537-52. doi: 10.1038/nrm.2016.74.
8. Zaffagnini G, Martens S. Mechanisms of Selective Autophagy. Journal of Molecular Biology. 2016;428(9Part A):1714-24. PubMed PMID: PMC4871809.
9. Sparrer KMJ, Gack MU. TRIM proteins: New players in virus-induced autophagy. PLOS Pathogens. 2018;14(2):e1006787. doi: 10.1371/journal.ppat.1006787.
10. Di Meo S, Reed TT, Venditti P, Victor VM. Role of ROS and RNS Sources in Physiological and Pathological Conditions. Oxidative Medicine and Cellular Longevity. 2016;2016:44.
11. Sallmyr A, Fan J, Rassool FV. Genomic instability in myeloid malignancies: Increased reactive oxygen species (ROS), DNA double strand breaks (DSBs) and error-prone repair. Cancer Letters. 2008;270(1):1-9.
12. Vilenchik MM, Knudson AG. Endogenous DNA double-strand breaks: Production, fidelity of repair, and induction of cancer. Proceedings of the National Academy of Sciences. 2003;100(22):12871-6.

13. Chang HHY, Pannunzio NR, Adachi N, Lieber MR. Non-homologous DNA end joining and alternative pathways to double-strand break repair. Nat Rev Mol Cell Biol. 2017;18(8):495-506.

14. Tubbs A, Nussenzweig A. Endogenous DNA Damage as a Source of Genomic Instability in Cancer. Cell. 2017;168(4):644-56. doi: 10.1016/j.cell.2017.01.002.

15. Hewitt G, Carroll B, Sarallah R, Correia-Melo C, Ogrodnik M, Nelson G, et al. SQSTM1/p62 mediates crosstalk between autophagy and the UPS in DNA repair. Autophagy. 2016;12(10):1917-30. doi: 10.1080/15548627.2016.1210368.

16. Wang Y, Zhang N, Zhang L, Li R, Fu W, Ma K, et al. Autophagy Regulates Chromatin Ubiquitination in DNA Damage Response through Elimination of SQSTM1/p62. Molecular Cell. 2016;63(1):34-48. doi: http://doi.org/10.1016/j.molcel.2016.05.027.

17. Schwertman P, Bekker-Jensen S, Mailand N. Regulation of DNA double-strand break repair by ubiquitin and ubiquitin-like modifiers. Nat Rev Mol Cell Biol. 2016;17(6):379-94. doi: 10.1038/nrm.2016.58.

18. Santana-Codina N, Mancias JD, Kimmelman AC. The Role of Autophagy in Cancer. Annual Review of Cancer Biology. 2017;1(1):19-39. doi: 10.1146/annurev-cancerbio-041816-122338.

19. O'Connor Mark J. Targeting the DNA Damage Response in Cancer. Molecular Cell. 2015;60(4):547-60. doi: https://doi.org/10.1016/j.molcel.2015.10.040.

20. Mathew R, White E. Autophagy, Stress, and Cancer Metabolism: What Doesn't Kill You Makes You Stronger. Cold Spring Harbor Symposia on Quantitative Biology. 2011;76:389-96. doi: 10.1101/sqb.2012.76.011015.

21. Sui X, Chen R, Wang Z, Huang Z, Kong N, Zhang M, et al. Autophagy and chemotherapy resistance: a promising therapeutic target for cancer treatment. Cell Death & Disease. 2013;4(10):e838. doi: 10.1038/cddis.2013.350. PubMed PMID: PMC3824660.

22. Lee DY, Sugden B. The latent membrane protein 1 oncogene modifies B-cell physiology by regulating autophagy. Oncogene. 2008;27(20):2833-42.

23. Ditzel M, Broemer M, Tenev T, Bolduc C, Lee TV, Rigbolt KTG, et al. Inactivation of effector caspases through nondegradative polyubiquitylation. Mol Cell. 2008;32(4):540-53.

24. Cavignac Y, Esclatine A. Herpesviruses and Autophagy: Catch Me If You Can! Viruses. 2010;2(1):314. PubMed PMID: doi:10.3390/v2010314.

25. Pratt ZL, Sugden B. How human tumor viruses make use of autophagy. Cells. 2012;1(3):617-30.

26. Hau P, Tsao S. Epstein–Barr Virus Hijacks DNA Damage Response Transducers to Orchestrate Its Life Cycle. Viruses. 2017;9(11):341. PubMed PMID: doi:10.3390/v9110341.

27. Whitehurst CB, Vaziri C, Shackelford J, Pagano JS. Epstein-Barr Virus BPLF1 Deubiquitinates PCNA and Attenuates Polymerase η Recruitment to DNA Damage Sites. Journal of Virology. 2012;86(15):8097-106. doi: 10.1128/jvi.00588-12.

28. Hafez A, Luftig M. Characterization of the EBV-Induced Persistent DNA Damage Response. Viruses. 2017;9(12):366. PubMed PMID: doi:10.3390/v9120366.

29. De Leo A, Colavita F, Ciccosanti F, Fimia GM, Lieberman PM, Mattia E. Inhibition of autophagy in EBV-positive Burkitt's lymphoma cells enhances EBV lytic genes expression and replication. Cell Death &Amp; Disease. 2015;6:e1876. doi: https://www.nature.com/articles/cddis2015156#supplementary-information.

30. Masucci MG. Epstein-Barr virus oncogenesis and the ubiquitin-proteasome system. Oncogene. 2004;23(11):2107-15.

31. Dantuma NP, Masucci MG. The ubiquitin/proteasome system in Epstein-Barr virus latency and associated malignancies. Sem Cancer Biol. 2003;13(1):69-76.

32. Shackelford J, Pagano J. Role of the ubiquitin system and tumor viruses in AIDS-related cancer. BMC Biochem. 2007;8(Suppl 1):S8.

33. Wang L, Wang Y, Zhao J, Ren J, Hall KH, Moorman JP, et al. LUBAC modulates LMP1 activation of NFκB and IRF7. Journal of Virology. 2017;91(4):e1138-16.

34. Wang L, Riggs K, Kohne C, Yohanon JU, Foxler DE, Sharp TV, et al. LIMD1 Is Induced by and Required for LMP1 Signaling, and Protects EBV-Transformed Cells from DNA Damage-Induced Cell Death. Oncotarget. 2018;9(5):6282-97. doi: doi.org/10.18632/oncotarget.23676. .

35. Lussignol M, Esclatine A. Herpesvirus and Autophagy: "All Right, Everybody Be Cool, This Is a Robbery!". Viruses. 2017;9(12):372. PubMed PMID: doi:10.3390/v9120372.

36. Barth S, Glick D, Macleod KF. Autophagy: assays and artifacts. The Journal of pathology. 2010;221(2):117-24. doi: 10.1002/path.2694. PubMed PMID: PMC2989884.

37. Matsumoto G, Wada K, Okuno M, Kurosawa M, Nukina N. Serine 403 phosphorylation of p62/SQSTM1 regulates selective autophagic clearance of ubiquitinated proteins. Mol Cell. 2011;44. doi: 10.1016/j.molcel.2011.07.039.

38. Taguchi K, Fujikawa N, Komatsu M, Ishii T, Unno M, Akaike T, et al. Keap1 degradation by autophagy for the maintenance of redox homeostasis. Proceedings of the National Academy of Sciences. 2012;109(34):13561-6. doi: 10.1073/pnas.1121572109.

39. Kurz EU, Douglas P, Lees-Miller SP. Doxorubicin Activates ATM-dependent Phosphorylation of Multiple Downstream Targets in Part through the Generation of Reactive Oxygen Species. Journal of Biological Chemistry. 2004;279(51):53272-81. doi: 10.1074/jbc.M406879200.

40. Hussain T, Mulherkar R. Lymphoblastoid Cell lines: a Continuous in Vitro Source of Cells to Study Carcinogen Sensitivity and DNA Repair. International Journal of Molecular and Cellular Medicine. 2012;1(2):75-87. PubMed PMID: PMC3920499.

41. Liu EY, Xu N, O'Prey J, Lao LY, Joshi S, Long JS, et al. Loss of autophagy causes a synthetic lethal deficiency in DNA repair. Proceedings of the National Academy of Sciences. 2015;112(3):773-8.

42. Kim SB, Pandita RK, Eskiocak U, Ly P, Kaisani A, Kumar R, et al. Targeting of Nrf2 induces DNA damage signaling and protects colonic epithelial cells from ionizing radiation. Proceedings of the National Academy of Sciences of the United States of America. 2012;109(43):E2949-E55. Epub 10/08. doi: 10.1073/pnas.1207718109. PubMed PMID: 23045680.

43. Ding W-X, Ni H-M, Gao W, Yoshimori T, Stolz DB, Ron D, et al. Linking of Autophagy to Ubiquitin-Proteasome System Is Important for the Regulation of Endoplasmic Reticulum Stress and Cell Viability. The American Journal of Pathology. 2007;171(2):513-24. doi: 10.2353/ajpath.2007.070188. PubMed PMID: PMC1934546.

44. Jacquemont C, Taniguchi T. Proteasome Function Is Required for DNA Damage Response and Fanconi Anemia Pathway Activation. Cancer Research. 2007;67(15):7395-405. doi: 10.1158/0008-5472.can-07-1015.

45. Martinez-Lopez N, Athonvarangkul D, Singh R. Autophagy and aging. Advances in experimental medicine and biology. 2015;847:73-87. doi: 10.1007/978-1-4939-2404-2_3. PubMed PMID: 25916586.

46. Matsuzawa-Ishimoto Y, Hwang S, Cadwell K. Autophagy and Inflammation. Annual Review of Immunology. 2018;36(1):73-101. doi: 10.1146/annurev-immunol-042617-053253. PubMed PMID: 29144836.

47. Hansen M, Rubinsztein DC, Walker DW. Autophagy as a promoter of longevity: insights from model organisms. Nature Reviews Molecular Cell Biology. 2018;19(9):579-93. doi: 10.1038/s41580-018-0033-y.

48. Marinković M, Šprung M, Buljubašić M, Novak I. Autophagy Modulation in Cancer: Current Knowledge on Action and Therapy. Oxidative Medicine and Cellular Longevity. 2018;2018:18. doi: 10.1155/2018/8023821.

49. Bouwman P, Jonkers J. The effects of deregulated DNA damage signalling on cancer chemotherapy response and resistance. Nat Rev Cancer. 2012;12(9):587-98.

50. Gruhne B, Sompallae R, Masucci MG. Three Epstein-Barr virus latency proteins independently promote genomic instability by inducing DNA damage, inhibiting DNA repair and inactivating cell cycle checkpoints. Oncogene. 2009;28(45):3997-4008.

51. Cerimele F, Battle T, Lynch R, Frank DA, Murad E, Cohen C, et al. Reactive oxygen signaling and MAPK activation distinguish Epstein–Barr Virus (EBV)-positive versus EBV-negative Burkitt's lymphoma. Proceedings of the National Academy of Sciences of the United States of America. 2005;102(1):175-9.

52. Gruhne B, Sompallae R, Marescotti D, Kamranvar SA, Gastaldello S, Masucci MG. The Epstein-Barr virus nuclear antigen-1 promotes genomic instability via induction of reactive oxygen species. Proceedings of the National Academy of Sciences. 2009;106(7):2313-8.

53. Raab-Traub N. Epstein-Barr virus in the pathogenesis of NPC. Seminars in Cancer Biology. 2002;12(6):431-41.

54. Kim SM, Hur DY, Hong SW, Kim JH. EBV-encoded EBNA1 regulates cell viability by modulating miR34a-NOX2-ROS signaling in gastric cancer cells. Biochem Biophys Res Commun. 2017;494(3-4):550-5. doi: 10.1016/j.bbrc.2017.10.095. PubMed PMID: 29061308.

55. Duran A, Linares JF, Galvez AS, Wikenheiser K, Flores JM, Diaz-Meco MT, et al. The Signaling Adaptor p62 Is an Important NF-κB Mediator in Tumorigenesis. Cancer Cell. 2008;13(4):343-54.

56. Umemura A, He F, Taniguchi K, Nakagawa H, Yamachika S, Font-Burgada J, et al. p62, Upregulated during Preneoplasia, Induces Hepatocellular Carcinogenesis by Maintaining Survival of Stressed HCC-Initiating Cells. Cancer Cell. 2016;29(6):935-48. doi: http://dx.doi.org/10.1016/j.ccell.2016.04.006.

57. Yang J, Peng H, Xu Y, Xie X, Hu R. SQSTM1/p62 (sequestosome 1) senses cellular ubiquitin stress through E2-mediated ubiquitination. Autophagy. 2018;14(6):1072-3. doi: 10.1080/15548627.2017.1332566. PubMed PMID: 28614034; PubMed Central PMCID: PMCPMC6103393.

58. Thompson HGR, Harris JW, Wold BJ, Lin F, Brody JP. p62 overexpression in breast tumors and regulation by prostate-derived Ets factor in breast cancer cells. Oncogene. 2003;22(15):2322-33.

59. Roodman GD, Hiruma Y, Kurihara N. p62: A Potential Target for Blocking Microenvironmental Support of Myeloma. Clinical Lymphoma and Myeloma. 2009;9:S25-S6. doi: 10.3816/CLM.2009.s.004.

60. Ichimura Y, Waguri S, Sou YS, Kageyama S, Hasegawa J, Ishimura R, et al. Phosphorylation of p62 activates the Keap1-Nrf2 pathway during selective autophagy. Mol Cell. 2013;51. doi: 10.1016/j.molcel.2013.08.003.

61. Zhao J, Dang X, Lam N, Cao D, Wu X, Zheng M, et al. Insufficiency of DNA repair enzyme ATM promotes naïve CD4 T cell loss in chronic hepatitis C virus infection. Cell Discovery. 2018;4(16):doi:10.1038/s41421-018-0015-4.

62. Nguyen LN, Zhao J, Cao D, Dang X, Wang L, Lian J, et al. Inhibition of TRF2 accelerates telomere attrition and DNA damage in naïve CD4 T cells during HCV infection. Cell Death and Diseases. 2018;9:900.

63. Pankiv S, Lamark T, Bruun J-A, Øvervatn A, Bjørkøy G, Johansen T. Nucleocytoplasmic Shuttling of p62/SQSTM1 and Its Role in Recruitment of Nuclear Polyubiquitinated Proteins to Promyelocytic Leukemia Bodies. Journal of Biological Chemistry. 2010;285(8):5941-53. doi: 10.1074/jbc.M109.039925.

64. Dellaire G, Bazett-Jones DP. PML nuclear bodies: dynamic sensors of DNA damage and cellular stress. BioEssays. 2004;26(9):963-77. doi: 10.1002/bies.20089.

65. Kaushik S, Cuervo AM. The coming of age of chaperone-mediated autophagy. Nature Reviews Molecular Cell Biology. 2018;19(6):365-81.

66. Chen S, Wang C, Sun L, Wang D-L, Chen L, Huang Z, et al. RAD6 Promotes Homologous Recombination Repair by Activating the Autophagy-Mediated Degradation of Heterochromatin Protein HP1. Molecular and Cellular Biology. 2015;35(2):406-16. doi: 10.1128/mcb.01044-14.

67. Park C, Suh Y, Cuervo AM. Regulated degradation of Chk1 by chaperone-mediated autophagy in response to DNA damage. Nature Communications. 2015;6:6823.

68. Eliopoulos AG, Havaki S, Gorgoulis VG. DNA Damage Response and Autophagy: A Meaningful Partnership. Frontiers in Genetics. 2016;7:204. PubMed PMID: PMC5116470.

69. Islam MA, Sooro MA, Zhang P. Autophagic Regulation of p62 is Critical for Cancer Therapy. International Journal of Molecular Sciences. 2018;19(5):1405. doi: 10.3390/ijms19051405. PubMed PMID: PMC5983640.

70. Hinuma Y, Konn M, Yamaguchi J, Wudarski DJ, Blakeslee JR, Grace JT. Immunofluorescence and Herpes-Type Virus Particles in the P3HR-1 Burkitt Lymphoma Cell Line. Journal of virology. 1967;1(5):1045-51.

71. Kozbor D, Lagarde AE, Roder JC. Human hybridomas constructed with antigen-specific Epstein-Barr virus-transformed cell lines. Proc Natl Acad Sci U S A. 1982;79(21):6651-5.

72. Zhou Y, Li GY, Ren JP, Wang L, Zhao J, Ning SB, et al. Protection of CD4+ T cells from hepatitis C virus infection-associated senescence via ΔNp63–miR-181a–Sirt1 pathway. Journal of Leukocyte Biology. 2016;100(5):1201-11. doi: 10.1189/jlb.5A0316-119RR.

73. Wang L, Zhao J, Ren JP, Wu XY, Morrison ZD, Elgazzar MA, et al. Expansion of myeloid-derived suppressor cells promotes differentiation of regulatory T cells in HIV-1+ individuals. AIDS. 2016;30(10):1521-31. doi: 10.1097/QAD.0000000000001083. PubMed PMID: 26959508.

74. Fan W, Tang Z, Chen D, Moughon D, Ding X, Chen S, et al. Keap1 facilitates p62-mediated ubiquitin aggregate clearance via autophagy. Autophagy. 2010;6(5):614-21. doi: 10.4161/auto.6.5.12189. PubMed PMID: 20495340; PubMed Central PMCID: PMCPMC4423623.

75. Ning S, Campos AD, Darnay B, Bentz G, Pagano JS. TRAF6 and the three C-terminal lysine sites on IRF7 are required for its ubiquitination-mediated activation by the Tumor Necrosis Factor Receptor family member Latent Membrane Protein 1. MolCellBiol. 2008;28(20):6536-46.

Figure Legend

Fig 1. p62-mediated selective autophagy is endogenously induced in viral latency and correlates with ROS-Keap1-NRF2 pathway activity

A. Endogenous ROS production in viral latency was measured by flow cytometry with the CellRox Green reagent (Invitrogen). Three independent repeats were conducted, and representative results are shown. Results are the mean ± standard error (SE) of duplicates for each sample. MFI=mean fluorescence intensity. **B**. The correlation of endogenous p62 protein levels and autophagy activity with the Keap1-NRF2 pathway activity in viral latency was evaluated by immunoblotting with indicated antibodies. **C**. The correlation of p62 expression with viral latency was evaluated at the transcription level by qPCR. The mRNA levels in SavI, P3HR1, and CEM were set to 1, and compared with the paired SavIII, JiJoye, and MT4 cell lines, respectively.

Fig 2. The ROS-Keap1-NRF2 pathway contributes to p62 expression and activation of p62-mediated autophagy in viral latency

A. SavI and SavIII cells, which were derived from the same patient, were treated with 2 µM of the topoisomerase II inhibitor doxorubicin HCl (Doxo) (UBPBio) for different time periods. **B-C**. SavI, P3HR1, and CEM were treated with 20 mM of the H_2O_2 catalase 3-amino-1,2,4-triazole (3-AT) (Fisher Scientific) or vehicle control for 48 h. Cells were then subjected to IB, qPCR, and flow cytometry analyses. **D**. IB4, LCL45, and MT4 were treated with 3 mM of the antioxidant N-acetylcysteine amide (NACA) (Sigma) or vehicle control for 30 h. The treated cells were then subjected to analyses for p62, autophagy, Keap1-NRF2 pathway, and ROS production. **E**. IB4 cells were treated with 20 mM 3-AT (Fisher Scientific) or vehicle control for

48 h, and analyzed for p62-autophagosome colocalization by confocal microscopy. Living

cells were first stained with a Cyto-ID Autophagy Detection kit (Enzo), and then fixed for

staining with the mouse p62 antibody (D-3) and Alexa 555 coupled anti-mouse antibody

(Invitrogen). Bar=3 μm. **F.** IB4 cells were treated with 50 μM H_2O_2 for 30 min and then

medium was replaced and continued in culture for 2 days, or treated with 20 mM 3-AT for 48

h, before subjected to analysis of autophagy flux by flow cytometry with the Cyto-ID

Autophagy Detection kit. MFI=mean fluorescence intensity.

Fig 3. Autophagy inhibition sensitizes EBV$^+$ cells to ROS-induced DNA damage that is associated with p62 accumulation

A. Cell lines with higher endogenous autophagy levels were treated with 0.4 μM of the

vacuolar ATPase inhibitor bafilomycin A1 (BafA1) (Sigma) or vehicle control for 24 h, and

then DNA damage (γH2AX) was evaluated by immunoblotting. **B-C**. The LCL lines IB4 and

LCL45 were treated with ionomycin (Iono) (Sigma) with indicated concentrations for 48 h plus

0.4 μM BafA1 or vehicle control for 24 h or plus 50 μM of the lysosome inhibitor chloroquine

(Chloro) (MP Biomedicals) or vehicle control for 6 h. p62, autophagy, and γH2AX were

analyzed by immunoblotting. **D**. IB4 cells were treated with 5 μg/ml Iono or vehicle control for

different time periods, and ROS production was measured by flow cytometry with the CellRox

Green reagent (Invitrogen). A representative result from three independent repeats is shown.

E. IB4 and LCL45 cells were treated with 5 μg/ml Iono plus 3 mM NACA or vehicle control for

different time periods (upper panel) or for 30 h (lower panel). p62, autophagy, and γH2AX

were analyzed by immunoblotting.

Fig 4. Autophagy inhibition promotes cell death

IB4 cells were treated with indicated concentrations of H_2O_2 in medium for 30 min. Then the medium was removed, and 0.4 µM BafA1 (Sigma) or vehicle control was added in freshly replaced medium for 48 h. **A**. p62, autophagy, and DNA damage were analyzed by immunoblotting. **B-C.** Cell death was analyzed by flow cytometry for Annexin V and 7-AAD expression. Results from a representative experiment of five independent repeats are shown (**B**), and statistical analysis results are expressed as mean ± standard error (SE) (**C**).

Fig 5. p62 accumulation upon autophagy inhibition destabilizes HR DNA repair proteins CHK1 and RAD51 in viral latency

A. Correlation of p62 with CHK1, RAD51 and 53BP1 protein levels in viral latency were analyzed in paired cell lines. **B**. Cell lines with higher levels of p62 protein were treated with 10 µM of the proteasome inhibitor MG132 for 6 h. Indicated proteins were probed by immunoblotting. **C**. Cell lines with higher levels of p62 protein were treated with 0.4 µM BafA1 (Sigma) for 48 h. Indicated proteins were probed by immunoblotting. **D**. Cell lines with lower levels of p62 protein were transfected with HA-p62 plasmids or vector control. Indicated proteins were analyzed by immunoblotting 48 h post-transfection. **E-G**. IB4 cells stably harboring p62 shRNA or shRNA control in 1 µg/ml puromycin were treated with 20 mM 3-AT or 0.4 µM BafA1 for 48 h before subjected to immunoblotting or flow cytometry. shRNA expression was induced by 1 µg/ml doxycycline for 2 days before the drug treatments. MFI=mean fluorescence intensity.

Fig 6. p62 depletion by RNA interference promotes RNF168-mediated chromatin ubiquitination and DNA repair in viral latency

IB4 cells stably expressing control or p62 shRNA were maintained with 1 µg/ml puromycin, and shRNA expression was induced by 1 µg/ml doxycycline for 3 days before subjected to confocal microscopy analysis for: **A**. RNF168, γH2AX and their colocalization in the nucleus with a rabbit RNF168 antibody (Millipore) and an Alexa 488 coupled anti-rabbit antibody (Invitrogen), and a mouse γH2AX(S139) antibody (BioLegend) and an Alexa 555 coupled anti-mouse antibody (BioLegend); **C**. chromatin ubiquitination, γH2AX and their colocalization in the nucleus with the mouse ubiquitin antibody clone FK2 (Millipore) and the Alexa 555 coupled anti-mouse antibody, and a rabbit γH2AX(S139) antibody (Cell Signaling Technol.) and the Alexa 488 coupled anti-rabbit antibody. Bar=6 µm. **B** and **D**. Quantification of RNF168/γH2AX foci or FK2/γH2AX foci in **A** and **C**, respectively. **E**. Interaction between RNF168 and p62 was assessed by immunoprecipitation. The indicated cell lines were treated with 0.4 µM BafA1 for 48h, and then cell lysates (0.5 mg total proteins for each) were subjected to immunoprecipitation with a rabbit RNF168 antibody (Proteintech Group Inc.), and immunoprecipitated proteins were analyzed by immunoblotting with the p62 antibody and the sheep RNF168 antibody (Invitrogen). **F**. Histone H3 ubiquitination was assessed in p62-depleted IB4 cells treated with 2.5 µg /ml ionomycin for 48 h. Cell lysates (0.5 mg total proteins for each) from IB4 cells stably expressing control or p62 shRNA were subjected to denatured immunoprecipitation with a rabbit H3 antibody (Invitrogen), and immunoprecipitated H3 was probed with the FK2 antibody and the mouse H3 antibody (1G1) (Santa Cruz).

Fig 7. p62 and autophagy are upregulated and correlate with ROS and DNA damage levels in B cell population of AIDS lymphoma patients

Endogenous p62, autophagy, ROS, and DNA damage levels in B cell population of AIDS lymphomas vs age-matched controls were evaluated by flow cytometry with corresponding antibodies and reagents. **A**. CD19-positive B cell population was analyzed with anti-human CD19-PE (eBioscience). **B-C**. p62 expression in B cell population was analyzed with the mouse p62 antibody (D-3) and Alexa 555 coupled anti-mouse antibody (Invitrogen). **D**. ROS levels in B cell population were analyzed with the CellRox Green reagent. **E**. Autophagy flux in B cell population was analyzed with the Cyto-ID Autophagy Detection kit (Enzo). **F**. DNA damage in B cell population was analyzed with γH2AX antibody. Statistical analyses were performed as detailed in Materials and Methods. MFI=mean fluorescence intensity.

Fig 8. A diagram showing the interaction of p62-mediated autophagy with DDR in EBV latency

EBV latent infection produces ROS, which further induce p62 expression through the Keap1-NRF2 pathway and activate p62-mediated selective autophagy. ROS also cause DNA damage, including double strand breaks (DSBs). p62 accumulated due to autophagy inhibition inhibits DSB repair through promoting proteasome-mediated degradation of CHK1 and RAD51 and interacting with RNF168. An appropriate level of endogenous p62-mediated autophagy in virus-transformed cells endows them with resistance to DNA damage. Loss of the p62-autophagy balance by exogenic stresses that inhibit autophagy or inducing ROS will exacerbate endogenous DNA damage.

Fig 1. p62-mediated selective autophagy is consititutively induced in viral latency and correlates with ROS-Keap1-NRF2 pathway activity

Fig 2. The ROS-Keap1-NRF2 pathway contributes to p62 expression and activation of p62-mediated autophagy in viral latency

Fig 3. Autophagy inhibition sensitizes EBV+ cells to ROS-induced DNA damage that is associated with p62 accumulation

Fig 4. Autophagy inhibition promotes cell death

Fig 5. p62 accumulation upon autophagy inhibition destabilizes HR DNA repair proteins CHK1 and RAD51 in viral latency

Fig 6. p62 depletion by RNA interference promotes RNF168-mediated chromatin ubiquitination and DNA repair in viral latency

Fig 7. p62 and autophagy are upregulated and correlates with ROS and DNA damage levels in B cell population of AIDS lymphoma patients

Fig 8. A diagram showing the interaction of p62-mediated autophagy with DDR in EBV latency