Estimating heritability and its enrichment in tissue-specific gene sets in admixed populations

Yang Luo[1-5,*], Xinyi Li[1-5,*], Xin Wang[6], Steven Gazal[3,7], Josep Maria Mercader[3,8], 23andMe

Research Team[6], SIGMA Type 2 Diabetes Consortium, Benjamin M. Neale[3,9], Jose C.

Florez[3,8,10], Adam Auton[6], Alkes L. Price[3,7,11], Hilary K. Finucane[3,#], Soumya Raychaudhuri[1-5,12,#]

Affiliations

[1]Division of Rheumatology, Immunology, and Allergy, Brigham and Women's Hospital, Harvard
Medical School, Boston, MA, USA
[2]Division of Genetics, Brigham and Women's Hospital, Harvard Medical School, Boston, MA,
USA
[3]Broad Institute of MIT and Harvard, Cambridge, MA, USA
[4]Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
[5]Center for Data Sciences, Brigham and Women's Hospital, Harvard Medical School, Boston,
MA, USA
[6]23andMe, Inc., Mountain View, California, USA
[7]Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston,
Massachusetts, USA
[8]Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital,
Boston, MA 02114, USA
[9]Analytic and Translational Genetics Unit, Massachusetts General Hospital and Harvard Medical
School, Boston, MA, USA
[10]Department of Medicine, Harvard Medical School, Boston, MA, USA
[11]Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, USA
[12]Arthritis Research UK Centre for Genetics and Genomics, Manchester Academic Health
Science Centre, University of Manchester, Manchester, UK
25

26 *: These authors contributed equally to this work.

27 #: Correspondence should be addressed to H.K.F. (finucane@broadinstitute.org) or S.R.

28 (soumya@broadinstitute.org).

Abstract

The increasing size and diversity of genome-wide association studies provide an exciting opportunity to study how the genetics of complex traits vary between diverse populations. Here, we introduce covariate-adjusted LD score regression (cov-LDSC), a method to accurately estimate genetic heritability (h_g^2) and its enrichment in both homogenous and admixed populations with summary statistics and in-sample LD estimates. In-sample LD can be estimated from a subset of the GWAS samples, allowing our method to be applied efficiently to very large cohorts. In simulations, we show that unadjusted LDSC underestimates h_g^2 by 10%- 60% in admixed populations; in contrast, cov-LDSC is robust to all simulation parameters. We apply cov-LDSC to genotyping data from approximately 170,000 Latino, 47,000 African American and 135,000 European individuals. We estimate h_g^2 and detect heritability enrichment in three quantitative and five dichotomous phenotypes respectively, making this, to our knowledge, the most comprehensive heritability-based analysis of admixed individuals. Our results show that most traits have high concordance of h_g^2 and consistent tissue-specific heritability enrichment between different ethnic groups. For example, estimates of h_g^2 for BMI are 0.22 ± 0.01, 0.23 ± 0.03 and 0.22 ± 0.01 in Latino, African American and European populations respectively. However, for age at menarche, we observe population-specific heritability differences with estimates of h_g^2. We observe consistent patterns of tissue-specific heritability enrichment across populations, for example in the limbic system for BMI, the per-standardized-annotation effect size τ^* are 0.16 ± 0.04, 0.28 ± 0.11 and 0.18 ± 0.03 in Latino, African American and European populations respectively. Our results demonstrate

50 **that our approach is a powerful way to analyze genetic data for complex traits from**

51 **underrepresented populations.**

Introduction

It is important for human geneticists to study how genetic variants that influence phenotypic

variability act across different populations worldwide[1,2]. With increasingly large and diverse

genetic studies, it is now becoming feasible to assess how the genetic mechanisms of complex

traits act across populations. However, to date, most genome-wide association studies (GWAS)

have been focused on relatively homogenous continental populations and, in particular those of

European descent[3]. Non-European populations, particularly those with mixed ancestral

backgrounds such as African Americans and Latinos, have been underrepresented in genetic

studies. Many statistical methods to analyze genetic data assume homogeneous populations. In

order to ensure that the benefits of GWAS are shared beyond individuals of homogeneous

continental ancestry, statistical methods for admixed populations are needed[4].

Among methods to analyze polygenic complex traits in homogeneous populations, summary

statistics-based methods such as linkage disequilibrium score regression (LDSC)[5,6] and its

extensions[5–8] have become particularly popular due to their computational efficiency, relative

ease of application, and their applicability without raw genotyping data[9]. These methods can be

used to estimate SNP heritability, the proportion of phenotypic variance explained by genotyped

variants[5,10–12], distinguish polygenicity from confounding[5], establish relationships between

complex phenotypes[7] , and model genome-wide polygenic signals to identify key cell types and

regulatory mechanisms of human diseases[6,13,14].

Summary statistics-based methods for polygenic analysis frequently rely on accurate linkage

disequilibrium (LD) calculations, which can be obtained on a random subset of the individuals

without phenotypic information. Consequently, LDSC, SumHer[15] and other summary

statistics-based methods have not easily been expanded to admixed populations. The LD score

for a SNP is the sum of its pairwise correlations (r^2) with all other SNPs. For most homogenous

populations, LD scores are easy to calculate for SNPs in local windows since r^2 between SNPs

more than one centimorgan (cM) is rarely nonzero. In admixed populations however, LD scores

are confounded since pairwise r^2 between SNPs is inflated due to admixture itself. Furthermore,

in homogenous populations LD scores can be easily calculated in reference panels since

representative data is widely available. For admixed populations, even when reference panels

are available, they may not be representative of the precise populations used in the genetic

study. For example Latino populations in different regions worldwide can be the result of

admixture between the same continental populations, but with dramatic differences in admixture

proportions and timing of the admixture event[16]. A generic reference panel cannot easily capture

these differences and hence cannot produce accurate LD scores that can be widely used for all

Latino populations. Thus, LDSC has only been recommended to be applied in homogeneous

populations.

In this work, we first examine the performance of LDSC in admixed populations and

demonstrate that LDSC yields severely downwardly biased estimates of SNP heritability. Next,

we extend the LDSC-based methods to admixed populations by introducing covariate-adjusted

LDSC (cov-LDSC). In cov-LDSC, for each variant we regress the global genotypic PCs,

obtained within the GWAS samples, out of the raw genotypes to obtain covariate-adjusted

genotypes. We then compute LD scores on the adjusted genotypes and use LDSC to estimate

heritability. Using covariate-adjusted in-sample LD to compute LD scores removes the potential

concerns of reference panel mismatch, long-distance admixture-LD, and covariate effects listed

above, and produces accurate estimates of heritability with summary statistics (**Methods,**

Figure 1). Furthermore, heritability can be partitioned to identify key gene sets that have

disproportionately high heritability. While access to the genotype data of the GWAS samples is

required to compute the covariate-adjusted LD scores, LD can be estimated on a random

subset of the individuals, preserving the computational efficiency of LDSC and allowing for its

application to very large studies. Individual cohorts can also release the in-sample

covariate-adjusted LD scores as well as the summary statistics to avoid privacy concerns

associated with genotype-level information to facilitate future studies.

After demonstrating that cov-LDSC is robust to a wide range of simulation scenarios, we apply it

to 8,124 Latinos from a type 2 diabetes study (the Slim Initiative in Genomic Medicine for the

Americas, SIGMA)[17] as well as 161,894 Latino, 46,844 African American, and 134,999

European research participants from a personal genetics company (23andMe). We analyze

three quantitative phenotypes (body mass index, height, and age at menarche), and five

dichotomous phenotypes (type 2 diabetes (available in the SIGMA cohort only), left

handedness, morning person, motion sickness, and nearsightedness).

One powerful component of LDSC is that it can be used to test whether a particular genome

annotation -- for example, sets of genes that are specifically expressed within a candidate tissue

or cell type -- capture more heritability than expected by chance[12,13]. We demonstrate that

cov-LDSC can be applied in the same way to identify trait-relevant tissue and cell types in

admixed and homogenous populations. We examine height, BMI and morning person since

these traits had sufficient statistical power for cell-type enrichment analyses in the personal

genetics cohort. We observe a high level of consistency among enriched tissue types,

highlighting that the underlying biological processes are shared among studied populations.

This analysis of hundreds of genome annotations in cohorts of over 100,000 individuals would

have been impossible with existing genotype-based methods[18,19].

Material and Methods

Mathematical framework of cov-LDSC

125   In the standard polygenic model on which LDSC is based, genotype vectors are represented as

126    $x_1, ..., x_N \in \mathbb{R}^M$, with each SNP normalized to mean zero and variance one in the population. In

127   the absence of covariates, we model the phenotypes y_i

$$y_i = x_i \cdot \beta + \epsilon_i \tag{1}$$

where $\epsilon_1, ..., \epsilon_N \overset{iid}{\sim} \mathcal{N}(0, \sigma_e^2)$ and $\beta \in \mathbb{R}^M$ is a vector of per-normalized-genotype effect sizes,

which we model as random with mean zero. In standard LDSC, the variance of β_j is $h_g{}^2/M$; in

stratified LD score regression the variance of β_j depends on a set of genome annotations. We

assume, without loss of generality, that y_i has mean zero and variance one in the population.

Let $\chi_j{}^2$ denote the chi-square statistic for the j-th SNP, approximately equal to $(X_j^T Y)^2/N$,

where $X_j = (x_{1j}, ..., x_{Nj})^T$ and $Y = (y_1, ..., y_n)^T$. The main equation on which LDSC is based

is:

$$\mathbb{E}[\chi_j^2] \approx 1 + Na + \frac{h_g^2}{M}\ell(j). \tag{2}$$

138   Where the LD score, $\ell(j)$, is:

$$\ell(j) = \sum R_{jk}^2,$$

$R_{jk}{}^2$ is the correlation between SNPs j and k in the underlying population. We estimate $h_g{}^2$ via

weighted regression of χ_j^2 on our estimates of $\ell(j)$, evaluating significance with a block

jackknife across SNPs[6].

In the absence of covariates, the LD scores can be estimated from an external reference panel

such as 1000 Genomes, as long as the correlation structure in the reference panel matches the

correlation structure of the sample. In most homogeneous populations, we can also assume that

the true underlying correlation is negligible outside of a 1-cM window.

Next, suppose that we would like to include covariates in our model, such that Equation (1) is

replaced with

$$y_i = x_i \cdot \beta + c_i \cdot \eta + \epsilon_i \tag{3}$$

152   or in matrix form,

$$Y = X\beta + C\eta + \epsilon_i \tag{4}$$

154   where c_i is the vector of covariate values for individual i, η is a vector of effect sizes of the

155   covariates, X is the genotype matrix with $X_{ij} = x_{ij}$, and C is a matrix of covariates with

156    $C_{ij} = c_{ij}$. We can project the covariates out of this equation by multiplying

157    $P = I - C(C^T C)^{-1} C^T$ on the left to get

$$\tilde{Y} = \tilde{X}\beta + \tilde{\epsilon}, \tag{5}$$

where $\tilde{Y} = PY$, $\tilde{X} = PX$ and $\tilde{\epsilon} = P\epsilon$ (if the covariates are genotype principal components,

then $P = 1 - CC^T$). When sample size is large relative to the number of covariates, then the $\tilde{\epsilon}$

are still approximately independent and normally distributed, and the variance of $\tilde{y}_i$ is still

approximately one, so we can proceed as before, replacing our phenotype vector Y with $\tilde{Y}$ and

our genotype matrix X with $\tilde{X}$.

The summary statistics in this covariate-adjusted model are the standard summary statistics of

a GWAS with covariates, and the relevant LD is the correlation structure of $\tilde{X}$, not X. If X is a

homogeneous population, then the correlation structure of $\tilde{X}$ will be similar to the correlation

structure of X, and well-approximated by a reference panel. However if X is the genotype

matrix from an admixed population and the covariates include PCs, then the correlation

structure of $\tilde{X}$ is no longer well-approximated either by X or by a reference panel. Thus, in

cov-LDSC, we compute LD scores directly from $\tilde{X}$ or a random subsample thereof. We call

them the covariate-adjusted LD scores.

Using genotype data to compute LD scores means that the model being fit is based on the joint

effects of a sparser set of SNPs, e.g. the genotyped SNPs, than when sequence data is used to

compute LD scores. For estimating total SNP heritability, this means that cov-LDSC estimates

the same estimand as GCTA ($h_g{}^2$) and not the usual estimand of LDSC ($h_{common}{}^2$; see below).

For partitioned heritability, the density of reference panel SNPs can be important because the

joint effect of a SNP in an annotation can include the tagged effect of an untyped SNP that is not

in the annotation, deflating estimates of enrichment. Thus, we recommend using cov-LDSC only

on annotations made of large contiguous regions, such as gene sets. Moreover, we urge

caution when interpreting quantitative estimates of heritability enrichment. Here, we look at the

significance of the conditional enrichment (i.e., regression coefficient) of gene sets for our

tissue-specific analysis (see below).

Window size and number of PCs in LD score calculations

In addition to computing LD from the covariate-adjusted genotypes, we also investigate the

appropriate window size for estimating LD scores. To do this, we examine the effect of varying

the genomic window size for both simulated and real data sets. We determine that LD score

estimates were robust to the choice of window size if the increase in the mean LD score

estimates was less than 1% per cM beyond a given window. Using this criterion, we use window

sizes of 5-cM and 20-cM for the simulated and real genotypes respectively (**Supplementary**

Table 1-3). We also calculate the squared correlations between LD score estimates using the

chosen window size and other LD score estimates with window sizes larger than the chosen

window. The squared correlations were greater than 0.99 in all cases (**Supplementary Table**

4-6) indicating the LD score estimates were robust at the chosen window sizes.

Similarly, to determine the number of PCs needed to be included in the GWAS association tests

and cov-LDSC calculations, we examine the effect of varying the genomic window size using

different numbers of PCs. The number of PCs that needed to be included for covariate

adjustment depended on the population structure for different datasets.

Genotype simulations

We evaluate the performance of LDSC and cov-LDSC with simulated phenotypes and both

simulated and real genotypes. For the simulated genotypes, we used msprime[20] version 0.6.1 to

simulate population structure with mutation rate 2×10^{-8} and recombination maps from the

HapMap Project[21]. We adapt the demographic model from Mexican migration history[22] using

parameters that were previously inferred from the 1000 Genomes Project[23]. We assume the

admixture event happened approximately 500 years ago to mirror the European colonization of

the Americas. We set different admixture proportions to reflect different admixed populations. In

each population, we simulate 10,000 individuals after removing second degree related samples

(kinship>0.125) using KING[24].

Slim Initiative in Genomic Medicine for the Americas (SIGMA) Type 2 Diabetes (T2D) cohort

8,214 Mexican and other Latin American samples were genotyped with Illumina HumanOmni2.5

array. We further filter the genotyped data to be MAF >5% and remove SNPs in high LD regions

(**Supplementary Table 7**). After QC, a total of 8,214 individuals and 943,244 SNPs remain. We

estimate the in-sample LD score with a 20-cM window and 10 PCs in all scenarios.

We use these genotypes for simulations. We also analyze three phenotypes from the SIGMA

cohort: height, BMI, and type 2 diabetes (T2D). For T2D, we assume a reported prevalence in

Mexico of 0.144[17]. For each phenotype, we include age, sex, and the first 10 PCs as fixed

effects in the association analyses.

Phenotype simulations

We simulate phenotypes with two different polygenic genetic architectures, given by GCTA[19]

and the baseline model[6] respectively. In the GCTA model, all variants are equally likely to be

causal independent of their functional or minor allele frequency (MAF) structure, and the

standardized causal effect size variance is constant, i.e. $var(\beta_j) = h_g^2/M$. In contrast, the

baseline model incorporates functionally dependent architectures. Briefly, it includes 53

overlapping genome-wide functional annotations (e.g. coding, conserved, regulatory). It models

$var(\beta_j) = \sum_C \alpha_c(j)\tau_c$ where $\alpha_c(j)$ is the value of annotation α_c at variant j and τ_c represents

the per-variant contribution, of one unit of the annotation α_c, to heritability. We generate all

causal variants among common observed variants with MAF >5% (~40,000 SNPs in simulated

genotypes and 943,244 SNPs in the SIGMA cohort). To represent environmental stratification,

similar to previously described[5], we add 0.2 * standardized first principal component to the

standardized phenotypes.

We simulate both quantitative and case-control traits with both GCTA and baseline model

genetic architectures, using both simulated and real genotypes, varying the number of causal

variants, the true heritability, and environmental stratification. For case-control simulations, we

adopt a liability threshold model with disease prevalence 0.1. We obtain 5,000 cases and 5,000

controls for each simulation scenario.

To obtain summary statistics for the simulated traits, we apply single-variant linear models for

quantitative traits and logistic models for binary trait both with 10 PCs as covariates in

association analyses using PLINK 1.90[25].

23andMe cohort

All participants were drawn from the customer base of 23andMe, Inc., a direct to consumer

genetics company. Participants provided informed consent and participated in the research

online, under a protocol approved by the external AAHRPP-accredited IRB, Ethical &

Independent Review Services (www.eandireview.com). Samples from 23andMe are then

chosen from consented individuals who were genotyped successfully on an Illumina Infinium

Global Screening Array (~640,000 SNPs) supplemented with ~50,000 SNPs of custom content.

We restrict participants to those who have European, African American, or Latino ancestry

determined through an analysis of local ancestry[26].

To compute LD scores, we use both genotyped and imputed SNPs. We filter genotyped variants

with a genotype call rate $\leq$ 90%, non-zero self-chain score, strong evidence of Hardy Weinberg

disequilibrium ($p > 10^{-20}$), and failing a parent-offspring transmission test. For imputed variants,

we use a reference panel that combined the May 2015 release of the 1000 Genomes Phase 3

haplotypes[23] with the UK10K imputation reference panel[27]. Imputed dosages are rounded to the

nearest integer (0, 1, 2) for downstream analysis. We filter variants with imputation r-squared $\leq$

0.9. We also filter genotyped and imputed variants for batch effects and sex dependent effects.

To minimize rounding inaccuracies, we prioritize genotyped SNPs over imputed SNPs in the

merged SNP set. We restrict the merged SNP set to HapMap3 variants with MAF $\geq$ 0.05. We

measure LD scores in a subset of African Americans (61,021) and Latinos (9,990) on

chromosome 2 with different window sizes from 1-cM to 50-cM (**Supplementary Table 3**) and

squared correlation between different window sizes (**Supplementary Table 6**). We compute all

LD scores with a 20-cM window.

In genome-wide association analyses, for each population, we choose a maximal set of

unrelated individuals for each analysis using a segmental identity-by-descent (IBD) estimation

algorithm[28]. We define individuals to be related if they share more than 700-cM IBD.

We perform association tests using linear regression model for quantitative traits and logistic

regression model for binary traits assuming additive allelic effects. We include covariates for

age, sex and the top 10 PCs to account for residual population structure. We list details of

phenotypes and genotypes in **Supplementary Table 8.**

Heritability Estimation

We calculate in-sample LD scores using both a non-stratified LD score model and the baseline

model[6]. In simulated phenotypes generated with the GCTA model, we use non-stratified LDSC

to estimate heritability. In simulated phenotypes generated using the baseline model, we use

LDSC-baseline to estimate heritability. We use the 53 non-frequency dependent annotations

included in the baseline model to estimate h_g^2 in the 23andMe research database and the

SIGMA cohort real phenotypes. We recognize recent studies have shown that genetic

heritability can be sensitive to the choice of LD-dependent heritability model[8,11]. However,

understanding the LD- and MAF-dependence of complex trait genetic architecture is an

important but complex endeavor potentially requiring both modeling of local ancestry as well as

large sequenced reference panels that are currently unavailable. We thus leave this complexity

for future work.

h_g^2 versus h_{common}^2

The quantity (h_g^2) we reported in the main analysis is defined as heritability tagged by

HapMap3 variants with MAF $\geq 5\%$, including tagged causal effects of both low-frequency and

common variants. This quantity is different from h_{common}^2, the heritability casually explained by

all common SNPs excluding tagged causal effects of low-frequency variants, reported in the

original LDSC[5]. In Europeans and other homogeneous populations, it is possible to estimate

h_{common}^2, since reference panels, such as 1000 Genome Project, are available which include

>99% of the SNPs with frequency >1%[23]. However, in-sample sequence data is usually not

available for an admixed GWAS cohort, and so cov-LDSC can only include genotyped SNPs in

the reference panel, and thus can only estimate the heritability tagged by a given set of

genotyped SNPs. In order to compare the same quantity across cohorts, we use common

HapMap3 SNPs (MAF $\geq$ 5%) for in-sample LD reference panel calculation, since most of them

should be well imputed for a genome-wide genotyping array. To quantify the difference between

h_g^2 and h_{common}^2, we pre-phase the genotype data in the SIGMA cohort using SHAPEIT2[29]. We

use IMPUTE2[30] to impute genotypes at untyped genetic variants using the 1000 Genomes

Project Phase 3[23] dataset as a reference panel. We merge genotyped SNPs and all well

imputed (INFO>0.99) SNPs (~6.9 million) in the SIGMA cohort as a reference panel and

reported h_{common}^2, to approximate what the estimate of h_{common}^2 would have been with a

sequenced reference panel (**Supplementary Table 9**).

Tissue Type Specific Analyses

Following Finucane et al[13], we extend cov-LDSC so that we can assess enrichment in and

around sets of genes that are specifically expressed in tissue and cell-types (cov-LDSC-SEG).

We annotate the genes with the same set of tissue specific expressed genes identified

previously[13] using the Genotype–Tissue Expression (GTEx) project and a public dataset made

available by the Franke lab. We calculate within-sample stratified cov-LD scores in the 23andMe

cohort for each of these gene sets. We obtain regression coefficients $\hat{\tau}_c$ from the model and

normalize them as

307
$$\tau_c^* = \frac{M_{h_g^2} \cdot sd_c}{h_g^2} \hat{\tau}_c,$$

Where $M_{h_g^2}$ $M_{h_g^2}$ is the number of SNPs used to calculate h_g^2 and sd_c is the standard

deviation (sd) of annotation a_c[8]. We interpret τ_c^* as the proportional change of averaged

per-SNP heritability by one sd increase in value of the annotation of each cell type, conditional

on other 53 non-cell type specific baseline annotations. We calculate a one-tailed p-value for

each coefficient where the null hypothesis is that the coefficient is non-positive[13]. All the

significant enrichments are reported with false discovery rate < 5% $(-log_{10}(p) > 2.75)$. We

perform fixed-effect inverse variance weighting meta-analysis using τ_c^* and normalized standard

error (se) across populations.

Software Availability

316 An open-source software implementation of covariate-adjusted LD score regression is publicly

317 available (see Web Resources).

Results

Robustness of LD score estimation

To demonstrate the effect of admixture on the stability of LD score estimates, we first calculated

LD scores with genomic window sizes ranging from 0-50 cM in both European (EUR, N=503)

and admixed American (AMR, N=347) populations from the 1000 Genomes Project[23]. As

window size increases, we expect the mean LD score to plateau because LD should be

negligible for large enough distance. If the mean LD score does not plateau, but continues to

rise with increasing window size, then one of two possibilities may apply: (1) the window is too

small to capture all of the LD; (2) the LD scores are capturing long-range pairwise SNP

correlations arising from admixture. If this increase is non-linear then there is non-negligible

distance-dependent LD, violating LDSC assumptions. Examining unadjusted LD scores, we

observed that in the EUR population[5], the mean LD score estimates plateaued at windows

beyond 1-cM in size, as previously reported. However, in the AMR population the mean LD

score estimates continued to increase concavely with increasing window size. In contrast, when

we applied cov-LDSC with 10 PCs to calculate covariate adjusted LD scores, we observed that

LD score estimates plateaued for both EUR and AMR at a 1-cM and 20-cM window size

respectively (<1% increase per cM, **Supplementary Table 10**). This suggested that cov-LDSC

was able to correct the long-range LD due to admixture and yielded stable estimates of LD

scores (**Method, Supplementary Figure 1**), and also that cov-LDSC was applicable in

homogeneous populations (**Supplementary Table 10**). The larger window size for the AMR

population was needed due to residual LD caused by recent admixture. We next tested the

sensitivity of the LD score estimates with regard to the number of PCs included in the

cov-LDSC. We observed that in the AMR panel, LD score estimates were unaffected by adding

PCs and by increasing window sizes above 20-cM (**Supplementary Figure 2**).

Simulations with simulated genotypes

To assess whether cov-LDSC produces unbiased estimates of h_g^2, we first simulated

genotypes of admixed individuals (**Methods**). We simulated genotypes of 10,000 unrelated

diploid individuals for approximately 400,000 common SNPs on chromosome 2 in a coalescent

framework using msprime[20]. First, we tested LDSC and cov-LDSC with different admixture

proportions between two ancestral populations, and a quantitative phenotype with a h_g^2 of 0.4

using an additive model (**Methods**). We observed that as the proportion of admixture increased,

$\hat{h}_g^2$ for LDSC increasingly underestimated true h_g^2 by as much as 18.6%. In marked contrast,

cov-LDSC produced consistently unbiased estimates regardless of admixture proportion

(**Supplementary Figure 3a**).

Second, we varied the percentage of causal variants from 0.01% to 50% in a polygenic

quantitative trait with $h_g{}^2 = 0.4$ in a population with a fixed admixture proportion of 50%. LDSC

again consistently underestimated $h_g{}^2$ by 12%-18.6%. In contrast, cov-LDSC yielded unbiased

estimates regardless of the percentage of causal variants (**Supplementary Figure 3b**).

Third, we assessed the robustness of LDSC and cov-LDSC for different assumed total $h_g{}^2$

(0.05, 0.1, 0.2, 0.3, 0.4 and 0.5). At each $h_g{}^2$ value, LDSC underestimated by 11.5%-19.6%. For

cov-LDSC, we observed that the standard error increased with $h_g{}^2$, but point estimates

remained unbiased (**Supplementary Figure 3c**).

Fourth, we included an environmental stratification component aligned with the first PC of the

genotype data (**Methods**), and concluded that cov-LDSC was also robust to confounding

(**Supplementary Figure 3d**).

Finally, to assess the performance of cov-LDSC in polygenic binary phenotypes, we simulated

genotype data for a binary trait with a prevalence of 0.1 assuming a liability threshold model

(**Methods**). We showed that cov-LDSC provided unbiased estimates in case-control studies

with the same four simulation scenarios (**Supplementary Figure 4**). In contrast, LDSC

underestimated heritability for binary phenotypes in the same way as it did for quantitative

phenotypes.

Simulation results with real genotypes

We next examined the performance of both unadjusted LDSC and cov-LDSC on real genotypes

of individuals from admixed populations. We used genotype data from SIGMA, which includes

8,214 Mexican and other Latino individuals. Using ADMIXTURE[31] and populations from the

1000 Genomes Project as reference panels, we observed that each individual in the SIGMA

cohort had different admixture proportions (**Supplementary Figure 5**). As in the AMR panel, we

observed that using a 20-cM window, LD score estimates plateaued in the SIGMA cohort

(**Supplementary Figure 6, Supplementary Table 2**), and were unaffected by different numbers

of PCs (**Supplementary Figure 7**). When we assumed a non-infinitesimal, additive model with

1% of all SNPs to be causal and h_g^2 =0.4, we observed that cov-LDSC h_g^2 estimates produced

unbiased estimates using a 20-cM window with 10 PCs (**Supplementary Figure 8**). We

subsequently used a 20-cM window and 10 PCs in all simulations.

We observed that cov-LDSC yielded unbiased h_g^2 estimates in simulated traits where we varied

the number of causal variants and total heritabilities (**Figure 2a-b**). In contrast, LDSC

underestimated heritability by as much as 62.5%. To examine the performance of cov-LDSC in

the presence of environmental confounding factors, we simulated an environmental stratification

component aligned with the first PC of the genotype data, representing European v.s. Native

American ancestry. In this simulation scenario, cov-LDSC still provided unbiased h_g^2 estimates

(**Figure 2c**). Intercepts of all the simulation scenarios were close to 1, suggesting that we had

adequately controlled for confounding from population stratification and cryptic relatedness

(**Supplementary Figure 9a-c**). We observed unbiased h_g^2 estimates with both the GCTA model

and baseline model (**Supplementary Table 11**).

Thus far, we have used cov-LDSC by calculating LD scores on the same set of samples that

were used for association studies (in-sample LD scores). In practical applications, computing LD

scores on the whole data set can be computationally expensive and difficult to obtain, so we

investigated computing LD scores on a subset of samples. To investigate the minimum number

of samples required to obtain accurate in-sample LD scores, we computed LD scores on

subsamples of 100, 500, 1,000 and 5,000 individuals from a GWAS of 10,000 simulated

genotypes (**Supplementary Figure 10**). We repeated these analyses in simulated phenotypes

in the SIGMA cohort. We subsampled the SIGMA cohort, and obtained unbiased estimates

when using as few as 1,000 samples (**Figure 2d**). We therefore recommend computing

in-sample LD scores on a randomly chosen subset of at least 1,000 individuals from a GWAS in

our approach.

Application to SIGMA and 23andMe cohorts

We next used cov-LDSC to estimate h_g^2 of height, BMI and T2D phenotypes, measured within

the SIGMA cohort (**Methods, Table 1**). We estimated h_g^2 of height, BMI and T2D to be 0.38 ±

0.08, 0.25 ± 0.06 and 0.26 ± 0.08 respectively. These results were similar to reported values

from UK Biobank[32] and other studies[10,33] for European populations. Although estimands differed

in different studies (**Methods**), we noted that without cov-LDSC, we would have obtained

severely deflated estimates (**Table 1**). To confirm that our reported heritability estimates were

robust under different model assumptions, we applied an alternative approach based on REML

in the linear mixed model framework implemented in GCTA[19]. To avoid biases introduced from

calculating genetic relatedness matrices (GRMs) in admixed individuals, we obtained a GRM

based on an admixture-aware relatedness estimation method REAP[34] (**Methods**). GCTA-based

results were similar to reported h_g^2 estimates from cov-LDSC, indicating our method was able

to provide reliable h_g^2 estimates in admixed populations (**Table 1**). We noted, however, that the

GCTA-based results would be impossible to obtain on the much larger datasets, for example the

23andMe cohort described below.

We then applied both LDSC and cov-LDSC to 161,894 Latino, 46,844 African American and

134,999 European research participants from 23andMe. We analyzed three quantitative and

four dichotomous phenotypes (**Methods**). In this setting, summary statistic methods to estimate

heritability were essential since the dataset would be too large for computationally expensive

genotype-based strategies. We used a 20-cM window and 10 PCs in LD score calculations for

both populations (**Supplementary Figure 11**). LDSC and cov-LDSC produced similar

heritability estimates in the European population, whereas in the admixed populations, LDSC

consistently provided low estimates of $h_g{}^2$. Heritability estimates were similar between

non-stratified and baseline models (**Supplementary Table 12**). For each phenotype, we

estimated $h_g{}^2$ using the same population-specific in-sample LD scores. For most phenotypes,

the reported $h_g{}^2$ was similar among the three population groups with a notable exception for

age at menarche (**Figure 3**). This suggested possible differences ($p = 7.1 \times 10^{-3}$ between

Latinos and Europeans) in the genetic architecture of these traits between different ethnic

groups. It has been long established that there is population variation in the timing of

menarche[35,36]. Early menarche might influence the genetic basis of other medically relevant

traits since early age at menarche is associated with a variety of chronic diseases such as

childhood obesity, coronary heart disease and breast cancer[37,38]. These results highlighted the

importance of including diverse populations in genetic studies in order to enhance our

understanding of complex traits that show differences in their genetic heritability.

Tissue type specific analysis

We applied stratified cov-LDSC to sets of specifically expressed genes[13] (SEG) to identify

trait-relevant tissue and cell types in traits included in the 23andMe cohort across European,

Latino, and African American populations. We only tested height, BMI and morning person,

which were the three traits that had heritability z-scores larger than seven in at least two

populations[6] (**Supplementary Table 13**). We also performed inverse-variance weighting

meta-analysis across the three populations (**Supplementary Table 14**). Across different

populations, BMI showed consistent enrichment in central nervous system gene sets . In the

European population, most of the enrichments recapitulated the results from the previous

analysis using UK Biobank[13]. We found similar but fewer enrichments in Latinos and African

Americans, most likely due to smaller sample sizes. The most significantly enriched tissue types

for BMI in all three populations were limbic system ($\tau^{*}_{EUR} = 0.18$, $\tau^{*}_{LAT} = 0.16$, $\tau^{*}_{AA} = 0.28$,

$\tau^{*}_{meta} = 0.18$), entorhinal cortex ($\tau^{*}_{EUR} = 0.18$, $\tau^{*}_{LAT} = 0.15$, $\tau^{*}_{AA} = 0.24$, $\tau^{*}_{meta} = 0.17$), and

cerebral cortex ($\tau^{*}_{EUR} = 0.16$, $\tau^{*}_{LAT} = 0.14$, $\tau^{*}_{AA} = 0.15$, $\tau^{*}_{meta} = 0.15$); none of the three effects

were significantly different across populations. When we compared the enrichments for all of the

tissues between population pairs, we observed that they were similar with regression slopes

close to one (European vs. Latinos: slope=0.69, se=0.036, $p = 3.1 \times 10^{-48}$; European vs.

African Americans: slope=0.67, se=0.12, $p = 3.4 \times 10^{-8}$) (**Figure 4(a)-(e)**). The sizes of these

three brain structures have been shown to be correlated with BMI using magnetic resonance

imaging data[39]. The midbrain and the limbic system are highly involved in the food rewarding

signals through dopamine releasing pathway[40]. Furthermore, the hypothalamus in the limbic

system releases hormones that regulate appetite, energy homeostasis and metabolisms, like

leptin, insulin, and ghrelin[40,41]. For height, similar to previously reported associations[13], we also

identified enrichments in the gene sets derived from musculoskeletal and connective tissues. In

the meta-analysis, the three most significant enrichments were cartilage ($\tau^{*}_{EUR} = 0.21$,

$\tau^{*}_{LAT} = 0.19$, $\tau^{*}_{AA} = 0.24$, $\tau^{*}_{meta} = 0.20$), chondrocytes ($\tau^{*}_{EUR} = 0.21$, $\tau^{*}_{LAT} = 0.15$, $\tau^{*}_{AA} = 0.11$,

$\tau^*_{meta} = 0.17$), and uterus ($\tau^*_{EUR} = 0.17$, $\tau^*_{LAT} = 0.15$, $\tau^*_{AA} = 0.16$, $\tau^*_{meta} = 0.16$) (**Figure 4(f)-(j)**).

A heterogeneity test revealed no difference across three populations (I^2 < 70% and p-value >

0.05). The slopes of the regression of the two population coefficients were close to 1 (European

vs. Latinos: slope = 0.76, se=0.020, $p = 4.9 \times 10^{-94}$; European vs. African Americans: 0.59,

se=0.055, $p = 1.4 \times 10^{-21}$). The importance of these tissues and their roles in height have been

addressed in the previous pathway analysis, expression quantitative trait loci (eQTLs) and

epigenetic profiling[42,43]. Previous studies have shown that the longitudinal growth of bones is

partly controlled by the number and proliferation rate of chondrocytes on the growth plate which

is a disc of cartilages[44]. For the morning person phenotype, we found enrichments in many brain

tissues in Europeans, concordant with a previous study[45]. Entorhinal cortex ($\tau^*_{EUR} = 0.16$,

$\tau^*_{LAT} = 0.22$, $\tau^*_{meta} = 0.18$), cerebral cortex ($\tau^*_{EUR} = 0.15$, $\tau^*_{LAT} = 0.22$, $\tau^*_{meta} = 0.18$), and brain (

$\tau^*_{EUR} = 0.17$, $\tau^*_{LAT} = 0.19$, $\tau^*_{meta} = 0.18$) were enriched in both Latinos and Europeans.

Evidence showed that circadian rhythm was controlled by the suprachiasmatic nucleus, the

master clock in our brain, and also the circadian oscillator that resides in neurons of the cerebral

cortex [46,47,48]. We also found unique enrichments of esophagus muscularis and the esophagus

gastroesophageal junction in the Latino populations, but the heterogeneity test showed that the

difference is not significant (I^2 = 0.49 and 0.50 respectively). We observed that the slope of

regression between Latino coefficients and European coefficients across gene sets was 0.70

(se=0.062, $p = 1.9 \times 10^{-23}$) (**Figure 4(k)-(n)**). Compared to the original LDSC-SEG,

cov-LDSC-SEG increased the statistical power in African Americans and Latinos

(**Supplementary Figure 12-14**).

Discussion

As we expand genetic studies to explore admixed populations around the world, extending

statistical genetics methods to make inferences within admixed populations is crucial. This is

particularly true for methods based on summary statistics, which are dependent on the use of

LD scores that we showed to be problematic in admixed populations. In this study, we confirmed

that original LDSC and other summary statistics-based methods, such as PCGCs[49] and

SumHer[50] that were originally designed for homogenous populations, should not be applied to

admixed populations. We introduced cov-LDSC which regresses out global PCs on individual

genotypes during the LD score calculation, and showed it can yield unbiased LD scores,

heritability estimates and its enrichment, such as trait-relevant cell and tissue type enrichments,

in homogenous and admixed populations.

Although our work provides a novel, efficient approach to estimate genetic heritability and to

identify trait-relevant cell and tissue types using summary statistics in admixed populations, it

has a few limitations (**Methods**). First, covariates included in the summary statistics should

match the covariates included in the covariate-adjusted LD score calculations (**Supplementary**

Figure 15), and $h_g{}^2$ estimates in admixed populations are more sensitive to potentially

unmatched LD reference panels. Unmatched reference panels are likely to produce biased

estimates[51,52]. We examined the performance of using an out-of-sample reference panel in

admixed populations (See **Appendix**) and caution that when using 1000 Genomes or any

out-of-sample reference panels for a specific admixed cohort, users should ensure that the

demographic histories are shared between the reference and the study cohort. Large

sequencing projects such as TOPMed[53] that include large numbers (N>1,000) of admixed

samples can potentially serve as out-of-sample LD reference panels, although further

investigations are needed to study their properties. We therefore advise to compute in-sample

LD scores from the full or a random subset of data (N>1,000) used to generate the admixed

GWAS summary statistics when possible. We also strongly encourage cohorts to release their

summary statistics and in-sample covariate-adjusted LD scores at the same time to facilitate

future studies. Second, when applying cov-LDSC to imputed variants, particularly those with

lower imputation accuracy (INFO <0.99), we caution that the heritability estimates can be

influenced by an imperfect imputation reference panel, especially in Latino populations[54,55]. To

limit the bias in varying genotyping array and imputation quality in studied admixed cohorts, we

recommend restricting the heritability analyses to common HapMap3 variants. Any extension to

a larger set of genetic variants, especially across different cohorts should be performed with

caution. Third, when we evaluated the performance of cov-LDSC in case-control studies, we

assumed no presence of binary covariates with strong effects and demonstrated that cov-LDSC

can yield robust $h_g{}^2$ estimates. However, it has been shown that LDSC can provide biased

estimates in the presence of extreme ascertainment for dichotomous phenotypes[49]. Adapting

cov-LDSC into case-control studies under strong binary effects remains a potential avenue for

future work. Fourth, recent studies have shown that heritability estimates can be sensitive to the

choice of the LD- and frequency-dependent heritability model[8,11,12,15]. Since our approach can

flexibly add annotations to estimate heritability under the model that is best supported by the

data, we believe it provides a good foundation for addressing the question of how to incorporate

ancestry-dependent frequencies in the LD-dependent annotation in the future (**Methods**).

Despite these limitations, in comparison with other methods, such as those based on restricted

maximum likelihood estimation (REML)[19] with an admixture-aware GRM, for estimating $h_g{}^2$ in

admixed populations or those with intra-population structure, cov-LDSC has a number of

attractive properties. First, covariate-adjusted in-sample LD scores can be obtained with a

subset of samples, enabling analysis of much larger cohorts than was previously possible.

Second, LD scores only need to be calculated once per cohort; this is particularly useful in large

cohorts such as 23andMe and UK Biobank[56], where multiple phenotypes have been collected

per individual and per-trait heritability can be estimated based on the same LD scores. Second,

as a generalized form of LDSC, it is robust to population stratification and cryptic relatedness in

both homogenous and admixed populations. Third, similar to the original LDSC methods,

cov-LDSC can be extended to perform analyses such as estimating genetic correlations,

partitioning h_g^2 by functional annotations, identifying disease-relevant tissue and cell types and

multi-trait analysis[6,57,58].

By applying cov-LDSC to approximately 344,000 individuals from European, African American,

and Latin American ancestry, we observed evidence of heritability differences across different

populations. These differences highlight the importance of studying diverse populations. How

these differences may correspond to differences in biological mechanisms may lead to

mechanistic insights about the phenotype. One strategy to do this, which we explored by

extending cov-LDSC, is to partition heritability by different cell type- and tissue-specific

annotations to dissect the genetic architecture in admixed populations. Our results

demonstrated that although there are some cases of nominal heterogeneity across populations

among tested tissue-types, most of the tissue-specific enrichments are consistent among the

populations studied here. This is consistent with the previous findings that show strong

correspondence in functional and cell type enrichment between Europeans and Asians[59,60].

Seeing the same tissue-type for a single trait emerge in multiple populations can give us more

confidence that this tissue may account for polygenic heritability. Larger sample sizes are

needed to increase the power of our current analyses and to enhance our understanding of how

genetic variants that are responsible for heritable phenotypic variability differ among

populations.

As the number of admixed and other diverse GWAS and biobank data become readily

available[1,53,61], our approach provides a powerful way to study admixed populations.

Appendix

In-sample versus out-of-sample LD

To test the reliability of using an out-of-sample reference LD panel for cov-LDSC applications,

we first examined the performance of out-of-sample LD scores obtained from 1,000 samples

with a perfectly matching demographic history in the simulated genotypes. cov-LDSC yielded

unbiased estimates when using 1,000 samples in an out-of-sample reference panel with a

perfectly matching population structure (**Supplementary Figure 10**). Next, we tested the

accuracy of heritability estimates when using 1000 Genome Project to obtain out-of-sample LD

scores. When using the AMR panel as a reference panel for the SIGMA cohort, we observed an

unbiased h_g^2 estimate ($p = 0.33$, **Figure 2d**). This suggested that the AMR panel included in

the 1000 Genomes Project has similar demographic history compared to the SIGMA cohort

(**Supplementary Figure 5**). However, as we decreased the number of samples included in the

subsampling, the cov-LDSC regression intercepts deviated further from 1 (**Supplementary

Figure 9d**). This is probably due to attenuation bias from noisily estimated LD scores at

N<1,000. Next, we explored the feasibility of applying an out-of-sample reference panel in cell

and tissue type specific analyses. To this end, we used 1000 Genomes AMR samples (N=347)

to obtain stratified LD scores and applied it on summary statistics obtained from 23andMe. In

contrast to using in-sample LD scores, we discovered no significant tissue type enrichment

(**Supplementary Figure 16**) when using LD scores obtained from the 1000 Genomes. This

suggested an out-of-sample reference panel may significantly reduce the power for detecting

trait relevant tissue types in admixed populations.

We therefore caution that when using 1000 Genomes or any out-of-sample reference panels for

a specific admixed cohort, users should ensure that the demographic histories are shared

between the reference and the study cohort. We highly recommend computing in-sample LD

scores on a randomly chosen subset of at least 1,000 individuals from a GWAS. We also

strongly encourage cohorts to release their summary statistics and in-sample covariate-adjusted

LD scores at the same time to facilitate future studies.

Web Resources

cov-LDSC software and tutorials, https://github.com/immunogenomics/cov-ldsc

msprime, https://pypi.python.org/pypi/msprime;

GCTA, http://cnsgenomics.com/software/gcta/;

LDSC, https://github.com/bulik/ldsc/;

PLINK 1.90, https://www.cog-genomics.org/plink2;

REAP v1.2, http://faculty.washington.edu/tathornt/software/REAP/download.html;

ADMIXTURE v1.3.0, http://www.genetics.ucla.edu/software/admixture/download.html;

Acknowledgements

594 The study was supported by the National Institutes of Health (NIH) TB Research Unit Network,

595 Grant U19 AI111224-01. The content is solely the responsibility of the authors and does not

596 necessarily represent the official views of the NIH.

598 We thank the research participants and employees of 23andMe for their contribution to this

599 study.

Author contributions

Y.L., X.L, H.K.F and S.R. conceived and supervised the study. Y.L. and X.L. analyzed data. X.W.

and A.A. contributed and analyzed the 23andMe data. S.G., B.M.N. and A.L.P. gave critical

feedback on LDSC and statistical models. J.M.M. and J.C.F. contributed the SIGMA study. All

authors contributed to the writing of this manuscript.

Members of the 23andMe Research Team:

Michelle Agee, Babak Alipanahi, Robert K. Bell, Katarzyna Bryc, Sarah L. Elson, Pierre

Fontanillas, Nicholas A. Furlotte, Barry Hicks, David A. Hinds, Karen E. Huber, Ethan M. Jewett,

Yunxuan Jiang, Aaron Kleinman, Keng-Han Lin, Nadia K. Litterman, Matthew H. McIntyre,

Kimberly F. McManus, Joanna L. Mountain, Elizabeth S. Noblin, Carrie A.M. Northover, Steven

J. Pitts, G. David Poznik, J. Fah Sathirapongsasuti, Janie F. Shelton, Suyash Shringarpure,

Chao Tian, Joyce Y. Tung, Vladimir Vacic, and Catherine H. Wilson.

Competing Financial interests

612 X.W., A.A. and members of the 23andMe Research Team are employees of 23andMe, Inc., and

613 hold stock or stock options in 23andMe.

References

1. Sirugo, G., Williams, S.M., and Tishkoff, S.A. (2019). The Missing Diversity in Human Genetic Studies. Cell *177*, 26–31.

2. Martin, A.R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B.M., and Daly, M.J. (2019). Clinical use of current polygenic risk scores may exacerbate health disparities. Nat. Genet. *51*, 584–591.

3. Popejoy, A.B., and Fullerton, S.M. (2016). Genomics is failing on diversity. Nature *538*, 161–164.

4. Seldin, M.F., Pasaniuc, B., and Price, A.L. (2011). New approaches to disease mapping in admixed populations. Nat. Rev. Genet. *12*, 523–528.

5. Bulik-Sullivan, B.K., Loh, P.-R., Finucane, H.K., Ripke, S., Yang, J., Schizophrenia Working Group of the Psychiatric Genomics Consortium, Patterson, N., Daly, M.J., Price, A.L., and Neale, B.M. (2015). LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. Nat. Genet. *47*, 291–295.

6. Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.-R., Anttila, V., Xu, H., Zang, C., Farh, K., et al. (2015). Partitioning heritability by functional annotation using genome-wide association summary statistics. Nat. Genet. *47*, 1228–1235.

7. Bulik-Sullivan, B., Finucane, H.K., Anttila, V., Gusev, A., Day, F.R., Loh, P.-R., ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, Duncan, L., et al. (2015). An atlas of genetic correlations across human diseases and traits. Nat. Genet. *47*, 1236–1241.

8. Gazal, S., Finucane, H.K., Furlotte, N.A., Loh, P.-R., Palamara, P.F., Liu, X., Schoech, A., Bulik-Sullivan, B., Neale, B.M., Gusev, A., et al. (2017). Linkage disequilibrium-dependent architecture of human complex traits shows action of negative selection. Nat. Genet. *49*, 1421–1427.

9. Pasaniuc, B., and Price, A.L. (2017). Dissecting the genetics of complex traits using summary association statistics. Nat. Rev. Genet. *18*, 117–127.

10. Yang, J., Benyamin, B., McEvoy, B.P., Gordon, S., Henders, A.K., Nyholt, D.R., Madden, P.A., Heath, A.C., Martin, N.G., Montgomery, G.W., et al. (2010). Common SNPs explain a large proportion of the heritability for human height. Nat. Genet. *42*, 565–569.

11. Speed, D., Cai, N., UCLEB Consortium, Johnson, M.R., Nejentsev, S., and Balding, D.J. (2017). Reevaluation of SNP heritability in complex human traits. Nat. Genet. *49*, 986–992.

12. Gazal, S., Marquez-Luna, C., Finucane, H.K., and Price, A.L. Reconciling S-LDSC and LDAK models and functional enrichment estimates.

13. Finucane, H.K., Reshef, Y.A., Anttila, V., Slowikowski, K., Gusev, A., Byrnes, A., Gazal, S.,
Loh, P.-R., Lareau, C., Shoresh, N., et al. (2018). Heritability enrichment of specifically
expressed genes identifies disease-relevant tissues and cell types. Nat. Genet. *50*, 621–629.

14. Guo, J., Yang, J., and Visscher, P.M. (2018). Leveraging GWAS for complex traits to detect
signatures of natural selection in humans. Curr. Opin. Genet. Dev. *53*, 9–14.

15. Speed, D., and Balding, D.J. (2019). SumHer better estimates the SNP heritability of
complex traits from summary statistics. Nat. Genet. *51*, 277–284.

16. Homburger, J.R., Moreno-Estrada, A., Gignoux, C.R., Nelson, D., Sanchez, E., Ortiz-Tello,
P., Pons-Estel, B.A., Acevedo-Vasquez, E., Miranda, P., Langefeld, C.D., et al. (2015). Genomic
insights into the ancestry and demographic history of South America. PLoS Genet. *11*,
e1005602.

17. SIGMA Type 2 Diabetes Consortium, Williams, A.L., Jacobs, S.B.R., Moreno-Macías, H.,
Huerta-Chagoya, A., Churchhouse, C., Márquez-Luna, C., García-Ortíz, H., Gómez-Vázquez,
M.J., Burtt, N.P., et al. (2014). Sequence variants in SLC16A11 are a common risk factor for
type 2 diabetes in Mexico. Nature *506*, 97–101.

18. Zaitlen, N., Pasaniuc, B., Sankararaman, S., Bhatia, G., Zhang, J., Gusev, A., Young, T.,
Tandon, A., Pollack, S., Vilhjálmsson, B.J., et al. (2014). Leveraging population admixture to
characterize the heritability of complex traits. Nat. Genet. *46*, 1356–1362.

19. Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a tool for
genome-wide complex trait analysis. Am. J. Hum. Genet. *88*, 76–82.

20. Kelleher, J., Etheridge, A.M., and McVean, G. (2016). Efficient Coalescent Simulation and
Genealogical Analysis for Large Sample Sizes. PLoS Comput. Biol. *12*, e1004842.

21. International HapMap Consortium (2003). The International HapMap Project. Nature *426*,
789–796.

22. Gutenkunst, R.N., Hernandez, R.D., Williamson, S.H., and Bustamante, C.D. (2009).
Inferring the joint demographic history of multiple populations from multidimensional SNP
frequency data. PLoS Genet. *5*, e1000695.

23. 1000 Genomes Project Consortium, Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P.,
Kang, H.M., Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., et al. (2015). A global
reference for human genetic variation. Nature *526*, 68–74.

24. Manichaikul, A., Mychaleckyj, J.C., Rich, S.S., Daly, K., Sale, M., and Chen, W.-M. (2010).
Robust relationship inference in genome-wide association studies. Bioinformatics *26*,
2867–2873.

25. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., and Lee, J.J. (2015).
Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience *4*,
7.

26. Durand, E.Y., Do, C.B., Mountain, J.L., and Michael Macpherson, J. (2014). Ancestry

Composition: A Novel, Efficient Pipeline for Ancestry Deconvolution.

27. UK10K Consortium, Walter, K., Min, J.L., Huang, J., Crooks, L., Memari, Y., McCarthy, S.,
Perry, J.R.B., Xu, C., Futema, M., et al. (2015). The UK10K project identifies rare variants in
health and disease. Nature *526*, 82–90.

28. Henn, B.M., Hon, L., Macpherson, J.M., Eriksson, N., Saxonov, S., Pe'er, I., and Mountain,
J.L. (2012). Cryptic distant relatives are common in both isolated and cosmopolitan genetic
samples. PLoS One *7*, e34267.

29. O'Connell, J., Gurdasani, D., Delaneau, O., Pirastu, N., Ulivi, S., Cocca, M., Traglia, M.,
Huang, J., Huffman, J.E., Rudan, I., et al. (2014). A general approach for haplotype phasing
across the full spectrum of relatedness. PLoS Genet. *10*, e1004234.

30. Howie, B., Fuchsberger, C., Stephens, M., Marchini, J., and Abecasis, G.R. (2012). Fast and
accurate genotype imputation in genome-wide association studies through pre-phasing. Nat.
Genet. *44*, 955–959.

31. Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based estimation of
ancestry in unrelated individuals. Genome Res. *19*, 1655–1664.

32. Ge, T., Chen, C.-Y., Neale, B.M., Sabuncu, M.R., and Smoller, J.W. (2017). Phenome-wide
heritability analysis of the UK Biobank. PLoS Genet. *13*, e1006711.

33. Xue, A., Wu, Y., Zhu, Z., Zhang, F., Kemper, K.E., Zheng, Z., Yengo, L., Lloyd-Jones, L.R.,
Sidorenko, J., Wu, Y., et al. (2018). Genome-wide association analyses identify 143 risk variants
and putative regulatory mechanisms for type 2 diabetes. Nat. Commun. *9*, 2941.

34. Thornton, T., Tang, H., Hoffmann, T.J., Ochs-Balcom, H.M., Caan, B.J., and Risch, N.
(2012). Estimating Kinship in Admixed Populations. Am. J. Hum. Genet. *91*, 122–138.

35. Demerath, E.W., Liu, C.-T., Franceschini, N., Chen, G., Palmer, J.R., Smith, E.N., Chen,
C.T.L., Ambrosone, C.B., Arnold, A.M., Bandera, E.V., et al. (2013). Genome-wide association
study of age at menarche in African-American women. Hum. Mol. Genet. *22*, 3329–3346.

36. Fernández-Rhodes, L., Malinowski, J.R., Wang, Y., Tao, R., Pankratz, N., Jeff, J.M.,
Yoneyama, S., Carty, C.L., Setiawan, V.W., Le Marchand, L., et al. (2018). The genetic
underpinnings of variation in ages at menarche and natural menopause among women from the
multi-ethnic Population Architecture using Genomics and Epidemiology (PAGE) Study: A
trans-ethnic meta-analysis. PLoS One *13*, e0200486.

37. Canoy, D., Beral, V., Balkwill, A., Wright, F.L., Kroll, M.E., Reeves, G.K., Green, J., Cairns,
B.J., and Million Women Study Collaborators* (2015). Age at menarche and risks of coronary
heart and other vascular diseases in a large UK cohort. Circulation *131*, 237–244.

38. Bodicoat, D.H., Schoemaker, M.J., Jones, M.E., McFadden, E., Griffin, J., Ashworth, A., and
Swerdlow, A.J. (2014). Timing of pubertal stages and breast cancer risk: the Breakthrough
Generations Study. Breast Cancer Res. *16*, R18.

39. Taki, Y., Kinomura, S., Sato, K., Inoue, K., Goto, R., Okada, K., Uchida, S., Kawashima, R.,

and Fukuda, H. (2008). Relationship Between Body Mass Index and Gray Matter Volume in 1,428 Healthy Individuals. Obesity *16*, 119–124.

40. Berthoud, H.-R., Münzberg, H., and Morrison, C.D. (2017). Blaming the Brain for Obesity: Integration of Hedonic and Homeostatic Mechanisms. Gastroenterology *152*, 1728–1738.

41. Clemmensen, C., Müller, T.D., Woods, S.C., Berthoud, H.-R., Seeley, R.J., and Tschöp, M.H. (2017). Gut-Brain Cross-Talk in Metabolic Control. Cell *168*, 758–774.

42. Guo, M., Liu, Z., Willen, J., Shaw, C.P., Richard, D., Jagoda, E., Doxey, A.C., Hirschhorn, J., and Capellini, T.D. (2017). Epigenetic profiling of growth plate chondrocytes sheds insight into regulatory genetic variation influencing height. Elife *6*,.

43. Wood, A.R., Esko, T., Yang, J., Vedantam, S., Pers, T.H., Gustafsson, S., Chu, A.Y., Estrada, K., Luan, J. 'an, Kutalik, Z., et al. (2014). Defining the role of common variation in the genomic and biological architecture of adult human height. Nat. Genet. *46*, 1173–1186.

44. Villemure, I., and Stokes, I.A.F. (2009). Growth plate mechanics and mechanobiology. A survey of present understanding. J. Biomech. *42*, 1793–1803.

45. Jones, S.E., Lane, J.M., Wood, A.R., van Hees, V.T., Tyrrell, J., Beaumont, R.N., Jeffries, A.R., Dashti, H.S., Hillsdon, M., Ruth, K.S., et al. (2019). Genome-wide association analyses of chronotype in 697,828 individuals provides insights into circadian rhythms. Nat. Commun. *10*, 343.

46. Bering, T., Carstensen, M.B., Wörtwein, G., Weikop, P., and Rath, M.F. (2018). The Circadian Oscillator of the Cerebral Cortex: Molecular, Biochemical and Behavioral Effects of Deleting the Arntl Clock Gene in Cortical Neurons. Cereb. Cortex *28*, 644–657.

47. Gnocchi, D., and Bruscalupi, G. (2017). Circadian Rhythms and Hormonal Homeostasis: Pathophysiological Implications. Biology *6*,.

48. Potter, G.D.M., Skene, D.J., Arendt, J., Cade, J.E., Grant, P.J., and Hardie, L.J. (2016). Circadian Rhythm and Sleep Disruption: Causes, Metabolic Consequences, and Countermeasures. Endocr. Rev. *37*, 584–608.

49. Weissbrod, O., Flint, J., and Rosset, S. (2018). Estimating SNP-Based Heritability and Genetic Correlation in Case-Control Studies Directly and with Summary Statistics. Am. J. Hum. Genet. *103*, 89–99.

50. Speed, D., and Balding, D. (2018). Better estimation of SNP heritability from summary statistics provides a new understanding of the genetic architecture of complex traits.

51. Yang, J., Bakshi, A., Zhu, Z., Hemani, G., Vinkhuyzen, A.A.E., Nolte, I.M., van Vliet-Ostaptchouk, J.V., Snieder, H., Lifelines Cohort Study, Esko, T., et al. (2015). Genome-wide genetic homogeneity between sexes and populations for human height and body mass index. Hum. Mol. Genet. *24*, 7445–7449.

52. Yang, J., Zeng, J., Goddard, M.E., Wray, N.R., and Visscher, P.M. (2017). Concepts, estimation and interpretation of SNP-based heritability. Nat. Genet. *49*, 1304–1310.

53. Taliun, D., Harris, D.N., Kessler, M.D., Carlson, J., Szpiech, Z.A., Torres, R., Gagliano
Taliun, S.A., Corvelo, A., Gogarten, S.M., Kang, H.M., et al. (2019). Sequencing of 53,831
diverse genomes from the NHLBI TOPMed Program.

54. Martin, A.R., Gignoux, C.R., Walters, R.K., Wojcik, G.L., Neale, B.M., Gravel, S., Daly, M.J.,
Bustamante, C.D., and Kenny, E.E. (2017). Human Demographic History Impacts Genetic Risk
Prediction across Diverse Populations. Am. J. Hum. Genet. *100*, 635–649.

55. Nelson, S.C., Stilp, A.M., Papanicolaou, G.J., Taylor, K.D., Rotter, J.I., Thornton, T.A., and
Laurie, C.C. (2016). Improved imputation accuracy in Hispanic/Latino populations with larger
and more diverse reference panels: applications in the Hispanic Community Health Study/Study
of Latinos (HCHS/SOL). Hum. Mol. Genet. *25*, 3245–3254.

56. Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P.,
Green, J., Landray, M., et al. (2015). UK biobank: an open access resource for identifying the
causes of a wide range of complex diseases of middle and old age. PLoS Med. *12*, e1001779.

57. Turley, P., Walters, R.K., Maghzian, O., Okbay, A., Lee, J.J., Fontana, M.A., Nguyen-Viet,
T.A., Wedow, R., Zacher, M., Furlotte, N.A., et al. (2018). Multi-trait analysis of genome-wide
association summary statistics using MTAG. Nat. Genet.

58. Gusev, A., Lee, S.H., Trynka, G., Finucane, H., Vilhjálmsson, B.J., Xu, H., Zang, C., Ripke,
S., Bulik-Sullivan, B., Stahl, E., et al. (2014). Partitioning heritability of regulatory and
cell-type-specific variants across 11 common diseases. Am. J. Hum. Genet. *95*, 535–552.

59. Kichaev, G., and Pasaniuc, B. (2015). Leveraging Functional-Annotation Data in
Trans-ethnic Fine-Mapping Studies. Am. J. Hum. Genet. *97*, 260–271.

60. Kanai, M., Akiyama, M., Takahashi, A., Matoba, N., Momozawa, Y., Ikeda, M., Iwata, N.,
Ikegawa, S., Hirata, M., Matsuda, K., et al. (2018). Genetic analysis of quantitative traits in the
Japanese population links cell types to complex human diseases. Nat. Genet. *50*, 390–400.

61. Matise, T.C., Ambite, J.L., Buyske, S., Carlson, C.S., Cole, S.A., Crawford, D.C., Haiman,
C.A., Heiss, G., Kooperberg, C., Marchand, L.L., et al. (2011). The Next PAGE in understanding
complex traits: design for the analysis of Population Architecture Using Genetics and
Epidemiology (PAGE) Study. Am. J. Epidemiol. *174*, 849–859.

787
788
789
790

Figure and table legends

Figure 1. Overview of the covariate-adjusted LD score regression. (a) As input, cov-LDSC

takes raw genotypes of collected GWAS samples and their global principal components.

(b) cov-LDSC regresses out the ancestral components based on global principal components

from the LD score calculation and corrects for long-range *admixture* LD. Black and red lines

indicate estimates before and after covariate adjustment respectively (c) Adjusted heritability

estimation based on GWAS association statistics (measured by χ^2) and covariate-adjusted LD

scores. (d) Estimation of heritability enrichment in tissue-specific gene sets.

Figure 2. Estimates of heritability (h_g^2) under different simulation scenarios using the SIGMA cohort. LDSC (orange) underestimated h_g^2 and cov-LDSC (blue) yielded robust h_g^2 estimates under all settings. Each boxplot represents the mean LD score estimate from 100 simulated phenotypes using the genotypes of 8,214 unrelated individuals from the SIGMA cohort. For cov-LDSC, a window size of 20-cM with 10 PCs were used in all scenarios. A true polygenic quantitative trait with $h_g^2 = 0.4$ is assumed for scenarios (a), (c) and (d) and 1% causal variants are assumed for scenarios (b)-(d). (a) h_g^2 estimation with varying proportions of causal variants (0.01% - 30%). (b) h_g^2 estimation with varying heritabilities (0, 0.05, 0.1, 0.2, 0.3, 0.4 and 0.5). (c) h_g^2 estimation when an environmental stratification component aligned with the first PC of the genotype data was included in the phenotype simulation. (d) h_g^2 estimation when using a subset of the cohort to obtain LD score estimates and using out-of-sample LD score estimates obtained from Admixed Americans included in the 1000 Genomes Project.

Figure 3. Estimates of heritability (h_g^2) of three quantitative and four dichotomous traits in two admixed populations in the 23andMe research cohort. For seven selected non-disease phenotypes (body mass index (BMI), height, age at menarche, left handedness, morning person, motion sickness and nearsightedness) in the 23andMe cohort, we reported their estimated genetic heritabilities and intercepts (and their standard errors) using the baseline model. LD scores were calculated using 134,999, 161,894, 46,844 individuals from 23andMe European, Latino and African American individuals respectively. For each trait, we reported the sample size in obtained summary statistics used in cov-LDSC. For BMI and height, we also reported the h_g^2 estimates from the SIGMA cohort.

Figure 4. Results of multiple-tissue analysis for height, BMI and morning person. Each point represents a tissue type from either the GTEx data set or the Franke lab data set as defined in Finucane et al[13]. From top to bottom, (a)-(d) show multiple-tissue analysis for BMI in the cross-population meta-analysis and in Europeans, Latinos and African Americans respectively. (e) shows the scatter plot of the estimated per-standardized-annotation effect size τ^* which represents the proportional change of averaged per-SNP heritability by one standard deviation increase in value of the annotation of each cell type, conditional on other 53 non-cell type specific baseline annotations, in the three populations for all tested tissue types (**Methods**). The x-axis shows the τ^* in European populations and the y-axis shows either τ^* in Latinos (blue) or African Americans (orange). We reported the slope and p-value when we regress Latinos (blue) and African Americans (organe) τ^* on Europeans τ^* for all tissue types. Error bars indicate standard errors of τ^*. Similarly, the results are shown in (f)-(j) for height and (k)-(n) for morning person. The significance threshold in plots (a)-(d), (f-i) and (k-m) is defined by the FDR < 5% cutoff, $-log_{10}(p) = 2.75$. Numerical results are reported in **Supplementary Table 14**.

Table 1. $h_g{}^2$ **estimates of height, BMI and type 2 diabetes using different heritability estimation methods.** Reported values are estimates of $h_g{}^2$ (with standard deviations in brackets) from LDSC using a 20-cM window, cov-LDSC using a 20-cM window and 10 PCs, and GCTA using REAP to obtain the genetic relationship matrix with adjustment by 10 PCs. The final column provides reported $h_g{}^2$ estimates in European populations from various studies[10,32,33].

Phenotype	LDSC (baseline)	cov-LDSC (baseline)	GCTA (REAP w/ 10pc)	Public
Height	0.159 (0.037)	0.379 (0.079)	0.450 (0.042)	0.45-0.685[10,32]
BMI	0.113 (0.030)	0.248 (0.061)	0.235 (0.041)	0.246-0.27[32]
T2D	0.121 (0.035)	0.263 (0.073)	0.376 (0.046)	0.139-0.414[32,33]

801