A Novel Uranium Bioremediation Approach using Radioresistance *Deinocoocus*

***radiodurans* Biofilm**

T. Manobala[1], Sudhir K. Shukla[2], T. Subba Rao[2*] and M. Dharmendira Kumar[1]

[1]Department of Applied Science and Technology, Anna University, Chennai,

Tamil Nadu 600 025, India

[2]Biofouling and Thermal Ecology Section, Water and Steam Chemistry Division

BARC Facilities, Kalpakkam, Tamil Nadu 603 102, India.

*Corresponding author Email: subbarao@igcar.gov.in

Abstract

Deinococcus radiodurans, the most radiation resistant organism ever known, has

gained importance in recent years as a potential candidate for bioremediation of heavy

metals, especially radioactive ones. This study investigates the efficiency of a recombinant *D.*

radiodurans (DR1-bf[+]) strain with an ability to form biofilm for uranium remediation. In this

study, a modified Arsenazo III dye method was used to estimate the uranium concentration.

Uranyl nitrate aqueous solution, wherein U exists as uranyl ion (UO_2^{+2}), is generated during

the operation of nuclear fuel reprocessing. The *D. radiodurans* biofilm (DR1-bf[+]) grown in

the presence of 20 mM Ca^{2+} ion showed phenomenal ability of uranyl ion removal. DR1-bf[+]

(+Ca^{2+}) biofilm removed ~75 ± 2% of 1000 mg/L uranium within 30 minutes post treatment

from uranyl nitrate aqueous solution. U removal rate was also found to be directly

proportional to biofilm age. This study discusses the extraordinary ability of *D. radiodurans*

biofilm in uranium removal.

Key words: *Deinococcus radiodurans*, radio-tolerance, biofilm, Uranium, Arsenazo,

bioremediation

Graphical abstract

1. Introduction

In a natural environment, almost 99% of all microorganisms live as surface-attached

communities known as biofilms (Costerton et al. (1987). A biofilm is an assemblage of

microbial communities, in which the microbial cells are irreversibly attached to a surface and

embedded in an extracellular polymeric matrix (EPS) produced by these cells (Donlan, 2002;

Das et al., 2012). Biofilms apart from enhancing the organism survival, improve the rate and

extent of contaminant transformation as compared to that of pure and planktonic cultures

(Mangwani et al., 2014). Remediation using microorganism (termed as bioremediation) is

emerging *in situ* technology for the clean-up of environmental pollutants. The economical

factor and inefficiency of some physico-chemical remediation methods has made this

biological treatment method, an improved and preferred alternative (Paul et al., 2005).

Heavy metals are released into the environment through various processes, leading to

contamination of air, water, and atmosphere (Akpor and Muchie, 2010). These heavy metals

need to be converted from one form to less hazardous form or its bioavailability should be

decreased (Wall and Krumholz, 2006). Among the many heavy metals, naturally occurring

and comparatively less abundant uranium has drawn attention due to the surge in

anthropogenic activities involving uranium and its potential hazards due to an accidental

release into the environment. The use of uranium as fuel in the nuclear reactors has increased

its worldwide demand. Radioactive uranyl nitrate ($UO_2(NO_3)_2.6H_2O$) is generated during the

operation of nuclear fuel reprocessing (Baumgärtner, 2009). In aqueous medium, uranium

exists as uranyl ion (UO_2^{2+}) i.e. UO_2^{+2} is highly soluble and mobile in the subsurface

environments and could be hazardous being a heavy metal and radioactive (Shukla and Rao,

2015). The intake of these ions can cause detrimental health hazards (Keith et al., 2013). The

chemical methods used for uranium removal have their own limitations due to very high cost

and low feasibility for *in situ* bioremediation (Kulkarni et al., 2013).

Bioremediation of uranium using microbes can be a very promising and more

efficient approach as microbes are nature's creative recyclers (N'Guessan et al., 2008).

Biofilms are ideal for bioremediation purpose as they need not to be separated from the bulk

liquid waste, thus making the whole process more economical and feasible (Shukla et al.,

2017). Biofilm mediated bioremediation of UO_2^{2+} would be an ideal process by a radio-

tolerant microbial biofilm to ease its further downstream processing. Also conversion of

aqueous U to insoluble uranyl phosphate precipitates are not susceptible for reoxidation in

contrast to the 'reduced' uranium mineral like uraninite which has the tendency to reoxidize

back to more soluble U (Kulkarni et al., 2013). In such cases, inherently radio-tolerant

bacterium *Deinococcus radiodurans* can be of use. *Deinococcus radiodurans* has an

extraordinary capability of surviving under high radiation stress (Battista, 1997), even under

low nutrient conditions (Shukla et al., 2014). The bioremediation potential of *D. radiodurans*

has gained importance in recent times. Engineered *D. radiodurans* cells have been used to

detoxify mercury, degrade toluene (Brim et al., 2000) and chromium reduction (Brim et al.,

2006). However, use of planktonic cells for bioremediation of heavy metals makes the

downstream process costly or less efficient.

A genetically modified strain of *D. radiodurans,* harboured with a constitutively expressing *gfp* plasmid (GenBank accession no. KF975402), was used in this study and found to be a biofilm forming strain (Shukla and Rao, 2017a). We have investigated whether this biofilm forming strain of *D. radiodurans* (hereafter denoted as DR-bf$^+$) can be used for the biofilm- mediated uranium bioremediation from the uranyl nitrate aqueous solution.

2. Materials and Methods

2.1 Microorganism and culture conditions

Deinococcus radiodurans R1 wild type (DR1-WT) and *gfp*-harbouring, kanamycin resistant, biofilm forming strain of *D. radiodurans* (DR1-bf$^+$) was used in this study. TGY medium consisting of tryptone (5 g/L), yeast extract (3 g/L), glucose (1 g/L) was used as growth medium. The DR1-bf$^+$ bacterium was grown for 48 h in TGY medium at 30^{o}C in an orbital shaker at 100 rpm until mid-log phase. Kanamycin antibiotic was added in the final concentration of 5µg/mL for the growth of *D. radiodurans* (DR1-bf$^+$). For most of the experiments mid-log phase cultures were used.

2.2. Biofilm estimation using crystal violet assay

Biofilm assay was performed in 96 well microtiter plates as per the modified classical crystal violet assay (Shukla and Rao, 2017b). Late log phase *D. radiodurans* culture (grown for 48 h) were diluted in sterile TGY medium in 1:40 ratio and 200 µl of this diluted culture was transferred to the pre-sterilized 96 well flat bottom polystyrene microtiter plates. After the prescribed time as per the experimental plan, planktonic cells were aspirated and the biofilm grown in 96 well plate was washed twice with 200 µl of sterile phosphate buffer saline (PBS), air dried and stained with 0.2% crystal violet solution. For dissolution of bound crystal violet, 95% ethanol was used. Biofilm growth was estimated in terms of absorbance at 570 nm using a multimode micro-plate reader (BioTek, USA).

2.3 Fluorescence microscopy study

The biofilm samples were grown in petri dishes for fluorescence microscopy study. The cells were grown in TGY media at 30°C and 150 rpm in 15 mL falcon tubes. 100 µl mid log phase planktonic cells of DR1-bf$^+$ (with OD of 0.7) were transferred to petri dishes (35 mm) with 4 ml of TGY media in the presence of calcium (20 mM) and required amount of kanamycin antibiotic with a final concentration of 5 µg/mL. The used up medium was aspirated and the biofilm was rinsed with sterile PBS to remove loosely attached cells. The petri plate was viewed under a fluorescence microscope (Carl Zeiss Axioscope I, Germany) equipped with a fluorescence illumination system and with appropriate filter sets. Images were obtained from randomly selected fields using 20X objective.

2.4 Biofilm growth of DR1-bf$^+$ for uranyl nitrate treatment

The biofilm forming strain of *D. radiodurans* (DR1-bf$^+$) was used in this biofilm-mediated bioremediation study (Shukla and Rao, 2015). The cells were grown in TGY media at 30°C and 150 rpm in 15 mL falcon tubes. 100 µl mid log phase planktonic cells of DR1-bf$^+$ (with OD of 0.7) were transferred to petri dishes (35 mm) with 4 ml of TGY media (with or without 20 mM Ca^{2+}) and kanamycin with a final concentration of 5 µg/mL. These plates were incubated in the static condition at 30°C for the required period of time and the used up media was replaced after every 48 h. After given time period, the planktonic cells were aspirated and biofilm was gently washed twice with PBS, before the treatment with filter sterilized uranyl nitrate solution (pH 3.61).

2.5 Uranium estimation protocol Arsenazo III dye method and by Inductively Coupled Plasma-Atomic Emission Spectroscopy (ICP-AES)

Uranium in the aqueous sample was estimated by modified protocol of Arsenazo III method. In this study, the standard arsenazo dye method (Savvin, 1961) was modified for the uranium estimation. Briefly this method involves simple treatment of the uranium solution

with a buffer Potassium Hydrogen Pthalate buffer (1.5 mM PHP in 0.025 N HCl) and 0.1%

Arsenazo III dye. To 1 ml of PHP buffer, 900 µl of the sample and 100 µl of Arsenazo III

reagent was added. It was mixed by vortexing and absorbance was measured at 653 nm

(absorption maxima for the U-Arsenazo III complex) after 5 minutes. The change in the color

is directly proportional to the amount of uranium present in the given solution. Suitable

dilution factors and arsenazo dye concentration were used to estimate uranium in wide range.

The samples were also analyzed for uranium estimation using Inductively Coupled Plasma-

Atomic Emission Spectroscopy (ICP-AES, model: Ultima-2, Horiba JovinYvon, France) to

substantiate the modified Arsenazo III dye method. Two ml of uranyl nitrate sample was

collected and taken for ICP-AES analysis. Standard graph was plotted for both modified

Arsenazo III dye method and ICP-AES to validate the Arsenazo III dye method (Savvin,

1961).

2.6 Estimation of uranium removal

Uranyl nitrate [UO_2 $(NO_3)_2.6H_2O$] solution (1000 mg/L) was prepared in ultrapure

water and filter sterilized. The pH of the aqueous solution was 3.61. DR1-bf$^+$ biofilms were

grown as described above [section 2.4] for the prescribed time of growth as per the

experiment plan. TGY medium along with planktonic cells was aspirated from the petri dish

and discarded. Biofilm was gently washed twice with sterile PBS. Only TGY medium with

Ca^{2+} and kanamycin (excluding bacterial cells) was also included as blank. Washed biofilms

were treated with of 1000 mg/L of Uranyl nitrate solution. Sample was collected at different

time intervals up to 2 hours for the estimation of uranium as per above described method.

2.7 Scanning Electron Microscopy (SEM)

DR1_bf$^+$ biofilm was grown as explained in the section 2.1 in the slides without

agitation. Biofilm were fixed with 2.5% glutaraldehyde (in PBS, pH 7.2) at 4°C overnight.

The biofilm was scrapped out, washed with PBS thrice and dehydrated with a series of

ethanol solutions (10, 20, 30, 40, 50, 60, 70, 80, 90 and 100 % w/v, with 10 minutes

incubations in each). The slides were coated with gold and were analyzed using scanning

electron microscopy (HELIOS SEG-SEM). Analysis of SEM images revealed the presence of

dense biomass embedded in EPS matrix.

2.8 Fourier transform infrared spectroscopy (FTIR)

147 The 4 day old DR1_bf$^+$ biofilm was grown as explained in (section 2.4) and scrapped

148 out (before and after treating with uranium) in to an eppendroff. The biomass was lyophilised

149 and taken for FTIR analysis. The infrared spectra of lyophilised bacterial biomass samples

150 (before and after uranium treatment) were recorded within a range 400–4,000 cm^{-1} in an ABB

151 MB3000 FTIR Spectrometer. The samples were prepared as KBr pellets.**3. Results**

3.1 Standardisation of uranium estimation protocol

Figure 1 shows that this method gave a linear correlation up to 20 mg/L concentration

of uranium using modified method of uranium estimation with arsenazo III with 0.01%

arsenazo II dye solution using potassium hydrogen phthalate buffer. The concentration

estimates derived by this method were cross checked for its accuracy by using ICP-AES

analysis. The concentrations of uranium estimated using arsenazo III method gave linear

relationship in the range of 1-20 mg/L of uranium. By changing the concentration of arsenazo

III dye the concentration estimated can go up to 1000 mg/L. Uranium estimation using ICP-

AES analysis values corroborated with the values obtained with modified arsenazo III

method proving the efficiency and accuracy of the modified arsenazo III protocol. Therefore,

this modified method was used for uranium estimation in this study.

3.2 Biofilm growth by *D. radiodurans* R1 (DR1-bf$^+$) strain

The crystal violet assay showed that the recombinant *D. radiodurans* (DR1-bf$^+$) has a

very good ability to form biofilm in contrast to its wild type counterpart (Figure 2a).

Fluorescence microscopic study also confirmed the biofilm formation (Figure 2b). The

maximum biofilm formation was observed at 4 days and calcium in the growth medium

enhanced the biofilm production in DR1-bf$^+$ proportionally. This shows the effect of Ca^{2+} on

EPS production and relative increase in thickness of biofilm (Shukla and Rao, 2015). 20 mM

concentration of Ca^{2+} was shown to have optimum effect on the extracellular matrix

production of DR1-bf$^+$ biofilm compared to other concentrations of calcium ion (Shukla and

Rao, 2017a). Therefore this concentration of Ca^{2+} was used for rest of the experiments.

3.3 Bioremediation of uranium using DR1-bf$^+$ biofilm and planktonic cells of DR1-WT

and DR1-bf$^+$

The biofilm was grown for 4 days and treated with aqueous uranium solution as

described in the sections 2.4 and 2.6. A control biofilm grown in the absence of Ca^{2+} was also

evaluated for comparison. Similarly the planktonic cells, both of *D. radiodurans* R1 wild type

(DR1-WT) and *D. radiodurans* (DR1-bf$^+$) strains were grown in test tubes, maintaining the

identical growth conditions but in the absence of Ca^{2+}. The assay was performed after 4 days

of biofilm growth. Required amount of sample was collected at different time intervals and

the reduction in uranium concentration was estimated. Results showed that the uranium

remediation by DR1-bf$^+$ biofilm in the presence of Ca^{2+} was very quick in the first 5 min post

treatment, which removed nearly 25% of initial concentration of uranium (1000 mg/L).

Beyond which the removal rate slowly decreased and reached a maximum of 75 ± 2%

removal within 30 min of the treatment. With increase in time, further removal of uranium

was not observed, which might be due to saturation of its adsorption capacity of the biofilm.

As shown in the Figure 3A (by Arsenazo dye method) and Figure 3B (by ICP-AES)

removal of uranyl ion by DR1-bf$^+$ biofilm when grown in the presence of Ca^{2+}, was

significantly high as compared to DR1 planktonic cells, DR1-bf$^+$ planktonic cell or DR1-bf$^+$

control biofilm (grown without Ca^{2+}). Results showed that the removal efficiency of DR1-bf+

(+ Ca^{2+}) biofilm was four times more as compare to DR1-bf+ planktonic cells. A visible

yellow precipitate of uranyl nitrate by the biofilm was observed after the remediation process

(Figure 3a) that indicates the conversion of aqueous form of U(VI) to an insoluble precipitate

U(IV), making the removal of uranium more competent than other methods. Although initial

experiments showed that the mechanism of uranium bioremediation appears to be an

adsorption phenomenon, the possibility of enzyme mediated process may not be ruled out, at

this stage of study.

3.4 The effect of biofilm age on uranium bioremediation

To study the effect of biofilm's age on the efficiency of uranium removal, a day-wise

U removal assay was performed. Results showed that there was a continuous increase in

uranium removal efficiency as the DR1-bf$^+$ (+Ca^{2+}) biofilm grew older (see Figure 4). The

removal rate was found to be higher after 4 days of biofilm growth. Without much increase in

biofilm biomass, this suggests that there is a continuous change in the biofilm matrix

composition over a period of time (Table 1).

Biofilm's age	Biofilm biomass [mg]
Day 1	29.24
Day 2	29.97
Day 3	29.43
Day 4	27.38
Day 5	29.38
Day 6	28.03
Day 7	32.43

Table 1: Table showing the amount of biomass produced with the increse in the age of the

bioiflm. These biofilm were grown as described in section 2.4. The biofilm was scrapped and

dried to get the weight.

Our previous studies showed that there was an enhanced exopolysaccharide and

eDNA production in the biofilm matrix in the presence of Ca^{2+} (Shukla and Rao, 2014;

Shukla and Rao, 2015). In view of this, it is speculated that uranyl ion (UO_2^{2+}) removal is by

biosorption process where negatively charged exopolysaccharide and eDNA play a critical

role. However, visible precipitation and faster removal of UO_2^{2+} by older biofilm also suggest

that this phenomenon can also be aided by enzymatic reduction.

3.5 Adsorption is the major method of uranium sequestration

FTIR spectroscopic study for the control biofilm (uranium-free) and the uranium

treated biomass were performed to elucidate the chemical groups involved in uranium

binding onto the bacterial biomass (Figure 6). The highly complex IR spectra was analysed to

assign the important characteristic peaks which illustrates the involvement of main functional

groups present in the bacterial biomass. The amino group (N–H stretching) peak lies in the

spectrum region occupied by a broad and strong band (3,600–3,000 cm^{-1}). This may be due to

hydroxyl groups that are hydrogen bonded to various degrees (Pagnanelli et al., 2000). In the

control spectrum, absorption peaks at 2,900–3,000 cm^{-1} are ascribed to the asymmetric

stretching of C–H bond of the –CH_2 groups combined with that of the CH_3 groups. In the

control spectrum, the $\gamma C=O$ of amide I and $\delta NH/\gamma C=O$ combination of the amide II bond

were present at 1,647 and 1,546 cm^{-1}, respectively, indicating the presence of carboxyl

groups (Kazy et al., 2006). In the uranium loaded samples, these peaks were shifted from

1,647 cm^{-1} to 1,650 cm^{-1} and 1,546 cm^{-1} to 1,540 cm^{-1} region, indicating an of interaction

between uranium and carboxyl group. Especially, 1540 cm^{-1} peak indicates the scissoring of

CH_2 group. Sharp spectra from 1456-1481 cm^{-1} gesture of carboxyl group. A decrease in the

intensity and a gradual shift of the peak at 1214 cm^{-1}, in the control sample to a lower energy

in uranium (1204 cm^{-1}) loaded samples clearly indicates the weakening of the P = O

character as a result of radionuclide binding to the phosphates. A sharp peak at 1387 cm^{-1} is

indicative of symmetrical bending of CH_3 group (Stuart, 2005). In uranium loaded sample,

distinct peak at 927 cm^{-1} and changes in peak positions and intensity around 1000 cm^{-1} region

can be assigned to asymmetric stretching vibrations of $\gamma 3\ UO_2^{2+}$ and stretching vibrations of

weekly bounded oxygen ligands with uranium (γ U-O ligand) (Kazy et al., 2009). The overall

spectral analysis strongly supports the major role of carboxyl, amide and phosphate groups in

uranium binding by the bacterial biomass.

4. Discussion

The present study involved the use of a recombinant *D. radiodurans* biofilm for the

bioremediation of uranium from aqueous solution. Although planktonic forms of these

species have been used for some treatment process, their drawbacks restrict them to be used

for a wider application. For example, earlier studies involving U remediation using *D.*

radiodurans, showed good removal efficiency up to 95% removal when treated with

uranium concentration ranging from 5-100 μM, but with an incubation period of 21 days,

which greatly affected the efficiency of the system (Fredrickson et al., 2000). Though in few

studies involving few bacterial cultures such as *Desulfosporosinus orientis* DSM 765,

Desulfosporosinus spp. P3 (Suzuki et al., 2004), *Clostridium sp.* (Francis et al., 1994) and

Clostridium sphenoides ATCC 19403 (Francis et al., 2004) have been shown a significant

uranium removal under anaerobic environment, but maintaining the anaerobic condition is

another challenging task and a major bottle neck while scaling up the bioremediation process

at industrial scale. (Marsili et al., 2005) reported the use of sulphate reducing bacterial

biofilm for uranium removal with preformed biofilm grown for 3 weeks in the reactor. The

efficiency of removal was estimated to be 30.4% in the reactor supplied with 3 mg/L of U

and 73.9% in the reactor supplied with 30 mg/L of U. The reactor performance was not up to

the expectation, which was hypothetically ascribed to the differences in biofilm sloughing

events in the reactor. Although these reports were evident of uranium removal process by

bacteria, none of the systems has been proved to be efficient enough for the large-scale

applications. Application of such bacteria in U bioremediation (low active waste) will be

further difficult as they do not have the capability to survive in the presence of radioactivity.

Deinococcus radiodurans is known for its extreme radiation tolerance (Krisko and

Radman, 2013). *D. radiodurans* does not have innate ability to form biofilm. Although

Deinococcus geothermalis (Kolari et al., 2002) has shown the ability to form biofilm, no

study has indicated its potential application in bioremediation. This study involves the use of

a genetically modified *D. radiodurans biofilm* for uranium bioremediation purpose. The

genetically modified strain of *D. radiodurans* (DR1-bf$^+$), when grown in the presence of 20

mM Ca^{2+} was found to produce very good biofilm thus DR1-bf$^+$ biofilm, grown in the

presence Ca^{2+} was tested for the uranium bioremediation in this study. Ca^{2+} plays an

important role in the adhesion of bacterial cells to the surfaces. They are involved both in

specific (actions which cannot be replaced by other cations) and non-specific interactions

(between cells and substratum) in biofilms (Shukla and Rao, 2013; Mangwani et al., 2014).

Also, the interaction of protein and polysaccharide at the cell surface is influenced by Ca^{2+}

(Geesey et al., 2000). In the biofilm, extracellular Ca^{2+} ions involve in the cross-linking of

extracellular polymeric substances, which   gives strength to the biofilm matrix and regulates

production of many extracellular enzymes. It also plays an important role in maintaining the

integrity of the biofilm (Shukla and Rao, 2014). In a way, Ca^{2+} acts as an important ionic

cross-bridging molecule for negatively charged bacterial polysaccharides. It was also shown

in our previous study that production of extracellular matrix was enhanced in *D. radiodurans*

biofilm when biofilm was grown in the presence of Ca^{2+} in the growth medium as compared

to that of biofilm grown in the absence of Ca^{2+} (Shukla and Rao, 2015). Among the various

concentrations of Ca^{2+} used in the study, 20 mM   showed a significant improvement in the

biofilm production, as compared to other concentrations (Shukla and Rao, 2015), therefore,

20 mM concentration was chosen in this study. It is possible that the Ca^{2+} can influence the

D. radiodurans biofilm growth in multiple ways but as of now, the exact mechanism is not

clear. The involvement of biosorption by negatively charged exopolysaccharide and eDNA in

the biofilm matrix of *D. radiodurans* has been speculated to be the reason for this

remediation process also the visible precipitate formation and detection of acid phosphatase

activity cannot rule out the possibility of enzyme-mediated reduction of U(VI) to U(IV)

(Francis et al.).

In this study, our results showed that 4 day old DR1-bf$^+$ biofilm had significant

capability to remove UO_2^{+2} ions that too at a very high rate which was ~75% removal within

30 mins (Figure 3A). This observation implicates the potential of *D. radiodurans* for the

development of biofilm-based bioremediation process for UO_2^{2+} removal from radioactive

aqueous waste solutions. The biofilm mode of *D. radiodurans* has shown tolerance to high

concentrations of uranium, up to 1000 mg/L. The DR1-bf$^+$ biofilm could withstand very high

concentrations of uranyl nitrate solution i.e. 1000 mg/L not only due to the fact that *D.*

radiodurans has significant heavy metal tolerance (Brim et al., 2000) but also of the fact that

biofilm mode of life reduces the toxicity of heavy metals to microbes as compared to that of

planktonic cells. This DR1-bf$^+$ biofilm-mediated uranium removal method showed significant

higher efficiency in terms of both performance as well as time (75% uranium removal within

30 min post-treatment) as compared to its planktonic counterparts as well as *D. radiodurans*

wild type strain. No requirement of special conditions such as maintaining an anaerobic

condition in this removal process and tolerance to high concentrations of uranium make *D.*

radiodurans biofilm-mediated U removal process highly promising. Further investigations

into the form of uranium which is precipitated and the mechanisms involved in the microbial

reduction of this U has to be evaluated, as such studies may provide insights into strategies

for promoting the microbial reduction in uranium-contaminated sites.

5. Conclusion:

Deinococcus radiodurans (DR1-bf[+]) biofilm was tested for its potential to remove

uranium from aqueous solution. The study showed that when DR1-bf[+] is grown in the

presence of 20 mM Ca^{2+}, it acquired an extraordinary capacity of UO_2^{+2} removal as compared

to that of its planktonic counterparts and control biofilm. In this study, it was shown that 4

days old DR1-bf[+] biofilm when grown in the presence of 20 mM Ca^{2+}, significantly removes

UO_2^{+2} ions (~75% removal) within few minutes. There was an increase in the uranium

removal capacity with increase in the age of the biofilm. Interestingly there was not much

variation in the biofilm biomass with the age. This speculated that uranyl ion (UO_2^{2+})

removal is a biosorption process where negatively charged exopolysaccharide and eDNA

play a critical role. Without significant change in the biofilm biomass, age-dependent

increase in U bioremediation was observed. FTIR analysis strongly supports the involvement

of carboxyl, amide and phosphate groups in uranium binding by the bacterial biomass. This

observation implicates the potential of *D. radiodurans* biofilm-based bioremediation process

for UO_2^{2+} precipitation.

6. References:

Akpor O, Muchie M. 2010. Remediation of heavy metals in drinking water and wastewater
treatment systems: Processes and applications. Int J Phy Sci 5:1807-1817.
Battista JR. 1997. Against all odds: the survival strategies of *Deinococcus radiodurans*. Annu
Rev Microbiol. 51:203-224.
Baumgärtner M. 2009. The use of reprocessed uranium in light water reactors: Problem
identification and solution finding. Report no. IAEA-TECDOC-CD--1630
Brim H, McFarlan SC, Fredrickson JK, Minton KW, Zhai M, Wackett LP, Daly MJ. 2000.
Engineering *Deinococcus radiodurans* for metal remediation in radioactive mixed waste
environments. Nat Biotechnol 18:85-90.

Brim H, Osborne JP, Kostandarithes HM, Fredrickson JK, Wackett LP, Daly MJ. 2006.
Deinococcus radiodurans engineered for complete toluene degradation facilitates Cr
(VI) reduction. Microbiology 152:2469-2477.

Costerton JW, Cheng K, Geesey GG, Ladd TI, Nickel JC, Dasgupta M, Marrie TJ. 1987.
Bacterial biofilms in nature and disease. Annu Rev Microbiol. 41:435-464.

Das N, Geetanjali Basak L, Salam JA, Abigail M. 2012. Application of biofilms on
remediation of pollutants–an overview. J Microbiol Biotechnol Res 2:783-790.

Donlan RM. 2002. Biofilms: microbial life on surfaces. Emerg Infect Dis 8.

Francis A, Dodge C, Gillow J, Halada G, Honeyman B. 2004. Reductive precipitation and
stabilization of uranium complexed with organic ligands by anaerobic bacteria. DOE-
NABIR PI Workshop.

Francis AJ, Dodge CJ, Lu F, Halada GP, Clayton CR. 1994. XPS and XANES studies of
uranium reduction by *Clostridium* sp. Environ Sci Technol 28.

Fredrickson JK, Kostandarithes HM, Li S, Plymale AE, Daly M. 2000. Reduction of fe (III),
cr (VI), U (VI), and tc (VII) by *Deinococcus radiodurans* R1. Appl Environ Microbiol
66:2006-2011.

Geesey GG, Wigglesworth-Cooksey B, Cooksey K. 2000. Influence of calcium and other
cations on surface adhesion of bacteria and diatoms: a review. Biofouling 15:195-205.

Kazy SK, D'souza S, Sar P. 2009. Uranium and thorium sequestration by a Pseudomonas sp.:
mechanism and chemical characterization. J Hazard Mater 163:65-72.

Kazy SK, Das SK, Sar P. 2006. Lanthanum biosorption by a Pseudomonas sp.: equilibrium
studies and chemical characterization. J Ind Microbiol Biotechnol.33:773-783.

Keith S, Faroon O, Roney N, Scinicariello F, Wilbur S, Ingerman L, Llados F, Plewak D,
Wohlers D, Diamond G. 2013. Toxicological profile for uranium. Agency for Toxic
Substances and Disease Registry (US), Atlanta (GA).

Kolari M, Schmidt U, Kuismanen E, Salkinoja-Salonen M. 2002. Firm but slippery
attachment of *Deinococcus geothermalis*. J Bacteriol 184:2473-2480.

Krisko A, Radman M. 2013. Biology of extreme radiation resistance: the way of *Deinococcus*
radiodurans. Cold Spring Harb Perspect Biol 5: a012765.

Kulkarni S, Ballal A, Apte SK. 2013. Bioprecipitation of uranium from alkaline waste
solutions using recombinant Deinococcus radiodurans. J Hazard Mater 262:853-861.

Mangwani N, Shukla S, Kumari S, Rao T, Das S. 2014. Characterization of
Stenotrophomonas acidaminiphila NCW-702 biofilm for implication in the degradation
of polycyclic aromatic hydrocarbons. J Appl Microbiol 117:1012-1024.

Marsili E, Beyenal H, Di Palma L, Merli C, Dohnalkova A, Amonette JE, Lewandowski Z.
2005. Uranium removal by sulfate reducing biofilms in the presence of carbonates. Wat
Sci Tech 52:49-55.

N'Guessan AL, Vrionis HA, Resch CT, Long PE, Lovley DR. 2008. Sustained removal of
uranium from contaminated groundwater following stimulation of dissimilatory metal
reduction. Environ Sci Technol 42:2999-3004.

Pagnanelli F, Petrangeli Papini M, Toro L, Trifoni M, Veglio F. 2000. Biosorption of metal
ions on Arthrobacter sp.: biomass characterization and biosorption modeling. Environ
Sci Technol 34:2773-2778.

Paul D, Pandey G, Pandey J, Jain RK. 2005. Accessing microbial diversity for
bioremediation and environmental restoration. Trends Biotechnol 23:135-142.

Savvin S. 1961. Analytical use of arsenazo III: determination of thorium, zirconium, uranium
and rare earth elements. Talanta 8:673-685.

Shukla SK, Mangwani N, Karley D, Rao TS. 2017. Bacterial Biofilms and Genetic
Regulation for Metal Detoxification. *In* Surajit Das, Dash HR, eds. Handbook of Metal-
Microbe Interactions and Bioremediation, 317-330

Shukla SK, Rao TS. 2013. Effect of calcium on *Staphylococcus aureus* biofilm architecture: a
confocal laser scanning microscopic study. Colloids Surf B 103:448-454.

Shukla SK, Rao TS. 2014. Calcium-Mediated Modulation of Staphylococcal Bacterial
Biofilms. Indian J Mar Sci 143:2107-2117

Shukla SK, Rao TS.2015. Heavy metals-bioremediation by highly radioresistant *Deinococcus*
radiodurans biofilm prospective use in nuclear reactor decontamination. Proceedings of
the symposium on water chemistry and corrosion in nuclear power plants in Asia-2015.

Shukla SK, Rao TS. 2017a. The first recorded incidence of *Deinococcus radiodurans* R1
biofilm formation and its implications in heavy metals bioremediation. bioRxiv:234781.

Shukla SK, Rao TS. 2017b. An Improved Crystal Violet Assay for Biofilm Quantification in
96-Well Microtitre Plate. bioRxiv:100214.

Shukla SK, Sankar GG, Paraneeiswaran A, Rao TS. 2014. Differential radio-tolerance of
nutrition-induced morphotypes of *Deinococcus radiodurans* R1. Curr Microbiol. 68:247-
253.

Stuart BH. 2005. Biological Applications. Pages 137-165. Infrared Spectroscopy:
Fundamentals and Applications, John Wiley & Sons, Ltd.

Suzuki Y, Kelly SD, Kemner KM, Banfield JF. 2004. Enzymatic U (VI) reduction by
Desulfosporosinus species. RADIOCHIM ACTA. 92:11-16.

Wall JD, Krumholz LR. 2006. Uranium reduction. Annu Rev Microbiol 60:149-166.

412
413

                                Legends to the Figures

Figure 1: Standard graph showing linear correlation between concentration of uranyl ion and

absorbance at 653 nm using modified Arsenazo II dye method.

Figure 2: Biofilm growth by *D. radiodurans* (DR1-bf[+]) strain. (a) CV assay conforming the

biofilm growth in the recombinant biofilm strain (DR1-bf[+]) compared to the wild type. (b)

Florescence microscope image showing the attachment of DR1-bf[+] cells, forming a biofilm.

DR1-bf[+] showing the green fluorescence due to the presence of constitutively expressing

GFP.

Figure 3: Comparison of uranium removal by *D. radiodurans* biofilm (grown with or

without Ca^{2+}) and planktonic cells of *D.radiodurans*-wild type and DR1-bf[+]. Uranium

concentrations were estimated by (A) Arsenazo dye method and (B) ICP – AES analysis. (C)

Visible precipitation of uranium (yellow color) was observed in the biofilm. Control DR1-

bf[+] biofilm biomass (2) DR1-bf[+] biofilm biomass treated with uranium aqueous solution

(1000 mg/L) for 5 min.

Figure 4: Uranium removal by different day old DR1-bf[+] biofilm grown in the presence of

20 mM Ca^{2+}

Figure 5: Scanning Electron Microscopic images of *Deinoccoccus radiodurans* (DR1_bf[+])

biofilm.

Figure 6: Fourier transformed IR spectra of *D. radiodurans* (DR1_bf[+]) biofilm biomass:

before (a), and after uranium treatment (b).

Figure 1

452   **Figure 2**

458   (**2**)

459   **Figure 3**

Figure 4

Figure 5

Figure 6