The Bayesian Superorganism III: externalised memories facilitate distributed

sampling

Edmund R. Hunt[1], Nigel R. Franks[1], Roland J. Baddeley[2]

[1]School of Biological Sciences, University of Bristol, Life Sciences Building, 24 Tyndall Avenue, Bristol,
BS8 1TQ, UK
[2]School of Experimental Psychology, University of Bristol, 12a Priory Road, Bristol, BS8 1TU, UK

ORCIDS: ERH, 0000-0002-9647-124X; NRF, 0000-0001-8139-9604; RJB, 0000-0002-7431-6580

Authors for correspondence: email: edmund.hunt@bristol.ac.uk, roland.baddeley@bristol.ac.uk

Abstract

A key challenge for any animal is to avoid wasting time by searching for resources in places it has

already found to be unprofitable. This challenge is particularly strong when the organism is a central

place forager – returning to a nest between foraging bouts – because it is destined repeatedly to

cover much the same ground. Furthermore, this problem will reach its zenith if many individuals

forage from the same central place, as in social insects. Foraging performance may be greatly

enhanced by coordinating movement trajectories such that each ant visits separate parts of the

surrounding (unknown) space. In this third of three papers, we find experimental evidence for an

externalised spatial memory in *Temnothorax albipennis* ants: chemical markers (either pheromones

or other cues such as cuticular hydrocarbon footprints) that are used by nestmates to mark explored

space. We show these markers could be used by the ants to scout the space surrounding their nest

more efficiently through indirect coordination. We also develop a simple model of this marking

behaviour that can be applied in the context of Markov chain Monte Carlo methods (see part two of

this series). This substantially enhances the performance of standard methods like the Metropolis–

Hastings algorithm in sampling from sparse probability distributions (such as those confronted by

the ants) with little additional computational cost. Our Bayesian framework for superorganismal

behaviour motivates the evolution of exploratory mechanisms such as trail marking in terms of

enhanced collective information processing.

Keywords: exploration, trail markers, spatial memory, Markov chain Monte Carlo, ants,

superorganism

Introduction

Exploring an unfamiliar, changing environment in search of valuable resources such as food or potential nest sites is a challenge for many organisms. A memory of where one has already explored, to avoid revisiting unprofitable locations, would generally seem to be an advantage. Nevertheless, while having a spatial memory of e.g. foraging locations is clearly adaptive, developing such a memory is associated with physiological costs. These include the metabolic overhead of a bigger brain (memory storage capacity) and the cost of encoding and retrieving memories (brain activity); these costs have to be traded-off against the benefit of improved foraging performance (Fagan et al. 2013). One way to circumvent the cost of carrying memories internally is to store the information externally in the environment. Indeed, such externalised information storage may have been the historical precursor to the development of internalised memory storage and retrieval (Chung and Choe 2009; Chung et al. 2009). Use of external markers as memory has been demonstrated in arguably simple organisms, including the slime mould *Physarum polycephalum* (Reid et al. 2012). Markers may be left in the environment to signify the presence or absence of good foraging or nest-making prospects at that location, so that when an animal returns it can make an appropriate and timely decision about the expenditure of its efforts.

An analogous problem to animals searching for food sources in an unfamiliar environment is that of sampling efficiently from unknown probability distributions. Markov chain Monte Carlo (MCMC) methods have been developed to generate such samples but suffer from the same problem of potentially revisiting the same area of probability space repeatedly. Although methods such as the Metropolis–Hastings algorithm show random walk behaviour in their sampling trajectory, they are still popular techniques because of their ease of implementation and computational simplicity. As discussed in the second paper of our series on the 'Bayesian superorganism' (Baddeley et al. 2018), MCMC methods can be used as a model of animal movement that enacts the optimal probability matching strategy for collective foraging ('gambling') (Kelly 1956). There, we drew a parallel with advances in efficiency in MCMC methods, to more adaptive animal behaviours emerging in natural

history: for example, from the random-walk type behaviour of the Metropolis-Hastings algorithm

(Metropolis et al. 1953; Hastings 1970) to the correlated random walks produced by Hamiltonian

Monte Carlo (Duane et al. 1987). The performance boost provided by externalised memories,

discussed here, may similarly be understood as an evolutionary advancement in collective

information processing capabilities.

Here, we present two main findings: (1) evidence that *Temnothorax albipennis* worker ants

avoid the footprints of nestmates during the reconnaissance of an unfamiliar space, and that this

enhances the efficiency of the colony-level collective exploration; (2) we develop a bio-inspired trail

avoidance method for the purpose of Markov chain Monte Carlo sampling that similarly uses a

memory of where the walker has visited to obtain a representative sample more rapidly. This

method is easy to implement and computationally simple, and offers significant improvements in

performance for the sort of sparse distributions that ants and other organisms also have to sample

from routinely. In the next section we introduce some more biological background on chemical

markers (externalised memories), but the reader familiar with this topic may proceed to the

Methods section.

Further biological background on chemical markers

In addition to expediting the process of environmental search, external markers can also

have the purpose of communicating one's presence to friendly or competitor conspecifics. Some

animals are well-known for marking territory: for instance wolves (*Canis lupus*) use scent marks to

maintain their territories (Peters and Mech 1975) while mice (*Mus musculus*) use olfactory cues to

help establish their territorial boundaries (Mackintosh 1973). In the Hymenoptera, the honeybee

Apis mellifera ligustica marks visited flowers with a scent and rejects flowers that have been recently

visited; it responds more strongly to its own markers than that of its nestmates (Giurfa 1993). Such

repellent scent-marks should help the bee to forage more efficiently. As in the ants, these marks

may simply be hydrocarbon footprints rather than costly signals (Eltz 2006). Ants are known to

engage in two distinguishable approaches to marking the area around their nest. In territorial

marking, ants will defend the marked area against intra- and inter-specific intruders. By contrast,

home-range markers label areas that the colony knows to be hospitable and available for foraging,

but such areas are not defended against other ant colonies (Hölldobler and Wilson 1990). Major

workers of the African weaver ant *Oecophylla longinoda* mark their territories with persistent

pheromones that are distinguishable to the ants at the colony level (Hölldobler and Wilson 1977),

whereas workers of the leafcutter ant *Atta cephalotes* deposit 'nest exit pheromones' in the vicinity

of their nest entrances, helping to orient workers and hence enhancing the efficiency of leaf

gathering (Hölldobler and Wilson 1986). *Lasius niger* ants have been found to engage in home-range

marking, which rather than territorial marking may help them to enlarge their potential foraging

space alongside other colonies, in a 'shared information' strategy (Devigne and Detrain 2002).

Apart from pheromones, which are signals laid deliberately by ants at a cost to themselves,

to influence the behaviour of other ants, ants have also been shown to respond to cues such as the

cuticular hydrocarbon footprints left by other ants. These can also be used to distinguish between

different colonies (Akino and Yamaoka 2005), and potentially also to recognize different castes, as in

a study of *Reticulitermes* termites (Bagnères et al. 1998). They can also be detected by ant

predators, and so this deposition is not without costs (Cárdenas et al. 2012). The response ants make

to detecting such markers depends on whether the cue is from a nestmate, a competing colony, or a

different species altogether, because depending on the relationship between sender and receiver, it

could represent a threat or useful information about the location of foraging resources (Wüst and

Menzel 2016). The role of the ant species (dominant or submissive) in the local community can be

key in this respect (Binz et al. 2014).

In our study organism, the house-hunting rock ant *T. albipennis*, there is evidence for

pheromone use in navigating to a new nest site (Pratt et al. 2001; McLeman et al. 2002) as well as in

T. curvispinosus (Bowens et al. 2013). Other *Temnothorax* species are also known to use individual-

specific trail pheromones for orientation outside their nests: *T. affinis* (Maschwitz et al. 1986) and *T.*

unifasciatus (Aron et al. 1988). *T. albipennis* is thought to use an individual-specific pheromone trail

during the nest-site assessment process: to measure the floor area using the 'Buffon's needle'

algorithm, an ant experiences a lower crossing rate of its own trail when the nest is larger. If its trail

was not individual-specific only one ant could do this before the trail was confused with those of

other scouting nestmates (Mallon and Franks 2000). *T. albipennis* colonies are also found to avoid

conspecific colonies when choosing where to locate its nest, as a result of odour cues left by foreign

colonies (Franks et al. 2007). Pheromones are also probably used by the ants when scouting

potential new nest sites, since when previously assessed nest sites are replaced with clean nest sites,

colonies are more likely to pick the poor option, having lost their reconnaissance information (Franks

et al. 2007). *T. rugatulus* was found to use an alarm pheromone that elicits two different behaviours

depending on the context: in an unfamiliar nest it causes ants to reject it as a potential new home,

to avoid danger; while when released near the ants' home nest it attracts them to defend against a

threat (Sasaki et al. 2014).

Chemical marks left by harvester ants (*Pogonomyrmex*) during exploration for new food

have been suggested as a way of avoiding redundant repeat searches in that area (Holldobler and

Wilson 1970; Carroll and Janzen 1973). The usefulness of an avoidance behaviour during the

searching phase for food was commented upon by Nonacs in a study of *Lasius pallitarsis* ants. Such a

behaviour was predicted to be more helpful for solitary foragers, than for species with mass

recruitment to prey items that need teams of workers to carry them home. Experimental

investigation found that the ants show a state-dependent avoidance of the path of previous

nestmates (Nonacs 1991). This question has recently been revisited when evidence was found that

Myrmica rubra (Myrmicinae) ants avoid the footprints of their nestmates, which suggests that they

are using the information it conveys to inform their foraging or scouting decisions (Binz et al. 2014).

However, this behaviour was not observed in the four other ant species in the study: *Formica*

polyctena (Formicinae), *Formica rufibarbis* (Formicinae), *Lasius niger* (Formicinae), and *Tetramorium*

caespitum (Myrmicinae). *T. albipennis* is also in the subfamily Myrmicinae, and we predicted that it

too uses cuticular hydrocarbon footprints left by previous exploring nestmates to scout an unknown

area surrounding its nest more efficiently. Here, we present evidence that *T. albipennis* indeed

avoids the footprints of nestmates during the reconnaissance of an unfamiliar space, and for the first

time show that it enhances the efficiency of the colony-level collective exploration.

Methods

Data processing and analysis

We examine data from a previous study of *T. albipennis* ant movement patterns (Hunt et al. 2016).

These data present two experimental regimes, one in which the foraging arena was entirely novel to

exploring ants, because it was cleaned by the experimenter in advance, and one in which previous

traces of the ants' activities remained. We call these treatments 'No cleaning' (NC) and 'Cleaning'

(C). These were both applied to six ants each from three colonies, for a total of 36 ants. The colonies

inhabited small artificial nests with a 2mm diameter entrance hole drilled into the centre of the top.

A colony was placed into a 90×90cm arena and a 15×15cm white paper mask with a 1×1cm central

hole affixed over the top, with a further 1.5×1.5cm white paper removable cover to block the nest

entrance. This created a smooth surface (with a raised area over the nest) to allow continuous

tracking of a single ant. The paper cover was removed to allow a single ant out of the nest, and the

entrance cover was then replaced, so that the ant would explore freely in isolation for 45 minutes

before being removed to a separate dish for the duration of the experiment. The exploration

trajectory of the ant was recorded by a camera mounted on a motorised gantry system, which

followed the ant's movements and recorded its path as a sequence of (x, y) points spaced by 0.1

seconds. In the NC treatment, six ants were consecutively released from the same colony on the

same day, and pheromone signals and/or other cues such as cuticular hydrocarbon footprints were

allowed to accumulate. In the C treatment, the arena surface was cleaned before the subsequent

ant was allowed into the arena, to prevent chemical communication between successive ants (see

the 'Methodology and apparatus' section of (Hunt et al. 2016) for a more detailed description of the

method).

In the previous study (Hunt et al. 2016) the focus of the analysis was on the characteristics of

movement 'events' – a period of non-zero speed followed by stopping – and the effect of treatments

on metrics such as average speed and the correlation between the average speed of consecutive

movement events. Here, we examine the effect of the treatment on exploration behaviour – how

efficiently the ants reconnoitre an unfamiliar experimental arena – with the hypothesis that the ants

use chemical information from previous exploring nestmates to do this more quickly, by avoiding

moving through regions of the space that have already been somehow marked as explored. We test

this hypothesis of enhanced exploration efficiency in the following way. In nature the ants might

benefit from sampling preferentially from surrounding regions of higher quality, in the sense of

being more likely to contain valuable resources such as food (Baddeley et al. 2018) or a potential

new nest site (Hunt et al. 2018); and would want to spend less time in 'empty' regions of the space

that contain little of relevance to their reproductive fitness. High-quality regions are likely to be

associated with cues that help the ants navigate toward them: for instance, a good potential nest

site may be more likely to be located in brighter regions of space than darker regions, because it

could benefit from warmer ambient temperatures. Therefore, in a natural environment we would

not expect the ants to explore uniformly in the region around their nest; and while we might naively

suppose that an empty experimental arena represents a uniform environment, from an ant's

perspective it may be somewhat heterogeneous in terms of lighting and so forth, despite an

experimenter's best efforts. Therefore, rather than measure the ants' exploration efficiency by

reference to how quickly they fill *uniform* space, we take the 'target distribution' that ants are trying

to sample from as being approximated by the actual exploratory trajectories taken by all 36 ants.

Although this may seem an unconventional approach, it will account for any heterogeneity in terms

of the lighting environment (though best efforts were taken to control it) and also differential

responses to surrounding landmarks in terms of lateralized behaviour. For instance, a preference for

viewing landmarks with the right eye has been suggested (Basari et al. 2013), which may affect

exploration behaviour, and also a leftward turning bias when exploring nest sites (Hunt et al. 2014),

which may relate to these ants having slightly asymmetric eyes (Hunt et al. 2018).

We create a target distribution for the exploring ants in the following way, using MATLAB

2015b. The exploration trajectories for all 36 ants are combined (a total exploration time of 27

hours), and trajectory points located inside the central region where the 15 × 15 cm paper mask is

located are removed, because many of the ants spend a significant proportion of their time

familiarising themselves with their immediate surroundings and engaging in thigmotaxis (edge

following) around the mask before beginning reconnaissance bouts away from the central region.

The (x, y) coordinates were then transformed from a Cartesian to a polar system (θ, r), where

$-\pi \leq \theta \leq \pi$ and $4.3 \leq \ln r < 6.5$ (the size of the arena in mm excluding the central area). A

periodic boundary for angle θ was created by adding the points in the region $[-\pi, 0]$ into a new

region $[\pi, 2\pi]$ and similarly copying $[0, \pi]$ into $[-2\pi, -\pi]$ to create a larger space $-2\pi \leq \theta \leq 2\pi$.

The coordinates from all 36 trajectories were counted using a histogram on a grid 881 × 1257 (θ step

size 0.01, $\ln r$ step size 0.0025). The periodic boundary prevented 'ringing' effects at the edge when

a gaussian blur was then applied over all the points in the space, to create a continuous distribution.

The gaussian blur has two parameters, filter size and σ which controls the width of the blur. In the

results presented a filter of size 200 × 300 was used, to be roughly a quarter of the size. Since the

ants are small (only 1mm wide) and hence only leave trails detectable in a surrounding space

perhaps 3mm wide, around 0.3% of the arena width, a fairly large blur is necessary to transform

their exploration trajectories into a continuous distribution. The σ used was set equal to 30, around

3% of the width of the new space, or $\approx$10 × the estimated trail width. After the blur was applied, the

distribution was then truncated back to its original size such that $-\pi \leq \theta \leq \pi$ again, and normalised

such that the sum of all the points was equal to 1 , as in a probability distribution where the value at

each position represents the likelihood of an ant wanting to visit it.

To measure how quickly each treatment converged toward the final distribution, the

exploration trajectories were combined by colony from all ants up to ant 1,2, … 6 out of the nest. For

instance, to measure the progress by ant 2 of the NC treatment, the trajectories of 2 ants were

combined (the first ant, and the second) to see its cumulative progress toward the goal. The process

described above was then repeated to obtain a probability distribution, which was then compared

against the final distribution using the cross-entropy (CE) as a measure of their similarity, where a

lower CE signifies greater progress toward the equilibrium distribution of a well-explored arena (see

also (Baddeley et al. 2018)). In the last step the three colony-level CE measures were averaged to

obtain a mean cross-entropy for the NC treatment by ant 2. Our hypothesis that the ants use

chemical information from preceding nestmates to explore more efficiently would correspond to a

CE that falls faster in the NC treatment than the C treatment.

While the NC and C treatments measure the difference to exploration made by chemical

information from nestmates, it is also quite possible that individual ants make use of their own

pheromone markers or other cues left behind, like their own footprints, to avoid wasting time

revisiting regions of space they have already walked through. To assess the benefit of individual-level

memory (both internal and external), we constructed simulated trajectories by randomly sampling

from a vector of the movements of all 36 ants. These 'Markov' ants provided a memoryless

benchmark of exploration performance, as ants that do not avoid their own trails, because there is

neither externalised memory (chemical markers) nor internal memory (in their brain) of where they

have been. A trajectory of 45 minutes of exploration was constructed by concatenating 1 second

periods of movement that stayed within the arena boundary (a 90×90cm space). The performance of

100 'colonies' of these Markov ants (6 per colony, as with the real colonies) was compared in the

same way.

To assess the statistical significance of one treatment converging more quickly on the final

distribution than the other, a permutation test was used, on ants 2-6 out of the nest. The first ant is

not included in the test for statistical significance since it is expected that the two treatments are

equivalent at this stage: the first ant explores an arena empty of any chemical information. A non-

parametric test is appropriate since the distribution of the CE for each treatment is unknown, and

relatively few data points are being compared (15 from each treatment). A permutation test was

also used to assess whether the average distance travelled by each exploring ant is different in the

NC and C treatments.

Trail avoidance model

In previous research (Baddeley et al. 2018; Hunt et al. 2018) we have proposed that a central

challenge that an ant colony confronts, that of trying to identify regions of value in an unknown

surrounding environment, is analogous to the problem found in statistics, physics, etc. of efficiently

sampling from complex probability distributions or computing numerical integrations. This is

because in order to sample from a probability distribution efficiently, one must identify regions of

high probability density, but these are not known *a priori* but only after evaluating the probability

density function at those locations. Similarly, an ant needs to find regions of high quality, but does

not know where they are unless it expends effort (time, energy) in exploration (Baddeley et al. 2018;

Hunt et al. 2018). An ant colony has a key advantage over a solitary animal: it is composed of many

workers, which can cooperate to benefit the colony as a whole, working together as one

'superorganism' (Hölldobler and Wilson 2009). Each worker could set off in a different direction to

that of its nestmates, and hence all of the surrounding area could be quickly explored. However, the

ants lack any central controller to direct their movements so as not to overlap and waste time

exploring the same region several times; instead they must make their decisions about where to

move based only on locally available information (Camazine et al. 2001). A key feature of the Markov

chain Monte Carlo (MCMC) movement models we have proposed is the use of local quality gradient

information in deciding where to move next: it would seem reasonable to expect ants (and other

organisms) preferentially to move toward regions of increasing quality (Baddeley et al. 2018). What

we have not yet considered is that this gradient may be modified by the ants themselves, by making

use of their sophisticated chemical communication systems (Hölldobler and Wilson 1990). We

suggest that ants may leave behind chemical markers, either deliberately (i.e. pheromones) or

incidentally (such as hydrocarbons from their footprints), that convey information about where they

and their nestmates have already explored. Such markers may be thought of as 'negative', in the

sense that an ant seeking to explore the environment will seek to avoid them, and as reducing the

quality gradient in that direction, such that other alternative directions have a relatively higher

gradient.

In a simple encapsulation of this exploration and trail avoidance behaviour, we can say that

an ant trying to sample from unknown resource quality distribution P also leaves behind a consistent

marker at every step that it takes, which corresponds to its subjective experience or model of the

world M (but which does not require any internal memory in the ant itself). M starts off as a uniform

prior of the world (e.g. a distribution of ones for each location) but then is added to (e.g. $+1$) at each

visited location. This is equivalent to Bayesian updating with a Dirichlet prior, because this is the

conjugate to the multinomial distribution (Frigyik et al. 2010). M can be renormalised at each time

step to be a probability distribution that sums to 1. Over time, as the ant more thoroughly explores

the distribution P, this model will converge $M \rightarrow P$, with more markers accumulating in regions of

higher probability. Since the objective of the ant is to sample from regions of high probability first,

one way to prioritise such regions is to sample from P^2, since this will substantially increase the

relative value of high probability regions over low probability regions. For instance, an ant would

rather visit a region of 0.2 probability over a region of 0.1, with the former being of $2 \times$ the quality

as the latter. If this is squared, this becomes a relative value of $\frac{0.2^2}{0.1^2} = \frac{0.04}{0.01} = 4 \times$ higher, making this

region of P space much more attractive, with a sharper gradient toward it too. Therefore, an

effective objective for ant exploration (or indeed MCMC methods) is to sample from:

286
$$P \times \frac{P}{M}$$

which converges in time to sample from P as $P/M \rightarrow 1$, but which should sample from high

probability regions first as a combination of the P^2 (attractive) and M (repulsive) terms. Such an

objective function for the ant or MCMC method to sample from does not require a change in the

method itself: it can be applied in conjunction with the most basic approaches such as Metropolis–

Hastings, which are computationally simple. A 'pseudocode' representation of our approach is

shown in Table 1.

Pseudocode for self-avoiding MCMC method applied to M–H algo.
input: target probability distribution P, α
output: ant's model of world M
M = α × ones (size (P)) *Dirichlet prior of world, e.g. α =1*
for t=1: simulation time
x _proposed = x _current + random (-1,0,1) *(same for y _prop.)*
ratio = P (x _prop, y _prop) / P (x _curr, y _curr) *(classic M–H criterion)*
R1 = P (x _prop, y _prop) / M (x _prop, y _prop)
R2 = P (x _old, y _old) / M (x _old, y _old)
ratio = ratio * (R1/R2) *(this obtains P 2 /M)*
if ratio > 1, accept = 1; elseif ratio > rand(1), accept = 1; else accept = 0
if accept = 1, (x,y) = proposed, else (x,y) = current *(unchanged)*
M(x,y) = M(x,y) + 1 *deposit 'negative pheromone'*
M=M/ ∑ M *renormalise to probability distribution summing to 1*
end for loop

294
295   Table 1. Pseudocode for the self-avoiding MCMC method.
296

Simulations of this 'trail-MCMC' model of ant exploration (or sampling approach) are presented in

the results, using a Metropolis–Hastings (M–H) method to sample from both a random distribution

(corresponding to a noisy, sparse world) and a distribution with three patches of differing qualities.

The random distribution was Dirichlet-distributed, generated by sampling from a gamma distribution

of shape k and scale parameter $\theta = 1$. The probability density function (p.d.f.) for this distribution is

shown in Figure 1. This results in mostly low-probability cells with a few higher-probability cells,

which corresponds to the ants' ecology where most space is unsuitable for locating the colony's

nest.

Figure 1. Probability density function for the gamma distribution with $k = 1, \theta = 1$.

The distribution with three patches had values of increasing relative magnitude of 1, 2, and 3 which
were given a Gaussian blur ($\sigma = 10$, filter size 100×100) and combined in equal proportion with a
gamma-distributed background noise, of $k = 0.2, \theta = 1$. Shape $k = 0.2$ results in a very sparse
distribution, with a few cells of higher probability. This background noise was given a Gaussian blur
($\sigma = 3$, filter size 100×100), which gave some local structure to the background noise (around the
higher probability cells). This resulted in the same distribution of 'nests' as featured in previous
papers on decision-making (Hunt et al. 2018) and foraging (Baddeley et al. 2018).

The M–H method samples from the distributions using periodic boundary conditions. To assess the

effect of cleaning (that is, removing the externalised spatial memory of the ants) the simulations are

also run under a 'cleaning' condition where the memory is reset five times, to correspond to the five

cleaning bouts in the experimental study (after each ant has explored, for six ants). To ease this

comparison the simulations are run for T=60,000 time steps, or 10,000 time steps per 'ant'. The

performance is compared to a M–H method sampling from the standard objective distribution P by

examining the rate of fall in cross-entropy between the sample and the target P (as previously

discussed in (Baddeley et al. 2018)).

Results

Ant exploration trajectories

The rate of convergence of the exploring ants toward the final distribution is shown in Figure 2; the

ants in the no cleaning treatment converge significantly more quickly than those in the cleaning

treatment (p=0.0161, permutation test on ants 2-6 across the 3 colonies. The analysis was repeated

for Gaussian blur size $\sigma = 20$, resulting in p=0.0394, and $\sigma = 40$, resulting in p=0.0098). This is

despite the average distance travelled by the ants from each treatment not being significantly

different. The average distance is 18.58 ± 0.77 m (s.e. of the mean) in the NC treatment and 17.57 ±

0.77 m in the C treatment (p=0.21, permutation test). As expected, without any memory

(externalised or internal) the simulated Markov ants converge on the target more slowly than either

of the experimental treatments, because they are essentially engaged in a random walk around the

arena.

Figure 2. The ants in the no cleaning treatment converge fastest toward the final distribution,
indicating that they benefit from chemical information left by preceding nest mates. The error bars
indicate one standard error of the mean; a permutation test on ants 2-6 indicate that the cross-
entropy is significantly lower in NC than C (p=0.0161). The 'Markov ant' converges slowest, because
it benefits neither from the chemical markers (externalised spatial memory) nor internal memory
about where it has already visited. The cross-entropy minimum is indicated by the dashed black line.
343

Trail avoidance model simulations

An example simulation of the models (standard M–H, trail-avoidance M–H with and without

cleaning) for the sparse distribution is shown in Figure 3. Such a distribution corresponds to a region

of empty space where the ants would wish to avoid sampling from the same parts of the distribution

repeatedly. It might be thought of as a 'zoomed-in' patch of empty space in the ants' environment,

as opposed to the distribution of three patches in Figure 4, which is 'zoomed-out' and shows all

available resources. Figure 3 shows that the standard M–H method converges slowly to the target

distribution when all the space needs to be sampled from roughly equally: with a random walk

considerable time is wasted going over previous ground. By contrast, the trail-avoidance M–H

methods converge much more quickly, as seen in a more rapidly falling cross-entropy. In a way

similar to the experimental result shown in Figure 2, the advantage of not cleaning the pheromone

trail is not immediately apparent but becomes obvious by 'ant 3' ($t = 3 \times 10^4$ on the x -axis),

because more pheromone has been accumulated for the no cleaning ant to avoid.

358
Figure 3. The trail avoidance model converges much more quickly to the target distribution, while
cleaning slows its progress somewhat. The target (cross-entropy minimum) is shown by the dashed
black line.

The result of a set of simulations sampling from three patches of different quality is shown in Figure

4. This does not show such a large improvement in performance for trail-avoidance M–H over

standard M–H, because significant portions of the probability mass are more readily accessible to

random walk-like behaviour. Nevertheless, the standard M–H method is seen to have spent more

time in the medium quality patch than the highest quality patch, in keeping with the general

problem the ants also face of prematurely favouring a local, rather than a global, quality maximum.

In comparison, the no cleaning trail-avoidance M–H samples from the three patches in the correct

proportion to their quality in this example, and converges most quickly to the target distribution as a

result. The cleaning simulation shows intermediate performance, sampling from the two main

patches but not exploring the whole distribution as effectively as when the pheromone memory is

allowed to accumulate.

Figure 4. The trail avoidance model again converges more quickly to the target. The M–H model is more likely to get 'stuck' in higher probability regions and take longer to get to other important parts of the distribution. However, the performance gain is less obvious than in the preceding gamma distribution example.

Discussion

Ants and other organisms confront the challenge of having to explore unknown environments while avoiding wasting time and incurring other costs and risks, which would occur if there are repeated visits to the same unprofitable areas. One strategy to avoid time-wasting would be for the animal to maintain an internal memory of where it has been, by using techniques such as learning the configuration of local landmarks (allothetic navigation), or counting steps in a certain direction from the nest (path integration, or other idiothetic mechanisms such as optic flow). However, the ants are pre-eminent manipulators of terrestrial chemical information (Hölldobler and Wilson 1990). Analysis of experimental data (Hunt et al. 2016) presented here provides evidence that *Temnothorax albipennis* ants use chemical depositions – either pheromones or cues such as cuticular hydrocarbon footprints – to explore an unknown arena more efficiently. This is because the ants in the treatment where chemical information is allowed to accumulate converge on the colony's target distribution more quickly than when that information is systematically cleaned away.

Our finding of collective exploration complements previous research on another Myrmicine ant, *Myrmica rubra*, that found evidence for nestmate footprint avoidance though did not demonstrate enhanced search efficiency (Binz et al. 2014). The avoidance behaviour is seen to be such that the trajectories of two successive ants will nevertheless intersect several times: persistent avoidance would, in effect, paint an ant into a corner. Instead the suggestion of a previous ant's reconnaissance leads to a reweighting of exploration toward a different region of space. In practice, where an ant holds private information about a source of food, for instance, that conflicts with social information (an exploratory marker) it is likely to dominate the decision about whether to return to the space when it is held with high confidence (probability) (Grüter et al. 2011). When there is a coherence between private memory and social information, this may help to boost ants' 'confidence' in their own memories and walk more directly (straighter and faster) toward their goal (Czaczkes et al. 2011).

The experimental evidence for path avoidance of nestmates helped to inspire a new

approach to tackling an analogous problem in the world of MCMC methods: slow convergence times

owing to repeated sampling of the same region of probability space. The original MCMC method,

Metropolis–Hastings, is well known for its random walk behaviour, yet is still a popular method

because of its ease of implementation. We have shown that its efficiency can be substantially

enhanced if it samples from the distribution $P \times {}^P\!/_M$, where M is a model (distribution) of where

the MCMC walker has visited in the space, rather than simply P (the target distribution). The

performance gain is particularly noticeable when sampling from sparse distributions, where the

need to avoid revisiting previous parts of the space is especially acute. *T. albipennis* ants live in just

such environments – where most of their surroundings hold little of relevance in terms of fitness,

either food or potential nest sites – and hence such a mechanism should indeed be favoured by

evolution. A shared substrate may serve effectively as a collective memory store, boosting the

information-processing capability of the colony beyond the bottleneck of a single ant's memory.

Here we have just examined the enhancement to sampling performance for a single walker (ant) – of

course in real colonies there will be many simultaneous walkers (ants). More sophisticated MCMC

methods that deploy parallel walkers can be compared to our approach, e.g. Wang and Landau

(2001).

The model presented here retains a perpetual memory of where the ant has walked,

although ongoing movement and pheromone deposition will lessen the relative weighting of past

trails as pheromone accumulates elsewhere. In reality, pheromones will decay in strength over time.

Although this is unavoidable, in a dynamic foraging environment it may well be preferable that

repellent markers decay over time in any event, because they may quickly lose their relevance and

even be maladaptive if they inhibit an ant from returning to an area where food has become

available. Similar considerations may apply as with animals deciding how to weigh past

observational memories with current ones; an exponential weighting is suitable in a Bayesian

analysis (McNamara and Houston 1987). It would be straightforward to introduce a reduction to the

potency of the markers over time by multiplying the model by a decay factor, as in $e^{-\lambda t}M$, where λ

is the decay constant. Theoretical consideration may be given to the optimal decay constant for

different environments (or probability distributions), with sparse environments with a lower chance

of opportunistic (randomly appearing) food favouring a lower constant, i.e. a slower decay. Such

predictions may be readily tested in different ant species associated with different ecologies.

Theoretical consideration of self-organised behaviours such as ant collective foraging had

predicted the value of inhibitory (negative) signals, before their empirical confirmation (Britton et al.

1998). In certain circumstances they could make the foraging process much more efficient (Stickland

et al. 1999). As anticipated, a 'no entry' signal has been found in Pharaoh's ants *Monomorium*

pharaonis to mark the unrewarding branch on a trail bifurcation. Concentrated at decision points –

where the path branches – it could be used to complement attractive trail pheromones or to

prevent an attractive pheromone becoming too strong through runaway positive feedback

(Robinson et al. 2005). These ants are thought to have at least three types of foraging trail

pheromone: one long-lasting attractive pheromone, thought to mark out the overall foraging

network, and two short-lived attractive and repellent pheromones, to mark out short term food

locations. The repellent pheromone was found to last more than twice as long as the short-lived

attractive pheromone (Robinson et al. 2008). Simulations indicate that repellent pheromone

increases the robustness and flexibility of the colony's overall foraging response (Robinson et al.

2008). The usefulness of such negative or repellent markers has also been demonstrated in a swarm

robotics context, in simulation (Fossum et al. 2015; Schroeder et al. 2017) or digitally on the robots

themselves (Payton et al. 2001; Van Dyke Parunak et al. 2005). Just as theoretical models of self-

organisation predicted the value of negative markers before they were empirically confirmed, it is

possible that advancements in Markov chain Chain Monte methods may inadvertently explain, or

pre-empt the discovery of, certain behavioural mechanisms. Similarly, the biology of collective

exploration is potentially a rich source of bio-inspiration for MCMC.

Conclusion

Many organisms face the challenge of identifying and exploiting food sources in an unfamiliar and changing environment. A common pitfall in the search process is revisiting previously explored regions of space that do not contain anything of value. An evolutionarily ancient way of avoiding this problem is to leave an externalised 'memory' of the previous visit by depositing a chemical marker that can be used to make a quick decision not to re-enter that space when it is encountered later. For superorganisms like ants, that seek to maximise their foraging performance at the level of the colony, pheromone signals or cues such as cuticular hydrocarbon footprints may be used to coordinate the movement decisions of their nestmates (i.e. through stigmergy) such that an unfamiliar space is quickly explored. Here, we presented evidence for avoidance of the paths of previous nestmates in *Temnothorax albipennis* ants, solitary foragers who would indeed particularly benefit from such a behaviour. We also demonstrated that this path avoidance phenomenon enhances the exploration efficiency of the ants at the colony level.

In addition to empirical findings, after noting the analogy between ant exploration and sampling from unknown probability distributions, we developed an ant-inspired enhancement to Markov chain Monte Carlo methods, whereby a memory is kept of where a walker has moved through the probability space. Such a method was shown to significantly enhance the efficiency of the Metropolis–Hastings method (Metropolis et al. 1953; Hastings 1970) when sampling from sparse probability distributions, because the random walk type behaviour of the method is reduced, but at little computational cost in comparison to more advanced momentum-based methods such as Hamiltonian Monte Carlo (Duane et al. 1987). When the 'superorganism' (Hölldobler and Wilson 2009) is examined using our Bayesian framework (Baddeley et al. 2018; Hunt et al. 2018), behavioural mechanisms such as individual trail markers become better understood in an ultimate sense (Tinbergen 1963) in relation to their facilitation of adaptive collective information-processing capabilities.

Author contributions. ERH drafted the paper and ran the simulations, NRF advised on social insect biology, RJB originated and advised on the theoretical concepts; all authors contributed to the present draft.

Funding statement. E.R.H. thanks the UK Engineering and Physical Sciences Research Council (grant no. EP/I013717/1 to the Bristol Centre for Complexity Sciences).

Ethics statement. The ant *Temnothorax albipennis* is not subject to any licencing regime for use in experiments. The ants were humanely cared for throughout the experiment.

Competing interests. We have no competing interests.

References

Akino T, Yamaoka R. 2005. Trail discrimination signal of *Lasius japonicus* (Hymenoptera: Formicidae). Chemoecology. 15(1):21–30. doi:10.1007/s00049-005-0288-6.

Aron S, Deneubourg JL, Pasteels JM. 1988. Visual cues and trail-following idiosyncrasy in L*eptothorax unifasciatus*: An orientation process during foraging. Insectes Soc. 35(4):355–366. doi:10.1007/bf02225811.

Baddeley RJ, Franks NR, Hunt ER. 2018. The Bayesian Superorganism II: optimal foraging and the information theory of gambling. bioRxiv.:497198. doi:10.1101/497198.

Bagnères A-G, Rivière G, Clément J-L. 1998. Artificial neural network modeling of caste odor discrimination based on cuticular hydrocarbons in termites. Chemoecology. 8(4):201–209. doi:10.1007/s000490050026.

Basari N, Bruendl AC, Hemingway CE, Roberts NW, Sendova-Franks AB, Franks NR. 2013. Landmarks and ant search strategies after interrupted tandem runs. J Exp Biol. 217(6):944–954. doi:10.1242/jeb.087296.

Binz H, Foitzik S, Staab F, Menzel F. 2014. The chemistry of competition: Exploitation of heterospecific cues depends on the dominance rank in the community. Anim Behav. 94:45–53. doi:10.1016/j.anbehav.2014.05.024.

Bowens SR, Glatt DP, Pratt SC. 2013. Visual Navigation during Colony Emigration by the Ant *Temnothorax curvispinosus*. PLoS One. 8(5). doi:10.1371/journal.pone.0064367.

Britton NF, Stickland TR, Franks NR. 1998. Analysis of Ant Foraging Algorithms. J Biol Syst. 06(04):315–336. doi:10.1142/S0218339098000212.

Camazine S, Deneubourg JL, Franks NR, Sneyd J, Theraulaz G, Bonabeau E. 2001. Self-organization in biological systems. Princeton University Press.

Cárdenas M, Jiroš P, Pekár S. 2012. Selective olfactory attention of a specialised predator to intraspecific chemical signals of its prey. Naturwissenschaften. 99(8):597–605. doi:10.1007/s00114-012-0938-9.

Carroll CR, Janzen DH. 1973. Ecology of Foraging by Ants. Annu Rev Ecol Syst. 4(1):231–257. doi:10.1146/annurev.es.04.110173.001311.

Chung JR, Choe Y. 2009. Emergence of Memory-like Behavior in Reactive Agents Using External Markers. In: 2009 21st IEEE International Conference on Tools with Artificial Intelligence. p. 404–408.

Chung JR, Kwon J, Choe Y. 2009. Evolution of recollection and prediction in neural networks. Proc Int Jt Conf Neural Networks:571–577. doi:10.1109/IJCNN.2009.5179065.

Czaczkes TJ, Grüter C, Jones SM, Ratnieks FLW. 2011. Synergy between social and private information increases foraging efficiency in ants. Biol Lett. 7(4):521–524. doi:10.1098/rsbl.2011.0067.

Devigne C, Detrain C. 2002. Collective exploration and area marking in the ant *Lasius niger*. Insectes Soc. 49(4):357–362. doi:10.1007/pl00012659.

Duane S, Kennedy AD, Pendleton BJ, Roweth D. 1987. Hybrid Monte Carlo. Phys Lett B. 195(2):216–

222. doi:10.1016/0370-2693(87)91197-x.

Van Dyke Parunak H, Brueckner SA, Sauter J. 2005. Digital Pheromones for Coordination of Unmanned Vehicles. In: Weyns D, Van Dyke Parunak H, Michel F, editors. Environments for Multi-Agent Systems. Berlin, Heidelberg: Springer Berlin Heidelberg. p. 246–263.

Eltz T. 2006. Tracing pollinator footprints on natural flowers. J Chem Ecol. 32(5):907–915. doi:10.1007/s10886-006-9055-6.

Fagan WF, Lewis MA, Auger-Methe M, Avgar T, Benhamou S, Breed G, LaDage L, Schlaegel UE, Tang W, Papastamatiou YP, et al. 2013. Spatial memory and animal movement. Ecol Lett. 16(10):1316–1329. doi:10.1111/ele.12165.

Fossum F, Montanier JM, Haddow PC. 2015. Repellent pheromones for effective swarm robot search in unknown environments. 2014 IEEE Symp Swarm Intell Proc.:243–250. doi:10.1109/SIS.2014.7011802.

Franks NR, Dornhaus A, Hitchcock G, Guillem R, Hooper J, Webb C. 2007. Avoidance of conspecific colonies during nest choice by ants. Anim Behav. 73:525–534. doi:10.1016/j.anbehav.2006.05.020.

Franks NR, Hooper JW, Dornhaus A, Aukett PJ, Hayward AL, Berghoff SM. 2007. Reconnaissance and latent learning in ants. Proc R Soc London B Biol Sci. 274(1617):1505–1509. doi:10.1098/rspb.2007.0138.

Frigyik BA, Kapila A, Gupta MR. 2010. Introduction to the Dirichlet Distribution and Related Processes. Department of Electrical Engineering, University of Washington. UWEETR-2010-0006.

Giurfa M. 1993. The repellent scent-mark of the honeybee *Apis mellifera tigustica* and its role as communication cue during foraging. Insectes Soc. 40(1):59–67. doi:10.1007/bf01338832.

Grüter C, Czaczkes TJ, Ratnieks FLW. 2011. Decision making in ant foragers (*Lasius niger*) facing conflicting private and social information. Behav Ecol Sociobiol. 65(2):141–148. doi:10.1007/s00265-010-1020-2.

Hastings WK. 1970. Monte Carlo sampling methods using Markov chains and their applications. Biometrika. 57(1):97-. doi:10.2307/2334940.

Holldobler B, Wilson EO. 1970. Recruitment Trails in the Harvester Ant *Pogonomyrmex Badius*. Psyche (Stuttg). 77(4):385–399. doi:10.1155/1970/38470.

Hölldobler B, Wilson EO. 1977. Colony-specific territorial pheromone in the African weaver ant Oecophylla longinoda (Latreille). Proc Natl Acad Sci. 74(5):2072–2075.

Hölldobler B, Wilson EO. 1986. Nest area exploration and recognition in leafcutter ants (Atta cephalotes). J Insect Physiol. 32(2):143–150. doi:10.1016/0022-1910(86)90133-2.

Hölldobler B, Wilson EO. 1990. The Ants. Harvard University Press.

Hölldobler B, Wilson EO. 2009. The superorganism: the beauty, elegance, and strangeness of insect societies. WW Norton & Company.

Hunt ER, Baddeley RJ, Worley A, Sendova-Franks AB, Franks NR. 2016. Ants determine their next move at rest: motor planning and causality in complex systems. R Soc Open Sci. 3(1):150534. doi:10.1098/rsos.150534.

Hunt ER, Baddeley RJ, Worley A, Sendova-Franks AB, Franks NR. 2016. Data from: Ants determine their next move at rest: motor planning and causality in complex systems. R Soc Open Sci. 3(1):150534. doi:10.1098/rsos.150534.

Hunt ER, Dornan C, Sendova-Franks AB, Franks NR. 2018. Asymmetric ommatidia count and behavioural lateralization in the ant *Temnothorax albipennis.* Sci Rep. 8(1). doi:10.1038/s41598-018-23652-4.

Hunt ER, Franks NR, Baddeley RJ. 2018. The Bayesian Superorganism I: collective probability estimation. bioRxiv.:468942. doi:10.1101/468942.

Hunt ER, O'Shea-Wheller T, Albery GF, Bridger TH, Gumn M, Franks NR. 2014. Ants show a leftward turning bias when exploring unknown nest sites. Biol Lett. 10(12). doi:10.1098/rsbl.2014.0945.

Kelly JL. 1956. A new interpretation of information rate. Bell Syst Tech J. 35(4):917–926.

Mackintosh JH. 1973. Factors affecting the recognition of territory boundaries by mice (*Mus musculus*). Anim Behav. 21(3):464–470. doi:10.1016/S0003-3472(73)80006-5.

Mallon EB, Franks NR. 2000. Ants estimate area using Buffon's needle. Proc R Soc B Biol Sci.
267(1445):765–770. doi:10.1098/rspb.2000.1069.
Maschwitz U, Lenz S, Buschinger A. 1986. Individual specific trails in the ant *Leptothorax affinis*
(Formicidae: Myrmicinae). Experientia. 42(10):1173–1174. doi:10.1007/bf01941298.
McLeman MA, Pratt SC, Franks NR. 2002. Navigation using visual landmarks by the ant *Leptothorax*
albipennis. Insectes Soc. 49(3):203–208. doi:10.1007/s00040-002-8302-2.
McNamara JM, Houston AI. 1987. Memory and the efficient use of information. J Theor Biol.
125(4):385–395. doi:10.1016/S0022-5193(87)80209-6.
Metropolis N, Rosenbluth AW, Rosenbluth MN, Teller AH, Teller E. 1953. Equation of state
calculations by fast computing machines. J Chem Phys. 21(6):1087–1092. doi:10.1063/1.1699114.
Nonacs P. 1991. Exploratory behavior of *Lasius pallitarsis* ants encountering novel areas. Insectes
Soc. 38(4):345–349. doi:10.1007/BF01241870.
Payton D, Daily M, Estowski R, Howard M, Lee C. 2001. Pheromone Robotics. Auton Robots.
11(3):319–324. doi:10.1023/A:1012411712038.
Peters RP, Mech LD. 1975. Scent-Marking in Wolves. Am Sci. 63(6):628–637.
doi:http://www.jstor.org/stable/27845779.
Pratt SC, Brooks SE, Franks NR. 2001. The Use of Edges in Visual Navigation by the Ant *Leptothorax*
albipennis. Ethology. 107(12):1125–1136. doi:10.1046/j.1439-0310.2001.00749.x.
Reid CR, Latty T, Dussutour A, Beekman M. 2012. Slime mold uses an externalized spatial "memory"
to navigate in complex environments. Proc Natl Acad Sci. 109(43):17490–17494.
doi:10.1073/pnas.1215037109.
Robinson EJH, Green KE, Jenner EA, Holcombe M, Ratnieks FLW. 2008. Decay rates of attractive and
repellent pheromones in an ant foraging trail network. Insectes Soc. 55(3):246–251.
doi:10.1007/s00040-008-0994-5.
Robinson EJH, Jackson DE, Holcombe M, Ratnieks FLW. 2005. Insect communication: "No entry"
signal in ant foraging. Nature. 438(7067):442. doi:10.1038/438442a.
Robinson EJH, Ratnieks FLW, Holcombe M. 2008. An agent-based model to investigate the roles of
attractive and repellent pheromones in ant decision making during foraging. J Theor Biol.
255(2):250–258. doi:10.1016/j.jtbi.2008.08.015.
Sasaki T, Hölldobler B, Millar JG, Pratt SC. 2014. A context-dependent alarm signal in the ant
Temnothorax rugatulus. J Exp Biol. 217(18):3229–3236. doi:10.1242/jeb.106849.
Schroeder A, Ramakrishnan S, Kumar M, Trease B. 2017. Efficient spatial coverage by a robot swarm
based on an ant foraging model and the Lévy distribution. Springer US.
Stickland TR, Britton NF, Franks NR. 1999. Models of information flow in ant foraging: the benefits of
both attractive and repulsive signals. In: Detrain C, Deneubourg JL, Pasteels JM, editors. Information
Processing in Social Insects. Basel: Birkhäuser Basel. p. 83–100.
Tinbergen N. 1963. On aims and methods of Ethology. Z Tierpsychol. 20(4):410–433.
doi:10.1111/j.1439-0310.1963.tb01161.x.
Wang F, Landau DP. 2001. Efficient, Multiple-Range Random Walk Algorithm to Calculate the Density
of States. Phys Rev Lett. 86(10):2050–2053. doi:10.1103/PhysRevLett.86.2050.
Wüst M, Menzel F. 2016. I smell where you walked – how chemical cues influence movement
decisions in ants. Oikos 126(1).. doi:10.1111/oik.03332.
625