1 **Plasticity and the role of mass-scaling in allocation, morphology and anatomical**
2 **trait responses to above and belowground resource limitation in cultivated**
3 **sunflower (_Helianthus annuus_ L.)**
4

5 Yan Wang[1,2], Lisa A. Donovan[2] , Andries A. Temme[2*]

6 1. College of Forestry, Shandong Agriculture University, Taian, Shandong, 271018, China

7 2. Department of Plant Biology, Franklin College of Arts and Sciences, The University of Georgia,
8 Athens, GA, 30602, USA

9 *Correspondence author: atemme@uga.edu (+1 706 207 4020)

Abstract

- In the face of resource limitations, plants show plasticity in multiple trait categories, including biomass allocation, morphology and anatomy, yet inevitably also grow less. The extent to which passive mass-scaling plays a role in trait responses that contribute to increased potential for resource acquisition are poorly understood. Here we assessed the role of mass-scaling on the direction, magnitude and coordination of trait plasticity to light and/or nutrient limitation in cultivated sunflower (*Helianthus annuus*).

- We grew seedlings of ten sunflower genotypes for three weeks in a factorial of light (50% shade) and nutrient (10% supply) limitation in the greenhouse and measured a suite of allocational, morphological and anatomical traits for leaves, stems, fine roots, and tap roots.

- Under resource limitation, plants were smaller and more biomass was allocated to the organ capturing the most limiting resource, as expected. Traits varied in the magnitude of plasticity and the extent to which the observed response was passive (scaled with plant mass) and/or had an additional active component. None of the allocational responses were primarily passive. Plastic changes to specific leaf area and specific root length were primarily active, and adjusted towards more acquisitive trait values under light and nutrient limitation, respectively. For many traits, the observed response was a mixture of active and passive components, and for some traits the active adjustment was antagonistic to the direction of passive adjustment, e.g. stem height, and tap root and stem theoretical hydraulic conductance. Passive scaling with size played a major role in the coordinated response to light, but correcting for mass clarified that the active responses to both limitations were more similar in magnitude, although still resource and organ specific.

- Our results demonstrate that both passive plasticity and active plasticity can contribute to increased uptake capacity for limiting resources in a manner that is resource, organ and trait specific. Indeed, passive adjustments (scaling with mass) of traits due to resource stress extends well beyond just mass allocation traits. For a full understanding of plants response to environmental stress both passive and active plasticity needs to be taken into account.

- **Keywords:** allocation, morphology, anatomy, allometry, phenotypic plasticity, K_s, SLA, SRL

Introduction

The extent of plant trait adjustment in response to a changed environment is generally considered as the plant's phenotypic plasticity (Valladares, Gianoli, & Gomez, 2007; Nicotra et al., 2010). According to theory, this plasticity serves to optimize/maximize the uptake of the most limiting resource (Bloom, Chapin, & Mooney, 1985; Gedroc, McConnaughay, & Coleman, 1996; Shipley & Meziane, 2002; Poorter et al., 2012). For example, increased mass allocation to leaves under shade, and to roots under nutrient limitation, alleviates some of the stress caused by the resource limitation (Shipley & Meziane, 2002; Sugiura & Tateno, 2011). However, plants are inevitably smaller under resource stress, raising the question of the role of mass-scaling in individual and coordinated traits shifts in response to resource limitation (McConnaughay & Coleman 1999; Shipley & Meziane, 2002; Weiner 2004; Poorter et al. 2012; Osnas, Lichstein, Reich, & Pacala, 2013; Reich, 2014 and 2018).

There is a long tradition of investigating mass-based scaling of biomass allocation to different plant parts (Weiner 2004). Generally, smaller plants allocate proportionally more mass to leaves than to roots (Poorter et al., 2012). Trait responses that scale with mass, which have been variously called allometric scaling, "passive" plasticity, or "apparent" plasticity, could predispose the plant to align its traits with resource availability and demand (McConnaughay & Coleman, 1999; Weiner 2004; van Kleunen & Fisher, 2005; Nicotra *et al.*, 2010; Poorter et al., 2012). However, there is evidence that additional "active" or "true" plasticity in biomass allocation can further increase the capacity for acquiring the most limiting resource (McConnaughay & Coleman, 1999; Shipley & Meziane, 2002; Poorter et al. 2012). There is also evidence of active plasticity in other traits related to resource uptake. For example, under low light conditions greater plant height and greater specific leaf area for a given plant mass aids in light uptake (Rice & Bazzaz, 1989; Freschet, Swart, & Cornelissen, 2015; Freschet, Violle, Bourget, Scherer-Lorenzen, & Fort, 2018; Reich, 2018). Thus, both passive and active trait adjustments combine in the realized response to resource stress.

The role of passive and active contributions to observed responses can be investigated by including analyses that correct responses for plant size (Poorter et al., 2012; Reich, 2018). In this framework, passive adjustments in traits associated with size cannot be a priori regarded as or ruled out as adaptive (Nicotra et al., 2010; Poorter et al., 2012 and 2019). If traits responses that are consistent with greater ability to take up the most limiting resource have both passive and active components, it will be important to consider both the magnitude and alignment of both components when evaluating evidence for functional and putatively adaptive responses.

Among plant traits, anatomical traits are often overlooked due to time and budget constraints. However, variation in anatomical traits underlies or contributes to variation in morphological and physiological traits that have received more attention (Kong et al., 2014; Scoffoni et al., 2015; John et al., 2017). For example, palisade parenchyma thickness is positively correlated to leaf thickness (Catoni, Gratani, Sartori, Varone, & Granata, 2015) and photosynthetic rate (Chatelet, Clement, Sack, Donoghue, & Edwards, 2013). Additionally, a thicker cortex could provide a relative larger site for mycorrhizal infection and higher resource uptake in thicker roots, especially for the arbuscular mycorrhiza (Kong et al., 2014). Root cortex thickness, due to the size of cortical cells (Eissenstat & Achor, 1999) strongly affects fine root diameter (Gu, Xu, Dong, Wang, & Wang, 2014; Guo et al., 2008), and a wider stele and/or xylem conduit greatly affects hydraulic conductivity (McElrone, Pockman, Martínez-Vilalta, & Jackson, 2004; Rico, Pitterman, Polley, Aspinwell, & Fay, 2013; Tyree & Ewers, 1991). Thus, xploring how passive and active responses in anatomical traits align with those of other traits will enhance our understanding of how plants adjust to changing environmental conditions from tissue, to organ, to architecture.

Across species, plant functional traits are thought to form a spectrum of resource use strategies
from fast to slow (Wright et al., 2004; Fortunel, Fine, & Baraloto, 2012; Reich, 2014; Díaz et al., 2016).
The "whole plant economic spectrum" suggests that values of traits related to resource use,
including mass allocation, morphology and transport, should be coordinated across all organs and
resources, with more resource conservative traits associated with adaptation to light and nutrient
limited habitats (Reich, 2014). For closely related species, there is mixed evidence of correlated
evolution among some traits related to resource acquisition and processing (Mason & Donovan,
2015; Bowsher, Mason, Goolsby, & Donovan 2016; Pilote & Donovan, 2016; Medeiros, Burns,
Nicolson, Rogers, & Valvarde-Barrantes, 2017; Muir, Conesa, Roldán, Molins, & Galmés, 2017).
However, much remains unclear about how the coordinated traits across 'whole plant economic
spectrum' relate to trait plasticity in response to resource limitations (Reich, 2014; Anderegg et al.,
2018; Agrawal, 2020). Thus, it is interesting to assess the coordination of responses across organs
and a broad range of traits in terms of the role of passive and active plasticity.

To add to our understanding of plant responses to resource limitation, we examined trait responses
to light and nutrient limitation of traits across different trait categories (biomass allocation,
morphology and anatomy) and organs (leaf, stem and root) in cultivated sunflower. Prior research
has shown strong plastic responses to resource limitation and other environmental factors in *H.*
annuus (Rico, Pitterman, Polley, Aspinwell, & Fay, 2013; Donovan, Mason, Bowsher, Goolsby, &
Ishibashi, 2014; Bowsher et al., 2017; Masalia, Temme, Torralba, & Burke, 2018; Temme et al., 2019).
Specifically, we sought to answer the following questions:

1) How do mass allocation, organ morphology, and anatomy change with above and below
ground resource limitation, and what role does size scaling of traits play in this?
2) How do traits compare for magnitude of plasticity and what role does size scaling of traits
play in this?
3) Do traits show a coordinated shift due to resource limitation across all organs and what role
does size scaling of traits play in this?

Material and Methods

Experimental design

To address these questions, we selected a set of 10 cultivated sunflower genotypes (Supplemental
Table S1), varying broadly in biomass based on prior work, from a larger diversity panel used for
genomic dissection of traits (Mandel, Dechaine, Marek, & Burke, 2011; Nambeesan et al., 2015;
Masalia, Temme, Torralba, & Burke, 2018). We conducted a factorial design of two nutrient
treatments (rich and poor) and two light treatments (sun and shade) at the Botany greenhouses of
The University of Georgia, Athens GA, USA, in March 2018. Achenes were sown in seedling trays and
allowed to grow for seven days, after which each seedling was transplanted to 5 liter (1.3 gallon) pot
filled with a 3:1 sand:calcinated clay mixture (Turface MVP, Turface Athletics, Buffalo Grove, IL). Pots
were arranged in a split plot design of 6 replicate blocks. The light treatment was applied as the
whole-plot factor, with 2 sub-plots in each plot randomly assigned to unshaded or 50% shade
generated with high density woven polyethylene cloth (Supplemental Figure S1). Within each
subplot, two pots of each genotype were randomly distributed and supplied with either 40g or 4g
fertilizer (Osmocote Plus 15-9-12 with micronutrients, Scotts, Marysville, OH, USA), totaling 240 pots
(plants). Greenhouse temperature controls were set to maintain 18–24 °C, and natural sunlight was
supplemented with sodium halide lighting (20-25 μmol m^{-2} s^{-1}) to maintain a 15/9-h photoperiod.

Plant harvest and trait measurements

132 Plants were harvested 3 weeks after transplanting (4 weeks after germination). At harvest, stem

height (ST-Hgt, from soil surface to top of apical meristem) and stem diameter (ST-Dia, midway
between cotyledons and first leaf pair) were measured. Plants were separated into root, leaf
(including cotyledons), and stem (including bud if present—rarely) for biomass and other
measurements. We assigned all measured traits to one of three categories (allocational,
morphological, anatomical) in order to compare the relative magnitude of adjustments to these
broad categories. While we believe our assignment of traits to categories is defensible, we
acknowledge that this is somewhat arbitrary and that different groupings could influence results
associated with comparisons among categories.

Each replicate plant was sampled for leaf, stem and root tissue for anatomical traits. One recently
matured fully expanded leaf was sampled for a 1×0.5 cm rectangle cut out of the leaf center. The
stem was sampled for a 5 mm length segment centered between the cotyledon and the first leaf pair.
The root was sampled for both tap root and fine root tissue. For the tap root, a 1 cm segment was
cut 4 cm below the root/stem junction. For a single lateral root attached to the tap root near to the
root/stem junction with an intact root tip, a 1 cm fine root segment was cut 2 cm from the apex of
the root. All tissue subsamples were weighed for fresh mass and then fixed in formalin–acetic acid-
alcohol solution, FAA (50% ethanol (95%), 5% glacial acetic acid, 10% formaldehyde (37%) and 35%
distilled water).

Fixed subsamples were processed for anatomy at the University of Georgia Veterinary Histology
Laboratory. Each sample was embedded and gradually infiltrated with paraffin, sliced with a sledge
microtome, mounted to a slide, and stained with safranin and fast green dye. Slides were imaged
with a camera mounted Zeiss light microscope using ZEN software (Carl Zeiss Microscopy,
Oberkochen, Germany). For leaf, stem, and fine and tap roots, the dimensions of anatomical
features were traced using Motic Images Advanced 3.2 software (Motic Corporation, Xiamen, China).
The organs dimensions included leaf thickness LF-Th), fine root diameter (FR-Dia) and tap root
diameter (TR-Dia). The tissue dimensions included leaf palisade thickness (LF-PT), Leaf spongy
thickness (LF-ST), stem cortex thickness (ST-CT), Stem xylem thickness (ST-XT), tap root cortex
thickness (TR-CT), tap root stele diameter (TR-SDia), fine root cortex thickness (FR-CT), and fine root
stele diameter (FR-SDia). The theoretical hydraulic conductivity (Ks, $kg \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$) per vascular
area for leaf (LF-Ks), stem (ST-Ks), fine root (FR-Ks), and tap root (TR-Ks) tissue was calculated (Sperry,
Hacke, & Pittermann, 2006), based on the Hagen-Poseuille equation (Tyree & Ewers, 1991):
$ks = (\pi\rho/128\eta A_w) \sum_{i=1}^{n} d_i^4$. where ρ is the density of water (988.3 $kg \cdot m^{-3}$ at 20 °C); η is the viscosity
of water (1.002×10^{-9} MPa·s at 20 °C); A_w is the stele (vascular) cross-section area, d is the diameter of
the i^{th} vessel and n is the number of conduits in the xylem.

After anatomical samples were collected, the remaining plant material was used to determine
morphological traits. The leaf, stem, and fine and tap roots for each plant were weighed for fresh
mass. Root tissue, stem tissue, and the most recently fully expanded leaf was scanned at 300 dpi
with an Epson Expression1680 scanner (Seiko Epson Corporation, Nagano-ken, Japan) and saved as a
TIF image. Total root length and volume of scanned fine root, tap root and stem, as well as leaf area
were measured using WinRhizo (v. 2002c, Regent Instruments, Quebec, Canada). Subsequently, the
leaf, stem, and fine roots and tap roots were dried at 60°C for 48 h and weighed. Specific leaf area
(LF-SLA, $cm^2 \cdot g^{-1}$) was calculated as the ratio of leaf area to leaf dry mass. Specific root length ($m \cdot g^{-1}$)
was calculated as the ratio of root length to root dry mass for the tap root and fine roots (TR-SRL, FR-
SRL, respectively). Tissue density ($g \cdot cm^{-3}$) was calculated as the ratio of dry mass to volume for stem,
tap toot, and fine roots (ST-Den, TR-Den, FR-Den, respectively). Leaf dry matter content, measured
as leaf dry mass divided by leaf fresh mass, was used as a proxy for leaf tissue density (LF-Den)
(Kramer-Walter et al., 2016; Wilson, Thompson, & Hodgson, 1999).

For allocational traits, total plant dry mass was calculated as the sum of all plant parts, including the
subsamples for anatomical and morphological traits. The fresh biomass of subsamples for
anatomical analysis was converted to dry biomass based on the ratio of fresh/dry biomass of for the
morphological traits. The mass fractions for each tissue were calculated as proportions of total plant
dry mass ($g \cdot g^{-1}$).

Data analysis
The statistical analysis for the phenotypic data was performed using R v3.5.1 (R Core Team). To
obtain genotype means from our split plot design, a mixed effects model was fitted using the
package *lme4* (Bates et al., 2018) with genotype, light and nutrient level and all their interactions as
fixed effects and light treatment within block as random factor. Least-square (LS) means of all trait
values without random factor were estimated from this model using the R package *emmeans* (Lenth
et al., 2018). To test the effect of genotype and treatment on measured traits we fitted a less
expansive mixed effects model with genotype, light and nutrient level as well as the interaction
between light and nutrient level as fixed effects (following Freschet, Violle, Bourget, Scherer-
Lorenzen, & Fort, 2018) and light treatment within block as random factor. From this model, fixed
effects were then tested using a Walds Chi-square test in a type III ANONA using the package *car*
(Fox et al., 2018). As we were interested in the interactive effects among genotype, light and
nutrient supply on the proportional changes in functional traits, rather than on their absolute
changes, we performed all analyses on log-transformed data (Freschet, Swart, & Cornelissen, 2015).
For each trait, differences among treatments were tested using Tukey's HSD (p=0.05) corrected for
multiple comparisons. We then estimated the influence of plant size on the significance of nutrient
and light limitation effects on traits by adding (log-transformed) plant biomass as a fixed factor to
both models and recalculating means and significance (Ryser & Eek, 2000; Wahl, Ryser, & Edwards,
2001).

To quantify the plastic response of each trait to each resource limitation treatment, we calculated
the relative distance plasticity index (RDPI, Valladares, Sanchez-Gomez, & Zavala, 2006, Scoffoni et
al., 2015) as $(x_i' - x_i)/(x_i' + x_i)$, where x_i and x_i' are the mean trait values of genotypes grown
under control (high-light and high-nutrient) and resource limited condition, respectively. Additionally,
the RDPI for each trait in each resource limitation treatment was recalculated after correcting for
plant size (total biomass). Significant values of RDPI (difference from zero, no plasticity) were
determined using T-test on genotype averages.

To determine major axes of variation across multiple traits and identify whether there were
concerted trait adjustments to limitation in above or belowground resources, we conducted a
principal component analysis (PCA) on the trait data before and after correcting for size. Differences
between treatments were tested using, Bonferroni corrected, Hotellings-t test on the first two
principal components. Data visualizations were made using *ggplot2* (Wickham et al., 2018).

Results
<u>How do plant mass allocation, organ morphology, and anatomy respond to above and below ground</u>
<u>resource limitation, and what role does size scaling of traits play in this?</u>

Across all genotypes, total plant biomass decreased substantially in response to light and nutrient
limitation (Figure 1 and Table 1). The biomass response to light limitation (54.7% decline) was
stronger than the response to the nutrient limitation (24.7% decline), and there was no interaction
of light and nutrient limitation on biomass when combined (67.1%% decline). Of the 31 allocational,
morphological and anatomical traits measured, all but two responded to at least one resource

limitation (Supplemental Table S2). Responses to light limitation were more prevalent than
responses to nutrient limitation, and there were relatively few interactive effects of combined light
and nutrient limitation.

For allocational traits, resource limitations affected all 6 traits, but traits were affected in contrasting
ways by light and nutrient limitation. The RatioLF-FRmass (ratio of leaf mass to fine root mass)
increased under light limitation due to increased LF-MF (leaf mass fraction) and decreased FR-MF
(fine root mass fraction) (Figures 2b-d and Table 1). Conversely, RatioLF-FRmass decreased under
nutrient limitation, due to decreased LF-MF and increased FR-MF (Figure 2f). A significant interaction
of light and nutrient limitation was found for LF-MF, ST-MF (stem mass fraction), FR-MF, and
RatioLF-FRmass. The RatioSLA-SRL (ratio of LF-SLA (specific leaf area) to FR-SRL (fine root specific
root length)), which additionally considers surface area per unit mass of acquisitive organ, paralleled
that of the RatioLF-FRmass, with a trend of increasing under light limitation and decreasing under
nutrient limitation (Figure 2e).

For morphological traits, light limitation affected 11 of the 12 traits, all except for FR-Dia (fine root
diameter) (Table 1). The responses led to increases in acquisitive values for resource acquiring traits
such as LF-SLA (Figure 2a), FR-SRL (Figure 2c), TR-SRL (tap root specific root length), and ST-Hgt (stem
height). There were also strong decreases in organ dimensions, e.g., LF-Th (leaf thickness), ST-Dia
(stem diameter), and TR-Dia (tap root diameter). LF-Den (leaf density), ST-Den (stem density), and
FR-Den (fine root density) also declined. Contrasting with the effect of light limitation, nutrient
limitation affected only four out of 12 morphological traits (Table 1), leading to decreased ST-Dia and
FR-Dia, and increased FR-SRL and TR-Den (tap root density).

For anatomical traits, limiting resources affected 12 of the 13 traits, and responses again differed by
limiting resource (Figure 3 and Table 1). Light limitation affected anatomical traits of all three organs.
Aboveground, light limitation decreased LF-PTh (leaf palisade parenchyma layer thickness), ST-CTh
(stem cortex thickness), ST-VTh (stem vascular bundle thickness), and ST-XTh (stem xylem thickness).
Belowground, light limitation decreased TR-CTh (tap root cortex thickness) and TR-SDia (tap root
stele diameter). Vascular tissue adjustment to light limitation led to changes in theoretical hydraulic
conductivity, with decreased LF-Ks (leaf hydraulic conductivity) but increased ST-Ks (stem hydraulic
conductivity). In contrast to the effects of light limitation, nutrient limitation predominately affected
fine root anatomy, leading to decreased FR-CTh (fine root cortex thickness) and FR-SDia (fine root
stele diameter).

To assess the role of passive mass-based scaling in the responses of resource limitations, trait
responses were reanalyzed with mass as a covariate in the model (Table 1, Supplemental Figure S3),
effectively comparing treatments again for trait values of plant of the same size. Contrasting the
results of the two models (including mass as covariate in Table 1), we designated the observed
response as being primarily passive (scaling with mass), primarily active (adjusted independent of
mass scaling), or a mixture of both (Table 1).

For six traits, the responses to resource limitation were primarily passive responses, with effects no
longer evident when mass was included as covariate). This included two morphological traits (FR-
Den and TR-SRL) and four anatomical traits (LF-Ks, ST-CTh, ST-XTh, and TR-CTh). For these traits, the
observed response to light limitation primarily was associated with smaller plant size.

For 13 traits, the responses were primarily active, with effects not changing when mass was included
in the model, indicating that the response was still evident after correcting for plant size. This
included 3 allocational traits (ST-MF, FR-MF, and TR-MF), six morphological traits (LF-Th, LF-SLA, LF-

Den, ST-Den, FR-Den, and FR-SRL) and four anatomical traits (LF-PT, LF-ST, FR-CT, and FR-SD). For
these traits, resource limitation effects were not associated with smaller plant size.

For 11 traits, the responses were a mixture of passive and active responses to resource limitation,
with the effects differed somewhat when mass was included in the model, indicating that scaling
with mass accounted for a portion of the response. This included 3 allocational traits (LF-MF,
RatioLF-FRmass, and RatioSLA-SRL), four morphological traits (ST-Dia, ST-Hgt, TR-Dia, and TR-Den)
and four anatomical traits (ST-VTh, ST-Ks, TR-SDia, and TR-Ks).

<u>How do traits compare for magnitude of relative plasticity and what role does size scaling of traits</u>
<u>play in this?</u>

The magnitude and direction of trait plasticity in response to resource limitation was assessed with a
relative distance plasticity index (RDPI, Figure 4a). The RDPI for at least one of the treatments was
significantly different from control for all of the traits expect FR-Ks. Visually comparing RDPI per trait
category showed that allocational traits had the greatest plastic responses, followed by morphology,
and then anatomy. Although the treatment that combined light and nutrient limitation ("shade-poor"
in Figure 4a) resulted in the largest decline in biomass, it was not consistently the treatment that
induced the greatest magnitude of plasticity across all of the traits.

When trait responses to resource limitation were corrected for plant mass (Figure 4b), the extent to
which RDPI values were affected was variable. Since this RDPI analysis is based on genotype means
instead of individual plants there are slight differences in the extent of active vs passive responses
though results broadly line up with the prior analysis (Table 1). For traits where responses were
designated as primarily passive (Table 1), correcting for plant mass resulted in substantially lower
RDPI (e.g., comparing RDPI in 4b to 4a for TR-SRL, LF-Ks, and ST-XT). For traits where responses were
designated as primarily active, correcting for plant mass had little effect on RDPI (e.g., comparing 4b
to 4a for ST-MF, FR-MF, LF-SLA, and LF-PT). For traits where responses were designated as a
combination of active and passive, correcting for plant mass had a variable effect on RDPI. For some
traits, RDPI was reduced for at least one treatment when mass was factored out (e.g., LF-MF,
RatioLF-FRmass, and ST-Dia) indicating that the passive and active components that response were
complementary in direction, resulting in a greater overall magnitude. However, for a few traits such
as ST-Hgt, ST-Ks and TR-Ks, correcting for plant mass resulted in a greater RDPI, indicating active and
passive responses to be in opposite directions, or antagonistic (Supplemental Figure S2).

<u>Do traits show a coordinated shift due to resource limitation across all organs and what role does</u>
<u>size scaling of traits play in this?</u>

When the genotypes means for the observed traits in all three traits categories (allocation,
morphology, and anatomy) and all treatments were included in a PCA the correlations among many
of the traits become evident. PC1 and PC2 explained 34.4% and 13.4% of the variation, respectively
(Figure 5a). Light limitation resulted in a shift of a key set of traits to be more acquisitive along the
first axis, with higher LF-SLA, TR-SRL, and FR-SRL under shade associated with lower values of other
morphological traits (LF-Den, ST-Dia, TR-Dia, TR-SDia) and lower anatomical trait values (ST-XTh, ST-
VTh). Nutrient limitation resulted in a smaller shift in traits, oriented more on PC2, and dominated
by higher TR-MF, TR-Den, and TR-Ks and lower RatioSLA-SRL, ST-Hgt, FR-Dia, FR-SDia (Suplemental
Figure S4).

When the trait values corrected for mass were included in a PCA, reflecting primarily active
responses, the correlations among traits changed and less of the variation was explained (PC1 and
PC2 explained 19.7% and 17.0% of the variation, respectively). Responses to light and nutrient

limitation were more similar in scale, although still resource specific (Figure 5b). Thus, while the
passive mass-effect of light limitation might be greater, the active responses to above and below
ground resource limitation were similar in magnitude.

Discussion

Here we assessed the direction, magnitude, and coordination of cultivated *H. annuus* responses to
light or/and nutrient limitation for traits in multiple categories (allocation, morphology and anatomy)
and across all organ (leaves, stem, and roots), using plants harvest at a similar age and
developmental stage. We found that more biomass was allocated to the organs acquiring the most
limiting resource (roots for nutrient stress and leaves for shade) and that there were additional
morphological and anatomical trait adjustments that generally led to more acquisitive trait values
(e.g. higher LF-SLA in shade and higher FR-SRL in nutrient stress). We explored the role of mass-
scaling in these responses by looking at the effect of including whole plant dry mass as a covariate to
correct for plant size. The trait responses to resource limitations varied in the extent to which they
were primarily passive (i.e., associated with lower biomass of stressed plants), primarily active (not
or only marginally associated with lower biomass of stressed plants) or a combination thereof. The
variable magnitude and direction of active plasticity played an appreciable role in shaping the
individual trait responses and the overall coordinated response that was unique to each resource
limitation.

<u>Allocation, morphology, and anatomy responses to above and below ground resource limitations</u>

Light limitation resulted in increased relative investment in aboveground plant parts and nutrient
limitation resulted in increased relative investment in belowground parts, consistent with the
expectation that plasticity serves to optimize/maximize the uptake of the most limiting resource
(Figure 1 and Table 1) (Bloom, Chapin, & Money, 1985; Shipley & Meziane 2002; Poorter et al., 2012).
This allocation pattern has been confirmed in other species and growth forms (e.g., grasses,
Siebenkäs, Schumacher, & Roscher, 2015; shrubs, Valladares, Wright, Lasso, Kitajima, & Pearcy, 2000;
trees, Reich, Tjoelker, Walters, Vanderklein, & Bushena, 1998, Poorter et al. 2012; Kramer-Walter &
Laughlin, 2017). Additionally, morphological and anatomical adjustments resulted in a higher LF-SLA
and FR-SRL, under light and nutrient limitations, respectively, decreasing the ratio of mass invested
to resource uptake potential (Figures 2,3). Anatomical adjustments in leaf and root, such as
decreased LF-PTh and FR-CTh, also reduced tissue metabolic and maintenance costs (Guo, Cao, & Xu,
2006, Galindo-Castañeda, Brown, & Lynch, 2018; Jaramillo, Nord, Chimungu, Brown, & Lynch, 2013).
Thus, for *H. annuus*, greater capacity for resource acquisition was coupled with reduced costs, and
this was achieved by a combination of passive and active plasticity.

The traits varied in the extent to which passive plasticity played a role in explaining the observed
response to resource limitations (Figure 4 and Table 2). The response to both light and nutrient
limitation appeared primarily passive for several morphological and anatomical traits, but not for
any of the allocational responses. And, the extent of passive adjustment differed by resource. Similar
to Poorter et al., 2012, we found that responses to light limitation were more often passive than for
nutrient limitation. For many traits there was evidence of a mixture of passive and active responses,
which could be complementary or antagonistic and thus influence the magnitude of the
overall/observed response (Figure 4, Supplemental figure S2). However, after correcting for plant
mass, treatment effects remained significant for all of the allocation traits and many morphological
and anatomical traits, such as LF-SLA (Freschet, Violle, Bourget, Scherer-Lorenzen, & Fort, 2018;
Reich, 2018; this study), FR-SRL, LF-PTh and FR-CTh (this study), indicating active adjustments of
these key traits for obtaining the most limited above or belowground resource.

 <u>Magnitude and direction of plasticity</u>

Consistent with other resource limitation studies, the observed plasticity of mass allocational traits
was largest, followed by morphological traits (Kramer-Walter & Laughlin, 2017, Valladares, Wright,
Lasso, Kitajima, & Pearcy, 2000), and smallest in anatomical traits (Catoni, Gratani, Sartori, Varone, &
Granata, 2015; Cai, Ji, Yan, Jiang, & Fang, 2017; Xu, Wang, Liu, Lu, & Guo, 2015). It should be noted
that differences in the extent of plasticity for different categories of traits may be species specific or
based on the traits included. For example, oak (*Quercus robur)* seedlings were more plastic in
physiological traits under shade, yet beech (*Fagus sylvatica)* was more plastic in morphological traits
(Valladares et al., 2002). Differences among trait categories for the magnitude of plasticity may
reflect an inherent hierarchy originating from the internal component traits of any specific organ.
For example, leaf thickness is highly correlated with palisade parenchyma (Scoffoni et al.*,* 2015) and
root diameter with cortex and/or stele thickness (Kong et al., 2014). Thus, small shifts in individual
component anatomical traits could add up to larger shifts in morphological traits which in turn affect
allocational traits.

For traits where there was evidence of a mixture of passive and active plasticity, the alignment of
both components (i.e. complementary or antagonistic in direction) affected the direction and
magnitude of the observed response (Figure 4, Supplemental Figure S2). For RatioLF-FRmass, the
passive and active responses to light limitation combined to enhance the relative investment in
leaves, increasing the capacity for light acquisition. For other traits, correcting for mass revealed
substantive active plasticity that was antagonistic to passive plasticity. For example, the active
plasticity for ST-Hgt, and ST-Ks and TR-Ks was of greater magnitude than the observed response. The
active adjustments counteracted the passive scaling with mass and resulted in shaded plants with
higher conducting capacity (TR-Ks, ST-Ks) as well as high axial transportation distance and total
transpiring surface (ST-Hgt and LF-SLA). This suggests that a coordinated increase in axial root and
stem hydraulic transport offset greater resistance due to a longer transportation distance (Plavcová
& Hacke, 2012), facilitating efficient movement of carbon to roots and nutrients and water to leaves
(Maurel, Simonneau, & Sutka, 2010; Rodríguez-Gamir, Primo-Millo, & Forner-Giner, 2016; Wahl,
Ryser, & Edwads, 2001). Thus, both passive and active trait adjustments to resource limitation serve
to increase resource uptake capacity and maintain (optimal) functioning of altered organs.

Where passive and active trait adjustments are antagonistic this could indicate a departure from
"safe" trait values under non-limiting conditions. For example, increased ST-Ks may come at the cost
of increased risk of xylem embolism in shaded plants (Tyree & Zimmermann, 2002). The anatomical
dataset collected in this work provides an excellent resource for anatomical water flow models
(Couvreur et al., 2018) to further shed light on the consequences of these anatomical trait
adjustments for plant hydraulics and how plants balance resource uptake demands with stress
safety margins.

<u>Coordinated trait shifts</u>

Resource limitation led to a coordinated shift amongst many correlated traits, with an overall shift
towards greater potential for resource acquisition, consistent with other resource manipulation
studies for commonly measured traits (Freschet, Swart, & Cornelissen, 2015). The coordinated trait
shifts differed by limiting resource, with light limitation dominated by higher LF-SLA and associated
with lower FL-Den and shifts in other stem and tap root traits, while nutrient limitation dominated
by thinner FR-SDia, FR-Dia, and TR-MF and associated with other root and stem traits. (Figure 5 and
Table 1). Correcting traits for plant size altered the magnitude of responses and relationships among
traits, and revealed that the active effects of light and nutrient limitation were still resource specific,

but more similar in magnitude.

The coordinated trait plasticity in response to resource limitations was generally a resource specific
shift in traits towards more acquisitive strategy, that was not evident across all organs. It is
interesting to note that these patterns are not consistent with the "whole plant economic spectrum"
expectation of correlated evolution of conservative traits across all organs and resource limitations
(Reich 2014, Díaz et al., 2016). This plasticity is an example of extensive responses to resource
limitations and other abiotic or biotic factors that undoubtedly contribute to variation captured in
broader surveys of plants specialized in different habitats (Anderegg et al., 2018). As such, these
findings contribute to ongoing efforts to understand the scale dependence of trait covariation and
the role that trait plasticity plays at both ecological and evolutionary scales.

<u>Conclusions</u>
Few studies have assessed the phenotypic response of whole plants in terms of biomass allocation,
organ morphology and anatomy simultaneously, and even less under multiple resource limitation.
Here, our research demonstrates that major traits from all three categories shift in response to
resource limitation. Many of these shifts are passive, i.e. scaling with decreased mass of plants under
stressed conditions. Moreover, these passive shifts extend well beyond mass allocational traits and
include both morphological and anatomical traits. The magnitude and direction of individual traits
responses, and their coordination are driven not only by passive scaling with plant mass, but by an
additional active component that can be complementary or antagonistic to the effect of mass scaling.
Thus, for a full understanding of plants response to environmental stress both this passive and active
plasticity needs to be taken into account.

Acknowledgements

We would like to thank M. Boyd and K. Tarner for assistance with plant growth and data collection, K.
Bettinger, S. Chhajed, G. Manning, J. Parrilli, N. Reisinger and J. Kobylanski for assistance with
experimental setup and data collection, and the greater Donovan lab group for comments that
improved this study and manuscript. Additionally, we would like to thank the reviewers for their
time and constructive comments. This work was financially supported by grant NSF1444522 to LAD
and a China Scholarship Council (CSC) grant to YW.

Author contributions

430 YW, AT, and LD designed the experiment. YW carried out the experiment and took all measurements.
431 YW and AT analyzed the results. YW wrote the initial manuscript draft with revisions by AT and LD.
432 All authors contributed to manuscript revision, and read and approved the submitted version

Figure and table captions

Table 1. Trait response to a factorial of light and nutrient limitation. Light vs Shade designates direct
sun vs 50% shade treatment. Rich vs Poor designates high nutrients vs 10% nutrient concentration
treatment. For each trait, the ANOVA effects of genotype (G), light (L) and nutrient (N) stresses and
their interactions (L×N) are presented, without (w/o) or with plant total biomass (Mass) included as
a covariate: ***: $p < 0.001$, **: $p < 0.01$, *: $p < 0.05$. Based on the difference between the models with
and without biomass as covariate we assigned a treatment response as either passive (primarily
scaling with mass), active (adjusted independent of mass), or a combination thereof. For each
treatment combination, the mean±SE of 10 genotypes (estimated marginal mean from the ANOVA,
based on 5-6 replicates per genotype) is presented for each trait. Different lower case letters
represent significant ($p < 0.05$) Tukey post hoc differences among treatments for each trait from the

ANOVA without biomass as a covariate.

Figure 1. Biomass and mass allocation of leaf, stem, tap root, and fine root tissue across a factorial of
446   light and nutrient limitation. Light/Shade, direct sun vs 50% shade. Rich/poor, high nutrients vs 10%
447   nutrient concentration. Whole plant biomass (mean±SE) of 10 genotypes (estimated marginal mean
448   from the ANOVA, based on 5-6 replicates per genotype), stacked by average tissue contribution.
449   Different lower-case letters represent significant (p<0.05) Tukey post hoc differences between
450   treatments for whole plant biomass.

Figure 2. Leaf and fine root morphology across a factorial of light and nutrient limitation.
Light/Shade, direct sun vs 50% shade. Rich/poor, high nutrients vs 10% nutrient concentration. **(a)**
specific leaf area (LF-SLA, cm^2 g^{-1}), **(b)** leaf mass fraction (LF-MF, %), **(c)** specific root length (FR-SRL,
cm g^{-1}), **(d)** fine root mass fraction (FR-MF, %), **(e)** ratio of leaf area to root length (RatioSLA-SRL), **(f)**
ratio of leaf mass to fine root mass (RatioLF-FRmass). Points indicate genotype (n=10) mean (n=5-6)
at a given treatment. Boxplots show distribution of values. Different letters represent significant
(p<0.05) Tukey post hoc differences between treatments for each trait.

Figure 3. Typical anatomical structure of leaf, stem, tap root and fine root of genotype HA $^-$ across a
459   factorial of light and nutrient limitation. Light/Shade, direct sun vs 50% shade. Rich/poor, high
460   nutrients vs 10% nutrient concentration. PA, palisade tissue; SP, spongy tissue; CO, cortex; VB,
461   vascular bundle; XY, xylem; ST, stele.

Figure 4. Trait plasticity in response to resource limitation. Points indicate the average (n=10)
relative distance plasticity index (RDPI) in response to light limitation (triangles, Shade-Rich) or
nutrient limitation (circles, Light-Poor), or combined (squares, Shade-Poor). Symbol fill represents a
T-test significance (p<0.05) of RDPI being different from zero (filled symbols) or not (open symbols).
Traits are ordered based on their treatment response, primarily passive (scaling with mass),
primarily active (adjusted independent of mass), or a combination thereof. **(a)** RDPI values taken
from base measurements. **(b)** RDPI values when correcting trait values for biomass. Trait
abbreviations as in Table 1.

Figure 5. Principal component analysis (PCA) of leaf, stem and roots traits in response to resource
471   limitations. Light/Shade, direct sun vs 50% shade. Rich/poor, high nutrients vs 10% nutrient
472   concentration. Panels represent **(a)** trait values as measured, **(b)** trait values after correcting for size
473   scaling (allometric effects). Different letters indicate significant Hotellings-T test, after Bonferroni
474   correction, between treatments on the first two principal components. Trait abbreviations and units
475   as in Table 1.

Supplemental materials

Supplemental Table S1. List of genotypes used in this study, it's common name, the corresponding
478   plant ID from the USDA GRIN Database for each genotype, and the market type of each genotype.

Supplemental Table S2. Genotype means for all traits discussed

Supplemental Figure S1. Comparison of light levels between shaded and unshaded treatment. On
481   average the low-light treatment received 50% of the photon flux density of the high-light treatment.
482   Light intensities in each treatment were measured with a handheld light meter (LI-189; LI-COR,
483   Lincoln, NE). Readings were taken from 8:00 AM to 7:00 PM. The light sensor was held near the soil
484   level in each plot. We presented representative data from May 15th, a cloud-free day.

Supplemental Figure S2. Effect of accounting for size scaling in traits when calculating relative

distance plasticity index (RDPI). RDPI changes are shown per treatment (Light/Shade, full sun/50%
shade; Rich/Poor, full nutrients/10% nutrients) and whether RDPI measures were positive or
negative when initially compared to control (Light-Rich). For positive RDPI values (traits that increase
in trait value with stress) a positive change when taking size scaling into account shows that size
scaling decreased apparent (as measured initially) plasticity (i.e taking size scaling into account
shows increased plasticity). A negative change on the other hand shows that the apparent plasticity
was enhanced by size scaling (i.e. taking size scaling into account shows decreased plasticity. For
negative RDPI values (traits that decrease in trait value with stress) the effect of the sign of change is
reversed. Symbols indicate whether RDPI significance (different from zero) is gained (circles), lost
(squares), or remains unchanged (triangles, when taking biomass into account.

Supplemental Figure S3. Scaling relationship between biomass and trait value among and between
genotypes and treatments. **(a)** Scaling relationships between individual plants per genotypes and
treatment (dotted lines) and **(b)** scaling relationships between genotypes per treatment.

Supplemental Figure S4. Trait contributions to principal components of figure 5.

References

Agrawal. A.A. (2020). A scale-dependent framework for tradeoffs, syndromes, and specialization in
organismal biology. Ecology, 101, e02924.

Anderegg, L.D.L., Berner, L.T, Badgley, G., Sethi, M.L., Law, B.E., HilliRisLambers, J. (2018). Within-
species patterns challenge our understanding of the leaf-economics spectrum. *Ecology Letters, 21,*
734-744

Bates, D., Maechler, M., Bolker, B., Christensen, R.H.B., Singmann, H., Dai, B., Scheipl, F.,
Grothendieck, G. & Green, P. (2018). Package 'lme4'.

Bloom, A.J., Chapin, F.S. & Mooney, H.A. (1985) Resource limitation in plants - an economic analogy.
Annual Review of Ecology and Systematics, *16*, 363-392.

Bowsher, A.W., Mason, C.M. Goolsby, E.W. & Donovan, L.A. (2016). Fine root tradeoffs between
nitrogen concentration and xylem vessel traits precludes unified whole-plant resource strategies
in *Helianthus*. *Ecology and Evolution, 6,* 1016-1031

Bowsher, A.W., Shelby, K.C., Ahmed, I., Krall, E., Reagan, D.J., Najdowski, M.N. & Donovan, L.A. (2017).
Genotype Rankings for Nutrient Stress Resistance are Unrelated to Stress Severity in Cultivated
Sunflower (*Helianthus annuus* L.). *Journal of Agronomy and Crop Science*, *203*, 241-253.

Cai, Q., Ji, C., Yan, Z., Jiang, X. & Fang, J. (2017). Anatomical responses of leaf and stem of *Arabidopsis*
thaliana to nitrogen and phosphorus addition. *Journal of Plant Research*, *130*, 1035-1045.

Catoni, R., Gratani, L., Sartori, F., Varone, L. & Granata, M.U. (2015). Carbon gain optimization in five
broadleaf deciduous trees in response to light variation within the crown: correlations among
morphological, anatomical and physiological leaf traits. *Acta Botanica Croatica*, *74*, 71-94.

Chapin, F.S.I. (1991). Integrated responses of plants to stress a centralized system of physiological
responses. *Bioscience*, *41*, 29-36.

Chatelet, D.S., Clement, W.L., Sack, L., Donoghue, M.J. & Edwards, E.J. (2013). The evolution of
photosynthetic anatomy in *Viburnum* (Adoxaceae). *International Journal of Plant Sciences*, *174*,
1277-1291.

Coleman, J.S., McConnaughay, K.D.M. & Ackerly, D.D. (1994). Interpreting phenotypic variation in
plants. *Trends in Ecology & Evolution*, *9*, 187-191.

Couvreur, V., Faget, M., Lobet, G., Javaux, M., Chaumont, F. & Draye, X. (2018). Going with the flow:
multiscale insights into the composite nature of water transport in roots. *Plant Physiology*, *178*,
1689-1703.

Díaz, S., Kattge, J., Cornelissen, J.H.C., Wright, I.J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C.,
Prentice, I.C., Garnier, E., Bönisch, G., Westoby, M., Poorter, H., Reich, P.B., Moles, A.T., Dickie, J.,
Gillison, A.N., Zanne, A.E., Chave, J., Wright, J.S., Sheremet'ev, S.N., Jactel, H., Baraloto, C.,
Cerabolini, B., Pierce, S., Shipley, B., Kirkup, D., Casanoves, F., Joswig, J.S., Günther, A., Falczuk, V.,
Rüger, N., Mahecha, M.D. & Gorné, L.D. (2016). The global spectrum of plant form and function.
Nature, *529*, 167-171.

Donovan, L.A., Mason, C.M., Bowsher, A.W., Goolsby, E.W. & Ishibashi, C.D.A. (2014). Ecological and
evolutionary lability of plant traits affecting carbon and nutrient cycling. *Journal of Ecology*, *102*,
302-314.

Eissenstat, D.M. & Achor, D.S. (1999). Anatomical characteristics of roots of citrus rootstocks that
vary in specific root length. *New Phytologist, 141*, 309-321.

Fortunel, C., Fine, P.V.A. & Baraloto, C. (2012). Leaf, stem and root tissue strategies across 758
Neotropical tree species. *Functional Ecology, 26*, 1153-1161.

Fox, J., Weisberg, S., Price, B., Adler, D., Bates, D., Baud-Bovy, G., Bolker, B., Ellison, S., Firth, D.,
Friendly, M., Gorjanc, G., Graves, S., Heriberger, R. & Laboissiere, R. (2018) Package 'car'.

Freschet, G.T., Swart, E.M. & Cornelissen, J.H.C. (2015). Integrated plant phenotypic responses to
contrasting above- and below-ground resources: key roles of specific leaf area and root mass
fraction. *New Phytologist, 206*, 1247-1260.

Freschet, G.T., Violle, C., Bourget, M.Y., Scherer-Lorenzen, M. & Fort, F. (2018). Allocation,
morphology, physiology, architecture: the multiple facets of plant above- and below-ground
responses to resource stress. *New Phytologist, 219*, 1338-1352.

Galindo-Castañeda, T., Brown, K.M. & Lynch, J.P. (2018). Reduced root cortical burden improves
growth and grain yield under low phosphorus availability in maize. *Plant, Cell & Environment, 41*,
1579-1592.

Gedroc, J., McConnaughay, K.D.M. & Coleman, J.S. (1996). Plasticity in root/shoot partitioning:
optimal, ontogenetic, or both?. *Functional Ecology, 1*, 44-50.

Gu, J., Xu, Y., Dong, X., Wang, H. & Wang, Z. (2014). Root diameter variations explained by anatomy
and phylogeny of 50 tropical and temperate tree species. *Tree Physiology, 34*, 415-425.

Guo, D., Xia, M., Wei, X., Chang, W., Liu, Y. & Wang, Z. (2008). Anatomical traits associated with
absorption and mycorrhizal colonization are linked to root branch order in twenty-three Chinese
temperate tree species. *New Phytologist, 180*, 673-683.

Guo, X.R., Cao, K.F. & Xu, Z.F. (2006). Acclimation to irradiance in seedlings of three tropical rain
forest Garcinia species after simulated gap formation. *Photosynthetica, 44*, 193-201.

Jaramillo, R.E., Nord, E.A., Chimungu, J.G., Brown, K.M. & Lynch, J.P. (2013). Root cortical burden
influences drought tolerance in maize. *Annals of Botany, 112*, 429-437.

John, G.P., Scoffoni, C., Buckley, T.N., Villar, R., Poorter, H. & Sack, L. (2017). The anatomical and
compositional basis of leaf mass per area. *Ecology Letters, 20*, 412-425.

Kong, D., Ma, C., Zhang, Q., Li, L., Chen, X., Zeng, H. & Guo, D. (2014). Leading dimensions in
absorptive root trait variation across 96 subtropical forest species. *New Phytologist, 203*, 863-872.

Kramer-Walter, K.R. & Laughlin, D.C. (2017). Root nutrient concentration and biomass allocation are
more plastic than morphological traits in response to nutrient limitation. *Plant and Soil, 416*, 539-
550.

Kramer-Walter, K.R., Bellingham, P.J., Millar, T.R., Smissen, R.D., Richardson, S.J. & Laughlin, D.C.
(2016). Root traits are multidimensional: specific root length is independent from root tissue
density and the plant economic spectrum. *Journal of Ecology, 104*, 1299-1310.

Lenth, R., Singmann, H., Love, J., Buerkner, P., Herve, M. & Abd Allah, E.F. (2018). Package 'emmeans'.

Mandel, J.R., Dechaine, J.M., Marek, L.F. & Burke, J.M. (2011). Genetic diversity and population
structure in cultivated sunflower and a comparison to its wild progenitor, *Helianthus annuus* L.
Theoretical and Applied Genetics, *123*, 693-704.

Masalia, R.R., Temme, A.A., Torralba, N.D.L. & Burke, J.M. (2018). Multiple genomic regions influence
root morphology and seedling growth in cultivated sunflower (*Helianthus annuus* L.) under well-
watered and water-limited conditions. *PLOS One*, *13*, e204279.

Mason, C.M. & Donovan, L.A. (2015) Evolution of the leaf economic spectrum in herbs: evidence
from environmental divergences in leaf physiology across *Helianthus*. *Evolution, 69*, 2705-2720.

Maurel, C., Simonneau, T. & Sutka, M. (2010). The significance of roots as hydraulic rheostats. *Journal*
of Experimental Botany, *61*, 3191-3198.

McConnaughay, K.D.M. & Coleman, J.S. (1999). Biomass allocation in plants: ontogeny or optimality a
test along three resource gradients. *Ecology*, *8*, 2581-2593.

Medeiros, J.S., Burns, J.H., Nicolson, J., Roger, L. & Valverde-Barrantes, O. (2017). Decoupled leaf and
root carbon economics is a key in the ecological diversity and evolutionary divergence of
deciduous and evergreen lineages of genus *Rhodendron*. *American Journal of Botany*, *104*, 803-
816.

McElrone, A.J., Pockman, W.T., Martínez-Vilata, J., & Jackson, R.B. (2004). Variation in xylem
structure and function in stems and roots of trees to 20 m depth. *New Phytologist, 163,* 507-517.

Muir, C.D., Conesa, M.A., Roldán, E.J., Molins, A. & Galmés, J. (2017). Weak coordination between
leaf structure and function among closely related tomato species. *New Phytologist, 213,* 1642-
1653.

Nambeesan, S.U., Mandel, J.R., Bowers, J.E., Marek, L.F., Ebert, D., Corbi, J., Rieseberg, L.H., Knapp,
S.J. & Burke, J.M. (2015). Association mapping in sunflower (*Helianthus annuus* L.) reveals
independent control of apical vs. basal branching. *BMC Plant Biology*, *15*, 84.

Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E.J., Mathesius, U., Poot, P.,
Purugganan, M.D., Richards, C.L., Valladares, F. & van Kleunen, M. (2010). Plant phenotypic
plasticity in a changing climate. *Trends in Plant Science*, *15*, 684-692.

Osnas, J.L.D., Lichstein, J.W., Reich, P.B. & Pacala, S.W. (2013). Global leaf trait relationships: mass,
area, and the leaf economics spectrum. *Science*, *340*, 741-744.

Pilote, A.J. & Donovan, L.A. (2016). Evidence of correlated evolution and adaptive differentiation of
stem and elf functional traits in the herbaceous genus, *Helianthus. American Journal of Botany,*
103, 2096-2014.

Plavcová, L. & Hacke, U.G. (2012). Phenotypic and developmental plasticity of xylem in hybrid poplar
saplings subjected to experimental drought, nitrogen fertilization, and shading. *Journal of*
Experimental Botany, *18*, 6481-6491.

Poorter, H., Niinemets, Ü., Ntagkas, N., Siebenkäs, A., Mäenpää, M., Matsubara, S. & Pons, T.L. (2019).
A meta-analysis of plant responses to light intensity for 70 traits ranging from molecules to whole
plant performance. *New Phytologist*, *223*, 1073-1105.

Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P. & Mommer, L. (2012). Biomass allocation to
leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New
Phytologist, 193*, 30-50.

Reich, P.B. (2014). The world-wide 'fast-slow' plant economics spectrum: a traits manifesto. *Journal
of Ecology, 102*, 275-301.

Reich, P.B. (2018). Do plants increase resource acquisition potential in the face of resource shortfalls,
and if so, how? *New Phytologist, 219*, 1142-1144.

Reich, P.B., Tjoelker, M.G., Walters, M.B., Vanderklein, D.W. & Bushena, C. (1998). Close association
of RGR, leaf and root morphology, seed mass and shade tolerance in seedlings of nine boreal tree
species grown in high and low light. *Functional Ecology, 12*, 327-338.

Rice, S.A. & Bazzaz, F.A. (1989). Quantification of plasticity of plant traits in response to light intensity
- comparing phenotypes at a common weight. *Oecologia, 78*, 502-507.

Rico, C., Pitterman, J., Polley, H.W., Aspinwell, M.J., & Fay, P.A. (2013). The effect of subambient to
elevated CO_2 concentration on vascular function in *Helianthus annuus*: Implications for plant
response to climate change. *New Phytologist, 199*, 956-965.

Rodríguez-Gamir, J., Primo-Millo, E. & Forner-Giner, M.Á. (2016) An integrated view of whole-tree
hydraulic architecture. does stomatal or hydraulic conductance determine whole tree
transpiration? *PLOS One, 11*, e155246.

Ryser, P. & Eek, L. (2000). Consequences of phenotypic plasticity vs. interspecific differences in leaf
and root traits for acquisition of aboveground and belowground resources. *American Journal of
Botany, 3*, 402-411.

Scoffoni, C., Kunkle, J., Pasquet-Kok, J., Vuong, C., Patel, A.J., Montgomery, R.A., Givnish, T.J. & Sack, L.
(2015). Light-induced plasticity in leaf hydraulics, venation, anatomy, and gas exchange in
ecologically diverse Hawaiian lobeliads. *New Phytologist, 207*, 43-58.

Shipley, B. & Meziane, D. (2002). The balanced-growth hypothesis and the allometry of leaf and root
biomass allocation. *Functional Ecology, 16*, 326-331.

Siebenkäs, A., Schumacher, J. & Roscher, C. (2015). Phenotypic plasticity to light and nutrient
availability alters functional trait ranking across eight perennial grassland species. *AoB Plants, 7.*
plv029.

Sperry, J.S., Hacke, U.G. & Pittermann J. (2006). Size and function in conifer tracheids and
angiosperm vessels. *American Journal of Botany, 93*:1490-1500.

Sugiura, D. & Tateno, M. (2011). Optimal leaf-to-root ratio and leaf nitrogen content determined by
light and nitrogen availabilities. *PLOS One, 6*, e22236.

Temme, A.A., Kerr, K.L., & Donovan, L.A. (2019) Vigour/tolerance trade-off in cultivated sunflower
(*Helianthus annuus*) response to salinity stress is linked to lead elemental composition. *Journal of
Agronomy and Crop Science, 205*, 508-518.

Tyree, M.T., & Zimmermann, M.H. (2002). Xylem structure and the ascent of sap. Springer.

Tyree, M.T. & Ewers, F.W. (1991) The hydraulic architecture of trees and other woody plants. *New
Phytologist, 119*, 345-360.

Valladares, F., Chico, J., Aranda, I., Balaguer, L., Dizengremel, P., Manrique, E. & Dreyer, E. (2002). The
greater seedling high-light tolerance of *Quercus robur* over *Fagus sylvatica* is linked to a greater
physiological plasticity. *Trees*, *16*, 395-403.

Valladares, F., Gianoli, E. & Gómez, J.M. (2007). Ecological limits to plant phenotypic plasticity. *New
Phytologist*, *176*, 749-763.

Valladares, F., Sanchez-Gomez, D. & Zavala, M.A. (2006) Quantitative estimation of phenotypic
plasticity: bridging the gap between the evolutionary concept and its ecological applications.
Journal of Ecology, *94*, 1103-1116.

Valladares, F., Wright, S.J., Lasso, E., Kitajima, K. & Pearcy, R.W. (2000). Plastic phenotypic response
to light of 16 congeneric shrubs from a Panamanian rainforest. *Ecology*, *81*, 1925-1936.

Van Kleunen, M. & Fisher, M. (2005) Constraints on the evolution of adaptive phenotypic plasticity in
plants. *New Phytologist, 166,* 49-60.

Wahl, S., Ryser, P. & Edwards, P. (2001). Phenotypic plasticity of grass root anatomy in response to
light intensity and nutrient supply. *Annals of Botany*, *88*, 1071-1078.

Weiner, J. (2004). Allocation, plasticity and allometry in plants. *Perspectives in Plant Ecology,
Evolution and Systematics*, *6*, 207-215.

Wickham, H., Chang, W., Henry, L., Pederson, T.L., Takahashi, K., Wikle, C. & Woo, K. (2018). Package
'ggplot2'.

Wilson, P.J., Thompson, K. & Hodgson, J.G. (1999). Specific leaf area and leaf dry matter content as
alternative predictors of plant strategies. *New Phytologist*, *143*, 155-162.

Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-Bares, J.,
Chapin, T., Cornelissen, J.H., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., Gulias, J., Hikosaka, K.,
Lamont, B.B., Lee, T., Lee, W., Lusk, C., Midgley, J.J., Navas, M.L., Niinemets, U., Oleksyn, J., Osada,
N., Poorter, H., Poot, P., Prior, L., Pyankov, V.I., Roumet, C., Thomas, S.C., Tjoelker, M,G,,
Veneklaas, E.J., Villar, R. (2004). The worldwide leaf economics spectrum. *Nature 428*, 821-827.
Xu, N., Wang, R., Liu, J., Lu, P. & Guo, W. (2015). Hierarchy of plasticity traits in responses of *Quercus
aliena* to light conditions and water availability. *Dendrobiology*, *74*, 169-180.