Salivary proteins of a gall-inducing aphid and their impact on early gene responses of susceptible and

resistant host-plant genotypes

Luis Portillo Lemus [1, 2] *, Jessy Tricard [1, 2] *, Jérôme Duclercq [2], Quentin Coulette [2], David Giron [3],

Christophe Hano [1], Elisabeth Huguet [3], Frédéric Lamblin [1], Anas Cherqui [2]°, Aurélien Sallé [1]°

[1] Laboratoire de Biologie des Ligneux et des Grandes Cultures, INRA, Université d'Orléans, 45067,

Orléans, France

[2] Ecologie et Dynamique des Systèmes Anthropisés, EDYSAN UMR CNRS-UPJV 7058, Université de

Picardie Jules Verne, Amiens, France

[3] Institut de Recherche sur la Biologie de l'Insecte, UMR 7261, CNRS/Université François-Rabelais de

Tours, Tours, France

* L. Portillo Lemus and J. Tricard should be considered joint first author

° A. Cherqui and A. Sallé should be considered joint senior author

Funding

This study has been supported by the Région Centre-Val de Loire Project no. 2014 00094521

(InsectEffect) coordinated by D. Giron.

Abstract

Plant manipulation by herbivores requires fine-tuned reprogramming of host metabolism, mediated by

effector molecules delivered by the parasite into its host. While plant galls may represent the epitome of

plant manipulation, secretomes of gall-inducers and their impact on host-plants have been rarely

studied. We characterized, with transcriptomic and enzymatic approaches, salivary glands and saliva of a

gall-inducing aphid, *Phloeomyzus passerinii*. Early responses to aphid saliva of plant genes belonging to

different metabolic and signaling pathways were assessed *in vivo*, with poplar protoplasts, and *in planta*,

in a heterologous *Arabidopsis* system. Several effectors potentially interfering with plant signaling have

been identified, including binding proteins, oxidoreductases, and phosphatidylinositol phosphate

kinases. Compatible interactions between protoplasts of a susceptible poplar genotype and the saliva of

P. passerinii led to an overall downregulation of defense-related genes while an upregulation was

observed during both incompatible interactions, with a resistant poplar genotype, and non-host

interactions, with the saliva of *Myzus persicae*, an aphid which does not feed on poplars. Compatible

interactions affected both auxin transport and homeostasis potentially leading to an intracellular

accumulation of auxin, which was further supported by *in planta* assays. Our results support the

hypothesis that effectors interfere with downstream signaling and phytohormone pathways.

Keywords

Effector, *Myzus persicae*, *Phloeomyzus passerinii*, plant-insect interaction, *Populus*, resistance, RT-qPCR

Acknowledgements

We thank Léa Fléchon for her assistance with transcriptome analyses.

1 Introduction

Nutritional imbalance and defense mechanisms are major hurdles hampering plant resources

exploitation by animals (Schoonhoven, Van Loon, & Dicke, 2005). To cope with this, some herbivores

have evolved strategies consisting in remodeling plant tissues to turn them into optimal substrates for

their development and fitness (Lieutier et al., 2017). The epitome of such plant manipulation is certainly

gall induction, triggering sometimes spectacular and complex tissue reorganization resulting in new plant

organs, within which parasites feed and grow (Stone & Schönrogge, 2003; Shorthouse, Wool & Raman,

2005; Giron, Huguet, Stone & Body, 2016). To adapt plant tissues to nutritional requirements of parasites

and to circumvent plant defenses, gall development implies fine-tuned metabolism reprogramming, and

direct modulations of both primary and secondary metabolisms of host plants have been reported for

gall-inducing organisms (Giron et al., 2016). More specifically, since they are important regulators of

plant growth, differentiation and defense, phytohormones are classical targets of plant-manipulating

organisms, and considered key factors intimately involved in the success or failure of gall differentiation

(Tooker & Helms, 2014; Giron et al., 2016).

Host metabolism hijacking by plant-manipulating organisms is considered to be mediated by effector

molecules delivered into the host-plant (Hogenhout & Bos, 2011; Giron et al., 2016). Effector proteins

secreted by herbivores in general have various functions, including suppression of plant defense,

alteration of plant development, and manipulation of plant resources (Hogenhout & Bos, 2011; Giron et

al., 2016). For gall-inducing organisms, similar functions are expected, especially disruption or diversion

of hormone-dependent pathways (Tooker & Helms, 2014) and inactivation of second messengers

involved in stress signaling like Ca^{2+}, reactive oxygen species (ROS) and extracellular ATP for instance

(Will, Tjallingii, Thönnessen & van Bel, 2007; Guiguet et al., 2016). While there is a growing literature

dealing with the identification of these effectors in herbivores, like aphids (e.g. Harmel et al., 2008;

Nicholson, Hartson & Puterka, 2012; Vandermoten et al., 2014; Boulain et al., 2018), few investigations

have been performed on secretions of gall-inducing organisms, with notable exceptions of the Hessian

fly (*Mayetiola destructor*, Say) and root-knot nematodes (*Meloidogyne* spp.) for instance (e.g. Zhao et al.,

2015; Favery, Quentin, Jaubert-Possamai & Abad, 2016). So far, the Hessian fly is the only gall-inducing

insect with a sequenced genome, and whose salivary gland transcriptome and proteome have been

studied (Stuart, Chen, Shukle & Harris, 2012; Zhao et al., 2015). Candidate genes with effector functions

have been detected and gene-for-gene interaction between insect strains and host-plant genotypes has

been demonstrated (Stuart et al., 2012; Zhao et al., 2015; 2016). How these effectors contribute to gall

induction is still unclear, but it is hypothesized that they interfere with downstream signaling and

phytohormone pathways (Zhao et al., 2016).

To validate the relevance of candidate effectors for plant-insect interactions, several approaches have

been used. *In planta* modulation of candidate effectors expression allowed to highlight impacts of

effector proteins on physiological traits and behavior of insects and plant responses (e.g. Mutti et al.,

2008; Atamian et al., 2013; Zhao et al., 2016). Alternatively, direct applications of oral secretions and

saliva infiltration into host plants also gave conclusive results (e.g. De Vos & Jander, 2009; Chaudhary,

Atamian, Shen, Briggs & Kaloshian, 2014). Nonetheless, probably because of their complex life cycles and

generally concealed endophytic development, very few functional validation experiments have been

performed with gall-inducing insects (e.g. Zhao et al., 2016). In-depth functional molecular approaches

would allow to unravel the mechanisms that contribute to plant manipulation by these organisms and

could possibly lead to identification of convergent mechanisms among them (Giron et al., 2016).

Here, we report the characterization of salivary proteins of *Phloeomyzus passerinii* Sign., a gall-inducing

aphid colonizing poplars (Sallé, Pointeau, Bankhead-Dronnet, Bastien & Lieutier, 2017). As this insect can

be easily handled under laboratory conditions, and its saliva can be collected on artificial medium, we

performed innovative functional validation assays to address the impact of the saliva of a gall-inducing

organism on gene responses of its host-plant. Our objectives were (i) to characterize with transcriptomic

and enzymatic approaches the salivary glands and saliva, respectively, of *P. passerinii*, (ii) to assess *in*

vivo, with a RT-qPCR approach, early responses of poplar genes belonging to different metabolic and

signaling pathways using protoplasts exposed to aphid saliva, and (iii) to investigate the impact of

salivary extracts on specific gene expression *in planta*, in a heterologous *Arabidopsis* system. For all these

steps comparative approaches have been used. The determined salivary proteins of *P. passerinii* were

compared with those of a sap-feeding aphid *Myzus persicae* (Sulzer) (Harmel et al., 2008; De Vos &

Jander, 2009) in order to identify proteins common to both aphid species as well as proteins specific to *P.*

passerinii. To get insight into how *P. passerinii* manipulates host metabolism and stress responses to

perform gall-induction, the impact of salivary extracts on gene expression has been assessed with poplar

genotypes either susceptible or resistant to *P. passerinii*, i.e. during compatible and incompatible

interactions, respectively. These poplar genotypes have also been exposed to salivary extracts of *M.*

persicae, which led to non-host interactions since *M. persicae* does not feed on poplar.

2 Materials and Methods

2.1 Plant and insect material

Phloeomyzus passerinii is a specialist of poplars, inducing open galls in the cortical parenchyma of its

host-trees and feeding on cell contents (Pointeau et al., 2012; Dardeau et al., 2014; Sallé et al., 2017).

During gall induction in susceptible hosts, cell multiplication in cortical parenchyma is visible one week

after the onset of aphid probing, and three weeks later thin-walled hypertrophied cells differentiate

while vacuolar phenolic compounds disappear from the galled tissues (Dardeau et al., 2014). In resistant

host genotypes, lignin, tannins and flavanols accumulate at the probing site one week after the onset of

probing and no gall differentiation occurs subsequently. As a consequence, *P. passerinii* cannot develop

on these genotypes (Dardeau et al., 2014). All individuals of *P. passerinii* used either for transcriptomic

analyses of salivary glands or saliva collection originated from the same monoclonal colony established

from an apterous parthenogenetic female, collected in 2013 in Brézé (France). The colony was

maintained in the laboratory on potted stem cuttings of I-214, a *Populus x canadensis* Moench.

genotype, under 20 ± 1 °C, 70 ± 10% relative humidity and 16/8 h light/dark cycles. The green peach

aphid, *Myzus persicae* is a sap-feeder and a generalist aphid. Nonetheless, its host range does not

include poplars, and preliminary establishment attempts confirmed that it cannot settle and develop on

poplars (data not shown). Therefore non-host interactions are expected between this aphid species and

poplar genotypes. Individuals of *M. persicae* originated from a monoclonal colony established from an

apterous parthenogenetic female collected in 1999 on a potato plant in Loos-en-Gohelle (France). The

colony was maintained under the same controlled conditions as *P. passerinii*, on turnips (Vilmorin).

For the protoplasts production, stem-cuttings of two *P. x canadensis* genotypes commonly planted in

France, I-214 and Koster, were used for the experiments. I-214 is highly susceptible to *P. passerinii*,

whereas aphids cannot settle on Koster which is consequently considered to be highly resistant to *P.*

passerinii (Sallé et al., 2017). Consequently, compatible and incompatible interactions are expected

between *P. passerinii* and I-214 and Koster, respectively, and non-host interactions should take place

between *M. persicae* and both poplar genotypes. Stem cuttings (ca. 25 cm long, 2 cm diameter) were

provided by the experimental nursery of Guéméné-Penfao (France). They were collected in the autumn

of 2016, and kept at 2°C, in dry conditions until use. In January 2017, the stem-cuttings were removed

from storage and planted in 0.4 L pots, filled with a sterile sand-compost (50:50) mixture (Klasmann

substrate 4 no. 267). The cuttings were then transferred to a growth chamber (20 ± 1°C, 70 ± 10%

relative humidity, 16/8 h light/dark photoperiod, 2.65 kLx, and watered three times a week).

For *in planta* functional validation of the effects of salivary proteins, two *Arabidopsis thaliana* (L.) Heynh.

transgenic lines were used, the auxin-responsive reporter *pIAA2*::GUS (Bishopp et al., 2011) and the

cytokinins-responsive reporter *pARR16*::GUS (European Arabidopsis Stock Centre). Seeds were sterilized

with chloral gas, sown in Petri dishes on 0.8% (w/v) agar with 1% (w/v) sucrose-containing 0.5 Murashige

and Skoog medium (MS), stored for 2 days at 4°C, and grown on vertically oriented plates in growth

chambers under a 16/8 h light/dark photoperiod at 18°C.

2.2 Salivary transcriptome

2.2.1 Sample collection, RNA isolation and *de novo* transcriptome assembly

About 500 adults of apterous parthenogenetic *P. passerinii* aphids were dissected to collect pairs of

salivary glands. Total RNA was extracted using the GeneJET RNA Purification kit (Thermo Fischer

Scientific), according to manufacturer's instructions. RNA was DNase treated using RNase-Free DNase Set

(Qiagen). RNA concentration was measured using the Qubit® RNA Assay Kit (Life Technologies) and a

Qubit® 2.0 Fluorometer (Invitrogen). Construction of cDNA-library and sequencing were performed by

Eurofins® Genomics using a MiSeq v3 Reagent Kit (600 Cycles PE, Illumina, USA) and a MiSeq sequencer

(Illumina). For the *de novo* transcriptome assembly, 15,453,942 pair-ended reads were sequenced and

assembled using Velvet (v1.2.10; Zerbino & Birney, 2008) and Oases (v0.2.8; Schulz, Zerbino, Vingron &

Birney, 2012) software tools (table S1). A multi-kmer approach was applied. Separate assemblies with

different kmer lengths have been conducted and the individual assemblies have been merged to a final

assembly. Kmer lengths of 69, 89, 109 and 129 were used. The separate assemblies were merged

following the filter1-CD-HIT-EST procedure proposed in Yang & Smith (2013). This Transcriptome

Shotgun Assembly project has been deposited at DDBJ/EMBL/GenBank under the accession

GHDF00000000. The version described in this paper is the first version, GHDF01000000.

2.2.2 Annotation, secreted proteins detection and identification

To perform comparisons with *M. persicae* the transcriptome of this aphid was retrieved on NCBI

(http://www.ncbi.nlm.nih.gov/genbank/, accession numbers: DW010205 - DW015017, EC387039 -

EC390992, EE570018 - EE572264, EE260858 - EE265165, ES444641 - ES444705, ES217505 - ES226848,

and ES449829 - ES451794). Salivary transcriptomes have been annotated using the pipeline described in

figure S1. Transcripts were first translated into amino acid sequences using Prodigal (v2.5; Hyatt et al.

2010). We then used the SignalP 4.0 Server (v4.1) to predict the presence of signal peptides and cleavage

sites in the amino acid sequences (Petersen, Brunak, von Heijne & Nielsen, 2011). To predict

transmembrane domains, we submitted each amino acid sequence with a signal peptide to the TMHMM

Server (v. 2.0; Ji et al., 2013). Putative proteins with a signal peptide and no transmembrane domain

were considered to be potential secreted proteins. The sequences of complete ORFs without signal

peptide were analyzed again with SecretomeP (v2.0; Bendtsen, Jensen, Blom, von Heijne & Brunak,

2004). To remove mitochondrial proteins with a signal peptide, which are not secreted in the saliva,

sequences were analyzed with TargetP (v1.1; Emanuelsson, Nielsen, Brunak, & von Heijne, 2000).

Likewise, to remove proteins of the endoplasmic reticulum with a signal peptide, sequences were

analyzed with PS-scan (Prosite pattern: PS00014), and with PredGPI (Pierleoni, Martelli & Casadio, 2008)

for glycosylphosphatidylinositol-anchor signals.

The remaining proteins were first mapped against the non-redundant protein sequences (nr) using

Blastp (v2.3.0, NCBI, accessed on 03/30/2016), with an E-value cutoff at 1^{e-3}. Protein domains were

annotated with Blast2Go (v3.3; Conesa et al., 2005), and InterProScan (v5.30-69; Jones et al., 2014).

Whenever possible, protein sequences were assigned to Gene ontology (GO) terms with an E-value

cutoff at 1^{e-6}, enzyme codes (EC) and KEGG pathways.

OrthoVenn (http://aegilops.wheat.ucdavis.edu/OrthoVenn; Wang et al., 2015) has been used to identify

orthologous proteins within and between salivary transcriptomes of the two aphids. Intraspecific

orthologous proteins are first grouped into clusters, which are then compared between species. Each

cluster has been annotated with the Uniprot database (http://www.uniprot.org; Pundir, Martin &

O'Donovan, 2017) and the nr peptide sequence database (NCBI, accessed on 03/30/2016).

To detect proteins orthologous to salivary effectors of aphids, protein sequences of known aphid

effectors, i.e. C002, ACE1, ACE2, ACYPI39568, ACYPI00346, MpC002, Mp1, Mp2, Mp42, Mp55, Me10,

Me23 (Guo et al., 2014; Jaouannet et al., 2014; Pan, Zhu, Luo, Kang & Cui, 2015), have been compared to

the salivary transcriptome of *P. passerinii* with Blastp (E-value ≤ 1^{e-3}).

2.3 Functional validation assays

2.3.1 Aphid saliva collection

Aphids secrete two types of saliva within their host-plants, liquid and solid saliva (Will, Steckbauer, Hardt

& van Bel, 2012). The solid saliva is secreted during probing. It hardens rapidly and forms a solid sheath

encasing aphid stylets within the host-plant, while the liquid saliva is secreted within cells and sieve

tubes (Will et al., 2012). Both types of saliva contain effectors (e.g. Will et al., 2012; Elzinga & Jander,

2013), and have been collected in our experiments. Because of the particular trophic substrates of *M.*

persicae and *P. passerinii* (i.e. sap and galled tissues, respectively), a special protocol has been used for

each species. The saliva of *P. passerinii* was collected after incubation of 30 to 40 individuals of 2^{nd} and

3^{rd} instars aphids on sachets of Parafilm$^{©}$ membranes containing an artificial diet (Cherqui & Tjallingii,

2000). The artificial diets were constituted by a disc of 0.5% (w/v) agar completed with 150 µL of 15%

(w/v) sucrose. The saliva of *M. persicae* was collected after incubation of 30 to 40 individuals, of 3^{rd} and

4^{th} instars, on artificial diet containing 120 µL of a 15% (w/v) sucrose as previously described by Cherqui

& Tjallingii (2000). Aphids were deposited in a feeding chamber during 24 h at 20°C, 60 ± 1% relative

humidity and a 16/8 h light/dark period with 2.65 kLx. Feeding chambers containing the artificial diets,

incubated in the absence of aphids, were used as control samples.

For *P. passerinii*, after 24 h aphid salivation, artificial diet discs were collected and transferred into 100

μL of TE buffer (10 mM Tris, 1 mM EDTA, pH 8). The salivary proteins were released from the artificial

diet according to Yang et al. (2010), with slight modifications. The tubes containing artificial diet discs

were frozen in liquid nitrogen for 1 min, immediately thawed at 70°C for 3 min and then centrifuged at

11,000 x g for 20 sec. To discard the excess of agar, salivary extracts were centrifuged in Sartorius tubes

with filters of 0.22 μm. The supernatant containing salivary proteins of *P. passerinii* were collected,

pooled and then stored at -20°C. For *M. persicae,* after 24 h salivation, aphid saliva was collected

according to Harmel et al. (2008). The artificial diet is collected containing soluble saliva. The solid saliva

was collected during the rinsing of each lower Parafilm membrane with TE buffer containing 0.1% (w/v)

of Tween 20 (TE/Tween). The extracts were centrifuged at 10,000 x g for 15 min. The salivary proteins in

the pellet were collected, pooled with the soluble saliva and then stored at -20°C.

The sample containing protein saliva extracts were concentrated using 2 mL Vivaspin[©] tube (Sartorius)

with 3kDa cut-off. The tubes were then centrifuged at 5,000 x g for 70 to 120 min according to sample

volumes, and proteins adhering to membranes were recovered by 100 μL of TE/Tween buffer. Control

samples were prepared with artificial diets from feeding chambers without aphids. The protein

quantification was performed by measuring absorbance at 280 nm with the NanoDrop[©] 1000

(ThermoScientific).

2.3.2 Enzyme activities

Several enzyme substrates were added to the previously described artificial diets with or without

agarose to detect enzymatic activities present in saliva excreted from the aphid. To visualize proteins in

the salivary sheaths, the lower Parafilm[©] membranes were stained by adding a drop of 0.01% (w/v)

Coomassie blue in 10% (v/v) glycerol for 2 h. Dihydroxyphenylalanine (DOPA), 0.1% (w/v) was added to

identify phenoloxidase activity (PO; catechol oxidase, EC 1.10.3.1). The enzymatic product, melanin,

should stain salivary sheaths and halos around the sheaths. To detect peroxidase (EC 1.11.1.7) activity,

artificial diets were immersed for some minutes in 0.1% (w/v) diaminobenzidine (DAB, Sigma) in 50 mM

Tris (pH 7.5) containing 0.1% (v/v) H_2O_2 (Sigma). The enzymatic product should induce reddish staining of

salivary sheaths and halos. For identification of pectinase activity, 0.1% (w/v) of pectin (Sigma) was

added to the medium. After exposure to aphids, the gel was transferred for 3 h into a Petri dish

containing 50 mM citrate-phosphate buffer, at pH 5.0 to detect pectin (methyl) esterase (PME, EC

3.1.1.11) and at pH 6.4 to detect polygalacturonase (PG, EC 3.1.1.15). The gel was then stained with a

solution of 0.01% (w/v) ruthenium red (Sigma) for 1 h, and then washed several times with distilled

water. At pH 6.4, red halos around the salivary sheaths indicate PME activity, while non-staining halos at

pH 5 in the pink pectin indicate PG activity. Finally, for proteinase activity (EC 3.4.99), 0.5% (w/v) of

gelatin (Sigma) was added to the medium. After exposure to aphids, the medium was incubated

overnight in a solution of 50 mM Tris (pH 8) containing 100 mM NaCl and 10 mM $CaCl_2$, then stained with

Coomassie blue. An absence of blue staining shows proteinase activity. All observations of proteins and

enzymatic activities were performed by light microscopy (Axioplan 2, Zeiss, Jena, Germany).

2.3.3 Poplar protoplast preparations and treatments

Mesophyll protoplasts of the two poplar genotypes were obtained from young leaves as described in Wu

et al. (2009). Leaves were cut into 1–2 mm fine strips in 0.3 M sorbitol and 66.67 mM $CaCl_2$ (pH 5.6) and

lysed in an enzyme solution (0.6 M mannitol, 0.25% (w/v) cellulase Onozuka R-10, 0.05% (w/v)

macerozyme R-10) in the dark for 16 h with gentle shaking (30 rpm) at room temperature. Protoplasts

were collected by filtering the lysis solution through a 70 μm cell strainer (Falcon®) and concentrated by

spinning down at ≈ 800 x g for 10 min at 4 °C. The pellet was washed twice with W5 buffer (154 mM

NaCl, 125 mM CaCl$_2$, 5 mM KCl, 5 mM glucose, 0.03% (w/v) MES, pH 5.8) and then resuspended in 0.6 M

mannitol to a final concentration of 1×10^6 protoplasts per mL. Protoplasts (1.10^6) were incubated at

20°C with gentle shaking (40 rpm) for 3 h with aphid salivary proteins or with protein extraction buffer

(control). Preliminary experiments investigating expression of 10 poplar genes, and conducted with 1,

10, 20, 40 or 80 µg of salivary proteins, indicated that the optimal response (i.e. the maximum fold

change) was observed with 1 and 10 µg of salivary proteins of *P. passerinii* and *M. persicae*, respectively.

Protoplast viability, before and after treatment with aphid saliva, was assessed using 0.005% (w/v)

fluorescein diacetate (FDA). After 5 min of incubation protoplasts were observed under blue light

epifluorescence, and cell viability was estimated as the percentage of fluorescent cells. Most protoplasts

were intact and viable after enzymatic digestion with cellulase and macerozyme (98%), as well as after

incubation with salivary proteins (95%; Fig. S2).

2.3.4 Quantitative RT-PCR

After the aphid saliva treatments, protoplasts were centrifuged at ≈ 800 x g for 2 min. Total RNAs were

extracted with the RNeasy® Plant Kit Mini Kit (Qiagen). A DNase treatment with the RNase-free DNase

Set (Qiagen) was carried out for 15 min at 25°C. Total RNA concentration was determined using a

Nanodrop ND-1000 spectrophotometer. All RNA samples were rejected if they did not reach a minimum

concentration of 100 ng µL^{-1}, a 260 nm/280 nm ratio between 1.8 and 2.0. Poly(dT) cDNA was prepared

from 1 µg total RNA using the iScriptTMcDNA Synthesis Kit (Bio-Rad) and quantified with a LightCycler

480 (Roche) and SYBR GREEN I Master (Roche), according to the manufacturer's instructions. PCR was

carried out in 384-well optical reaction plates heated for 10 min to 95°C to activate hot start Taq DNA

polymerase, followed by 40 cycles of denaturation for 60 sec at 95°C and annealing/extension for 60 sec

at 58°C. The distribution of the quantitative RT-PCR mix containing SYBR Green I Master (Roche), cDNAs

and primers was performed using the EVO150$^{©}$ (Tecan) pipetting robot in a 384-well plate. The

expression of 43 genes, belonging to eight different physiological processes or metabolic pathways (i.e.

auxin, cytokinins, jasmonates, ethylene, salicylic acid, phenolic compounds, reactive oxygen species

(ROS), cell cycle), was quantified with specific primer pairs designed by Quant-Prime (Arvidsson,

Kwasniewski, Riaño-Pachón & Mueller-Roeber, 2008) based on the *Populus trichocharpa* sequence (v3.0)

from Phytozome (https://phytozome.jgi.doe.gov/pz/portal.html; Table S2). Expression levels were

normalized to the levels of *PtUBIQUITIN10* (*PtUBQ10*), commonly used as a reference gene in plants (e.g.

Tong, Gao, Wang, Zhou & Zhang, 2009). All RT-qPCR experiments were done with three independent

biological replicates, with two technical replicates each. One of the biological replicates of the *M.*

persicae – Koster interaction has been excluded from the analyses because of technical issues during

quantification. Relative gene expression was calculated according to the ^{ΔΔ}Cp method, with protoplasts

incubated with protein extraction buffer as controls. Primers used for gene expression analysis are listed

in Table S2.

2.3.5 Histochemical analysis of GUS activity

Transgenic seedlings of *A. thaliana* (five-day-old and eight-day-old for *pIAA2*::GUS and *pARR16*::GUS,

respectively) were incubated with 2 mL of liquid MS containing 1 µg and 10 µg of aphid salivary proteins

(in TE/Tween buffer) of either *P. passerinii* or *M. persicae* for 3 h and 4 h for *pIAA2*::GUS and

pARR16::GUS, respectively. Positive controls were incubated with 20 µM of indole acetic acid (IAA)

(Sigma-Aldrich), and 20 µM of 6-benzylaminopurine (BAP) (Sigma-Aldrich). Negative controls were

incubated in liquid MS and corresponding volumes of TE/Tween buffer. Five seedlings were used for

each modality. Seedlings were then incubated in reaction buffer containing 0.1 M sodium phosphate

buffer (pH 7), 2 mM ferricyanide, 2 mM ferrocyanide, 0.1% (v/v) Triton X-100 and 1 mg ml$^{−1}$ X-Gluc for

1 up to 24 h in dark at 37 °C. Afterwards, chlorophyll was removed by destaining in 70% ethanol and

seedlings were cleared as described by Malamy and Benfey (1997). GUS expression was monitored by

differential interference contrast microscopy.

2.4. Data analysis

All tests were carried out with the statistical software R 2.11.0 (R Development Core Team, 2013). RT-

qPCR results have been expressed as fold-changes in gene expression compared to the reference gene

PtUBQ10. Fold-changes have been $\log_2$ - transformed. Following this transformation, fold-changes varied

between -∞ and +∞, with negative values corresponding to gene underexpression, positive values to

overexpression and zero, to no change in gene expression. Permutational multivariate anovas

(permanovas), using 1000 permutations, were computed to test the effect of aphid species and host

plant genotype on the simultaneous fold-changes in gene expression of the 43 considered poplar genes

using the R package vegan (Oksanen et al. 2013). To visualize similarities and differences in fold-changes

a heatmap was built, with genes in columns and modalities in lines. The heatmap was built with a Z-

score, i.e. $\log_2$ – transformed fold changes which have been normalized and centered per column.

Hierarchical clustering with Euclidean distances was added to the heatmap to visualize the proximity

among genes in columns and among modalities in lines.

A univariate model has also been used to analyze the effect of modalities (i.e. aphid species and host

genotype) on fold-changes of each gene. The model equation was:

$$\log_{10}(y_{ij}) = \alpha_i + \varepsilon_{ij}$$

where y_{ij} is the fold-change for the modality i and the biological replicate j, and α_i is the effect of the

modality i on relative gene expression. The model error follows a normal distribution, with a null mean

and a variance σ^2. There is no intercept in this model and 4 independent α_i parameters were estimated,

corresponding to each aphid – poplar genotype combination. These parameters were estimated with a

Bayesian approach, with $\alpha_i \sim N(0, 0.5^2)$ for prior. The parameters have been estimated with a Markov

Chain Monte Carlo algorithm and the package R brms (Buerkner, 2016). The maxima *a posteriori* and

95% credibility intervals were calculated, with downregulation and upregulation probabilities for each

gene. A credibility interval excluding 1 (i.e. the constitutive expression of genes) indicates a significant

effect of aphid saliva on gene expression. This approach was selected because it allows to perform

statistical analyses with a limited number of biological replications.

3 Results

3.1 Annotation, secreted proteins detection and identification

From 36,312 and 3,233 transcripts, 1,243 and 221 transcripts were predicted to encode for secreted

salivary proteins in *P. passerinii* and *M. persicae*, respectively. About half of them (604) have been

annotated for *P. passerinii* and 190 for *M. persicae*. Using OrthoVenn, 121 and 58 protein clusters have

been identified for *P. passerinii* and *M. persicae*, respectively. About 17% of these clusters were common

between the two aphids (Table S3).

Blast2GO determined that *P. passerinii* salivary proteins were predominantly binding proteins and

enzymes (Fig. 1, table S4). The most common enzymes were peptidases (especially serine-type and

cysteine-type endopeptidases), kinases (especially phosphatidylinositol phosphate (PIP) kinases) and

hydrolases. Several enzymes involved in the degradation of carbohydrates (i.e. cellulase, trehalase, β-

glucuronidase, mannosidase and glucosylceramidase), and of phenolic compounds (i.e. peroxidase and

oxidoreductase) were also identified (Fig. 1, table S4). Among binding proteins, dimerization protein,

nucleic acids binding (especially DNA binding), nucleotide binding (especially ATP binding) and cation-

binding (mostly calcium ion-binding and zinc-binding proteins) were the most commonly found (Fig. 1,

table S4). Proteins related to hormone activity were also identified. Glucose dehydrogenases were also

detected with OrthoVenn (table S3). Among the 12 aphid salivary effectors considered, five were

identified in *P. passerinii*, with low E-values (< 7 e^{-71}): Mp10, ARMET, ACE 1, ACE2 and ACE3.

3.2 Enzyme activities

Staining with Coomassie blue confirmed the protein nature of the salivary sheath material (Fig. 2A) and

DOPA staining indicated a phenoloxidase activity in the sheaths (Fig. 2B). Black halos were also observed

around some sheaths (Fig. 2B). Likewise, peroxidase activity was found in salivary sheaths and halos

around sheaths (Fig. 2C). However, no pectinesterase, polygalacturonase and proteinase activity was

detected.

3.3 Quantitative RT-PCR

The multivariate analysis based on the expression of the 43 genes showed a highly significant effect of

aphid species (pseudo-$F_{1, 10}$ = 12.07, P < 0.001), no effect of poplar genotype, but a significant aphid x

poplar genotype interaction (pseudo-$F_{1, 10}$ = 3.51, P = 0.038). This indicated that the salivary proteins

effect differed depending on aphid x poplar genotypes combinations. Fold changes across modalities are

presented in figure 3. The hierarchical clustering of modalities (left dendrogram) shows an arrangement

of modalities into two groups. In the upper group, most poplar genes were upregulated following

incubation with salivary proteins, while in lower group most genes were downregulated (Fig. 3). The

upper group gathered most biological replicates of incompatible (i.e. *P. passerinii* – Koster) and non-host

(i.e. *M. persicae* – Koster, *M. persicae* – I-214) interactions. The lower group gathered all replicates of

compatible interactions (i.e. *P. passerinii* – I-214) and an incompatible one. The hierarchical clustering of

genes (upper dendrogram) showed that genes could also be arranged into two groups. The group on the

left allowed to separate the effect of salivary proteins of *M. persicae*, with a general downregulation of

genes, from those of *P. passerinii*, with a general upregulation (Fig. 3). The group on the right allowed to

separate compatible interactions, with a general downregulation of genes, from other interactions

during which genes were generally upregulated (Fig. 3). There was no clustering of genes according to

physiological process or metabolic pathway.

For the auxin pathway, the compatible interaction was characterized by a downregulation of genes

related to auxin transport (i.e. *PtAUX1* and *PtPIN1* with downregulation probabilities of 62% and 86%,

respectively) and homeostasis (i.e. *GH3* with a downregulation probability >99%; Fig. 4A). However,

there was no effect on genes involved in auxin biosynthesis (i.e. *PtNIT1* and *PtYUCCA*). Reverse variations

were observed for non-host and incompatible interactions which led to an upregulation of *PtGH3*, with

upregulation probabilities >99% for *M. persicae* and 86% for *P. passerinii*, and upregulations of genes

related to auxin transport with probabilities always higher than 86% (Fig. 4A).

For the cytokinin pathway, the compatible interaction did not affect the expression of genes related to

cytokinin metabolism (i.e. *PtLOG5* and *PtIPT*), but downregulated cytokinin signaling genes (*PtAHK4* and

PtARR2), with downregulation probabilities >99% (Fig. 4B). Non-host interactions led to upregulations of

all genes but *PtLOG5*, with probabilities >95% (Fig. 4B). Response of genes to the incompatible

interaction followed a similar trend, with upregulation of all genes, including *PtLOG5*, with probabilities

>95%.

Regarding biotic stress signaling (i.e. jasmonates, salicylic acid, ethylene and ROS), the compatible

interaction was characterized by null or downregulation responses for all the considered genes (Fig. S3).

For ROS genes weak downregulations were also observed for non-host interactions (>85% and >64%

downregulation probabilities for *PtSOD* and *PtCAT*, respectively) and a weak upregulation for the

incompatible interaction (91% and 77% upregulation probabilities for *PtSOD* and *PtCAT*, respectively).

For salicylic acid, jasmonates and ethylene, non-host interactions induced a strong upregulation of most

genes (>99% upregulation probabilities in most cases), except *PtPR5* for salicylic acid which was

downregulated (>98% downregulation probabilities), *PtAOS* for jasmonates and *PtEIN2* for ethylene

which did not respond. The incompatible interaction resulted in an upregulation of all genes involved in

jasmonates and ethylene pathways but to a lesser extent than non-host interactions (>81% upregulation

probabilities), except *PtJAZ1* which was downregulated (73% downregulation probability) and *PtJAR1*,

PtERF1 which were not affected. For salicylic acid, weak upregulations were observed for *PtNPR1*, *PtPR5*

and *PtNDR1* (>62% upregulation probabilities), and no effect was detected for *PtPR1* and *PtEDS1*.

For genes involved in the phenolic compounds pathway, *PtF5H* and *PtANT* were downregulated during

the compatible interaction (>85% downregulation probability), while all genes were upregulated during

non-host and incompatible interactions (>89% upregulation probability in all cases; Fig. S3). Finally,

regarding genes involved in cell cycle, *PtCYCD5*, *PtMCM2* and *PtRBR* tended to be downregulated during

compatible interactions (>72% downregulation probability). Conversely, most genes, except *PtCAK1* and

PtMCM2, were upregulated during non-host interactions (>94% upregulation probabilities in most

cases), and most genes were not affected during incompatible interactions, except *PtCDK5*, *PtCDK20* and

PtCAK1 which were upregulated (>95% upregulation probabilities; Fig. S3).

3.4 Histochemical analysis of GUS activity

Salivary proteins of *P. passerinii* increased *pIAA2*::GUS signals (Fig. 5E, 5F, 5I, 5J), which were similar to

those caused by an exogenous application of auxin (Fig. 5A and 5B). Incubation with salivary proteins of

M. persicae resulted in faint colorations (Fig. 5G, 5H, 5K, 5L), similar to those of negative controls (Fig. 5C

and 5D).

Positive controls of *pARR16*::GUS were characterized by a strong staining in the middle part of root

central cylinder (Fig. 6A and 6B), which was weak in negative controls as well as with *M. persicae* salivary

proteins (Fig. 6G, 6H, 6K, and 6L). No coloration was visible in the roots of seedlings incubated with

salivary proteins of *P. passerinii* (Fig. 6E, 6F, 6I and 6J).

4 Discussion

Plant manipulation by parasites requires a finely tuned reprogramming of host metabolism, achieved

through secretion of effectors into the host plant (Hogenhout & Bos, 2011; Giron et al., 2016). Using

both *in vivo* and *in planta* approaches, we confirmed that salivary proteins impact gene transcription of

its host tree, strongly suggesting they mediate the interactions between *P. passerinii* and its host-tree.

Salivary extracts of *P. passerinii* significantly affected gene expression of the two host-plant genotypes

considered, in a markedly different way. Compatible interactions were characterized by a general trend

of gene downregulation in all the metabolic pathways and physiological processes investigated, including

genes involved in biotic stress signaling. This supports the hypothesis that the saliva of *P. passerinii*

includes effectors proteins affecting plant signaling and defense mechanisms.

Several of the detected salivary proteins may interfere with plant signaling. PIP kinases were quite

common in the saliva of *P. passerinii*, but have not been previously reported from any aphid saliva. These

enzymes catalyze phosphorylation of phosphatidyl-inositol into phosphatidylinositol-4-5-biphosphate

(PIP_2). Hydrolysis of PIP_2 produces secondary messengers like diacylglycerol and inositol-1-4-5-

triphosphate (IP_3), which can in turn be hydrolyzed into phosphatidic acid (PA), considered an important

signaling molecule in plants, triggered in response to various biotic and abiotic stresses (Testerink &

Munnik, 2005). Likewise, PIP_2 and IP_3 can affect cellular oscillations of Ca^{2+} and are involved in multiple

processes including cell cycle and phytohormone regulation (Xue, Chen & Mei, 2009). Other proteins,

frequently detected in the saliva of *P. passerinii*, may also interfere with secondary messengers like

calcium-binding, ATP-binding, and GTP-binding proteins or with hormone signaling like hormone-binding

proteins (Vandermoten et al., 2014; Giron et al., 2016). Likewise, it has been hypothesized that trehalase

may interfere with trehalose-based defense responses in *A. thaliana* (Nicholson et al., 2012), and

cellulases may also contribute to the degradation of oligogalacturonides involved in damage signaling

pathways (Cherqui & Tjallingii, 2000). Nucleic acid-binding proteins and protein-binding proteins could

also participate in the manipulation of host-plant metabolism (Vandermoten et al., 2014). Finally,

oxidoreductases, especially peroxidases and phenoloxidases could degrade phenolic compounds of the

host plant (Cherqui & Tjallingii, 2000; Carolan, Fitzroy, Ashton, Douglas & Wilkinson, 2009). Gall

induction by *P. passerinii* in susceptible host genotypes is characterized by transient accumulation of

phenolic compounds (Dardeau et al., 2014), and oxidoreductases could help aphids to cope with these

secondary metabolites. Glucose dehydrogenases may similarly help aphids to detoxify defensive

compounds of the host-plant (Carolan et al., 2011; Nicholson et al., 2012).

Several proteins orthologous to salivary effectors of *M. persicae* and *A. pisum* have been identified.

Likewise, many of the proteins detected in our study (e.g. calcium-binding, DNA-binding, ATP-binding,

GTP-binding proteins, glucose dehydrogenases, oxidoreductases, trehalases and phosphatases) have also

been previously identified in the saliva of different aphid species, which were not gall-inducers (Harmel

et al., 2008; Nicholson et al., 2012; Elzinga & Jander, 2013; Vandermoten et al., 2014). Therefore,

although these proteins probably contribute to plant manipulation by aphids (Elzinga & Jander, 2013),

they are probably not specifically involved in gall induction. Interestingly, calcium-binding proteins are

supposedly key components of sap-feeding aphids' saliva, preventing the plugging of sieve tubes (Will et

al., 2007). Since *P. passerinii* does not feed on sap (Pointeau et al., 2012), it suggests that these proteins

also play other crucial roles during aphid-plant interactions. Several proteins detected in the saliva of *P.*

passerinii such as serine proteases, acid phosphatases, lipases and metalloproteases have been proposed

as potential effectors involved in formation and/or maintenance of fig wasps galls (Martinson, Hackett,

Machado & Arnold, 2015).

In situ biochemical assays confirmed the presence of peroxidases in both solid and soluble saliva of *P.*

passerinii. The phenoloxidase activity is also congruent with the numerous oxidoreductase sequences

identified in the salivary transcriptome. Likewise, the absence of pectin (methyl) esterase and

polygalacturonase activity during *in situ* bioassays is consistent with their absence in the salivary

transcriptome. However, while numerous proteases were identified among the salivary proteins of *P.*

*passerin*ii, no activity was observed during *in situ* bioassays. Gelatin was probably not the adequate

substrate to detect the protease activity of *P. passerinii*. Additional *in situ* assays could be conducted to

detect, in *P. passerinii* saliva, the activity of the proteases and possibly other enzymes like cellulases

(Cherqui & Tjallingii, 2000). A proteomic analysis of salivary extracts should also confirm and

complement the predictions of our transcriptomic approach (Carolan et al., 2011; Boulain et al., 2018).

To knock-down or divert stress signaling and/or ensure finely-tuned reprogramming of host metabolism

and anatomical transformation of host tissues, plant manipulation by herbivores or pathogens generally

requires reconfiguration of phytohormone pathways and signaling (Giron, Frago, Glevarec, Pieterse &

Dicke, 2013; Tooker & Helms, 2014). Both RT-qPCR experiments and histochemical assays confirmed that

P. passerinii can actively manipulate phytohormone pathways during compatible interactions. This is in

line with the numerous proteins potentially interfering with plant signaling and metabolism found in the

aphid saliva. Salivary extracts of *P. passerinii* did not affect the auxin biosynthesis during compatible

interactions, but downregulated auxin transporter genes such as *PtPIN1* and *PtAUX1*, as well as *PtGH3*,

which is involved in the homeostasis of auxin active forms (Park et al., 2007). These downregulations

could lead to intracellular accumulation of active auxin forms, and result in the targeted cell hypertrophy

and multiplication commonly observed during gall differentiation by *P. passerinii* (Dardeau et al., 2014).

The activation of the auxin-responsive promoter *IAA2* during *in planta* assays with transgenic seedlings

of *A. thaliana* further supports this hypothesis of an intracellular accumulation of auxin during gall

initiation by *P. passerinii*. Similar auxin accumulation, as a result of a reduction in both *PtGH3* activity and

auxin transport, is also probably involved in the initiation and development of root galls by cyst and root-

knot nematodes (Karczmarek, Overmars, Helder & Goverse, 2004), suggesting a potentially convergent

manipulation strategy between these organisms. As genes related to cytokinin biosynthesis (*PtIPT*) and

activation (*PtLOG5*) were not affected by salivary proteins of *P. passerinii*, the strong downregulation of

cytokinin signaling genes (*PtAHK4*, *PtARR2* and *PtARR16*) upon treatment could correspond to an auxin

accumulation-induced regulation loop (Jones et al., 2010; Schaller, Bishopp, & Kieber, 2015). In this

aspect, *P. passerinii* could manipulate auxin and cytokinin plant responses to promote the host division

cycle leading to the formation of galls (Giron et al., 2013), which provide to the insects food and shelter

at the expense of the host plant (Tooker & De Moraes, 2008).

Apart from the Mp10 effector, the saliva of *P. passerinii* and *M. persicae* shared few similarities. The

transcriptomes of *P. passerinii* and *M. persicae* have been obtained with a different sequencing and

assembly methods (Ramsay et al., 2007). This probably explains the difference in transcripts amounts

gathered for both aphids and several salivary proteins of *M. persicae* might be missing in its

transcriptome, leading to an apparently low similarity between secretomes. Previous comparisons

among salivary proteins of aphid species also indicated that aphids with different host species and / or

feeding strategies exhibited very different salivary protein profiles (Cooper, Dillwith & Puterka, 2011;

Vandermoten et al., 2014). This is in agreement with the highly significant effect of aphid species on

gene expression profiles in both poplar genotypes. Nonetheless, gene expression profiles of poplar

protoplasts during incompatible interactions shared many similarities with those observed during non-

host interactions. Both types of interactions were characterized by an overall upregulation of host genes,

which was generally more important during non-host interactions than during incompatible ones,

probably because one of the three biological replicates of incompatible interactions shared similarities

with compatible interactions. During both non-host and incompatible interactions, most genes involved

in jasmonates, ethylene and salicylic acid pathways were upregulated. All of these pathways can be

activated following aphid feeding, together or separately, depending on the aphid – plant interaction

system considered (Morkunas, Mai & Gabryś, 2011; Kerchev, Fenton, Foyer & Hancock, 2012; Louis &

Shah, 2013). Due to the salicylic acid - jasmonates cross-talk, a differential gene expression between the

two pathways could be expected (Zarate, Kempema & Walling, 2007), but was not observed here.

However, this may occur later during the interaction, since in our experiments we only considered early

gene responses (Kerchev et al., 2012). As a consequence of activation of plant defense signaling, genes

related to secondary metabolism, i.e. *PtF3'5'H, PtANT* and *PtF5H* were also upregulated during both

incompatible and non-host interactions, while they were unaffected or downregulated during

compatible interactions. This is in agreement with previous histochemical analyses showing a light and

transient accumulation of phenolic compounds followed by their marked disappearance in galled tissues

during a compatible interaction, while their accumulation was strong and continuous during an

incompatible interaction (Dardeau et al., 2014).

Interestingly, most of the few genes that were differentially expressed during non-host and incompatible

interactions were related to biotic stress signaling, and all genes but *PtJAZ1* were upregulated during

incompatible interactions while they were unaffected or downregulated during non-host interactions.

These genes included ROS-related genes, *PtSOD* and *PtCAT*, the expression of which was slightly

downregulated during non-host interactions. A similar repression of *SOD* and *CAT* genes, leading to an

increase in H_2O_2 content, together with an upregulation of *JAZ1* and *PR1*, has been reported in potato

leaves infested by *M. persicae* (Kerchev et al., 2012). Some phytohormones-related genes were also

upregulated during incompatible interactions only, like *PtAOS* for jasmonates, *PtPR5* for salicylic acid,

PtLOG5 for cytokinins, *PtNIT1* for auxin, and finally *PtEIN2* for ethylene. This latter gene is involved in

transduction of ethylene signaling and can be up- or downregulated during both compatible and

incompatible interactions in different plant-aphid systems and plays equivocal roles during host-plant

resistance to *M. persicae* (Morkunas et al., 2011; Louis & Shah, 2013). Moreover, the putative

intracellular accumulation of auxin during compatible interactions could also interfere with salicylic acid

signaling and defense responses (Park et al., 2007), which is congruent with the overall downregulation

of salicylic acid pathway related genes. The downregulation of *PtAHK4* and *PtARR2* could also interfere

with stress signaling by preventing both accumulation of jasmonates and activation of *PtPR1* (O'Brien &

Benkova, 2013), which is consistent with our observations during compatible interactions. As a result,

the effectors of *P. passerinii* saliva, in addition to modifying phytohormonal contents, down-regulate the

plant defenses to allow gall formation.

In conclusion, our transcriptomic analysis of the saliva of *P. passerinii* and *M. persicae* showed that the

gall-inducing aphid probably secretes a highly peculiar saliva, filled with potential effectors that may

interfere with several plant secondary messengers and signaling pathways. Our *in vivo* and *in planta*

approaches confirmed the ability of salivary extracts of the gall-inducing insects to manipulate host

response during compatible interactions. As expected phytohormones pathways were strongly affected,

probably to impair biotic stress signaling but also to reconfigure host metabolism and anatomy. Although

the saliva of *P. passerinii* and *M. persicae* were very different, incompatible and non-host interactions led

to similar host responses, with a different intensity however, and few differences in biotic stress

signaling. Additional modalities including different populations of *P. passerinii*, different poplar

genotypes with intermediate resistance levels, leading for instance to semi-compatible interactions, and

additional host metabolic pathways could be considered in future experiments and give further insights

on the molecular processes underpinning failed and successful host manipulation by a gall-inducer.

References

Arvidsson S., Kwasniewski M., Riaño-Pachón D.M. & Mueller-Roeber B. (2008). QuantPrime--a flexible

tool for reliable high-throughput primer design for quantitative PCR. *BMC Bioinformatics*, 9, 465.

Atamian H.S., Chaudhary R., Dal Cin V., Bao E., Girke T. & Kaloshian I. (2013) *In planta* expression or

delivery of potato aphid *Macrosiphum euphorbiae* effectors Me10 and Me23 enhances aphid fecundity.

Molecular Plant-Microbe Interactions, 26, 67–74.

Bendtsen J.D., Jensen L.J., Blom N., Von Heijne G. & Brunak, S. (2004). Feature-based prediction of non-

classical and leaderless protein secretion. *Protein Engineering Design and Selection*, 17, 349-356.

Bishopp A., El-Showk S., Weijers D., Scheres B., Friml J., Benková E., … Helariutta Y. (2011). A mutually

inhibitory interaction between auxin and cytokinin specifies vascular pattern in roots. *Current Biology*,

21, 917-926.

Boulain H., Legeai F., Guy E., Morlière S., Douglas N.E., Oh J., … & Sugio A. (2018). Fast evolution and

lineage-specific gene family expansions of aphid salivary effectors driven by interactions with host-

plants. *Genome Biology and Evolution*, 10, 1554-1572.

Buerkner P.C. (2016) brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical*

Software, 80, 1-28.

Carolan J.C., Fitzroy C.I.J., Ashton P.D., Douglas A.E. & Wilkinson T.L. (2009). The secreted salivary

proteome of the pea aphid *Acyrthosiphon pisum* characterised by mass spectrometry. *Proteomics*, 9,

2457–2467.

Carolan J.C., Caragea D., Reardon K.T., Mutti N.S., Dittmer N., … Edwards O.R. (2011). Predicted effector

molecules   in   the   salivary   secretome   of   the   pea   aphid   (*Acyrthosiphon   pisum*):   a   dual

transcriptomic/proteomic approach. *Journal of Proteome Research*, 10, 1505–1518.

Chaudhary R., Atamian H.S., Shen Z., Briggs S.P. & Kaloshian I. (2014). GroEL from the endosymbiont

Buchnera aphidicola betrays the aphid by triggering plant defense. *Proceedings of the National Academy*

of Sciences, 111, 8919-8924.

Cherqui A. & Tjallingii W.F. (2000). Salivary proteins of aphids, a pilot study on identification, separation

and immunolocalisation. *Journal of Insect Physiology*, 46, 1177–1186.

Conesa A., Götz S., García-Gómez J.M., Terol J., Talón M. & Robles M. (2005). Blast2GO: a universal tool

for annotation, visualization and analysis in functional genomics research. *Bioinformatics*, 21, 3674–

3676.

Cooper W.R., Dillwith J.W. & Puterka G.J. (2011). Comparisons of salivary proteins from five aphid

(Hemiptera: Aphididae) species. *Environmental Entomology*, 40, 151-156.

Dardeau F., Deprost E., Laurans F., Lainé V., Lieutier F. & Sallé, A. (2014). Resistant poplar genotypes

inhibit pseudogall formation by the wooly poplar aphid, *Phloeomyzus passerinii* Sign. *Trees*, 28, 1007-

1019.

De Vos M. & Jander G. (2009). *Myzus persicae* (green peach aphid) salivary components induce defence

responses in *Arabidopsis thaliana. Plant Cell & Environment*, 32, 1548–1560.

Elzinga D.A. & Jander G. (2013). The role of protein effectors in plant–aphid interactions. *Current Opinion*

in Plant Biology, 16, 451-456.

Emanuelsson O., Nielsen H., Brunak S. & von Heijne G. (2000). Predicting subcellular localization of

proteins based on their N-terminal amino acid sequence. *Journal of Molecular Biology*, 300, 1005-1016.

Favery B., Quentin M., Jaubert-Possamai S. & Abad P. (2016). Gall-forming root-knot nematodes hijack

key plant cellular functions to induce multinucleate and hypertrophied feeding cells. *Journal of insect*

physiology, 84, 60-69.

Giron D., Frago E., Glevarec G., Pieterse C.M. & Dicke M. (2013). Cytokinins as key regulators in plant–

microbe–insect interactions: connecting plant growth and defence. *Functional Ecology*, 27, 599-609.

Giron D., Huguet E., Stone G.N. & Body M. (2016). Insect-induced effects on plants and possible effectors

used by galling and leaf-mining insects to manipulate their host-plant. *Journal of Insect Physiology*, 84,

70-89.

Guiguet A., Dubreuil G., Harris M.O., Appel H., Schultz J.C., Pereira M.H. & Giron D. (2016). Shared

weapons of blood- and plant-feeding insects: surprising commonalities for manipulating hosts. *Journal of*

Insect Physiology, 84, 4–21.

Guo K., Wang W., Luo L., Chen J., Guo Y. & Cui F. (2014). Characterization of an aphid-specific, cysteine-

rich protein enriched in salivary glands. *Biophysical Chemistry*, 189, 25–32.

Harmel N., Létocart E., Cherqui A., Giordanengo P., Mazzucchelli G., Guillonneau F., … Francis F. (2008).

Identification of aphid salivary proteins: a proteomic investigation of *Myzus persicae*. *Insect Molecular*

Biology, 17, 165–174.

Hogenhout S.A. & Bos J.I. (2011). Effector proteins that modulate plant–insect interactions. *Current*

Opinion in Plant Biology, 14, 422-428.

Hyatt D., Chen G.L., LoCascio P.F., Land M.L., Larimer F.W. & Hauser L.J. (2010). Prodigal: prokaryotic

gene recognition and translation initiation site identification. *BMC Bioinformatics*, 11:119.

Jaouannet M., Rodriguez P.A., Thorpe P., Lenoir C.J.G., MacLeod R., Escudero-Martinez C. & Bos J.I.B.

(2014). Plant immunity in plant–aphid interactions. *Frontiers in Plant Science*, 5, 663.

Ji R., Yu H., Fu Q., Chen H., Ye W., Li S. & Lou Y. (2013). Comparative transcriptome analysis of salivary

glands of two populations of rice brown planthopper, *Nilaparvata lugens*, that differ in virulence. *PloS*

one, 8, e79612.

Jones B., Gunnerås S.A., Petersson S.V., Tarkowski P., Graham N., May S., … Ljung, K. (2010). Cytokinin

regulation of auxin synthesis in *Arabidopsis* involves a homeostatic feedback loop regulated via auxin and

cytokinin signal transduction. *The Plant Cell*, 22, 2956-2969.

Jones P., Binns D., Chang H.Y., Fraser M., Li W., McAnulla C., … Pesseat S. (2014). InterProScan 5:

genome-scale protein function classification. *Bioinformatics*, 30, 1236-1240.

Karczmarek A., Overmars H., Helder J. & Goverse A. (2004). Feeding cell development by cyst and

root-knot nematodes involves a similar early, local and transient activation of a specific auxin-inducible

promoter element. *Molecular Plant Pathology*, 5, 343-346.

Kerchev P.I., Fenton B., Foyer C.H. & Hancock R.D. (2012). Infestation of potato (*Solanum tuberosum* L.)

by the peach-potato aphid (*Myzus persicae* Sulzer) alters cellular redox status and is influenced by

ascorbate. *Plant, Cell & Environment*, 35, 430-440.

Lieutier F., Bermudez-Torres K., Cook J., Harris M.O., Legal L., Sallé A., ... Giron D. (2017). From Plant

Exploitation to Mutualism (eds N. Sauvion, D. Thiéry & P. A. Calatayud), pp. 55-109. Elsevier.

Louis J. & Shah J. (2013). *Arabidopsis thaliana*—*Myzus persicae* interaction: shaping the understanding of

plant defense against phloem-feeding aphids. *Frontiers in Plant Science*, 4, 213.

Malamy J.E. & Benfey P.N. (1997). Organization and cell differentiation in lateral roots of *Arabidopsis*

thaliana. *Development*, 124, 33-44.

Martinson E.O., Hackett J.D., Machado C.A. & Arnold A.E. (2015). Metatranscriptome analysis of fig

flowers provides insights into potential mechanisms for mutualism stability and gall induction. *PloS One*,

10, e0130745.

Morkunas I., Mai V.C. & Gabryś B. (2011). Phytohormonal signaling in plant responses to aphid feeding.

Acta Physiologia Plantarum, 33, 2057–2073.

Mutti N.S., Louis J., Pappan L.K., Pappan K., Begum K., Chen M.S., ... Reeck G.R. (2008). A Protein from the

salivary glands of the pea aphid, *Acyrthosiphon pisum*, is essential in feeding on a host plant. *Proceedings*

of the National Academy of Sciences, 105, 9965-9969.

Nicholson S.J., Hartson S.D. & Puterka G.J. (2012). Proteomic analysis of secreted saliva from Russian

Wheat Aphid (Diuraphis noxia Kurd.) biotypes that differ in virulence to wheat. *Journal of Proteomics*, 75,

2252–2268.

O'Brien J.A. & Benková E. (2013). Cytokinin cross-talking during biotic and abiotic stress responses.

Frontiers in Plant Science, 4, 451.

Oksanen J., Blanchet F.G., Kindt R., Legendre P., Minchin P.R., O'hara R.B., ... Wagner H. (2013). Package

'vegan'. Community Ecol Package Version 2.

Pan Y., Zhu J., Luo L., Kang L. & Cui F. (2015). High expression of a unique aphid protein in the salivary

glands of *Acyrthosiphon pisum. Physiological and Molecular Plant Pathology*, 92, 175–180.

Park J.E., Park J.Y., Kim Y.S., Staswick P.E., Jeon J., Yun J., ... Park C.M. (2007). GH3-mediated auxin

homeostasis links growth regulation with stress adaptation response in *Arabidopsis. Journal of Biological*

Chemistry, 282, 10036–10046.

Petersen T.N., Brunak S., von Heijne G. & Nielsen H. (2011). SignalP 4.0: discriminating signal peptides

from transmembrane regions. *Nature Methods*, 8, 785.

Pierleoni A., Martelli P.L. & Casadio, R. (2008). PredGPI: a GPI-anchor predictor. *BMC Bioinformatics*, 9,

392.

Pointeau S., Ameline A., Laurans F., Sallé A., Rahbé Y., Bankhead-Dronnet S. & Lieutier F. (2012).

Exceptional plant penetration and feeding upon cortical parenchyma cells by the woolly poplar aphid.

Journal of Insect Physiology, 58, 857–866.

Pundir S., Martin M.J. & O'Donovan C. (2017). Uniprot protein knowledgebase. *Protein Bioinformatics:*

From Protein Modifications and Networks to Proteomics, 41-55.

R Core Team (2013). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.

Ramsey J.S., Wilson A.C.C., de Vos M., Sun Q., Tamborindeguy C., Winfield A., Malloch G., Smith D.M.,

Fenton B., Gray S.M., Jander G. (2007). Genomic resources for *Myzus persicae*: EST sequencing, SNP

identification, and microarray design. *BMC Genomics*, 8, 423–423.

Sallé A., Pointeau S., Bankhead-Dronnet S., Bastien C. & Lieutier, F. (2017). Unraveling the tripartite

interactions among the woolly poplar aphid, its host tree, and their environment: a lead to improve the

management of a major tree plantation pest? *Annals of Forest Science*, 74, 79.

Schaller G.E., Bishopp A., & Kieber J.J. (2015). The yin-yang of hormones: cytokinin and auxin interactions

in plant development. *The Plant Cell*, 27, 44-63.

Schoonhoven L.M., Van Loon J.J.A. & Dicke M. (2005). Insect–Plant Biology. Oxford University Press.

Schulz M.H., Zerbino D.R., Vingron M. & Birney E. (2012). Oases: robust *de novo* RNA-seq assembly

across the dynamic range of expression levels. *Bioinformatics*, 28, 1086-1092.

Shorthouse J.D., Wool D., & Raman A. (2005). Gall-inducing insects – nature's most sophisticated

herbivores. *Basic and Applied Ecology*, 6, 407-411.

Stuart J.J., Chen M.S., Shukle R. & Harris M.O. (2012). Gall midges (Hessian flies) as plant pathogens.

Annual Review of Phytopathology, 50, 339-357.

Testerink C. & Munnik T. (2005). Phosphatidic acid: a multifunctional stress signaling lipid in plants.

Trends in Plant Science, 10, 368-375.

Tong Z., Gao Z., Wang F., Zhou J. & Zhang Z. (2009). Selection of reliable reference genes for gene

expression studies in peach using real-time PCR. *BMC Molecular Biology,* 10, 71.

Tooker J.F. & De Moraes C.M. (2008). Gall insects and indirect plant defenses: A case of active

manipulation? *Plant Signaling & Behavior*, 3, 503-504.

Tooker J.F. & Helms A.M. (2014). Phytohormone dynamics associated with gall insects, and their

potential role in the evolution of the gall-inducing habit. *Journal of Chemical Ecology*, 40, 742–753.

Vandermoten S., Harmel N., Mazzucchelli G., De Pauw E., Haubruge E. & Francis F. (2014). Comparative

analyses of salivary proteins from three aphid species. *Insect Molecular Biology*, 23, 67-77.

Will T., Tjallingii W.F. Thönnessen A. & van Bel A.J. (2007). Molecular sabotage of plant defense by aphid

saliva. *Proceedings of the National Academy of Sciences*, 104, 10536-10541.

Wu F.H., Shen S.C., Lee L.Y., Lee S.H., Chan M.T. & Lin C.S. (2009). Tape-*Arabidopsis* Sandwich-a simpler

Arabidopsis protoplast isolation method. *Plant Methods*, 5, 16.

Xue H.W., Chen X. & Mei Y. (2009). Function and regulation of phospholipid signalling in plants.

Biochemical Journal, 421, 145–156.

Yang J.L., Yang R., Cheng A.C., Jia R.Y., Wang M.S. & Zhang S.H. (2010). Five-minute purification of PCR

products by new-freeze-squeeze method. *Journal of Food, Agriculture & Environment*, 8, 32–33.

Yang Y. & Smith S.A. (2013). Optimizing *de novo* assembly of short-read RNA-seq data for phylogenomics.

BMC Genomics, 14, 328.

Zarate S.I., Kempema L.A. & Walling L.L. (2007). Silverleaf whitefly induces salicylic acid defenses and

suppresses effectual jasmonic acid defenses. *Plant Physiology*, 143, 866-875.

Zerbino D. & Birney E. (2008). Velvet: algorithms for de novo short read assembly using de Bruijn graphs.

Genome Research, gr-074492.

Zhao C., Escalante L.N., Chen H., Benatti T.R., Qu J., Chellapilla S., … & Richards S. (2015). A massive

expansion of effector genes underlies gall-formation in the wheat pest *Mayetiola destructor*. *Current*

Biology, 25, 613-620.

Zhao C., Shukle R., Navarro Escalante L., Chen M., Richards S., & Stuart J.J. (2016). Avirulence gene

mapping in the Hessian fly (*Mayetiola destructor*) reveals a protein phosphatase 2C effector gene family.

Journal of Insect Physiology, 84, 22-31.

Figure captions

Figure 1: Gene Ontology treemap for the salivary transcriptome of *Phloeomyzus passerinii*. The box size

correlates to the number of sequences isolated. Numbers between brackets indicate the number of

sequences identified. Green boxes indicate binding proteins, purple boxes indicate enzymes, red boxes

indicate structural constituents, the blue box indicates transporters and the brown box molecular

transducers. A Detailed list of the proteins can be found in the table S4.

Figure 2: Representative salivary sheaths secreted in artificial diets by *Phloeomyzus passerinii*. Sheaths

stained and observed after 24 h probing in an agarose diet: (A) sheath stained with Coomassie blue; (B)

black stained sheaths in diet containing 0.1% DOPA, indicating a phenoloxidase activity, note the dark

halo surrounding the upper sheath; (C) reddish stained sheath in diet immersed with 0.1% DAB and 0.1%

H_2O_2, indicating a peroxidase activity. Black bars represent 10 μm.

Figure 3: Heatmap of $\log_2$-fold changes of 43 poplar genes belonging to eight different physiological

processes or metabolic pathways (lower left box), after incubation of poplar protoplasts of two poplar

genotypes (Koster and I-214) with salivary proteins of two aphids (*Myzus persicae* (*M. p.*) and

Phloeomyzus passerinii (*P. p.*)). Non-host interactions are expected between both poplar genotypes and

M. persicae, while incompatible interactions are expected between *P. passerinii* and Koster and

compatible interactions between *P. passerinii* and I-214. Downregulation appears in blue and

upregulation in red. Gene code is presented below the heatmap, modalities (i.e. aphid x poplar genotype

combinations are presented on the right of the heatmap). Hierarchical clustering has been built with

Euclidean distances.

Figure 4: Fold changes of genes involved in auxin (left) and cytokinin (right) pathways of poplar

protoplasts collected from two poplar genotypes (I-214 and Koster), after incubation with salivary

proteins of two aphids (*Myzus persicae* and *Phloeomyzus passerinii*). Dots represent the average fold

change value, and bars represent credibility intervals.

Figure 5: Representative GUS assays of transgenic seedlings of *Arabidopsis thaliana pIAA2*::GUS, showing

whole plants (A, C, E, G, I and K), and root tips (B, D, F, H, J and L), after 3 h of incubation in 20 μM of IAA

(A and B), TE/Tween buffer (C and D), and 1 and 10 μg of salivary proteins of *Phloeomyzus passerinii* (E, F,

I and J) or *Myzus persicae* (G, H, K and L). Black bars represent 1 mm for whole plants (A, C, E, G, I and K)

and 10 μm for root tips (B, D, F, H, J and L). Five seedlings were used for each modality.

Figure 6: Representative GUS assays of transgenic seedlings of *Arabidopsis thaliana pARR16*::GUS,

showing whole plants (A, C, E, G, I and K), and root tips (B, D, F, H, J and L), after 4 h of incubation in 20

μM of BAP (A and B), TE/Tween buffer (C and D), and 1 and 10 μg of salivary proteins of *Phloeomyzus*

passerinii (E, F, I and J) or *Myzus persicae* (G, H, K and L). Black bars represent 1 mm for whole plants (A,

C, E, G, I and K) and 10 μm for root (B, D, F, H, J and L). Five seedlings were used for each modality.