TITLE PAGE

Experimental colitis drives enteric alpha-synuclein accumulation and Parkinson-like brain

pathology

Stefan Grathwohl[1], Emmanuel Quansah[2], Nazia Maroof[l], Jennifer A. Steiner[2], Liz Spycher[1], Fethallah

Benmansour[3], Gonzalo Duran-Pacheco[4], Juliane Siebourg-Polster[4], Krisztina Oroszlan-Szovik[1], Helga

Remy[1], Markus Haenggi[1], Marc Stawiski[1], Matthias Sehlhausen[4], Pierre Maliver[4], Andreas Wolfert[5],

Thomas Emrich[5], Zachary Madaj[2], Martha L. Escobar Galvis[2], Christoph Mueller[6], Annika

Herrmann[4], Patrik Brundin[2], and Markus Britschgi[1]

[1] Roche Pharma Research and Early Development, Neuroscience Discovery, Roche Innovation Center

Basel, F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, Basel, Switzerland

[2] Center for Neurodegenerative Science, Van Andel Research Institute, 333 Bostwick Ave. NE, Grand

Rapids, MI, USA

[3] Roche Pharma Research and Early Development, pREDi, Roche Innovation Center Basel, F.

Hoffmann-La Roche Ltd, Grenzacherstrasse 124, Basel, Switzerland

[4] Roche Pharma Research and Early Development, Pharmaceutical Sciences, Roche Innovation

Center Basel, F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, Basel, Switzerland

[5] Roche Pharma Research and Early Development, Pharmaceutical Sciences, Roche Innovation Center

Munich, Roche Diagnostics GmbH, Nonnenwald 2, 82377 Penzberg, Germany

[6] Institute of Pathology, University of Bern, Murtenstrasse 31, Bern, Switzerland

Corresponding authors:

Markus Britschgi, Roche Pharma Research and Early Development, Neuroscience Discovery, Roche

Innovation Center Basel, F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, Basel, Switzerland

Tel: +41 61 6879116

Email: markus.britschgi@roche.com

Patrik Brundin, Van Andel Research Institute, 333 Bostwick Ave. NE, Grand Rapids, MI 49503,

USA.

Tel: +1 616.234.5312

Email: patrik.brundin@vai.org

Abstract

Intraneuronal accumulation of α-synuclein (αSyn) is key in the pathogenesis of Parkinson's disease

(PD). Published studies suggest that this process begins in the enteric nervous system (ENS) and

propagates into the brain decades before clinical diagnosis of PD. The triggers and mechanisms

underlying the accumulation of αSyn remain unknown but evidence is growing, that immune

pathways and in particular colitis may play a critical role. Here we demonstrate that patients with

inflammatory bowel disease (IBD) exhibit αSyn accumulation in their colon. We then confirmed in an

experimental model of IBD that intestinal inflammation can trigger αSyn accumulation in the ENS of

wildtype and αSyn transgenic mice. We discovered that the type and degree of inflammation

modulates the extent of αSyn accumulation in the colon and that macrophage-related signaling limits

this process. Remarkably, experimental colitis at three months of age exacerbated the accumulation of

aggregated phospho-Serine 129 αSyn in the midbrain, including the substantia nigra, in 21-month but

not 9-month-old αSyn transgenic mice. This was accompanied by loss of nigral tyrosine hydroxylase-

immunoreactive neurons, another neuropathological hallmark of PD. Together, our data suggest a

critical role for intestinal inflammation in the initiation and progression of PD.

Introduction

Parkinson's disease (PD) is a progressively debilitating neurodegenerative disease affecting 1% of the population above 60 years (1). Typical symptoms are motor impairments including muscle rigidity, tremor, and bradykinesia. Neuropathologically, PD is hallmarked by dopaminergic cell loss in the substantia nigra (SN), a resultant loss of striatal dopaminergic signaling (2, 3), and the presence of intraneuronal inclusions called Lewy bodies and neurites (4, 5). Lewy pathology is enriched in α-synuclein (αSyn), a presynaptic protein that tends to misfold, aggregate, and become phosphorylated under pathological conditions (2, 3). Alpha-synuclein is also genetically linked to rare familial forms of PD and certain SNPs close to the αSyn gene (*SNCA*) locus are associated with an increased risk for developing PD (6). These findings in humans make αSyn a focal point of biomarker and drug development programs for PD.

Several years before the first appearance of motor features, many patients exhibit a variety of non-motor symptoms including constipation, sleep disorder, depression, and hyposmia (7–9). Co-occurrence of some of these non-motor symptoms increases the risk of progressing to PD even in otherwise healthy individuals in population-based and cohort studies (10–13). Constipation may be a particularly important non-motor feature of preclinical PD, with 28-61% of patients often exhibiting gastrointestinal dysfunction years before progressing to motor symptoms (9, 12, 14). This suggests an early connection between the status of the intestinal environment and the progression to PD (15). In line with this connection and due to the discovery of αSyn immunoreactive inclusions in neurons of the submucosal plexus in people with PD, Braak and colleagues have suggested an early involvement of the enteric nervous system (ENS) in the pathogenesis of PD (4, 5, 16, 17). It is currently unclear, whether occurrence of αSyn aggregates in the peripheral nervous system is specific for PD since individuals with dementia with Lewy bodies and normal people can also show such inclusions in various tissues (18–21). Yet based on their findings, Braak and colleagues hypothesized that αSyn immunoreactive inclusions first appear in the ENS and then progress over time to the brainstem regions, including the dorsal motor nucleus of the vagus nerve and midbrain areas (4, 5, 16, 17). Several studies in preclinical models have recently demonstrated a potential propagation of αSyn

pathology in the gut and the brain (22–26). Intriguingly, a clinical study reported that vagotomy in a

Danish population decreases the risk of PD (27). Determining the critical factors for αSyn pathology

in the ENS could provide insights into disease etiology, enabling the development of novel treatment

paradigms to slow disease progression. A key factor that could trigger ENS αSyn accumulation and

pathology spread is inflammation.

Interestingly, a recent finding in children with gastrointestinal inflammation suggests an immune

regulatory function of αSyn (28). Immune pathways are indeed activated in the brain and colon of PD

cases (29, 30). Also, several genes associated with an increased risk of developing PD have an

immune system-related function (31), and a recent genome-wide association study identified common

genetic pathways linking PD and autoimmune disorders (32). Most prominently, LRRK2, a major

genetic risk factor for PD also confers increased risk for developing inflammatory bowel disease

(IBD) (33) and is known to modulate the function of monocytes, macrophages and other immune cells

(34, 35). Intriguingly, IBD is now confirmed to be associated with an increased risk for developing

PD and specifically blocking the TNF pathway reduces this risk (36–38).

Here we tested the hypothesis that intestinal inflammation (e.g. colitis) triggers αSyn aggregation in

the ENS and the subsequent development of αSyn pathology in the brain. We discovered that patients

with IBD exhibited increased αSyn accumulation in the submucosal region of the colon. Experimental

colitis in wild type and αSyn transgenic mice demonstrated that the type and degree of inflammation

modulates the extent of αSyn accumulation in the colon and that macrophage-related signaling limits

this process. Remarkably, experimental colitis at 3 months of age exacerbated the accumulation of

aggregated αSyn in midbrain, including the substantia nigra, in 21-month but not 9-month-old αSyn

transgenic mice. This was accompanied by loss of nigral tyrosine hydroxylase-immunoreactive

neurons. Together, our data suggest a critical role for intestinal inflammation in the initiation and

progression of PD.

Results

IBD patients show αSyn accumulation in the ENS and local macrophages

Recent epidemiological data links inflammatory bowel disease (IBD) to an increased risk of

developing PD (36–38). In order to explore if IBD is associated with enteric αSyn accumulation we

performed immunohistochemistry for αSyn in cryo-sections from colonic biopsies of patients with

ulcerative colitis (UC, n = 11, mean age 31 years), Crohn's disease (CD, n = 11; mean age 35 years),

and from normal controls (NC, n = 8; mean age 51 years). We observed in eight UC cases various

degrees of αSyn accumulation, mostly in neuritic structures (**Figure 1A-C**). Interestingly, the eight

UC cases, and four patients with CD also showed marked intracellular αSyn staining in many

infiltrating monocytic cells. In contrast, only one NC subject showed a few immunoreactive cells.

This finding in human tissue suggests a potential role of local inflammation in the development of

enteric αSyn accumulation.

Experimental IBD exacerbates αSyn load in submucosal plexus of αSyn transgenic and wildtype
mice

During the process of further characterizing a (Thy1)-h[A30P]αSyn transgenic mouse line (39) we

detected human αSyn accumulation in all innervated organs that were analyzed (**Supplemental**

Figure 1). This included the myenteric and submucosal plexuses of the ENS, where it co-localized

with peripherin, a specific marker for peripheral nerves (**Figure 2A).** We observed an age-dependent

increase of baseline αSyn inclusion in both plexuses between the age of three and twelve months

(**Figure 2B**). Since this transgenic model has the capacity to increase the αSyn load in the ENS over

time we wanted to test the hypothesis whether IBD- related inflammation in the colon may trigger or

exacerbate local αSyn accumulation. Administration of dextran sodium sulfate (DSS) in the drinking

water in acute or chronic paradigms is a well-established mouse model of IBD colitis, resulting in

infiltration of leukocytes into the submucosa with various degrees of destruction of the colonic tissue

and the mucosa (40). We administered DSS at different concentrations and durations to (Thy1)-

h[A30P]αSyn transgenic mice (**Figure 2C**), resulting in leukocyte infiltration in a dose-dependent

manner (**Figure 2D and 3A**). Intriguingly, in the acute paradigm, 2.5% but not 1% DSS triggered

intracellular accumulation of αSyn in the submucosal plexus **(Figure 3A and 3B)**. In the chronic DSS

paradigm, a dose-dependent increase of αSyn load in the submucosal plexus was observed **(Figure**

3A and 3B). To confirm that this finding was not related to a transgene effect, an acute and a chronic

(consistent dose) DSS paradigm was respectively applied also in wildtype mice. In both paradigms,

we observed aggregation of endogenous murine αSyn in the submucosal plexus **(Figure 3C and 3D)**.

A separate experiment also confirmed that the observed effects of DSS could not be attributed to

increased gene expression of murine or the transgenic human αSyn **(Supplemental Figure 2)**.

Together, these results confirmed the validity of this experimental IBD paradigm to test the effect of

inflammation on αSyn accumulation in the ENS.

Colitis induced by peroral DSS but not by peritoneal administration of LPS aggravates αSyn

accumulation in colonic submucosal plexus of αSyn transgenic mice

In order to explore effects of different approaches to induce inflammation in or nearby the gut in

(Thy1)-h[A30P]αSyn transgenic mice, we compared the outcome of an acute 5% DSS in drinking

water with acute intraperitoneal LPS administration (**Figure 2C**). At d7, both agents had induced

variable degrees of leukocyte infiltration in the submucosa of the colon with a marked destruction of

the mucosa induced by only DSS (**Figure 2D and 4A**). Here, the DSS-exposed mice presented with

increased mean accumulation of αSyn in the ganglia of the submucosal plexus, higher than in the

2.5% DSS-exposed mice in the previous experiment (**Figure 4B-F**). In contrast, no change in αSyn

load was detected in the myenteric plexus, consistent with lack of leukocyte infiltration in this

anatomical region of the colon. Interestingly, LPS-induced inflammation did not increase αSyn

accumulation in the colonic nervous plexuses (**Figure 4C and E**). The two inflammatory insults

resulted, however, in a differential expression of cytokines, and consistent with leukocyte recruitment,

CCL2 was elevated in both (**Figure 4G and H**). Interestingly, in the LPS paradigm, mRNA for IL-10

was markedly elevated, whereas DSS strongly increased IL-6 and to some extent also IL-1β but not

IL-10. Together these results indicate that certain types of local inflammation can increase the

intracellular accumulation of aggregated αSyn in the periphery (i.e. in the colon).

Lack of Cx3cr1 signaling during DSS colitis aggravates αSyn load in the submucosal plexus of

αSyn transgenic mice

In both the IBD patients as well as the (Thy1)-h[A30P]αSyn transgenic mice that experienced acute

DSS colitis, we observed αSyn positive infiltrating cells (**Figure 5C and Figure 5D-F**). In the mice

these infiltrating cells were positive for the macrophage marker Iba-1 (**Figure 5G**). In order to explore

further the role of monocytes/macrophages in the accumulation of αSyn in our DSS model, we added

an experimental arm with (Thy1)-h[A30P]αSyn transgenic mice crossed with mice that have a

deletion of Cx3cr1 induced by an insertion of GFP (Cx3cr1-GFP knock-in mice). The CX3CR1-

CX3CL1 axis plays an important role in maintaining the function of the lamina propria macrophage

population of the gastrointestinal wall and lack of this signaling pathway in experimental colitis

models may either aggravate or ameliorate the induced pathology (41–43). In our experiment

exposure to DSS resulted in leukocyte infiltration in the Cx3cr1-deficient αSyn transgenic mice at a

minimally higher mean percent leukocyte area covered than for the Cx3cr1-competent group (**Figure**

3A). We observed, however, a significantly higher mean level of αSyn accumulated in the

submucosal plexus in αSyn transgenic mice lacking Cx3cr1 compared to αSyn transgenic mice

expressing Cx3cr1 ($p = 0.001$, two-way ANOVA with Tukey HSD post-hoc analysis; **Figure 3F**).

Consistent with the Cx3cr1-sufficient αSyn transgenic mice, no overt increased accumulation of αSyn

was observed in the myenteric plexus **(Figure 3D)**. These results indicate a potential link between

monocyte/macrophage signaling and αSyn accumulation in ENS in this experimental IBD model.

Systemic IL-10 ameliorates DSS-induced colitis and associated enteric αSyn accumulation in αSyn

transgenic mice

Interleukin-10 is an important regulator of monocytes/macrophages, and genetic ablation of IL-10

signaling or blocking IL-10 with specific antibodies was shown to enhance DSS colitis (44, 45).

Interestingly, we observed above that LPS-induced colonic leukocyte infiltration did not result in

increased accumulation of αSyn in the ENS of the colon and that the main difference in cytokine

expression between the DSS and LPS paradigms was increased IL-10 expression in the LPS group

(**Figure 3G and H**). To mimic the effect of higher levels of IL-10 in an acute model of DSS colitis,

we administered intravenously murine IL-10 (mIL10), which was recombinantly engineered onto two

different murine IgG variants to extend the half-life of mIL-10 in circulation (mIgG1(v1)-mIL10 and

mIgG1(v2)-mIL10, respectively). As shown previously, DSS induced a marked increase in leukocyte

infiltration and αSyn accumulation, which was similar in the untreated and control IgG treated group

(**Figure 4**). Both mIgG1(v1)-mIL10 and mIgG1(v2)-mIL10 significantly reduced leukocyte

infiltration despite administration of DSS (p<0.0001, one-way ANOVA with Tukey HSD post-hoc

analysis; **Figure 4A**). Both mIgG1-mIL10 variants also had reduced mean levels of human αSyn in

the submucosal plexus whereas mIgG1(v2)-mIL10 reached significance (p=0.02, one-way ANOVA

with Tukey HSD post-hoc analysis; **Figure 4B**). Interestingly, the significantly reduced αSyn

accumulation was associated with detectable serum exposure of mIgG1(v2)-mIL10 whereas

mIgG1(v1)-mIL10 was no longer detectable at the end of the in vivo phase, after 7d (**Figure 4**). These

results underline further an important role for the IL-10 pathway in keeping αSyn accumulation at a

reduced level throughout the course of experimental IBD. Together, the genetic and pharmacological

modulation of DSS colitis corroborates a relevant role for monocyte/macrophage pathways in the

development of αSyn accumulations in the ENS of the colon.

Experimental colitis-induced enteric αSyn accumulation persists for months and exacerbates brain

pathology and dopaminergic cell loss in αSyn transgenic mice

The previously highlighted hypothesis by Braak and colleagues associates αSyn brain pathology with

previous development of αSyn pathology in the ENS (5, 46). However, it is currently unknown

whether previous inflammatory conditions propel this ENS αSyn accumulation that may further

worsen αSyn aggregation in the brain. To test this, we first explored αSyn accumulation in the

submucosal plexus of (Thy1)-h[A30P]αSyn transgenic mice that had experienced DSS colitis

previously but had fully recovered two months later. As expected, leukocyte infiltration in the

submucosal plexus had returned to normal levels following the two-month recovery period

(**Supplemental Figure 3A**). Remarkably however, αSyn accumulation in the ganglia of the

submucosal plexus was still almost two-fold higher when compared to αSyn transgenic mice that were

not exposed to DSS, and this was exacerbated in αSyn transgenic mice deficient for Cx3cr1

(**Supplemental Figure 3B**).

To assess development of brain αSyn pathology, we exposed 3-month old hemizygous (Thy1)-

h[A30P]αSyn transgenic mice to DSS or normal drinking water (chronic increasing dose paradigm)

and then returned all to normal drinking water (**Figure 1C**). After aging up to 9 or 21 months of age

(i.e. mice aged for an additional 6 or 18 months after the chronic DSS paradigm), various brain

regions were analyzed for pathologically accumulated αSyn (proteinase K resistant, pSer129-αSyn

immunoreactive inclusions). Twenty-one month old mice that only received water during their

lifetime showed low levels of pathological αSyn inclusions in the brain (**Figure 6 and Supplemental**

Figure 4), which is consistent with previous observations in the brain of these hemizygous (Thy1)-

h[A30P]αSyn transgenic mice (47). In marked contrast, the majority of 21-month-old and previously

DSS-exposed αSyn transgenic mice presented with higher degrees of pathological αSyn in the same

brain areas where mice on water displayed much lower pSer129-positive immunoreactivity.

Intriguingly, mice that were aged to 9 months presented with extremely low levels of pathological

αSyn inclusions in the brain (**Figure 6 and Supplemental Figure 4**). For the mice that aged up to 21

months, the accumulated αSyn pathology recorded in the SN was accompanied by a significant loss of

tyrosine hydroxylase (TH) and Nissl positive cells ($p \leq 0.05$, Student's T-test; **Figure 7**). Together,

DSS colitis at a young age caused an age-dependent exacerbation of αSyn inclusion pathology in the

brain and a loss of nigral dopaminergic cells, similar to the human neuropathological hallmarks of PD.

Discussion

Currently, there is no therapy for PD available to slow or stop its progression and an obstacle in the

quest to develop one is that we do not understand the triggers of PD (48). Intraneuronal accumulation

of aggregated αSyn (i.e. Lewy bodies and neurites) is a key neuropathological hallmark, thus the

distribution of Lewy pathology in postmortem brain tissues has been used for disease staging in PD

(2, 4, 46). Abnormal accumulation of αSyn has also been observed in the peripheral nervous system in

PD and in some individuals at risk of developing the disease (49, 50). Similar to this finding in

people, αSyn immunoreactive inclusions have also been detected in the ENS of a transgenic mouse

model prior to changes in the CNS (51). It is also suggested, based on preclinical models and

postmortem pathology in various organs including the brain, that αSyn pathology propagates in a

temporospatial and prion-like manner (2, 4, 5, 52, 53). However, the initial factors triggering αSyn

aggregation are yet to be established (48). Moreover, the involvement of peripheral stimuli in the

aggregation and pathogenic spread of αSyn is only beginning to unravel.

In this study, we provide evidence that patients with IBD have increased αSyn accumulation in their

ENS (**Figure 1**) and that a IBD-like inflammation (DSS colitis) triggers αSyn accumulation in the

ENS of wildtype mice and in a transgenic mouse model of PD (**Figure 3**). Interestingly, in IBD

patients and in the mouse model of IBD, macrophages filled with αSyn could be observed in the

inflamed area (**Figure 5**). Aggravation of enteric αSyn accumulation in αSyn transgenic mice lacking

Cx3cr1 signaling and amelioration of inflammation and enteric αSyn load by systemic IL-10 suggest a

modulatory role of monocytes/macrophages in this process (**Figure 4 and 6**). Remarkably, we further

observed that the aggravation of αSyn accumulation in the ENS persisted even after two months of

recovery from DSS colitis, indicating that the observed effect is not a transient phenomenon

(**Supplemental Figure 3**). Most importantly, 18 months but not 6 months post DSS colitis, αSyn

trasngenic mice had developed a Parkinson-like pathology in their brain (**Figure 7 and 8 and**

Supplemental Figure 4). This included elevated proteinase K resistant pSer129-αSyn pathology in

the midbrain in the midbrain, including the substantia nigra, and other brain regions and an average

decrease of 30-50% of TH- and Nissl positive cells in the SN. Together, this colitis mouse model

recapitulated the proposed accumulation of αSyn in the ENS long before clinical diagnosis of PD in

humans and the subsequent age-related development of αSyn pathology in the brain, together with the

loss of nigral dopaminergic neurons.

We established that a mechanism by which peripheral inflammation promote αSyn accumulation in

the colon potentially involves monocytes and macrophages. Interestingly, both peroral DSS and

peritoneal LPS administration provoked strong local immune reactions resulting in leukocyte

infiltration into the submucosa of the colon. This inflamed region is anatomically right next to the

submucosal plexus and is separated by a muscular layer from the myenteric plexus (**Figure 2**). This

might explain why the effect of αSyn accumulation was only observed in the nerves of the

submucosal plexus but not in the ganglia of the myenteric plexus in our (Thy1)-h[A30P]αSyn

transgenic mouse model of colitis. The mechanism underlying how intraperitoneally administered

LPS leads to submucosal leukocyte infiltration probably involves the monocyte attractant chemokine

CCL2 (**Figure 4**), but the specifics remain unclear. CCL2 was also upregulated in the colon of our

DSS model. However, in contrast to peritoneal LPS, where infiltration of macrophages followed a

patchy pattern, DSS-related macrophage infiltration was distributed across the whole submucosa.

Also, the primary mechanism of peroral administered DSS is to destruct the mucosa of the colon,

similar to some forms of ulcerative colitis, resulting in the transient disintegration of the intestinal

epithelial barrier. The subsequent immune response to the infiltration of commensal bacteria evoked

also in our (Thy1)-h[A30P]αSyn transgenic mouse model the expression of cytokines such as IL-1β

and IL-6. This upregulation was absent in the LPS paradigm that had an intact mucosa. By acting on

tight junctions, IL-1β and IL-6 can increase intestinal barrier permeability (gut leakiness), facilitating

the recruitment of additional immune cells to the site of the inflammatory signal eventually

culminating in widespread immune activation (54, 55). Consistent with the breach of barrier

permeability in our mouse model, some PD patients express increased colonic cytokines such as IL-

1β, IL-6 and TNF, occurring together with increased intestinal permeability (29, 56). In this context, it

is striking that IBD patients on anti-TNF therapy have a reduced risk of developing PD (37). Here we

demonstrate that patients with IBD also present with αSyn accumulations in the ENS as well as in

infiltrating leukocytes nearby. In this regard, it is interesting to note that mucosal macrophages with

intralysosomal αSyn content were previously described in the intact human appendix (57). These

macrophages were in close proximity to the axonal varicosities of the vermiform appendix which

showed an enriched staining for αSyn in the mucosal plexus. Indeed, we recently demonstrated that

the vermiform appendix contains aggregated and truncated αSyn that has the propensity to aggregate

recombinant αSyn *in vitro* (18). Also, monomeric and oligomeric αSyn reportedly act as

chemoattractants for neutrophils and monocytes, enhancing the maturation of dendritic cells in the

ENS (28, 58). With such a role in intestinal immunity therefore, it is possible that the strong tissue

destruction induced by DSS in the present study released αSyn, which perhaps served as a

chemoattractant for monocytes. It is likely that increased abundance of extracellular αSyn and altered

intestinal permeability, along with the DSS-evoked inflammatory response may have provided an

enabling milieu allowing further αSyn accumulation in the ENS of the colon (59). However, the

pathways leading to αSyn accumulation in macrophages remain to be elucidated. Mechanistically, it

has been shown that excess endogenous αSyn compromises phagocytic activity in human iPSC-

derived macrophages (60). Macrophages and other immune cells are also regulated by LRRK2, an

established risk gene for PD and IBD. It will be interesting to explore in our colitis paradigm the

effect of LRRK2 (e.g. its effect on autophagy in macrophages (34, 35, 61)) and other pathways in

macrophages on αSyn accumulation.

Most profoundly, besides the accumulation of αSyn in the submucosal plexus, we found that a single

chronic DSS paradigm at young age exacerbated αSyn pathology in the CNS of (Thy1)-h[A30P]αSyn

transgenic mice later in life (**Figure 7**). But how does this severe αSyn inclusion pathology develop in

the brain of these mice? One hypothesis is that the brain αSyn pathology observed in this study could

be due to direct effects of peripheral immune activation on the brain and brain vasculature.

Hypothetically, colitis induced by DSS or LPS could compromise both active and passive components

of the blood-brain barrier, which may render the CNS vulnerable to immune cells from the periphery

(62, 63). Alternatively, certain peripheral triggers can directly affect microglial activity in the brain.

For instance, short-chain fatty acids derived from gut microbiota appear to regulate function and

maturation of microglia in the mouse CNS (64) and inflammatory mediators from gut microbiota

could be carried in the bloodstream and induce pathological and behavioral changes in an αSyn

transgenic mouse model (65). Moreover, rats and nematodes develop αSyn inclusions after exposure

to the bacterial amyloid protein curli, a protein which stimulates microgliosis, astrogliosis, and

secretion of IL-6 and TNF (66). Intriguingly, a recent study reported that peripherally applied

inflammatory stimuli induce acute immune training (that exacerbates β-amyloid pathology) and

immune tolerance in the brain that reprograms microglia, an effect which can persist for at least six

months (67).

Another hypothesis is that the brain αSyn pathology observed may have accumulated following the

transfer of pathogenic αSyn seeds from the gut via the vagal nerve. It is already established in rodents

and non-human primates that αSyn pathology in the CNS can propagate from cell to cell, perhaps in a

prion-like fashion (52, 53, 68, 69) and that certain pathogenic forms of αSyn may serve as templates

in recipient cells, seeding and inducing further aggregation of endogenous αSyn in these cells (68, 70–

72). Interestingly, several studies in preclinical *in vivo* models have demonstrated that pathogenic

αSyn seeds can be transferred from the peripheral nervous system to the CNS. Indeed, aggregated

recombinant αSyn injected intraperitoneally, intramuscularly or into the gastric wall of certain mouse

models of PD results in αSyn inclusions in the CNS (24, 73), with the latter being transmitted via the

vagal nerve (74). Another study showed that vagotomy in mice prevents propagation of pathology

into the CNS (74), a finding that is parallel to a peripheral rotenone-related αSyn rat model in which

vagotomy reduced propagation of αSyn pathology (75), and to a human study that reported that

vagotomy in a Danish population also decreases the risk of PD (27). Similarly, after injection of PD

brain lysate and recombinant αSyn into the intestinal wall αSyn pathology trafficked through the

vagal nerve to the dorsal motor nucleus of the vagus (23). Others have shown that virus-based

expression of human αSyn in the medulla oblongata resulted in a transfer of αSyn towards rostral

brain regions in a rat model (76). The findings in these previous studies, along with the present

findings recall the Braak hypothesis and highlight the critical connection of the vagal nerve. Indeed, in

the present study, αSyn pathology was much more prominent in the reticular nucleus (including the

vagal area) and midbrain areas (compared to the rostral areas) 18 months post DSS colitis, suggesting

the potential involvement of this nerve in the accumulation of αSyn in the CNS.

Here, we report that individuals with IBD exhibit αSyn accumulation in the colon concomitant with

infiltrating monocytes/macrophages positive for αSyn. Employing a experimental model of IBD, we

provide proof-of-principle demonstrating that αSyn accumulated in the colon of (Thy1)-h[A30P]αSyn

transgenic and wildtype mice upon DSS colitis and that this process can be modulated by

monocyte/macrophage-related signaling. We further demonstrate that chronic DSS colitis in young

(Thy1)-h[A30P]αSyn transgenic mice robustly exacerbated αSyn accumulation in the aged brain

which was accompanied by reduced numbers of TH- and Nissl positive neurons in the nigra. The

present findings are in consonance with retrospective and cohort studies demonstrating a link between

IBD and PD in various populations including those in Taiwan, Denmark and the US (36, 37, 77),

although this may not be the case in all populations (78). The exact mechanisms connecting the gut

pathology to the pathology in the brain including the role of monocytes/macrophages, remain to be

established. In summary, this novel mouse model of a potential preclinical stage of PD recapitulated

the accumulation of αSyn in the ENS and the subsequent age-related development of brain pathology.

Together, our data suggest a critical role for intestinal inflammation in the initiation and progression

of PD.

Methods

Mice

Male C57BL/6 wild type mice (Jackson Laboratories, Bar Harbor, USA), hemizygous Tg(Thy1-

SNCA*A30P)18Pjk ((Thy1)-h[A30P]αSyn) (39) and Tg(Thy1-SNCA*A30P)18Pjk crossed with

Cx3cr1tm1Litt ((Thy1)-h[A30P]αSyn /CX3CR1-def; homozygous for Cx3cr1-GFP knock-in allele;

(79) transgenic mice were used for the study. (Thy1)-h[A30P]αSyn transgenic mice express mutant

human αSyn under the neuron selective Thy1 promoter. (Thy1)-h[A30P]αSyn transgenic mice were

crossed to Cx3cr1-def transgenic mice which express eGFP replacing fractalkine gene expression. All

mice were maintained on a C57BL/6 background for more than 10 generations and under specific

pathogen-free conditions. To the extent possible, littermates were used in the experiments. Health

status was monitored daily during experiments.

Human Subjects

Samples from patients with Crohn's disease (CD), ulcerative colitis (UC) or neither were provided by

the tissue bank, Institute of Pathology, University of Bern. Briefly, specimens were obtained from

patients who underwent surgical procedures at the University Hospital (Inselspital) in Bern,

Switzerland between 2004 and 2011. Three selected male patients previously clinically diagnosed

with UC with a reported disease duration > 6 year (n=3) and undergoing steroid therapy combined

with either metronidazole or mesalazine. CD patients were of mixed gender and aged 22-56 years

ranging from 2 months to 11 years post disease diagnosis undergoing treatment with either infliximab

or mesalazine in combination with steroids. Healthy subjects were of mixed gender with no report of

inflammatory bowel disease, aged 40-59. All samples contained the mucosa and submucosa regions

including minor parts of the circular muscle layer. Following surgical removal, tissue samples were

immediately immersed in O.C.T. compound (VWR International GmbH, Dietikon, Switzerland),

frozen in liquid nitrogen and stored at -80°C. Diagnosis of disease status was made according to

established criteria for histopathological analysis.

Experimental IBD in mice with DSS and LPS

Paradigms for the induction of inflammation were either 1 week (acute) or 3-4 weeks (chronic) with

or without an incubation phase of 2, 6 or 18 months post application (**Figure 2**). Acute systemic

inflammation was induced by intraperitoneal Lipopolysaccharide (LPS) application (80) of 0.5 mg/kg

in 100 µl injection volume on day 1 and 4 (Sigma-Aldrich Chemie GmbH, Steinheim, Germany, LPS

055:B5). Acute colitis was induced by application of 36-50kDa Dextran Sodium Sulfate (DSS) (81)

(160110, MP Biomedicals, LLC, Illkirch, France) at 0%, 1%, 2.5% or 5% in autoclaved drinking

water for 5 continuous days respectively, followed by 2 days of water (1 DSS application cycle).

Chronic colitis was achieved by 4 repeating DSS application cycles. The DSS concentration during 4

weeks of chronic colitis was either 1% or 2.5% for 4 weeks or 2.5%-4% raised 0.5% every week for 4

weeks. Mice from same littermate group were randomized per cage into vehicle and inflammation

inducing agent.

IL-10 treatment and exposure measurement

Two different forms of mouse IgG bound murine IL-10 (mIgG(v1)-mIL10 and mIgG(v2)-mIL10)

were diluted in pre-prepared sterile formulation buffer comprised of 0.5% mouse serum supplemented

with 25mM citrate, 300mM arginine to a final concentration of 0.75mg/ml and the pH adjusted to 6.7

on the day of application. Each mouse was treated once with 150 µg i.p concurrently with the

initiation of the acute colitis paradigm with 5% DSS. The concentrations of mIgG-mIL10 fusion

proteins in murine serum samples were determined by enzyme-linked immunosorbent assays (ELISA)

specific for the Fab moiety of the administered mIgG-mIL10 fusion protein. Biotinylated mIgG-

mIL10-specific target molecules were used for capturing, goat anti-mIg IgG-HRP conjugate and

peroxidase substrate ABTS was used for quantitative detection of mIgG-mIL10 fusion proteins.

Immunohistochemistry

Animals were injected with a lethal dose of pentobarbital 150 mg/kg. Upon full anesthesia, mice

received transcardial perfusion with room temperature phosphate buffered saline (PBS). For

biochemical and immunohistochemical analysis, one section of either the proximal colon was fresh

frozen and stored at -80°C or post-fixed in 4% paraformaldehyde (PFA) solution for 24 h. Following

post-fixation, organs were incubated in 30% sucrose/PBS at 4°C for at least 48 h before further

processing. Subsequently, enteric tissue was cryotome-sectioned to 35 μm thick longitudinal sections

(approx. 1 cm length). The brain was collected and post-fixed for 24h in 4% PFA followed by 30%

sucrose in phosphate buffer until cryo-sectioning of floating sections at 40 μm. Histological analysis

of the colon was performed using standard hematoxylin staining. Immunohistochemical staining was

accomplished using the Vectastain Elite ABC Kits and Peroxidase Substrate Kit SK-4100 (Vector

Laboratories, Burlingame, CA, USA) or fluorescently labelled secondary antibodies (Alexa coupled

to dye 488, 555 or 647, Life Technologies, Zug, Switzerland). The following primary antibodies have

been used for overnight incubation at a dilution of 1:1000: monoclonal antibody to human α-synuclein

(sc-12767, Santa Cruz Biotechnology, Heidelberg, Germany), polyclonal antibody to murine α-

synuclein (Syn1, BD Transduction Laboratories, Allschwil, Switzerland), polyclonal antibody to the

peripheral neuronal marker Peripherin (Millipore Corporation, Billerica, MA, USA), and polyclonal

antibody to macrophage marker Iba1 (Wako Chemical GmbH, Neuss, Germany). To detect

phosphorylated αSyn (pSer129 pathology) in the free-floating brain sections, monoclonal antibody

(ab51253, Abcam, Cambridge, USA) to human αSyn was used at a dilution of 1:10000. Prior to the

pSer129 staining, the free-floating brain sections were incubated for 10 min at room temperature in a

phosphate buffered saline solution containing 10 μg/mL proteinase K (Cat # 25530015; Invitrogen,

California, USA). TH-immunoreactive cells were detected using a polyclonal antibody (657012,

Millipore Sigma) at a dilution of 1:1000. To measure the density of Nissl-positive cells, the TH-

stained cells were counter-stained with Cresyl violet. The slides were incubated in 0.1% Cresyl violet

solution for 9 min and then dehydrated in 95% and 100% ethanol and then xylene prior to

coverslipping with Cytoseal 60 mounting media (Thermo Fisher Scientific). Quantifications of the

blind-coded TH/Nissl stained slides were done using Stereoinvestigator (version 2017.01.1; MBF

Bioscience, Williams, VT, USA) on Imager M2 microscope (ZEISS) coupled to a computer. We

analyzed 5-7 nigral sections per animal, and a total of 7-8 animals per treatment group. We outlined

the substantia nigra pars compacta and counted every TH-immunoreactive and Nissl-positive cell in

that area and computed the number of cells per section, generating the mean cell density per animal.

We then calculated the mean density of cells per treatment group and analyzed the data using

unpaired Student's T-test after confirming normality and homoscedasticity in Prism 7.0 (GraphPad

Software).

Imaging and stereological quantification of αSyn deposits in enteric nervous system

Imaging and stereological quantification was performed on a Zeiss Axio Imager Z2 fluorescence

microscope (Carl Zeiss AG, Jena, Germany). Leica TCS SP5 confocal system using an HCX PL APO

CS 40x 1.3 oil UV or an HCX PL APO LB 63x 1.4 oil UV objective was utilized for image recording.

Accumulation of αSyn in the ENS was assessed on a random set of 3 adjacent 35 μm thick, αSyn-

immunostained sections comprising the myenteric and submucosal neuronal plexuses. Analysis was

performed with the aid of Stereologer software (Stereo Investigator 10, MBF Bioscience, Williams,

VT, USA) as described previously (82). In the myenteric plexus ganglion volume was defined by

multiple outlined plexuses containing a range of 5-20 neuronal cells and quantified by the optical

fraction fractionator technique. In contrast to the myenteric plexus, the submucosa consists of

compact plexuses with 1-5 cells including interconnecting neurites. Therefore, the entire submucosa

was set as region of interest, analyzed with the area fraction fractionator technique. Results of the

submucosal plexus are displayed by percent area containing αSyn deposits. For the IL-10 experiment,

αSyn positive granules from immunofluorescence images were counted for each image. Granules

were filtered based on having a size between 12 and 50000 pixels and a minimal intensity value

greater than 300. The filtering step was included to exclude small background particles and

macrophages (very large spots). The counts were then aggregated to the animal level by summing the

granule counts of all images per animal and then normalizing for (i.e. dividing by) the number of

images for a given animal. Upon exploratory data analysis two animals were excluded: one mouse

because it only had one image and another due it being an outlier, based on its infiltration score and

image data.

Blinding

For analyses of colon and brain tissue on slides, a second individual assigned unique codes to

stained slides. Therefore, the experimenter conducted the analyses blinded to the identity of the

mice. For randomization of treatment groups see above.

Quantification of leukocytes infiltration

To determine the leukocyte covered area in the colon after LPS or DSS application, three adjacent

hematoxylin stained sections were quantified. Total area of colon sections and localizations of

leukocyte assemblies within the tissue architecture were identified and outlined utilizing Stereologer

Software (Stereo Investigator 6, MBF Bioscience, Williams, VT, USA). Percentage of leukocyte

covered area has been set in proportion to total area of the analyzed colon section. For the IL-10

experiment, hematoxilin stained colon slices were examined by an expert pathologist blinded to

treatment conditions. A score of 0-3 was assigned to each section for each of the 3 layers lamina

propria, submucosa and muscularis based on the degree of inflammatory infiltration. A score of 0

denoted no inflammation and a score of 3 indicated extensive infiltration. The mean of the values for

all 3 layers was taken as the final measure of leukocyte infiltration per mouse.

Quantification of αSyn/Iba1 double positive macrophages

The number of α-syn+/Iba1+ positive cells was evaluated by quantification of 10 random regions in 2

adjacent sections of the proximal colon. The region of interest was set to contain the myenteric

plexus/circular muscle layer and the submucosal plexus. Cells were assessed for positive αSyn

staining and concomitant co-localization with the macrophage marker Iba1 was quantified.

Scoring of pSer129 pathology and brain heatmap

We evaluated pSer129 pathology on a full series of immunostained coronal sections from 10 mice per

treatment group (i.e. water vs. DSS-treated groups) on blind-coded slides using a previously described

method (83). We visualized pathology from one hemisphere of all brain sections (apart from the

olfactory area) using NIKON Eclipse Ni-U microscope and assigned scores ranging from 0 to 4 to

each brain area based on the relative abundance of PK-resistant pSer129-positive inclusions (i.e. cell

bodies and neurites). In this case, 0 = no aggregates, 1 = sparse, 2 = mild, 3= dense, 4 = very dense.

For the heatmap, we obtained the average score values of each brain area for each treatment group.

The average data for each treatment group (n=10/ group) was then represented as a heatmap in a

sagittal mouse brain background (http://atlas.brain-

map.org/atlas?atlas=2#atlas=2&structure=771&resolution=16.75&x=

7755.7470703125&y=3899.625&zoom=-3&plate=100883867&z=5).

Densitometry of pSer129 αSyn brain pathology

The density of pSer129 pathology in 12 major brain areas (reticular nucleus, pontine reticular nucleus,

periaqueductal gray, gray and white layer, reticular formation, substantia nigra, ventral tegmental

area, thalamus, hypothalamus, central amygdala, pallidum and striatum) were determined in the water

and DSS-treated animals. NIKON Eclipse Ni-U microscope was used to acquire 20x magnification

images (without condenser lens) from all the indicated brain areas, using the same exposure time for

all images. In all cases, images were acquired on three sections separated by 420 μm intervals

(localized between Bregma). We then processed the acquired images using Image J64 (84), created a

mask (to exclude background) that redirects to the original image for analysis, measured the total area

and the mean grey value of the area that had inclusions. For brain areas such as periaqueductal gray

that do not fill the entirety of the field to be analyzed, we drew a contour of the area and the analysis

was performed only within that contoured area. We subsequently calculated the grey value of the area

per square pixels for each image (i.e. $A.U./px^2$ = mean grey value x area stained/total area assessed).

Based on this, we calculated the average grey value per square pixels for each brain area for each

animal (n = 6 mice/group), and then extended this calculation to determine the average grey value per

square pixels for each treatment group and each of the twelve brain areas of interest.

mRNA expression

To assess mRNA expression levels from the proximal colon, RNA was extracted from fresh frozen

tissue with MagnaLyser green beads (Roche Diagnostics, Mannheim, Germany) and Qiazol Lysis

(Reagent cat.no.79306, Hilden, Germany) purified on MagnaPure LC (HP Kit no.03542394001, F.

Hoffmann - La Roche AG, Rotkreuz, Switzerland) and amplified via real-time PCR (4ng

RNA/reaction; Lightcycler 480, Roche Diagnostics Corporation, Indianapolis, USA). Amplification

of mRNA was performed by using TaqMan probes for human or murine specific α-synuclein and for

selected cytokines/chemokines (Applied Biosystems Europe B.V., Zug, Switzerland). Target mRNA

was normalized to tissue-specific murine GAPDH levels and displayed as relative expression after 30

amplification cycles.

Statistics

Statistical analysis of gut pathology and inflammation was performed using GraphPad Prism 6.04 or

7.0 software (GraphPad Software, Inc. La Jolla, CA, USA). The results are expressed as mean values

± standard errors of the mean (SEM). Student's T-test (or Welch's T-test for unequal variances) was

used to compare two groups and ANOVA was used for multi-comparison of groups followed by

Tukey HSD post-hoc analysis. For the statistical analysis of the pSer129 αSyn brain pathology,

negative-binomial mixed-effects models with a random intercept for each sample were used to

analyze the dataset via the 'lme4' (http://lme4.r-forge.r-project.org/) package in R v 3.4.4. To analyze

the pSer129 αSyn cell count dataset, an offset for the total area examined was included to model the

densities. Linear contrasts with false discovery rate (FDR) adjustments were then used to test our

hypotheses and account for multiple testing (for brain area and experimental group). Like the pSer129

dataset, the Iba-1/αSyn-double positive dataset were analyzed using negative-binomial regression and

Tukey HSD adjusted contrasts to test our hypotheses.

For the mRNA expression analysis, data quality was assessed by inspecting the distribution of Cp

values of reference endogenous genes across samples, by inspecting the level of Cp variation between

technical replicates and by exploring the samples multivariate signal distribution as in a principal

component analysis. Relative gene expression levels were expressed as $2^{-(Cpgene - CpRef)}$. Statistical

analyses to assess the effect of the experimental conditions on the log2 gene expression levels were

done with linear models using the *limma* package (Bioconductor/R, Smyth, 2005). These analyses

were implemented in R v3.1.1.

For the statistical modelling of the effects of the IL-10 treatment on αSyn counts, as well as

infiltration scores, the levels for IgG1(v1)-IL10 and IgG1(v2)-IL10 treatment were compared to the

positive (vehicle/DSS) control. Additionally, since levels of the control antibody treatment (IgG1(v1))

were very similar to the positive control, the two groups were pooled in further contrasts in which

effects of individual antibodies or control IgG was assessed. For αSyn counts, a linear model on the

treatment groups with one-degree freedom contrasts was applied. For the infiltration score a Kruskal-

Wallis test, with the same contrasts was used.

Study approval

The human subjects' study was conducted with the approval of the local Ethical Committee in Bern

No. 47/04. Written informed consent was obtained from each patient. The animal experiments were

approved by a Roche internal review board and the local authorities.

Author contributions

S.G., N.M. and L.S. planned and performed the in vivo experiments, colon immunostaining, analysis, and quantification; S.G. and N.M. drafted a first version of the manuscript; E.Q. performed, imaged, quantitated pSer129, TH and Nissl staining in the brain sections, and drafted a more advanced version of the manuscript with J.A.S., who also provided helpful discussion. F.B. and K.O.S. supported the image acquisition and image analysis for the colon samples; M.St. performed imaging and data analysis of experiments with wildtype mice; G.D.P. and J.S.P. performed statistical analysis of the DSS experiments; H.R. and M.H. performed mRNA analyses; M.Se. trained S.G. and L.S. on mouse necropsy and supported their work; P.M. performed expert pathology staging on leukocyte infiltration; T.E. and A.W. provided mIgG-mIL-10 fusion proteins and measured serum exposure; Z.M. performed statistical analysis for the pSer129 αSyn immunohistochemistry data. M.L.E.G. provided helpful discussion and project planning. A.H. co-mentored S.G. and N.M., performed expert pathology staging on leukocyte infiltration and contributed to experimental planning. C.M. trained S.G. on the colitis model, provided human tissue and expert input on the experimental IBD model. M.B. and P.B. co-mentored Roche Postdoc Fellows S.G. and N.M., conceived and oversaw the study, and performed experimental planning; M.B., P.B. and E.Q. wrote the final version of the manuscript.

Acknowledgments

We acknowledge the human donors for providing tissue used in this study. We thank Drs. L. Ozmen, A. Bergadano, and A. Su for their tremendous support in maintaining the mouse colony and establishing of relevant animal experiment licenses, and we are grateful to the animal care takers, veterinarians and many unnamed staff at Roche for their valuable work with the mice in this study. In addition, at Roche we thank Dr. K.G. Lassen for critical input to the paper, Dr. C. Ullmer for co-mentoring S.G. and providing scientific input, Dr. L. Collin for helping with confocal imaging and we are grateful to Dr. T. Kremer, N. Haenggi, D. Mona, A. Girardeau, and J. Messer for providing support in tissue dissections and G. Walker and R. Lauria for technical support. Ms. E. Schulz from VARI assisted with immunostaining of the brain tissue. We thank the Contract Research Organization Frimorfo for carefully sectioning the brains for this study. We acknowledge Drs. L. Gaudimier (née

Chicha) and F. Pan-Montojo for scientific discussions early in the project and Dr. W. Zago from

Prothena for valuable scientific input. P.B. reports relevant grants from NIH (R01DC016519-01 and

5R21NS093993-02), Department of Defense (W81XWH-17-1-0534), The Michael J. Fox Foundation

for Parkinson's Research, and Cure Parkinson's Trust. Finally, we thank the Roche Postdoctoral

Fellowship Program for supporting S.G. and N.M.

Conflicts of Interest

At the time of the study S.G. and N.M. were Roche Postdoctoral Fellows employed by Roche and

L.S., F.B., G.D.P., J.S.P., K.O.S., H.R., M.H., M.Se. M.St., P.M., A.W., T.E., A.H. and M.B. are or

were fulltime employees or trainees at Roche and they may additionally hold Roche stock/stock

options. S.G. and L.S. are currently employees of Neurimmune AG, Schlieren, Switzerland. P.B. has

received commercial support as a consultant from Renovo Neural, Roche, Teva, Lundbeck A/S,

AbbVie, NeuroDerm, Fujifilm Cellular Dynamics, Living Cell Technologies, IOS Press Partners, and

Axial Biotherapeutics. Additionally, P.B. has received commercial support as a consultant from

Renovo Neural, Roche, Teva, Lundbeck A/S, AbbVie, NeuroDerm, Fujifilm Cellular Dynamics,

Living Cell Technologies, IOS Press Partners, Axial Biotherapeutics and CuraSen. Additionally, P.B.

has received commercial support for grants/research from Renovo, Roche, Teva, and Lundbeck. P.B.

has ownership interests in AcouSort AB. The other authors do not have conflicts of interest with

regard to this research.

References

1. Tysnes O-B, Storstein A. Epidemiology of Parkinson's disease. *J Neural Transm (Vienna)*

2017;124(8):901–905.

2. Spillantini MG et al. Alpha-synuclein in Lewy bodies. *Nature* 1997;388(6645):839–840.

3. Spillantini MG, Goedert M. Neurodegeneration and the ordered assembly of α-synuclein. *Cell*

Tissue Res. 2017;373(1):137–148.

4. Braak H et al. Staging of the intracerebral inclusion body pathology associated with idiopathic

Parkinson's disease (preclinical and clinical stages). *J. Neurol.* 2002;249 Suppl 3:III/1-5.

5. Del Tredici K, Braak H. Lewy pathology and neurodegeneration in premotor Parkinson's disease.

Mov. Disord. 2012;27(5):597–607.

6. Polymeropoulos MH et al. Mutation in the alpha-synuclein gene identified in families with

Parkinson's disease. *Science* 1997;276(5321):2045–2047.

7. Schrag A, Horsfall L, Walters K, Noyce A, Petersen I. Prediagnostic presentations of Parkinson's

disease in primary care: a case-control study. *Lancet Neurol* 2015;14(1):57–64.

8. Gaenslen A, Swid I, Liepelt-Scarfone I, Godau J, Berg D. The patients' perception of prodromal

symptoms before the initial diagnosis of Parkinson's disease. *Mov. Disord.* 2011;26(4):653–658.

9. Pont-Sunyer C et al. The onset of nonmotor symptoms in Parkinson's disease (the ONSET PD

study). *Mov. Disord.* 2015;30(2):229–237.

10. Berg D et al. The PRIPS study: screening battery for subjects at risk for Parkinson's disease. *Eur.*

J. Neurol. 2013;20(1):102–108.

11. Postuma RB, Gagnon J-F, Bertrand J-A, Génier Marchand D, Montplaisir JY. Parkinson risk in

idiopathic REM sleep behavior disorder: preparing for neuroprotective trials. *Neurology*

2015;84(11):1104–1113.

12. Abbott RD et al. Frequency of bowel movements and the future risk of Parkinson's disease.

Neurology 2001;57(3):456–462.

13. Savica R et al. Medical records documentation of constipation preceding Parkinson disease: A

case-control study. *Neurology* 2009;73(21):1752–1758.

14. Mahlknecht P, Seppi K, Poewe W. The Concept of Prodromal Parkinson's Disease. *J Parkinsons*

Dis 5(4):681–697.

15. Houser MC, Tansey MG. The gut-brain axis: is intestinal inflammation a silent driver of

Parkinson's disease pathogenesis?. *npj Parkinson's Disease* 2017;3(1):3.

16. Braak H, Del Tredici K. Neuroanatomy and pathology of sporadic Parkinson's disease. *Adv Anat*

Embryol Cell Biol 2009;201:1–119.

17. Braak H, de Vos RAI, Bohl J, Del Tredici K. Gastric alpha-synuclein immunoreactive inclusions

in Meissner's and Auerbach's plexuses in cases staged for Parkinson's disease-related brain

pathology. *Neurosci. Lett.* 2006;396(1):67–72.

18. Killinger BA et al. The vermiform appendix impacts the risk of developing Parkinson's disease.

Sci Transl Med 2018;10(465):eaar5280.

19. Lee JM et al. The Search for a Peripheral Biopsy Indicator of α-Synuclein Pathology for

Parkinson Disease. *J. Neuropathol. Exp. Neurol.* 2017;76(1):2–15.

20. Corbillé A-G et al. Evaluation of alpha-synuclein immunohistochemical methods for the detection

of Lewy-type synucleinopathy in gastrointestinal biopsies. *Acta Neuropathol Commun* 2016;4:35.

21. Donadio V et al. A new potential biomarker for dementia with Lewy bodies: Skin nerve α-

synuclein deposits. *Neurology* 2017;89(4):318–326.

22. Phillips RJ, Walter GC, Wilder SL, Baronowsky EA, Powley TL. Alpha-synuclein-

immunopositive myenteric neurons and vagal preganglionic terminals: autonomic pathway implicated

in Parkinson's disease?. *Neuroscience* 2008;153(3):733–750.

23. Holmqvist S et al. Direct evidence of Parkinson pathology spread from the gastrointestinal tract to

the brain in rats. *Acta Neuropathol* 2014;128(6):805–820.

24. Breid S et al. Neuroinvasion of α-Synuclein Prionoids after Intraperitoneal and Intraglossal

Inoculation. *J. Virol.* 2016;90(20):9182–9193.

25. Sargent D et al. "Prion-like" propagation of the synucleinopathy of M83 transgenic mice depends

on the mouse genotype and type of inoculum. *J. Neurochem.* 2017;143(1):126–135.

26. Manfredsson FP et al. Induction of alpha-synuclein pathology in the enteric nervous system of the

rat and non-human primate results in gastrointestinal dysmotility and transient CNS pathology.

Neurobiol. Dis. 2018;112:106–118.

27. Svensson E et al. Vagotomy and subsequent risk of Parkinson's disease. *Ann Neurol.*

2015;78(4):522–529.

28. Stolzenberg E et al. A Role for Neuronal Alpha-Synuclein in Gastrointestinal Immunity. *JIN*

2017;9(5):456–463.

29. Devos D et al. Colonic inflammation in Parkinson's disease. *Neurobiol. Dis.* 2013;50:42–48.

30. Mogi M et al. Interleukin-1 beta, interleukin-6, epidermal growth factor and transforming growth

factor-alpha are elevated in the brain from parkinsonian patients. *Neurosci. Lett.* 1994;180(2):147–

150.

31. Brás J, Guerreiro R, Hardy J. SnapShot: Genetics of Parkinson's disease. *Cell* 2015;160(3):570-

570.e1.

32. Witoelar A et al. Genome-wide Pleiotropy Between Parkinson Disease and Autoimmune

Diseases. *JAMA Neurol* 2017;74(7):780–792.

33. Umeno J et al. Meta-analysis of published studies identified eight additional common

susceptibility loci for Crohn's disease and ulcerative colitis. *Inflamm. Bowel Dis.* 2011;17(12):2407–

2415.

34. Gardet A et al. LRRK2 is involved in the IFN-gamma response and host response to pathogens. *J.*

Immunol. 2010;185(9):5577–5585.

35. Hakimi M et al. Parkinson's disease-linked LRRK2 is expressed in circulating and tissue immune

cells and upregulated following recognition of microbial structures. *J Neural Transm (Vienna)*

2011;118(5):795–808.

36. Lin J-C, Lin C-S, Hsu C-W, Lin C-L, Kao C-H. Association Between Parkinson's Disease and

Inflammatory Bowel Disease: a Nationwide Taiwanese Retrospective Cohort Study. *Inflamm. Bowel*

Dis. 2016;22(5):1049–1055.

37. Peter I et al. Anti-Tumor Necrosis Factor Therapy and Incidence of Parkinson Disease Among

Patients With Inflammatory Bowel Disease. *JAMA Neurol* 2018;75(8):939–946.

38. Wan Q-Y, Zhao R, Wu X-T. Older patients with IBD might have higher risk of Parkinson's

disease. *Gut* 2018;gutjnl-2018-317103.

39. Kahle PJ et al. Subcellular localization of wild-type and Parkinson's disease-associated mutant

alpha -synuclein in human and transgenic mouse brain. *J. Neurosci.* 2000;20(17):6365–6373.

40. Chassaing B, Aitken JD, Malleshappa M, Vijay-Kumar M. Dextran sulfate sodium (DSS)-induced

colitis in mice. *Curr Protoc Immunol* 2014;104:Unit 15.25.

41. Weber B, Saurer L, Schenk M, Dickgreber N, Mueller C. CX3CR1 defines functionally distinct

intestinal mononuclear phagocyte subsets which maintain their respective functions during

homeostatic and inflammatory conditions. *Eur. J. Immunol.* 2011;41(3):773–779.

42. Medina-Contreras O et al. CX3CR1 regulates intestinal macrophage homeostasis, bacterial

translocation, and colitogenic Th17 responses in mice. *J. Clin. Invest.* 2011;121(12):4787–4795.

43. Kostadinova FI et al. Crucial involvement of the CX3CR1-CX3CL1 axis in dextran sulfate

sodium-mediated acute colitis in mice. *J. Leukoc. Biol.* 2010;88(1):133–143.

44. Kang S et al. Intestinal epithelial cell-derived semaphorin 7A negatively regulates development of

colitis via αvβ1 integrin. *J. Immunol.* 2012;188(3):1108–1116.

45. Li B, Alli R, Vogel P, Geiger TL. IL-10 modulates DSS-induced colitis through a macrophage-

ROS-NO axis. *Mucosal Immunol* 2014;7(4):869–878.

46. Braak H et al. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol.*

Aging 2003;24(2):197–211.

47. Neumann M et al. Misfolded proteinase K-resistant hyperphosphorylated alpha-synuclein in aged

transgenic mice with locomotor deterioration and in human alpha-synucleinopathies. *J. Clin. Invest.*

2002;110(10):1429–1439.

48. Johnson ME, Stecher B, Labrie V, Brundin L, Brundin P. Triggers, Facilitators, and Aggravators:

Redefining Parkinson's Disease Pathogenesis. *Trends Neurosci.* 2018;S0166-2236(18):30253–30254.

49. Shannon KM, Keshavarzian A, Dodiya HB, Jakate S, Kordower JH. Is alpha-synuclein in the

colon a biomarker for premotor Parkinson's disease? Evidence from 3 cases. *Mov. Disord.*

2012;27(6):716–719.

50. Lebouvier T et al. Colonic biopsies to assess the neuropathology of Parkinson's disease and its

relationship with symptoms. *PLoS ONE* 2010;5(9):e12728.

51. Kuo Y-M et al. Extensive enteric nervous system abnormalities in mice transgenic for artificial

chromosomes containing Parkinson disease-associated α-synuclein gene mutations precede central

nervous system changes. *Hum Mol Genet* 2010;19(9):1633–1650.

52. Luk KC et al. Pathological α-synuclein transmission initiates Parkinson-like neurodegeneration in

nontransgenic mice. *Science* 2012;338(6109):949–953.

53. Rey NL et al. Widespread transneuronal propagation of α-synucleinopathy triggered in olfactory

bulb mimics prodromal Parkinson's disease. *J Exp Med* 2016;213(9):1759–1778.

54. Al-Sadi RM, Ma TY. IL-1beta causes an increase in intestinal epithelial tight junction

permeability. *J. Immunol.* 2007;178(7):4641–4649.

55. Capaldo CT, Nusrat A. Cytokine regulation of tight junctions. *Biochim Biophys Acta*

2009;1788(4):864–871.

56. Forsyth CB et al. Increased Intestinal Permeability Correlates with Sigmoid Mucosa alpha-

Synuclein Staining and Endotoxin Exposure Markers in Early Parkinson's Disease. *PLOS ONE*

2011;6(12):e28032.

57. Gray MT, Munoz DG, Gray DA, Schlossmacher MG, Woulfe JM. Alpha-synuclein in the

appendiceal mucosa of neurologically intact subjects. *Mov. Disord.* 2014;29(8):991–998.

58. Labrie V, Brundin P. Alpha-Synuclein to the Rescue: Immune Cell Recruitment by Alpha-

Synuclein during Gastrointestinal Infection. *JIN* 2017;9(5):437–440.

59. Kelly LP et al. Progression of Intestinal Permeability Changes and Alpha-Synuclein Expression in

a Mouse Model of Parkinson's Disease. *Mov Disord* 2014;29(8):999–1009.

60. Haenseler W et al. Excess α-synuclein compromises phagocytosis in iPSC-derived macrophages.

Sci Rep 2017;7(1):9003.

61. Takagawa T et al. An increase in LRRK2 suppresses autophagy and enhances Dectin-1–induced

immunity in a mouse model of colitis. *Science Translational Medicine* 2018;10(444):eaan8162.

62. Erickson MA, Hansen K, Banks WA. Inflammation-induced dysfunction of the low-density

lipoprotein receptor-related protein-1 at the blood-brain barrier: protection by the antioxidant N-

acetylcysteine. *Brain Behav. Immun.* 2012;26(7):1085–1094.

63. Cardoso FL et al. Exposure to Lipopolysaccharide and/or Unconjugated Bilirubin Impair the

Integrity and Function of Brain Microvascular Endothelial Cells. *PLOS ONE* 2012;7(5):e35919.

64. Erny D et al. Host microbiota constantly control maturation and function of microglia in the CNS.

Nat. Neurosci. 2015;18(7):965–977.

65. Sampson TR et al. Gut Microbiota Regulate Motor Deficits and Neuroinflammation in a Model of

Parkinson's Disease. *Cell* 2016;167(6):1469-1480.e12.

66. Chen SG et al. Exposure to the Functional Bacterial Amyloid Protein Curli Enhances Alpha-

Synuclein Aggregation in Aged Fischer 344 Rats and *Caenorhabditis elegans*. *Scientific Reports*

2016;6:34477.

67. Wendeln A-C et al. Innate immune memory in the brain shapes neurological disease hallmarks.

Nature 2018;556(7701):332–338.

68. Steiner JA, Quansah E, Brundin P. The concept of alpha-synuclein as a prion-like protein: ten

years after. *Cell Tissue Res.* 2018;373(1):161–173.

69. Recasens A et al. Lewy body extracts from Parkinson disease brains trigger α-synuclein pathology

and neurodegeneration in mice and monkeys. *Ann. Neurol.* 2014;75(3):351–362.

70. Hansen C et al. α-Synuclein propagates from mouse brain to grafted dopaminergic neurons and

seeds aggregation in cultured human cells. *J Clin Invest* 2011;121(2):715–725.

71. Angot E et al. Alpha-Synuclein Cell-to-Cell Transfer and Seeding in Grafted Dopaminergic

Neurons In Vivo. *PLOS ONE* 2012;7(6):e39465.

72. Desplats P et al. Inclusion formation and neuronal cell death through neuron-to-neuron

transmission of α-synuclein. *Proc Natl Acad Sci U S A* 2009;106(31):13010–13015.

73. Sacino AN et al. Intramuscular injection of α-synuclein induces CNS α-synuclein pathology and a

rapid-onset motor phenotype in transgenic mice. *Proc Natl Acad Sci U S A* 2014;111(29):10732–

10737.

74. Uemura N et al. Inoculation of α-synuclein preformed fibrils into the mouse gastrointestinal tract

induces Lewy body-like aggregates in the brainstem via the vagus nerve. *Molecular*

Neurodegeneration 2018;13:21.

75. Pan-Montojo F et al. Environmental toxins trigger PD-like progression via increased alpha-

synuclein release from enteric neurons in mice. *Sci Rep* 2012;2:898.

76. Ulusoy A et al. Neuron-to-neuron α-synuclein propagation in vivo is independent of neuronal

injury. *Acta Neuropathol Commun* 2015;3(13):1–13.

77. Villumsen M, Aznar S, Pakkenberg B, Jess T, Brudek T. Inflammatory bowel disease increases

the risk of Parkinson's disease: a Danish nationwide cohort study 1977-2014. *Gut* 2018;68(1):18–24.

78. Fujioka S et al. Occurrence of Crohn's disease with Parkinson's disease. *Parkinsonism & Related*

Disorders 2017;37:116–117.

79. Jung S et al. Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green

fluorescent protein reporter gene insertion. *Mol. Cell. Biol.* 2000;20(11):4106–4114.

80. Kitazawa M, Oddo S, Yamasaki TR, Green KN, LaFerla FM. Lipopolysaccharide-induced

inflammation exacerbates tau pathology by a cyclin-dependent kinase 5-mediated pathway in a

transgenic model of Alzheimer's disease. *J. Neurosci.* 2005;25(39):8843–8853.

81. Schenk M, Bouchon A, Seibold F, Mueller C. TREM-1--expressing intestinal macrophages

crucially amplify chronic inflammation in experimental colitis and inflammatory bowel diseases. *J.*

Clin. Invest. 2007;117(10):3097–3106.

82. Grathwohl SA et al. Formation and maintenance of Alzheimer's disease β-amyloid plaques in the

absence of microglia. *Nat Neurosci* 2009;12(11):1361–1363.

83. Rey NL et al. Spread of aggregates after olfactory bulb injection of α-synuclein fibrils is

associated with early neuronal loss and is reduced long term. *Acta Neuropathol* 2017;1–19.

84. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat.*

Methods 2012;9(7):671–675.

Figures and figure legends

Grathwohl et al., Figure 1

Figure 1

Alpha-Synuclein inclusions in the enteric nervous system and in macrophages of patients with

inflammatory bowel disease. (A, B) Immunofluorescence images of αSyn inclusions (Syn1) in the

submucosal region of 10 μm cryo-sections from colons of patients with colitis ulcerosa. Arrow heads

point to neuritic features indicating presence of inclusions in enteric nerves. Scale bar 50 μm. **(C)**

Close-up of a colonic region with active leukocyte infiltration in a patient with ulcerative colitis.

Immunoreactivity for αSyn (red) was observed in neuritic features (arrow heads) as well as in

individual leukocytes (arrows). Nuclei are shown in blue (DAPI). Scale bar 20 μm.

Figure 2

Age dependent increase of intracellular αSyn accumulation in enteric nervous system of

heterozygous (Thy1)-h[A30P]αSyn transgenic mice and setup of the experimental colitis

paradigms. (A) Confocal microscopy imaging of the granular inclusions of human αSyn (red,

antibody clone 211) within the ganglia of the submucosal plexus (green, peripherin; blue,

DAPI/nuclei) of heterozygous (Thy1)-h[A30P]αSyn transgenic mice. Scale bar, 100 μm. **(B)**

Stereological quantification of normally occurring human αSyn inclusions in the myenteric and

submucosal plexuses of 3 and 12 months old heterozygous (Thy1)-h[A30P]αSyn transgenic mice (n =

4 per group; mean and S.E.M. are shown; Student t-test between the two age groups in each region).

(C) Setup of experimental colitis paradigms employing dextran sodium sulfate (DSS, per os in

drinking water). Additionally, peripheral inflammation was induced by bacterial lipopolysaccharide

(LPS, intraperitoneal injection). For some DSS paradigms mice were sacrificed after an additional 2, 9

or 18-month of aging time on normal water **(D)** Hematoxylin staining of 35 μm thick colon sections

of 3 months old heterozygous (Thy1)-h[A30P]αSyn transgenic mice. Organizational layers of the

intact colon (left panel). Representative images of various severity degrees of DSS-driven colitis from

weak leukocyte infiltration (top panel of acute DSS) to mucosal ulceration (lowest panel of acute

DSS). Note the different appearance of enteric inflammation in acute LPS-driven peripheral

inflammation compared with DSS; e.g., confined immune cell clustering and lymphoid hyperplasia;

intact mucosal layer. Scale bar 50 μm (intact colon), 100 μm (acute DSS), and 200 μm (LPS).

A

aSyn Tg mice

B

C

D

Figure 3

Colitis severity and duration-dependent aggravation of accumulation of αSyn inclusions in the

colonic submucosal plexus of heterozygous (Thy1)-h[A30P]αSyn transgenic and wild type mice.

(A) DSS dose-dependent increase of leukocyte infiltration in the acute and chronic paradigm. The

highest acute dose (2.5%) and the two constant chronic doses led to an increase of αSyn inclusions in

the submucosal plexus (stereological quantification of αSyn inclusions in the submucosal plexus of 3

and 6 months old heterozygous (Thy1)-h[A30P]αSyn transgenic mice (n = 5-7 per group; mean and

s.e.m. are shown). **(B)** Representative 2D stacks of confocal images of increasing abundance of αSyn

inclusions (red, clone 211 antibody) in a ganglion of the submucosal plexus (green, peripherin) with

cellular nuclei in blue (DAPI). Scale bar 200 μm. **(C)** Overview of colonic region of 3-month-old

wildtype mice (top row) exposed to water or acute DSS (5%) with immunofluorescence analysis of

murine aSyn load in the colon performed immediately after colitis or exposed to constant chronic DSS

(2.5%) and analysis after aging on normal water for another 2 months. White dotted rectangles

indicate area that was zoomed out below illustrating in more detail the granular murine αSyn

inclusions (red, syn1 antibody) after DSS colitis in the submucosal plexus (green, peripherin

antibody). The lower panel shows DAPI and αSyn inclusions without the peripherin channel. Scale

bar for the lower two panels 200 μm. **(D)** Stereological quantification of granular murine αSyn

inclusions in the submucosal plexus of wildtype mice right after acute DSS colitis or after 2 months of

recovery from a 4-week chronic DSS colitis (n = 5 per group). Note the immunoreactivity for the

physiological, non-granular αSyn in the enteric nerves of the water group. The significant increase in

the DSS groups is evident without a statistical test.

Figure 4

Colitis induced by peroral DSS but not peritoneal LPS enhances αSyn accumulation in the

colonic submucosal plexus of heterozygous (Thy1)-h[A30P]αSyn transgenic mice and can be

aggravated by lack of Cx3cr1 signaling. (A) Administration of DSS (acute 5%) induces leukocyte

infiltration in wildtype, (Thy1)-h[A30P]αSyn transgenic (αSyn Tg) and Cx3cr1- deficient (Thy1)-

h[A30P]αSyn transgenic mice (αSyn Tg Cx3cr1-def). Relevant increase compared to water group is

evident without a statistical test. **(B)** Representative 2D stacks of confocal images of intracellular

granular αSyn inclusions (red, clone 211 antibody) in a ganglion of the myenteric plexus (green,

peripherin) with cellular nuclei in blue (DAPI). Scale bar 50 μm. Due to the clear distinction of the

large ganglia in the myenteric plexus granular features of the αSyn inclusions could be stereologically

quantified as particle load per ganglion **(C, D)**, whereas the αSyn inclusions in the elongated and more

diffuse appearing ganglia of the submucosal plexus were stereologically quantified as % area **(E, F)**.

Note the slight but significant mean increase of αSyn inclusions in Cx3cr1- deficient (Thy1)-

h[A30P]αSyn transgenic mice compared to (Thy1)-h[A30P]αSyn transgenic mice (Two-way ANOVA

with Tukey post hoc test). Gene expression analysis of selected cytokines in the colon of (Thy1)-

h[A30P]αSyn transgenic mice that received either acutely LPS **(G)** or DSS **(H)** compared to their

respective vehicle or water controls. Note the strong increase in IL-6 and the lack of elevation of IL-

10 in the DSS paradigm compared to the LPS paradigm indicating a different inflammatory colonic

milieu despite the abundant leukocyte infiltration in both paradigms. n = 5-8 per group; mean and

s.e.m.; Student's t-test between inflammatory agent and vehicle for individual cytokines.

Figure 5

Alpha-synuclein co-localizes with ENS and macrophages upon DSS colitis in αSyn transgenic

mice. (A, B) Immunofluorescence image of αSyn staining in colonic region of (Thy1)-h[A30P]αSyn

transgenic mice on water **(A)** or after acute DSS colitis (2.5%) **(B)**. Note the small dotted structures of

the typical αSyn inclusions in the submucosal plexus (arrow heads) and the large features of

immunoreactivity which localize to apparent infiltrating leukocytes (arrows), similar to what was

observed in IBD patients in Figure 1. Scale bar 100 μm. **(C)** 2D stacks and close-up of confocal

images co-localizing αSyn (red) with the macrophage marker Iba-1 (green) in the colon of a (Thy1)-

h[A30P]αSyn transgenic mouse after DSS colitis. Note the dotted structures of the typical αSyn

inclusions in the submucosal plexus (arrow heads). Scale bar 40 μm and 13 μm for the close-up. **(D)**

Quantification of numbers of Iba-1/αSyn-double positive macrophages (n = 3 per group; mean and

S.E.M.; difference in submucosal plexus of DSS group vs water is obvious without a statistical test).

Figure 6

Systemic IL-10 ameliorates DSS colitis and associated local αSyn accumulation in (Thy1)-

h[A30P]αSyn transgenic mice. Two different recombinantly engineered and murine IgG1-fused

forms of murine IL-10 (mIgG1(v1)-mIL10 and mIgG1(v2)-mIL10) were administered i .p. in an acute

DSS paradigm (5%) with (Thy1)-h[A30P]αSyn transgenic mice. Vehicle and the mIgG1(v1) alone

served as untreated controls. **(A)** Leukocyte infiltration was assessed by visual scoring and **(B)**

inclusion features of αSyn were stereologically and semi-automatically quantified and result log

scaled for statistical analysis. Both the vehicle group and the mIgG1(v1) group had similar levels of

905 leukocyte infiltration and αSyn inclusions and were merged for the statistical analysis to compare with

906 the IL-10 treated groups. Both forms of IL-10 ameliorated leukocyte infiltration whereas mIgG1(v2)-

907 mIL10 also blocked the appearance of αSyn inclusions significantly (n = 3-6 per group; mean and

908 s.e.m.; one-way ANOVA and Tukey post hoc test). **(C)** Persistent exposure mIgG1(v2)-mIL10 versus

909 mIgG1(v1)-mIL10 (lower limit of detection is indicated) as measured in serum at the end of the in

910 vivo phase corresponds with beneficial treatment effects on αSyn readout observed above. The

911 mIgG1(v1) was only measured in two mice.

Figure 7

A single chronic DSS colitis insult causes an age-dependent accumulation of proteinase K

resistant pSer129-αSyn in various brain regions of (Thy1)-h[A30P]αSyn transgenic mice. A 3-

week chronic increasing dose DSS paradigm was performed with 3-month old (Thy1)-h[A30P]αSyn

transgenic mice. After recovering and further aging, various brain regions were analyzed for

proteinase K resistant pSer129-αSyn immunoreactivity in 9-month **(A-F)** and 21-month old **(G-L)**

mice, respectively. Densitometric quantification of pSer129-αSyn immunoreactivity in different brain

regions in 9-month **(M-O)** and 21-month old mice **(R-T)** (n=6 mice per group). Statistical analyses

were performed using negative-binomial mixed-effects models adjusting for multiple comparisons.

Representative heatmap of the average distribution scores of pSer129-αSyn immunoreactivity for

each treatment group in varying brain regions in all the 9-month **(P-Q)** and 21-month old **(U-V)** mice

was generated in a sagittal mouse brain (n=10 mice per group). Scale bars: 500 μm.

Figure 8

L**oss of tyrosine hydroxylase and Nissl positive cells in the substantia nigra of (Thy1)-**

h[A30P]αSyn transgenic mice at 18 months post recovery from DSS colitis. (Thy1)-h[A30P]αSyn

transgenic mice that were exposed to a chronic DSS-colitis paradigm at 3 months and were aged to 21

months showed a significant loss of mean count of cells with tyrosine hydroxylase (TH)

immunoreactivity and cellular Nissl staining in the substantia nigra compared to age-matched

littermate mice in the group that did not experience DSS colitis (water). **(A)** Representative images of

two levels of the substantia nigra in one mouse per group. **(B)** Stereological quantification of cells

positive for TH or Nissl (n=7-8 mice per group). Statistical analyses of the TH dataset was performed

using Student's T-test, while Welch's T-test was used for the Nissl dataset to adjust for unequal

variances. Scale bar: 500 μm.