Affective flexibility without perceptual awareness

Philipp Homan[1], H. Lee Lau[1], Ifat Levy[2], Candace M. Raio[3], Dominik R. Bach[4,5],
David Carmel[6,7*], Daniela Schiller[1*]

[1]Department of Psychiatry, Department of Neuroscience, and Friedman Brain Institute,
Icahn School of Medicine at Mount Sinai
1470 Madison Ave, NY, NY 10029, USA

[2]Departments of Comparative Medicine, Neuroscience and Psychology
Yale University, New Haven, CT, USA

[3]Department of Psychology, New York University
6 Washington Place, NY, NY 10003, USA

[4]Clinical Psychiatry Research
Department of Psychiatry, Psychotherapy, and Psychosomatics
University of Zurich, Zurich, Switzerland

[5]Wellcome Centre for Human Neuroimaging,
12 Queen Square, London WC1N 3BG, UK

[6]School of Psychology, Victoria University of Wellington,
Easterfield Building, Kelburn Parade, Wellington 6012, New Zealand

[7]Psychology Department, University of Edinburgh,
7 George Square, Edinburgh EH8 9JZ, UK

*These authors contributed equally; address correspondence to
daniela.schiller@mssm.edu or david.carmel@vuw.ac.nz

Abstract

In an ever-changing environment, survival depends on learning which stimuli represent threat,

and also on updating such associations when circumstances shift. Humans can acquire

physiological responses to threat-associated stimuli even when they are unaware of them, but the

role of awareness in *updating* threat contingencies remains unknown. This complex process –

generating novel responses while simultaneously suppressing learned ones – relies on distinct

neural mechanisms from initial learning, and has only been shown with awareness. Can it occur

unconsciously? Here we show that it can. Participants underwent classical threat conditioning to

visual stimuli that were suppressed from their awareness. One of two images was paired with an

electric shock; halfway through the experiment, contingencies were reversed and the shock was

paired with the other image. We found that physiological responses reflected changes in

stimulus-threat pairings independently of stimulus awareness, demonstrating the sophistication

of unconscious affective flexibility.

Introduction

Flexible responses to environmental threats are essential for adaptive behavior. Cues that predict threat constantly change - new threats may arise while old ones cease to pose a risk. When consciously perceiving such cues, we are able to flexibly update and shift threat responses from one cue to another (1-3). But can we update our reaction to stimuli that predict danger when we are not aware of them? It is known that threat-conditioned stimuli that are perceived without awareness can still elicit defensive physiological reactions (4-7), and that new threat associations can be formed through classical conditioning even without any awareness of the conditioned stimuli (8-10). Updating threat associations when contingencies change, however, is an entirely different matter: it involves a complex process of creating novel responses while simultaneously suppressing acquired ones. To date, such updating has only been shown in humans who were aware of the stimuli (2), and in animals under conditions where stimuli were fully available for perceptual processing (11); these studies have shown, furthermore, that the neural substrates of threat updating differ from those of the initial learning. It is thus unknown whether the sophisticated re-evaluation involved in such affective flexibility requires awareness, or can be accomplished without it. Here we show that it can, and furthermore, that stimulus awareness does not seem to play a substantial role in such affective flexibility.

To examine this, we employed the reversal paradigm, a laboratory model that requires flexible updating of threat contingencies (2). In an initial acquisition phase, participants encounter two conditioned stimuli (CSs) and learn that only one of them predicts an electric shock. Halfway through the experiment, with no warning, these contingencies flip, initiating the reversal phase: Participants must flexibly learn that the formerly safe CS now predicts the shock and that the old one no longer does. To assess learning, participants' physiological arousal is recorded throughout the experiment, typically (and here) by measuring their skin conductance

responses. Appropriate response reversal requires a sophisticated form of updating, in that one

must learn to respond to a cue that now predicts threat while simultaneously inhibiting responses

to the previously threatening cue that is now safe.

To see whether reversal of conditioned threat requires awareness, we had a large group of

participants ($N = 86$) undergo reversal learning with the CSs suppressed from awareness by

continuous flash suppression (CFS), a technique commonly used to examine unconscious

perception (10, 12-14): The CSs were visual images presented monocularly, while the other eye

was shown a high-contrast, dynamic image (the CFS mask) at the corresponding retinal location

(See Figure 1 for a description of the design and procedure).

CFS can suppress images from awareness for several seconds. However, it is also known that

its effectiveness may vary across trials and individuals, and the suppressed stimulus may "break

through" the suppression (15). Over the last decade, a growing body of work has raised concerns

that the standard approach - removing from analysis data (participants and trials) in which

breakthrough had occurred - may bias the findings (16, 17; See Supplementary Methods for

further details of these issues.) Here, we adopt a number of methodological approaches to ensure

our results are robust to these potential concerns.

Specifically, we remove no data and instead incorporate individual levels of reported

stimulus awareness, as well as response patterns that might reflect residual awareness, into a

regression model accounting for physiological responses. The model also adjusts for baseline

anxiety (which has been previously shown to correlate with unconscious learning; (10)).

Additionally, we use a Bayesian approach to establish that a model in which participants were

updating their learning provides a better account for the findings than a model in which they

were simply (and independently of the stimulus) predicting the probability of a shock on the next

trial (18). Finally, in order to verify that our procedure is able to induce reversal learning when

participants are aware of the stimuli (so that if no learning or updating were found under CFS,

we could rule out the possibility this may be simply due to an ineffective procedure) we ran a no-

CFS group ($N = 12$), in which participants also viewed the CSs monocularly (as the CFS group

did), but were aware of them as no CFS masks were presented to their other eye.

We hypothesized that physiological responses to threat can be flexibly reversed without

perceptual awareness. We find that reversal indeed occurs independently of CS awareness, and

that there is strong evidence for the reversal of threat learning even in its complete absence.

Results

Overall assessment of physiological reversal learning

To assess the physiological arousal evoked by CSs, we used a model-based approach (19) to

estimate the amplitude of anticipatory sudomotor nerve activity (SNA) from skin conductance

data recorded during stimulus presentation. A variational Bayes approximation was employed to

invert a forward model that describes how hidden SNA translates into observable SCRs (see

Materials and Methods). Previous work has shown that this approach is more sensitive than

conventional SCR peak-to-peak analysis (19-21). Figure 2A shows the time course of evoked

SNA to Spiders A and B, separately for the CFS and no-CFS groups. In both groups, responses

to Spider A relative to Spider B were larger during the acquisition phase and smaller during the

reversal phase. To quantify the magnitude of physiological reversal learning, we calculated a

reversal learning index for each participant (see Materials and Methods). The reversal learning

index was positive and significantly greater than zero for both the CFS and no-CFS groups

(Figure 2B). A linear mixed model (see Materials and Methods for details) revealed a significant

interaction of stage and spider in both groups (CFS: $\beta = 0.27$, $t(2935) = 4.23$, $P = < 0.001$; no-

CFS: $\beta = 1.23$, $t(2935) = 7.29$, $P = < 0.001$). Note that a significant interaction is formally

equivalent to a significant reversal learning index. On its own, however, it simply reveals a

difference in the comparative magnitude of responses to the two CSs across the two halves of the

experiment; follow-up tests show that this difference is indeed due to reversal: Spider A evoked

greater responses than Spider B in the acquisition phase (CFS: t (341.9) = 3.0, P = 0.003; no-

CFS: t (201.1) = 4.6, P < 0.001) and the pattern was reversed in the reversal phase (CFS: t

(341.9) = 2.8, P = 0.005; no-CFS: t (341.9) = 3.6, P = 0.0003). These results indicate that

reversal learning was evident in both groups. Although Figure 2 shows that it was more

pronounced in the no-CFS group, we note that this difference is not straightforwardly

interpretable because the no-CFS group (a control, intended to rule out an ineffective

manipulation if no effect was found for the CFS group) was substantially smaller; furthermore,

as addressed in detail below, suppression from awareness was very heterogenous in the CFS

group.

As previous work has found a negative association between anxiety and threat acquisition

with and without awareness (10), we also calculated correlations between the CFS group's

baseline anxiety measures (STAIT, STAIS, FSQ) and the reversal learning index. Overall,

reversal learning decreased significantly with increasing levels of state and trait anxiety, and to a

lesser but non-significant extent for spider phobia (Figure 2C).

Reversal learning and perceptual awareness

The CFS manipulation reduced awareness of the CSs; as expected, however, it was differentially

effective in doing so across participants, precluding an overall conclusion that all learning under

CFS happened non-consciously. The CFS group showed significantly lower accuracy in response

to the "which seen?" question (M = 0.46, SD = 0.29) compared to the no-CFS group (M = 0.86,

SD = 0.16; t (22.77) = -7.24, P < 0.001), and accuracy in the CFS group was not significantly

different from the 50% random-response level (t (85) = -1.21, P = 0.229). The CFS group also

showed lower confidence (M = 1.73, SD = 0.65) than the no-CFS group (M = 2.83, SD = 0.08; t

(95.38) = -15.05, P < 0.001).

However, group differences in accuracy and confidence, and even random-level response

accuracy, are not sufficient to establish an absence of perceptual awareness in the CFS group.

Notably, average confidence of correct responses in this group was low but significantly greater

than the minimum value of 1 (t (77) = 10.79, P < 0.001), suggesting that at least some

participants were aware of some of the CSs; learning might thus have arisen from a subset of

trials and/or participants where such awareness occurred. To address this, we quantified CS

awareness by calculating an awareness index for each participant, ranging in possible values

from 0 for no awareness to 1 for full awareness (see Materials and Methods). Although the

awareness index of the CFS group (M = 0.28, SD = 0.34) was significantly lower than the no-

CFS group's (M = 0.92, SD = 0.18; t (23.93) = -10.19, P < 0.001), it was still significantly

higher than zero (t (85) = 7.59, P < 0.001).

Therefore, in order to test our main hypothesis that the reversal of acquired threat responses

can be achieved without perceptual awareness, we characterized the quantitative relation

between the level of awareness and the magnitude of reversal learning in the CFS group. To

control for possible artifacts of regression to the mean (see Supplementary Methods), we

followed the recommendation (16) to first examine the correlation between two independent

estimates of the awareness index, one calculated from even-numbered trials, the other from odd-

numbered trials. Because noise at the measurement level might occasionally yield extreme (i.e.,

very low or very high) awareness index scores, an association of such randomly-extreme scores

with reversal learning (specifically, low awareness with intact learning) could be an artifact.

However, it is highly unlikely that across participants, random noise would yield consistent (and

similarly extreme) measurements in separate estimates. Due to regression to the mean, if random

extreme values occur in one of the two estimates, they are less likely to occur in the other,

resulting in a considerable attenuation of any correlation between the two. We found, however,

that the two measures were strongly correlated (r (84) = 0.96, $P < 0.001$; Figure 3A);

participants' awareness level in one set of trials was overwhelmingly predictive of their

awareness in the other set, confirming their reliability as estimates of awareness.

Next, we examined the association between the awareness index and the reversal learning

index, using values of both indices obtained separately from even (Figure 3B) and odd (Figure

3C) trials. As the color-coding of Figure 3 shows, the relation between individual participants'

awareness and their reversal learning was highly consistent across these separate measurements.

In light of this, we pooled the data from all trials and regressed the reversal learning index on the

perceptual awareness index (Figure 3D). The parameter of interest was the intercept, i.e., the

magnitude of reversal learning at zero perceptual awareness. The intercept was positive and

significantly different from zero. Furthermore, the awareness index regressor did not contribute

significantly to prediction of reversal learning; importantly, this finding was even stronger in

models that accounted for STAIT scores and a binary factor indicating whether participants were

tracking the stimuli with their responses (see Materials and Methods; Figure 3E and Table 1).

Comparing learning and expectation-based accounts

Well-controlled lab-based conditioning procedures require strict constraints that preclude

complete randomization of the number and order of different CSs; this comes with a cost:

participants are able to develop expectations with above-chance validity, based on the sequence

of trials so far, about the likelihood of a shock on any upcoming trial (18). Even without any

awareness of the CSs, a participant should have been able to distinguish two types of trials:

reinforced (with shock) and non-reinforced (no-shock). In a study with two CSs and a 100%

reinforcement rate like ours, such expectations would correspond to an anticipated pattern of

alternating trial-types (shock/no-shock or vice versa), with an increase in shock anticipation after

every no-shock trial. The question, therefore, was whether the physiological responses we had

measured might simply reflect participants' pattern-based anticipation of shock, rather than

learning of the contingencies associated with the CSs.

To answer this question, we used a Bayesian approach to compare the probability of our

findings being accounted for by a classic Rescorla-Wagner learning model (22) and a trial-

sequence model. We hypothesized that successful threat reversal without perceptual awareness

should be better explained by the Rescorla-Wagner learning model, whereas simple pattern-

based expectation would be better explained by the trial-sequence learning model. We used

maximum likelihood estimation to assess the log likelihood and calculate the Bayesian

Information Criterion (BIC) of each model (See Materials and Methods for details of each model

and calculation of the BIC). A smaller BIC indicates a better model, and BIC values can thus be

compared by calculating the difference between them and interpreting the resulting Δ BIC as

providing evidence against the higher BIC. The Rescorla-Wagner model (BIC: 562.1)

outperformed the pattern-based expectation model (BIC: 584.9), with the difference (Δ BIC:

22.9) greater than 10, suggesting that the evidence against the trial switch model is very strong

(23). Repeating this comparison for just the participants with zero mean awareness confirmed the

lower BIC for the Rescorla-Wagner model (BIC: 114.3) compared to the pattern-based

expectation model (BIC: 125.7), with the difference again greater than 10 (Δ BIC: 11.3; see also

Figure S2). This model comparison provides convincing evidence that a classical Rescorla-

Wagner learning model explains our findings better than an alternative expectation-based model.

Discussion

These results indicate that participants updated their defensive physiological responses independently of their awareness of threat-related cues. Previous studies have shown that new threat associations can be formed without perceptual awareness of the conditioned stimuli (5, 9-10). However, until now it was unknown whether the far more complex process of threat reversal - shifting reactions from a stimulus that no longer predicts danger to one that now does - can be accomplished without awareness. The independence of reversal learning from perceptual awareness, reported here, suggests that separate processes underlie affective flexibility and conscious processing (24). Conversely, the negative correlation between reversal learning and anxiety suggests that the various impairments caused by anxiety are not limited to the systems underlying conscious processes.

Previous studies have pointed out the limitations of using accuracy and confidence measures to assess perceptual awareness, and suggested remedies including the calculation of metacognitive sensitivity measures (25), Bayesian statistics (26), or parametric variation of the experimental manipulation (27). The present study addresses an issue not covered in previous discussions, by showing that a trial-wise analysis may reveal hints for incomplete suppression that analyses relying on average measures might easily miss. Future studies that rely on forced-choice questions for awareness assessment should thus examine response patterns across trials in addition to collecting aggregate measures.

Notably, a previous study (10) that used CFS to investigate acquisition of threat responses without awareness of the stimuli found that such acquisition can occur, but is rapidly forgotten. The present study again showed that such acquisition can occur (and, additionally, be reversed), but did not find the same rapid forgetting. The reasons for this are unclear, but we speculate that the difference may be due to specific aspects of the stimuli, design and procedure: our use of

pictures of spiders (rather than the faces used in the previous study) and a 100% (rather than

50%) reinforcement protocol may have altered the temporal characteristics of acquisition.

Similarly, the temporal profile of reversal may change if the stimuli and reinforcement regime

are different.

The present results add to a growing body of findings distinguishing functions that do and do

not require awareness. Such distinctions are important in guiding research into the neural

mechanisms of conscious and non-conscious processing. Previous research hints at the

mechanism underlying the non-conscious affective flexibility reported here, although it remains

to be elucidated: The ability to reverse conditioned responses depends on the integrity of

circuitry spanning several neural regions, particularly the ventromedial prefrontal cortex

(vmPFC) and its connections with the amygdala (1) where threat associations are formed (28).

Consistent with this, it is known that patients with anxiety disorders often show rigid and

inflexible threat responses in conjunction with prefrontal cortex dysfunction (29, 30).

Indeed, the real-life settings that people with anxiety disorders find challenging often require

the updating and shifting of threat responses. Deficits in affective flexibility may thus explain the

threat learning and extinction deficits seen in such disorders (31): Compared to healthy controls,

patients are less able to distinguish between safe and unsafe stimuli in threat learning (when it is

adaptive to do so), and distinguish between them to a greater extent during extinction (when it is

non-adaptive). Threat learning without perceptual awareness is also negatively correlated with

baseline state anxiety in healthy participants (10). Our new finding that baseline anxiety is

negatively correlated with affective flexibility suggests a potential use for reversal learning as a

model paradigm for investigating how anxiety modulates various processes in a variety of

disorders, including, for example, posttraumatic stress disorder, in which there is an impairment

of threat inhibition (32).

Methods

Participants

Ninety-eight healthy participants (mean age = 29.97; range 18-65) were assigned to one of the two groups: reversal learning with CFS (CFS group; $N = 86$, 48 female) or without CFS (no-CFS group; $N = 12$, 5 female). Assignment was random until each group reached a size of 12; subsequent participants were assigned to the CFS group. Measures of trait and state anxiety (Spielberger Trait-State Anxiety Inventory (33); STAIT and STAIS, respectively) and spider phobia (Fear of Spider Questionnaire; FSQ (34)) were taken prior to participation and did not differ between the groups (Table S1). The experiment was approved by the Institutional Review Board of the Icahn School of Medicine at Mount Sinai. All participants provided written informed consent and were financially compensated for their participation.

Experimental procedure

Participants viewed the stimuli monocularly, through a mirror stereoscope (StereoAids, Australia) placed at a distance of 45 cm from a 17-inch Dell monitor. The CSs (schematic low-contrast images of spiders), presented to the left eye only, were suppressed from awareness in the CFS group: while the left eye saw them, the right eye was presented with "Mondrians" - arrays of high contrast, multi-colored, randomly generated rectangles alternating at 10 Hz. Both the CSs and the CFS masks were flanked by identical textured black and white bars, to facilitate stable ocular vergence. The no-CFS group viewed identical CSs (also presented monocularly), but with no Mondrians presented to the other eye.

The experiment consisted of 16 acquisition trials followed by 16 reversal trials. One of two spider images was presented on each trial. The spider images were schematic and had similar low-level features. During acquisition, spider A always terminated with a shock and spider B

never did. Reversal occurred halfway through the experiment: spider B now terminated with a

shock and spider A did not. The spider stimuli were presented for 6 s each in pseudorandomized

order. One of four possible trial orders was used for each participant. Orders were generated by

imposing specific constraints on the trial order, such that the first trial was always reinforced and

no more than two of the same trial type ever occurred consecutively.

Trial order and spider identity were counterbalanced across participants. To assess the

effectiveness of the awareness manipulation (35), 1 s after the offset of every CS participants

were shown the question "Which seen?" (1 = flower, 2 = spider; notably, flowers were never

shown, meaning the question addressed detection rather than discrimination as it could be

answered correctly even with a brief glimpse). This was followed by the question "How

confident?" (1 = guess to 3 = sure; participants were instructed to indicate how confident they

were of the flower/spider answer they had just given). Both questions were presented binocularly

(1.5 - 2 s each, during which responses had to be given by pressing number keys on a standard

keyboard). The second question was followed by an 8 to 10 s inter-trial interval.

Psychophysiological stimulation and measurement

Mild electric shocks were delivered using a Grass Medical Instruments SD9 stimulator and

stimulating bar electrode attached to the participant's right wrist. Shocks (200 ms; 50 pulse/s)

were delivered at a level determined individually by each participant as "uncomfortable but not

painful" (maximum of 60V), during a work-up procedure prior to the experiment.

Skin conductance responses (SCR) were measured with Ag-AgCl electrodes, filled with

standard isotonic NaCl electrolyte gel, and attached to the middle phalanges of the second and

third fingers of the left hand. SCR signals were sampled continuously at a rate of 200 Hz,

amplified and recorded with a MP150 BIOPAC Systems skin conductance module connected to

a PC.

Analysis of physiological responses

Model-based analysis

We estimated SNA from SCR data with a model-based variational Bayes approximation (19),

inverting a forward model that describes how (hidden) SNA translates into (observable) SCR. A

unit increase in SNA corresponds to an increase in SCR of 1 micro Siemens. The model assumes

that the observed SCR can be decomposed into different components including anticipation,

evocation, and spontaneous fluctuations, each of which are generated by bursts of SNA driven by

changes in sympathetic arousal. The generative (forward) model thus describes how sympathetic

arousal, the physiological measure that is taken as an index of the psychological process of

threat, translates into sudomotor nerve bursts which then generate the observable SCR (19).

Using Bayesian inference, the forward model can then be reversed in order to estimate the most

likely underlying SNA given the observed SCR:

$$p(\theta|y) = \frac{p(\theta)p(y|\theta)}{p(y)},$$

(1)

where the most likely parameter vector Θ (corresponding to the SNA) given the observed

outcome y (corresponding to the SCR) is given by the prior estimate of Θ weighted by the

likelihood of y given Θ. Solving this equation involves integration over the model evidence p(y)

which is analytically hard to compute (and possibly intractable). This can be resolved by

replacing this integration problem by an optimization problem, which can be approximated with

Variational Bayes procedures (36), where the log of the model evidence can be framed as the

sum of the Kullback-Leibler divergence and the Free Energy. By maximizing the Free Energy

the Kullback-Leibler divergence is minimized, and a lower bound to the log model evidence can

be derived iteratively.

The SNA estimates were computed using previously developed software package PsPM (19)

implemented in MATLAB R2016b (The Mathworks Inc, Natick, MA, USA). The statistical

analyses were conducted with the R software (37) (R version 3.4.2 (2017-09-28)) and the

libraries lme4 (38) and lsmeans (39). Welch's t-tests were used instead of two sample t-tests

when groups had unequal variances.

Reversal Learning Index

An estimate of SNA was obtained for each trial. We expected Spider A to evoke greater SNA

than Spider B during the acquisition phase, and Spider B to evoke greater SNA than Spider A

during the reversal phase. The strength of reversal learning can thus be quantified by calculating,

separately for the acquisition and reversal phases, the difference between the average SNA

evoked by each spider. To quantify the degree of reversal (which is formally equivalent to the

interaction of phase and stimulus), the reversal learning index was calculated by subtracting the

difference between mean SNAs evoked by each spider during reversal from the difference during

acquisition (the larger the index, the greater the magnitude of reversal learning):

$$\text{Reversal learning index} = \Delta\text{Acquisition} - \Delta\text{Reversal}$$
$$\Delta\text{Acquisition} = [\text{mean(Spider A)} - \text{mean(Spider B)}]_{\text{Acquisition}}$$
$$\Delta\text{Reversal} = [\text{mean(Spider A)} - \text{mean(Spider B)}]_{\text{Reversal}}$$

310   (2)

To formally test for group differences in the strength of reversal learning, we computed a

linear mixed model using the lme4 library in R. We used the skin conductance response

(converted to a model-based measure of sudomotor nerve activity, SNA) as the dependent

variable and entered group (CFS, no-CFS), stage (acquisition, reversal), and spider (spider A,

spider B) as well as a continuous variable for trial (to account for habituation) as predictors. The

random structure of the model included an intercept and slopes for stage and spider.

Assessments of perceptual awareness

Perceptual awareness index

To characterize participants' reported awareness of CSs, each trial was assigned a perceptual

awareness score, defined by a combination of detection and confidence responses: Correct

answers with a confidence rating of 1 (guess) and incorrect answers irrespective of confidence

were assigned an awareness score of 0; correct answers with a confidence rating of 2 (medium)

were assigned a score of 0.5, and correct answers with a confidence rating of 3 (high) were

assigned an awareness score of 1. A perceptual awareness index was calculated for each

participant by averaging awareness scores across all trials.

Stimulus-response association patterns ("tracking")

We also assessed response patterns across trials, to see whether participants were able to track

stimuli with their responses, accurately discriminating the images despite not being able to label

them. We plotted individual trial-by-trial responses to the question "Which seen?", overlaid on

the trial-by-trial presentation of spiders (spider A, spider B; Figure S1A). We then calculated the

number of consecutive "hits", defined as the number of consecutive trials where these two time-

courses were either identical or consistently in opposition, suggesting that there was a possible

association between the stimulus and the response during those trials. The probability of such

consecutive hits occurring by chance alone can be derived as follows:

Let $p = 0.5$ be the probability of a hit, k the number of consecutive hits, n the number of trials

left, i the number of consecutive hits already observed; the chance of observing k consecutive

hits for the remaining n trials can then be formulated as a recursive problem:

$$f_{p,k}(i, n) = p f_{p,k}(i+1, n-1) + (1-p) f_{p,k}(0, n-1),$$

338   (3)

which can be solved analytically with dynamic programming or recursion. Trivially, $f_{p,k}(k, n) =$

1 for $n \geq 0$ since k consecutive hits have already been observed, and $f_{p,k}(i, n) = 0$ for $k - i > n$

since there are not enough trials left to observe k consecutive hits.

For example, assuming we want to know how likely it is to observe $k = 8$ consecutive hits

within $n = 32$ trials given $p = 0.5$, i.e., $f_{0.5, 8}(0, 32)$, we find that this yields a probability of

0.050.

Alternatively, the probability can be derived by simulation for all possible numbers of

consecutive hits within 32 trials (i.e., from 1 to 31). For each possible number, we thus also

simulated 10^5 draws of a binomial distribution and calculated the average probability of that

number of hits being consecutive. As can be seen in Figure S1B, the result for 8 consecutive hits

(0.04991) was very close to the analytical solution. Fifteen participants showed evidence of

tracking the spiders or the shocks with their responses (8 or more consecutive hits); notably, 3 of

these participants appeared to have a perceptual awareness index of zero. We thus adjusted our

subsequent analysis with an additional binary covariate, indicating whether participants did or

did not show 8 or more consecutive hits.

Comparing learning and expectation-based models

The Rescorla-Wagner model (22) describes how the prediction for each trial is updated according to a prediction error and learning rate:

$$V_{n+1}(x_n) = V_n(x_n) + \alpha\delta_n$$
$$\delta_n = r_n - V_n(x_n),$$

$$(4)$$

where x_n is the conditioned stimulus on trial n (Spider A or Spider B), and δ_n is the punishment prediction error that measures the difference between the expected and the actual shock (r_n) on trial n. The learning rate α for the value update is a constant free parameter. The value for the CS not observed on trial n remains unchanged. To derive the best fits for the Rescorla-Wagner model, we assumed that $V_0 = 0.5$, reflecting the assumption that getting a shock or not was equally likely for the first trial.

For the alternative trial-sequence learning model, we assumed that a participant expecting a strict sequence of alternating trial types (shock/no shock or vice versa) would update this expectation according to the actually encountered trial types and a constant learning rate:

$$V'_{n+1} = V'_n + \alpha'\delta'_n$$
$$\delta'_n = r'_n - V'_n$$
$$\tau_n = |(r'_{n-1} - 1)|,$$

$$(5)$$

where V'_{n+1} is the expected trial type switch at trial $n+1$ (if V'_{n+1} is larger than 0.5, a trial switch is expected), α' is the learning rate, and δ'_n is the prediction error. The prediction error corresponds to the difference between the actual trial type switch for trial n (r'_n; coded as one for a trial type switch and zero for an equal trial type) and the expectation for trial n. A changing trial type for trial n was tracked by τ_n, which was one if the preceding trial was zero and zero if

373   the preceding trial type was one. To map these expectations onto expected values, we assumed

374   that

$$V_{n+1} = \begin{cases} V'_{n+1} \cdot \tau_n + (1 - V'_{n+1})(1 - \tau_n), & \text{if } V' > 0.5 \\ V'_{n+1}, & \text{otherwise,} \end{cases}$$

375   (6)

where the expected value for trial $n+1$ was calculated according to whether a trial type switch

was expected ($V' > 0.5$) or not.

We performed a formal model comparison between the conventional Rescorla-Wagner

model and the trial switch model for our data set (Figure S2), using maximum likelihood

estimation and non-linear optimization (implemented with the fmincon function in MATLAB

R2016b (The Mathworks Inc, Natick, MA, USA). Using the log likelihood, we calculated the

Bayesian Information Criterion (BIC) to compare the two models as follows:

$$\text{BIC} = \log(n)\, k - 2 \cdot \log(\hat{L}),$$

383   (7)

384   where n is the number of data points, k is the number of regressors, and $\hat{L}$ is the maximized

385   value of the likelihood function.

References

1. Morris JS, Dolan RJ (2004) Dissociable amygdala and orbitofrontal responses during reversal fear conditioning. *Neuroimage* 22(1):372–80.

2. Schiller D, et al. (2008) From fear to safety and back: Reversal of fear in the human brain. *Journal of Neuroscience* 28(45):11517–25.

3. Fleming SM, Dolan RJ, Frith CD (2012) Metacognition: computation, biology and function. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 367(1594):1280–6.

4. Critchley HD, Mathias CJ, Dolan RJ (2002) Fear conditioning in humans: the influence of awareness and autonomic arousal on functional neuroanatomy. *Neuron* 33(4):653–63.

5. Morris JS, Ohman A, Dolan RJ (1998) Conscious and unconscious emotional learning in the human amygdala. *Nature* 393(6684):467–70.

6. Ohman A, Soares JJ (1994) "unconscious anxiety": phobic responses to masked stimuli. *Journal of Abnormal Psychology* 103(2):231–40.

7. Whalen PJ, et al. (1998) Masked presentations of emotional facial expressions modulate amygdala activity without explicit knowledge. *Journal of Neuroscience* 18(1):411–8.

8. Katkin ES, Wiens S, Ohman A (2001) Nonconscious fear conditioning, visceral perception, and the development of gut feelings. *Psychol Sci* 12(5):366–70.

9. Manns JR, Clark RE, Squire LR (2002) Standard delay eyeblink classical conditioning is independent of awareness. *Journal of Experimental Psychology: Animal Behavior Processes* 28(1):32–7.

10. Raio CM, Carmel D, Carrasco M, Phelps EA (2012) Nonconscious fear is quickly acquired but swiftly forgotten. *Current Biology* 22(12):R477–9.

410   11.   Izquiredo A, Brigman JL, Radke AK, Rudebeck PH, Holmes A (217) The neural basis of

411   reversal learning: An updated perspective. *Neuroscience* 345:12–26.

412   12.   Tsuchiya N, Koch C (2005) Continuous flash suppression reduces negative afterimages.

413   *Nature Neuroscience* 8(8):1096–101.

414   13.   Stein T, Hebart MN, Sterzer P (2011) Breaking continuous flash suppression: A new

415   measure of unconscious processing during interocular suppression? *Frontiers in Human*

416   *Neuroscience* 5:167.

417   14.   Carmel D, Arcaro M, Kastner S, Hasson U (2010) How to create and use binocular rivalry.

418   *J Vis Exp* (45).

419   15.   Gayet S, Stein T (2017) Between-subject variability in the breaking continuous flash

420   suppression paradigm: Potential causes, consequences, and solutions. *Frontiers in*

421   *Psychology* 8:437.

422   16.   Shanks DR (2016) Regressive research: The pitfalls of post hoc data selection in the study

423   of unconscious mental processes. *Psychonomic Bulletin & Review* 24:752–775.

424   17.   Stein T, Sterzer P (2014) Unconscious processing under interocular suppression: getting

425   the right measure. *Frontiers in Psychology*, *5*, 387.

426   18.   Wiens S, Katkin ES, Ohman A (2003) Effects of trial order and differential conditioning

427   on acquisition of differential shock expectancy and skin conductance conditioning to

428   masked stimuli. *Psychophysiology* 40(6):989–97.

429   19.   Bach DR, Daunizeau J, Friston KJ, Dolan RJ (2010) Dynamic causal modelling of

430   anticipatory skin conductance responses. *Biological Psychology* 85(1):163–70.

431   20.   Bach DR (2014) A head-to-head comparison of scralyze and ledalab, two model-based

432   methods for skin conductance analysis. *Biological Psychology* 103:63–8.

433   21.   Staib M, Castegnetti G, Bach DR (2015) Optimising a model-based approach to inferring

434   fear learning from skin conductance responses. *Journal of Neuroscience Methods* 255:131–

435   8.

436   22.   Rescorla R, Wagner A (1972) *A theory of Pavlovian conditioning: Variations in the*

437   *effectiveness of reinforcement and nonreinforcement*, eds. Black A, Prokasy W. (Appleton-

438   Century-Crofts, New York), pp. 64–99.

439   23.   Raftery AE (1995) Bayesian model selection in social research. *Sociological methodology*

440   pp. 111–163.

441   24.   Lau H, Rosenthal D (2011) Empirical support for higher-order theories of conscious

442   awareness. *Trends in Cognitive Sciences* 15(8):365 – 373.

443   25.   Fleming SM, Lau HC (2014) How to measure metacognition. *Frontiers in Human*

444   *Neuroscience* 8:443.

445   26.   Dienes Z (2015) *How Bayesian statistics are needed to determine whether mental states*

446   *are unconscious*, ed. Overgaard M. (Oxford University Press, Oxford), pp. 199–221.

447   27.   Schmidt T (2015) Invisible stimuli, implicit thresholds: Why invisibility judgments cannot

448   be interpreted in isolation. *Adv Cogn Psychol* 11(2):31–41.

449   28.   Roy M, Shohamy D, Wager TD (2012) Ventromedial prefrontal-subcortical systems and

450   the generation of affective meaning. *Trends in Cognitive Science* 16(3):147–56.

451   29.   Ressler KJ, Mayberg HS (2007) Targeting abnormal neural circuits in mood and anxiety

452   disorders: from the laboratory to the clinic. *Nature Neuroscience* 10(9):1116–24.

453   30.   Rauch SL, Shin LM, Phelps EA (2006) Neurocircuitry models of posttraumatic stress

454   disorder and extinction: human neuroimaging research–past, present, and future. *Biological*

455   *Psychiatry* 60(4):376–82.

31.   Duits P, et al. (2015) Updated meta-analysis of classical fear conditioning in the anxiety

disorders. *Depression and Anxiety* 32(4):239–253.

32.   Jovanovic T, Norrholm SD (2011) Neural mechanisms of impaired fear inhibition in

posttraumatic stress disorder. *Frontiers in Behavioral Neuroscience* 5:44.

33.   Spielberger CD (1983) State-trait anxiety inventory for adults. *PsycTESTS Dataset*.

34.   Szymanski J, O'Donohue W (1995) Fear of spiders questionnaire. *PsycTESTS Dataset*.

35.   Merikle PM, Smilek D, Eastwood JD (2001) Perception without awareness: perspectives

from cognitive psychology. *Cognition* 79(1-2):115–34.

36.   Friston K, Mattout J, Trujillo-Barreto N, Ashburner J, Penny W (2006) Variational free

energy and the laplace approximation. *NeuroImage* 34(1):220–234.

37.   R Core Team (2016) *R: A Language and Environment for Statistical Computing* (R

Foundation for Statistical Computing, Vienna, Austria).

38.   Bates DM (2005) Fitting linear mixed models in R. *R News* 5:27–30.

39.   Lenth RV (2016) Least-squares means: The R package lsmeans. *Journal of Statistical

Software* 69(1):1–33.

Acknowledgments

We thank Patrik Vuilleumier who created and shared the spider stimuli. This work was supported in part through the computational resources and staff expertise provided by Scientific Computing at the Icahn School of Medicine at Mount Sinai. Funding was provided by NIMH grant MH105515 and a Klingenstein-Simons Fellowship Award in the Neurosciences to D.S.; ERC Advanced Grant XSPECT-DLV-692739 to D.C. (Co-I); and Swiss National Science Foundation grant SNF 161077 to P.H.

Author Contributions

PH carried out the computational modeling and statistical analysis, interpreted the results, and drafted the manuscript. HL prepared materials, collected the data and critically revised the manuscript. IL contributed to the conception of the study, the computational modeling, the interpretation of the results, and critically revised the manuscript. CMR contributed to the interpretation of the results and critically revised the manuscript. DRB contributed to the computational modeling and critically revised the manuscript. DS conceived, designed and coordinated the study, contributed to data analysis and interpretation, and critically revised the manuscript. DC contributed to the conception of the study, data analysis and interpretation of results, and drafted the manuscript in its final form. All authors gave final approval for publication.

Conflict of Interests

All authors declare no conflicts of interest with regard to the current study.

Figures and Tables

Figure 1: Schematic description of experimental design and procedure. In each trial of the acquisition phase, participants were presented with one of two stimuli (schematic pictures of spiders, presented monocularly for 6 sec and suppressed from awareness by a CFS mask shown to the other eye). One image (spider A) always terminated with a mild electric shock to the wrist, whereas the other (spider B) never did. Halfway through the experiment, with no warning, the contingencies flipped and the reversal phase began: the formerly safe stimulus (spider B) now predicted the shock, and the old threat-associated one (spider A) was now safe. Each spider was shown 8 times in each phase. Trial order was pseudorandomized (see Materials and Methods) and spider identity (A and B) was counterbalanced across participants. To assess the success of the awareness manipulation, participants answered the questions "Which seen?" (1=flower, 2=spider) and "How confident?" (1=guess to 3=sure), presented binocularly (1.5 - 2 s each), beginning 1 s after the offset of every CS, and followed by an 8-10 s inter-trial interval (the questions are only shown here for the first depicted trial, but were repeated in all trials). Participants who underwent the same procedure without CFS were shown identical CSs, but the CFS mask was absent.

508
509

Figure 2: Physiological reversal learning. A. Time courses reveal reversal of threat responses with and without continuous flash suppression. Data points represent trial-wise mean responses to spider A (the CS+ during acquisition) and spider B (the CS- during acquisition). Both groups showed reversal learning, as indicated by greater responses to Spider A during the acquisition phase and greater responses to Spider B during the reversal phase. Error bars represent standard errors. **B. Mean reversal learning index for each group.** Error bars represent 95% confidence intervals, indicating that the interaction of stage and stimulus and thus the magnitude of reversal learning in both groups was significantly greater than zero. **C. Heightened anxiety is associated with impaired reversal learning under CFS.** A negative correlation between baseline anxiety measures and the strength of threat reversal learning is evident for state and trait anxiety. Blue lines show linear fits of each score to the reversal index, and ribbons around lines indicate bootstrapped 95% confidence intervals around the estimate. *Abbreviations:* STAIS/STAIT, state/trait anxiety subscale of the Spielberger State-Trait Anxiety Inventory; FSQ, Fear of Spider Questionnaire, ~, $P < .1$; *, $P < .05$.

528
529
Figure 3: Characterizing the relation between perceptual awareness and reversal learning
in the CFS group. A. Correlation between the awareness index of even and odd-numbered
trials. Each data point represents an individual participant. The strong positive correlation
between these independent measures of awareness demonstrates that individual participants'
awareness ratings - even those with extreme values of zero or one - are unlikely to be due to
measurement noise. For illustrative purposes, the color scheme marks all participants with an
awareness index of 0 in even trials in red (UA, unaware, $N = 27$) and classifies the rest of the
CFS group in 3 tertiles (T1-T3). Note that some data points overlap. **B. Reversal learning**
plotted against perceptual awareness for individual participants, for data obtained from
even-numbered trials. The color scheme is the same as in Panel A. **C. Reversal learning**
plotted against perceptual awareness for individual participants, for data obtained from
odd-numbered trials. Individual participants are marked with the same color as in the previous
panels; the overall distribution of participants is highly similar across panels. **D. Reversal**
learning as a function of perceptual awareness in the CFS group, using data pooled from
all trials. The intercept, indicating the magnitude of reversal learning in the absence of
awareness, is positive and significantly different from zero. **E. Reversal Index intercepts and**
their 95% confidence intervals in a series of regression models. Model 1 depicts the intercept
(the value of the reversal index when the awareness index equals zero) shown in Panel D. Model
2 shows the intercept when the regression model includes STAIT scores in addition to the
perceptual awareness index. Model 3 regresses the reversal index onto the perceptual awareness
index, STAIT and tracking scores. (Excluding the potential outlier in the top left corner of panel
D weakens significance of the intercept in model 1, $P = 0.07$; the intercepts of model 2 and 3
remain significant after removal of this outlier). Blue lines show linear fits, and ribbons around
lines indicate bootstrapped 95% confidence intervals around the estimate.

Model	Predictor	Beta	SE	*t*	*P*
1	Intercept	0.3	0.2	2.1	0.035
1	Awareness index	-0.1	0.4	-0.4	0.692
2	Intercept	1.4	0.5	3	0.004
2	STAIT	0	0	-2.3	0.024
2	Awareness index	-0.2	0.4	-0.5	0.596
3	Intercept	1.5	0.5	3.1	0.003
3	STAIT	0	0	-2.4	0.021
3	Tracking score	-0.3	0.3	-1	0.318
3	Awareness index	-0.2	0.4	-0.5	0.597

554 **Table 1: Regression coefficients for all awareness index models.** Reversal learning was the
555 dependent variable in all models. Model 1 included an intercept and the perceptual awareness
556 index; model 2 additionally included STAIT scores; model 3 additionally included STAIT and
557 tracking scores.