A BIRD'S WHITE-EYE VIEW ON AVIAN SEX CHROMOSOME EVOLUTION

Thibault Leroy[1], Yoann Anselmetti[1], Marie-Ka Tilak[1], Sèverine Bérard[1], Laura Csukonyi[1,2], Maëva Gabrielli[2], Céline Scornavacca[1], Borja Milá[3], Christophe Thébaud[2], Benoit Nabholz[1]*

[1] *ISEM, Univ Montpellier, CNRS, IRD, EPHE, Montpellier, France*
[2] *Laboratoire Evolution et Diversité Biologique (EDB), UMR 5174 CNRS – Université Paul Sabatier – IRD, Toulouse, France*
[3] *National Museum of Natural Sciences (MNCN), Spanish National Research Council (CSIC), Madrid, Spain*

* Corresponding author: benoit.nabholz@umontpellier.fr

ABSTRACT

Chromosomal organization is relatively stable among avian species, especially with regards to sex chromosomes. Members of the large Sylvioidea clade however have a pair of neo-sex chromosomes which is unique to this clade and originate from a parallel translocation of a region of the ancestral 4A chromosome on both W and Z chromosomes. Here, we took advantage of this unusual event to study the early stages of sex chromosome evolution. To do so, we sequenced a female (ZW) of two Sylvioidea species, a *Zosterops borbonicus and a Z. pallidus*. Then, we organized the *Z. borbonicus* scaffolds along chromosomes and annotated genes. Molecular phylogenetic dating under various methods and calibration sets confidently confirmed the recent diversification of the genus *Zosterops* (1-3.5 million years ago), thus representing one of the most exceptional rates of diversification among vertebrates. We then combined genomic coverage comparisons of five males and seven females, and homology with the zebra finch genome (*Taeniopygia guttata*) to identify sex chromosome scaffolds, as well as the candidate chromosome breakpoints for the two translocation events. We observed reduced levels of within-species diversity in both translocated regions and, as expected, even more so on the neoW chromosome. In order to compare the rates of molecular evolution in genomic regions of the autosomal-to-sex transitions, we then estimated the ratios of non-synonymous to synonymous polymorphisms (π_N/π_S) and substitutions (d_N/d_S). Based on both ratios, no or little contrast between autosomal and Z genes was observed, thus representing a very different outcome than the higher ratios observed at the neoW genes. In addition, we report significant changes in base composition content for translocated regions on the W and Z chromosomes and a large accumulation of transposable elements (TE) on the newly W region. Our results revealed contrasted signals of molecular evolution changes associated to these autosome-to-sex transitions, with congruent signals of a W chromosome degeneration yet a surprisingly weak support for a fast-Z effect.

1 Introduction

Spontaneous autosomal rearrangements are
common across higher metazoan lineages (Choghlan et
al. 2005). By contrast, sex chromosome architectures are
much more conserved, even across distant lineages
(Murphy et al. 1999; Raudsepp et al. 2004; Fraïsse et al.
2017). A growing number of studies have recently
observed departures from this pattern of evolutionary
conservation by detecting changes in the genomic
architecture of sex chromosomes in some particular
lineages - so-called neo-sex chromosomes – that are
mainly generated by fusion or translocation events of at
least one sex chromosome with an autosome (*e.g.* Kitano
& Peichel, 2012; Zhou & Bachtrog, 2012). Considering the
long-term conservation of sex chromosome synteny,
neo-sex chromosomes provide opportunities to
investigate the processes at work during the early stages
of sex chromosome evolution (Charlesworth et al. 2005).
Previous detailed studies have for example investigated
its important role for divergence and speciation (*e.g.*
Kitano et al. 2009; Yoshida et al. 2014; Bracewell et al.
2017).

From a molecular evolution point of view, an
important consequence of the transition from autosomal
to sex chromosome is the reduction in effective
population size (N_e). Assuming a 1:1 sex ratio, N_e of
transposed regions on the Y (or W) and X (or Z)
chromosomes are expected to decrease by three-fourths
and one-fourths, respectively (see Ellegren, 2009 for
details). According to the neutral theory, nucleotide
diversity is then expected to be reduced proportionally
to the reduction in N_e due to increased drift effects
(Vicoso & Charlesworth, 2006; Pool & Nielsen, 2007). N_e
reduction also induces a change in the balance between
selection and drift, with drift playing a greater role after
the translocation, thus reducing the efficacy of natural
selection to purge deleterious mutations from
populations. Mutations - including deleterious ones -
may also drift to fixation at a faster rate in sex-
chromosomes than in autosomes, thus generating
expectations for faster evolution at X and Y

chromosomes (so-called fast-X or fast-Y effects) (Mank et
al. 2007; Rousselle et al. 2016). In addition, given that
mutations are on average recessive, positive selection is
expected to be more efficient in the heterogametic sex
(Hvilsom et al. 2012; Nam et al. 2015). Suppression of
recombination is also expected to initiate a degenerative
process on the Y chromosome, that may result in the
accumulation of nonsynonymous deleterious
substitutions owing to a series of processes acting
simultaneously: Muller's ratchet, the Hill-Robertson
effect, and positive and background selection
(Charlesworth and Charlesworth, 2000). For the same
reason, transposable elements (TEs) are also expected to
accumulate soon after the cessation of recombination in
the Y chromosomes (Charlesworth 1991; Charlesworth et
al. 1994).

Except for few reported examples (de Oliveira et
al. 2005; Nanda et al. 2006; Kapusta & Suh, 2017;
O'Connor et al. 2018), all birds share a high degree of
synteny conservation across autosomal chromosomes
(Griffin et al. 2007; Nanda et al. 2008; Ellegren, 2010;
Völker et al. 2010; Warren et al. 2010; Ellegren, 2013)
and even more at the Z chromosome (Nanda et al. 2008).
A notable exception is the neo-sex chromosome of
Sylvioidea species, a superfamily of passerine birds in
which a translocation of a large part of the Zebra finch 4A
chromosome onto both the W and the Z chromosomes,
as characterized by genetic mapping (hereafter
translocations of the neoW-4A and neoZ-4A on ancestral
neoW-W and neoZ-Z respectively; Pala et al. 2012a).
These two autosome-to-sex chromosome transitions are
present in reed warblers (Acrocephalidae), old-world
warblers (Sylviidae) and larks (Alaudidae) (see also
Brooke et al. 2010), and therefore likely occurred in the
common ancestor of all present-day Sylvioidea (Pala et
al. 2012a,b; Sigeman et al. 2018), between 15 and 30
million years ago (Myrs) (Ericson et al. 2014; Prum et al.
2015; Nabholz et al. 2016). These two sex chromosome
translocations provide unique opportunities to
investigate the early stages of the W and Z chromosome
evolution.

Based on phylogenetic trees calibrated using
geological events (Moyle et al. 2009), *Zosterops* species
of the family Zosteropidae (more commonly referred to

as white-eyes) are considered to have emerged around the Miocene/Pliocene boundary. Considering both this recent emergence and the remarkable high diversity currently observed in this genus (more than 80 species), this group appears to have one of the highest diversification rates reported to date for vertebrates and is therefore considered as one of the 'great speciator' examples (Diamond et al. 1976; Moyle et al. 2009). White-eye species are typical examples of taxa spanning the entire "grey zone" of speciation (Roux et al. 2016). As a consequence of these different degrees of reproductive isolation between taxa, white-eyes have long been used as models to study bird speciation (*e.g.* Clegg and Philimore 2010; Melo et al. 2011; Oatley et al. 2012; Oatley et al. 2017). Among all white-eyes species, the Reunion grey white-eye *Zosterops borbonicus* received considerable attention over the last 50 years. This species is endemic from the volcanic island of Reunion and shows an interesting pattern of microgeographical variation, with five distinct colour variants distributed over four specific regions across the 2,500 km² of island surface. Both plumage color differentiation data (*e.g.* Gill et al. 1973; Milá et al. 2010; Cornuault et al. 2015) and genetic data (*e.g.* Milá et al. 2010; Delahaie et al. 2017) support this extensive within-island diversification. Despite its important role in the understanding of the diversification of *Zosterops* species, no genome sequence is currently available for this species. More broadly, only one *Zosterops* species has been sequenced to date (the silvereye *Z. lateralis,* Cornetti et al. 2015).

Here, we obtained detailed genome data from *Z. borbonicus*, a member of the Zosteropidae family and then arranged scaffolds into chromosomes to provide insights into the evolutionary processes that may have contributed to the early stages of the sex chromosome evolution. We also generated a more fragmented genome sequence for *Z. pallidus* for molecular dating and molecular evolution analyses. We found similar breakpoint locations for both translocated regions suggesting evolution from the same initial gene sets, and studied the molecular evolution of the two newly sex-associated regions. By comparing levels of within-species nucleotide diversity at autosomal and sex chromosomes, we found support for a substantial loss of diversity on both translocated regions, largely consistent with expectations under neutral theory. We then compared patterns of polymorphisms and divergence at neo-sex chromosome genes and found support for a considerable fast-W effect, but surprising weak support for a fast-Z effect. Investigations of candidate changes in base led to the identification of specific signatures associated with abrupt changes in recombination rates (reduction or cessation) of the two neo-sex regions. Finally, we reported larger transposable elements (TE) content on the newly W than on the newly Z regions, suggesting ongoing neoW chromosome degeneration.

RESULTS

ZOSTEROPS BORBONICUS GENOME ASSEMBLY

Using a strategy combining long-read sequencing with PacBio and short-read Illumina sequencing with both mate-pairs and paired-end reads, we generated a high-quality reference genome for a female Reunion grey white-eye captured during a field trip to Reunion (Mascarene archipelago, southwestern Indian Ocean). The 1.22 gigabase genome sequence comprises 97,537 scaffolds (only 3,760 scaffolds after excluding scaffolds smaller than 10kbp), with a N50 of 1.76 Mb (Fig. 1, Table S1). The BUSCO completeness statistic seems particularly good (93.0%). Among all investigated avian species, the 'GRCg6a' chicken genome assembly is the only one exceeding this value (93.3%) (Fig. S1). Compared to the other species, our *Z. borbonicus* reference assembly also exhibits the lowest proportions of 'missing' (2.5%, a value only observed for the reference chicken genome) and 'fragmented' genes (4.3%, a value which is 0.1% higher than for the reference chicken assembly).

Using the PASA pipeline combined with EVidenceModeler (Haas et al. 2008; Haas et al. 2011) and several *in silico* tools trained by the RNAseq data, a total of 22,565 gene models were predicted. Among these 22,565 genes, half of the "hidden" avian genes identified by Botero-Castro et al. (2017), *i.e.* a large fraction of avian genes in GC-rich regions missing from most avian genome assemblies, were recovered (1,072 out of 2,132).

Figure 1: Z. borbonicus *genome assembly and gene model summary statistics. The reported numbers of reads and the total numbers of bases sequenced correspond to the trimmed reads. Photo credits: Maëva Gabrielli & Laurent Brillard (http://laneb.re)*

The vast majority of the 22,565 CDS (83.3%) are supported by at least one RNA-seq read (18,793 CDS), including 17,474 with a FPKM above 0.01. Among the 22,565 gene models, TE content is globally low (8.5%), but 4,941 genes exhibit a TE content greater than 0.25, including 1,275 genes predicted to be TE over the total length of the gene (Fig. S2). Still more broadly, the level of expression is strongly negatively correlated with the within-gene TE content (r^2=0.038, p<2.2x10^{-16}, after excluding 709 genes with FPKM>100).

Additionally, we generated a genome assembly of a female Orange River white-eye (*Z. pallidus*) using mate-pair and paired-end reads. This assembly is much more fragmented (170,557 scaffolds and N50 = 375 Kb, Table S1) than the *Z. borbonicus*. Both *Z. borbonicus* and *Z. pallidus* assemblies are available from Figshare repository URL: https://figshare.com/s/122efbec2e3632188674.

[GenBank accessions will be provided upon publication].

REFERENCE-ASSISTED GENOME ORGANIZATION

We then anchored scaffolds using the v3.2.4 reference genome of the zebra finch (Warren et al. 2010) assuming synteny. We used the zebra finch as a pivotal reference since this reference sequence is of high-quality, with 1.0 of 1.2 gigabases physically assigned to 33 chromosomes including the Z chromosome, plus three additional linkage groups based on genetic linkage and BAC fingerprint maps (Warren et al. 2010). We anchored 928 among the longest scaffolds to the zebra finch chromosomal-scale sequences, thus representing a total of 1.01 Gb among the 1.22 Gb of the *Z. borbonicus* assembly (82.8%).

In parallel to the zebra finch-oriented approach, we used DeCoSTAR (Duchemin et al. 2017), a tool that improves the assembly of several fragmented genomes by proposing evolutionary-induced adjacencies between scaffolding fragments containing genes. To perform this analysis, we used the reference sequences of 27 different avian species (Table S2) with associated gene tree phylogeny of 7,596 single copy orthologs (trees are available from the following Figshare repository, URL: https://figshare.com/s/122efbec2e3632188674).

Among the 97,537 scaffolds (800 containing at least one orthologous gene), DeCoSTAR organized 653 scaffolds into 188 super-scaffolds for a total of 0.837 Gb (68.5% of the *Z. borbonicus* assembly), thus representing a 2.59-fold improvement of the N50 statistic (4.56Mb). Interestingly, among the 465 scaffold junctions, 212 are not only supported by gene adjacencies within the other species, but are also supported by at least one *Z. borbonicus* paired-end read. From a more global point of view, DeCoSTAR not only improved the *Z. borbonicus* genome, but also those of 25 other species (mean gain in N50 over 11Mb, representing 3.30-fold improvement on average, range: 1.0-6.02). The only exception is the already well-assembled chicken genome reference. For all these species, the proposed genome organization ("agp file") were made available at the following URL: https://figshare.com/s/122efbec2e3632188674.

We then combined zebra finch-oriented and DeCoSTAR approaches for the *Z. borbonicus* genome, by guiding DeCoSTAR using the a priori information of the zebra finch-oriented approach to get beyond two limitations. First, DeCoSTAR is a gene-oriented strategy, and thus cannot anchor scaffolds without genes that have orthologous analogues in the other species, which is generally the case for short scaffolds. Second, the zebra-finch oriented approach assumes a perfect synteny conservation between *T. guttata* and *Z. borbonicus*, which is unlikely. By combining both approaches, we were able to anchor 1,082 scaffolds, including 1,045 scaffolds assigned to chromosomes representing a total 1.047 Gb (85.7% of the *Z. borbonicus* assembly). In addition, DeCoSTAR helped propose more reliable *Z. borbonicus* chromosomal organizations for these 1,045 scaffolds by excluding some *T. guttata*-specific intra-chromosomal inversions.

ASSIGNING SCAFFOLDS TO W AND Z CHROMOSOMES

To identify sex chromosome scaffolds, we first mapped trimmed reads from males and females *Z. borbonicus* individuals which were previously sequenced by Bourgeois et al. (2017) and then computed median per-site coverage over each scaffold for males and females (Fig. 2). After taking into account differences in coverage between males and females, we then identified scaffolds that significantly deviated from 1:1 and identified neoW and neoZ scaffolds (see methods section). This strategy led to the identification of 218 neoW (7.1 Mb) and 360 neoZ scaffolds (91.8 Mb) among the 3,443 scaffolds longer than 10 kb (Fig. 2). Among the 360 neoZ scaffolds assigned by coverage, 339 scaffolds were already anchored to the neoZ chromosome based on the synteny-oriented approach, thus confirming the accuracy of our previous assignation and suggesting that we have generated a nearly complete Z chromosome sequence.

Due to the absence of a chromosomal-scale W sequence for a passerine bird species, we are unable to provide a chromosomal structure for the 218 neoW scaffold. We however assigned 174 among the 218 neoW scaffolds to the neoW-4A region. These scaffolds representing a total length of 5.48 Mb exhibited high levels of homology with the zebra finch 4A chromosome. We found support for neoW-4A scaffolds aligning between positions 21,992 and 9,603,625 of the *T. guttata* 4A chromosome (thus representing 57% of the corresponding 9.6 Mb 4A region). Leaving aside the difficulty of sequencing and assembling the neoW chromosome (Tomaszkiewicz et al. 2017), a large part of the difference between the total assembled size and the corresponding region in *T. guttata* is likely due to a 1.75 Mb chromosomal deletion on the neoW-4A. Indeed, we found no neoW scaffold with a homology to the large 4A *T. guttata* region between positions

1,756,080 and 3,509,582. Remaining neoW scaffolds,
i.e. those having no homology with the 4A *T. guttata*
region, were considered as sequences belonging to the
ancestral W chromosome (hereafter neoW-W region),
except for six scaffolds showing reliable hits but only at
specific locations of the scaffold, and for which the
accuracy of any assignation was considered too low.

Figure 2: Identification of Z and W scaffolds using
coverages of male and female individuals.
Normalized sum of median per-site coverage of
trimmed reads over the 5 males (x-axis) and 7 females
(y-axis). Values were only calculated for scaffolds
longer than 10-Kb. Linear regressions for slope=1
(blue) and slope=0.5 (yellow) are shown. Scaffolds
were assigned following the decision rule described in
the material and methods section.

MOLECULAR DATING

The molecular phylogenetic analyses were
aimed at estimating the divergence time of the
Zosterops genus. Indeed, even if our focal dataset is
composed of only 3 species, the divergence between *Z.*
pallidus / Z. borbonicus and *Z. lateralis* represents the
first split within the genus except *Z. wallacei*, *i.e.* the
origin of clade B in Moyle et al. (2009). As a
consequence, our phylogeny (Fig. 3, S3) is expected to
provide an accurate estimate for the diversification of
the *Zosterops* lineage. For this molecular dating, we

added 3 species to the 27 species used in the previous
analyses and generated gene sequence alignments.
Given that these newly added species - namely the
silvereye, Orange River white-eye and willow warbler -
have no gene models available yet, we used the AGILE
pipeline (Hughes & Teeling 2018, see method) to
obtain orthologous sequences. Due to the inherent
computational burden of Bayesian molecular dating
analyses, we randomly selected 100 alignments among
the least GC rich single-copy orthologs and performed
10 replicated analyses (chronograms available at
FigShare URL:
https://figshare.com/s/122efbec2e3632188674).
Indeed, GC poor genes are known to be slowly and
more clock-like evolving genes as compared to the
other genes (Jarvis et al. 2014).

We used several combinations of fossil
calibrations and substitution models (Table S3). For the
radiation of Neoaves, all analyses led to molecular
dating consistent with Jarvis et al. (2014) and Prum et
al. (2015) with estimates around 67-70 Myrs, except for
the calibration set 4 (82 Myrs), albeit with large
confidence intervals (CI = 64 - 115 Myrs) (Table 1).
Calibration set 4 is very conservative with no maximum
calibration bound except for the Paleognathae /
Neognathae set to 140 Myrs. In contrast, calibration set
3 is the more constrained with the Suboscines / Oscines
split bounded between 28 - 34 Myrs. Unsurprisingly,
this calibration led to the youngest estimates, dating
the origins of passerines at 59 Myrs (56 - 63 Myrs) and
the Paleognathae / Neognathae at 65 Myrs (63 - 69
Myrs).

In all runs, our estimates of the origin of
Zosterops have lower limits of CI including 2.5 Myrs and
mean estimates are also often considerably lower than
this value (Table 1). Using the calibration set 1, the ten
replicated datasets gave a mean age estimate for the
origin of *Zosterops* around 2 Myrs (Table 1). More
broadly, our analysis are consistent with a recent origin
of *Zosterops*, with a crown clade age of less than 5
Myrs and probably between 1 and 3.5 Myrs (Table 1).

Table 1: Molecular dating analyses. *Mean dates are indicated and 95% confidence intervals (CI) are provided within*
parentheses

Datasets*	Calibration set**	Crown Zosterops	Zosterops / Phylloscopus	Corvus / Passerida	Passerines	Crown Neoaves
1	1	2.8 (1.7, 4.2)	17.2 (14.2, 20.4)	25.8 (22.6, 28.9)	62 (58.3, 65.5)	67.1 (63.2, 71.2)
1	2	3.1 (1.8, 4.6)	18.4 (15.2, 22.2)	27.3 (23.7, 31.5)	64.8 (58.5, 71.9)	70.1 (63.5, 77.5)
1	3	1.9 (1.3, 2.8)	12.7 (10.9, 14.7)	19.5 (17.5, 21.4)	59.1 (56.5, 62.9)	65 (63, 69.1)
1	4	3.5 (1.9, 6.1)	21.4 (15.4, 31.6)	31.9 (23.9, 45.2)	75.9 (59.4, 106.4)	82.1 (64.2, 114.9)
2	1	0.9 (0.5, 1.4)	7.8 (5.1, 10.3)	13.4 (9.7, 16.5)	62.9 (58.9, 66.3)	68 (63.7, 71.9)
3	1	2.0 (1.3, 3.2)	12.4 (9.7, 16.2)	19.5 (15.9, 23.5)	62 (58.5, 65.6)	67.1 (63.1, 71.1)
4	1	1.8 (1.2, 2.9)	11.9 (8.6, 15.4)	18.7 (14.1, 22.8)	61.1 (57.5, 65.3)	68.2 (64.2, 72.6)
5	1	1.7 (1, 2.6)	11 (8, 14.7)	17.3 (13.1, 21.8)	61.6 (57.7, 65.4)	68.1 (64, 72.3)
6	1	3.2 (1.9, 4.9)	16.8 (13.9, 19.9)	23.8 (20.8, 26.5)	62 (58.3, 65.4)	67.2 (63.1, 71.3)
7	1	2.2 (1.2, 3.5)	13.3 (9.8, 16.8)	20.6 (16.5, 24.3)	64.7 (62.9, 65.7)	71.8 (69.7, 73.9)
8	1	2.1 (1.3, 3.5)	13.2 (9.9, 17.2)	21.2 (16.9, 25.3)	64.6 (62.4, 65.7)	69.7 (67.3, 71.6)
9	1	1.5 (0.9, 2.5)	9.5 (6.6, 12.6)	16.9 (12.8, 20.9)	65.3 (63.5, 66.7)	72.3 (69.9, 74.5)
10	1	1.9 (1.3, 2.8)	12.4 (9.6, 14.9)	19.4 (15.6, 22.3)	61.7 (57.4, 65.5)	68.5 (63.8, 73.1)

364 *Different numbers indicate independent replicated datasets of 100 randomly selected orthologs*

365 ** *Number corresponds to the different calibration sets used and presented in Table S3*

Figure 3: Molecular phylogeny and dating of the investigated passerine birds. *Estimates of molecular dating are based on dataset 1 and calibration set 1 (Table S3). For greater clarity, the cladogram focuses on passerine birds only, but see Fig S3 for a cladogram of the inferred phylogeny for the whole set of investigated avian species.*

Additionally, we performed a completely independent molecular dating by applying the regression method proposed in Nabholz et al. (2016). Based on 9 full *Zosterops* mitochondrial genomes, we calculated molecular divergence. The divergence between *Z. lateralis* and the other *Zosterops* has a median of 0.205 subst./site (max = 0.220, min = 0.186) for the third codon position. Assuming a median body mass for the genus of 10.7g (Dunning, 2007), we estimated a divergence date between 2.3 and 6.2 Myrs. These estimates are in line with our previous dating based on nuclear data and with molecular dating based on fossil calibration confirming the extremely rapid diversification rate of white-eyes.

CHROMOSOMAL BREAKPOINTS

Our synteny-oriented approach using pairwise whole-genome alignments of *Z. borbonicus* and *T. guttata* sequences helped us identifying the chromosomal breakpoint of the neoZ sex chromosome. We reported a scaffold (scaffold329) with long sequence alignments with both the 4A and the Z chromosomes (Fig. 4). Considering the intervals between the last LastZ hit on the 4A and the first one on the Z chromosome, we estimated that this breakpoint occurred between positions 9,605,374 and 9,606,437 of the 4A zebra finch chromosome (genome version : v.3.2.4), which is fairly close to the estimate of 10Mb previously reported by Pala et al. (2012a). Based on the soft-masked version 3.2.4 of the zebra finch genome, this 1kb region is well assembled (no ambiguous "N" bases) and shows no peculiarities in TE or GC content (7.4% and 39.0%, respectively) as compared to the rest of the zebra finch 4A chromosome (18.7% and 43.7%).

To discard the potential confounding factor of a sequence artifact due to a chimerism in the *Z. borbonicus* assembly, we used the procedure for the sequence assembly of *Z. lateralis* (Cornetti et al. 2015) and identified a scaffold (LAII01000008.1) supporting the same chromosomal breakpoint. Using the *Z. lateralis* sequence, we estimated that this breakpoint occurred between positions 9,605,524 and 9,606,431.

Figure 4: Fine scale genome architecture of the Sylvioidea neo-sex chromosomes. Ancestral region of the neoW (neoW-W), ancestral autosomal 4A chromosome and ancestral Z chromosome (neoZ-Z) are shown in green, purple and yellow, respectively. Translocated regions of the ancestral 4A chromosome (neoW-4A and neoZ-4A) are shown with a purple outline. LastZ hits on scaffold supporting the chromosomal breakpoints are shown using the same color code. For ease of illustration, coordinates on the Z and 4A chromosomes are reversed (i.e. the white disk represents the start of the chromosome).

We then investigated the chromosomal breakpoint for the neoW (Fig. 4). We identified a candidate scaffold, *scaffold1768_size33910*, with alignment hits on both the 4A *T. guttata* and the W of *F. albicollis* (Smeds et al. 2015). Among all W scaffolds, this scaffold aligns at the latest positions of the 4A *T. guttata* chromosome (from 9,590,067 to 9,603,625), which is remarkably close to our estimation for the region translocated on the neoZ chromosome. Some other hits have been however observed, including a 2-kbp region at the end of this scaffold which is homologous to the beginning of the chromosome 4A (1,667,182-1,669,166), probably as a result of the high activity of transposable

elements after the translocation (see below). Considering also alignments against contigs of the *F. albicollis* W chromosome, we estimated that the chromosomal breakpoint probably occurred between positions 9,603,625 and 9,605,621.

CHROMOSOMAL-SCALE ESTIMATES OF NUCLEOTIDE DIVERSITY

We used sequencing data from 6 *Z. borbonicus* individuals (3 males and 3 females) sequenced by Bourgeois et al. (2017) to explore the genomic landscape of within-species diversity. Over all autosomal 10-Kb

windows, mean nucleotide diversity (Tajima's π) estimates were roughly similar in males and females (π_{males}=1.82e-3 and $\pi_{females}$=1.81e-3, respectively). The nucleotide diversity landscape greatly varies within and between chromosomes (A in Fig 5, Fig S4 & S5). In addition, we identified some series of 10-Kb windows with very low level of nucleotide diversity (red bars, Fig. 5A). Interestingly, some of these regions also exhibit the highest negative values of Tajima's D (red bars, Fig. 5B & S6; *e.g.* the end of the chromosome 2).

Figure 5: Intra- and inter-chromosomal variations in nucleotide diversity. *Variations of Tajima's π_{males} (A) and D_{males} (B) estimates along the 6 Z. borbonicus macrochromosomes, the autosomal 4A and the neo-Z chromosome. The two metrics were calculated in non-overlapping 10-Kb sliding windows. Top 2.5% and bottom 2.5% windows are shown in green and red, respectively. For both Tajima's π_{males} and Tajima's D_{males}, each bar shows the deviation from the mean genomic value over the whole genome (Tajima's π_{males} and Tajima's D_{males} baselines: 1.82e-3 and -0.26, respectively). C) Interchromosomal differences in Tajima's $\pi_{females}$ between autosomes and sex chromosomes. Fig. S5 and S6 show Tajima's pi and D along most* Z. borbonicus *chromosomes.*

Small interchromosomal differences in the
distribution of nucleotide diversity values were
detected for both female- and male-based estimates,
suggesting that both datasets give similar results at the
chromosomal level, except for the neoZ chromosome
for which substantial differences were observed
between the two datasets (Fig. S4). Even considering
this source of variability, the neoZ chromosome still
shows significant deviation from the mean autosomal
diversity for both datasets ($\pi_{females}$=1.04e-3 and
π_{males}=1.34e-3, respectively; p<2e-16 for both datasets),
thus representing 57.6% and 73.5% of the mean
autosomal diversity. This reduced level of nucleotide
diversity was similarly detected for the neoZ-Z and the
neoZ-4A regions of the neo-Z chromosome (C, Fig. 5).
Based on both datasets, a lower nucleotide diversity
was observed on neoZ regions as compared to the
autosomal chromosomes. Mean π_{males} was estimated to
1.21e-3 for the neoZ-4A region and 1.35e-3 for the
neoZ-Z region, corresponding to 66.6% and 74.2% of
the autosomal diversity. Mean $\pi_{females}$ values are
roughly similar with 1.37e-3 and 1.00e-3, thus
representing 76.2% and 55.5% of the autosomal
diversity, respectively.

Similarly, data from females only were used to
estimate the level of within-species diversity variation
on the neoW chromosome (C, Fig. 5). We similarly
observed a reduced level of diversity as compared to
the autosomal chromosomes (mean $\pi_{females}$=5.87e-4,
p<2e^{-16}), with only one-third (32.5%) of the mean
nucleotide diversity estimated for the autosomes.
Significant differences in Tajima's $\pi_{females}$ were
observed on the neoW-W region of the neoW
chromosome and on the neoW-4A translocated region
(p=0.009), with higher diversity on the neoW-W
(π=1.08e-3) as compared to the neoW-4A (π=4.65e-4).
Median values are however much more consistent
between the two regions (π=1.25e-4 and π=1.16e-4,
respectively), suggesting that few neoW-W windows
greatly contributed to this discrepancy (C, Fig. 5).

RATIOS OF NON-SYNONYMOUS TO
SYNONYMOUS POLYMORPHISMS (Π_N/Π_S)
AND SUBSTITUTIONS (D_N/D_S)
490

We computed π_N/π_S ratios among all genes in
autosomal, neoZ and neoW chromosomes. Estimated
π_N/π_S of chromosome 4A genes is slightly lower as
compared to the rest of autosomal chromosomes
(π_N/π_S= 0.212 vs. π_N/π_S = 0.170). NeoZ-Z exhibits slightly
higher values than both autosomal sets (π_N/π_S=0.282),
however no or very little difference has been observed
between neoZ-4A (π_N/π_S=0.181) and autosomal
chromosomes. On the contrary, π_N/π_S ratios on genes
of the neoW chromosome are very high, with
respectively π_N/π_S=0.418 for the neoW-4A and π_N/π_S=
0.780 for the neoW-W regions.

We estimated d_N/d_S ratios for *Z. borbonicus*
and 10 additional passerine species (all passerines
except *A. chloris* in Fig. 1) for a total of 6,339
alignments of single-copy orthologs, corresponding to
6,073 autosomal genes, 66 on the remaining autosomal
region of the 4A chromosome (hereafter autosome-
4A), 164 on the neoZ-Z, 54 on the neoZ-4A and 17 on
the neoW-4A. For the neoZ-4A and neoW-4A, we made
a special effort to identify gametologs (paralogous
sequences between neoZ-4A and neoW-4A genes,
which were previously excluded during the filtering of
1:1 orthologs).

We first compared to evolution of neoZ-4A
and neoZ-Z genes. To do that, we randomly
subsampled the data to match the number of genes in
neoZ-4A region (54) before concatenation and then
computed d_N/d_S. The variability in d_N/d_S is evaluated by
bootstrapping genes and creating repeated
concatenation of 54 genes for each genomic region.

For all species, the d_N/d_S ratio reaches
significantly higher values for neoZ-Z genes than for
autosomal genes (Fig. 6). Surprisingly, we found no
evidence for an increase in d_N/d_S ratios in the neoZ-4A
when compared to autosomes (Fig. 6, species in the red
frame). For both the willow warbler and *Zosterops*
species, d_N/d_S ratios are lower in neoZ-4A genes than in

Figure 6: Variation in d_N/d_S ratios among chromosomal regions for the 11 investigated species. *Estimates are performed at the genus level, i.e. on branches before the split of the two* Corvus *(C. cornix and* C. brachyrhynchos*) and the three* Zosterops *species (Z. lateralis, Z. pallidus and Z. borbonicus). The red box shows species with the translocated neoZ-4A region For these species, the 4A:0-9.6Mb region corresponds to the neoZ-4A and the Z corresponds to the neoZ-Z.*

autosomal regions (willow warbler: neoZ-4A d_N/d_S =
0.09 (95% CI=0.07-0.11) *vs.* autosomal d_N/d_S = 0.11
(95% CI=0.08-0.15);   for white-eyes: neoZ-4A d_N/d_S =
0.10 (95% CI = 0.08-0.13) *vs.* autosomal d_N/d_S = 0.14
(95% CI=0.09-0.20)). This also holds true when we
compare genes of the ancestral 4A chromosome
translocated on the neoZ chromosome (neoZ-4A,
"chromosome 4A:0-9.6Mb" in Fig. 6) and genes of the
4A chromosome ("chromosome 4A:9.6-20.7Mb"). Even
if we report slightly higher d_N/d_S for the translocated
region as compared to the rest of the 4A chromosome,
such a difference between the two ancestral 4A regions
are also observed in several other species that do not
have the translocation (e.g., *M. vitellinus*, *T. gutatta* or
Z. albicollis; Fig. 6), including a much bigger difference
for *T. guttata* (Fig. 6). As a consequence, our d_N/d_S
analysis did not provide any support for a higher d_N/d_S
ratio associated to the autosomal-to-Z translocation.

Next, we computed the d_N/d_S of the branch
leading to the neoZ-4A and neoW-4A copies in the
Zosterops borbonicus genome. In this case, the d_N/d_S of
the neoW-4A genes were significantly higher than the

d_N/d_S of the neoZ-4A copies (mean d_N/d_S = 0.531,
sd = 0.417 for neoW-4A genes; mean d_N/d_S = 0.194, sd
= 0.234 for neoZ-4A genes; Wilcoxon signed rank test,
p-value =7.7e-5, Fig. 7). The increase in d_N/d_S is
particularly strong, including some neoW-4A genes
with d_N/d_S close or slightly higher than 1 suggesting
ongoing pseudogenization.

Figure 7: The d_N/d_S ratios of the neoW-4A and neoZ-4A copies (gametologs) of the* Zosterops borbonicus *genomes. *For each gene, horizontal lines linked values of the two gametologs.*

GC, GC* and Transposable Elements (TE) contents

Next, we investigated the potential change in
base composition after the translocation events, due to
changes in recombination rates and more precisely, the
recombination-associated effect of GC-biased gene
conversion (gBGC, Duret & Galtier 2009, Nabholz et al.
2011, Weber et al. 2014). For this purpose, we
computed the GC content over 10kb sliding windows
(Fig. S7) and the GC content at equilibrium for
orthologous sequences (Fig. S8). As expected,
differences in GC contents are observed between
chromosomes, with a higher GC content in short
chromosomes. Among all chromosomes, the neoZ

chromosome showed the second lowest median GC
rate. Interestingly, GC content is lower in the neoZ-4A
regions as compared to the autosomal 4A chromosome
(p=2.2e-4) or as compared to the neoW-4A, although
this difference is only marginally significant (p=0.059).
Slight differences in GC content are observed between
neoW-4A (p=0.066) and neoW-W (0.096) as compared
to the autosomal-4A chromosome suggesting that the
GC content of the neoW-4A likely decreased after the
translocation, but to a lesser extent than for the neoZ-
4A (Fig. 8). We also evaluated variation in GC content at
equilibrium (GC*) for the newly gametologs, *i.e.*
paralogous genes in the neoW-4A and neoZ-4A regions

(Fig. 9A). For these 17 genes, all Passerida species
without the neo-sex regions exhibited a GC* between
0.7 and 0.8 (Fig. 9A). Passerida species with these neo-
sex chromosomes showed a slightly reduced GC* at
neoZ-4A genes (mean GC* = 0.57 and 0.68 for the
willow warblers and the white-eyes respectively, and a
strongly reduced GC* content at neoW-4A genes
(mean=0.4, 95% CI = 0.30-0.51, Fig. 9A). In contrast, the
GC* of the non-translocated region of the ancestral
chromosome 4A (position 9.6-20.7Mb) apparently
remains unchanged between the Sylvioidea and the
other Passerida (GC* between 0.71 and 0.86, Fig. 9B).

Figure 8: Variation in GC (top) and TE (below) contents between chromosomes. *GC and TE contents were estimated along the whole genome using non-overlapping 10-kb sliding windows. Significance was evaluated using multiple t-tests. Thresholds: . < 0.1, * < 0.05, ** < 0.01, *** < 0.001*

Figure 9: *Phylogenetic relationships and estimation of GC equilibrium (GC*). A) Estimation using the 17 genes coded by the translocated region of the ancestral chromosome 4A (0-9.6Mb). Bold font indicates NeoW-4A sequences. B) Estimation using the 79 genes coded by the non-translocated region of the ancestral chromosome 4A (9.6-20.7Mb). Branch color and numbers above branches indicated GC*, mean and 95% confidence interval obtained by bootstrap are in parenthesis. Number in blue in front of the node indicates ultra-fast bootstrap values lower than 100%.*

We also found support for a higher abundance
of transposable elements (TE) in the W chromosome as
compared to all other chromosomes (Fig. 8 & S9).
Overall, 45.1% of the cumulative size of the scaffolds
assigned to the neoW-W is composed of transposable
elements, which represents a 3.31- to 9.95-fold higher
content than on the autosomal chromosomes.
Interestingly, the TE content over 10-kb windows (Fig.
8) is also much higher on the neoW-4A than on the
autosomal 4A chromosome, the other autosomes or,
even more interesting, the neoZ-4A region (p<2e-16 for
all these comparisons). Even if RepeatModeler was
unable to classify most *Z. borbonicus*-specific TE ("no
category" in Fig S9), Class I LTR elements seem to have
greatly contributed to this higher content in the neoW-
W, as well as in the neoW-4A chromosome (Fig S9).

DISCUSSION

A NEW HIGH QUALITY REFERENCE GENOME FOR SYLVIOIDEA AND *ZOSTEROPS*

620
621

Using a combination of short Illumina and long
PacBio reads, we have generated a high-quality bird
assembly of *Z. borbonicus* with a N50 scaffold
exceeding one megabase, which is comparable to the
best passerine reference genomes available to date
(Kapusta & Suh, 2017; Peona et al. 2018). Similar
conclusions can be drawn by comparing BUSCO
analyses between this assembly and a set of other 26
extant avian genome assemblies (Fig. S1). This
assembly is of equivalent quality to the well-assembled
congeneric species *Z. lateralis* (Cornetti et al. 2015),
thus jointly representing important genomic resources
for *Zosterops*, a bird lineage exhibiting one of the
fastest rates of species diversification among
vertebrates (Moyle et al. 2009).

Avian GC-rich regions are notoriously difficult to assemble (Botero-Castro et al. 2017; Tilak et al. 2018; Peona et al. 2018). Combining moderate coverage (10x) PacBio and Illumina sequencing technologies, we have generated gene models for half of the "hidden" genes of Botero-Castro et al. (2017). The use of long PacBio reads seems therefore a promising solution to partially address the under-representation of GC-rich genes in avian genomes. In the future, it will be interesting to combine long read technologies such as PacBio or Oxford Nanopore with the Illumina library preparation proposed by Tilak et al. (2018).

To further improve the contiguities of the *Z. borbonicus* sequence, we used 26 avian species as pivotal resources to chromosomally organized scaffolds of the *Z. borbonicus* assembly. This strategy has led to a 1.047 Gb chromosome-scale genome sequence for *Z. borbonicus* (Table S1). We were able to obtain assembly statistics comparable to some assemblies using high coverage long-read data (Weissensteiner et al. 2017) or a pedigree linkage map (Kawagami et al. 2014). Importantly, the application of the DeCoSTAR strategy has not only improved the *Z. borbonicus* genome, but also those of 25 other species. Indeed, at the notable exception of the chicken reference genome, all genome assemblies have been improved using DeCoSTAR (4.19-fold improvement of the median N50), thus demonstrating the utility of the inclusion of other genome assemblies, even fragmented, to polish the genome assembly of a species of interest (Duchemin et al. 2017; Anselmetti et al. 2018).

Confirming *Zosterops* as great speciators

The availability of the *Z. borbonicus* genome sequence is also an important step to study the evolution of the *Zosterops* lineage, which was described as the "Great Speciator" (Moyle et al. 2009; Cornetti et al. 2015). Indeed, the availability of sequence data for two new species (*Z. borbonicus* and *Z. pallidus*), in addition with the sequence of *Z. lateralis* (Cornetti et al. 2015) helped us to validate the recent origin of this taxon. Moyle et al. (2009) estimated that the white-eyes genus (except *Z. wallacei*) originated in the early pleistocene (~2.5 Myrs). With more that 80 species, this clade exhibits an exceptional rate of diversification compared to other vertebrates (net rate of diversification without extinction (r) : 1-2.5 species per Myr, Magallon and Sanderson, 2001 cited in Harmon, 2018).

The divergence time estimated by Moyle et al. (2009) was based on a biogeographic calibration obtained from the ages of the Solomon Islands. These biogeographic calibrations should however be interpreted with care as previous phylogenetic studies found evidence for older lineages than the emergence ages of the islands for which they are endemic (Heads, 2005; Heads, 2011). This could be the consequence of extinction of mainland relatives leading to long branches of some island species (Heads, 2005; Heads, 2011). More recently, Cai et al. (2019) have obtained similar dates using a larger phylogeny but, again, have applied a biogeographic calibration. In this study, we took the opportunity to combine genetic information of these three *Zosterops* with the other investigated bird species to obtain new estimates using independent fossil calibrations to reassess the conclusions of Moyle et al. (2009) regarding the recent and extensive diversification of *Zosterops*. Using two independent datasets (mitochondrial and nuclear) and methods, we confirmed the recent origin of the genus. Our analyses are consistent with a diversification of white-eyes over less than 5 Myrs and that could be as young as 1 Myr. With the exception of the African Great Lakes cichlids (Genner et al. 2007), this represents one of the most exceptional diversification rates among vertebrates (Lagomarsino et al. 2016). As an example, it is more than ten times higher than the average diversification rate estimated across all bird species by Jetz et al. (2009). Even the large and relatively recent radiation of the Furnariidae (ovenbirds and woodcreepers) has a net rate of diversification markedly lower than the white-eyes (r = 0.16; Derryberry et al. 2010).

NO EVIDENCE FOR A FAST-NEOZ EFFECT

Taken all together, our analyses are surprisingly consistent with a pattern of substantial reduction of nucleotide diversity, but a low impact of the autosome-to-Z translocation on the molecular evolution of *Z. borbonicus*.

First, using pairwise genome alignments of the zebra finch genome with either the *Z. borbonicus* or the *Z. lateralis* genomes, we found support for a narrow candidate region of 1kb around position 9.606 Mb of the v.3.2.4 zebra finch 4A chromosome, in which the chromosomal breakpoint likely occurred. This result is consistent and fairly close to the estimate of 10Mb suggested by Pala and collaborators (2012a) who first demonstrated the translocation of approximately a half of the zebra finch 4A on the Z chromosome using an extended pedigree of the great reed warbler, a Sylvioidea species. A recent article reported a similar estimate (9.6 Mb) in another Sylvioidea species, the common whitethroat (Sigeman et al. 2018). To get this estimate, these latter authors used a very similar approach to ours (H. Sigeman, personal communication).

Second, relative estimates of within-species diversity on both sides of this chromosomal breakpoint (*i.e.* neoZ-4A and neoZ-Z regions) were estimated. As compared to all autosomes, neoZ-4A and neoZ-Z regions of the neo-Z chromosome exhibit reduced levels of within-species diversity in both the ancestral Z chromosome (*i.e.* neoZ-Z:autosomes = 0.605) as well as in the newly translocated region (neoZ-4A:autosomes=0.782), consistent with a substantial loss of diversity associated with this autosome-to-sex transition, following the expected effects of changes in effective population sizes. The neoZ-4A:autosomal-4A nucleotide diversity reported is slightly higher than 0.75 but is in line with a previous report for the common whitethroat (0.82, Pala et al. 2012b). Strongest deviations have however been reported in two other Sylvioidea species, namely the great reed warbler and the skylark (0.15 and 0.42, respectively) but some of the variation might be explained by the moderate number of loci analyzed (Pala et al. 2012b).

Sex ratio imbalance or selection are known to contribute to strong deviations from neutral equilibrium expectations of three-fourths (reviewed in Ellegren 2009 and Wilson Sayres, 2018).

Third, we found no support for a fast-Z evolution in the neoZ-4A region, neither an elevated ratio of non-synonymous to synonymous polymorphisms (π_N/π_S) nor substitutions (d_N/d_S) at neoZ-4A genes when compared with autosomal-4A sequences. This result is intriguing as the decrease of nucleotide diversity observed on the neoZ-4A is expected to reflect a decrease in N_e and, therefore, a decrease in the efficacy of natural selection. This should result in an increase of the frequency of slightly deleterious mutations (Ohta 1992; Lanfear et al. 2014). The increase in d_N/d_S of avian Z-linked genes compared to autosomes - a pattern that we recovered well in our analyses - has often been interpreted in that way (Mank et al. 2010; Wright et al. 2015). We propose several potential hypotheses to explain the absence of fast-Z on the neoZ-4A regions. First, the hemizygous status of neoZ-4A regions could help to purge the recessive deleterious mutation and, therefore, limit the increase of π_N/π_S and d_N/d_S as reported in Satyrinae butterflies (Rousselle et al. 2016). Second, the intensity of purifying selection is not only determined by N_e but also by gene expression (Drummond & Wilke 2008; Nguyen et al. 2015) and recombination rate (Hill & Roberston 1966). It is therefore possible that the expression pattern of neoZ-4A genes has changed after the translocation. The change in recombination rate, however, seems to go in the opposite direction as GC and GC* decrease in the neoZ-4A region, suggesting a decrease in recombination rate and, therefore, a decrease in the efficacy of natural selection.

Finally, base composition has changed after the translocation to the Z chromosome. We indeed found evidence for lower GC content and GC* in the neoZ-4A region than in the remaining autosomal region of the 4A chromosome. In birds, as in many other organisms, chromosome size and recombination rate are negatively correlated (Backström et al. 2010), probably because one recombination event occurs per

chromosome arm per meiosis. Given that GC content
highly correlates with chromosome size (Eyre-Walker,
1993; Pessia et al. 2012), the observed difference in GC
content is a likely consequence of changes in the
intensity of the recombination-associated effect of
gBGC (Duret & Arndt, 2008; Duret & Galtier, 2009),
resulting from the instantaneous changes in
chromosome sizes due to the translocation. From the
translocated region of the ancestral 4A chromosome
point of view, the chromosomal context has drastically
changed from an ancestral ~21 Mb 4A chromosome to
a ~90 Mb neo-Z chromosome probably with a lower
recombination rate. Similarly, from the remaining 4A
chromosome point of view, the chromosomal context
has drastically changed too, from a ~21Mb to a ~11Mb
chromosome. As a consequence, we can expect that
base compositions have evolved in the opposite
direction with an increase in GC. However, this
hypothesis must be qualified since we were unable to
find any change in GC* at autosomal 4A genes,
suggesting that the chromosome-scale recombination
rate might not have been impacted by the
translocation. This could be possible if the translocation
occurred close to the centromere (*i.e.* a nearly whole-
arm translocation). In this case, the overall
recombination rate is not expected to change.

834 CONVERGENT CHROMOSOMAL FOOTPRINTS
835 OF A NEOW DEGENERATION

Degeneration of the non-recombining
chromosome, *i.e.* the chromosome only present in the
heterogametic sex, has been explored in depth in a
variety of species (*e.g.* Bachtrog & Charlesworth, 2002;
Papadopulos et al. 2015). Long-term gradual
degeneration through the accumulation of deleterious
mutations, partial loss of adaptive potential and gene
losses is expected to start soon after species cease to
recombine due to a series of factors including Muller's
ratchet, linked selection and the Hill-Robertson effect
(Charlesworth & Charlesworth, 2000; Bachtrog, 2005;
Sun & Heitman 2012).

To investigate this degeneration, we have first
identified 218 scaffolds with a female-specific pattern
in read coverage, for a total of 7.1 Mb. Among these
scaffolds, we have assigned 174 scaffolds to the neoW-
4A region, because of a high level of homology with the
zebra finch 4A chromosome, thus representing a
neoW-4A of a total length of 5.48 Mb. The absence of
any neoW scaffold homologous to the 4A *T. guttata*
chromosome between positions 1,756,080 and
3,509,582 suggests a large chromosomal deletion. The
total length of the neoW-W region we have assigned is
low, only representing 1.23 Mb of sequence, which is
far from the 6.94 Mb sequence that Smeds and
collaborators (2015) identified in the collared flycatcher
genome or the 6.81 Mb sequence reported in the
reference Chicken genome (GRCg6a version,
International Chicken Genome Sequencing Consortium
2004; Warren et al. 2017). It is however important to
specify that our objective was not to be exhaustive, but
rather to focus on the longest scaffolds for which both
estimates of the median coverage and alignments
against the zebra finch chromosome 4A were
considered reliable enough to be confident with its
assignation to the W chromosome, particularly in a
context of the intense activity of transposable
elements. Given the known difficulty to sequence and
assemble the W chromosome (*e.g.* Tomaszkiewicz et al.
2017), such a reduced-representation of the W
chromosome was reasonably expected. Our intent was
to get sufficient information to study the molecular
evolution of neoW-W specific genes too. Obtaining a
full sequence of the neoW chromosome for the *Z.*
borbonicus species, while possible, would require a
considerable additional sequencing effort to be
achieved.

Then, by aligning *Z. borbonicus* assembly
against the zebra finch 4A chromosome, we found
support for a candidate scaffold supporting the
chromosomal breakpoint. Alignments on both ends of
this scaffold suggest a potential chromosomal
breakpoint occurring around positions 9.603-9.605 Mb
of the v.3.2.4 zebra finch 4A chromosome, which is
remarkably close to our estimate for the translocation
on the neoZ-4A. Importantly, such an observation

therefore supports an evolution of neoW-4A and neoZ-
4A regions from initially identical gene sets.
Interestingly, we have also identified a large
chromosomal deletion on the W chromosome, which
represents another expected early signature of the W
degeneration (Charlesworth, 1991).

We have found support for a highly reduced
level of nucleotide diversity in the neoW chromosome
as compared to autosomes. This also holds true for the
neoW-4A region (mean neoW-4A:autosomal nucleotide
diversity = 0.36), which is in broad agreement with the
hypothesis of a three-fourth reduction in effective
population size associated to an autosomal-to-W or
autosomal-to-Y translocation. Our overall result of low
within-species diversity on the W chromosome is
however nothing comparable with the dramatically
reduced diversity observed by Smeds and collaborators
(2015) on the W chromosome of several flycatcher
species, with a W:autosomal diversity ranging from
0.96% to 2.16% depending on populations and species.
Non-recombinant W chromosome of *Z. borbonicus* also
exhibits elevated π_N/π_S in the neoW-W (0.78) as well as
in the neoW-4A regions (0.42), representing 3.68-fold
and 1.97-fold higher ratios than for the autosomal
genes, respectively (4.59-fold and 2.46-fold higher
ratios when compared with autosomal-4A genes only).
Higher d_N/d_S at neoW-4A genes as compared to their
neoZ-4A gametologs were also observed. This higher
d_N/d_S ratio is in agreement with the pattern observed in
another Sylvioidea species, the common whitethroat
(Sigeman et al. 2018) and, more broadly, with other
young sex chromosome systems (*e.g.* Marais et al.
2008). Sigeman et al. (2018) also reported an
association between amino acid and gene expression
divergences for the neoW-4A. All together, these
results are consistent with an accumulation of
deleterious mutations associated with the strong
reduction of the net efficacy of natural selection to
purge deleterious mutations (Charlesworth &
Charlesworth, 1997).

TE accumulation on W or Y chromosome is
suspected to play a particularly important role in the
first phases of the evolution of chromosome

differentiation (Bachtrog, 2003). To investigate this, we
have *de novo* identified *Z. borbonicus*-specific TE and
have analyzed distribution and abundance of TE. This
has led to the identification of a high TE load in the
ancestral W chromosome (45.1%), which is the same
order of magnitude as the reported value for the W
chromosome of *Ficedula albicollis* (48.5%, Smeds et al.
2015). Interestingly, we found support for a high TE
load in the translocated region too (21.6%), which is
approximately twice the observed TE content of any
other autosomal chromosome, including the autosomal
4A chromosome. TE classification, albeit incomplete,
supports an important contribution of class I LTR
elements to the overall TE load. LTR elements seem to
be particularly active in the zebra finch too (Kapusta
and Suh, 2017) suggesting that the recent burst of LTR
elements on autosomes may have facilitated the
accumulation of TEs on the neoW-4A chromosomes.
Although much more data and work will be needed in
the future to analyze in greater depth this
accumulation of TEs, and particularly the determinants
of this accumulation, our results suggest that LTR
elements may have virtually played a major role in
altering gene content, expression and/or chromosome
organization of the newly translocated region of the
Sylvioidea neo-W chromosome.

962 CONCLUSION

In this study, we generated a high quality
reference genome for *Z. borbonicus* that has provided
us unique opportunities to investigate the molecular
evolution of neo-sex chromosomes, and more broadly
to improve our understanding of avian sex
chromosome evolution. Since this species belong to the
Sylvioidea, one of the three major clades of passerines,
comprising close to 1,300 species, we can reasonably
suspect that this chromosomal-scale assembly will
serve as a reference for a large diversity of genome-
wide analyses in the Sylvioidea lineage itself, and more
generally in passerine birds.

Through detailed analyses of the evolution of the newly sex chromosome-associated regions, we found evolutionary patterns that were largely consistent with the classic expectations for the evolution of translocated regions on sex chromosomes, including evidence for reduction of diversity, ongoing neoW chromosome degeneration and base composition changes. A notable exception was the neoZ region for which no fast-neoZ effect was identified. Although most of the analyses are congruent, our report is based on a limited number of individuals and from only one population of *Z. borbonicus*. Further investigations based on a complementary and extensive dataset will probably help to fine-tune the conclusions, especially regarding the lack of any fast-Z effect or the drastic increase of TE on the neoW-4A. Lastly, given the huge difference in diversity between autosomes and sex chromosomes, we emphasize the importance of taking into account sex chromosomes for local adaptation studies, at least by scanning autosomes and sex chromosomes separately (see Bourgeois et al. 2018 for an example).

MATERIALS & METHODS

DNA AND RNA EXTRACTION, SEQUENCING

We extracted DNA from fresh tissues collected on a *Z. borbonicus* female individual (field code: 15-179), which died accidentally during fieldwork in May 2015 at Pas de Bellecombe (Gîte du volcan, Réunion; coordinates: S: -21.2174, E: 55.6872; elevation: 2246m above sea level). Sampling was conducted under a research permit (#602) issued to Christophe Thébaud by the Centre de Recherches sur la Biologie des Populations d'Oiseaux (CRBPO) – Muséum National d'Histoire Naturelle (Paris).

We also extracted DNA of a *Zosterops pallidus* individual collected in February 2015 in South Africa, Free state province, Sandymount Park, 10 kms from Fauresmith (coordinates: S: -29.75508, E: 25.17733). The voucher is stored at the Museum National d'Histoire Naturelle (MNHN), Paris, France, under the code BN72.

For both samples, 9µg of total genomic DNA were extracted from liver and/or muscle, using DNeasy Blood and Tissue kit (QIAGEN) following the manufacturer instructions. Each of these samples was sequenced as followed: one paired-end library with insert sizes of 300bp and 3 mate-pair libraries (3kb, 5kb and 8kb) using Nextera kit. Libraries and sequencing were performed by platform "INRA plateformes Génomes et transcriptomes (GeT-PlaGe)", Toulouse, France. Illumina sequencing was also performed at the platform GeT-PlaGe using Illumina HiSeq 3000 technology.

To improve genome assembly of Z. *borbonicus*, an additional sequencing effort was made by generating 10X coverage of PacBio long reads data. 20µg of high molecular weight DNA were extracted from muscle using MagAttract HMW DNA kit (QIAGEN) following manufacturer instructions. The PacBio librairies and sequencing were performed at Genome Québec (Centre d'innovation Génome Québec et Université McGill, Montréal, Quebec, Canada).

Total RNA of *Z. borbonicus* individual was extracted from brain and kidney tissues using RNeasy Plus Mini Kit (Qiagen) following manufacturer's instructions and then stabilized using RNAlater (Sigma). Both the RNAseq library preparation and the Hiseq2500 sequencing (1 line) were performed at the genomic platform MGX-Montpellier GenomiX from an equimolar pool of two different RNA extracts

GENOME ASSEMBLY

The paired-end reads were filtered using Trimmomatic (v0-33; Bolger et al., 2014) using the following parameters: "ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:50". The mate-pair reads were cleaned using NextClip (v1.3.1; Leggett et

al. 2013) using the following parameters : "--min_length 20 --trim_ends 0 --remove_duplicates".

Paired-end and mate-pair reads were assembled using SOAPdenovo (v2.04; Luo et al 2012) with parameters "-d 1 -D 2". Several k-mers (from 27 to 37-mers) were tested and we chose the assembly maximizing the N50 scaffold length criteria. Next, we applied Gapcloser v1.10 (a companion program of SOAPdenovo) to fill the gap in the assemblies.

Given that the PacBio technology produces long reads but with quite high sequencing error rates, we used LoRDEC (v0.6; Salmela & Rivals 2014) to correct the PacBio reads of the *Z. borbonicus* individual, using the following parameters: "-k 19 -s 3". In brief, LoRDEC corrects PacBio reads (both insert/deletions and base call errors) by the use of Illumina paired-end reads, a technology producing short reads only, but with a much higher base call accuracy and depth of coverage. The corrected PacBio reads were then used to scaffolds the SOAPdenovo assembly using SSPACE-LongRead (v1.1; Boetzer & Pirovano 2014). For *Z. borbonicus*, we also used MaSuRCA (v3.2.4; Zimin et al. 2017) to perform a hybrid assembly with a mixture of short and long reads. MaSuRCA produced an assembly with similar quality but slightly shorter than SOAPdenovo+SSPACE-LongRead (Table S1).

Several statistics were computed using assemblathon_stats.pl script (Bradnam et al. 2013; https://github.com/ucdavis-bioinformatics/assemblathon2-analysis) to evaluate the different genome assemblies. These statistics include genome size, number of scaffolds, N50, N90 and the proportion of missing data (N%). Additionally, we used BUSCO (v3.0.2; Waterhouse et al. 2017), a commonly used tool for evaluating the genome completeness based on the content in highly conserved orthologous genes (options: "-m genome -e 0.001 -l aves_odb9 -sp chicken").

The mitochondrial genome was assembled using mitobim (v1.8; Hahn et al. 2013) with the mitochondrial genome of *Z. lateralis* (accession : KC545407) used as reference (so-called "bait"). The mitochondrial genome was automatically annotated using the web server mitos2 (http://mitos2.bioinf.uni-leipzig.de; Bernt et al. 2013). This annotation was manually inspected and corrected using alignment with the other *Zosterops* mitochondrial genomes available in genbank.

GENOME-GUIDED DE NOVO TRANSCRIPTOME ASSEMBLY.

RNA-seq reads were used to reconstruct a transcriptome. First, RNA-seq reads were filtered with trimmomatic (v0-33; Bolger et al., 2014) using the following parameters : "ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 SLIDINGWINDOW:4:5 LEADING:5 TRAILING:5 MINLEN:25". Second, the filtered RNA-seq reads were mapped onto the reference genome using HISAT2 (Kim et al. 2015). HISAT2 performed a splice alignment of RNA-Seq reads, outperforming the spliced aligner algorithm implemented in TopHat (Kim et al. 2015). Finally, two methods were used to assemble the transcripts from the HISAT2 output bam file: i) Cufflinks (Trapnell et al. 2010) using the following parameters: "-q -p 10 -m 300 -s 100" and ii) Trinity (v2.5.0; Haas et al. 2013) using the following parameters : "--genome_guided_max_intron 100000".

Protein-coding Gene Annotation

Gene annotation was performed using the PASA pipeline combined with EVidenceModeler (Haas et al. 2008; Haas et al. 2011; https://github.com/PASApipeline/PASApipeline/wiki). The complete annotation pipeline involved the following steps:

1. *ab initio* gene finding with Augustus (Stanke and Waack 2003 ; http://bioinf.uni-greifswald.de/augustus/) using the parameter : "--species=chicken".
2. Protein homology detection and intron resolution using genBlastG (She et al. 2011;

http://genome.sfu.ca/genblast/download.html).
Protein sequences of several passerines were used as
references, namely zebra finch (*Taeniopygia guttata*;
assembly taeGut3.2.4; Warren et al. 2010), collared
flycatcher (*Ficedula albicollis*; assembly FicAlb_1.4;
Ellegren et al. 2012), medium ground-finch (*Geospiza*
fortis; assembly GeoFor_1.0; (Zhang et al. 2012) and
hooded crow (*Corvus cornix*; accession number
JPSR00000000.1; Poelstra et al. 2014). Genblast
parameters were : "-p genblastg -c 0.8 -r 3.0 -gff -pro -
cdna -e 1e-10".
3. PASA alignment assemblies based on
overlapping transcript from Trinity genome-guided *de*
novo transcriptome assembly (see above) (Haas et al.
2003). This step involved the so-called PASAPipeline
(v2.2.0;
https://github.com/PASApipeline/PASApipeline/) used
with the following parameters : "-C -r -R --ALIGNERS
blat,gmap".
4. Next, EVidenceModeler (v1.1.1; Haas et al.
2008) was used to compute weighted consensus gene
structure annotations based on the previous steps (1 to
3). We used the following parameters : "--segmentSize
500000 --overlapSize 200000" and an arbitrary weight
file following the guidelines provided at
http://evidencemodeler.github.io/ .
5. Finally, we used the script
"pasa_gff3_validator.pl" of PASA add UTR annotations
and models for alternatively spliced isoforms.
1165

1166 Finally, StringTie (Pertea et al. 2015) was also used to
1167 estimate the proportion of annotated CDS with RNA-
1168 seq information support.

1169

ORTHOLOGY DETECTION

We used the available passerine genomes
(namely, zebra finch, collared flycatcher, white-
throated sparrow and hooded crow) plus the high-
coverage genomes (>100x) of Zhang et al. (2014) (Table
S2). Orthology detection was performed using
OrthoFinder (v2.2.6; Emms & Kelly 2015). Single copy
(one-to-one) orthologs were extracted from
OrthoFinder results to perform multi-species alignment
with TranslatorX (v1.1; Abascal et al. 2010) using

MAFFT (Katoh et al. 2002) to build the alignment.
Alignments were inspected by HMMCleaner (v1.8;
Amemiya et al. 2013; Philippe et al. 2017) to exclude
badly aligned sites. Next, dubious, highly divergent,
sequences were excluded using trimAl (Capella-
Gutiérrez et al. 2009) option "-resoverlap 0.60 -
seqoverlap 80". We also used genome assemblies of 3
passerine species for which no gene annotation sets
were publicly available, namely the silvereye, *Z.*
lateralis (Cornetti et al. 2015), the Orange River white-
eye *Z. pallidus* (this study) and the willow warbler
Phylloscopus trochilus (Lundberg et al. 2017). For these
genomes, gene orthology detection was conducted
using AGILE (Hughes & Teeling 2018). AGILE is a
pipeline for gene mining in fragmented assembly
overcoming the difficulty that genes could be located in
several scaffolds. We applied AGILE using *Z. borbonicus*
single copy orthologs as query genes.

1198

FILTERING SCAFFOLDS ORIGINATING FROM AUTOSOMES, W AND Z CHROMOSOMES

Given that we have sequenced a female
genome, we likely assembled contigs from W, Z and
autosomal chromosomes. To identify scaffolds
originating from sex chromosomes or autosomes, we
mapped 7 females and 5 males reads from *Z.*
borbonicus birds published in Bourgeois et al. (2017)
onto our female genome assembly. We first mapped all
raw reads against the *Z. borbonicus* reference genome
using BWA mem v. 0.7.5a (Li, 2013), we then removed
duplicates with Picard 1.112
(http://broadinstitute.github.io/picard). We then used
Mosdepth (Pedersen & Quinlan, 2018) to compute
median per-site coverage for each scaffold, following
the same strategy than in Smeds et al. (2015). For each
i scaffold, total coverage for males and females were
computed as the sum of coverage of all individuals.
Then, we compute a normalized coverage per scaffolds
for male as :

Normalized(Cov. Scaffold $_i$(Males)) =

*Cov. Scaffold $_i$(Males) * $\frac{MeanCov.(Females)}{MeanCov.(Males)}$*

Where "*Mean Cov. (Females)*" and "*Mean Cov.*
(Males)" corresponds to the median coverages for
males and females across all scaffolds longer than 1Mb.

This normalization is intended to take into account the
different number of male and female individuals.

Next, we used the normalized median per-site
coverage to detect W-linked scaffolds (zero and above
5X in males and females, respectively), Z-linked (female
with less than 0.75 the male coverage and male with >
9X coverage) and autosomes (all the remaining
scaffolds corresponding female and male with a
roughly similar median coverage). To decrease the
probability of false identification due to the mapping in
repeat-rich regions, this approach has only been
performed using coverage data from scaffolds longer
than 10 kb (3,443 / 97,537 scaffolds, >96% of the
assembly size).

1238

PSEUDO-CHROMOSOME ASSEMBLY
We first aligned soft-masked *Z. borbonicus*
scaffolds on the soft-masked Zebra Finch genome
(*Taeniopygia guttata*) using LastZ v. 1.04.00 (Schwartz
et al. 2003). For *Z. borbonicus*, *de novo* transposable
elements prediction and genome assembly soft-
masking have been performed using Repeat Masker v.
4.0.3 (Smit et al. 2013-2015). For *T. guttata*, we used
the soft-masked genome v. 3.2.4 (taeGut3.2.4) made
available by the Zebra Finch genome consortium. This
soft-masking procedure was put in place in order to
exclude these regions for the LastZ's seeding stage, and
thereby to avoid finding alignments in these highly
repeated regions. We then filtered LastZ alignments
hits in order to only keep reliable hits (5% longest hits
and with sequence identities greater than the median
over all alignments). For each scaffold, we then defined
syntenic blocks as adjacencies of several reliable hits. A
syntenic block represents a homologous region
between zebra finch and *Z. borbonicus* starting at the
first reliable hit and ending at the last one. Syntenic
blocks covering at least 80% of a *Z. borbonicus* scaffold
size and 80-120% the corresponding homologous
region in the Zebra Finch genome were automatically
anchored to its *T. guttata* chromosome position,
assuming complete synteny between the zebra finch
and *Z. borbonicus*. Unanchored scaffolds were then

manually identified by visually inspecting the summary
statistics and positions of all raw alignments. In case of
chimeric scaffolds, these scaffolds were cut into two or
several new scaffolds assuming that this chimerism is
due to a contigging or a scaffolding artifact.

Second, we used DeCoSTAR, a computational
method tracking gene order evolution from
phylogenetic signal by inferring gene adjacencies
evolutionary histories, in order to not only improve the
genome assembly of *Z. borbonicus*, but also 26
additional extant avian assemblies including the high
coverage (>100X) of Zhang et al. 2014 and the other
passerine genomes (Table S2). The gene trees were
obtained for phylogenies of single-copy orthologs (see
above) estimated using IQ-TREE model GTR+G4 and
1000 so-called "ultra-fast bootstrap" (v1.6.2; Nguyen et
al. 2014). A snakemake pipeline
(https://github.com/YoannAnselmetti/DeCoSTAR_pipel
ine) were used to apply DeCoSTAR on gene orders and
phylogenies.

Finally, scaffold orders given by LASTZ and
DeCoSTAR were manually reconciled to determine the
consensus scaffold orders along the *Z.borbonicus*
chromosomes.

1290

MOLECULAR DATING
We performed two independent molecular
dating analyses. First, we used relaxed molecular clock
analysis based on nuclear sequences and fossil
calibrations applied on the full bird phylogeny. Second,
we applied the approach of Nabholz et al. (2016) using
white-eyes body mass to estimate their mitochondrial
substitution rate.

In the first approach, molecular dating
analyses were performed using the 27 species selected
for the DeCoSTAR analysis plus the silvereye, the
Orange River white-eye and the willow warbler leading
to a total of 30 species. We restricted our analyses to
single-copy orthologs with low GC content. These
genes are known to evolve slowing and more clock-like
than GC rich genes (Nabholz et al. 2011; Jarvis et al.

2014). To do that, we randomly selected 100 single-
copy orthologs coded by chromosome 1 and 2
excluding genes at the beginning and at the end of the
chromosomes (minimum distance from each
chromosome tip: 10 Mbp). This step was replicated ten
times to evaluate the robustness of the inferences
among different sets of genes.

We used several fossil calibrations sets
summarized in Table S3. Our different calibration sets
reflect the current uncertainty surrounding bird
diversification dates. We used a conservative set with
only one maximum bound at 140 Myrs for the origin of
Neornithes (set 4). At the other end, we used another
set with a narrow constraint between 28 and 34 Myrs
for the Suboscines / Oscines divergence (set 3). This
constraint could turn out to be incorrect with future
advances in the bird fossil records. For all the
calibration sets, the Neognathae / Palaeognathae
divergence minimal age was set to 66 Myrs using
Vegavis iaai fossil (Benton et al. 2009; Mayr, 2013;
Ksepka & Clarke, 2015). The maximum age of this node
is much more difficult to select. We have opted for two
maximal ages. The first one at 86.5 Myrs following the
rationale of Prum et al. (2015) based of the upper
bound age estimate of the Niobrara Formation (set 1, 2
and 3). We used another very conservative maximum
bound at 140 Myrs (set 4). This is the maximum age
estimated for the origin of Neornithes by Lee et al.
(2014) using an extensive morphological clock analysis.
Additionally, in sets 1 and 3, we constrained the
divergence between Passeriformes / Psittaciformes to
be between 53.5 and 65.5 Myrs, assuming that the
complete absence of passerine bird species during the
Cretaceous. In sets 2 and 4, we only used a minimum
bound on the first fossil occurrence of passerines in the
Eocene (Boles, 1997) and on the stem Psittaciformes
fossil *Pulchrapollia gracilis* (Dyke & Cooper, 2000). In
calibration set 3, we constrained the
Oscines/Suboscines split between 28 and 34 Myrs,
following the rationale of Mayr (2013) assuming crown
Oscines and Suboscines originated in the early
Oligocene (28 Myrs, Mayr & Manegold 2006) and
based on the absence of Eocene fossil records
discovered so far (Eocene/Oligocene limit is at 34

Myrs). All the other minimum-bound calibrations we
used followed the suggestions of Ksepka & Clarke
(2015) and are presented in Table S3.

Molecular dating was performed using
Phylobayes (v4.1; Lartillot et al. 2009) using a CAT-GTR
substitution model. For the relaxed clock model, we
used the log-nomal auto-correlated rates (ln) model.
We also tested the uncorrelated gamma multipliers
(ugam) model that gave similar results than the ln
model (results not shown). We used uniform prior on
divergence times, combined with soft calibrations
(Rannala and Yang, 2007; Yang and Rannala, 2006). The
MCMC were run for at least 8,000 cycles. MCMC
convergence were diagnosed by running two
independents MCMC and by visually checking the
evolution of the likelihood and other parameters along
the Markov chain (in ".trace" files).

1368

Additionally, we applied an independent
molecular dating using the method proposed by
Nabholz et al. (2016). Seven complete mitochondrial
Zosterops genomes were downloaded from Genbank,
including *Z. erythropleuros* (KT194322), Z. japonicus
(KT601061), *Z. lateralis* (KC545407), *Z. poliogastrus*
(KX181886), *Z. senegalensis* (KX181887), *Z. senegalensis*
(KX181888). We also included the sequences of the
Réunion grey white-eye and the Orange River white-
eye assembled in the present study. The mitochondrial
sequences of *Yuhina diademata* (KT783535) and
Zoothera dauma (KT340629) were used as outgroups.
Body mass for all *Zosterops* species were obtained from
Dunning (2007) and the median of these body masses
was computed. Phylogenetic relationship and branch
length were estimated using IQ-TREE with a HKY+G4
substitution model using third codon positions only.
Then, we applied the formula of Nabholz et al. 2016 to
derive the substitution rate (substitution per site per
Myr) as $\frac{10^{-0.141*log10(BodyMass)+0.367}}{100}$ and
$\frac{10^{-0.243*log10(BodyMass)+0.905}}{100}$ for the minimal and maximal
rate where "*Body Mass*" is the median body mass in
grams (logarithm of base 10). Next, we computed the

median divergence time between the silver-eye (*Z. lateralis)* and all the other white-eye species to obtain an estimate of the crown *Zosterops* clade age. Finally, we divided this divergence time by the rate obtained with the formula above to obtain divergence dates in Myrs.

GENOME-WIDE ESTIMATES OF NUCLEOTIDE DIVERSITY

Based on the previously generated BAM files (see '*Filtering scaffolds originating from autosomes, W and Z chromosomes*' section*)*, we then use GATK to generate the gvcf from the 6 parents of the 3 progenies described in Bourgeois et al. 2017 (3 males and 3 females). Joint genotyping, as well as all subsequent analyses, was then performed separately for the 3 males and the 3 females. We followed all the GATK best practices under the GATK suite, except for variant filtration which was performed using a custom script to speed up computations. This step however followed the same procedures than under GATK, assuming the following thresholds: QD>2.0, FS<60.0, MQ>40.0, MQRANKSUM>-2.0, READPOSRANKSUM>-2.0 and RAW_MQ > 45000.

For each individual, we then reconstructed two genomic sequences. At each position of the genome, the position (reference or alternate if any) was added to the sequence if the coverage at the position was between 3 and 50. In any other case, a "N" character was added in order to keep the sequence length the same. We then computed the Tajima's π estimator of nucleotides diversity and the GC content over non-overlapping 10 kb windows. To be highly conservative in the analysis of the neoW chromosome, all windows associated to W scaffolds and found covered in the male (ZZ) dataset were excluded from the analysis of the female (ZW) dataset, even if this non-zero coverage was observed at a single base over the genomic window.

ESTIMATES OF NON-SYNONYMOUS AND SYNONYMOUS DIVERGENCES AND GC EQUILIBRIUM

For the neoZ-4A and neoW-4A, we made a specific effort to retrieve the paralogous sequences. Assuming that the two translocated regions evolved from the same gene sets (see results), each gene located on the neoZ-4A are expected to have a paralog on the neoW-4A (so called "gametologs", Pala et al. 2012a,b). As a consequence, most gametologs have been eliminated during our selection of the single-copy orthologs. We visually inspected all the alignments containing a neoZ-4A gene and then tried to identify the corresponding copy in a scaffold assigned to the neoW chromosome.

Given the drastically different number of genes between autosomes, neoZ-Z and neoZ-4A regions, we subsampled the data to match the category with the lowest number of genes (i.e., the neoZ-4A region). Then, we concatenated genes and computed d_N and d_S as the sum of non-synonymous and synonymous branch lengths respectively. This will decrease the variance of the estimated d_N/d_S ratios and limit problems associated to low d_S values (Wolf et al. 2009). Finally, the variability in d_N/d_S was evaluated by bootstrapping genes 1000 times within each genomic region. For GC equilibrium (GC*), we used the nonhomogeneous model T92 (Galtier and Gouy 1998) implemented in the BPPSUITE package (Dutheil and Boussau 2008, http://biopp.univ-montp2.fr/wiki/index.php/BppSuite) based on the Bio++ library (Guéguen et al. 2013) to infer GC* at third codon position for each branch. We used a similar bootstrapping strategy (1000 times within each genomic region) to evaluate the variation in GC*.

For the comparison between neoZ-4A and neoW-4A, we used the d_N/d_S and GC* of neoZ-4A and neoW-4A genes for the branch leading to the Reunion grey white-eye. When the neoW-4A sequences of the Reunion grey white-eye was very closely related to a sequence of the Orange River white-eye, we assumed that the Orange River white-eye sequence also came from the neoW-4A regions and we computed the d_N/d_S

of the ancestral branch of these two species. We evaluated the different in d_N/d_S using a paired-samples Wilcoxon test (also known as Wilcoxon signed-rank test).

GC and TE contents

We used the automated approach implemented in RepeatModeler Open (v.1.0.11, Smit & Hubley, 2008) to *de novo* detect *Z.borbonicus*-specific TE consensus. The generated list of *de novo* TE sequences was merged to the chicken repetitive sequences publicly available in Repbase (Jurka et al. 2005). We then used this set of sequences as a custom library for RepeatMasker (v.open-4.0.3, Smit et al. 2013) to generate a softmasked version of the *Z. borbonicus* genome assembly. Then, we used a non-overlapping 10-kbp sliding windows approach to calculate the GC and TE contents along the whole genome. All genomic windows composed of more than 50% of Ns were excluded to ensure accurate estimates of local TE and GC contents.

All statistical analyses were performed using R v. 3.4.4 (R core Team, 2018). Some analyses and graphics were performed using several additional R packages: APE (Paradis & Strimmer, 2004), beanplot (Kampstra, 2008), circlize (Gu et al. 2014), cowplot (Wilke, 2016), ggplot2 (Wickham, 2016), phytools (Revell, 2012) and plotrix (Lemon, 2006).

Data availability

Nuclear and mitochondrial genome sequences, scripts and programs used are available at the following FigShare repository: https://figshare.com/s/122efbec2e3632188674. Raw reads and genome sequence are available on SRA and GenBank (XXXXX and XXXXX).

Acknowledgments

This research was funded by the French ANR (BirdIslandGenomic project, ANR-14-CE02-0002). The analyses benefited from the Montpellier Bioinformatics Biodiversity (MBB) platform services and the genotoul bioinformatics platform Toulouse Midi-Pyrenees (Bioinfo Genotoul). We are grateful to Jérôme Fuchs for providing the mitochondrial alignments, to Fabien Condamine for constructive discussions on molecular dating and Anna-Sophie Fiston-Lavier for providing advice on the analysis of transposable elements. We thank the Reunion National Park for granting us permission to conduct fieldwork in Pas de Bellecombe, Reunion. We are grateful for the logistic support provided by the field station of Marelongue, funded by the P.O.E., Reunion National Park and OSU Reunion.

References

Abascal F, Zardoya R, Telford MJ. **2010**. TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Research* **38**: W7–W13.

Amemiya CT, Alföldi J, Lee AP, Fan S, Philippe H, MacCallum I, Braasch I, Manousaki T, Schneider I, Rohner N, *et al.* 2013. The African coelacanth genome provides insights into tetrapod evolution. *Nature* **496**: 311.

Anselmetti Y, Duchemin W, Tannier E, Chauve C, Bérard S. **2018**. Phylogenetic signal from rearrangements in 18 Anopheles species by joint scaffolding extant and ancestral genomes. *BMC Genomics* **19**: 96.

Bachtrog D. **2005**. Sex chromosome evolution: molecular aspects of Y-chromosome degeneration in Drosophila. *Genome research* **15**: 1393–1401.

Bachtrog D, Charlesworth B. **2002**. Reduced adaptation of a non-recombining neo-Y chromosome. *Nature* **416**: 323.

Backström N, Forstmeier W, Schielzeth H, Mellenius H,
Nam K, Bolund E, Webster MT, Ost T, Schneider M,
Kempenaers B, et al. 2010. The recombination
landscape of the zebra finch Taeniopygia guttata
genome. *Genome research* **20**: 485–495.
Benton M, Donoghue P, Asher R. 2009. Calibrating and
constraining molecular clocks. In: The timetree of Life.
Oxford University Press, 35–86.
Bernt M, Donath A, Jühling F, Externbrink F, Florentz
C, Fritzsch G, Pütz J, Middendorf M, Stadler PF. 2013.
MITOS: Improved de novo metazoan mitochondrial
genome annotation. *Mitogenomics and Metazoan*
Evolution **69**: 313–319.
Boetzer M, Pirovano W. 2014. SSPACE-LongRead:
scaffolding bacterial draft genomes using long read
sequence information. *BMC Bioinformatics* **15**: 211.
Boles W. 1997. *Fossil Songbirds (Passeriformes) from*
the Early Eocene of Australia.
Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a
flexible trimmer for Illumina sequence data.
Bioinformatics **30**: 2114–2120.
Botero-Castro F, Figuet E, Tilak M-K, Nabholz B, Galtier
N. 2017. Avian Genomes Revisited: Hidden Genes
Uncovered and the Rates versus Traits Paradox in Birds.
Molecular Biology and Evolution **34**: 3123–3131.
Bourgeois YXC, Delahaie B, Gautier M, Lhuillier E,
Malé P-JG, Bertrand JAM, Cornuault J, Wakamatsu K,
Bouchez O, Mould C, et al. 2017. A novel locus on
chromosome 1 underlies the evolution of a melanic
plumage polymorphism in a wild songbird. *Royal*
Society open science **4**: 160805–160805.
Bourgeois Y, Ruggiero R, Manthey J, Boissinot S. 2018.
Recent secondary contacts, background selection and
variable recombination rates shape genomic diversity
in the model species Anolis carolinensis. *bioRxiv*.
Bracewell RR, Bentz BJ, Sullivan BT, Good JM. 2017.
Rapid neo-sex chromosome evolution and incipient
speciation in a major forest pest. *Nature*
Communications **8**: 1593

Bradnam KR, Fass JN, Alexandrov A, Baranay P,
Bechner M, Birol I, Boisvert S, Chapman JA, Chapuis G,
Chikhi R, et al. 2013. Assemblathon 2: evaluating de
novo methods of genome assembly in three vertebrate
species. *GigaScience* **2**: 2047-217X-2–10.
Brooke M de L, Welbergen JA, Mainwaring MC, van
der Velde M, Harts AMF, Komdeur J, Amos W. 2010.
Widespread Translocation from Autosomes to Sex
Chromosomes Preserves Genetic Variability in an
Endangered Lark. *Journal of Molecular Evolution* **70**:
242–246.
Cai T, Cibois A, Alström P, Moyle RG, Kennedy JD,
Shao S, Zhang R, Irestedt M, Ericson PGP, Gelang M, et
al. 2019. Near-complete phylogeny and taxonomic
revision of the world's babblers (Aves: Passeriformes).
Molecular Phylogenetics and Evolution **130**: 346–356.
Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T.
2009. trimAl: a tool for automated alignment trimming
in large-scale phylogenetic analyses. *Bioinformatics* **25**:
1972–1973.
Charlesworth B. 1991. The evolution of sex
chromosomes. *Science* **251**: 1030.
Charlesworth B, Charlesworth D. 1997. Rapid fixation
of deleterious alleles can be caused by Muller's ratchet.
Genetics Research **70**: 63–73.
Charlesworth B, Charlesworth D. 2000. The
degeneration of Y chromosomes. *Philosophical*
transactions of the Royal Society of London. Series B,
Biological sciences **355**: 1563–1572.
Charlesworth B, Sniegowski P, Stephan W. 1994. The
evolutionary dynamics of repetitive DNA in eukaryotes.
Nature **371**: 215.
Clegg SM, Phillimore AB. 2010. The influence of gene
flow and drift on genetic and phenotypic divergence in
two species of Zosterops in Vanuatu. *Philosophical*
transactions of the Royal Society of London. Series B,
Biological sciences **365**: 1077–1092.
Coghlan A, Eichler EE, Oliver SG, Paterson AH, Stein L.
2005. Chromosome evolution in eukaryotes: a multi-

kingdom perspective. *Trends in Genetics* **21**: 673–682.

Cornetti L, Valente LM, Dunning LT, Quan X, Black RA,
Hébert O, Savolainen V. **2015**. The Genome of the
"Great Speciator" Provides Insights into Bird
Diversification. *Genome Biology and Evolution* **7**: 2680–
2691.

Cornuault J, Delahaie B, Bertrand JAM, Bourgeois YXC,
Milá B, Heeb P, Thébaud C. **2014**. Morphological and
plumage colour variation in the Réunion grey white-eye
(Aves: Zosterops borbonicus): assessing the role of
selection. *Biological Journal of the Linnean Society* **114**:
459–473.

Delahaie B, Cornuault J, Masson C, Bertrand JAM,
Bourgeois YXC, Milá B, Thébaud C. **2017**. Narrow
hybrid zones in spite of very low population
differentiation in neutral markers in an island bird
species complex. *Journal of Evolutionary Biology* **30**:
2132–2145.

Derryberry EP, Claramunt S, Derryberry G, Chesser RT,
Cracraft J, Aleixo A, Pérez-Emán J, Remsen J JV,
Brumfield RT. **2011**. Lineage diversification and
morphological evolution in a large-scale continental
radiation: the neotropical ovenbirds and woodcreepers
(Aves: Furnariidae). *Evolution* **65**: 2973–2986.

Diamond JM, Gilpin ME, Mayr E. **1976**. Species-
distance relation for birds of the Solomon Archipelago,
and the paradox of the great speciators. *Proceedings of*
the National Academy of Sciences of the United States
of America **73**: 2160–2164.

Drummond DA, Wilke CO. **2008**. Mistranslation-
Induced Protein Misfolding as a Dominant Constraint
on Coding-Sequence Evolution. *Cell* **134**: 341–352.

Duchemin W, Anselmetti Y, Patterson M, Ponty Y,
Bérard S, Chauve C, Scornavacca C, Daubin V, Tannier
E. **2017**. DeCoSTAR: Reconstructing the Ancestral
Organization of Genes or Genomes Using Reconciled
Phylogenies. *Genome Biology and Evolution* **9**: 1312–
1319.

Dunning J. **2007**. *CRC Handbook of Avian Body Masses*.

Boca Raton.

Duret L, Arndt PF. **2008**. The Impact of Recombination
on Nucleotide Substitutions in the Human Genome.
PLOS Genetics **4**: e1000071.

Duret L, Galtier N. **2009**. Biased Gene Conversion and
the Evolution of Mammalian Genomic Landscapes.
Annual Review of Genomics and Human Genetics **10**:
285–311.

Dutheil J, Boussau B. **2008**. Non-homogeneous models
of sequence evolution in the Bio++ suite of libraries and
programs. *BMC evolutionary biology* **8**: 255–255.

Dyke GJ, Cooper JH. **2008**. A new psittaciform bird
from the London Clay (Lower Eocene) of England.
Palaeontology **43**: 271–285.

Ellegren H. **2009**. The different levels of genetic
diversity in sex chromosomes and autosomes. *Trends in*
Genetics **25**: 278–284.

Ellegren H. **2010**. Evolutionary stasis: the stable
chromosomes of birds. *Trends in Ecology & Evolution*
25: 283–291.

Ellegren H. **2013**. The Evolutionary Genomics of Birds.
Annual Review of Ecology, Evolution, and Systematics
44: 239–259.

Ellegren H, Smeds L, Burri R, Olason PI, Backström N,
Kawakami T, Künstner A, Mäkinen H, Nadachowska-
Brzyska K, Qvarnström A, *et al.* 2012. The genomic
landscape of species divergence in Ficedula flycatchers.
Nature **491**: 756.

Emms DM, Kelly S. **2015**. OrthoFinder: solving
fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy.
Genome Biology **16**: 157.

Ericson PG, Klopfstein S, Irestedt M, Nguyen JM,
Nylander JA. **2014**. Dating the diversification of the
major lineages of Passeriformes (Aves). *BMC*
Evolutionary Biology **14**: 8.

Eyre-Walker A. **1993**. Recombination and Mammalian
Genome Evolution. *Proceedings: Biological Sciences*
252: 237–243.

Fraïsse C, Picard MAL, Vicoso B. **2017**. The deep
conservation of the Lepidoptera Z chromosome
suggests a non-canonical origin of the W. *Nature*
Communications **8**: 1486.
Galtier N, Gouy M. **1998**. Inferring pattern and process:
maximum-likelihood implementation of a
nonhomogeneous model of DNA sequence evolution
for phylogenetic analysis. *Molecular Biology and*
Evolution **15**: 871–879.
Genner MJ, Seehausen O, Lunt DH, Joyce DA, Shaw
PW, Carvalho GR, Turner GF. **2007**. Age of Cichlids:
New Dates for Ancient Lake Fish Radiations. *Molecular*
Biology and Evolution **24**: 1269–1282.
Gill FB. **1973**. Intra-Island Variation in the Mascarene
White-Eye Zosterops borbonica. *Ornithological*
Monographs: iii–66.
Griffin DK, Robertson LBW, Tempest HG, Skinner BM.
2007. The evolution of the avian genome as revealed
by comparative molecular cytogenetics. *Cytogenetic*
and Genome Research **117**: 64–77.
Gu Z, Gu L, Eils R, Schlesner M, Brors B. **2014**. circlize
implements and enhances circular visualization in R.
Bioinformatics **30**: 2811–2812.
Guéguen L, Gaillard S, Boussau B, Gouy M, Groussin
M, Rochette NC, Bigot T, Fournier D, Pouyet F, Cahais
V, *et al.* 2013. Bio++: Efficient Extensible Libraries and
Tools for Computational Molecular Evolution.
Molecular Biology and Evolution **30**: 1745–1750.
Haas BJ, Delcher AL, Mount SM, Wortman JR, Smith
RK Jr, Hannick LI, Maiti R, Ronning CM, Rusch DB,
Town CD, *et al.* 2003. Improving the Arabidopsis
genome annotation using maximal transcript alignment
assemblies. *Nucleic acids research* **31**: 5654–5666.
Haas BJ, Papanicolaou A, Yassour M, Grabherr M,
Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber
M, *et al.* 2013. De novo transcript sequence
reconstruction from RNA-seq using the Trinity platform
for reference generation and analysis. *Nature Protocols*
8: 1494.

Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis
J, White O, Buell CR, Wortman JR. **2008**. Automated
eukaryotic gene structure annotation using
EVidenceModeler and the Program to Assemble Spliced
Alignments. *Genome Biology* **9**: R7.
Haas BJ, Zeng Q, Pearson MD, Cuomo CA, Wortman
JR. **2011**. Approaches to Fungal Genome Annotation.
Mycology **2**: 118–141.
Hahn C, Bachmann L, Chevreux B. **2013**.
Reconstructing mitochondrial genomes directly from
genomic next-generation sequencing reads—a baiting
and iterative mapping approach. *Nucleic Acids Research*
41: e129–e129.
Harmon LJ. **2018**. *Phylogenetic Comparative Methods:*
Learning from Trees. CreateSpace Independent
Publishing Platform.
Heads M. **2005**. Dating nodes on molecular
phylogenies: a critique of molecular biogeography.
Cladistics **21**: 62–78.
Heads M. **2011**. Old Taxa on Young Islands: A Critique
of the Use of Island Age to Date Island-Endemic Clades
and Calibrate Phylogenies. *Systematic Biology* **60**: 204–
218.
Hill WG, Robertson A. **1966**. The effect of linkage on
limits to artificial selection. *Genetics Research* **8**: 269–
294.
Hughes GM, Teeling EC. **2018**. AGILE: an assembled
genome mining pipeline. *Bioinformatics*: bty781–
bty781.
Hvilsom C, Qian Y, Bataillon T, Li Y, Mailund T, Sallé B,
Carlsen F, Li R, Zheng H, Jiang T, *et al.* 2012. Extensive
X-linked adaptive evolution in central chimpanzees.
Proceedings of the National Academy of Sciences **109**:
2054.
International Chicken Genome Sequencing
Consortium, Hillier LW, Miller W, Birney E, Warren W,
Hardison RC, Ponting CP, Bork P, Burt DW, Groenen
MAM, *et al.* 2004. Sequence and comparative analysis
of the chicken genome provide unique perspectives on

vertebrate evolution. *Nature* **432**: 695.

Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, Ho SYW, Faircloth BC, Nabholz B, Howard JT, *et al.* 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science* **346**: 1320.

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. **2012**. The global diversity of birds in space and time. *Nature* **491**: 444.

Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. **2005**. Repbase Update, a database of eukaryotic repetitive elements. *Cytogenetic and Genome Research* **110**: 462–467.

Kampstra P. **2008**. Beanplot: A Boxplot Alternative for Visual Comparison of Distributions. *Journal of Statistical Software, Code Snippets* **28**: 1–9.

Kapusta A, Suh A. **2017**. Evolution of bird genomes—a transposon's-eye view. *Annals of the New York Academy of Sciences* **1389**: 164–185.

Katoh K, Misawa K, Kuma K, Miyata T. **2002**. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic acids research* **30**: 3059–3066.

Kawakami T, Smeds L, Backström N, Husby A, Qvarnström A, Mugal CF, Olason P, Ellegren H. **2014**. A high-density linkage map enables a second-generation collared flycatcher genome assembly and reveals the patterns of avian recombination rate variation and chromosomal evolution. *Molecular Ecology* **23**: 4035–4058.

Kim D, Langmead B, Salzberg SL. **2015**. HISAT: a fast spliced aligner with low memory requirements. *Nature Methods* **12**: 357.

Kitano J, Peichel CL. **2012**. Turnover of sex chromosomes and speciation in fishes. *Environmental Biology of Fishes* **94**: 549–558.

Kitano J, Ross JA, Mori S, Kume M, Jones FC, Chan YF, Absher DM, Grimwood J, Schmutz J, Myers RM, *et al.* 2009. A role for a neo-sex chromosome in stickleback speciation. *Nature* **461**: 1079.

Ksepka D, Clarke J. **2015**. *Phylogenetically vetted and stratigraphically constrained fossil calibrations within Aves*.

Lagomarsino LP, Condamine FL, Antonelli A, Mulch A, Davis CC. **2016**. The abiotic and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae). *New Phytologist* **210**: 1430–1442.

Lanfear R, Kokko H, Eyre-Walker A. **2014**. Population size and the rate of evolution. *Trends in Ecology & Evolution* **29**: 33–41.

Lartillot N, Lepage T, Blanquart S. **2009**. PhyloBayes 3: a Bayesian software package for phylogenetic reconstruction and molecular dating. *Bioinformatics* **25**: 2286–2288.

Lee MSY, Cau A, Naish D, Dyke GJ. **2014**. Morphological Clocks in Paleontology, and a Mid-Cretaceous Origin of Crown Aves. *Systematic Biology* **63**: 442–449.

Leggett RM, Clavijo BJ, Clissold L, Clark MD, Caccamo M. **2014**. NextClip: an analysis and read preparation tool for Nextera Long Mate Pair libraries. *Bioinformatics* **30**: 566–568.

Lemon J. **2006**. Plotrix: A package in the red light district of R. *R-News* **6**: 8–12.

Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*.

Lundberg M, Liedvogel M, Larson K, Sigeman H, Grahn M, Wright A, Åkesson S, Bensch S. **2017**. Genetic differences between willow warbler migratory phenotypes are few and cluster in large haplotype blocks. *Evolution Letters* **1**: 155–168.

Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, *et al.* 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *GigaScience* **1**: 2047-217X-1–18.

Magallon S, Sanderson MJ. **2001**. Absolute diversification rates in angiosperm clades. *Evolution* **55**: 1762–1780.

Mank JE, Axelsson E, Ellegren H. **2007**. Fast-X on the Z:

Rapid evolution of sex-linked genes in birds. *Genome Research* **17**: 618–624.

Mank JE, Nam K, Ellegren H. **2010**. Faster-Z Evolution Is Predominantly Due to Genetic Drift. *Molecular Biology and Evolution* **27**: 661–670.

Marais GAB, Nicolas M, Bergero R, Chambrier P, Kejnovsky E, Monéger F, Hobza R, Widmer A, Charlesworth D. **2008**. Evidence for Degeneration of the Y Chromosome in the Dioecious Plant Silene latifolia. *Current Biology* **18**: 545–549.

Mayr G. **2009**. *A small suboscine-like passeriform bird from the early Oligocene of France*.

Mayr G. **2013**. The age of the crown group of passerine birds and its evolutionary significance – molecular calibrations versus the fossil record. *Systematics and Biodiversity* **11**: 7–13.

Melo M, Warren BH, Jones PJ. **2011**. Rapid parallel evolution of aberrant traits in the diversification of the Gulf of Guinea white-eyes (Aves, Zosteropidae). *Molecular Ecology* **20**: 4953–4967.

Milá B, Warren BH, Heeb P, Thébaud C. **2010**. The geographic scale of diversification on islands: genetic and morphological divergence at a very small spatial scale in the Mascarene grey white-eye (Aves: Zosterops borbonicus). *BMC Evolutionary Biology* **10**: 158.

Moyle RG, Filardi CE, Smith CE, Diamond J. **2009**. Explosive Pleistocene diversification and hemispheric expansion of a "great speciator". *Proceedings of the National Academy of Sciences* **106**: 1863.

Murphy WJ, Sun S, Chen Z-Q, Pecon-Slattery J, O'Brien SJ. **1999**. Extensive Conservation of Sex Chromosome Organization Between Cat and Human Revealed by Parallel Radiation Hybrid Mapping. *Genome Research* **9**: 1223–1230.

Nabholz B, Künstner A, Wang R, Jarvis ED, Ellegren H. **2011**. Dynamic Evolution of Base Composition: Causes and Consequences in Avian Phylogenomics. *Molecular Biology and Evolution* **28**: 2197–2210.

Nabholz B, Lanfear R, Fuchs J. **2016**. Body mass-corrected molecular rate for bird mitochondrial DNA. *Molecular Ecology* **25**: 4438–4449.

Nam K, Munch K, Hobolth A, Dutheil JY, Veeramah KR, Woerner AE, Hammer MF, Mailund T, Schierup MH. **2015**. Extreme selective sweeps independently targeted the X chromosomes of the great apes. *Proceedings of the National Academy of Sciences* **112**: 6413.

Nanda I, Karl E, Volobouev V, Griffin DK, Schartl M, Schmid M. **2006**. Extensive gross genomic rearrangements between chicken and Old World vultures (Falconiformes: Accipitridae). *Cytogenetic and Genome Research* **112**: 286–295.

Nanda I, Schlegelmilch K, Haaf T, Schartl M, Schmid M. **2008**. Synteny conservation of the Z chromosome in 14 avian species (11 families) supports a role for Z dosage in avian sex determination. *Cytogenetic and Genome Research* **122**: 150–156.

Nguyen L-P, Galtier N, Nabholz B. **2015a**. Gene expression, chromosome heterogeneity and the fast-X effect in mammals. *Biology letters* **11**: 20150010–20150010.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. **2015b**. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution* **32**: 268–274.

Oatley G, De Swardt DH, Nuttall RJ, Crowe TM, Bowie RCK. **2017**. Phenotypic and genotypic variation across a stable white-eye (Zosterops sp.) hybrid zone in central South Africa. *Biological Journal of the Linnean Society* **121**: 670–684.

Oatley G, Voelker G, Crowe TM, Bowie RCK. **2012**. A multi-locus phylogeny reveals a complex pattern of diversification related to climate and habitat heterogeneity in southern African white-eyes. *Molecular Phylogenetics and Evolution* **64**: 633–644.

O'Connor RE, Romanov MN, Kiazim LG, Barrett PM, Farré M, Damas J, Ferguson-Smith M, Valenzuela N,

Larkin DM, Griffin DK. 2018. Reconstruction of the diapsid ancestral genome permits chromosome evolution tracing in avian and non-avian dinosaurs. *Nature Communications* 9: 1883.

Ohta T. 1992. The Nearly Neutral Theory of Molecular Evolution. *Annual Review of Ecology and Systematics* 23: 263–286.

de Oliveira EHC, Habermann FA, Lacerda O, Sbalqueiro IJ, Wienberg J, Müller S. 2005. Chromosome reshuffling in birds of prey: the karyotype of the world's largest eagle (Harpy eagle, Harpia harpyja) compared to that of the chicken (Gallus gallus). *Chromosoma* 114: 338–343.

Pala I, Naurin S, Stervander M, Hasselquist D, Bensch S, Hansson B. 2012a. Evidence of a neo-sex chromosome in birds. *Heredity* 108: 264–272.

Pala I, Hasselquist D, Bensch S, Hansson B. 2012b. Patterns of Molecular Evolution of an Avian Neo-sex Chromosome. *Molecular Biology and Evolution* 29: 3741–3754.

Papadopulos AST, Chester M, Ridout K, Filatov DA. 2015. Rapid Y degeneration and dosage compensation in plant sex chromosomes. *Proceedings of the National Academy of Sciences* 112: 13021.

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics* 20: 289–290.

Pedersen BS, Quinlan AR. 2018. Mosdepth: quick coverage calculation for genomes and exomes. *Bioinformatics* 34: 867–868.

Peona V, Weissensteiner MH, Suh A. 2018. How complete are "complete" genome assemblies?—An avian perspective. *Molecular Ecology Resources* 18: 1188–1195.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology* 33: 290.

Pessia E, Popa A, Mousset S, Rezvoy C, Duret L, Marais GAB. 2012. Evidence for widespread GC-biased gene conversion in eukaryotes. *Genome biology and evolution* 4: 675–682.

Philippe H, Vienne D, Ranwez V, Roure B, Baurain D, Delsuc F. Pitfalls in supermatrix phylogenomics. *European Journal of Taxonomy* 283: 1–25.

Poelstra JW, Vijay N, Bossu CM, Lantz H, Ryll B, Müller I, Baglione V, Unneberg P, Wikelski M, Grabherr MG, et al. 2014. The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. *Science* 344: 1410.

Pool JE, Nielsen R. 2007. Population size changes reshape genomic patterns of diversity. *Evolution; international journal of organic evolution* 61: 3001–3006.

Prum RO, Berv JS, Dornburg A, Field DJ, Townsend JP, Lemmon EM, Lemmon AR. 2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature* 526: 569.

R core team. 2018. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.

Rannala B, Yang Z. 2007. Inferring Speciation Times under an Episodic Molecular Clock. *Systematic Biology* 56: 453–466.

Raudsepp T, Santani A, Wallner B, Kata SR, Ren C, Zhang H-B, Womack JE, Skow LC, Chowdhary BP. 2004. A detailed physical map of the horse Y chromosome. *Proceedings of the National Academy of Sciences of the United States of America* 101: 9321.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3: 217–223.

Rousselle M, Faivre N, Ballenghien M, Galtier N, Nabholz B. 2016. Hemizygosity Enhances Purifying Selection: Lack of Fast-Z Evolution in Two Satyrine Butterflies. *Genome Biology and Evolution* 8: 3108–3119.

Roux C, Fraïsse C, Romiguier J, Anciaux Y, Galtier N,

Bierne N. **2016**. Shedding Light on the Grey Zone of
Speciation along a Continuum of Genomic Divergence.
PLOS Biology **14**: e2000234.
Salmela L, Rivals E. **2014**. LoRDEC: accurate and
efficient long read error correction. *Bioinformatics* **30**:
3506–3514.
Schwartz S, Kent WJ, Smit A, Zhang Z, Baertsch R,
Hardison RC, Haussler D, Miller W. **2003**. Human–
Mouse Alignments with BLASTZ. *Genome Research* **13**:
103–107.
She R, Chu JS-C, Uyar B, Wang J, Wang K, Chen N.
2011. genBlastG: using BLAST searches to build
homologous gene models. *Bioinformatics* **27**: 2141–
2143.
Sigeman H, Ponnikas S, Videvall E, Zhang H, Chauhan
P, Naurin S, Hansson B. **2018**. Insights into Avian
Incomplete Dosage Compensation: Sex-Biased Gene
Expression Coevolves with Sex Chromosome
Degeneration in the Common Whitethroat. *Genes* **9**.
Smeds L, Warmuth V, Bolivar P, Uebbing S, Burri R,
Suh A, Nater A, Bureš S, Garamszegi LZ, Hogner S, *et*
al. **2015**. Evolutionary analysis of the female-specific
avian W chromosome. *Nature Communications* **6**:
7330.
Smit A, Hubley R. **2008**. *RepeatModeler Open-1.0*.
Smit A, Hubley R, Green P. **2013**. *RepeatMasker Open-*
4.0.
Stanke M, Waack S. **2003**. Gene prediction with a
hidden markov model and a new intron submodel.
Bioinformatics **19**: 215–225.
Sun S, Heitman J. **2012**. Should Y stay or should Y go:
the evolution of non-recombining sex chromosomes.
BioEssays : news and reviews in molecular, cellular and
developmental biology **34**: 938–942.
Tilak M-K, Botero-Castro F, Galtier N, Nabholz B. **2018**.
Illumina Library Preparation for Sequencing the GC-Rich
Fraction of Heterogeneous Genomic DNA. *Genome*
Biology and Evolution **10**: 616–622.
Tomaszkiewicz M, Medvedev P, Makova KD. **2017**. Y

and W Chromosome Assemblies: Approaches and
Discoveries. *Trends in Genetics* **33**: 266–282.
Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan
G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L.
2010. Transcript assembly and quantification by RNA-
Seq reveals unannotated transcripts and isoform
switching during cell differentiation. *Nature*
biotechnology **28**: 511–515.
Vicoso B, Charlesworth B. **2006**. Evolution on the X
chromosome: unusual patterns and processes. *Nature*
Reviews Genetics **7**: 645.
Völker M, Backström N, Skinner BM, Langley EJ,
Bunzey SK, Ellegren H, Griffin DK. **2010**. Copy number
variation, chromosome rearrangement, and their
association with recombination during avian evolution.
Genome research **20**: 503–511.
Warren WC, Clayton DF, Ellegren H, Arnold AP, Hillier
LW, Künstner A, Searle S, White S, Vilella AJ, Fairley S,
et al. **2010**. The genome of a songbird. *Nature* **464**:
757.
Warren WC, Hillier LW, Tomlinson C, Minx P,
Kremitzki M, Graves T, Markovic C, Bouk N, Pruitt KD,
Thibaud-Nissen F, *et al.* **2016**. A New Chicken Genome
Assembly Provides Insight into Avian Genome
Structure. *G3 (Bethesda, Md.)* **7**: 109–117.
Waterhouse RM, Seppey M, Simão FA, Manni M,
Ioannidis P, Klioutchnikov G, Kriventseva EV, Zdobnov
EM. **2017**. BUSCO applications from quality
assessments to gene prediction and phylogenomics.
Molecular biology and evolution **35**: 543–548.
Weber CC, Boussau B, Romiguier J, Jarvis ED, Ellegren
H. **2014**. Evidence for GC-biased gene conversion as a
driver of between-lineage differences in avian base
composition. *Genome biology* **15**: 549–549.
Weissensteiner MH, Pang AWC, Bunikis I, Höijer I,
Vinnere-Petterson O, Suh A, Wolf JBW. **2017**.
Combination of short-read, long-read, and optical
mapping assemblies reveals large-scale tandem repeat
arrays with population genetic implications. *Genome*

Research **27**: 697–708.

Wickham H. **2016**. *ggplot2: Elegant Graphics for Data*
Analysis. Springer-Verlag New York.

Wilke C. **2016**. *cowplot: Streamlined Plot Theme and*
Plot Annotations for 'ggplot2'.

Wilson Sayres MA. **2018**. Genetic Diversity on the Sex
Chromosomes. *Genome Biology and Evolution* **10**:
1064–1078.

Wolf JBW, Künstner A, Nam K, Jakobsson M, Ellegren
H. **2009**. Nonlinear dynamics of nonsynonymous (dN)
and synonymous (dS) substitution rates affects
inference of selection. *Genome biology and evolution* **1**:
308–319.

Wright AE, Harrison PW, Zimmer F, Montgomery SH,
Pointer MA, Mank JE. **2015**. Variation in promiscuity
and sexual selection drives avian rate of Faster-Z
evolution. *Molecular ecology* **24**: 1218–1235.

Yang Z, Rannala B. **2006**. Bayesian Estimation of
Species Divergence Times Under a Molecular Clock
Using Multiple Fossil Calibrations with Soft Bounds.
Molecular Biology and Evolution **23**: 212–226.

Yoshida K, Makino T, Yamaguchi K, Shigenobu S,
Hasebe M, Kawata M, Kume M, Mori S, Peichel CL,
Toyoda A, *et al.* **2014**. Sex Chromosome Turnover
Contributes to Genomic Divergence between Incipient
Stickleback Species. *PLOS Genetics* **10**: e1004223.

Zhang G, Li C, Li Q, Li B, Larkin DM, Lee C, Storz JF,
Antunes A, Greenwold MJ, Meredith RW, *et al.* **2014**.
Comparative genomics reveals insights into avian
genome evolution and adaptation. *Science* **346**: 1311.

Zhang G, Parker P, Li B, Li H, Wang J. **2012**. The
genome of Darwin's Finch (Geospiza fortis).
GigaScience.

Zhou Q, Bachtrog D. **2012**. Sex-Specific Adaptation
Drives Early Sex Chromosome Evolution in *Drosophila*.
Science **337**: 341.

Zimin AV, Puiu D, Luo M-C, Zhu T, Koren S, Marçais G,
Yorke JA, Dvořák J, Salzberg SL. **2017**. Hybrid assembly
of the large and highly repetitive genome of Aegilops
tauschii, a progenitor of bread wheat, with the
MaSuRCA mega-reads algorithm. *Genome research* **27**:
787–792.

2141

Supplementary Information

2145 Table S1: Summary statistics of the *Zosterops* genome assemblies as computed by Assemblathon2. Values in bold correspond
2146 to the summary statistics of the publicly available sequences.

Species	Assembler	Reference	Assembly size	# scaffolds	Longest scaff.	N50 scaff.	L50 scaff.	N%	# contigs
Z. borbonicus	SOAPdenovo	this study	1188818773	130278	2861829	477419	714	6.43	213846
Z. borbonicus	**SOAPdenovo + SSPACE**	**this study**	**1222501712**	**97537**	**11984413**	**1762991**	**174**	**8.53**	**207715**
Z. borbonicus	SOAPdenovo + SSPACE + Synteny	this study	1222650938	96503	155428647	71485074	6	8.54	207709
Z. borbonicus	MaSuRCA	this study	1077182487	4487	11330673	1864885	153	0.41	9504
Z. pallidus	**SOAPdenovo**	**this study**	**1163666926**	**170557**	**3877680**	**374698**	**785**	**4.57**	**227542**
Z. lateralis	**AllPath_LG**	**Cornetti et al. 2015**	**1036003386**	**2933**	**15146312**	**3581248**	**83**	**3.32**	**55972**

2158   Table S2: List of the 27 avian species used in DeCoSTAR. Number of CDS translated to protein and used for orthology detection.

Species	Accession (NCBI) or URL	# Proteins
Acanthisitta chloris	http://dx.doi.org/10.5524/101015	15052
Anas platyrhynchos	http://dx.doi.org/10.5524/101001	15053
Aptenodytes forsteri	http://dx.doi.org/10.5524/100005	14767
Calypte anna	http://dx.doi.org/10.5524/101004	14543
Chaetura pelagica	http://dx.doi.org/10.5524/101005	14111
Charadrius vociferous	http://dx.doi.org/10.5524/101007	14465
Columba livia	http://dx.doi.org/10.5524/100007	14982
Corvus brachyrhynchos	http://dx.doi.org/10.5524/101008	14927
Corvus cornix	GCA_000738735.2_ASM73873v2	13622
Cuculus canorus	http://dx.doi.org/10.5524/101009	14727
Egretta garzetta	http://dx.doi.org/10.5524/101002	14127
Falco peregrinus	http://dx.doi.org/10.5524/101006	14839
Ficedula albicollis	GCA_000247815.2_FicAlb1.5	15382
Gallus gallus	ftp://climb.genomics.cn/pub/10.5524/100001_101000/101000/chicken/	14728
Geospiza fortis	http://dx.doi.org/10.5524/100040	14180
Haliaeetus leucocephalus	http://dx.doi.org/10.5524/101027	15212
Manacus vitellinus	http://dx.doi.org/10.5524/101010	16402
Melopsittacus undulates	http://dx.doi.org/10.5524/100059	14231
Nipponia Nippon	http://dx.doi.org/10.5524/101003	15018
Opisthocomus hoatzin	http://dx.doi.org/10.5524/101011	13333
Picoides pubescens	http://dx.doi.org/10.5524/101012	14136
Pygoscelis adeliae	http://dx.doi.org/10.5524/100006	13734
Struthio camelus	http://dx.doi.org/10.5524/101013	14577
Taeniopygia guttata	ftp://climb.genomics.cn/pub/10.5524/100001_101000/101000/zebrafinch/	15693
Tinamus guttatus	http://dx.doi.org/10.5524/101014	15723
Zonotrichia albicollis	GCF_000385455.1_Zonotrichia_albicollis-1.0.1	14376
Zosterops borbonicus	this study	22565

2159

2160

2161

2162

2163

2164

2165

2166   Table S3: Fossil calibration combinations used in the molecular dating analyses

Calibration sets	Node	Maximum bound (Myr)	Minimum bound (Myr)
1,2,3	Neognathae /Palaeognathae	86.5	66
1,2,3,4	Galloanserae / Neoaves	Free	66
1,2,3,4	Apodidae / Trochilidae	Free	51
1,2,3,4	Sphenisciformes / Threskiornithidae	Free	60.5
1,3	Passerines / Psittaciformes	65.5	53.3
2,4	Passerines / Psittaciformes	Free	53.3
3	Oscines / Suboscines	34	28
4	Neognathae / Palaeognathae	140	66

2181

A   B

2182

Figure S1: A) Phylogenetic tree based on all investigated avian species. The red star indicates the origin of the two neo-sex
chromosomes (ancestor of Sylvioidea). B) Summary statistics of the BUSCO analysis based on the 27 genome assemblies used
as input for DeCoSTAR. Blue, light blue, orange and red colors indicate single copy, duplicated, fragmented, missing genes,
respectively. The blue dotted line corresponds to the cumulative proportion of complete genes (single copy and duplicated
ones) found in *Z. borbonicus*. For unknown reasons, BUSCO analysis failed for the *F. albicollis* assembly.

2188

2189

2190

2191

2192

2193

2194

2195

2196

2197

2198

2199

2200

2201

2202 **Figure S2: Distribution of the TE content observed in the 22,565** *Z. borbonicus* **gene models.**

2203

2204

2205

2206

2207

2208

2209

2210

2211

2212

2213

2214

2215 **Figure S3:** *Molecular dating of the 30 birds species.* *Molecular dating estimates are based on the dataset 1 with CAT GTR*
2216 *substitution model, log-normal molecular rate model and calibration set 1 (Table S3).*

2217

2218

2219

2220

2221

2222

2223

2224

2225

2226

2227

2247 **Figure S4: Interchromosomal and interdataset variation in Tajima's $\pi_{females}$ (light blue) and π_{males} (orange) over non-**
2248 **overlapping 10-kb sliding windows.**

2249

2250

2251

2252

2253

Figure S5: Nucleotide diversity variations along 30 *Z. borbonicus* chromosomes as estimated using genetic information from 3 males (π_males, orange) and 3 females (π_females, light blue). For each dataset, top 2.5% and bottom 2.5% of π values among windows scanning all chromosomes are shown in green and red, respectively. The chromosomal breakpoint on the neoZ chromosome is shown with a red line.

2254
2255
2256
2257

2258

2259

Figure S6: Tajima's D variations along 30 *Z. borbonicus* chromosomes as estimated using genetic information from 3 males (orange) and 3 females (light blue). Top 2.5% and bottom 2.5% of D values among windows scanning all chromosomes are shown in green and red, respectively (baseline for bars is for D=0). The chromosomal breakpoint on the neoZ chromosome is shown with a red line.

2264

2265
2266 **Figure S7: Variation in G+C content along 30 *Z. borbonicus* chromosomes.** The red dotted line indicated the median GC value
2267 over non-overlapping 10kb sliding windows for scaffolds assigned to chromosomes only. As expected, strong departures from
2268 this median values are observed on minichromosomes.

2269

Figure S8: Variations in GC content at equilibrium (GC*) at third codon positions. Variability was obtained by bootstrapping genes within each genomic region.

2286

2287   **Figure S9: TE density for each chromosome and for each RepBase TE family.** Class I TE elements were categorized following
2288   the two subclasses: LTR and non-LTR retrotransposons.

2289

2290

2291

2292

2293

2294