The Genomic Landscape of Centromeres in Cancers

Anjan K. Saha[1,2,3], Mohamad Mourad[3], Mark H. Kaplan[3], Ilana Chefetz[4], Sami N. Malek[3], Ronald Buckanovich[5], David M. Markovitz [2,3,6,7], and Rafael Contreras-Galindo[3,*]

[1]Medical Scientist Training Program, University of Michigan, Ann Arbor, MI, 48109, USA

[2]Program in Cancer Biology, University of Michigan, Ann Arbor, MI, 48109, USA

[3]Department of Internal Medicine, University of Michigan, Ann Arbor, MI, 48109, USA

[4]The Hormel Institute, University of Minnesota, Austin, MN, 55912, USA

[5]Magee-Womens Research Institute, University of Pittsburgh, Pittsburgh, PA, 15213, USA

[6]Program in Cell and Molecular Biology, University of Michigan, Ann Arbor, MI, 48109, USA

[7]Program in Immunology, University of Michigan, Ann Arbor, MI, 48109, USA

*Correspondence: Rafael Contreras-Galindo

                      Department of Internal Medicine

                      Division of Infectious Diseases

                      University of Michigan, Ann Arbor, MI

                      48109-5640, USA

                      Tel#: (734) 936-8185

                      Fax#: (734) 764-0101

                      E-mail: rafaelc@med.umich.edu

ABSTRACT

Centromere genomics remain poorly characterized in cancer, due to technologic limitations in sequencing and bioinformatics methodologies that make high-resolution delineation of centromeric loci difficult to achieve. We here leverage a highly specific and targeted rapid PCR methodology to quantitatively assess the genomic landscape of centromeres in cancer cell lines and primary tissue. PCR-based profiling of centromeres revealed widespread heterogeneity of centromeric and pericentromeric sequences in cancer cells and tissue as compared to healthy counterparts. Quantitative reductions in select centromeric core and pericentromeric markers were observed in neoplastic samples as compared to healthy counterparts. Subsequent phylogenetic analysis of a pericentromeric endogenous retrovirus amplified by PCR revealed possible gene conversion events occurring at numerous pericentromeric loci in the setting of malignancy. Our findings collectively represent the first look into centromere genetics in the setting of malignancy, providing valuable insight into the evolution and reshuffling of centromeric sequences in cancer development and progression.

INTRODUCTION

 The centromere is essential to eukaryotic biology due to its critical role in genome inheritance (Cleveland et al.,

 2003; Hayden, 2012). The nucleic acid sequences that dominate the human centromeric landscape are α-satellites,

 arrays of ~171 base-pair monomeric units arranged into higher-order arrays throughout the centromere of each

 chromosome (Cleveland et al., 2003; Hayden, 2012; Henikoff et al., 2015). These α-satellites underlie a

 hierarchical network of proteins that collectively make up the kinetochore, a large multimeric structure that serves

 as a molecular bridge between chromosomes and microtubule polymers from the mitotic spindle during cell

 division. The interaction between centromeres, kinetochores and microtubule polymers lies at the nexus of

 metaphase and anaphase, ensuring faithful separation of the sister chromatids during mitosis.

 Centromeres are thus critical to maintaining the fidelity of chromosomal segregation in proliferating tissues.

 While much is known about the hierarchical network of proteins that epigenetically compartmentalizes

 centromeres, the genomic foundation of the centromere remains largely uncharted. Centromeres remain a genetic

 black box that encompasses 2-5% of the human genome (Aldrup-Macdonald and Sullivan, 2014). Despite

 advancements in next-generation sequencing (NGS) technologies, full assemblies of centromeric loci are still

 unavailable within the latest builds of the human genome, with the exception of a linear assembly of the

 centromere of chromosome Y (Jain et al., 2018). Low complexity genomic regions, characterized by the

 contiguous arrangement of repetitive sequences, present computational challenges owing to nonunique

 alignments that are impractical for current informatics pipelines to navigate. Low complexity regions like

 centromeric loci are consequently excluded from most downstream bioinformatics analyses.

 Methodologies that can add resolution to the genomic landscape of the centromere will thus play an integral

 role in developing a more nuanced understanding of its contribution to health and disease. Recent efforts at

 overcoming the technical shortcomings of NGS approaches have focused on more conventional molecular

 biology techniques, including extended chromatin fiber analysis, fluorescent *in-situ* hybridization (FISH),

 Southern blotting, and polymerase chain-reaction (PCR) based approaches (Aldrup-MacDonald et al., 2016;

 Contreras-Galindo et al., 2017; Li et al., 1995; Liehr, 2013). Chromatin fiber analysis, FISH, and Southern

 blotting, while effective for qualitatively characterizing localization and size of given centromeric proteins and

sequences, do not quantitatively assess the genomic content of the centromere. Moreover, PCR-based approaches offer expedited evaluation of the centromeric content within any given sample, making it more scalable than chromatin fiber analysis and hybridization-based approaches when evaluating samples derived from human cell lines and tissue. Corroboration of the specificity and sensitivity of PCR-approaches by a number of orthogonal methodologies suggests that using rapid centromere targeted PCR methodologies is a viable strategy for studying centromere genetics (Contreras-Galindo et al., 2017; Liehr, 2013; Miga et al., 2014; Shepelev et al., 2015).

Applying scalable PCR-based approaches to the assessment of centromere size and structure in different biological settings is therefore critical to contextualizing our knowledgebase on centromere genetics. Diseases of cell division, particularly cancer, remain largely unexplored within the realm of centromere genetics (Vig et al., 1989). Gaining deeper insight into the contribution of centromere genetics to tumorigenesis and cancer progression thus has the potential to inform novel therapeutic strategies capable of improving long-term outcomes. Unfortunately, the oncogenic potential of centromeric sequences remains undetermined, due to the shortcomings of sequencing methodologies.

Here we report global heterogeneity in the centromeric landscape in cancer cell lines and tissues. Both solid and hematologic tumors demonstrated marked alterations in the primary structure of centromeres and pericentromeres, as measured by a previously described quantitative centromere-specific PCR assay that targets core centromeric α-satellite DNA as well as pericentromeric human endogenous retrovirus (HERV) DNA (Contreras-Galindo et al., 2017). Phylogenetic analysis of HERV sequences in several cancer cell lines suggests centromeric sequences undergo aberrant recombination during tumorigenesis and/or disease progression. Strikingly, centromere contraction is a feature present across cancer tissue types, including primary tissue samples, providing further substantiation to the notion that genomic instability in centromeres is a ubiquitous occurrence in cancer. Evaluation of the centromeric landscape in the setting of malignancy thus reveals previously overlooked genetic alterations that may reflect novel pathophysiologic contributions to the development and progression of cancer.

RESULTS

Cancer Cell Lines Demonstrate Heterogeneous Losses in Centromeric DNA

NGS approaches to interrogating genetic alterations in cancer have repeatedly demonstrated ubiquitous genomic instability that is a hallmark of malignancy. However, the lack of an end-to-end assembly of centromeric loci prevents mapping of representative centromeric reads to a standardized reference. We have thus employed a rapid PCR-based approach that we previously described to evaluate the genomic landscape of centromeres in several human cancer cell lines (Figure 1) (Contreras-Galindo et al., 2017). These cell lines are representative of a variety of different tissue types, originating from both solid and hematologic malignancies. Our PCR-based methodology unveils significant heterogeneity in the centromeric and pericentromeric content in all 24 chromosomes. This heterogeneity extends to HERVs, such as HERV K111, that we have previously shown to reside in pericentromeric regions. Unsupervised hierarchical clustering of the chromosome specific repeats demonstrates a striking organization to the patterns in centromere heterogeneity, differentiated by the region of the centromere (core or pericentromere) to which each repeat localizes. Similar clustering analysis applied across the different cell lines revealed that heterogeneity in centromeric content is tissue type agnostic, with the exception of healthy peripheral blood lymphocytes (PBLs) that demonstrate higher relative concordance. The heterogeneity observed reflects a preference for contractions in centromeric content, consistent across numerous tissue types (Supplementary Figure S1A-S1E). Consistent with the loss of Chr Y previously reported in teratocarcinoma cells, we noted also that the centromere array of chromosome Y (DYZ3) was missing in teratocarinoma cell lines derived from male patients in this study (Supplementary Figure S1E) (Machiela et al., 2017; Nathanson et al., 2005). Of note, K111 contractions stood out as markedly ubiquitous across all evaluated cell lines. Collectively comparing normal PBMCs to cancer cell lines, grouped by tissue type, revealed marked reductions in centromeric material, using K111 copy number as a surrogate for pericentromeric content (Supplementary Figure S2) (Contreras-Galindo et al., 2013; Zahn et al., 2015).

A more focused analysis on breast cancer cell lines allowed us to cross-reference the observed heterogeneity in centromeric DNA against known molecular classifications for each cell line (Figure 2) (Taherian-Fard et al., 2015). Strikingly, the centromeric content demonstrated heterogeneity across the four molecular subtypes for breast cancer (Basal, HER2, Luminal A, and Luminal B); unsurprisingly, healthy PBLs clustered together. Similar

to other tissue types tested, breast cancer cell lines also demonstrated a predilection for contracted centromeres compared to healthy PBLs (Supplementary Figure S3). K111 again demonstrated robust contractions relative to other markers. The strong reduction in DYZ3 (α-satellite on chromosome Y) provided validation for the specificity of the rapid PCR-based approach to evaluating centromeric content, given the absence of Y-chromosomes in breast cancer cell lines derived from females. Taken together, marked heterogeneity in centromeric DNA is observed in cancer cell lines, with a predilection towards centromeric contraction when comparing cancer cell lines to healthy PBLs.

Gene Conversion of Centromeric Sequences in Cancer Cell Lines

The genomic landscape of the centromere is characterized by thousands of copies of repetitive elements arranged in tandem to form higher order arrays (Cleveland et al., 2003). Repetitive genomic regions are thus subject to recombination events due to sequence homology (Blanco et al., 2000; Klein, 1995). Intrachromosomal recombination is one form these events can take, leading to deletions that reduce the number of repeat units or gene conversion events that genetically homogenize the sequences of repeat units. Interestingly, in contrast to healthy PBLs, we identified drastic reductions in centromeric K111 sequences across all evaluated cancer cell lines (Figures 1 and 2). While real-time PCR demonstrates global deletion of centromeric material in cancer cell lines, purely quantitative assessments do not provide insight into other recombination events, such as gene conversion. Furthermore, sequence analysis of α-satellites is unreliable for identifying gene conversion events. We thus conducted phylogenetic analysis on the sequences of real-time PCR amplicons from cancer cell lines to identify gene conversion events within K111 loci, given ubiquitous loss of K111 across all cancer cell lines (Figure 3A). Our previous work has shown that divergence in K111 sequence similarity is dependent on chromosomal location of K111 loci (Contreras-Galindo et al., 2013; Zahn et al., 2015). We now show that K111 copies identified in breast cancer cell lines demonstrate cell line dependent sequence convergence towards K111 subtypes that organize into distinct clades (Figure 3B). The K151 cell line (pink) remarkably produced distinct clades that emerged in close proximity relative to each other from the same ancestral sequence. Sequences amplified from the K151 cell line were notably not distributed heterogeneously throughout the tree. Three

additional breast cancer cell lines (MDA-MB-435, DT-13, and HCC1599) formed two exclusive subtypes that

were also separated by phylogenetic analysis.

Phylogenetic analysis was also conducted in adult T-cell leukemia (ATL) cell lines and revealed similar

patterns as in breast cancer (Supplementary Figure S4). ATL26 alone formed three exclusive subtypes that diverge

in homology from normal K111 clades. Of note, K111 clades arising from ATL43 and ATL16 demonstrated

strong homology to K111 Solo LTRs, suggesting intrachromosomal recombination that is deleting centromeric

material. ATL43 and ATL16 indeed demonstrate the strongest reductions in K111 copy number relative to other

ATL cell lines (Supplementary Figure S2). As Solo LTRs are the result of homologous recombination between

the LTRs flanking endogenous retroviral sequences, ATL cell lines having *de novo* K111 sequences with higher

relative homology to Solo LTR sequences suggested that pericentromeric K111 sequences served as templates

for gene conversion. Taken together, cell line dependent sequence convergence of HERV-K111 in cancer cell

lines suggests that gene conversion events are driving sequence evolution within the pericentromeres of cancer

cell lines.

Heterogeneous Loss of Centromere DNA in Cancer Tissue

Human cancer cell lines are useful models for evaluating cancer biology and genetics in an *in vitro* setting.

Indefinite cellular propagation, however, results in clonal selection for cells that have a fitness advantage for

growing *ex vivo*. Such a fitness advantage is sometimes conferred by abnormal karyotypes (aneuploidy), a

cytogenetic feature that can influence the results of PCR based analyses. Evaluation of cancer tissue itself is thus

most reflective of the tissue architecture and microenvironment that modulate cancer biology and genetics. We

thus applied our rapid PCR-based approach to DNA isolated from primary cancer tissue. Profiling the centromeric

landscape in 9 different ovarian cancer samples against matched peripheral blood mononuclear cells (PBMCs)

revealed similarly significant heterogeneity observed in cell lines (Figure 4). Indeed, quantitative assessment of

this heterogeneity again revealed global centromeric contraction, similar to findings noted in cell lines

(Supplementary Figure S5). Strikingly, a drastic reduction in the centromere of chromosome 17 (D17Z1) was

seen in ovarian cancer tissue when compared to healthy tissue (Supplementary Figure S5), corroborating previous

reports of chromosome 17 anomalies in ovarian cancer (Tavassoli et al., 1993). No changes were seen in the single copy gene *GAPDH* found in the arm of chromosome 12.

While matched blood samples provide reliable non-malignant references to their malignant counterparts, comparisons between primary ovarian cancer tissue and matched blood does not sufficiently deconvolute tissue specific heterogeneity that may be present in normal biologic settings. To expand upon our findings, we profiled the centromeres of B-cells and T-cells that were separated by cell-surface marker selection from chronic lymphocytic leukemia (CLL) primary samples. CLL is a malignancy that arises in B-cells as opposed to T-cells within the bone marrow. Applying our methodology to compare patient matched B-cells and T-cells from CLL samples, both cells of lymphocytic lineage, thus eliminates the confounding contributions of normal development and tissue specificity to genetic heterogeneity in the centromere. Intriguingly, unsupervised hierarchical cluster analysis across patient samples cleanly separates healthy cells from diseased cells based on chromosome specific α-satellite abundance (Figure 5). We show contraction of numerous centromeres in malignant CD19 positive B-cells as compared to their normal CD3 positive T-cell counterparts, whereas no changes were seen in the housekeeping gene *GAPDH* found in the arm of chromosome 12 (Supplementary Figure S6). Strikingly, we see no such centromeric differences between B-cells and T-cells separated from blood samples derived from healthy individuals. Taken together, centromeric contraction is a characteristic that is present in primary cancer samples, consistent with our data in cancer cell lines.

DISCUSSION

The importance of centromeres to cell division provides a strong rationale for interrogating the genetics of the centromere in cancers. The challenges associated with studying the genomic landscape of centromeres, owing to the informatics impracticalities of evaluating low complexity regions, have however hindered meaningful progress in understanding the contributions of centromere genetics to tumorigenesis and cancer progression. Only one previous study reported the loss of centromere DNA in leukemia cells using fluorescent in situ hybridization (FISH)(MacKinnon and Campbell, 2011). We demonstrate, for the first time, that centromeres display heterogeneous alterations in the setting of malignancy, both in cancer cell lines and primary samples. We show

that these heterogeneous alterations reflect global reductions and gene conversions of repetitive elements and HERVs, suggesting that oncogenic genomic instability selects against the presence of centromeric sequences. While mechanistically uncharacterized, these findings have direct implications for our understanding of global genomic instability in cancer, given the importance of centromeres to faithful segregation of chromosomes. The loss of centromeric material in chromosome 17 described above is an example of the concordance between centromere instability and ovarian cancer pathogenesis, given the recurrent alterations in chromosome 17 that have been previously described in ovarian cancer (Tavassoli et al., 1993). Aberrant segregation of chromosomes has also been shown to result in the formation of micronuclei, a cellular event that precipitates catastrophic processes such as chromothripsis (Zhang et al., 2015). Gaining deeper insight into the mechanism driving gene conversion and centromere contraction may facilitate the development of novel therapeutic approaches that can prevent catastrophic and potentially oncogenic events like micronuclei formation and chromothripsis.

Though instability in centromeric loci was poorly described previously, in malignancy global genome instability is a well characterized hallmark of cancer (Hanahan and Weinberg, 2011). Subsets of breast and ovarian cancer have well studied DNA repair aberrations in homologous recombination proteins BRCA1/2 (Roy et al., 2011). Recent genomic profiling of several other malignancies has identified new disease subsets classified by molecular alterations in DNA repair genes and pathways (Robinson et al., 2015, 2017). It is conceivable that subsets of cancer that are dysfunctional in DNA repair may exhibit pronounced heterogeneity in centromeric content. Stratifying samples by DNA repair signatures prior to profiling the genomic landscape of centromeres may provide a strategy for identifying mechanistic contributors to centromere contraction in the setting of malignancy. Moreover, genomic profiling of centromeres in cancer tissue may produce signatures that are predictive of responders to therapies that target the DNA repair machinery, such as poly-ADP ribose polymerase (PARP) inhibitors.

In conclusion, we here provide quantitative resolution to the largely uncharacterized human centromere in the setting of cancer. We notably shed light on a genomic region that has been widely considered a black box and impervious to inquiry at the genomic level. The heterogeneous alterations observed in cancer cell lines and primary tissue provide a sound rationale to mechanistically interrogate the molecular machinery that is likely

driving the selection against centromeric material. Mechanistic characterization of genomic instability at centromeric loci has the potential to inform therapeutic approaches aimed at improving disease outcomes across several cancer types.

MATERIALS AND METHODS

Cell Lines and Cell Culture. Cell lines were cultured according to American Type Culture Collection (ATCC) recommendations. Cell lines were grown at 37 °C in a 5% CO_2 cell culture incubator and authenticated by short tandem repeat (STR) profiling for genotype validation at the University of Michigan Sequencing Core. ATL cell lines were cultured and authenticated as previously described (Maeda et al., 1985).

DNA Isolation. DNA extraction was performed on cell lines and tissue with the DNeasy Blood and Tissue Kit (QIAGEN) according to manufacturer's instructions. DNA was preserved at -20° C.

Blood and Tumor Cell Separation. Between January 2005 and September 2016 patients with chronic lymphocytic leukemia (CLL) evaluated at the University of Michigan Comprehensive Cancer Center were enrolled onto this study. The trial was approved by the University of Michigan Institutional Review Board (IRB no. HUM00045507). Patients consented for tissue donation in accordance with a protocol approved by the University of Michigan's IRB (IRB no. HUM0009149). Written informed consent was obtained from all patients before enrollment in accordance with the Declaration of Helsinki. CLL diagnostic criteria were based on the National Cancer Institute Working Group Guidelines for CLL. Eligible patients needed to have an absolute lymphocytosis (> 5000 mature lymphocytes/μL), and lymphocytes needed to express CD19, CD23, sIg (weak), and CD5 in the absence of other pan-T-cell markers. Peripheral blood mononuclear cells (PBMCs) were isolated by venipuncture and separated using Histopaque-1077 (Sigma). Cryopreserved PBMCs (frozen after Ficoll-gradient purification) from CLL blood specimens were prepared for FACS and sorted into CD19+ and CD3+ cells as previously described (Kujawski et al., 2008). Ovarian cancer DNA were isolated from Stage IIIc or Stage IV ovarian carcinomas. Tumor samples were obtained from the operating room and immediately taken to the

laboratory for processing. Tissue was maintained in RPMI/10% FBS throughout processing. Fresh $4 \times 4 \times 2$–mm

tumor slices were rinsed several times to remove all loosely attached cells. The tissue was then placed in a tissue

culture dish and DNA was extracted as described above.

Rapid Centromere Target PCR Assay. PCR was conducted on DNA samples from cell lines and primary cancer

samples according the previously described conditions (Contreras-Galindo et al., 2017). Briefly, copy numbers

for each centromeric array, proviruses K111/K222, and single-copy genes were measured by qPCR using specific

primers and PCR conditions as described. PCR amplification products were confirmed by sequencing. The qPCR

was carried out using the Radiant Green Low-Rox qPCR master mix (Alkali Scientific) with an initial enzyme

activation step for 10 min at 95°C and 16–25 cycles consisting of 15 sec of denaturation at 95°C and 30 sec of

annealing/extension.

PCR for 5' and 3' K111 LTR Insertions. K111 insertions were amplified by PCR using the Expand Long Range

dNTPack PCR kit (Roche Applied Science, Indianapolis, IN) as described. K111 5' and 3' LTRs and

accompanying flanking regions were amplified. PCR was performed using an initial step of 94 °C for 2 min

followed by 35 cycles consisting of denaturation at 94 °C for 30 sec, annealing at 55 °C for 30 sec, and extension

at 68 °C for 5 min. The amplification products were cloned into the topo TA vector (Invitrogen, Carlsbad, CA)

and sequenced.

Phylogenetic Analysis. The K111-related LTR sequences obtained from the DNA of cell lines, and DNA from

human/rodent chromosomal cell hybrids were subjected to BLAST analysis against the NCBI nucleotide

database. Sequences were aligned in BioEdit and exported to the MEGA5 matrix. LTR trees were generated using

Bayesian inference (MrBayes v 3.2 (Ronquist and Huelsenbeck, 2003)) with four independent chains run for at

least 1,000,000 generations until sufficient trees were sampled to generate more than 99% credibility.

Statistics and Data Analysis. All heatmaps were generated using the R statistical programming language. Data were log$_2$ normalized to the median values of healthy samples. Tests of statistical significance employed two-sided student t-tests, with level of significance denoted on appropriate plots.

ACKNOWLEDGEMENTS

The authors are grateful to Debra Buck and Maureen Legendre for assistance and Dr. Gil Omenn for inspiring discussions. ATL cell lines were kindly provided by Dr. Mikulus Popovic from the Institute of Human Virology at the University of Maryland.

COMPETING INTEREST

Anjan K. Saha is an equity holder in Deoxylytics, LLC.

REFERENCES

Aldrup-MacDonald ME, Kuo ME, Sullivan LL, Chew K, Sullivan BA. 2016. Genomic variation within alpha satellite DNA influences centromere location on human chromosomes with metastable epialleles. *Genome Res* **26**:1301–1311. doi:10.1101/gr.206706.116

Aldrup-Macdonald ME, Sullivan BA. 2014. The past, present, and future of human centromere genomics. *Genes (Basel)* **5**:33–50.

Blanco P, Shlumukova M, Sargent CA, Jobling MA, Affara N, Hurles ME. 2000. Divergent outcomes of intrachromosomal recombination on the human Y chromosome: male infertility and recurrent polymorphism. *J Med Genet* **37**:752–758.

Cleveland DW, Mao Y, Sullivan KF. 2003. Centromeres and kinetochores: from epigenetics to mitotic checkpoint signaling. *Cell* **112**:407–421.

Contreras-Galindo R, Fischer S, Saha AK, Lundy JD, Cervantes PW, Mourad M, Wang C, Qian B, Dai M, Meng F, Chinnaiyan A, Omenn GS, Kaplan MH, Markovitz DM. 2017. Rapid molecular assays to study human centromere genomics. *Genome Res* **27**:2040–2049. doi:10.1101/gr.219709.116

Contreras-Galindo R, Kaplan MH, He S, Contreras-Galindo AC, Gonzalez-Hernandez MJ, Kappes F, Dube D,

Chan SM, Robinson D, Meng F, Dai M, Gitlin SD, Chinnaiyan AM, Omenn GS, Markovitz DM. 2013.

HIV infection reveals widespread expansion of novel centromeric human endogenous retroviruses.

Genome Research **23**:1505–1513. doi:10.1101/gr.144303.112

Hanahan D, Weinberg RA. 2011. Hallmarks of cancer: the next generation. *Cell* **144**:646–674.

doi:10.1016/j.cell.2011.02.013

Hayden KE. 2012. Human centromere genomics: now it's personal. *Chromosome Research* **20**:621–633.

doi:10.1007/s10577-012-9295-y

Henikoff JG, Thakur J, Kasinathan S, Henikoff S. 2015. A unique chromatin complex occupies young -satellite

arrays of human centromeres. *Science Advances* **1**:e1400234–e1400234. doi:10.1126/sciadv.1400234

Jain M, Olsen HE, Turner DJ, Stoddart D, Bulazel KV, Paten B, Haussler D, Willard HF, Akeson M, Miga KH.

2018. Linear assembly of a human centromere on the Y chromosome. *Nat Biotechnol* **36**:321–323.

doi:10.1038/nbt.4109

Klein HL. 1995. Genetic control of intrachromosomal recombination. *Bioessays* **17**:147–159.

doi:10.1002/bies.950170210

Kujawski L, Ouillette P, Erba H, Saddler C, Jakubowiak A, Kaminski M, Shedden K, Malek SN. 2008.

Genomic complexity identifies patients with aggressive chronic lymphocytic leukemia. *Blood*

112:1993–2003. doi:10.1182/blood-2007-07-099432

Li X, Tsuji T, Wen S, Mimura Y, Wang Z, Sasaki K, Shinozaki F. 1995. A fluorescence in situ hybridization

(FISH) analysis with centromere-specific DNA probes of chromosomes 3 and 17 in pleomorphic

adenomas and adenoid cystic carcinomas. *J Oral Pathol Med* **24**:398–401.

Liehr T. 2013. Benign and Pathological Chromosomal Imbalances: Microscopic and Submicroscopic Copy

Number Variations (CNVs) in Genetics and Counseling. Academic Press.

Machiela MJ, Dagnall CL, Pathak A, Loud JT, Chanock SJ, Greene MH, McGlynn KA, Stewart DR. 2017.

Mosaic chromosome Y loss and testicular germ cell tumor risk. *J Hum Genet* **62**:637–640.

doi:10.1038/jhg.2017.20

MacKinnon RN, Campbell LJ. 2011. The Role of Dicentric Chromosome Formation and Secondary Centromere

Deletion in the Evolution of Myeloid Malignancy. *Genetics Research International* **2011**:1–11.

doi:10.4061/2011/643628

Maeda M, Shimizu A, Ikuta K, Okamoto H, Kashihara M, Uchiyama T, Honjo T, Yodoi J. 1985. Origin of

human T-lymphotrophic virus I-positive T cell lines in adult T cell leukemia. Analysis of T cell receptor

gene rearrangement. *Journal of Experimental Medicine* **162**:2169–2174. doi:10.1084/jem.162.6.2169

Miga KH, Newton Y, Jain M, Altemose N, Willard HF, Kent WJ. 2014. Centromere reference models for

human chromosomes X and Y satellite arrays. *Genome Res* **24**:697–707. doi:10.1101/gr.159624.113

Nathanson KL, Kanetsky PA, Hawes R, Vaughn DJ, Letrero R, Tucker K, Friedlander M, Phillips K-A, Hogg

D, Jewett MAS, Lohynska R, Daugaard G, Richard S, Chompret A, Bonaïti-Pellié C, Heidenreich A,

Olah E, Geczi L, Bodrogi I, Ormiston WJ, Daly PA, Oosterhuis JW, Gillis AJM, Looijenga LHJ,

Guilford P, Fosså SD, Heimdal K, Tjulandin SA, Liubchenko L, Stoll H, Weber W, Rudd M, Huddart R,

Crockford GP, Forman D, Oliver DT, Einhorn L, Weber BL, Kramer J, McMaster M, Greene MH, Pike

M, Cortessis V, Chen C, Schwartz SM, Bishop DT, Easton DF, Stratton MR, Rapley EA. 2005. The Y

deletion gr/gr and susceptibility to testicular germ cell tumor. *Am J Hum Genet* **77**:1034–1043.

doi:10.1086/498455

Robinson D, Van Allen EM, Wu Y-M, Schultz N, Lonigro RJ, Mosquera J-M, Montgomery B, Taplin M-E,

Pritchard CC, Attard G, Beltran H, Abida W, Bradley RK, Vinson J, Cao X, Vats P, Kunju LP, Hussain

M, Feng FY, Tomlins SA, Cooney KA, Smith DC, Brennan C, Siddiqui J, Mehra R, Chen Y, Rathkopf

DE, Morris MJ, Solomon SB, Durack JC, Reuter VE, Gopalan A, Gao J, Loda M, Lis RT, Bowden M,

Balk SP, Gaviola G, Sougnez C, Gupta M, Yu EY, Mostaghel EA, Cheng HH, Mulcahy H, True LD,

Plymate SR, Dvinge H, Ferraldeschi R, Flohr P, Miranda S, Zafeiriou Z, Tunariu N, Mateo J, Perez-

Lopez R, Demichelis F, Robinson BD, Schiffman M, Nanus DM, Tagawa ST, Sigaras A, Eng KW,

Elemento O, Sboner A, Heath EI, Scher HI, Pienta KJ, Kantoff P, de Bono JS, Rubin MA, Nelson PS,

Garraway LA, Sawyers CL, Chinnaiyan AM. 2015. Integrative Clinical Genomics of Advanced Prostate

Cancer. *Cell* **161**:1215–1228. doi:10.1016/j.cell.2015.05.001

Robinson DR, Wu Y-M, Lonigro RJ, Vats P, Cobain E, Everett J, Cao X, Rabban E, Kumar-Sinha C, Raymond

V, Schuetze S, Alva A, Siddiqui J, Chugh R, Worden F, Zalupski MM, Innis J, Mody RJ, Tomlins SA,

Lucas D, Baker LH, Ramnath N, Schott AF, Hayes DF, Vijai J, Offit K, Stoffel EM, Roberts JS, Smith

DC, Kunju LP, Talpaz M, Cieślik M, Chinnaiyan AM. 2017. Integrative clinical genomics of metastatic

cancer. *Nature* **548**:297–303. doi:10.1038/nature23306

Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models.

Bioinformatics **19**:1572–1574.

Roy R, Chun J, Powell SN. 2011. BRCA1 and BRCA2: different roles in a common pathway of genome

protection. *Nat Rev Cancer* **12**:68–78. doi:10.1038/nrc3181

Shepelev VA, Uralsky LI, Alexandrov AA, Yurov YB, Rogaev EI, Alexandrov IA. 2015. Annotation of

suprachromosomal families reveals uncommon types of alpha satellite organization in pericentromeric

regions of hg38 human genome assembly. *Genom Data* **5**:139–146. doi:10.1016/j.gdata.2015.05.035

Taherian-Fard A, Srihari S, Ragan MA. 2015. Breast cancer classification: linking molecular mechanisms to

disease prognosis. *Brief Bioinformatics* **16**:461–474. doi:10.1093/bib/bbu020

Tavassoli M, Ruhrberg C, Beaumont V, Reynolds K, Kirkham N, Collins WP, Farzaneh F. 1993. Whole

chromosome 17 loss in ovarian cancer. *Genes Chromosomes Cancer* **8**:195–198.

Vig BK, Sternes KL, Paweletz N. 1989. Centromere structure and function in neoplasia. *Cancer Genet*

Cytogenet **43**:151–178.

Zahn J, Kaplan MH, Fischer S, Dai M, Meng F, Saha AK, Cervantes P, Chan SM, Dube D, Omenn GS,

Markovitz DM, Contreras-Galindo R. 2015. Expansion of a novel endogenous retrovirus throughout the

pericentromeres of modern humans. *Genome Biol* **16**:74. doi:10.1186/s13059-015-0641-1

Zhang C-Z, Spektor A, Cornils H, Francis JM, Jackson EK, Liu S, Meyerson M, Pellman D. 2015.

Chromothripsis from DNA damage in micronuclei. *Nature* **522**:179–184. doi:10.1038/nature14493

FIGURES AND LEGENDS

Figure 1. Heterogeneous loss of centromeres in cancer cell lines. Heatmap representing the abundance of α-satellites specific for each centromere array (Y-axis) obtained by qPCR in 50 ng of DNA from healthy cells and from cancer lines (X-axis). Relative abundance is denoted by the gradient legend (top left). Cancer type and α-satellite localization is depicted as indicated by the legend (bottom left). Data depicting α-satellite abundance are $\log_2$ normalized to healthy PBL median values (asterisks, red). The nomenclature of these α-satellites begins with the letter D, followed by their resident chromosome number (1–22, X or Y), followed by a Z, and a number indicating the order in which the sequence was discovered. The DYZ3 repeat was excluded from the analysis to reduce confounding from gender.

Figure 2. Genomic profiling of centromeres in breast cancer cell lines. Heatmap representing the abundance of α-satellites specific for each centromere array (Y-axis) obtained by qPCR in 50 ng of DNA from healthy cells and from breast cancer lines (X-axis). Relative abundance is denoted by the gradient legend (bottom left). Data depicting α-satellite abundance are log_2 normalized to healthy PBL median values (asterisks). Hormone receptor, *TP53* status, histologic, and molecular classifications are depicted as indicated by the legend (top left). The nomenclature of these α-satellites begins with the letter D, followed by their resident chromosome number (1–22, X or Y), followed by a Z, and a number indicating the order in which the sequence was discovered. The DYZ3 repeat was excluded from the analysis to reduce confounding from gender.

Figure 3. Gene conversion of HERV-K111 in breast cancer cell lines. A) Schematic outline of the experimental methodology employed to identify gene conversion events. B) Phylogenetic analysis conducted on K111 sequences amplified by PCR on breast cancer cell lines (T47D, BT549, HCC-1599, MD-MB-435, DT13, DT22,

K151, and SKBr3). Amplicons are labeled and color-coded along the edge of the phylogenetic tree according to the cell line that produced the amplicon. Amplicons from human-hamster hybrid cell lines are denoted numerically by the human chromosome present in each hybrid cell line. Amplicons from K111 5'LTR, 3'LTR, and Solo LTR are additionally denoted. An example of gene conversion is shown in the cell line K151, possessing clades (pink) that localize in close proximity relative to each other but are not found heterogeneously throughout the tree. Convergence on two distinct K111 subtypes can additionally be identified within the MDA-MB-435, DT-13, and HCC1599 cell lines.

Figure 4. Genomic profiling of centromeres in primary ovarian cancer tissue. Heatmap representation of rapid
PCR data from nine primary ovarian cancer tissue samples with matched PBMC DNA. Data depicting α-satellite
abundance are $\log_2$ normalized to PBMC median values. Relative abundance is denoted by the gradient legend
(bottom left). The nomenclature of these α-satellites begins with the letter D, followed by their resident
chromosome number (1–22, X or Y), followed by a Z, and a number indicating the order in which the sequence
was discovered.

Figure 5. CLL and tissue matched T-cells assessed for select centromeric α-satellite markers. Heatmap representation of rapid PCR data from six primary CLL and two healthy samples post-separation by indicated cell surface markers into B-cell (CD19+) and T-cell (CD3+) populations. Data depicting α-satellite abundance are $\log_2$ normalized to T-cell median values. Relative abundance is denoted by the gradient legend (bottom left). Lymphocyte characterization and disease status is depicted as indicated by the legend (top left). The nomenclature of these α-satellites begins with the letter D, followed by their resident chromosome number (1–22, X or Y), followed by a Z, and a number indicating the order in which the sequence was discovered.