**Strong Patterns of Intraspecific Variation and Local Adaptation in Great Basin Plants**

**Revealed Through a Review of 75 Years of Experiments**

Owen W. Baughman[1,2], Alison C. Agneray[1], Matthew L. Forister[3], Francis F. Kilkenny[4], Erin K.

Espeland[5], Rob Fiegener[6], Matthew E. Horning[7], Richard C. Johnson[8], Thomas N. Kaye[6], Jeffrey

E. Ott[4], J. Bradley St. Clair[9], Elizabeth A. Leger[1]

[1]Department of Natural Resources and Environmental Science, University of Nevada, 1664 N.

Virginia St., Reno, NV 89557 USA; [2]Corresponding author, owbaughman@gmail.com, current

address: The Nature Conservancy, 67826-A Hwy. 205, Burns, OR 97720 USA, [3]Department of

Biology, University of Nevada, Reno, 1664 N. Virginia St., Reno, NV 89557 USA; [4]USDA

Forest Service, Rocky Mountain Research Station, 322 E Front St. Suite 401, Boise, ID 83702

USA; [5]Pest Management Research Unit, USDA-Agricultural Research Service Northern Plains

Agricultural Laboratory, 1500 N Central Ave., Sidney, MT 59270 USA; [6]Institute for Applied

Ecology, 563 SW Jefferson Ave., Corvallis, OR 97333 USA; [7]USDA Forest Service Pacific

Northwest Region, Deschutes National Forest, 63095 Deschutes Market Rd., Bend, OR 97701;

[8]Washington State University, PO Box 646240, Pullman, WA 99164 USA; [9]USDA Forest

Service, Pacific Northwest Research Station, 3200 SW Jefferson Way, Corvallis OR, 97331

**Keywords:** Local adaptation, phenotypic traits, meta-analysis, common garden, reciprocal

transplant, natural selection, intraspecific variation, restoration

**Running head:** Local adaptation in Great Basin plants

**Abstract**

Variation in natural selection across heterogeneous landscapes often produces 1) among-

population differences in phenotypic traits, 2) trait-by-environment associations, and 3) higher

fitness of local populations. Using a broad literature review of common garden studies published

between 1941 and 2017, we documented the commonness of these three signatures in plants

native to North America's Great Basin, an area of extensive restoration and revegetation efforts,

and asked which traits and environmental variables were involved. We also asked, independent

of geographic distance, whether populations from more similar environments had more similar

traits. From 327 experiments testing 121 taxa in 170 studies, we found 95.1% of 305 experiments

reported among-population differences, and 81.4% of 161 experiments reported trait-by-

environment associations. Locals showed greater survival in 67% of 24 reciprocal experiments

that reported survival, and higher fitness in 90% of 10 reciprocal experiments that reported

reproductive output. A meta-analysis on a subset of studies found that variation in eight

commonly-measured traits was associated with mean annual precipitation and mean annual

temperature at the source location, with notably strong relationships for flowering phenology,

leaf size, and survival, among others. Although the Great Basin is sometimes perceived as a

region of homogeneous ecosystems, our results demonstrate widespread habitat-related

population differentiation and local adaptation. Locally-sourced plants likely harbor adaptations

at rates and magnitudes that are immediately relevant to restoration success, and our results

suggest that certain key traits and environmental variables should be prioritized in future

assessments of plants in this region.

**Introduction**

All plant species have limits to the range of conditions in which they can live, and all but the narrowest endemics grow across environments that vary in biotic and abiotic conditions. This natural complexity has significant impacts on individual survival and reproduction, and thus plant evolution (Loveless and Hamrick, 1984; Linhart and Grant, 1996; Ackerly *et al.*, 2000; Reich *et al.*, 2003). As plants are subject to different conditions associated with their local environment, populations of the same species will experience differential selection pressures (Turesson, 1922; Clausen, Keck and Hiesey, 1948; Antonovics and Bradshaw, 1968; Langlet, 1971), creating habitat-correlated intraspecific variation. When this intraspecific variation results in populations that are more fit in their home environment than foreign populations, these populations are considered to be locally adapted (Kawecki and Ebert, 2004; Blanquart *et al.*, 2013). The existence of local adaptation is well-established across different organisms and ecosystems, although our synthetic knowledge of this important topic rests on surprisingly few reviews of the subject (e.g. Leimu and Fischer, 2008; Hereford, 2009; Oduor, Leimu and van Kleunen, 2016). Here, we focus on a particular region and ask if plant species share patterns of intraspecific variation and local adaptation, and, across taxa, what functional traits and environmental variables are most important for such patterns in this region. The regional focus provides a strong test of expectations generated from more heterogeneous samples, facilitates comparison of the strength of selection among specific traits, and provides an opportunity to link basic evolutionary patterns with applied concerns.

The detection of local adaptation ideally involves reciprocal transplant experiments designed to test for a local advantage across environments (Blanquart *et al.*, 2013; Bucharova, Durka, *et al.*, 2017). However, patterns associated with local adaptation (hereafter, signatures)

can be detected in non-reciprocal comparisons of different populations of the same species

(Endler, 1986). When populations are locally adapted to environmental variables, we expect to

see three basic signatures from common garden experiments: 1) differences among populations

in fitness-related traits, 2) correlations between these trait values and environmental or other

habitat-related variables, and, if reciprocal transplants have been conducted, 3) higher fitness of

local over nonlocal populations in the local environment. Although population differences

(signature 1) are necessary for local adaptation, they alone are not sufficient evidence due to

factors such as genetic drift, high gene flow, and rapid environmental change, among other

factors (Kawecki and Ebert, 2004; Blows and Hoffmann, 2005). While fitness differences in

reciprocal transplant experiments (signature 3) are the "gold standard" for detecting local

adaptation, there are experimental trade-offs between the number of populations sampled and the

ability to do fully reciprocal transplants (Blanquart *et al.*, 2013). Thus, correlative approaches

(signature 2) are popular alternatives that can sample many more populations to infer local

adaptation (e.g. St Clair, Mandel and Vance-Borland, 2005), though spurious correlations, low

sample sizes, or high variability in trait values could over- or under-predict the degree of local

adaptation in wild populations using this approach. Given these considerations, separately

reporting all three signatures can give an overall picture of the likelihood of within-species

variation and potential local adaptation in a region, and is the first step towards a better

understanding of variation in the strength and consistency of natural selection (Siepielski,

Dibattista and Carlson, 2009).

The Great Basin Desert of North America is a ~540,000 km$^2$ cold desert landscape

characterized by hundreds of internally-draining basin and range formations, which create high

spatial and environmental heterogeneity and variability (Tisdale and Hironaka, 1981; Comstock

and Ehleringer, 1992). While these are the kinds of conditions that would be expected to result in

widespread local adaptation, the flora of the Great Basin is poorly represented in the relatively

few reviews on the subject (Leimu and Fischer, 2008; Hereford, 2009; Oduor, Leimu and van

Kleunen, 2016), and this has resulted in uncertainty as to the prevalence, magnitude, and

importance that local adaptation plays in this large and increasingly imperiled region (United

States. House of Representatives. Committee on Appropriations., 2014; Jones, Monaco and

Rigby, 2015; Chivers *et al.*, 2016). Gaining a better understanding of local adaptation in the

Great Basin is important not only because it is a large, relatively intact floristic region in the

Western US, but also because this information has direct impacts on conservation and restoration

efforts. Large-scale, seed-based restoration has been very common in the Great Basin for many

decades (Pilliod, Welty and Toevs, 2017), and trends in large destructive wildfires (Dennison *et*

*al.*, 2014) and other disturbances (Rowland, Suring and Michael, 2010; Davies *et al.*, 2011)

ensure even higher demand for restoration efforts in the future. Guided by the various national

policies and strategies dating from the 1960s (Richards, Chambers and Ross, 1998) to the present

National Seed Strategy (Plant Conservation Alliance, 2015) and Integrated Rangeland Fire

Management Strategy (USDOI, 2015), a growing majority of these efforts are using native

plants. However, few of the widely-available sources of commercially-produced seeds of native

species originate from populations within the Great Basin (Jones and Larson, 2005) or have been

selected based on their success in restoring Great Basin habitats (Leger and Baughman, 2015).

Further, demand for native seed has always exceeded supply (McArthur and Young, 1999;

Johnson *et al.*, 2010), which has resulted in the prioritization of seed quantity and uniformity

over population suitability and local adaptation (Meyer, 1997; Richards, Chambers and Ross,

1998; Leger and Baughman, 2015). Therefore, it is still uncommon for restorationists in this

region to prioritize or even have the option to prioritize the use of local populations, despite

growing support of the importance of such practices (Basey, Fant and Kramer, 2015; Espeland *et*

*al.*, 2017).

Though our understanding of the prevalence and scale of local adaptation in the Great

Basin is far from complete, there is an abundant literature of peer-reviewed studies on the plants

native to this region spanning over 75 years that have directly measured trait variation between

populations via laboratory, greenhouse, or field common gardens and reciprocal transplants.

Many of these studies have also tested for correlations between intraspecific variation and

environmental variables, and some were designed to detect local adaptation. This research

includes studies of germination patterns (e.g. McArthur, Meyer and Weber, 1987; Meyer *et al.*,

1995), large genecology experiments (e.g. Erickson, Mandel and Sorenson, 2004; Johnson,

Leger and Vance-Borland, 2017), and reciprocal transplants (e.g. Evans and Young, 1990;

Barnes, 2009), among other types of studies. This rich literature provides an opportunity to

summarize local adaptation and its associated patterns, or signatures (defined above), in this

region, as well as describe which phenotypic traits have the strongest signatures of local

adaptation.

Here, we present results of a broad literature review and subsequent meta-analysis using

published studies that compared phenotypic traits of multiple populations of native Great Basin

species in one or more common environments. Our first objective was to record published

instances of the three expected signatures of local adaptation (population variation, trait-by-

environment association, and greater local fitness) within grasses, forbs, shrubs, and deciduous

trees native to the Great Basin, asking how common these signatures are, as well as which

phenotypic traits and environmental variables were most commonly associated with these

signatures. We also present results by taxonomic group, lifeform, lifespan, distribution, and

mating system. This first objective encompassed all possible studies, including those that did not

provide sufficient details for formal meta-analysis, which allowed us to incorporate the broadest

range of studies, including older studies that provided minimal quantitative detail. Our second

objective was to examine links between the magnitude of trait and environmental divergence

(mean annual precipitation and mean annual temperature) among populations across multiple

taxa, for the subset of experiments amenable to this approach, asking whether populations from

more similar environments were more similar in phenotypic traits. We also used meta-analysis to

ask which traits and environmental variables showed the strongest patterns of association.

We expected to find widespread evidence of local adaptation and its signatures in the

plants of the Great Basin, and we hypothesize that phenological and size-based traits, which

show phenotypic variation in response to climate variation in both plants and animals (e.g.

Sheridan and Bickford, 2011; Anderson *et al.*, 2012) and have been observed to be under

selection in the Great Basin (Leger and Baughman, 2015), would be important indicators of

adaptation in this region. We discuss our results both as a contribution to our general

understanding of natural selection in plants, and as an example of evolutionary theory applied to

the management and restoration of a large geographic region, where active and ongoing

management can benefit from information on intraspecific variation and local adaptation.

**Methods**

*Literature search*

We began by using the search engines Google Scholar and Web of Science to search for

combinations of key terms (see additional methods in Supporting Information Appendix 1). In

order to be included in our review, a study had to meet all these criteria:

160           a)   Examined a species that is native within the floristic Great Basin

161           b)   Examined and compared more than one population of that species

162           c)   Measured at least one phenotypic, physiological, phenological, or other

163                potentially fitness-related trait (e.g. survival; hereafter, trait)

164           d)   Measured the trait(s) of the populations in at least one common environment

165                (including laboratories, growth chambers, greenhouses, or outside gardens;

166                hereafter, garden).

A plant was determined to be native to the Great Basin if the taxa had at least one

occurrence with native status within the floristic Great Basin according to occurrence

information from the USDA Plants Database (USDA and NRCS, 2018) and/or the U.S Virtual

Herbarium Online (Barkworth *et al.*, 2018). A total of 170 studies published between 1941 and

July 2017 were encountered that met these criteria.

*Categorization and scoring of literature*

All studies meeting our criteria were categorized and scored for each signature. The

coordinates of all gardens and populations in each study were recorded or, if possible, generated

from localities described in the studies (Supporting Information Appendix 1). For each study, we

then noted these 15 characteristics: the year published, year(s) of plant material collection,

year(s) of experimentation, number of years reported, taxa (genus, species, subspecies), life

history traits (taxonomic status, lifeform, geographic range, life span, breeding system),

experiment type (laboratory, greenhouse, common garden, reciprocal transplant), number of

gardens, number of populations tested, which generation of material was used, and whether or

not experimenters attempted to control for maternal effects prior to testing (Supporting

Information Appendix 1). Life history traits were compiled for each taxon from the USDA Plants

Database as well as from published literature (Supporting Information Appendix 1). Each taxon

(subspecies level, if given) was entered separately for studies addressing multiple taxa. In studies

where more than one experiment was performed, and the experiments differed in the experiment

type (defined above), the identity of the populations being compared, and/or the generation of

material used, they were entered as separate experiments. In cases where the list of tested

populations was identical among multiple published studies, and these materials came from the

same collections, these experiments were entered separately if the garden type or location(s)

differed among the studies or if authors separately published different traits from the same

gardens, ensuring that no trait was recorded twice for the same set of populations in the same

garden. In cases where the list of tested populations did not completely overlap between studies,

even if some from each study arose from the same collections, they were entered separately.

These methods carefully emphasized the inclusion of the greatest number of relevant

experiments and traits without duplication, but nonetheless resulted in some non-independence

between some experiments. A total 327 taxa-specific entries (hereafter, experiments) were

generated from the 170 published studies (Supporting Information Appendix 2).

The first two expected signatures of local adaptation were scored using a Yes/No

designation for each experiment which considered all measured phenotypic traits. A score of

"Yes", or, in the absence of supporting statistical evidence, "Authors claim Yes", was given

when at least one measured trait significantly demonstrated the signature for at least two

populations, and a score of "No" or "Authors claim No" was given when the signature was not

detected between any pair of populations (Supporting Information Appendix 1). In addition, each

of the measured and reported traits and environmental variables were scored (hereafter, trait

scores) in the same way for each signature. Of the 327 experiments, 305 (93.3%) met the criteria

to score for among-population variation (signature 1) and 161 (49.5%) met the criteria to score

for trait-by-environment association (signature 2). Pearson's chi-squared tests were used to

determine if there were differences in signatures 1 and 2 among plants with different life-history

traits, using totals from both "Yes/No" and "Authors Claim Yes/No" results, excluding any life

history groups represented by less than 10 experiments.

To score whether there was higher fitness of a local population in a common garden

(hereafter, signature 3), only experiments in which outdoor reciprocal transplants or common

gardens were performed using a local population in at least one garden were considered

(Supporting Information Appendix 1). Additionally, the experiment had to measure a fitness-

relevant response: survival, reproductive output (number of seeds or flowers, or other

reproductive output), a fitness index (a combination of several size and production traits), or total

aboveground biomass. Each experiment was assigned a composite score to fully capture

variation in the performance of each garden's local population, across multiple gardens as well

as through multiple sampling dates (Supporting Information Appendix 1). The five possible

composite scores were "Yes for all gardens at all times", "Yes for all gardens at some times",

"Yes for some gardens at all times", "Yes for some gardens at some times", and "No for all

gardens at all times". These scores refer only to those gardens within each experiment that

included their own local population. Of the 326 experiments, 27 (8.3%) were appropriate for this

scoring. This scoring provides an estimate of the commonness of higher local fitness, but it is not

a measure of the importance of the difference per se. For example, a fitness difference could

occur uncommonly, but have a large impact on population trajectories (i.e. large differences in

survival after a rare drought event).

Our dataset, which had uneven numbers of experiments representing each species,

contained the possibility of bias associated with highly-studied taxa influencing patterns more

than less-studied taxa. To ask how this affected overall results, we compared tallies of all scores

without correcting for multiple experiments per species to tallies using an average score for each

species for each signature. To generate these average scores for signature 1 and 2, we totaled all

"Yes" and "Authors claim Yes" scores for each species and divided by the total number of scores

(all Ys plus all Ns) for that species. For signature 3, all forms of "Yes" (all but "No for all

gardens at all times") were totaled into a Y and divided by the total number of scores. Then, we

averaged these per-species scores to re-calculate overall effects in which each species was

represented only once, and compared the results of the different averaging methods for each

signature.

*Quantitative comparison of trait-by-environment associations*

As a complement to the survey of author-reported results described above, we conducted

a further, quantitative analysis of trait and climate values. Specifically, to examine associations

between the differences in trait values and the differences in environmental and geographic

distance among population origins, we utilized experiments from which population-specific trait

data and geographic coordinates could be extracted or obtained through author contact. Data

from laboratory and greenhouse experiments were not considered for this extraction. First, we

identified the most commonly measured traits across studies, which were then manually

extracted from text, tables, or graphical data (Supporting Information Appendix 1). Next, we

extracted trait data from the latest sampling date for which the most populations at the most

gardens were represented, and if multiple treatments were used, we only extracted data for the

author-defined 'control' treatment. However, if no control was defined, we used the treatment

that was the most unaltered or representative of the garden environment (e.g. unweeded, or

unwatered). For each population/trait combination, we used either author-provided mean values

or calculated a mean trait value from available data. Rather than averaging values across gardens,

data, data from each garden location within each experiment was extracted separately and

considered its own sample. We did this because it is not uncommon for traits to be expressed

differently in different common garden locations (e.g. Johnson, Leger and Vance-Borland,

2017). Finally, we generated 30-year annual precipitation and mean annual temperature values

for each population's location of origin using the ClimateNA v5.10 software package based on

methodology described by Wang et al. (2016). These 30-year averages are calculated every 10

years (i.e. 1951-1980, 1961-1990, etc.). Because studies took place at many times over the last

75 years, we used the most proximate climate normal for each experiment that did not include or

surpass the years during which the experiment's populations were collected (Supporting

Information Appendix 2).

To reduce the likelihood of spurious correlations or false negative results, we limited this

dataset to traits measured in at least 5 populations in at least 20 common garden locations (mean

locations per trait: 34.4; range: 21-46), resulting in 81 locations (from 56 experiments) that

measured at least one of eight frequently-measured phenotypic traits (Table 1). Within each

location, we calculated pairwise Euclidean distances for each trait value, climate factor, and

geographic distance for every possible pair of populations. Geographic distances were generated

using the earth.dist function in fossil package (Vavrek, 2011) in the statistical computing

environment R (R Core Team, 2017). Then, partial Mantel tests were used to compare pairwise

trait and climate distances for each experiment while controlling for geographic distances, using

the vegan package (Oksanen *et al.*, 2018) in R (R Core Team, 2017). We used the metacor.DSL

function in the metacor package (Laliberté, 2011) to generate an overall effect size (partial

correlation) and upper and lower confidence intervals for each combination of trait and

environmental variable. Lastly, to better understand effect sizes for a subset of species, we ran

simple linear regression analyses for each location, comparing average trait values and

environmental values to generate a slope that estimated trait change per unit change in climate

factors. Experiments with $R^2$ values of 0.2 or less were excluded from this particular analysis,

and the median slope across experiments was retained as an estimate of the trait-by-environment

relationship. The arbitrary cutoff ($R^2 = 0.2$) for this step was used simply as a way to focus on

and report effect sizes from some of the stronger biological relationships that could be of

particular interest to managers, restoration practitioners and evolutionary ecologists. Due to

limited sample sizes for factors such as lifeform, mating system, geographic distribution, etc., we

did not include these factors in any of the quantitative analyses, but present lifeform (shrub,

grass, or forb) information for each trait response as additional results in the Supporting

Information Appendix 3.

**Results**

*Summary of reviewed literature*

Our literature search revealed 170 published studies that measured trait responses from

more than one population in at least one common environment, resulting in 327 separate

experiments involving 121 taxa of 104 species of grasses, shrubs, forbs, and deciduous trees

(Fig. 1). These experiments represent approximately 3,234 unique populations tested in

approximately 208 outdoor garden locations (Fig. 2) and 154 indoor lab or greenhouse

experiments. Grasses accounted for 21.0% of the taxa and 40.2% of the experiments, forbs

composed 50.8% of the taxa and 30.7% of experiments, shrubs 26.6% of the taxa and 28.5% of

experiments, and deciduous trees accounted for only 1.6% of taxa and 0.6% of experiments (Fig.

1A). Experiments were most commonly conducted in non-reciprocal outdoor common gardens

(47.5%) or in the laboratory (31.9%), with fewer conducted in greenhouses (15.3%) or in

reciprocal outdoor gardens (5.2%, Fig. 1B). For experiments in outdoor gardens, the median

number of gardens per experiment across lifeform ranged from 1 (grasses, shrubs, and trees) to 2

(forbs) for non-reciprocal gardens, and from 2 (grasses and forbs) to 4 (shrubs) for reciprocal

gardens . Overall, the median number of populations tested in each experiment was 5 (range= 2 -

193, IQR =  3 – 11.5, Fig. 1C), and was slightly lower for shrubs (median = 4, range = 2 – 111,

IQR = 2 - 8) than grasses (median = 6, range = 2 – 193, IQR = 3 - 12.25), forbs (median = 6,

range = 2 – 67, IQR = 3 – 10.25), and trees (median = 7, range = 5 – 9, IQR = 6 – 8).

Experiments took place between 1940 and 2015, with collections from native stands

occurring between 1938 and 2013 (Fig. 3A). One quarter of the experiments (24.5%) reported

only early germination and seedling stages of plants (generally less than 0.5 years), while the

remaining experiments (75.5%) reported study periods ranging from 0.5 to 17 years, with an

average of 2.1 years (Fig. 3B, C). Average pairwise geographic distance among populations per

experiment for the 91% of experiments for which coordinates were available was 351 km$\pm$20

SE, with a range from 610 m to 2,551 km.  Most experiments were conducted on taxa with

regional distributions, perennial species, grasses, and outcrossing species; very few annuals,

endemic species, or selfing species were represented (Fig. 4). Over half of experiments (58.6%)

tested plants grown directly from wild-collected seeds (or the seed of wild collected adults),

16.9% tested wild-collected adults, 13% tested materials with mixed generations since collection,

6.7% tested 1st or 2nd generation descendants of wild collected seeds, 0.3% tested only cultivars,

and 4.3% did not provide enough information to determine.

*Among-population variation*

Of the 305 experiments appropriate for addressing among-population trait variation

(signature 1), 290 (95.1%) experiments reported finding variation among populations in at least

one phenotypic trait, with 230 (75.4%) of these 290 reporting significant variation, and 60

(19.6%) claiming such variation in the absence of any supporting statistics (Fig. 4A). Only 12

(3.9%) experiments reported no such differentiation in any trait after statistically testing for it,

and 3 (1%) claimed no such variation without presenting statistical evidence. When categorized

by basic life history traits, several differences appeared among groups. Eudicots exceeded

monocots (the majority of which were grasses) in the degree of population differentiation($X_1^2 =$

7, $P = 0.0081$), and, similarly, forbs and shrubs had more population differentiation than grasses

($X_2^2 = 8.05$, $P = 0.0143$). There were no significant differences in signature 1 among plants with

different geographic distributions, life span, or breeding systems.

A total of 1,465 trait scores were recorded from the 305 experiments appropriate for

addressing signature 1. Frequently-measured traits (20 or more experiments) that had differences

between populations in over 75% of experiments (with or without supporting statistics) were

floral structure, vigor, emergence, plant size, number of leaves, plant structure, shoot biomass,

leaf structure, and number of inflorescences (Fig. 5).

*Trait-by-environment associations*

Of the 161 experiments appropriate for testing trait-by-environment associations

(signature 2), 131 (81.4%) reported associations for at least one comparison, with 81 (50.3%)

supported by statistical tests and 50 (31.1%) supported by claims in the absence of statistics (Fig.

4B). Conversely, 13 (8.1%) of experiments reported no such correlations after having

statistically tested for it, and 17 (10.6%) reported no such correlations but lacked any supporting

statistics. There were no significant differences in the commonness of trait-by-environment

associations for taxonomic status, lifeform, geographic distribution, or breeding system, but

perennials (both long-lived and short-lived) had more frequent correlations between traits and

environment than did annuals or short-lived perennials ($X_3^2 = 8.08$, $P = 0.0444$).

A total of 592 trait scores were recorded from the 161 experiments appropriate for

addressing signature 2 (Fig. 6A). Frequently-measured traits (20 or more experiments) that were

correlated with environmental variables in over 75% of experiments (with or without supporting

statistics) were multivariate trait axes, floral structure, and germination date. Every remaining

trait that was measured in >15 experiments was correlated with environmental characteristics in

over 50% of experiments, and many, including leaf length, survival, flowering date, and leaf

structure, were correlated with environmental variables in ≥70% of experiments.

A total of 426 environmental variable scores were recorded from the 161 experiments

appropriate for addressing signature 2 (Fig. 6B). Of the variables most frequently reported as

correlated with plant traits, many categorical variables or composite metrics made this list, with

seed zones, ecoregions, multivariate environmental axes, and habitat classifications topping the

list of important environmental variables (important in > 84% of experiments that reported

them). Additionally, derived climate metrics (such as climate continentality, heat/moisture index,

potential evapotranspiration, etc.), climate seasonality, and history of invasive species presence

were correlated with plant traits in over 75% of studies that reported them.

*Higher local performance in a local common garden*

The 27 experiments that were suitable for detecting higher fitness of a local population in

a local garden (signature 3) generated 39 scores (some experiments measured multiple fitness

traits), with 27 scores (69.2%) reporting signature 3 for at least one fitness trait in at least one of

the tested gardens during at least one sampling date, and the remaining 12 scores (30.8%) not

reporting signature 3 at any point (Fig. 4C). Thirty-two of the 39 scores (82%) were generated

from experiments with more than one garden. Survival was the most frequently measured fitness

trait in these experiments, reported in 24 of the 27 experiments, followed by reproduction (10),

biomass (3), and fitness indices (2). Incidence of the local-does-best pattern was highest in

experiments that directly measured reproductive output, with 90% reporting higher values for

locals at some point in an experiment, followed by survival (67%), fitness indices that

incorporated biomass (50%), and biomass measures (33%). For experiments in which only

"some" gardens showed local-does-best patterns (Fig. 4C, hashed bars), the percentage of

gardens showing this trend was 40%, 50%, and 40% for reproduction, survival, and biomass

traits, respectively (not shown). For experiments in which only "some" sampling dates showed

local-does-best patterns (gray bars), the percentage of sampling dates showing this trend was

56%, 47%, and 25% for reproduction, survival, and biomass traits, respectively (not shown).

*Considering possible biases: highly-studied species and maternal effects*

The number of experiments per species in our dataset ranged from 1 (52 species) to 25

(*Artemisia tridentata*), with a median of 1 (IQR = 1 – 4). The most highly-represented species

were *Artemisia tridentata* (25 experiments), *Elymus elymoides* (24), *Ericameria nauseosa* (17),

*Achnatherum hymenoides* (17), *Krascheninnikovia lanata* (13), *Pascopyrum smithii* (11),

*Atriplex canescens* (9), *Leymus cinereus* (9), and *Poa secunda,* (8). Results in which scores were

averaged for each species (see methods) were similar to uncorrected results: signature 1 was 4%

higher when corrected (98% vs. 94%), signature 2 was 1% lower when corrected (79% vs. 80%),

and signature 3 was 8% higher when corrected (78% vs. 70%). Thus, uncorrected calculations

were used throughout our study.

Only 19 experiments (5.8%) used an experimental design that could control for maternal

effects (e.g. growing all populations for a generation in a common environment before initiating

an experiment). An additional 30 experiments (9.2%) were unclear on this point, and the

remaining 278 (85%) experimented directly on populations differing in maternal environment.

The incidence of population differences (signature 1) was 100% in the 16 experiments that

moderated maternal effects, 95% for the 259 that did not make an attempt, and 97% for the 30

which were unclear. Too few of the experiments that attempted to control for maternal effects

were appropriate for measuring signature 2 (4 experiments) and signature 3 (1 experiment) to

compare incidences of these signatures.

*Quantitative comparison of trait-by-environment associations*

399            Overall, we found positive relationships between the magnitude of differences among

populations in all eight phenotypic traits and the magnitude of differences between MAT and

MAP at the collection locations (Fig. 7). The strongest relationship was observed between

differences in flowering time and differences in MAT, and leaf size also showed a strong

relationship with MAT. Multiple strong relationships were observed between trait/environment

divergence for MAP, with leaf size, survival, shoot mass, inflorescence number, and flowering

time all showing strongly positive relationships for grasses, forbs, and shrubs. (Fig. 7, Supporting

Information Appendix 3). Regression analyses demonstrated that, for the 15 common garden

locations in which strong flowering time and MAT relationships were observed, each degree

change in MAT was associated with a median change of 3.5 days (IQR = 1.2 - 5.3) in flowering

time. Small sample sizes (few experiments that could be included in the analyses) and challenges

with interpreting changes in physical traits across species of various shapes and sizes precluded

the presentation of estimates of this nature for the other trait-by-environment relationships.

**Discussion**

Our results represent the most extensive review of intraspecific variation and local

adaptation for plants native to the floristic Great Basin, a region comprised of largely continuous

but increasingly imperiled arid and semi-arid plant communities (Davies *et al.*, 2011; Finch *et*

*al.*, 2016). Additionally, they represent a significant addition to the noteworthy though relatively

small number of reviews investigating this topic in a manner that identifies individual traits and

environmental factors involved. We found that Great Basin plant species contain large amounts

of intraspecific diversity in a wide range of phenotypic traits, that differences in these phenotypic

traits are often associated with the heterogeneous environments of origin, and that differences

among populations are commonly relevant to outplanting fitness. The cascading importance of

intraspecific variation for the structure, functioning, and biodiversity of communities and

ecosystems can be considerable (Bolnick *et al.*, 2011; Bucharova *et al.*, 2016), and may equal or

exceed the importance of species diversity (Des Roches *et al.*, 2018). Our quantification of local

adaptation and trait-environment associations should serve as encouragement to seriously

consider intraspecific diversity in native plant materials used in restoration and conservation in

this region throughout the selection, evaluation, and development process (Basey, Fant and

Kramer, 2015). The results reported here should also serve as a cautionary note to restoration

approaches that focus on only a few specific traits or search for general-purpose genotypes. Our

results suggest that, in the absence of species-specific information to the contrary, it is reasonable

to assume that local adaptation is present in this region, and that locally-sourced populations

would outperform non-local populations a majority of the time.

Our investigation encompassed 170 studies published between 1941 and 2017 in which

over 3,230 unique populations of 104 native Great Basin plant species were compared in 327

experiments, ranging from laboratory germination trials to multiple-year common gardens and

reciprocal transplants. The great majority (95%) found differences between populations

(signature 1) in the majority of traits measured in a common environment, which indicates that

different traits are variable among populations, at both small and large geographic scales.

Additionally, a clear majority (81.4%) of experiments found trait-by-collection environment

associations (signature 2), suggesting that intraspecific variation is frequently an adaptive

outcome of natural selection in heterogeneous environments (Linhart and Grant, 1996; Reich *et*

*al.*, 2003). In experiments suitable for detecting local performance advantages (signature 3), local

populations had higher performance (measured by differences in reproductive output, survival,

and biomass) than nonlocal populations more often than not (69.2%), and this was particularly

true when researchers reported traits related to reproductive output (90%). We used a vote-

counting method to summarize results for our broadest pool of studies, allowing us to

incorporate a wealth of older studies for which quantitative details were not available. Results

from a vote-counting approach can sometimes differ from results of meta-analysis, as vote-

counting does not incorporate the same level of detail about factors such as study size or effect

size (Combs *et al.*, 2011). However, in our study, the overall incidence of "local does best" in the

Great Basin is similar to other reviews that have found local adaptation to be commonplace, but

not ubiquitous. In a review of local adaptation in plants that compared survival, reproduction,

biomass and germination traits in reciprocal transplants, Leimu and Fischer (2008) found that

local plants outperformed non-local ones in 71% of 35 published experiments. Similarly,

Hereford (2009) quantified local adaptation in 70 published studies (50 of them plants), reporting

only survival or reproductive traits, and found evidence of local adaptation in 65-71% of

experiments. Our results indicated that the strongest indication of local adaptation came from

experiments that directly measured reproductive output, and that using biomass as a fitness proxy

may not be an effective way to compare relative performance in the Great Basin. This is

consistent with a previous study that demonstrated selection for smaller, rather than larger,

individuals in disturbed arid systems (Kulpa and Leger, 2013). Literature reviews conducted

across biomes may occlude regionally-important trait differentiation and mask patterns of local

adaptation, as we might expect, for example, biomass to be more strongly linked to fitness in

regions where light is a contested resource (Espeland, Johnson and Horning, 2017).

There are many processes that can reduce or prevent the development of local adaptation,

such as the lack of divergent selection between sites, high gene flow, rapid or extreme

environmental change, high phenotypic plasticity, and/or low genetic diversity (Sultan and

Spencer, 2002; Kawecki and Ebert, 2004; Blows and Hoffmann, 2005). The high incidence of

intraspecific variation, much of it habitat-correlated, that we found in the literature confirms that

divergent selection by heterogeneous environments is the norm for species native to the Great

Basin, presumably outweighing the balancing effects of gene flow and genetic drift. Key

environmental factors in the Great Basin such as fire frequency, grazing regimes, resource

availability, and climate are certainly being altered to varying degrees by invasive species

introductions, changing land uses, and climate change, and it can be argued that such changes

could outpace the ability of local populations to remain adapted to their surroundings (Jones and

Monaco, 2009; Breed *et al.*, 2013; Havens *et al.*, 2015; Kilkenny, 2015). However, our analysis

also demonstrated relatively high instances of trait correlations with relatively recent

disturbances such as invasive species introductions. Rapid evolution in response to invasive

species (Oduor, 2013) and other anthropogenic changes (Hoffmann and Sgrò, 2011; Franks,

Weber and Aitken, 2014) has been documented for many species, indicating that local adaptation

can evolve rapidly in some circumstances.

482   Some traits and environmental characteristics stood out as particularly important

483 indicators of local adaptation and its signatures across the studied taxa. For example, in our

484 quantitative comparison of divergence in traits and environments, flowering phenology was

485 strongly affected by MAT, with a median change of 3.5 days in flowering time per degree

486 change in MAT of collection origin. Flowering phenology, along with germination phenology,

487 were also in the top tier of frequently measured traits that showed significant correlations with

488 environmental variables, consistent with other studies that have shown reproductive (Bucharova,

489 Michalski, *et al.*, 2017) and germination (Donohue *et al.*, 2010) phenology to be an important

490 response to environmental variation. Leaf size is also an important adaptive response to

491 differences in temperature globally (Wright *et al.*, 2017), and in concert with this, we saw overall

492 positive responses to MAP and MAT for leaf size in our analyses as well as frequent trait-by-

493 environment associations in the literature. Floral structure, which has important adaptive

494 significance for angiosperms (Harder and Barrett, 2007; Armbruster, 2014), was among the most

495 frequent traits scored for among-population variation and trait-by-environment interactions.

496 Seasonality of precipitation, which varies in this region depending on summer rainfall

497 (Comstock and Ehleringer, 1992), was more predictive of trait variation overall than was mean

498 annual precipitation (signature 2). In our quantitative comparisons, differences in MAP values

499 were important for multiple phenotypic traits, including leaf size, shoot mass, reproductive

500 output, and flowering phenology, in addition to being important for overall plant survival. Larger

501 scale environmental descriptors, such as ecoregions and seed transfer zones, universally

502 demonstrated signature 2, likely because they were developed based on climate/soil/vegetation

503 associations or, in the case of seed transfer zones, developed based on trait-by-environment

504 correlations. As found in other reviews (Geber and Griffen, 2003), physiological traits,

505    phytochemical traits, and root traits were not measured as frequently as other traits, and though

506    these did not show as frequent associations with environmental characteristics as other traits,

507    they are known to vary across environments in some systems (Reich *et al.*, 2003). Additional

508    studies of these traits in the Great Basin would be informative and could reveal different patterns

509    than those observed here.

510           As in any review and analysis of published papers, there are elements of our design that

were difficult to control. For example, consistent with other reviews (Gibson *et al.*, 2016), the

vast majority of studies involved wild-collected plants or seeds, and thus maternal environment

effects almost certainly affected some results (e.g. Bischoff and Müller-Schärer, 2010; Espeland

*et al.*, 2016). Additionally, though the majority of populations tested in the literature were from

western states, some of the populations compared in the literature were collected from well

outside of the Great Basin, which increased the likelihood of observing local adaptation in these

species. However, understanding patterns of intraspecific variation across the full range of the

species native to the Great Basin is pertinent because it has been common (and for some species,

ubiquitous) to utilize sources of native species originating from outside the Great Basin to use for

restoration within the Great Basin (Jones and Larson, 2005). Finally, the scores and percentages

for each of the signatures used throughout this study are uncorrected for phylogeny, as is our

pairwise trait/environment analysis, and calculated such that each experiment is weighed equally.

This introduces the possibility for phylogenetic biases, in which closely related taxa represented

by many experiments affect the results more than less frequently studied taxa or groups of taxa.

Though we did not conduct phylogenetic corrections for relatedness among taxa (Harvey and

Pagel, 1991; de Bello *et al.*, 2015), our results were essentially identical for signatures 1-3 when

we averaged results across species (scores differed by +3%, -1%, and +8%, respectively),

suggesting that our lack of phylogenetic corrections are not unduly affecting our results. We

present all species-specific information in Supporting Information Appendix 2 and available

datasets section of the electronic supplementary material for further review.

Current approaches to seed sourcing in restoration and conservation include genetic (e.g.

Williams, Nevill and Krauss, 2014), genecological (e.g. Johnson, Leger and Vance-Borland,

2017), local-only (e.g. Erickson *et al.*, 2017), predictive (e.g. Prober *et al.*, 2015), and agronomic

(e.g. United States. House of Representatives. Committee on Appropriations., 2014)) strategies,

as well as strategies mixing several of these viewpoints (i.e. Rice and Emery, 2003; Rogers and

Montalvo, 2004; Breed *et al.*, 2013; Havens *et al.*, 2015; Bucharova *et al.*, 2018). These

approaches vary in the degree to which they meet the needs of seed producers and land managers

while balancing population differences that stem from adaptive evolution in different

environments. The prevalence of local adaptation and its signatures found in our study justify

and support incorporating existing best-practices (e.g. Basey, Fant and Kramer, 2015; Espeland

*et al.*, 2017) for capturing and preserving important intraspecific variation into seed sourcing and

plant production systems. For example, our results demonstrated a strong relationship between

flowering time and MAT, so it would be wise to collect materials for research, evaluation, and

testing from populations that vary in MAT, to collect seeds at multiple times to fully capture

population variation in flowering time, and ensure that seeds are not transferred during

restoration among sites that differ strongly in these characteristics. On the production side, best

practices for seed harvesting should include methods that avoid inadvertent selection on

flowering time, either for reduced variation or for a directional shift away from the wild

condition. Similarly, emergence date was correlated with environmental variation in many

plants, so testing in common gardens should involve seeding trials in place of or in addition to

using transplants, and evaluation trials should guard against inadvertent selection on emergence

timing by randomly, rather than systematically, selecting individuals to use in transplant

experiments. These examples are not exhaustive, but demonstrate how evidence revealed by this

study regarding which traits and environmental factors are generally involved in adaptation in

this region can be used to improve approaches to seed sourcing and restoration. Finally, we

acknowledge that ours is not the first review and meta-analysis to affirm an abundance of

intraspecific variation and local adaptation in plants. However, our focus on the Great Basin is

important, because the large and frequent yet commonly unsuccessful restoration efforts

occurring in this region have lagged behind those of other regions with respect to recognizing the

importance of intraspecific variation and local adaptation on outplanting success.

**Conclusions**

Reestablishing and maintaining native plant communities in arid regions has proven

challenging (Svejcar *et al.*, 2017), and the lack of practical knowledge guiding more appropriate

selection of seed sources is a major barrier (Friggens *et al.*, 2012; Gibson *et al.*, 2016). The

forestry industry has long adopted the principles of local adaptation in their reforesting

guidelines with great success (Matyas, 1996; Johnson *et al.*, 2004; Aitken and Bemmels, 2016),

and similar approaches to restoration in the rangelands of the Great Basin may also increase

success as our data support similarly high levels of population differentiation within grass, forb

and shrub life history groups. Our results, including both a qualitative literature survey and a

quantitative meta-analysis, could benefit from future work using additional techniques to explore

spatial structure (e.g. Griffith and Peres-Neto, 2006) and the relative importance of geographic

distance and environmental variation, especially as additional studies become available in the

literature. Nevertheless, our results as they currently stand are in agreement with observations of

abundant local adaptation in plant populations world-wide, and further, we identified particular

phenotypic traits (flowering and germination phenology, floral structures, leaf size, biomass,

survival, and reproductive output), environmental characteristics (MAT, MAP, climate metrics,

seasonality), and habitat classifications and site history (seed zones, ecoregions, history of

invasive species) that were important predictors of local adaptation in plants native to the Great

Basin floristic region. Given the speed and severity with which natural communities are being

altered by anthropogenic factors, the application of an evolutionary perspective to restoration

ecology is more important than ever. Adjusting seed-selection priorities to account for the

existence of locally adapted, intraspecific variation in the Great Basin will promote the

maintenance and recovery of resilient, self-sustaining vegetation communities in this region

(Meyer, 1997; Lesica and Allendorf, 1999; Rogers and Montalvo, 2004; Broadhurst *et al.*, 2008;

Vander Mijnsbrugge, Bischoff and Smith, 2010).

**Acknowledgements**

We would like to acknowledge Vicki Thill and Sage Ellis for many hours of extracting

coordinates and trait data, as well as Susan Meyer, Andrea Kramer, Tom Jones, Vicky Erickson,

Allan Stevens, Dan Atwater, Clinton Shock, Jessica Irwin, Huixuan Liao, and David Solance

Smith for providing information and data not available in their publications. We also thank

several anonymous reviewers for their insightful suggestions that greatly improved this work.

**Dedication**

We would like to dedicate this paper to the memory of our co-author Dr. Erin K.

Espeland, friend and collaborator to all of us, who worked on this manuscript. Erin's light and

life will never be forgotten by those who knew her, and we want to recognize her creative

contributions to the field of plant ecology, including this effort. Erin is dearly missed.

**Funding**

This project was funded by a grant from the United States Department of the Interior

Great Basin Landscape Conservation Cooperative (2016-Kilkenny/Leger), which provided

support for EAL and OWB. Additionally, ACA was supported by a grant from the United States

Department of Agriculture National Institute of Food and Agriculture; EE was supported by

United States congressional appropriation 3032-21220-002-00-D; FFK and JEO were supported

by the Great Basin Native Plant Project, the United States Department of the Interior Bureau of

Land Management, and the United States Department of Agriculture Forest Service; JBS and

MEH were supported by the United States Department of Agriculture Forest Service; MLF was

supported by a Trevor James McMinn fellowship; RCJ was supported by the Great Basin Native

Plant Project; RF was supported by the Institute for Applied Ecology; and TMK was supported

by the United States Department of the Interior Bureau of Land Management and the Institute for

Applied Ecology

**Data accessibility**

Raw datasets and statistical code supporting this study (Baughman *et al.*, In Review) have been

deposited at Dryad, [DOI: TBD]

**Authors' contributions**

EAL, OWB, FFK, EKE, RF, TNK, and JBS conceived and designed the study; OWB

conducted the literature search; OWB, ACA, FFK, JEO, RCJ, and JBS categorized, compiled

and extracted data; OWB, EAL, FFK, ACA and MLF analyzed data; OWB, EAL, and ACA

drafted the manuscript; all authors critically revised the manuscript for important intellectual

content and approved of the version to be published.

**References**

Ackerly, D. D. *et al.* (2000) 'The Evolution of Plant Ecophysiological Traits : Recent Advances

and Future Directions', *AIBS Bulletin*, pp. 979–995. doi: 10.1641/0006-

3568(2000)050[0979:teopet]2.0.co;2.

Aitken, S. N. and Bemmels, J. B. (2016) 'Time to get moving: Assisted gene flow of forest

trees', *Evolutionary Applications*, pp. 271–290. doi: 10.1111/eva.12293.

Anderson, J. T. *et al.* (2012) 'Phenotypic plasticity and adaptive evolution contribute to

advancing flowering phenology in response to climate change', *Proceedings of the Royal Society*

*of London B: Biological Sciences*. doi: 10.1098/rspb.2012.1051.

Antonovics, J. and Bradshaw, A. (1968) 'Evolution in closely adjacent plant populations',

*Heredity*, 23(4), pp. 507–524. doi: 10.1038/hdy.1970.36.

Armbruster, W. (2014) 'Floral specialization and angiosperm diversity: phenotypic divergence,

fitness trade-offs and realized pollination accuracy', *AoB Plants*, 6.

Barkworth, M. *et al.* (2018) 'US Virtual Herbarium'. Available at: http://usvhproject.org/#/.

Barnes, M. (2009) 'The effect of plant source location on restoration success: a reciprocal

transplant experiment with winterfat (Krascheninnikovia lanata)', *Doctoral Dissertation. The*

*University of New Mexico*. Available at: http://digitalrepository.unm.edu/biol_etds/4 (Accessed:

8 January 2018).

Basey, A. C., Fant, J. B. and Kramer, A. T. (2015) 'Producing native plant materials for

restoration: 10 rules to collect and maintain genetic diversity', *Native Plants Journal*, 16(1), pp.

37–53. doi: 10.3368/npj.16.1.37.

Baughman, O. *et al.* (In Review) 'Data from: Strong Patterns of Intraspecific Variation and

Local Adaptation in Great Basin Plants Revealed Through 75 Years of Experiments', *Dryad*

*Digital Repository*. doi: TBD.

de Bello, F. *et al.* (2015) 'On the need for phylogenetic "corrections" in functional trait-based

approaches', *Folia Geobotanica*, 50, pp. 349–357. doi: 10.1007/s12224-015-9228-6.

Bischoff, A. and Müller-Schärer, H. (2010) 'Testing population differentiation in plant species -

How important are environmental maternal effects', *Oikos*, 119(3), pp. 445–454. doi:

10.1111/j.1600-0706.2009.17776.x.

Blanquart, F. *et al.* (2013) 'A practical guide to measuring local adaptation', *Ecology Letters*, pp.

1195–1205. doi: 10.1111/ele.12150.

Blows, M. W. and Hoffmann, A. A. (2005) 'A reassessment of genetic limits to evolutionary

change', *Ecology*, pp. 1371–1384. doi: 10.1890/04-1209.

Bolnick, D. I. *et al.* (2011) 'Why intraspecific trait variation matters in community ecology',

*Trends in Ecology and Evolution*. doi: 10.1016/j.tree.2011.01.009.

Breed, M. F. *et al.* (2013) 'Which provenance and where? Seed sourcing strategies for

revegetation in a changing environment', *Conservation Genetics*, 14(1), pp. 1–10. doi:

10.1007/s10592-012-0425-z.

Broadhurst, L. M. *et al.* (2008) 'Seed supply for broadscale restoration: Maximizing evolutionary

potential', *Evolutionary Applications*, 1(4), pp. 587–597. doi: 10.1111/j.1752-

4571.2008.00045.x.

Bucharova, A. *et al.* (2016) 'Plant ecotype affects interacting organisms across multiple trophic

levels', *Basic and Applied Ecology*. doi: 10.1016/j.baae.2016.09.001.

Bucharova, A., Durka, W., *et al.* (2017) 'Are local plants the best for ecosystem restoration? It

depends on how you analyze the data', *Ecology and Evolution*, 7(24), pp. 10683–10689. doi:

10.1002/ece3.3585.

Bucharova, A., Michalski, S., *et al.* (2017) 'Genetic differentiation and regional adaptation

among seed origins used for grassland restoration: lessons from a multispecies transplant

experiment', *Journal of Applied Ecology*, 54(1), pp. 127–136. doi: 10.1111/1365-2664.12645.

Bucharova, A. *et al.* (2018) 'Mix and match: regional admixture provenancing strikes a balance

among different seed-sourcing strategies for ecological restoration', *Conservation Genetics*. doi:

10.1007/s10592-018-1067-6.

Chivers, I. H. *et al.* (2016) 'The merits of artificial selection for the development of restoration-
ready plant materials of native perennial grasses', *Restoration Ecology*, 24(2), pp. 174–183. doi:
10.1111/rec.12323.

Clausen, J., Keck, D. and Hiesey, W. (1948) 'Experimental studies on the nature of species. III.
Environmental responses of climatic races of Achillea', *Carnegie Institution of Washington*, 581,
pp. 1–129.

Combs, J. G. *et al.* (2011) 'Assessing Cumulative Evidence within "Macro" Research: Why
Meta-Analysis Should be Preferred Over Vote Counting', *Journal of Management Studies*. doi:
10.1111/j.1467-6486.2009.00899.x.

Comstock, J. P. and Ehleringer, J. R. (1992) 'Correlating genetic variation in carbon isotopic
composition with complex climatic gradients.', *Proceedings of the National Academy of*
*Sciences*, 89(16), pp. 7747–7751. doi: 10.1073/pnas.89.16.7747.

Davies, K. W. *et al.* (2011) 'Saving the sagebrush sea: An ecosystem conservation plan for big
sagebrush plant communities', *Biological Conservation*, 144(11), pp. 2573–2584. doi:
10.1016/j.biocon.2011.07.016.

Dennison, P. E. *et al.* (2014) 'Large wildfire trends in the western United States, 1984-2011',
*Geophysical Research Letters*, 41(8), pp. 2928–2933. doi: 10.1002/2014GL059576.

Donohue, K. *et al.* (2010) 'Germination, postgermination adaptation, and species ecological
ranges', *Annual Review of Ecology, Evolution, and Systematics*, 41, pp. 293–319.

Endler, J. A. (1986) *Natural selection in the wild*, *Monographs in Population Biology*. doi:
10.2307/302397.

Erickson, T. E. *et al.* (2017) 'Benefits of adopting seed-based technologies for rehabilitation in
the mining sector: a Pilbara perspective', *Australian Journal of Botany*. doi: 10.1071/BT17154.

Erickson, V. J., Mandel, N. L. and Sorenson, F. C. (2004) 'Landscape patterns of phenotypic
variation and population structuring in a selfing grass, Elymus glaucus (blue wildrye)', *Canadian*
*Journal of Botany*, 82(12), pp. 1776–1789. doi: 10.1139/B04-141.

Espeland, E. K. *et al.* (2016) 'Perennial grass cultivars grown at different production farms

respond differently to storage and planting environments', *Crop Science*, 56, pp. 249–258. doi:

10.2135/cropsci2015.05.0318.

Espeland, E. K. *et al.* (2017) 'Evolution of plant materials for ecological restoration: Insights

from the applied and basic literature', *Journal of Applied Ecology*, 54, pp. 102–115. doi:

10.1111/1365-2664.12739.

Espeland, E. K., Johnson, R. C. and Horning, M. E. (2017) 'Plasticity in native perennial grass

populations: Implications for restoration', *Evolutionary Applications*, 00, pp. 1–10. doi:

10.1111/eva.12560.

Evans, R. A. and Young, J. A. (1990) 'Survival and Growth of Big Sagebrush (Artemisia

tridentata) Plants in Reciprocal Gardens', *Weed Science*, 38(3), pp. 215–219. doi:

10.2307/4045014.

Finch, D. M. *et al.* (2016) 'Conservation and Restoration of Sagebrush Ecosystems and Sage-

Grouse: An Assessment of USDA Forest Service Science', *General technical report*, RMRS-

GTR-3(US Department of Agriculture, Forest Service, Rocky Mountain Research Station).

Available at: https://www.fs.fed.us/rm/pubs/rmrs_gtr348.pdf.

Franks, S. J., Weber, J. J. and Aitken, S. N. (2014) 'Evolutionary and plastic responses to climate

change in terrestrial plant populations', *Evolutionary Applications*, 7(1), pp. 123–139. doi:

10.1111/eva.12112.

Friggens, M. M. *et al.* (2012) 'Decision support: Vulnerability, conservation, and restoration

(Chapter 8)', *USDA Forest Service, Rocky Mountain Research Station - General Technical*

*Report*, (RMRS-GTR-285), pp. 116–139. Available at:

http://www.scopus.com/inward/record.url?eid=2-s2.0-

84865151549&partnerID=40&md5=7e0d86d3567ac2ed6c834c0d831cbdca.

Geber, M. A. and Griffen, L. R. (2003) 'Inheritance and Natural Selection on Functional Traits',

*International Journal of Plant Sciences*, 164(S3), pp. S21–S42. doi: 10.1086/368233.

Gibson, A. L. *et al.* (2016) 'Can local adaptation research in plants inform selection of native

plant materials? An analysis of experimental methodologies', *Evolutionary Applications*, 9(10),

pp. 1219–1228. doi: 10.1111/eva.12379.

Griffith, D. A. and Peres-Neto, P. R. (2006) 'Spatial modeling in ecology: The flexibility of

eigenfunction spatial analyses', *Ecology*. doi: 10.1890/0012-

9658(2006)87[2603:SMIETF]2.0.CO;2.

Harder, L. D. and Barrett, S. C. H. (2007) *Ecology and evolution of flowers*. New York: Oxford

University Press.

Harvey, P. H. and Pagel, M. D. (1991) 'The comparative method in evolutionary biology',

*Oxford Series in Ecology and Evolution*, p. 239. doi: 10.1016/0169-5347(92)90117-T.

Havens, K. *et al.* (2015) 'Seed Sourcing for Restoration in an Era of Climate Change', *Natural*

*Areas Journal*, 35(1), pp. 122–133. doi: 10.3375/043.035.0116.

Hereford, J. (2009) 'A Quantitative Survey of Local Adaptation and Fitness Trade- Offs', *The*

*American Naturalist*, 173(5), pp. 579–588. doi: 10.1086/597611.

Hoffmann, A. A. and Sgrò, C. M. (2011) 'Climate change and evolutionary adaptation.', *Nature*,

470(7335), pp. 479–485. doi: 10.1038/nature09670.

Johnson, G. *et al.* (2004) 'Pacific Northwest Forest Tree Seed Zones: A Template for Native

Plants?', *Native Plants Journal*, 5, pp. 131–140. doi: 10.2979/NPJ.2004.5.2.131.

Johnson, R. *et al.* (2010) 'What are the best seed sources for ecosystem restoration on BLM and

USFS lands?', *Native Plants Journal*, 11(2), pp. 117–131. doi: 10.2979/NPJ.2010.11.2.117.

Johnson, R. C., Leger, E. A. and Vance-Borland, K. (2017) 'Genecology of Thurber's

Needlegrass (Achnatherum thurberianum [Piper] Barkworth) in the Western United States',

*Rangeland Ecology & Management*. the Society for Range Management, 70(4), pp. 509–517.

doi: 10.1016/j.rama.2017.01.004.

Jones, T. A. and Larson, S. R. (2005) 'Status and use of important native grasses adapted to

sagebrush communities', in *Shaw, Nancy L.; Pellant, Mike; Monsen, Stephen B., comps. Sage-*

*grouse habitat restoration symposium proceedings; 2001 June 4-7, Boise, ID. Proc. RMRS-P-38.*

Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research

Station., pp. 49–55.

Jones, T. A. and Monaco, T. A. (2009) 'A role for assisted evolution in designing native plant

materials for domesticated landscapes', *Frontiers in Ecology and the Environment*, 7(10), pp.

541–547. doi: 10.1890/080028.

Jones, T. A., Monaco, T. A. and Rigby, C. W. (2015) 'The potential of novel native plant

materials for the restoration of novel ecosystems', *Elementa: Science of the Anthropocene*, 3.

doi: 10.12952/journal.elementa.000047.

Kawecki, T. and Ebert, D. (2004) 'Conceptual issues in local adaptation', *Ecology Letters*, 7(12),

pp. 1225–1241.

Kilkenny, F. F. (2015) 'Genecological Approaches to Predicting the Effects of Climate Change

on Plant Populations', *Natural Areas Journal*, 35(1), pp. 152–164. doi: 10.3375/043.035.0110.

Kramer, A. T., Larkin, D. J. and Fant, J. B. (2015a) 'Assessing potential seed transfer zones for

five forb species from the Great Basin Floristic Region, USA', *Natural Areas Journal*, 35(1).

doi: 10.3375/043.035.0119.

Kramer, A. T., Larkin, D. J. and Fant, J. B. (2015b) 'Assessing Potential Seed Transfer Zones for

Five Forb Species from the Great Basin Floristic Region, USA', *Natural Areas Journal*, 35(1),

pp. 174–188. doi: 10.3375/043.035.0119.

Kulpa, S. M. and Leger, E. A. (2013) 'Strong natural selection during plant restoration favors an

unexpected suite of plant traits', *Evolutionary Applications*, 6(3), pp. 510–523. doi:

10.1111/eva.12038.

Laliberté, E. (2011) 'metacor: Meta-analysis of correlation coefficients', *R package 1.0-2*.

Available at: https://cran.r-project.org/package=metacor.

Langlet, O. (1971) 'Two Hundred Years Genecology', *Taxon*. International Association for Plant

Taxonomy (IAPT), 20(5/6), pp. 653–721. doi: 10.2307/1218596.

Larsen, E. C. (1947) 'Photoperiodic Responses of Geographical Strains of Andropogon

scoparius', *Botanical Gazette*, 109(2), pp. 132–149. doi: 10.1086/335463.

Leger, E. A. and Baughman, O. W. (2015) 'What seeds to plant in the Great Basin? Comparing
traits prioritized in native plant cultivars and releases with those that promote survival in the
field', *Natural Areas Journal*, 35(1), pp. 54–68. doi: 10.3375/043.035.0108.

Leimu, R. and Fischer, M. (2008) 'A meta-analysis of local adaptation in plants', *PLoS ONE*,
3(12), p. e4010. doi: 10.1371/journal.pone.0004010.

Lesica, P. and Allendorf, F. W. (1999) 'Ecological Genetics and the Restoration of Plant
Communities: Mix or Match?', *Restoration Ecology*, 7(1), pp. 42–50. doi: 10.1046/j.1526-
100X.1999.07105.x.

Linhart, Y. B. and Grant, M. C. (1996) 'Evolutionary Significance of Local Genetic
Differentiation in Plants', *Annual Review of Ecology and Systematics*, 27(1), pp. 237–277. doi:
10.1146/annurev.ecolsys.27.1.237.

Loveless, M. D. and Hamrick, J. L. (1984) 'Ecological Determinants of Genetic Structure in
Plant Populations', *Annual Review of Ecology and Systematics*, 15(1), pp. 65–95. doi:
10.1146/annurev.es.15.110184.000433.

Matyas, C. (1996) 'Climatic adaptation of trees: rediscovering provenance tests', *Euphytica*, Jan
1(92 (1-2)), pp. 45–54.

McArthur, E. D., Meyer, S. E. and Weber, D. J. (1987) 'Germination rate at low temperature:
Rubber rabbitbrush population differences', *Journal of Range Management*, 40(6), pp. 530–533.
doi: 10.2307/3898874.

McArthur, E. D. and Young, S. A. (1999) 'Development of native seed supplies to support
restoration of pinyon-juniper sites', in *S.B. Monsen and R. Stevens, compilers, Proceedings:*
*ecology and management of pinyon-juniper communities within the Interior West. Proc. RMRS-*
*P-9*. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT, pp. 327–330.

Meyer, S. E. *et al.* (1995) 'Germination ecophysiology of Leymus cinereus (Poaceae)',
*International Journal of Plant Sciences*, 156(2), pp. 206–215. doi: 10.1086/297242.

Meyer, S. E. (1997) 'Genecological Considerations in Grassland Restoration Using Wild-

Collected Seed Sources', in *Proceesings of the 18th International Grassland Congress,*

*Winnipeg and Saskatoon, Canada*, pp. 8–17. Available at:

http://www.internationalgrasslands.org/files/igc/publications/1997/iii-299.pdf (Accessed: 8

January 2018).

Meyer, S. E. and Monsen, S. B. (1991) 'Habitat-correlated variation in mountain big sagebrush

(Artemisia tridentata ssp. vaseyana) seed germination patterns', *Ecology*, 72(2), pp. 739–742.

doi: 10.2307/2937214.

Vander Mijnsbrugge, K., Bischoff, A. and Smith, B. (2010) 'A question of origin: Where and

how to collect seed for ecological restoration', *Basic and Applied Ecology*, 11(4), pp. 300–311.

doi: 10.1016/J.BAAE.2009.09.002.

Oduor, A. M. O. (2013) 'Evolutionary responses of native plant species to invasive plants: a

review', *New Phytologist*, 200(4), pp. 986–992. doi: Doi 10.1111/Nph.12429.

Oduor, A. M. O., Leimu, R. and van Kleunen, M. (2016) 'Invasive plant species are locally

adapted just as frequently and at least as strongly as native plant species', *Journal of Ecology*.

doi: 10.1111/1365-2745.12578.

Oksanen, J. *et al.* (2018) 'Vegan: community ecology package', *R package 2.4-6*. doi:

10.4135/9781412971874.n145.

Pilliod, D. S., Welty, J. L. and Toevs, G. R. (2017) 'Seventy-Five Years of Vegetation

Treatments on Public Rangelands in the Great Basin of North America', *Rangelands*, 39(1), pp.

1–9. doi: 10.1016/j.rala.2016.12.001.

Plant Conservation Alliance, P. (2015) 'National Seed Strategy for Rehabilitation and

Restoration 2015-2020'. Bureau of Land Management. Available at:

https://www.blm.gov/programs/natural-resources/native-plant-communities/national-seed-

strategy.

Prober, S. M. *et al.* (2015) 'Climate-adjusted provenancing: a strategy for climate-resilient

ecological restoration', *Frontiers in Ecology and Evolution*, 3, p. 65. doi:

10.3389/fevo.2015.00065.

R Core Team (2017) 'R: A Language and Environment for Statistical Computing', *R Foundation*
*for Statistical Computing*. Vienna. Available at: https://www.r-project.org/.

Reich, P. B. *et al.* (2003) 'The evolution of plant functional variation: traits, spectra, and
strategies', *International Journal of Plant Sciences*, 164(S3), pp. S143–S164. doi:
10.1086/374368.

Rice, K. J. and Emery, N. C. (2003) 'Managing microevolution: Restoration in the face of global
change', *Frontiers in Ecology and the Environment*, 1(9), pp. 469–478. doi: 10.1890/1540-
9295(2003)001[0469:MMRITF]2.0.CO;2.

Richards, R. T., Chambers, J. C. and Ross, C. (1998) 'Use of native plants on federal lands:
policy and practice', *Journal of Range Management*, 51(6), pp. 625–632. doi: 10.2307/4003603.

Des Roches, S. *et al.* (2018) 'The ecological importance of intraspecific variation', *Nature*
*Ecology & Evolution*, 2(1), pp. 57–64. doi: 10.1038/s41559-017-0402-5.

Rogers, D. L. and Montalvo, A. M. (2004) 'Genetically appropriate choices for plant materials to
maintain biological diversity', in *University of California. Report to the USDA Forest Service,*
*Rocky Mountain Region, Lakewood, CO.* doi: Genetic Considerations in Ecological Restoration.

Rowland, M. M., Suring, L. H. and Michael, J. (2010) 'Assessment of habitat threats to
shrublands in the Great Basin: a case study', *Advances in threat assessment and their application*
*to forest and rangeland management. General Technical Report. PNW-GTR-802*, pp. 673–685.
Available at: http://www.northern-ecologic.com/publications/54.pdf.

Sheridan, J. A. and Bickford, D. (2011) 'Shrinking body size as an ecological response to climate
change', *Nature Climate Change*, 1(8), pp. 401–406. doi: 10.1038/nclimate1259.

Siepielski, A. M., Dibattista, J. D. and Carlson, S. M. (2009) 'It's about time: The temporal
dynamics of phenotypic selection in the wild', *Ecology Letters*, pp. 1261–1276. doi:
10.1111/j.1461-0248.2009.01381.x.

St Clair, J. B., Mandel, N. L. and Vance-Borland, K. W. (2005) 'Genecology of Douglas fir in

Western Oregon and Washington', *Annals of Botany*, 96(7), pp. 1199–1214. doi:

10.1093/aob/mci278.

Sultan, S. E. and Spencer, H. G. (2002) 'Metapopulation structure favors plasticity over local

adaptation', *The American Naturalist*, 160(2), pp. 271–283. doi: 10.1086/341015.

Svejcar, T. *et al.* (2017) 'Challenges and limitations to native species restoration in the Great

Basin, USA', *Plant Ecology*, 218(1), pp. 81–94. doi: 10.1007/s11258-016-0648-z.

Tisdale, E. W. and Hironaka, M. (1981) 'The Sagebrush-Grass Region: A Review of the

Ecological Literature', *University of Idaho Forest, Wildlife, and Range Experiment Station*.

Available at: http://digital.lib.uidaho.edu/cdm/ref/collection/fwres/id/169 (Accessed: 5 January

2018).

Turesson, G. (1922) 'The genotypical response of the plant species to the habitat', *Hereditas*,

3(3), pp. 211–350. doi: 10.1111/j.1601-5223.1922.tb02734.x.

United States. House of Representatives. Committee on Appropriations. (2014) 'American Seed

Trade Association Statement by Mark Mustoe, Co-Owner and Manager of Clearwater Seed

Regarding Efficient Native Seed Use by the Bureau of Land Management'. Alexadria, VA.

Available at: http://docs.house.gov/meetings/AP/AP06/20140410/101762/HHRG-113-AP06-

Wstate-MustoeM-20140410.pdf (Accessed: 4 January 2018).

USDA and NRCS (2018) *The PLANTS Database, National Plant Data Team, Greensboro, NC*

*27401-4901 USA*. Available at: http://plants.usda.gov (Accessed: 8 January 2018).

USDOI (2015) *SO-3336: An Integrated Rangeland Fire Management Strategy. Final Report to*

*the Secretary of the Interior.* Available at:

https://www.forestsandrangelands.gov/rangeland/documents/IntegratedRangelandFireManageme

ntStrategy_FinalReportMay2015.pdf. [Accessed January 26, 2018].

Vavrek, M. J. (2011) 'Fossil: palaeoecological and palaeogeographical analysis tools',

*Palaeontologia Electronica*, 14(1), p. 16.

Wang, T. *et al.* (2016) 'Locally downscaled and spatially customizable climate data for historical

and future periods for North America', *PLoS ONE*, 11(6), p. e0156720. doi:

10.1371/journal.pone.0156720.

Ward, R. T. (1969) 'Ecotypic variation in Deschampsia caespitosa (L.) Beauv. from Colorado',
*Ecology*, 50(3), pp. 519–522. doi: 10.2307/1933914.

Williams, A. V., Nevill, P. G. and Krauss, S. L. (2014) 'Next generation restoration genetics:
Applications and opportunities', *Trends in Plant Science*, 19(8), pp. 529–537. doi:
10.1016/j.tplants.2014.03.011.

Wright, I. J. *et al.* (2017) 'Global climatic drivers of leaf size', *Science*, 357(6354), pp. 917–921.
doi: 10.1126/science.aa14760.

Table 1. Traits measured in outdoor common gardens or reciprocal transplants for at least 5

populations in at least 20 common garden locations, with data available from text, tables, author

contact, or extraction from figures. Note that in some cases, multiple highly similar measures

were grouped, as indicated in footnotes.

| Trait | Units | Locations |
|---|---|---|
| date – flowering[1] | # days | 34 |
| size – floral[2] | cm | 22 |
| height - plant | cm | 46 |
| size – leaf[3] | cm | 30 |
| mass – shoots[4] | g | 43 |
| number - inflorescence[5] | # | 36 |
| number – seeds[6] | # | 21 |
| survival | % | 43 |

[1]Flowering date or any other floral phenology

[2]Any size measurement of a floral structure

[3]Most frequently, leaf length; occasionally leaf width

[4] Any measure of aboveground biomass

[5] Counts of flowers or flowering structures

[6] Most frequently seed number, but also seed yield in mass and/or seed yield rating/rank

**Figure captions**

Figure 1. Summary of reviewed literature that compared traits among at least two populations in at least one common environment, by lifeform. Total counts of published studies, species, taxa, and taxa-specific experiments (A); types of experiments (B); means and standard errors of duration of the experiments that measured more than germination traits (C); total counts of experiments that measured only germination traits, (D); means and standard errors of number of populations tested in each experiment (E), and garden sites per experiment for outdoor reciprocal transplant and common garden experiments (F).

Figure 2. Map of 129 different outdoor common garden locations (A) and 2953 unique population collection sites (B) for the 80% of outdoor gardens and 91% of experiments for which coordinates could be obtained or generated, from 170 studies reviewed. The size of the marker in panel A represents the number of experiments in which each specific garden location was used, with larger symbols indicating garden locations used in more experiments. Although all species represented are native to the floristic Great Basin (white outline), many populations were collected and tested outside this region.

Figure 3. Summary of the years in which the collections of each experiment were made (A, left), the year each experiment was performed (A, right), and the average geographic distance among population collections sites in each experiment. The percent of 327 experiments that reported this information were 99% and 88% (respectively) for panel A, and 80% for panel B. Collection year and experiment year represent the average for each experiment, as it was common for materials to be collected and tested over multiple years for each experiment. Geographic distance is the

mean pairwise distance among populations in each experiment; note the noncontinuous vertical

axis.

Figure 4. Summary of among-population variation (A, signature 1) and trait-by-environment

associations (B, signature 2) for any measured trait, grouped by five life history traits. Summary

of local advantage (C, signature 3) for reproductive traits, survival traits, fitness indices, or

biomass. Data compiled from 327 experiments from 170 published studies on Great Basin plants

(see Supporting Information Appendix 2 and available datasets in electronic supplementary

material). For signatures 1 and 2, "Yes" and "No" represent statistical comparisons, while

"Authors claim "Yes"" and "Authors claim "No"" represent textual, claim-based results where

supporting statistics were not reported (common in older studies). For signature 3, most

experiments had multiple gardens, and many evaluated performance at multiple sampling dates,

leading to 5 different scores. These scores, from "All gardens, all times" to "No gardens at any

time" represent a gradient of incidence and frequency of this signature (see methods). For all

panels, numbers in parentheses, (x), indicate the number of experiments scored in a given

category, and the dashed gray lines indicate 50%.

Figure 5. Summary of 1,465 trait scores from the 305 experiments appropriate for detecting

signature 1 (differences between populations). Scores of "Yes" and "No" were supported by

statistical comparisons, while the "Authors claim…" scores represent textual, claim-based results

where supporting statistics were not reported (common in older studies). Numbers in

parentheses, (x), indicate the total experiments that measured each trait or reported each factor,

and dashed gray line indicates 50%.

Figure 6. Summary of scores for associations between 592 traits (A) and 426 environmental

factors (B) from the 161 experiments appropriate for detecting signature 2 (trait-by-environment

association), expressed by trait/factors, and an example from the literature (C, redrawn with

permission from (Meyer and Monsen, 1991)) in which date of germination for mountain big

sagebrush is correlated with a measure of monthly temperature (treatment: 2-week chill). Scores

of "Yes" and "No" were supported by statistical comparisons, while the "Authors claim…"

scores represent textual, claim-based results where supporting statistics were not reported

(common in older studies). For panels A and B, numbers in parentheses, (x), indicate the total

experiments that measured each trait or reported each factor, and the dashed gray lines indicate

50%.

Figure 7. Results of comparisons of pairwise trait and environmental distances for eight

frequently measured phenotypic traits and (A) the mean annual precipitation (MAP) or (B) mean

annual temperature (MAT) at the original collection location. Values are effect sizes and 95%

confidence intervals for each trait, averaged across all experiments for which data were available

(number of experiments in parentheses). Examples of the two strongest relationships are shown

for leaf size and MAP (C), where each line shows the correlation coefficient and confidence

intervals for an individual experiment, for which we calculated the relationship between

differences in percent survival and difference MAP at location of origin. Color indicates

functional groups: Green = grasses, blue = shrubs, orange = forbs. Examples are shown for the

two highest effect sizes: D), experiment 297A, (Kramer, Larkin and Fant, 2015), *Penstemon*

*deustus* and E), experiment 297A, (Kramer, Larkin and Fant, 2015), *Eriogonum microthecum*.

Similarly, flowering time and MAT (F) is shown, with examples of G) experiment 271A,

(Larsen, 1947), *Schizachyrium scoparium*, and H) experiment 245A, (Ward, 1969), *Deschampsia*

*caespitosa*. Full results for each trait/environment relationship are shown as additional results in

Supporting Information Appendix 3.

Figure 1.

Figure 2.

Figure 3.

    Figure 4.

991     Figure 5.

993    Figure 6.

998 Figure 7.

1000 **Appendix 1. Additional Methods**

1001

*Literature search*

Terms used to search the literature included 'plant', 'Great Basin', 'Intermountain West',

'western United States', 'local adaptation', 'ecotypic variation', 'phenotypic variation', 'genetic

variation', 'habitat-correlated variation', 'genecology', 'intraspecific variation', 'ecotype', 'seed zones',

'common garden', 'reciprocal garden', and 'transplant garden', as well as combinations of these terms.

Literature was obtained primarily using the World Wide Web as well as databases such as Google

Scholar, Web of Science, Academic Search Premier, JSTOR, Science Direct, and Wiley Online Library.

When digital copies were not available, they were obtained from academic libraries. The citations within

the resulting literature were also mined for additional literature that our first search had missed.

1011

*Geographic range categorization*

Four categories of geographic range were assigned from distributions in the USDA Plants

Database (https://plants.sc.egov.usda.gov), as follows. Widespread: found in majority of United States

(e.g. *Elymus elymoides* (Raf.) Swezey); Regional: common in the floristic Great Basin but not found

throughout the United States (e.g. *Atriplex confertifolia* (Torr. & Frém.) S. Watson); Narrow: limited to

specific, well-defined habitats within the Great Basin (e.g. *Penstemon confusus* M.E. Jones); Endemic:

restricted to several counties (e.g. *Allium passeyi* N.H. Holmgren & A.H. Holmgren).

1019

*Geographic coordinate generation*

Geographic coordinates and elevations for gardens and populations were recorded verbatim from

studies that contained precise coordinates, or were generated manually using Google Earth Pro (Google

Inc., 2018) with assistance from the Geographic Names Information System (US Geological Survey,

2018) when vague coordinates or textual localities were given. All coordinates were converted to decimal

degrees (WGS 84) and elevations were recorded in meters. Uncertainties in manually generated

coordinates were recorded in a measure of accuracy, either 'high' (confident to within a ~2 mile radius),

'fair' (confident to within a ~5 mile radius), or 'low' (confident to within a ~15 mile radius). Numeric

coordinates given in the studies were assumed to be accurate to within one mile. If elevations were given

for populations or gardens with vague localities, we utilized this information to increase the confidence of

our generated location. Coordinates were not generated for localities that were exceptionally vague or

studies which did not include localities. If a study utilized a named release or cultivar, the location of

origin was determined by locating the original published release notice, if available. Cultivars bred using

populations from multiple locations were not assigned origin coordinates.

1034

*Scoring experiments for each signature of local adaptation*

For among-population variation (signature 1), a score of 'Yes' was given when at least one

measured trait was reported to differ significantly between at least two populations, and a score of 'No'

was given when differences in any phenotypic trait were not detected between any pair of populations.

For trait-by-environment association (signature 2), a score of 'Yes' was given when authors reported a

significant association between at least one trait and one measure of the environment of origin, and a

score of 'No' was given when the author tested for but found no such relationship. In addition to a score

for each experiment, each of the measured and reported traits and environmental variables were scored

(hereafter, trait scores) in a manner that indicated which traits did or did not vary between populations, as

well as which traits and environmental variables were or were not correlated with each other (see

available datasets in electronic supplementary material). Some experiments met the criteria for both

signatures while others met only one or the other. In several studies, especially older studies or studies

whose analyses did not include among-population comparisons, the significance of variation and/or

correlation needed for scoring signatures 1 and 2 could not be determined because the authors provided

results without statistical analyses. In these cases, results were scored as 'Authors Claim Yes' or 'Authors

Claim No', and the scoring was done as described above, taking authors at their word in the absence of

published statistical evidence.

To score whether there was higher fitness of a local population in a common garden (signature 3),

only experiments in which outdoor reciprocal transplants or common gardens were performed using a

local population (identified as such by the author, or clearly collected from the common garden site) in at

least one garden were considered. Additionally, the experiment had to measure survival, reproductive

output (number of seeds or flowers, or other reproductive output), a fitness index (a combination of

several size and production traits), or total aboveground biomass. Each experiment was given a composite

score to fully capture variation in the performance of the local population across gardens (spatial), as well

as through different sampling dates (temporal). For the spatial component, 'Yes for all gardens' indicates

the highest values in each garden belonged to that garden's local population, 'Yes for some gardens'

indicates the highest value in at least one but not all of the gardens belonged to each garden's local

population, and 'No for all gardens' if the highest value never belonged to a garden's local population.

For the temporal component, the experiment was scored as 'Always' if the local population had the

highest value at all sampling dates, or 'Sometimes' if the local population had the highest value at one but

not all of the sampling dates. For "some" and "sometimes" scores, we calculated the number of

observations of higher fitness of local populations per garden and per time measured to understand what

proportion of gardens and sampling dates showed higher local fitness. This provides an estimate of the

frequency of higher local fitness, but it is not a measure of the importance of the difference per se. For

example, a fitness difference could occur at a low frequency, but have a large impact on population

trajectories (i.e. large differences in survival after a rare drought event).

1071

*Determining whether maternal effects were controlled*

Experiments which tested populations that had all shared one or more generations in the same

location prior to testing were considered to have attempted to control for maternal effects. We determined

the number of generation in common by carefully reading the methods for mentions of the populations'

lineages prior to testing. Some experiments supplied the original location of material collection but

indicated that all materials were collected from areas such as 'evaluation plots', 'seed fields', 'uniform

gardens', or 'increase fields', indicating that at least one generation was shared among all populations,

and therefore and attempt had been made to control maternal effects (intentional or not). Some complex

studied had to be split into multiple experiments because they used different generations of the same

populations in different tests. For example, a study which collected wild adults from their native habitats

and grew them in a common garden for the duration of the experiment before measuring traits of the

plants as well as traits of the seeds they produced were split into two experiments, one containing the

traits of the adult plants (which did not attempt to control for maternal effects, because the progenitors of

the measured material did not share a common location), and one for the seed traits (which did attempt to

control for maternal effects, because the progenitors of the measured material did share a common

location).

1088

*Extraction for quantitative comparison of trait-by-environment association*

To examine links between the variation in trait values and the variation in environmental and

geographic distance among the population's origins, we utilized experiments from which population-

specific trait data as well as geographic coordinates for at least one garden and at least two populations

could be extracted or obtained through author contact. Data from laboratory and greenhouse experiments

were not considered for this extraction, because the great majority of these experiments were not designed

to completely simulate natural growing conditions. Excluding these experiments reduced our pool from

325 to 161. Next, a list of priority fitness traits were developed (Table S1-1) based on traits that were

most commonly measured and potentially associated with plant fitness in the Great Basin (Bower, Clair,

and Erickson, 2014; Leger and Baughman, 2015). Any experiment that did not measure at least one

priority trait was omitted from next steps, and this further reduced our pool from 161 to 153.

Table S1-1. Priority traits targeted in the extraction for the dataset used in the quantitative comparison,

and the preferred units. Note that for several traits, several highly similar measures were included, as

indicated in footnotes.

| Trait | units | Trait | units |
|---|---|---|---|
| survival | % | number - inflorescence[4] | # |
| emergence | % | number – seeds[5] | # |
| germination | % | number – leaves | # |
| height - plant | cm | date – germination | # days |
| length – root | cm | date – regrowth/greenup | # days |
| length – leaf[1] | cm | date – emergence | # days |
| dimensions – floral[2] | cm | date – flowering[6] | # days |
| mass – roots | g | date – seed shatter | # days |
| mass – shoots[3] | g | date – senescence | # days |
| mass – seed | g/seed | | |

[1]If unmeasured, then leaf width was recorded, if available

[2]Any measure of a floral structure

[3]Any measure of aboveground biomass

[4]Any kind of count of flowers or flowering structures was recorded

[5]If no direct count was available, any measure of seed yield was recorded,
including total seed yield in weight and/or seed yield rating/rank

[6]If unmeasured, any other floral phenology was considered

The remaining studies were then examined for textual, tabular, or visual data that could be

extracted as mean values of priority traits for each population in each garden. Extracted values for were

recorded verbatim from tables and throughout the text where possible, and from figures using

WebPlotDigitizer (Rohatgi, 2017) when needed. Means for at least two populations in at least one garden

were required for extraction. If exact matches to certain priority traits were not reported in the studies,

similar measures that were likely to be strongly correlated to the given trait could be recorded as

surrogates if available, and a note was made (footnotes, Table S1-1). We extracted the latest date for

which the most populations at the most gardens were represented if studies presented data for multiple

dates throughout the experiment. In some cases, experiments were conducted with multiple treatments in

which growing conditions were altered to address study questions. In these cases, we only extracted data

for the author-defined 'control' treatment. However, if no control was defined, we used the treatment that

was the most unaltered or representative of the garden environment (e.g. unweeded, or unwatered).

**Appendix 2. Summary of literature and available datasets**

1115

The data collected and generated by this study (Baughman *et al.*, 2019), as well as the list of

publications that were involved in each part of this study, are provided so that additional questions may be

addressed and for other applications. We encourage such additional analyses.

*Summary of literature*

Appendix 2 Table 1. Summary, by species, of the literature included in this study, including

lifeform (F = forb, G = grass, S = shrub, T = tree), counts of studies, experiments, unique populations, and

experiments by type (LAB = laboratory, GH = greenhouse, CG = outdoor common garden, RT = outdoor

reciprocal transplant), the incidence of each signature of local adaptation (1 = differences among

populations, 2 = trait/environment correlations, 3 = higher performance of local than nonlocal population

in local's environment), counts of experiments used in the quantitative comparison of trait-by-

environment associations (QC), and a list of traits used in the QC. See footnotes for additional

information.

| Species | Lifeform | # Studies | # Expts. | # Pops[1] | # LAB expts. | # GH expts. | # CG expts. | # RT expts. | Signature 1[2] | Signature 2[2] | Signature 3[3] | # Expts. in QC | Traits in QC[4] |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| *Achnatherum thurberianum* | G | 2 | 2 | 68 | - | 1 | 1 | - | 2Y/0N | 2Y/0N | - | 1 | 1-7 |
| *Allium acuminatum* | F | 2 | 2 | 56 | 1 | - | 1 | - | 2Y/0N | 2Y/0N | - | 1 | 1,4-8 |
| *Allium brandegeei* | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| *Allium passeyi* | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| *Amelanchier utahensis* | S | 2 | 4 | 18* | - | - | 4 | - | 4Y/0N | 3Y/0N | - | - | - |
| *Andropogon scoparius* | G | 5 | 6 | 55* | - | 2 | 3 | 1 | 6Y/0N | 4Y/0N | - | - | - |
| *Artemisia tridentata* | S | 24 | 25 | 172* | 7 | - | 14 | 4 | 22Y/3N | 8Y/0N | 2Y/5N | - | - |
| *Astragalus filipes* | F | 1 | 1 | 67 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1-4,8 |
| *Atriplex canescens* | S | 9 | 9 | 79 | 4 | - | 5 | - | 9Y/0N | 3Y/0N | 1Y/2N | 2 | 1,2,8 |
| *Atriplex confertifolia* | S | 2 | 3 | 126 | 2 | - | 1 | - | 3Y/0N | 3Y/0N | - | - | - |
| *Bouteloua gracilis* | G | 5 | 6 | 66* | - | 1 | 4 | 1 | 6Y/0N | 4Y/0N | - | 3 | 5 |
| *Bromus carinatus* | G | 1 | 1 | 193 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1-6 |
| *Carex aquatilis* | G | 1 | 1 | 5 | - | - | - | 1 | 1Y/0N | 1Y/0N | 2Y/0N | - | - |
| *Cercocarpus montanus* | T | 1 | 1 | 9 | - | 1 | 1 | - | 1Y/0N | 0Y/1N | - | - | - |
| *Chaenactis douglasii* | F | 1 | 1 | 15 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1,2,4,8 |
| *Chrysothamnus viscidiflorus* | S | 4 | 4 | 8 | - | - | 4 | - | 2Y/2N | - | - | - | - |
| *Cistanthe umbellata* | F | 1 | 2 | 5* | 2 | - | - | - | - | 2Y/0N | - | - | - |
| *Claytonia perfoliata* | F | 1 | 2 | 2* | - | - | - | 2 | 2Y/0N | - | 3Y/0N | - | - |
| *Cleome lutea lutea* | F | 1 | 1 | 9 | - | 1 | - | - | 1Y/0N | 1Y/0N | - | - | - |
| *Coleogyne ramosissima* | S | 2 | 2 | 53 | 1 | - | 1 | - | 2Y/0N | 2Y/0N | - | 1 | 2,3 |
| *Cryptantha circumscissa* | F | 1 | 2 | 4* | 2 | - | - | - | - | 1Y/1N | - | - | - |
| *Dalea ornata* | F | 1 | 1 | 22 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1,3-5 |
| *Dalea searlsiae* | F | 1 | 1 | 20 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1,3,4 |
| *Delphinium nelsonii* | F | 1 | 1 | 3 | - | - | - | 1 | 1Y/0N | 1Y/0N | 2Y/0N | - | - |
| *Deschampsia cespitosa* | G | 2 | 2 | 22* | - | - | 2 | - | 2Y/0N | 2Y/0N | 0Y/1N | - | 1,2,5,6 |
| *Elymus canadensis* | G | 3 | 4 | 30* | - | 1 | 2 | 1 | 4Y/0N | 2Y/0N | - | 3 | 5 |
| *Elymus elymoides* | G | 19 | 24 | 170* | 5 | 11 | 8 | - | 21Y/1N | 6Y/4N | 2Y/1N | 3 | 1,3,5 |
| *Elymus glaucus* | G | 3 | 4 | 158* | 1 | 1 | 1 | 1 | 4Y/0N | 1Y/0N | 2Y/0N | 2 | 1,5-8 |
| *Elymus multisetus* | G | 7 | 7 | 36* | 2 | 4 | 1 | - | 6Y/0N | 3Y/1N | 1Y/0N | 1 | 1,5 |
| *Elymus trachycaulus* | G | 2 | 3 | 5 | 1 | 1 | 1 | - | 2Y/1N | - | - | - | - |
| *Encelia farinosa* | S | 1 | 1 | 2 | - | 1 | - | - | 1Y/0N | 1Y/0N | - | - | - |
| *Ephedra nevadensis* | S | 1 | 1 | 2 | - | - | 1 | - | 1Y/0N | - | 1Y/0N | - | - |
| *Epilobium densiflorum* | F | 1 | 1 | 22 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1,3,5-7 |
| *Ericameria nauseosa* | S | 16 | 17 | 64* | 9 | - | 8 | - | 15Y/2N | 3Y/0N | 0Y/1N | - | - |
| *Eriogonum microthecum* | S | 1 | 1 | 6 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1-4,6,8 |
| *Eriogonum ovalifolium* | S | 1 | 1 | 6 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | 1 | 1-4,6,8 |
| *Eriogonum umbellatum* | S | 1 | 1 | 5 | - | - | 1 | - | 1Y/0N | 0Y/1N | - | - | - |
| *Erysimum capitatum* | F | 2 | 4 | 9* | 2 | - | 2 | - | 2Y/0N | 1Y/1N | 1Y/0N | - | - |
| *Festuca idahoensis* | G | 5 | 6 | 43* | 2 | - | 4 | - | 6Y/0N | 2Y/1N | - | 3 | 1,3 |
| *Grayia spinosa* | S | 2 | 2 | 7 | 1 | - | 1 | - | 2Y/0N | - | - | - | - |
| *Gutierrezia sarothrae* | S | 3 | 4 | 6* | - | 3 | 1 | - | 4Y/0N | 0Y/1N | - | - | - |
| *Helianthus anomalus* | F | 1 | 1 | 8 | - | 1 | - | - | 1Y/0N | 1Y/0N | - | - | - |
| *Hesperostipa comata* | G | 5 | 5 | 10* | - | 2 | 3 | - | 3Y/2N | - | 1Y/0N | - | - |

1128

| | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Holodiscus discolor | S | 1 | 1 | 39 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | - | - |
| Koeleria macrantha | G | 4 | 5 | 29* | 1 | 2 | 2 | - | 4Y/1N | - | - | 2 | 1,5,6 |
| Krascheninnikovia lanata | S | 9 | 13 | 26* | 7 | 5 | - | 1 | 13Y/0N | 2Y/0N | 3Y/0N | 1 | 2,7 |
| Leptosiphon nuttallii | F | 1 | 1 | 3 | 1 | - | - | - | - | 1Y/0N | - | - | - |
| Leymus cinereus | G | 6 | 9 | 141* | 3 | 1 | 4 | 1 | 8Y/1N | 3Y/0N | 1Y/0N | 3 | 1-7 |
| Linum lewisii | F | 3 | 6 | 22* | 1 | 1 | 4 | - | 6Y/0N | 2Y/0N | - | 2 | 2,4,8 |
| Lomatium dissectum | F | 1 | 1 | 3 | - | - | 1 | - | 1Y/0N | - | - | - | - |
| Lotus utahensis | F | 1 | 1 | 14 | - | - | 1 | - | 1Y/0N | 0Y/1N | - | - | - |
| Lupinus latifolius | F | 1 | 1 | 53 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | - | - |
| Machaeranthera canescens | F | 1 | 1 | 9 | - | - | 1 | - | 1Y/0N | - | - | 1 | 1,2,8 |
| Mimulus cardinalis | F | 1 | 1 | 6 | - | - | 1 | - | 1Y/0N | - | - | 1 | - |
| Mimulus lewisii | F | 1 | 1 | 6 | - | - | 1 | - | 1Y/0N | - | - | 1 | - |
| Nicotiana attenuata | F | 1 | 1 | 43 | - | 1 | - | - | 1Y/0N | 0Y/1N | - | - | - |
| Oxyria digyna | F | 1 | 2 | 11 | 2 | - | - | - | - | 0Y/2N | - | - | - |
| Pascopyrum smithii | G | 5 | 11 | 42* | - | 2 | 9 | - | 10Y/1N | 1Y/0N | 0Y/1N | 5 | 1,3,4,6,8 |
| Penstemon acuminatus | F | 1 | 1 | 2 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon comarrhenus | F | 2 | 2 | 4* | 2 | - | - | - | 2Y/0N | Y/1N | - | - | - |
| Penstemon confusus | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon cyananthus | F | 1 | 1 | 10 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| Penstemon cyanocaulis | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| Penstemon deustus | F | 3 | 4 | 12* | 2 | - | 2 | - | 4Y/0N | 2Y/0N | - | 1 | 2-4,6-8 |
| Penstemon eatonii | F | 2 | 3 | 13* | 2 | - | 1 | - | 3Y/0N | 2Y/0N | - | - | - |
| Penstemon fruticosus | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon humilis | F | 1 | 1 | 6 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon leiophyllus | F | 1 | 1 | 4 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| Penstemon leonardii | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon linarioides | F | 1 | 1 | 2 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon newberryi | F | 1 | 2 | 2* | 2 | - | - | - | - | 1Y/1N | - | - | - |
| Penstemon pachyphyllus | F | 3 | 4 | 14* | 2 | - | 2 | - | 4Y/0N | 4Y/0N | - | 1 | 2-4,6-8 |
| Penstemon palmeri | F | 1 | 2 | 13 | 2 | - | - | - | 2Y/0N | 0Y/1N | - | - | - |
| Penstemon petiolatus | S | 1 | 1 | 2 | 1 | - | - | - | 1Y/0N | 0Y/1N | - | - | - |
| Penstemon rostriflorus | F | 3 | 5 | 16* | 3 | - | 2 | - | 5Y/0N | 4Y/1N | - | 1 | 2-4,6-8 |
| Penstemon rydbergii | F | 1 | 1 | 2 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon subglaber | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| Penstemon utahensis | F | 1 | 1 | 5 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Penstemon watsonii | F | 1 | 1 | 4 | 1 | - | - | - | 1Y/0N | 1Y/0N | - | - | - |
| Plantago ovata | F | 1 | 2 | 12* | 1 | - | 1 | - | 2Y/0N | 2Y/0N | - | - | - |
| Poa secunda | G | 7 | 8 | 158* | 2 | 2 | 3 | 1 | 8Y/0N | 3Y/0N | 2Y/2N | 2 | 1-7 |
| Polemonium viscosum | F | 1 | 2 | 2 | - | 1 | - | 1 | 2Y/0N | 1Y/0N | 2Y/0N | - | - |
| Polygonum viviparum | F | 1 | 1 | 3 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| Populus angustifolia | T | 1 | 1 | 5 | - | 1 | - | - | 1Y/0N | - | - | - | - |
| Potentilla pulcherrima | F | 1 | 1 | 15 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | - | - |
| Pseudoroegneria spicata | G | 6 | 7 | 138 | 1 | 1 | 5 | - | 6Y/1N | 1Y/0N | 1Y/0N | 3 | 1,3-7 |
| Purshia tridentata | S | 2 | 2 | 24 | 1 | - | 1 | - | 1Y/0N | 0Y/2N | - | - | - |
| Ranunculus flammula | F | 1 | 1 | 25 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | - | - |
| Saxifraga oregana | F | 1 | 1 | 10 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | - | - |
| Solidago gigantea | F | 1 | 1 | 7 | - | 1 | - | - | 1Y/0N | 0Y/1N | - | - | - |
| Solidago velutina | F | 1 | 1 | 6 | - | - | 1 | - | 1Y/0N | 1Y/0N | - | - | - |

| | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| *Sphaeralcea coccinea* | F | 1 | 2 | 2* | - | - | 2 | - | 2Y/0N | - | - | - | - |
| *Sphaeralcea grossulariifolia* | F | 1 | 1 | 3 | - | - | 1 | - | 1Y/0N | - | - | - | - |
| *Sphaeralcea munroana* | F | 1 | 2 | 49* | - | - | 2 | - | 2Y/0N | - | - | - | - |
| *Sphaeralcea parvifolia* | F | 1 | 1 | 7 | - | - | 1 | - | 1Y/0N | - | - | - | - |
| *Sporobolus airoides* | G | 4 | 4 | 10 | 1 | 1 | 2 | - | 3Y/1N | 3Y/0N | - | - | - |
| *Sporobolus cryptandrus* | G | 3 | 4 | 30* | 1 | - | 3 | - | 4Y/0N | 4Y/0N | - | 1 | 1,3,5 |
| *Stellaria longipes* | F | 1 | 2 | 7* | 1 | - | - | 1 | 2Y/0N | - | - | - | - |
| *Stephanomeria minor* | F | 1 | 2 | 3* | 2 | - | - | - | - | 1Y/1N | - | - | - |
| *Stipa hymenoides* | G | 15 | 17 | 397* | 3 | - | 13 | 1 | 16Y/1N | 3Y/4N | 0Y/1N | 2 | 1,3-8 |
| *Symphoricarpos oreophilus* | S | 1 | 1 | 25 | - | - | 1 | - | 1Y/0N | 0Y/1N | - | - | - |
| *Typha latifolia* | F | 1 | 1 | 2 | 1 | - | - | - | 1Y/0N | - | - | - | - |
| *Xanthium strumarium* | F | 1 | 2 | 49* | 1 | - | 1 | - | 2Y/0N | 2Y/0N | - | 1 | 5 |

[1]Numbers followed by a * indicates a reduced count that accounts for the same populations used in multiple experiments

[2]Count of experiments; Y = "Yes" and "Authors claim Yes" scores, N = "No" and "Authors claim No" scores.

[3]Count of experiments; Y = Local did best in at least one garden during at least one sampling date, N = Local never did best

[4]Indicates which traits were used for this species in the quantitative comparison for trait-by-environment associations; 1 = height, 2 = survival, 3 = shoot mass, 4 = number of inflorescences, 5 = flowering day, 6 = leaf size, 7 = floral size,  8 = seed number.

1130

*Available datasets*

Data have been uploaded to Dryad at DOI: TBD (Baughman *et al.*, In Review). Several datasets

are available. The "Summary and signature scores" dataset includes all of the studies and experiments and

summarizes literature categorization as well as scores and associated information for each of the

signatures of local adaptation. The "Trait scores" dataset includes basic study categorization as well as

information that indicated which phenotypic traits (for signatures 1 and 2) and environmental variables

(for signature 2) were involved in each of the signatures of local adaptation. The "Quantitative

comparison" dataset includes all of the data used to conduct the quantitative comparison of trait-by-

environment associations, and lists population-specific mean values for our priority traits for all studies

for which such data were available, the latitude and longitude of each population origin, and extensive

climate information for each origin generated with the ClimateNA v5.10 software package based on

methodology described by Wang et al. (2016). The "Location data" dataset lists all outdoor gardens and

population origin coordinates and elevations for which authors gave this information, as well as those for

which we could confidently generate it. For descriptions of each column in each of these datasets, refer to

the "Data Dictionary" file.

1146

*Bibliography of reviewed literature*

A list of all the literature used in any of the datasets is provided below. Following each citation is

a set of codes in brackets indicating which parts of our study the publication was used in. Codes S1, S2,

and S3 indicate that at least one of the "experiments" in the given publication was used to generate a

score for signatures 1, 2, and 3, and code QC indicates the publication (or the data summarized in it, even

if not available from the publication itself) was used in analyses for the quantitative comparison of trait-

by-environment associations. Note that some published studies were scored as multiple experiments for

multiple species.

Angert, A. L., and D. W. Schemske. 2005. "The Evolution of Species' Distributions: Reciprocal
Transplants across the Elevation Ranges of Mimulus Cardinalis and M. Lewisii." Evolution 59
(8):1671–84. https://doi.org/10.1554/05-107.1. [S1]

Atwater, Daniel Z., and Ragan M. Callaway. 2015. "Testing the Mechanisms of Diversity-
Dependent Overyielding in a Grass Species." Ecology 96 (12):3332–42.
https://doi.org/10.1890/15-0889.1. [S1, QC]

Barker, Jr, and Cm McKell. 1986. "Differences in Big Sagebrush (Artemisia Tridentata) Plant
Stature along Soil-Water Gradients: Genetic Components." Journal of Range Management 39
(2):147–51. http://www.jstor.org/stable/10.2307/3899288. [S1, QC]

Barnes, Melanie G, and Diane L Marshall. 2009. "The Effect of Plant Source Location on
Restoration Success: A Reciprocal Transplant Experiment with Winterfat (Krascheninnikovia
Lanata)." Ecology. https://doi.org/3390801. [S1, S2, S3, QC]

Baughman, Owen W., Susan E. Meyer, Zachary T. Aanderud, and Elizabeth A. Leger. 2016.
"Cheatgrass Die-Offs as an Opportunity for Restoration in the Great Basin, USA: Will Local
or Commercial Native Plants Succeed Where Exotic Invaders Fail?" Journal of Arid
Environments 124:193–204. https://doi.org/10.1016/j.jaridenv.2015.08.011. [S1, S3, QC]

Beckstead, Julie, Susan E. Meyer, and Phil S. Allen. n.d. "Effects of Afterripening on Cheatgrass
(Bromus Tectorum) and Squirreltail (Elymus Elymoides) Germination." In: Roundy, Bruce A.;
McArthur, E. Durant; Haley, Jennifer S.; Mann, David K., Comps. 1995. Proceedings:
Wildland Shrub and Arid Land Restoration Symposium; 1993 October 19-21; Las Vegas, NV.
Gen. Tech. Rep. INT-GTR-315. Ogden, UT: U.S.D.A. [S1, S2]

Bergum, Karin E., Ann L. Hild, and Brian A. Mealor. 2010. "Phenotypes of Two Generations of
Sporobolus Airoides Seedlings Derived from Acroptilon Repens-Invaded and Non-Invaded
Grass Populations." Restoration Ecology 20 (2):227–33. https://doi.org/10.1111/j.1526-
100X.2010.00754.x. [S1, S2]

Bhattarai, Kishor, B. Shaun Bushman, Douglas A. Johnson, and John G. Carman. 2010. "Phenotypic
and Genetic Characterization of Western Prairie Clover Collections from the Western United
States." Rangeland Ecology and Management 63 (6):696–706. https://doi.org/10.2111/REM-
D-10-00008.1. [S1, S2, QC]

Bhattarai, Kishor, Douglas A. Johnson, Thomas A. Jones, Kevin J. Connors, and Dale R. Gardner.
2008. "Physiological and Morphological Characterization of Basalt Milkvetch (Astragalus
Filipes): Basis for Plant Improvement." Rangeland Ecology and Management 61 (4):444–55.
https://doi.org/10.2111/08-011.1. [S1, S2, QC]

Bhattarai, Kishor, B. Shaun Bushman, Douglas A. Johnson, and John G. Carman. 2011. "Searls
Prairie Clover (Dalea Searlsiae) for Rangeland Revegetation: Phenotypic and Genetic
Evaluations." Crop Science 51 (2):716–27. https://doi.org/10.2135/cropsci2010.07.0387. [S1,
S2, QC]

Bleak, A.T., and Neil C. Frischknecht. 1965. "Problems in Artificial and Natural Revegetation of the
Arid Shadscale Vegetation Zone of Utah and Nevada." Journal of Range Management, 59–65.
[S1]

Bohmont, B.L., and Robert Lang. 1957. "Some Variations in Morphological Characteristics and
Palatability among Geographic Strains of Indian Ricegrass." Journal of Range Management,
127–31.[S1]

Booth, D. Terrance. 1992. "Seedbed Ecology of Winterfat: Imbibition Temperature Affects Post-
Germination Growth." Journal of Range Management 45 (2):159–64. [S1]

Booth, D. Terrance, Charles G. Howard, and Charles E. Mowry. 1980. "'Nezpar' Indian Ricegrass:
Description, Justification for Release, and Recommendations for Use." Rangelands Archives 2
(2):53–54. [S1]

Brabec, Martha M., Matthew J. Germino, Douglas J. Shinneman, David S. Pilliod, Susan K.
McIlroy, and Robert S. Arkle. 2015. "Challenges of Establishing Big Sagebrush (Artemisia
Tridentata) in Rangeland Restoration: Effects of Herbicide, Mowing, Whole-Community
Seeding, and Sagebrush Seed Sources." Rangeland Ecology and Management 68 (5):432–35.
https://doi.org/10.1016/j.rama.2015.07.001. [S1, S3, QC]

Brouillette, Larry C., Chase M. Mason, Rebecca Y. Shirk, and Lisa A. Donovan. 2014. "Adaptive
Differentiation of Traits Related to Resource Use in a Desert Annual along a Resource
Gradient." New Phytologist 201 (4):1316–27. https://doi.org/10.1111/nph.12628. [S1, S2]

Butterfield, Bradley J., and Troy E. Wood. 2015. "Local Climate and Cultivation, but Not Ploidy,
Predict Functional Trait Variation in Bouteloua Gracilis (Poaceae)." Plant Ecology 216
(10):1341–49. https://doi.org/10.1007/s11258-015-0510-8. [S1, S2]

Chabot, Brian F., and W. D. Billings. 1972. "Origins and Ecology of the Sierran Alpine Flora and
Vegetation." Ecological Monographs 42 (2):163–99. https://doi.org/10.2307/1942262. [S2]

Chaney, Lindsay, Bryce A. Richardson, and Matthew J. Germino. 2017. "Climate Drives Adaptive
    Genetic Responses Associated with Survival in Big Sagebrush (Artemisia Tridentata)."
    Evolutionary Applications 10 (4):313–22. https://doi.org/10.1111/eva.12440. [S1, S2, QC]

Chapin, F. S. III, and C. M. Chapin. 1981. "Ecotypic Differentiation of Growth Processes in Carex
    Aquatilis along a Latitudinal and Local Gradients." Ecology 62 (4):1000–1009.
    https://doi.org/10.2307/1936999. [S1, S2, S3, QC]

Clark, Lesley D., and Neil E. West. 1971. "Further Studies of Eurotia Lanata Germination in
    Relation to Salinity." The Southwestern Naturalist, 371–75. [S1]

Clary, Warren P. 1975. "Ecotypic Adaptation in Sitanion Hystrix." Ecology 56 (6):1407–15. [S1,
    S2, QC]

Clary, Warren P. 1979. "Variation in Leaf Anatomy and $CO_2$ Assimilation in Sitanion Hystrix
    Ecotypes." The Great Basin Naturalist, 427–32. [S1, S2]

Clauss, M. J., and D. L. Venable. 2000. "Seed Germination in Desert Annuals: An Empirical Test of
    Adaptive Bet Hedging." The American Naturalist 155 (2):168–86.
    https://doi.org/10.1086/303314. [S1, S2, QC]

Cook, Stanton A, and Michael P Johnson. 1968. "Adaptation to Heterogeneous Environments. I.
    Variation in Heterophylly in Ranunculus Flammula L." Evolution 22 (3):496–516.
    https://doi.org/10.1111/j.1558-5646.1968.tb03988.x. [S1, S2]

Dewey, Douglas R. 1960. "Salt Tolerance of Twenty-Five Strains of Agropyron." Agronomy
    Journal 52 (11):631–35. https://doi.org/10.2134/agronj1960.00021962005200110006x. [S1,
    QC]

Doede, David L. 2005. "Genetic Variation in Broadleaf Lupine (Lupinus Latifolius) on the Mt Hood
    National Forest and Implications for Seed Collection and Deployment." Native Plants Journal
    6 (1):36–48. https://doi.org/10.1353/npj.2005.0018. [S1, S2]

Doescher, P.S. 1983. "Phyto-Edaphic Relationships and Ecotypic Development of Festuca
    Idahoensis in Eastern Oregon Habitat Types of Artemisia Tridentata." Ph.D. Dissertation,
    Oregon State University, Corvallis. [S1, S2, QC]

Emery, R. J. N., C. C. Chinnappa, and J. G. Chmielewski. 1994. "Specialization, Plant Strategies,
    and Phenotypic Plasticity in Populations of Stellaria Longipes Along an Elevational Gradient."
    International Journal of Plant Sciences 155 (2):203–19. https://doi.org/10.2307/2995565. [S1,
    QC]

Erickson, Vicky J, Nancy L Mandel, and Frank C Sorenson. 2004. "Landscape Patterns of
    Phenotypic Variation and Population Structuring in a Selfing Grass, Elymus Glaucus (Blue
    Wildrye)." Canadian Journal of Botany 82:1776–89. https://doi.org/10.1139/B04-141. [S1, S2,
    QC]

Evans, Raymond A, and James A Young. 1990. "Survival and Growth of Big Sagebrush (Artemisia
    Tridentata) Plants in Reciprocal Gardens." Weed Science 38 (3):215–19.
    https://doi.org/10.2307/4045014. [S1, S3, QC]

Ferguson, Scot D., Elizabeth A. Leger, Jun Li, and Robert S. Nowak. 2015. "Natural Selection

Favors Root Investment in Native Grasses during Restoration of Invaded Fields." Journal of Arid Environments 116:11–17. https://doi.org/10.1016/j.jaridenv.2015.01.009. [S1]

Ferrero-Serrano, Ángel, Ann L. Hild, and Brian A. Mealor. 2011. "Can Invasive Species Enhance Competitive Ability and Restoration Potential in Native Grass Populations?" Restoration Ecology 19 (4):545–51. https://doi.org/10.1111/j.1526-100X.2009.00611.x. [S1, S2]

Fisk, Matthew R. 2016. "Dynamics of Cold Hardiness Accumulation and Loss in the Great Basin Native Species Eriogonum Umbellatum." Ph.D. Dissertation, University of Idaho, Boise. [S1, S2]

Fitzsimmons, Kevin, Cynthia Lovely, and Edward Glenn. 1998. "Growth Differences among Widely Separated Geographic Accessions of Fourwing Saltbush (Atriplex Canescens) in the Great Basin Desert, New Mexico, USA." Arid Soil Research and Rehabilitation 12 (2):87–94. https://doi.org/10.1080/15324989809381501. [S1, S2, QC]

Fonseca, Carolina, Erin Espeland, and James W. Baxter. 2014. "Patterns of Population Differentiation in Early Traits of Development in Elymus Glaucus: Implications for Restoration." Ecological Restoration 32 (4):388–95. https://doi.org/10.3368/er.32.4.388. [S1]

Galen, Candace, Joel S Shore, and Hudson Deyoe. 1991. "Ecotypic Divergence in Alpine Polemonium Viscosum: Genetic Structure, Quantitative Variation, and Local Adaptation." Evolution 45 (455):1218–28. https://doi.org/10.2307/2409729. [S1, S2, S3, QC]

Goergen, Erin M., Elizabeth A. Leger, and Erin K. Espeland. 2011. "Native Perennial Grasses Show Evolutionary Response to Bromus Tectorum (Cheatgrass) Invasion." PLoS ONE 6 (3). https://doi.org/10.1371/journal.pone.0018145. [S1, S2]

Hall, J.W., D.G. Stout, and B. Brooke. 1990. "Effect of Seed Source on Growth of Giant Wildrye (Elymus Cinereus) at Two Elevations in Interior British Columbia." Canadian Journal of Plant Science 70 (2):551–54. [S1, S3, QC]

Hardegree, Stuart P., Thomas A. Jones, Frederick B. Pierson, Patrick E. Clark, and Gerald N. Flerchinger. 2008. "Dynamic Variability in Thermal-Germination Response of Squirreltail (Elymus Elymoides and Elymus Multisetus)." Environmental and Experimental Botany 62 (2):120–28. https://doi.org/10.1016/j.envexpbot.2007.07.010. [S2]

Harmon, Dan, and Charlie D. Clements. 2016. "Characteristics That Determine a Successful Squirreltail (Elymus Elymoides)." In Poster Session Presented at the Society for Range Management, Corpus Christi, TX. [S1]

Hergert, Holden J., Brian A. Mealor, and Andrew R. Kniss. 2015. "Inter-and Intraspecific Variation in Native Restoration Plants for Herbicide Tolerance." Ecological Restoration 33 (1):74–81. https://doi.org/10.3368/er.33.1.74. [S1]

Hild, A L, J M Muscha, and N L Shaw. 2007. "Emergence and Growth of Four Winterfat Accessions in the Presence of the Exotic Annual Cheatgrass." Proceedings: Shrubland Dynamics-Fire and Water; 2004 August 10-12; Lubbock, TX., no. 47:0–147. [S1]

Hintz, Lisa, M.M. Eshelman, A. Foxx, T.E. Wood, and A. Kramer. 2016. "Population Differentiation in Early Life History Traits of Cleome Lutea Var. Lutea in the Intermountain West." Western North American Naturalist 76 (1):6–17. [S1, S2]

Horning, Matthew E., Theresa R. McGovern, Dale C. Darris, Nancy L. Mandel, and Randy Johnson. 2010. "Genecology of Holodiscus Discolor (Rosaceae) in the Pacific Northwest, U.S.A." Restoration Ecology 18 (2):235–43. https://doi.org/10.1111/j.1526-100X.2008.00441.x. [S1, S2]

Humphrey, L. David, and Eugene W. Schupp. 2002. "Seedling Survival from Locally and Commercially Obtained Seeds on Two Semiarid Sites." Restoration Ecology 10 (1):88–95. https://doi.org/10.1046/j.1526-100X.2002.10109.x. [S1, S3, QC]

Jaindl, Raymond G., Paul Doescher, Richard F. Miller, and Lee E. Eddleman. 1994. "Persistence of Idaho Fescue on Degraded Rangelands: Adaptation to Defoliation or Tolerance." Journal of Range Management 47 (1):54. https://doi.org/10.2307/4002841. [S1, S2, QC]

Johnson, R C, V J Erickson, N L Mandel, J Bradley St Clair, and K W Vance-Borland. 2010. "Mapping Genetic Variation and Seed Zones for Bromus Carinatus in the Blue Mountains of Eastern Oregon, USA." Botany 88 (8):725–36. https://doi.org/10.1139/B10-047. [S1, S2, QC]

Johnson, R C, B C Hellier, and K W Vance-Borland. 2013. "Genecology and Seed Zones for Tapertip Onion in the US Great Basin." Botany-Botanique 91 (10):686–94. https://doi.org/DOI 10.1139/cjb-2013-0046. [S1, S2, QC]

Johnson, R. C., M. J. Cashman, and K. Vance-Borland. 2012. "Genecology and Seed Zones for Indian Ricegrass Collected in the Southwestern United States." Rangeland Ecology and Management 65 (5):523–32. https://doi.org/10.2111/REM-D-11-00165.1. [S1, S2, QC]

Johnson, R. C., E. A. Leger, and Ken Vance-Borland. 2017. "Genecology of Thurber's Needlegrass (Achnatherum Thurberianum [Piper] Barkworth) in the Western United States." Rangeland Ecology and Management 70 (4):509–17. https://doi.org/10.1016/j.rama.2017.01.004. [S1, S2, QC]

Johnson, R. C., and Ken Vance-Borland. 2016. "Linking Genetic Variation in Adaptive Plant Traits to Climate in Tetraploid and Octoploid Basin Wildrye [Leymus Cinereus (Scribn. & Merr.) A. Love] in the Western U.S." PLoS ONE 11 (2). https://doi.org/10.1371/journal.pone.0148982. [S1, S2, QC]

Johnson, Richard C., Matthew E. Horning, Erin K. Espeland, and Ken Vance-Borland. 2015. "Relating Adaptive Genetic Traits to Climate for Sandberg Bluegrass from the Intermountain Western United States." Evolutionary Applications 8 (2):172–84. https://doi.org/10.1111/eva.12240. [S1, S2, QC]

Jones, T A, D C Nielson, J T Arredondo, and M G Redinbaugh. 2003. "Characterization of Diversity among 3 Squirreltail Taxa." Journal of Range Management 56 (5):474–82. https://doi.org/Doi 10.2307/4003839. [S1, QC]

Jones, T.A. 2004. "Registration of Ribstone Indian Ricegrass Germplasm." Crop Science 44 (3):1031–33. [S1]

Jones, T.A., D.C. Nielson, S.K. Caicco, G.A. Fenchel, and S. A. Young. 2005. "Registration of Star Lake Indian Ricegrass Germplasm." Crop Science 45 (4):1666–67. [S1]

Jones, Thomas A., S.R. Winslow, S.D. Parr, and K.L. Memmott. 2010. "Notice of Release of White River Germplasm Indian Ricegrass." Native Plants Journal 11 (2):133–36. [S1]

Kardol, P., J. R. De Long, and D. A. Wardle. 2014. "Local Plant Adaptation across a Subarctic
Elevational Gradient." Royal Society Open Science 1 (3):140141–140141.
https://doi.org/10.1098/rsos.140141. [S1]

Kim, Eunsuk, and Kathleen Donohue. 2013. "Local Adaptation and Plasticity of Erysimum
Capitatum to Altitude: Its Implications for Responses to Climate Change." Journal of Ecology
101 (3):796–805. https://doi.org/10.1111/1365-2745.12077. [S1, S3, QC]

Kitchen, Stanley G. n.d. "Return of the Native: A Look at Select Accessions of North American
Lewis Flax." In: Roundy, Bruce A.; McArthur, E. Durant; Haley, Jennifer S.; Mann, David K.,
Comps. 1995. Proceedings: Wildland Shrub and Arid Land Restoration Symposium; 1993
October 19-21; Las Vegas, NV. Gen. Tech. Rep. INT-GTR-315. Ogden, UT: U.S.D.A. [S1]

Kitchen, Stanley G., and Loren St. John. 1996. "Release Documentation for Maple Grove Lewis
Flax." USDA NRCS Aberdeen Plant Materials Center, Aberdeen, Idaho.
https://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/idpmcrn5639.pdf.
[S1, S2, QC]

Kramer, Andrea T. 2009. "Ecological Genetics of Penstemon in the Great Basin, USA." Ph.D.
Dissertation, University of Illinois at Chicago. [S1, S2]

Kramer, Andrea T., Daniel J. Larkin, and Jeremie B. Fant. 2015. "Assessing Potential Seed Transfer
Zones for Five Forb Species from the Great Basin Floristic Region, USA." Natural Areas
Journal 35 (1):174–88. [S1, S2, QC]

Larsen, Eugene C. 1947. "Photoperiodic Responses of Geographical Strains of Andropogon
Scoparius." Botanical Gazette 109 (2):132–49. [S1, S2, QC]

Leger, Elizabeth A. 2008. "The Adaptive Value of Remnant Native Plants in Invaded Communities:
An Example from the Great Basin." Ecological Applications 18 (5):1226–35. [S1, S2]

Li, Dapeng, Ian T. Baldwin, and Emmanuel Gaquerel. 2015. "Navigating Natural Variation in
Herbivory-Induced Secondary Metabolism in Coyote Tobacco Populations Using MS/MS
Structural Analysis." Proceedings of the National Academy of Sciences 112 (30):E4147–55.
[S1, S2]

Liao, Huixuan, Priscila C.S. Gurgel, Robert W. Pal, David Hooper, and Ragan M. Callaway. 2016.
"Solidago Gigantea Plants from Nonnative Ranges Compensate More in Response to Damage
than Plants from the Native Range." Ecology 97 (9):2355–63.
https://doi.org/10.1002/ecy.1481. [S1, S2]

Link, Steven O., Jeffrey L. Smith, Jonathan J. Halvorson, and Harvey Bolton. 2003. "A Reciprocal
Transplant Experiment within a Climatic Gradient in a Semiarid Shrub-Steppe Ecosystem:
Effects on Bunchgrass Growth and Reproduction, Soil Carbon, and Soil Nitrogen." Global
Change Biology 9 (7):1097–1105. https://doi.org/10.1046/j.1365-2486.2003.00647.x. [S1, S3,
QC]

Love, Stephen L, Robert R Tripepi, and Thomas Salaiz. 2014. "Influence of Harvest Timing and
Storage Interval on Rabbitbrush Seed Germination, Emergence, and Viability." Native Plants
Journal (University of Wisconsin Press) 15 (2):98–108. https://doi.org/10.1353/npj.2014.0017.
[S1]

Mann, Rebecca K. 2016. "Intraspecific Variation in the Response of Elymus Elymoides to Competition from Bromus Tectorum." Ph.D. Dissertation, Utah State University, Logan. [S1, S2]

McArthur, E. Durant, Susan E. Meyer, and Darrel J. Weber. 1987. "Germination Rate at Low Temperature: Rubber Rabbitbrush Population Differences." Journal of Range Management, 530–33. [S1, S2]

McArthur, E. Durant, Richard Stevens, and A. Clyde Blauer. 1983. "Growth Performance Comparisons among 18 Accessions of Fourwing Saltbush [Atriplex Canescens] at Two Sites in Central Utah." Journal of Range Management, 78–81. [S1, S3, QC]

McArthur, E. Durant, and Bruce L Welch. 1982. "Growth Rate Differences among Big Sagebrush [Artemisia Tridentata ] Accessions and Subspecies." Journal of Range Management 35 (3):396–401. https://doi.org/10.2307/3898327. [S1, QC]

McIntyre, Patrick J., and Sharon Y. Strauss. 2014. "Phenotypic and Transgenerational Plasticity Promote Local Adaptation to Sun and Shade Environments." Evolutionary Ecology 28 (2):229–46. https://doi.org/10.1007/s10682-013-9670-y. [S1, S3, QC]

McMillan, Calvin. 1957. "Nature of the Plant Community. III. Flowering Behavior within Two Grassland Communities under Reciprocal Transplanting." American Journal of Botany, 144–53. [S1, QC]

McMillan, Calvin. 1959a. "Nature of the Plant Community. V. Variation within the True Prairie Community-Type." American Journal of Botany, 418–24. [S1, QC]

McMillan, Calvin. 1959b. "The Role of Ecotypic Variation in the Distribution of the Central Grassland of North America." Ecological Monographs 29 (4):286–308. https://doi.org/Genetic Considerations in Ecological Restoration. [S1, S2, QC]

McNaughton, S.J. 1966. "Thermal Inactivation Properties of Enzymes from Typha Latifolia L. Ecotypes." Plant Physiology 41:1736–38. https://doi.org/10.2307/4260909. [S1]

Mealor, Brian A., and Ann L. Hild. 2007. "Post-Invasion Evolution of Native Plant Populations: A Test of Biological Resilience." Oikos 116 (9):1493–1500. https://doi.org/10.1111/j.2007.0030-1299.15781.x. [S1, S2]

Messina, Frank J., Susan L. Durham, James H. Richards, and E. Durant McArthur. 2002. "Trade-off between Plant Growth and Defense? A Comparison of Sagebrush Populations." Oecologia 131 (1):43–51. https://doi.org/10.1007/s00442-001-0859-3. [S1, QC]

Meyer, S. E., and S. G. Kitchen. 1994. "Life History Variation in Blue Flax (Linum Perenne: Linaceae): Seed Germination Phenology." American Journal of Botany 81 (5):528–35. https://doi.org/10.2307/2445726. [S1, S2, QC]

Meyer, S. E., S. G. Kitchen, and S. L. Carlson. 1995. "Seed Germination Timing Patterns in Intermountain Penstemon (Scrophulariaceae)." American Journal of Botany. https://doi.org/10.2307/2445584. [S1, S2]

Meyer, S. E., E. D. McArthur, and G. L. Jorgensen. 1989. "Variation in Germination Response to Temperature in Rubber Rabbitbrush (Chrysothamnus Nauseosus: Asteraceae) and Its

Ecological Implications." American Journal of Botany. https://doi.org/10.2307/2444519. [S1]

Meyer, S. E., and S. B. Monsen. 1992. "Big Sagebrush Germination Patterns: Subspecies and Population Differences." Journal of Range Management 45 (1):87–93. https://doi.org/10.2307/4002533. [S1, S2, QC]

Meyer, SE, J Beckstead, PS Allen, and H Pullman. 1995. "Germination Ecophysiology of Leymus Cinereus (Poaceae)." International Journal of Plant Sciences 156 (2):206–15. https://doi.org/10.1086/297242. [S1, S2, QC]

Meyer, Susan E. 1992. "Habitat Correlated Variation in Firecracker Penstemon (Penstemon Eatonii Gray: Scrophulariaceae) Seed Germination Response." Bulletin of the Torrey Botanical Club 119 (3):268–79. https://doi.org/10.2307/2996758. [S1, S2]

Meyer, Susan E. 1997. "Ecological Correlates of Achene Mass Variation in Chrysothamnus Nauseosus (Asteraceae)." American Journal of Botany 84 (4):471–77. https://doi.org/10.2307/2446023. [S1, S2, QC]

Meyer, Susan E., and Stephanie L. Carlson. n.d. "Seed Germination Biology of Intermountain Populations of Fourwing Saltbush (Atriplex Canescens: Chenopodiaceae)." In: Sosebee, Ronald E.; Wester, David B.; Britton, Carlton M.; McArthur, E. Durant; Kitchen, Stanley G., Comps. 2007. Proceedings: Shrubland Dynamics—fire and Water; 2004 August 10-12; Lubbock, TX. Proceedings RMRS-P-47. Fort Collins, CO: U.S.D.A. [S1, S2]

Meyer, Susan E., Stephanie L. Carlson, and Susan C. Garvin. 1998. "Seed Germination Regulation and Field Seed Bank Carryover in Shadscale (Atriplex Confertifolia: Chenopodiaceae)." Journal of Arid Environments 38 (2):255–67. https://doi.org/10.1006/jare.1997.0321. [S1, S2, QC]

Meyer, Susan E., and Stephen B. Monsen. 1991. "Habitat-Correlated Variation in Mountain Big Sagebrush (Artemisia Tridentata Ssp. Vaseyana) Seed Germination Patterns." Ecology 72 (2):739–42. [S1, S2]

Meyer, Susan E., Stephen B. Monsen, and E. Durant McArthur. 1990. "Germination Response of Artemisia Tridentata (Asteraceae) to Light and Chill: Patterns of Between-Population Variation." Botanical Gazette. https://doi.org/10.1086/337817. [S1, S2]

Meyer, Susan E, and Stanley G Kitchen. 1994. "Habitat-Correlated Variation in Seed Germination Response to Chilling in Penstemon Section Glabri (Scrophulariaceae)." American Midland Naturalist 132 (2):349–65. https://doi.org/10.2307/2426591. [S1]

Miller, Roy V. 1967. "Ecotypic Variation in Andropogon Scoparius and Bouteloua Gracilis." Ph.D. Dissertation, Colorado State University, Fort Collins. [S1, S2]

Miller, Stephanie A., Amy Bartow, Melanie Gisler, Kimiora Ward, Amy S. Young, and Thomas N. Kaye. 2011. "Can an Ecoregion Serve as a Seed Transfer Zone? Evidence from a Common Garden Study with Five Native Species." Restoration Ecology 19 (201):268–76. https://doi.org/10.1111/j.1526-100X.2010.00702.x. [S1, S2, QC]

Monaco, T. A., S. B. Monsen, B. N. Smith, and L. D. Hansen. 2005. "Temperature-Dependent Physiology of Poa Secunda, a Cool Season Grass Native to the Great Basin, United States." Russian Journal of Plant Physiology 52 (5):653–58. https://doi.org/10.1007/s11183-005-0096-

[4](underline). [S1]

Monson, R K, S D Smith, J L Gehring, W D Bowman, S R Szarek, British Ecological Society, and
Functional Ecology. 1992. "Physiological Differentiation within an Encelia Farinosa
Population along a Short Topographic Gradient in the Sonoran Desert." Functional Ecology 6
(6):751–59. [S1, S2]

Moyer, J.L., and R. L. Lang. 1976. "Variable Germination Response to Temperature for Different
Sources of Winterfat Seed." Journal of Range Management 29:320–21. [S1]

Mummey, Daniel L., M.E. Herget, K.M. Hufford, and L. Shreading. 2016. "Germination Timing
and Seedling Growth of Poa Secunda and the Invasive Grass, Bromus Tectorum, in Response
to Temperature: Evaluating Biotypes for Seedling Traits That Improve Establishment."
Ecological Restoration 34 (3):200–208. [S1, S2]

Munda, B.D., S.M. Lambert, and J.C. Garrison. 1990. "Registration of 'Santa Rita' Fourwing
Saltbush." Crop Science 30 (6). [S1]

Nasri, Mohamed, and Paul S. Doescher. 1995. "Effect of Temperature on Growth of Cheatgrass and
Idaho Fescue." Journal of Range Management, 406–9. [S1]

Orodho, A B, R L Cuany, and M J Trlica. 1998. "Previous Grazing or Clipping Affects Seed of
Indian Ricegrass." Journal of Range Management 51 (1):37–41. https://doi.org/none. [S1, S2]

Orodho, Apollo B., and M J Trlica. 1990. "Clipping and Long-Term Grazing Effects on Biomass
and Carbohydrate Reserves of Indian Ricegrass." Journal of Range Management, 52–57. [S1,
S2]

Par, Steve, and Marti Walsh. 2008. Notice of Release of Long Ridge Germplasm Utah Serviceberry.
Upper Colorado Environmental Plant Center, USDA NRCS, Colorado State Agricultural
Experiment Station.
https://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/copmcrn8043.pdf
[S1]

Parsons, Matthew C., Thomas A. Jones, Steven R. Larson, Ivan W. Mott, and Thomas A. Monaco.
2011. "Ecotypic Variation in Elymus Elymoides Subsp. Brevifolius in the Northern
Intermountain West." Rangeland Ecology and Management 64 (6):649–58.
https://doi.org/10.2111/REM-D-09-00143.1. [S1]

Parsons, Matthew C., Thomas A. Jones, and Thomas A. Monaco. 2011. "Genetic Variation for
Adaptive Traits in Bottlebrush Squirreltail in the Northern Intermountain West, United States."
Restoration Ecology 19 (4):460–69. https://doi.org/10.1111/j.1526-100X.2010.00705.x. [S1,
S2]

Pearcy, R W, and R T Ward. 1972. "Phenology and Growth of Rocky Mountain Populations of
Deschampsia Caespitosa at Three Elevations in Colorado." Ecology 53:1171–78.
https://doi.org/10.2307/1935431. [S1, S2, S3, QC]

Pendleton, B. K., and S. E. Meyer. 2004. "Habitat-Correlated Variation in Blackbrush (Coleogyne
Ramosissima: Rosaceae) Seed Germination Response." Journal of Arid Environments 59
(2):229–43. https://doi.org/10.1016/j.jaridenv.2003.12.009. [S1, S2]

Petersen, J L, D N Ueckert, R L Potter, and J E Huston. 1987. "Ecotypic Variation in Selected Fourwing Saltbush Populations in Western Texas USA." Journal of Range Management 40 (4):361–66. https://doi.org/Genetic Considerations in Ecological Restoration. [S1, S3, QC]

Phillips, Nathan C., Daniel T. Drost, William A. Varga, Leila M. Shultz, and Susan E. Meyer. 2010. "Germination Characteristics along Altitudinal Gradients in Three Intermountain Allium spp.(Amaryllidaceae)." Seed Technology 32 (1):15–25. http://www.jstor.org/stable/23433619. [S1, S2]

Polley, H. W., and J. K. Detling. 1988. "Herbivory Tolerance of Agropyron Smithii Populations with Different Grazing Histories." Oecologia 77 (2):261–67. https://doi.org/10.1007/BF00379196. [S1, S2]

Potter, R.L., D.N. Ueckert, and J L Petersen. 1986. "Germination of Fourwing Saltbush Seeds: Interaction of Temperature, Osmotic Potential, and pH." Journal of Range Management, 43–46. [S1]

Quinn, James A., and Richard T Ward. 1969. "Ecological Differentiation in Sand Dropseed (Sporobolus Cryptandrus)." Ecological Monographs 39 (1):61–78. [S1, S2, QC]

Quinn, James A, and Jeffrey D Wetherington. 2002. "Genetic Variability and Phenotypic Plasticity in Flowering Phenology in Populations of Two Grasses." Journal of the Torrey Botanical Society 129 (2):96–106. https://doi.org/10.2307/3088723. [S1, S2]

Ray, Peter M., and William E. Alexander. 1966. "Photoperiodic Adaptation to Latitude in Xanthium Strumarium." American Journal of Botany, 806–16. [S1, S2, QC]

Rice, Kevin J., and Eric E. Knapp. 2008. "Effects of Competition and Life History Stage on the Expression of Local Adaptation in Two Native Bunchgrasses." Restoration Ecology 16 (1):12–23. https://doi.org/10.1111/j.1526-100X.2007.00257.x. [S1, S3, QC]

Richardson, Bryce A., Stanley G. Kitchen, Rosemary L. Pendleton, Burton K. Pendleton, Matthew J. Germino, Gerald E. Rehfeldt, and Susan E. Meyer. 2014. "Adaptive Responses Reveal Contemporary and Future Ecotypes in a Desert Shrub." Ecological Applications 24 (2):413–27. https://doi.org/10.1890/13-0587.1. [S1, S2]

Richardson, Bryce A., Hector G. Ortiz, Stephanie L. Carlson, Deidre M. Jaeger, Nancy L. Shaw, and D. P.C. Peters. 2015. "Genetic and Environmental Effects on Seed Weight in Subspecies of Big Sagebrush: Applications for Restoration." Ecosphere 6 (10). https://doi.org/10.1890/ES15-00249.1. [S1, QC]

Robertson, Phillip A. 1976. "Photosynthetic and Respiratory Responses of Natural Populations of Koeleria Cristata Grown in Three Environmental Regimes." Botanical Gazette 137 (1):94–98. [S1]

Robertson, Phillip A., and Richard T Ward. 1970. "Ecotypic Differentiation in Koeleria Cristata (L.) Pers. from Colorado and Related Area." Ecology 51 (6):1083–87. [S1, S2, QC]

Rogler, George A. 1960. "Relation of Seed Dormancy of Indian Ricegrass (Oryzopsis Hymenoides (Roem. & Schult) Ricker.) to Age and Treatment 1." Agronomy Journal 52 (8):470–73. [S1]

Rowe, Courtney L J, and Elizabeth A. Leger. 2011. "Competitive Seedlings and Inherited Traits: A

Test of Rapid Evolution of Elymus Multisetus (Big Squirreltail) in Response to Cheatgrass Invasion." Evolutionary Applications 4 (3):485–98. https://doi.org/10.1111/j.1752-4571.2010.00162.x. [S1, S2]

Rowe, Courtney L. J., and Elizabeth A. Leger. 2012. "Seed Source Affects Establishment of Elymus Multisetus in Postfire Revegetation in the Great Basin." Western North American Naturalist 72 (4):543–53. https://doi.org/10.3398/064.072.0410. [S1, S3]

Rumbaugh, M.D., and B.M. Pendery. 1993. "Registration of ARS-2892 Munroe Globemallow Germplasm." Crop Science 33 (5). [S1]

Rumbaugh, M.D., B.M. Pendery, H.F. Mayland, and G.E. Shewmaker. 1993. "Registration of ARS-2936 Scarlet Globemallow Germplasm." Crop Science 33:1106–8. [S1]

Sanderson, S. C., H. C. Stutz, and E. D. McArthur. 1990. "Geographic Differentiation in Atriplex Confertifolia." American Journal of Botany. [S1, S2]

Schellenberg, M.P. 2003. "Germination Temperature Response of Two Ecotypes of Winterfat [Kraschenninikovia Lanata (Pursh) Guldenstaedt]." Canadian Journal of Plant Science 83 (1):65–68. [S1]

Shaw, Nancy L., Marshall R. Haferkamp, and Emerenciana G. Hurd. 1994. "Germination and Seedling Establishment of Spiny Hopsage in Response to Planting Date and Seedbed Environment." Journal of Range Management 47 (2):165–74. https://doi.org/10.2307/4002827. [S1, QC]

Shock, Clinton C., Erik B. Feibert, A. Rivera, Lamont D. Saunders, Nancy Shaw, and Francis F. Kilkenny. 2016. "Irrigation Requirements for Seed Production of Five Lomatium Species in a Semiarid Environment." HortScience 51 (10):1270–77. [S1]

Slauson, William L., and Richard T Ward. 1986. "Ecogenetic Patterns of Four Shrub Species in Semi-Arid Communities of Northwest Colorado." The Southwestern Naturalist, 319–29. [S1, S2]

Smith, David Solance, Matthew K. Lau, Ryan Jacobs, Jenna A. Monroy, Stephen M. Shuster, and Thomas G. Whitham. 2015. "Rapid Plant Evolution in the Presence of an Introduced Species Alters Community Composition." Oecologia 179 (2):563–72. https://doi.org/10.1007/s00442-015-3362-y. [S1, S2]

Smith, David Solance, Jennifer A. Schweitzer, Philip Turk, Joseph K. Bailey, Stephen C. Hart, Stephen M. Shuster, and Thomas G. Whitham. 2012. "Soil-Mediated Local Adaptation Alters Seedling Survival and Performance." Plant and Soil 352 (1–2):243–51. https://doi.org/10.1007/s11104-011-0992-7. [S1]

Springfield, H.W. 1968. "Germination of Winterfat Seeds under Different Moisture Stresses and Temperatures." Journal of Range Management, 314–16. [S1]

Springfield, H.W. 1966. "Germination of Fourwing Saltbush Seeds at Different Levels of Moisture Stress." Agronomy Journal 58 (2):149–50. [S1]

Springfield, H.W. 1964. "Some Factors Affecting Germination of Fourwing Saltbush." Rocky Mountain Forest and Range Experiment Station, Forest Service, US Department of Agriculture

25. [S1, S2]

St. Clair, John Bradley, Francis F. Kilkenny, Richard C. Johnson, Nancy L. Shaw, and George
Weaver. 2013. "Genetic Variation in Adaptive Traits and Seed Transfer Zones for
Pseudoroegneria Spicata (Bluebunch Wheatgrass) in the Northwestern United States."
Evolutionary Applications 6 (6):933–48. https://doi.org/10.1111/eva.12077. [S1, S2, QC]

Staub, Jack E, Matthew D Robbins, Yingmei Ma, and Paul G Johnson. 2014. "Phenotypic and
Genotypic Analysis of a US Native Fine-Leaved Festuca Population Reveals Its Potential Use
for Low-Input Urban Landscapes." Journal of the American Society for Horticultural Science
139 (6):706–15. [S1, QC]

Stettler, Jason M., Douglas A. Johnson, B. Shaun Bushman, Kevin J. Connors, Thomas A. Jones,
Jennifer W. Macadam, and David J. Hole. 2017. "Utah Lotus: North American Legume for
Rangeland Revegetation in the Southern Great Basin and Colorado Plateau." Rangeland
Ecology and Management 70 (6):691–99. https://doi.org/10.1016/j.rama.2017.06.002. [S1, S2]

Stevens, Allan R, Val Jo Anderson, and Rachel Fugal. 2014. "Competition of Squirreltail with
Cheatgrass at Three Nitrogen Levels *." American Journal of Plant Sciences 5:990–96. [S1,
QC]

Stinson, Kristina A. 2004. "Natural Selection Favors Rapid Reproductive Phenology in Potentilla
Pulcherrima (Rosaceae) at Opposite Ends of a Subalpine Snowmelt Gradient." American
Journal of Botany 91 (4):531–39. https://doi.org/10.3732/ajb.91.4.531. [S1, S2]

Tilley, Derek J. 2015. "Notice of Release of Amethyst Germplasm Hoary Tansyaster: Selected Class
of Natural Germplasm." Native Plants Journal 16 (1):54–60. [S1, QC]

Tilley, Derek J. 2015. Douglas' Dustymaiden Initial Evaluation Planting  Final Study Report. USDA
NRCS Aberdeen Plant Materials Center.
https://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/idpmcsr13140.pd
f. [S1, S2, QC]

Tisdale, E. W. 1961. "Intraspecific Variation in Festuca." Carnegie Institution of Washington
Yearbook 60:388–91. [S1]

Toole, Vivian K. 1941. "Factors Affecting the Germination of Various Dropseed Grasses
(Sporobolus Spp.)." Journal of Agricultural Research 62:691–715. [S1, S2]

USDA NRCS Bridger Plant Materials Center. 1996. Notice of Release of 'Rimrock' Indian
Ricegrass. USDA NRCS Bridger Plant Materials Center, Montana Agricultural Experiment
Station, Wyoming Agricultural Experiment Station, USDA Agricultural Research Station.
[S1]

USDA NRCS Los Lunas Plant Materials Center. 1973. Supporting Data For the Release of Arriba
Western Wheatgrass. New Mexico State University's Los Lunas Agricultural Science Center,
Colorado State University, New Mexico Department of Transportation, and the USDA Natural
Resources Conservation Service Los Lunas Plant Materials Center. [S1, QC]

Waldron, B L, K B Jensen, A J Palazzo, T J Cary, J G Robins, M D Peel, D G Ogle, and L St John.
2011. "'Recovery', a New Western Wheatgrass Cultivar with Improved Seedling
Establishment on Rangelands." Journal of Plant Registrations 5 (3):367–73.

https://doi.org/Doi10.3198/Jpr2010.09.0527crc. [S1]

Wan, Changgui, Ronald E. Sosebee, and Bobby L. McMichael. 1998. "Water Relations and Root
Growth of Two Populations of Gutierrezia Sarothrae." Environmental and Experimental
Botany 39 (1):11–20. https://doi.org/10.1016/S0098-8472(97)00021-X. [S1]

Wan, Changgui, Ronald E. Sosebee, and Bobby L. McMichael. 1995. "Water Acquisition and
Rooting Characteristics in Northern and Southern Populations of Gutierrezia Sarothrae."
Environmental and Experimental Botany 35 (1):1–7. https://doi.org/10.1016/0098-
8472(94)00038-7. [S1, S2]

Wan, Changgui, Ronald E. Sosebee, and Bobby L. McMichael. 1996. "Lateral Root Development
and Hydraulic Conductance in Four Populations of Gutierrezia Sarothrae." Environmental and
Experimental Botany 36 (2):157–65. https://doi.org/10.1016/0098-8472(96)01008-8. [S1]

Ward, Richard T. 1969. "Ecotypic Variation in Deschampsia Caespitosa (L.) Beauv. from
Colorado." Ecology 50 (3):519. https://doi.org/10.2307/1933914. [S1, S2, QC]

Waser, Nickolas M., and Mary V. Price. 1985. "Reciprocal Transplant Experiments with
Delphinium Nelsonii (Ranunculaceae): Evidence for Local Adaptation." American Journal of
Botany, 1726–32. [S1, S2, S3]

Wood, M. Karl, Robert W. Knight, and James A. Young. 1976. "Spiny Hopsage Germination."
Journal of Range Management, 53–56. [S1]

Workman, John P, and Neil E West. 1967. "Germination of Eurotia Lanata in Relation to
Temperature and Salinity." Ecology 48 (4):659–61. [S1]

Workman, John P, and Neil E West. 1969. "Ecotypic Variation of Eurotia Lanata Populations in
Utah." Botanical Gazette 130 (1):26–35. https://doi.org/10.2307/2473599. [S1, S2]

Young, J A, C D Clements, and T Jones. 2003. "Germination of Seeds of Big and Bottlebrush
Squirreltail." Journal of Range Management 56 (3):277–81. https://doi.org/10.2307/4003819.
[S1]

Young, JA, RA Evans, and DE Palmquist. 1989. "Big Sagebrush (Artemisia Tridentata) Seed
Production." Weed Science 37 (1):47–53. http://www.jstor.org/stable/10.2307/4044754. [S1,
S3, QC]

Young, James A., and Raymond A. Evans. 1989. "Reciprocal Common Garden Studies of the
Germination of Seeds of Big Sagebrush (Artemisia Tridentata)." Weed Science 37 (3):319–
325. https://doi.org/none. [S1, QC]

Young, James A, Raymond A Evans, and B L Kay. 1984. "Persistence and Colonizing Ability of
Rabbitbrush Collections in a Common Garden." Journal of Range Management 37 (4):373–77.
https://doi.org/10.2307/3898715. [S1, QC]

Zhang, Huarong, Laura E. DeWald, Thomas E. Kolb, and Dan E. Koepke. 2011. "Genetic Variation
in Ecophysiological and Survival Responses to Drought in Two Grasses: Koeleria Macrantha
and Elymus Elymoides." Western North American Naturalist 71 (1):25–32.
https://doi.org/10.3398/064.071.0104. [S1, S2]

1649

## Appendix 3. Additional Results

*Additional results of literature summary*

Dicots accounted for 23.6% of the taxa and 42.8% of the experiments in the final pool of reviewed literature. Regional taxa accounted for 47.2% of the taxa and 48.9% of experiments, widespread taxa accounted for 26.0% of taxa and 36.7% of experiments, narrow taxa accounted for 24.4% of taxa and 13.5% of experiments, and endemic taxa accounted for 2.4% of taxa and 0.9% of experiments. Perennials accounted for 46.3% of taxa and 32.1% of experiments, long-lived perennials accounted for 32.5% of taxa and 39.4% of experiments, short-lived perennials accounted for 14.6% of taxa and 25.1% of experiments, annuals accounted for 5.7% of taxa and 3.1% of experiments, and biennials accounted for 0.8% of taxa and 0.3% of experiments. Primarily outcrossing plants accounted for 71.4% of taxa and 72.2% of experiments, primarily selfing plants accounted for 11.1% of taxa and 14.1% of experiments, and plants with mixed mating accounted for 17.5% of taxa and 13.8% of experiments.

*Additional results for quantitative comparison of trait-by-environment associations*

Appendix 3 Figures 1-16. For each trait/environment correlation (16 combinations), the results of correlation coefficients (with 95% confidence intervals) for the pairwise comparisons, for each population in each experiment, between the difference in phenotypic trait and environmental characteristic at the collection location, while controlling for geographic distance among populations (see methods). MAT = Mean Annual Temperature, MAP = Mean Annual Precipitation. Study number identifies the particular experiment in the quantitative comparison dataset in the electronic supplementary material, "# pops." is the number of populations included in each study, and "dist." is the average pairwise distance between populations (in km). Grasses

1673    are shown in green, shrubs in blue, and forbs in orange. The overall effect size and confidence

1674    intervals, across all studies, is shown in gray at the bottom of each figure. See main text, Table 1,

1675    for descriptions of phenotypic traits.

1676

1677

1678

1679

1680

1681

1682

1683

1684

1685

1686

## MAP
### Height

1687

1688

1689

1690

1691