Title:

Interactions between non-structured domains of FG- and non-FG-nucleoporins coordinate the ordered

assembly of the nuclear pore complex in mitosis

Author affiliation:

Hide A. Konishi[§] and Shige H. Yoshimura[*]

Laboratory of Plasma Membrane and Nuclear Signaling, Graduate School of Biostudies, Kyoto

University, Yoshida Konoe, Sakyo-ku, Kyoto 606-8501, Japan

*Corresponding author:

Shige H. Yoshimura

Laboratory of Plasma Membrane and Nuclear Signaling,

Graduate School of Biostudies, Kyoto University,

Yoshida Konoe, Sakyo-ku, Kyoto 606-8501, Japan

Tel & Fax: +81-75-753-7906

E-mail: yoshimura@lif.kyoto-u.ac.jp

[§]Current address: Laboratory of Chromosome and Cell Biology, The Rockefeller University, New

York, NY 10065, USA

Summary

In this study, we examined how channel-forming subunits of the nuclear pore complex (NPC) are assembled into a selective channel within a highly structured scaffold ring during post-mitotic assembly. We focused on non-structured domains of the scaffold Nups and performed *in vitro* self-assembled particle assays with those derived from channel-forming FG-Nups. We found that non-structured domains of ELYS and Nup35N interacted with channel-forming FG-Nups to form a self-assembled particle. Sequential addition of FG-Nups into the scaffold particle revealed that ELYS, which initiates post-mitotic NPC reassembly, interacts with early assembling FG-Nups (Nups98 and 153) but not middle stage-assembling FG-Nups (Nups58 and 62). Nup35, which assembles between the early and middle stages, facilitated the assembly of Nup62 into the early assembling Nups both *in vitro* and *in vivo*. These results demonstrate that ELYS and Nup35 have a role of facilitator in the ordered assembly of channel-forming FG-Nups during mitosis.

Keywords

Nuclear pore complex, Intrinsically disordered region, Molecular crowding, Self-assembly

Introduction

One of the largest protein complexes in eukaryotic cells is the nuclear pore complex (NPC), which spans the nuclear membrane and functions as the gate for macromolecular trafficking between the cytoplasm and nucleoplasm (1). The NPC is composed of several copies of more than 30 different nucleoporin (Nup) subunits (2). The NPC comprises several structurally distinct domains, such as the nuclear basket, cytoplasmic fibril, scaffold, and central channel, each of which is composed of defined sets of Nups. A number of studies using crystallography and electron tomography have revealed the molecular structures of the Nups and sub-complexes that comprise the scaffold (3). The entire architecture of the scaffold was then composited by integrating all available structural information for each of the subunits and sub-complexes with information regarding protein-protein interactions (4). Despite the low degree of amino acid conservation between Nups among different species, the entire architecture of the scaffold turned out to be well conserved. The outer ring containing the Nup107-160 sub-complex (Y-complex) is located at both the cytoplasmic and nucleoplasmic rims, and the inner ring, which contains the Nup93 sub-complex, is located deeper in the pore and connects the membrane and the central channel (3–5). Interactions between structured Nups and sub-complexes is thought to be stereospecific, resulting in a rigid architecture in the nuclear envelope.

In contrast to the scaffold, the central channel is formed primarily by a number of non-structured domains of Nups, which assemble via promiscuous interactions to form a highly crowded environment within the pore (6). Nups containing a large number of phenylalanine-glycine (FG) motifs (FG-Nups) in a long intrinsically disordered region (IDR-Nups) (7, 8) form a molecular sieve-like environment that functions as a selective barrier with respect to molecular transport. Small molecules (<40 kDa) diffuse passively, but large molecules (>40 kDa) are repelled unless they are

carried by an appropriate transport receptor, such as a karyopherin (2, 9). Although several

experimental and/or schematic models have been proposed to explain the structure and function of

the barrier (10, 11), the molecular mechanism through which a large number of promiscuous

interactions among IDR-Nups constructs a functional barrier within the rigid scaffold remains poorly

understood.

Several lines of evidence have provided clues to enhance understanding of the mechanism

of channel formation. Several FG-Nups self-assemble and form a hydrogel (10, 12) via hydrophobic

interactions between Phe residues at a high protein concentration ($\approx$10 mM) (13). A recent study

reported that FG-domains of Nup98 spontaneously self-assemble into particles *in vitro* at physiologic

concentrations ($\approx$10 μM) (13). These findings suggest that the central channel of the NPC forms via

self-assembly of non-structured FG-Nups. Another study using a crowding-sensitive fluorescent

probe demonstrated that protein crowding is not homogeneous throughout the channel; "protein-rich"

domains lie at both rims of the channel but not in the central cavity (6). This *in vivo* evidence

suggests that even though FG-Nups self-assemble into protein-rich phases, several sets of Nups

preferentially assemble into distinct protein-rich phases, possibly one at the cytoplasmic rim and

another at the nucleoplasmic rim of the central channel. Furthermore, live-cell imaging of individual

Nups during post-mitotic assembly of the NPC demonstrated that individual FG-Nups assemble

orderly on the chromosome surface (6, 14, 15), suggesting that the assembly of FG-Nups is driven by

a specific molecular mechanism and does not proceed solely via a random self-assembly–driven

mechanism. Collectively, this evidence suggests that although FG-Nups self-assemble into

protein-rich phases, a specific molecular mechanism regulates and/or coordinates the assembly of

FG-Nups to construct distinct protein-rich phases within the central channel.

In this study, we focused on IDRs from FG-Nups in the central channel, as well as those

from non–FG-Nups in the scaffold, and examined how they assemble and interact using an *in vitro*

self-assembled particle assay. The results demonstrated that some non–FG-Nups, Nup35 and ELYS,

interact with distinct sets of FG-Nups in a self-assembled particle and facilitate interactions with

another set of FG-Nups. Our data suggest that IDRs in non–FG-Nups play an important role in the

ordered assembly of FG-Nups to construct the functional selective barrier in the central channel of

the NPC.

Results

Both FG-rich and non-FG IDRs form self-assembled particles in a crowded environment

Almost all FG-Nups contain a long stretch of IDRs (Figure S1A), each of which normally contains a

number of FG motifs. In contrast, most non–FG-Nups do not carry IDRs, with the exception of

Nup35, ELYS, and Tpr (Figure S1A). Nup35, which is found in the inner ring complex, contains a

long IDR at its N-terminus (a.a. 1-160). ELYS, a component of the Y-complex in the outer ring,

contains IDRs in the C-terminal half of the molecule (a.a. 1310-1559 and 1851-2275). Tpr is a

component of the nuclear basket and has an extremely high number of IDRs throughout the molecule

(especially in the C-terminal half). We focused on examining the interactions of these IDRs from

non–FG-Nups, together with a number of FG-rich IDRs from FG-Nups (Nups214, 153, 98, 62, 58,

and 50). IDR fragments examined in this study are summarized in Figure 1A and Table S1. Most

IDRs from FG-Nups contain a large number of FG motifs, whereas Nup214-C and Nup50FG have

only a few FG motifs and a relatively high content of charged residues (Figure 1A). The IDRs from

non–FG-Nups have almost no FG motifs and are extremely rich in charged residues (Figure 1A).

Purified IDR fragments were labeled with fluorescent dye, diluted into varying concentrations of

polyethylene glycol (PEG)-containing buffer, and then observed under a fluorescence microscope.

As summarized in Figures 1B-I, some of the IDR fragments self-assembled and form condensed

particles in a PEG-dependent manner. Among FG-rich IDRs, Nups214FG, 153FG, 98FG, and 62FG

self-assembled (Figures S1B-E), whereas Nups214-C, 58FG-N, 58FG-C, and 50FG did not (Figure

S1F-I). The ability of forming particles appeared to correlate well with the FG content (Figure 1A).

A FG-rich IDR from Nup214 (Nup214FG, containing 28 FG motifs) formed particles at a low PEG

concentration, whereas Nup214-C (4 FG motifs) and Nup50FG (2 FG motifs) did not form particles

even at a high PEG concentration (Figures 1D, S1H and S1I), suggesting that hydrophobic

interactions are involved in particle formation by FG-rich IDRs (see later section).

Interestingly, some of the non-FG IDRs (Nup35N, ELYS-C1, -C2, and Tpr-IDR1) also

self-assembled and form particles even though they do not contain FG motifs, but others did not

(ELYS-M and Tpr-IDR2) (Figures 1F-I, S1J-M, -N, and -O). Amino acid sequence analyses revealed

that all of these IDRs contained a number of charged residues, but only ELYS-M and Tpr-IDR2 had

a large negative net charge (>25, Figure 1A and Table S1). This result suggests that particle

formation by non-FG IDRs is driven predominantly by electrostatic interactions, but a large negative

net charge may interfere with particle formation. Analysis of the secondary structure revealed that

Tpr-IDR1, but not other IDRs, had a relatively high degree of coiled-coil structure (Figure S2B),

although all of them had a high disorder probability. Circular dichroism (CD) spectra of Tpr-IDR1

indeed exhibited peaks at 222 nm and 207 nm, which indicates the presence of an α-helix (Figure

S2C). These results suggest that coiled-coil interactions contribute to the self-assembly of Tpr-IDR1,

as demonstrated in a previous study (16).

FG-rich IDRs in self-assembled particle exhibited a very slow exchange rate with the

extra-particle environment and low mobility within the particle, as determined by fluorescence

recovery after photobleaching (FRAP) analysis (Figures S2D-F) (13). In contrast, non-FG IDRs

within particles exhibited faster recovery than FG-Nups (Figures S2G-I). This result is consistent

with the previous report describing the difference between hydrophobic and electrostatic interaction

in the self-assembly of proteins (17).

Self-assembly of FG-rich IDRs and some non-FG IDRs is driven by hydrophobic interactions

We next characterized the protein-protein interactions that induce self-assembly of IDR-Nups by

employing two reagents to quantify hydrophobic and electrostatic interactions: 1,6-hexanediol, an

aliphatic alcohol that weakens hydrophobic interactions (11); and NaCl, which strengthens

hydrophobic interactions but weakens electrostatic interactions. Self-assembly of FG-rich IDRs

(Nup98FG and Nup214FG) was significantly inhibited with increasing concentrations of

1,6-hexanediol (Figures 2A, B and S3A), indicating the involvement of hydrophobic interactions in

particle assembly. NaCl had less effect on FG-rich IDRs compared with 1,6-hexanediol; assembly of

Nup214FG increased slightly (15-20%), whereas that of Nup98FG declined by 35-40% with

increasing NaCl concentrations (Figures 2A, 2B, and S3B). These results suggest that the particle

formation of FG-rich IDRs is facilitated by hydrophobic interaction.

In a clear contrast, the self-assembly of non-FG-Nup (Tpr-IDR1) was increased by

1,6-hexanediol but reduced with increasing NaCl concentrations (Figures 2C, S3A, and S3B),

demonstrating that electrostatic interactions govern the self-assembly of Tpr-IDR1. Interestingly,

Nup35N, which is also a non-FG IDR but with a small number of charged residues (Figure 1A) and

high content of hydrophobic residues (Table S1), behaved as an FG-rich IDR; the particle mass

increased as the NaCl concentration increased and declined as the 1,6-hexanediol concentration

increased (Figures 2D, S3A, and S3B). Furthermore, ELYS-C1, a non-FG IDR containing a number

of charged residues (Figure 1A), exhibited both electrostatic and hydrophobic characteristics

(Figures 2E, S3A, and S3B). The assembly of ELYS-C1 was suppressed both by NaCl and

1,6-hexanediol, which could have been due to its unique amino acid sequence, which is characterized

by short stretches (~20 amino acids) of charged and hydrophobic residues alternating with each other

throughout the fragment (~200 amino acids) (Figure S3C). These results are summarized in Figure

2F. Most of the FG-rich IDRs self-assembled via hydrophobic interactions, but some non-FG IDRs

(ELYS-C1 and Nup35N) also self-assembled via hydrophobic interactions.

IDRs in ELYS and Nup35 interact with FG-IDRs

Two different IDRs, each labeled with a different fluorescent dye, were mixed in the same tube and

observed under a fluorescence microscope to examine their interaction (Figure 3). One IDR was

defined as the "host", and the partition coefficient (PC) of the other IDR (client) in the host particle

was then quantified (see Experimental Procedures). BSA was used as a standard of client. As

summarized in Figure 3A and B, IDRs from FG-Nups generally co-localized well with FG-rich IDRs,

although the individual PC value varied widely. FG-rich IDRs with a high FG content and a low

content of charged residues (Nups214FG, 62FG, 98FG, and 153FG) exhibited high PC values,

whereas IDRs with a low FG content and a high content of charged residues (Nups214-C, 58FG-C,

and 50FG) exhibited low PC values, which are comparable to the standard BSA (Figures 3A and C).

The interactions between non-FG IDRs (ELYS-C1, Nup35N, and Tpr-IDR1) and between

non-FG IDRs and FG-rich IDRs were also examined (Figures 3C and D). Tpr-IDR1, which

self-assembled primarily via electrostatic interactions (Figure 2C), did not co-localize with most of

the FG-rich IDRs except for Nup50FG (Figure 3C and D). In contrast, Nup35N, a non–FG-Nup that

self-assembled via hydrophobic interactions (Figure 2D), co-localized with most of the FG-rich IDRs

(Nups153FG, 98FG, 58FG-N, 62FG, and 214FG) (Figures 3C and D). Similarly, ELYS-C1

co-localized with FG-rich IDRs (Figures 3C and D). These results agree well with those shown in

Figures 2D and E, showing that Nup35 and ELYS-C1, both of which are derived from non–FG-Nups,

self-assembled via hydrophobic interactions, implying that Nup35 and ELYS-C1 play a role in the

linkage between FG-Nups in the central channel and non–FG-Nups in the scaffold.

ELYS-IDR interacts with early assembling but not middle stage–assembling FG-Nups

In mitosis, the NPC disassembles during prophase and reassembles during anaphase via the ordered

assembly of individual Nups. ELYS has been shown to play a critical role in the initial step of

reconstruction by binding to the chromosome surface (18). Nups located on the nucleoplasmic side

of the central channel (e.g., Nups153 and 98) and some of the scaffold Nups (e.g., Nups133 and 160)

(early assembling Nups) follow ELYS. Nups in the inner ring complex (Nups35 and 155) and

membrane-spanning Nups (Pom121 and Ndc1) then assemble, which is followed by Nups in the

central cavity (Nups54, 58, and 62) (14, 15) (Figure 4A). To investigate how the ordered assembly of

channel-forming FG-Nups is regulated, we sequentially added FG-rich IDRs as clients and examined

how they are incorporated into the ELYS-C1 particle as a host (Figure 4B, upper panel).

190   IDRs from early-assembling FG-Nups (Nups98FG and 153FG) were rapidly incorporated

into the ELYS-C1 particle after addition (Figures 4C and S4A), consistent with the results shown in

Figure 3D and F. Interestingly, lower amounts of 58FG-N (which assembles during the middle stage)

were incorporated into the ELYS particle, compared with the early assembling Nups (Figure 4C).

Furthermore, Nup62FG (which assembles during the middle to late stages) was not incorporated into

the ELYS particle (Figure 4C), despite generally co-localizing in the same particle when mixed

(Figures 3D and F). These results demonstrate a clear correlation between the interaction between

ELYS particle and the order of the FG-Nups during post-mitotic reassembly; ELYS, which binds to

the chromosome surface at the beginning of NPC reassembly, readily recruits Nups98 and 153 (early

assembling Nups) but not Nups58 and 62 (middle stage–assembling Nups).

Nup35 facilitates the assembly of middle stage–assembling Nups

In addition to ELYS, Nup35N also interacted with FG-rich IDRs in our particle assay (Figure 3). As

Nup35 appears on the chromosome surface during the early to middle stages of post-mitotic

reassembly (19) (Figure 4A) and interacts with the structured domain of Nup62 (4), it can be

hypothesized that it facilitates the assembly of middle stage–assembling FG-Nups. We therefore

tested the possibility of Nup35N to facilitate the incorporation of middle stage–assembling Nups into

ELYS particles. Nup35N, as 1st client, was pre-incorporated into the ELYS-C1 particle in part to

mimic the early to middle stages of post-mitotic reassembly (Figure 4B lower panel). As shown in

Figure 4D, Nup35N was incorporated into the ELYS-C1 particle at a rate comparable to that of other

early assembling Nups (Nups153FG and 98FG) (Figure 4C), consistent with results shown in

Figures 3D and F. The middle stage–assembling Nup58 or 62 was then added, and its incorporated

fluorescence signal in the particle was quantified. Interestingly, incorporation of Nup62FG into the

ELYS-C1 particle was enhanced when the particle contained Nup35N (Figure 4E). A similar effect

of Nup35N was observed with the incorporation of Nup58FG, although the effect was smaller

(Figure 4F). It should be noted that no such stimulatory effect was observed with Nup98FG, an early

assembling FG-Nup; pre-incorporation of Nup98FG into ELYS-C1 particle did not facilitate the

incorporation of Nups62 and 58 (Figures 4E and F). These results demonstrate that Nup35N

facilitates the interaction of middle stage–assembling Nups with early assembling Nups.

Finally, we examined the role of Nup35 in the post-mitotic NPC reassembly process *in*

vivo. An EGFP-fused FG-Nup (Nup62 or 153) was expressed in Nup35 (or control) knock-down

(KD) cells, and the appearance of fluorescence signals on the chromosome rim during telophase and

anaphase was observed by time-lapse imaging. Signal associated with Nup62 began to appear on the

chromosome surface ~10 min after anaphase onset and gradually increased for 30 min until it

reached a plateau (Figure 4G). In contrast, Nup35-KD cells exhibited a significant delay in Nup62

assembly; it began to appear 10 min after the anaphase onset, which is similar to control knock down,

increased slowly , and did not reach the plateau even after 50 min. It should be noted that the effect

of Nup35 KD was not observed in the assembly of Nup153, an early assembling Nup; it began to

appear 3-5 min after the anaphase onset, and reached the plateau at 12-15 min in both control and

Nup35-KD cells (Figure 4H). These results provide *in vivo* evidence that Nup35 facilitates the

assembly of middle stage–assembling Nups during post-mitotic NPC reassembly.

Discussion

In this study, we investigated the self-assembling properties of IDRs derived from FG-Nups and

non–FG-Nups as well as their interaction patterns using an *in vitro* self-assembled particle assay. The

results demonstrated that i) many FG-rich IDRs self-assemble via hydrophobic interactions and

co-assemble (co-localize) well in the same particle; ii) some IDRs from non–FG-Nups self-assemble

via electrostatic interactions, whereas others (Nup35N and ELYS-C1) assemble via hydrophobic

interactions and interact with FG-rich IDRs; iii) The IDR of ELYS (ELYS-C1) interacts with early

assembling FG-rich IDRs (Nups98FG and 153FG) but not with middle stage–assembling Nups

(Nups58 and 62); and iv) Nup35N facilitates the assembly of middle stage–assembling FG-rich IDRs

(Nups58FG and 62FG) into the ELYS particle. These results demonstrate the important roles of

IDRs from scaffold Nups in coordinating the assembly of FG-Nups. It can be hypothesized that

although self-assembly of Nups is generally driven via promiscuous interactions among IDRs,

"ordered" assembly of Nups and the construction of a functional selective channel during the

post-mitotic reassembly process could be mediated by a few "facilitator subunits," such as ELYS and

Nup35. Functional analyses of these mediator subunits could elucidate the mechanism underlying the

construction of the central channel, and together with structural information regarding the scaffold,

such analyses could elucidate the relationship between the structure and function of the mitotically

formed NPC.

250   The molecular architecture of the scaffold has been studied extensively using

crystallography, electron tomography, and mass spectroscopy (3, 20). Nup35 is a component of the

Nup93 complex, which is the major structural unit of the inner ring, and interacts with Nups in the

same complex (Nups93, 155, and 205) as well as other Nups (Nup98 and Ndc1) and the membrane

(4). A recent study reported the crystal structure of the Nup93 complex with structured domains of

the central channel components Nups62, 58, and 54 (21), demonstrating that the Nup93 complex

links the scaffold and central channel. Although the entire structure of Nup35 was not elucidated in

the crystallographic study, it is possible that the IDR of Nup35 associates with the IDRs of Nups62,

54, and 58. Our *in vitro* self-assemble particle assay indeed demonstrated that Nup35N, which has no

FG motif, interacted with most of the FG-rich IDRs examined (Figure 3D and F).

260   From a structural perspective, the interaction between Nup35N and FG-rich IDRs could be

attributed to its unique amino acid composition, which is rich in proline (≈20%) and glycine (≈11%)

(Figure S3D). A large number of proline residues could aid the formation of self-assembling

particles, as well as hydrophobic interactions with FG-rich IDRs. It was previously demonstrated that

interactions between proline-rich motifs in N-WASP and the SH3 domain of a partner molecule are

important for multivalent interactions between them (22). Proline is also known to break secondary

structures (α-helixes and β-sheets) (23–25) and confer a high degree of flexibility on polypeptides.

Such flexibility could increase the mobility of molecules within the particle, thus facilitating the

incorporation of other molecules into the host particle.

269   In addition to Nup35, a fragment of the ELYS IDR (ELYS-C1) also interacted with

FG-rich IDRs, indicating that it can also function as a facilitator between the scaffold and central

channel. ELYS contains several distinct functional domains in addition to its IDR and was shown to

play an important role in the initial step of post-mitotic NPC reassembly. ELYS binds to the

chromosome surface via its AT-hook domain (aa. 1971-1983) (18), which recruits early assembling

Nups such as the Nup107-160 complex and Nups98 and 153. The AT-hook domain is located within

the ELYS-C1 fragment examined in this study. Therefore, it is plausible that the C-terminal domain

of ELYS functions as a platform for the successive assembly of FG-Nups via hydrophobic

interactions among IDRs. In contrast, the structured N-terminal domain interacts directly with a

helical domain of Nup120 (human Nup160) in the Y-complex (26). Therefore, it seems likely that

the structured N-terminal domain of ELYS acts as a stereospecific binding site for the scaffold

Y-complex, whereas the non-structured C-terminal region binds to the chromosome surface and

initiates the ordered assembly of FG-Nups (Figure 4I).

Considering the facilitator function of ELYS and Nup35 in conjunction with the

self-assembling characteristics of FG-Nups, it seems likely that reconstruction of the functional NPC

during mitosis is driven by both promiscuous interactions among IDRs and stereospecific

interactions among scaffold Nups, which are linked by unique Nup IDRs such as ELYS and Nup35N

(Figure 4I). ELYS initiates the reassembly process by binding to the chromosome surface via its

AT-hook domain. ELYS then recruits both scaffold Nups (Nups in the Y-complex), primarily via

stereospecific interactions involving the N-terminus, and IDRs of early assembling FG-Nups

(Nups153 and 98), primarily via promiscuous interactions involving the C-terminal IDR. The middle

stage–assembling Nups58 and 62 cannot enter this IDR-rich phase at this stage. Nup35 eventually

assembles into the IDR-rich phase formed at the ELYS C-terminus and then recruits the middle- to

late-stage assembling Nups58 and 62. Simultaneously, the N-terminus of Nup35 appears to interact

with the scaffold via the Nup93 complex (4) to link the FG-rich IDR phase to the entire scaffold. In

the later stages, cytoplasmic Nups (Nups214 and 358) in addition to other Nups (Tpr, etc.) associate,

potentially inducing further rearrangements of the IDR phase to establish the functional selective

barrier within the NPC. It could be intriguing to investigate how phosphorylation/dephosphorylation

of Nups (such as the Nup107-160 complex (27), Nup35 (28), and Nup98 (29) is involved in the

regulation of Nup reassembly.

Materials and Methods

Amino acid sequence analyses

The disordered propensity was analyzed by DisProt server

(http://disorder.compbio.iupui.edu/metapredictor.php) with a PONDR-fit algorithm. The net charge

and the charge distribution were analyzed by a protein calculator (INNOVAGEN,

https://pepcalc.com/protein-calculator.php), and EMBOSS explorer

(http://www.bioinformatics.nl/cgi-bin/emboss/charge?_pref_hide_optional=1), respectively. The

hydrophobicity was analyzed by ProtScale server in ExPASy (https://web.expasy.org/protscale/) by

using Kyte & Doolittle method. COILS server in ExPASy

(https://embnet.vital-it.ch/software/COILS_form.html) was used for coiled-coil analysis.

DNA construction and protein purification

cDNA fragments encoding human Nup50FG (aa. 1–145), rat Nup62FG (aa. 1–268), human

Nup98FG (aa. 214–480), human Nup153FG (aa. 1190–1464), human Nup214FG (aa. 1786–2090),

human Nup214-C (aa. 1251–1550), human Nup35N (aa. 1–160), human Tpr-IDR1 (aa. 1451–1689),

and human Tpr-IDR2 (aa. 1690–1937) were amplified by PCR from the full-length cDNA and

sub-cloned into pET28-a (+) or pET28-b (+) (Novagen). Human Nup58FG-N (aa. 1–250), -C (aa.

438–599), human ELYS-IDR-M (aa. 1310-1559), human ELYS-IDR-C1 (aa. 1851-2033), and

human ELYS-IDR-C2 (aa. 2034-2275) were artificially synthesized with the codon usage optimized

for bacterial expression (Thermo Fisher Scientific), and sub-cloned into pET28-a (+). For IDRs

lacking endogenous cysteine residue, a single cysteine was added to the carboxyl-terminal end of the

fragment either by PCR or during artificial synthesis.

The expression of His$_6$-tagged IDR-fragment in *Escherichia coli* cells

(BL21-CodonPlus(DE3)-RIL, Agilent Technologies, Wilmington, DE) was induced by 0.5 mM

IPTG at 37°C for 3 hours. The cells were harvested by centrifugation (5,000x g, 4°C, 15 min), and

dissolved in Denaturing Buffer (8 M urea, 100 mM NaPO$_4$, 10 mM Tris-HCl, 5 mM

2-mercaptoethanol (2-Me) (Nacalai Tesque, Kyoto, Japan), 10 mM imidazole, pH 8.0) at 4°C for 16

hours with a gentle agitation. An insoluble fraction was separated by centrifugation (10,000x g, 4°C,

30 min), and the supernatant was then incubated with Ni-NTA agarose beads (QIAGEN) at 4°C for 2

hours or at room temperature for 30 min. The beads were washed for several times with Denaturing

Buffer containing 20 or 25 mM imidazole. His-tagged protein was eluted by 200 or 300 mM

imidazole in Denaturing Buffer.

Purified protein was dialyzed against 0.1% (v/v) trifluoroacetic acid (TFA) (Nacalai

Tesque) with 2 mM 2-Me at 4°C for 3 hours, and subsequently against 0.1% (v/v) TFA with 2 mM

2-Me at 4°C for ~20 hours. It was finally dialyzed against 0.05% (v/v) TFA, without 2-Me at 4°C for

2-3 hours, lyophilized (FDU-2200, EYELA), and stored at 4°C until use. Purified protein for CD

measurement (Nup58CC (aa. 256-381) and Nup58FG-N) was first dialyzed against a buffer

containing 100 mM L-arginine (50 mM KPO$_4$, 100 mM L-arginine, 150 mM NaCl, 2 mM 2-Me),

then against 50 mM L-arginine, and finally against 50 mM HEPES (pH 7.4) with 100 mM NaCl and

2 mM 2-Me at 4°C for 1 day, and stored -80°C by flash freezing with liquid nitrogen. The final

protein concentration was determined by using Coomassie brilliant blue (CBB) staining, with bovine

serum albumin (BSA) as a standard.

Protein labeling

Purified IDR-Nups' fragments were labeled by utilizing maleimide reaction with artificially added

cysteine in the carboxyl terminal of each IDR-fragment or endogenous cysteine for some Nups

(Nups214 and Nup58FG-C). Using ATTO dyes (ATTO390, 488, and 610, ATTO-TEC) dissolved in

anhydrous DMSO (Nacalai Tesque), the appropriate volume of the ATTO solution was added in the

protein stock buffer (final, 10 mM) before dissolving the lyophilized proteins. To stop the maleimide

reaction, final 5 mM of dithiothreitol (DTT, Nacalai Tesque) was added into the denaturing protein

stock solution and incubated at 4℃ for more than 12 hours.

In vitro self-assemble particle assay

Lyophilized proteins were dissolved in a small volume (~50 µL) of protein stock buffer (2 M

Guanidine hydrochloride, 100 mM Tris-HCl, 10 mM HEPES, pH 8.0) and stored at 4℃ for 1 day.

This stock solution (≈1200 µM) was quickly diluted (>100 times) into the assay buffer (50 mM

HEPES, 100 mM NaCl, pH 7.4) containing varying concentrations of crowding agents, salt, or

1,6-hexanediol (Nacalai Tesque), after removing aggregates by centrifugation (5 min at 10,000 x g).

The final protein concentration was ≈ 2 µM for the client proteins and ≈3 µM for the host protein.

Microscopic observation, image processing and analysis

Protein solution containing self-assembled particles was applied to a 96-well glass bottom plate

(IWAKI, Shizuoka, Japan) or a custom-made flow chamber. In the mixing assay described in Figure

3, images were taken 5-10 min after the dilution of mixing host and client proteins in the chamber. In

this time range, no significant change in the particle formation was observed. In the sequential

addition of two different clients, 1st client protein was first added to the host particle (ELYS-C1).

After 300 sec, fluorescence signal was captured by microscope, and 2nd client protein was added.

After another 300 sec, the fluorescence signal of the 2nd client was observed under the microscope

and subjected to the quantification. Microscopic observation was performed using FV3000 confocal

laser-scanning microscope (OLYMPUS, Tokyo, Japan) using $60 \times$ NA1.42 objective lens

(PLAPON60XO) with oil immersion. For the total particle mass analysis, 10 images were randomly

taken in 2x zoom field of $60 \times$ objective lens. The excitation wavelength against ATTO390, 488, and

610 was 405 nm, 488 nm, and 564 nm, respectively. Sampling speed was 8.0 µs per pixel.

For *in vivo* live cell imaging, EGFP-fused Nups62 or 153 was co-expressed with mPlum

(Clontech)-fused histone H3 in HeLa cells cultured in DMEM (SIGMA) supplemented with 10%

FBS (Hyclone). Before the microscopic observations, the culture medium was replaced with phenol

red-free medium. The stage-top chamber (Tokai-hit, Shizuoka, Japan) was filled with moisture using

distilled water and 5 % CO_2, and was maintained at 37°C. Images were captured every 1 min

between metaphase and early G1 phase. Signal intensity at chromosome rim was measured as

described below.

Microscopic images were exported as raw tiff 8-bit images (512x512 pixels) and analyzed

with particle analyzer plugin in FIJI macro (https://fiji.sc/). When analyzing self-assembled particles,

the sum of the RawIntDen (Raw Integrated Density) of each particle per image (total particle mass)

was measured. For obtaining the partition coefficient (PC) of a client protein, a threshold for the

particle size (>1 µm in Feret's diameter) was applied to eliminate non-self-assembled aggregations.

The ratio of the mean intensity of the client protein in the particle ($[I]_{particle}$) over that of the bulk

solution ($[\Pi]_{bulk}$) was calculated after the background subtraction and shown as common logarithm

values. For the sequential partitioning assay, the threshold for the particles' size was set between 5

and 15 µm in Feret's diameter to avoid both small and large particles.

For analyzing EGFP signal at the chromosome surface in mitotic HeLa cells, the

chromosome rim was detected in mPlum-H3 image and defined as a region of interest (ROI), which

is then transferred to the image of EGFP channel. EGFP signal intensity in the ROI was quantified,

subtracted by that of extrachromosomal space (background), and normalized by the value in

interphase. Obtained data were arranged in Excel (Microsoft) and R version 3.4.1.

FRAP analysis

The self-assembled particle was formed by ATTO488-labeled IDR-Nups as described above. A half

or a center of the particle was bleached by a 488 nm laser at the maximum output for 4 sec. After

bleaching, the time lapse observation was continued every 10 sec. Signal intensity of the bleached

region was quantified manually using the FIJI and the obtained values were processed as mentioned

above, and shown as a relative intensity to the signal intensity before bleaching.

CD spectra measurement

The CD spectra of purified IDR-Nups and a coiled-coil protein (Nup58CC) were measured using a

CD spectrophotometer (J-805, JASCO, Tokyo, Japan) with a 0.1 cm quartz cuvette. Data were

acquired every 0.1 nm between 200 and 260 nm. Samples were prepared in assay buffer (50 mM

HEPES pH 7.4, 100 mM NaCl, 2 mM 2-Me). The final protein concentration of Nup58FG-N,

Nup58CC, Tpr-IDR1, and Tpr-IDR2 were 0.2, 0.2, 0.3, and 0.3 mg/mL, respectively.

Nup35 knockdown

Small interference RNAs for Nup35 and control (luciferase) were purchased from Thermo (Silencer

Select siRNA #29400 for Nup35 and Stealth[TM] RNAi 12935-146 for Luciferase). It was introduced

into HeLa cells by Lipofectamine RNAi MAX (Invitrogen) by following the manufacturer's

protocol.

Statistical analysis

To analyze the statistics for all of obtained data in this study, Welch two sample *t* test was used for

normally distributed data, and Wilcoxon Rank-Sum test was used for non-normally distributed data.

A Shapiro-Wilk normality test was used to determine whether data were distributed normally. All of

statistical tests were done by using R version 3.4.1.

Author Contributions

H.A.K. planned the project, designed the experiments, performed all experiments, and wrote the

paper. S.H.Y. planned the project, designed the experiments, performed live cell imaging, and wrote

the paper.

Acknowledgements

We thank Prof. T. Haraguchi (human Nup98), Prof. K. Ullman (rat Nup62 and human Nup153), Prof.

J. Ellenberg (human Nup214) for providing cDNAs encoding nucleoporins. We also thank S. Dodo

for technical support, especially recombinant protein preparation and members of Yoshimura lab for

discussions. This study was supported financially by the Funding Program for Next Generation

World-leading Researchers (S.H.Y., No. LS076), and a Grant-in-Aid for Scientific Research (B)

(S.H.Y., No. 21370054) from the Japan Society for the Promotion of Science (JSPS).

Figure legends

Figure 1 Self-assembly of IDR-Nups

(A) Amino acid compositions of IDR fragments used in this study. Vertical lines indicate the

positions of positively charged (blue), negatively charged (red), and hydrophobic (gray) amino acids,

and FG motifs (black). For visibility, other amino acid residues (polar and non-polar) are not shown

here. Amino acid positions in the full-length protein are shown at both ends. The number of

FG-motifs, positively and negatively charged residues, as well as the ability of self-assembly are

shown on the right.

(B-I) DIC and fluorescence images of the self-assembled particles of Nup98FG (B), Nup153FG (C),

Nup214FG (D), Nup62FG (E), Tpr-IDR1 (F), Nup35N (G), ELYS-C1 (H), and ELYS-C2 (I) in the

presence or absence of 15 % PEG3350. Scale bar: 10 μm.

Figure 2 Self-assembly of IDR-Nups is driven by hydrophobic and/or electrostatic interactions

(A-E) Co-line plot of NaCl- and 1,6-hexanediol-dependency in self-assembly of IDR-Nups

(Nup98FG (A), Nup214FG (B), Tpr-IDR1 (C), Nup35N (D), ELYS-C1 (E)). Data are represented as

mean ± SEM from three independent experiments. Each value is shown as a relative value to the

standard condition (50 mM HEPES pH 7.4, 100 mM NaCl) in the presence or absence of 15 %

PEG3350. (*: $p<0.05$, **: $p<0.005$, ***: $p<0.0005$). P-values were obtained by appropriate statistical

tests (see Materials and Methods).

(F) A summary of self-assembling properties of IDR-Nups. Hydrophobic-based (gray) and

electrostatic-based (red) self-assembled IDR-Nups are highlighted.

Figure 3 Interaction diagram of IDR-Nups in a self-assembled particle

Interaction between FG-Nups (A and B), and between FG-Nup and non-FG-Nup (C and D) were

examined.

(A and C) Fluorescence images of the self-assembled particle. Images were taken 5-10 min after

mixing host and client proteins. Fluorescence signals of the host protein (labeled with ATTO390,

left), and the client protein (labeled with ATTO488, middle), as well as the merged image, (right) are

shown. Scale bar: 2 µm.

(B and D) Tile plot of partition coefficient (PC) of client proteins against host proteins. Data are

presented as an averaged value of the PC. An asterisk (*) indicates that client did not colocalize with

host in the same particle as shown in Figure S3E. All of PC values are shown in Table S2.

Figure 4 Nup35 facilitates the assembly of middle-assembling FG-Nups during the post-mitotic NPC

reassembly

(A) The order of Nup reassembly in the post-mitotic NPC reassembly. Early-, middle-, and late-stage

assembling Nups are plotted along the time after anaphase onset.

(B) Experimental schemes of the *in vitro* sequential addition assay. A fluorescently-labeled client

protein was added to the host particle (ELYS-C1) solution and the fluorescence signal was observed

under the microscope 300 sec after the addition (upper panel). Alternatively, a fluorescently-labeled

1st client was first added to the host particle solution (t = 0), and then the 2nd client, which was

labeled with different fluorescent dye, was added 300 sec after the addition of the 1st client (t = 300).

476 The fluorescence signal of the 2nd client was observed under the microscope 300 sec after the

477 addition of the 2nd client (t = 600) (lower panel).

478 (C) A summary of sequential addition assay as depicted in the upper panel of (B). DIC image of the

479 particle (300 sec after the addition) and the fluorescence images of the client (Nups98FG, 153FG,

480 58FG-N, and 62FG) (0 and 300 sec) are shown (upper panels). The fluorescence intensity of the

481 client in the host particle was quantified and plotted in the bottom panel. Data are represented as

482 mean ± SEM from three independent experiments. Multi-color images of both the host and the

483 clients are shown in Figure S4A. Scale bar: 10 μm.

484 (D) Fluorescence images of Nup35N, which had been incorporated into the host particle (ELYS-C1).

485 The fluorescence intensity was quantified as in (C), and plotted in the right panel.

486 (E and F) Fluorescence images of the 2nd client proteins in the sequential addition assay. As

487 described in B (bottom panel), the 1st client (Nup35N or Nup98FG) was first added to the host

488 particle (ELYS-C1) (t =0) (images are shown in Figure S4B), and then the 2nd client (Nup62FG (E),

489 Nup58FG-N (F)) was added (t = 300). Images of the 2nd client were taken 300 sec after the addition

490 (t =600), and are shown in (B, lower panel). The fluorescence intensity of the 2nd client protein

491 (Nup62FG or Nup58FG-N) was quantified and plotted as a relative value to the value without the 1st

492 client (none). Data are represented as mean ± SEM from three independent experiments. Scale bar:

493 10 μm. *: $p<0.05$, **: $p<0.005$, ***: $p<0.0005$, n.s.: no significance. P-values were obtained by

494 appropriate statistical tests (see Materials and Methods).

495 (G and H) The effect of Nup35-knockdown on the post-mitotic assembly of Nup153 and Nup62.

496 EGFP-fused Nup62 (G) or Nup153 (H) was expressed in Nup35-knocked-down HeLa cells,

497 together with mPlum-fused histone H3, and was subjected to the time-lapse imaging during mitosis.

498 Fluorescence signal at the chromosome rim was quantified as described in Materials and Methods,

and presented as relative values to the maximum value.　Data are shown as mean ± SEM. The

number of the analyzed cells is also indicated. *: $p < 0.05$ and n.s.: no significance. P-values were

obtained by appropriate statistical tests (see Materials and Methods). Scale bar: 10 μm.

(I) Schematic model of the post-mitotic NPC reassembly. See text for detail.

Table S1 Amino acid compositions and characteristics of IDR-Nups used in this study

Table S2 Partition coefficient of client protein against host protein in the assay shown in Figure 3.

Data are summarized and represented as mean ± SD from more than two independent experiments. n

indicates the number of analyzed particles. NA: not assigned.

Figure S1 Self-assembly of IDR-Nups

(A) Disorder probability of Nups used in this study. The results of PONDR-fit algorithm prediction

are shown against amino acid position.

(B-E and J-M) DIC and fluorescence images of self-assembled particles formed by IDRs from

FG-Nups (B-E) and non-FG-Nups (J-M) in different concentrations of PEG3350 (left). The total

particle mass (see Experimental Procedures) was measured and summarized in the right panel. Data

are presented as mean ± SEM, from three independent experiments. (*: $p < 0.05$, **: $p < 0.005$, ***:

$p < 0.0005$). P-values were obtained by appropriate statistical tests (see Experimental Procedures).

(F-I, N and O) DIC and fluorescence images obtained from non-self-assembling Nups in the

presence of 15% PEG3350. Scale bar: 10 μm.

Figure S2 Characterization of the self-assembling IDR-Nups

(A) Disorder probability of Tpr-IDRs. The result of PONDR-fit algorism prediction is shown.

(B) Co-line plot of the coiled-coil propensity (red) and the helix propensity (blue) of Tpr-IDRs.

(C) CD spectra of Tpr-IDR1 and Tpr-IDR2, Nup58FG-N and Nup58 (aa. 25-381). It was previously

reported that Nup58FG-N is disordered and Nup58CC forms a stable coiled-coil (30).

(D-I) FRAP analysis of self-assembled particles (Nup98FG (D), Nup153FG (E), Nup214FG (F),

Tpr-IDR1 (G), Nup35N (H), ELYS-C1 (I)). Fluorescence images before and after (0, 60, and 300

sec) bleaching (left panels), and signal intensity plot over time after the bleaching (right panel) are

shown. Two different bleaching patterns (half; blue, centre; red) were tested. Data are represented as

mean ± SD from three independent experiments.

Figure S3 Investigation of the driving force of self-assembly

(A and B) Fluorescence images of self-assembled particles in different concentrations of

1,6-hexanediol (A) and NaCl (B).

(C) Co-line plot of charge (red) and hydrophobicity (black) of ELYS-C1 against the amino acid

position.

(D) Amino acid sequence of Nup35N fragment. Proline and glycine residue are labeled in green and

red, respectively.

(E) Fluorescence images of non-colocalized particles shown as an asterisk (*) in Figure 3D. Scale

bar: 2 µm.

Figure S4 Sequential assembly of IDR-Nups into a self-assembled particle of ELYS-C1

(A) DIC and fluorescence images of the host (ELYS-C1, blue) and the client proteins (green).

Images were taken 3 min after the addition of the client. Scale bar: 20 µm.

(B) DIC and fluorescence images of the host (ELYS-C1, blue) and the 1st client (Nup35N or

Nup98FG, red). The incorporation of the 1st client into the host particle was confirmed before the

addition of the 2nd client protein. Scale bar: 20 µm.

(C) RNAi of Nup35. Small interference RNA for human Nup35 or luciferase was introduced into

HeLa cells. The cell lysate was subjected to immunoblot analysis using anti-Nup35 and anti-β-actin

antibodies. The position of Nup35 (~35 kDa) is indicated by an arrow.

Declaration of Interests

553 The authors declare no competing financial interests.

References

1. Maximiliano A. D'Angelo and Martin W. Hetzer. (2008) Structure, dynamics and function of

nuclear pore complexes. *Trends Cell Biol.* **18**, 456–466

2. Elad, N., Maimon, T., Frenkiel-Krispin, D., Lim, R. Y., and Medalia, O. (2009) Structural

analysis of the nuclear pore complex by integrated approaches. *Curr. Opin. Struct. Biol.* **19**,

226–232

3. von Appen, A., Kosinski, J., Sparks, L., Ori, A., DiGuilio, A. L., Vollmer, B., Mackmull,

M.-T., Banterle, N., Parca, L., Kastritis, P., Buczak, K., Mosalaganti, S., Hagen, W.,

Andres-Pons, A., Lemke, E. a., Bork, P., Antonin, W., Glavy, J. S., Bui, K. H., and Beck, M.

(2015) In situ structural analysis of the human nuclear pore complex. *Nature* **526**, 140–143

4. Fischer, J., Teimer, R., Amlacher, S., Kunze, R., and Hurt, E. (2015) Linker Nups connect the

nuclear pore complex inner ring with the outer ring and transport channel. *Nat. Struct. Mol.*

Biol. **22**, 774–781

568 5. Ullman, K. S. and Powers, M. A. (2015) Locking down the core of the pore. *Science (80-.).*

569   **350**, 33–34

570 6. Konishi, H. A., Asai, S., Watanabe, T. M., and Yoshimura, S. H. (2017) In vivo analysis of

571   protein crowding within the nuclear pore complex in interphase and mitosis. *Sci. Rep.* **7**, 5709

572 7. Rabut, G., Doye, V., and Ellenberg, J. (2004) Mapping the dynamic organization of the

573   nuclear pore complex inside single living cells. *Nat. Cell Biol.* **6**, 1114–1121

574 8. Denning, D. P., Patel, S. S., Uversky, V., Fink, A. L., and Rexach, M. (2003) Disorder in the

575   nuclear pore complex: the FG repeat regions of nucleoporins are natively unfolded. *Proc. Natl.*

576   *Acad. Sci. U. S. A.* **100**, 2450–2455

577 9. Mohr, D., Frey, S., Fischer, T., Güttler, T., and Görlich, D. (2009) Characterisation of the

578   passive permeability barrier of nuclear pore complexes. *EMBO J.* **28**, 2541–2553

579 10. Frey, S. and Görlich, D. (2007) A saturated FG-repeat hydrogel can reproduce the

580   permeability properties of nuclear pore complexes. *Cell* **130**, 512–523

581 11. Patel, S. S., Belmont, B. J., Sante, J. M., and Rexach, M. F. (2007) Natively unfolded

582   nucleoporins gate protein diffusion across the nuclear pore complex. *Cell* **129**, 83–96

583 12. Frey, S., Richter, R. P., and Görlich, D. (2006) FG-rich repeats of nuclear pore proteins form a

584   three-dimensional meshwork with hydrogel-like properties. *Science* **314**, 815–817

585 13. Schmidt, H. B. and Görlich, D. (2015) Nup98 FG domains from diverse species

586   spontaneously phase-separate into particles with nuclear pore-like permselectivity. *Elife* **4**,

587   1–30

588 14. Dultz, E., Zanin, E., Wurzenberger, C., Braun, M., Rabut, G., Sironi, L., and Ellenberg, J.

589   (2008) Systematic kinetic analysis of mitotic dis- and reassembly of the nuclear pore in living

590   cells. *J. Cell Biol.* **180**, 857–865

591   15.   Otsuka, S., Szymborska, A., and Ellenberg, J. (2014) Imaging the assembly, structure, and

592   function of the nuclear pore inside cells. *Methods Cell Biol.* **122**, 219–238

593   16.   Woodruff, J. B., Wueseke, O., Viscardi, V., Mahamid, J., Ochoa, S. D., Bunkenborg, J.,

594   Widlund, P. O., Pozniakovsky, A., Zanin, E., Bahmanyar, S., Zinke, A., Hong, S. H., Decker,

595   M., Baumeister, W., Andersen, J. S., Oegema, K., and Hyman, A. a. (2015) Centrosomes.

596   Regulated assembly of a supramolecular centrosome scaffold in vitro. *Science* **348**, 808–812

597   17.   Iwashita, K., Handa, A., and Shiraki, K. (2018) Coacervates and coaggregates: Liquid–liquid

598   and liquid–solid phase transitions by native and unfolded protein complexes. *Int. J. Biol.*

599   *Macromol.* **120**, 10–18

600   18.   Gillespie, P. J., Khoudoli, G. a, Stewart, G., Swedlow, J. R., and Blow, J. J. (2007)

601   ELYS/MEL-28 chromatin association coordinates nuclear pore complex assembly and

602   replication licensing. *Curr. Biol.* **17**, 1657–1662

603   19.   Otsuka, S. and Ellenberg, J. (2017) Mechanisms of nuclear pore complex assembly - two

604   different ways of building one molecular machine. *FEBS Lett.* **592**, 475–488

605   20.   Kim, S. J., Fernandez-Martinez, J., Nudelman, I., Shi, Y., Zhang, W., Raveh, B., Herricks, T.,

606   Slaughter, B. D., Hogan, J. A., Upla, P., Chemmama, I. E., Pellarin, R., Echeverria, I.,

607   Shivaraju, M., Chaudhury, A. S., Wang, J., Williams, R., Unruh, J. R., Greenberg, C. H.,

608   Jacobs, E. Y., Yu, Z., de la Cruz, M. J., Mironska, R., Stokes, D. L., Aitchison, J. D., Jarrold,

609   M. F., Gerton, J. L., Ludtke, S. J., Akey, C. W., Chait, B. T., Sali, A., and Rout, M. P. (2018)

610   Integrative structure and functional anatomy of a nuclear pore complex. *Nature* **555**, 475–482

611   21.   Chug, H., Trakhanov, S., Hülsmann, B. B., Pleiner, T., and Görlich, D. (2015) Crystal

612   structure of the metazoan Nup62•Nup58•Nup54 nucleoporin complex. *Science* **350**, 106–110

22.   Li, P., Banjade, S., Cheng, H.-C., Kim, S., Chen, B., Guo, L., Llaguno, M., Hollingsworth, J.

V, King, D. S., Banani, S. F., Russo, P. S., Jiang, Q.-X., Nixon, B. T., and Rosen, M. K.

(2012) Phase transitions in the assembly of multivalent signalling proteins. *Nature* **483**,

336–340

23.   Piela, L., Némethy, G., and Scheraga, H. A. (1987) Proline-induced constraints in α-helices.

Biopolymers **26**, 1587–1600

24.   O'Neil, K. T. and DeGrado, W. F. (1990) A thermodynamic scale for the helix-forming

tendencies of the commonly occurring amino acids. *Science* **250**, 646–651

25.   Rauscher, S., Baud, S., Miao, M., Keeley, F. W., and Pomès, R. (2006) Proline and Glycine

Control Protein Self-Organization into Elastomeric or Amyloid Fibrils. *Structure* **14**,

1667–1676

26.   Bilokapic, S. and Schwartz, T. U. (2012) Molecular basis for Nup37 and ELY5/ELYS

recruitment to the nuclear pore complex. *Proc. Natl. Acad. Sci.* **109**, 15241–15246

27.   Glavy, J. S., Krutchinsky, A. N., Cristea, I. M., Berke, I. C., Boehmer, T., Blobel, G., and

Chait, B. T. (2007) Cell-cycle-dependent phosphorylation of the nuclear pore Nup107-160

subcomplex. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3811–3816

28.   Linder, M. I., Köhler, M., Boersema, P., Weberruss, M., Wandke, C., Marino, J., Ashiono, C.,

Picotti, P., Antonin, W., and Kutay, U. (2017) Mitotic Disassembly of Nuclear Pore

Complexes Involves CDK1- and PLK1-Mediated Phosphorylation of Key Interconnecting

Nucleoporins. *Dev. Cell* **43**, 141–156

29.   Laurell, E., Beck, K., Krupina, K., Theerthagiri, G., Bodenmiller, B., Horvath, P., Aebersold,

R., Antonin, W., and Kutay, U. (2011) Phosphorylation of Nup98 by multiple kinases is

crucial for NPC disassembly during mitotic entry. *Cell* **144**, 539–550

30.   Solmaz, S., Chauhan, R., Blobel, G., and Melčák, I. (2011) Molecular architecture of the

transport channel of the nuclear pore complex. *Cell* **147**, 592–602

Konishi HA et al. Figure 1

Konishi HA et al. Figure 2