RMI-chip: a microfluidics setup for functional imaging of microbial interactions with tree roots

Marie-Francoise S. Gros[a], Shalaka V. Shinde[a,†], Chase Akins[a], Jessica L. Johnson[a], Sarah Zerbs[a], Rosemarie

Wilton[a], Ken Kemner[a], Philippe Noirot[a], Gyorgy Babnigg[a]*

Abstract

Coupling microfludics with microscopy has emerged as a powerful approach to study dynamics in plant

physiology at cellular resolution and at higher throughput than with conventional methods. Most devices

have been designed to study the model plant *Arabidopsis thaliana*, including root-microbe interactions.

However, there is a need for microfluidic devices which enable *in vivo* studies of root development and

root-microbe interactions in woody plants. Here, we developed the RMI-chip, a microfluidic setup that

allows for continuous microscopic scale observation of real-time interactions between live *Populus*

tremuloides (aspen tree) seedling roots and the rhizobacterium *Pseudomonas fluorescens* over an

extended period longer than a month. Colonization dynamics in the RMI-chip are consistent with previous

observations in a different experimental set-up. Also, we find that biosensors based on the rhizobacterium

Bacillus subtilis can be used to monitor compositional changes in root exudates but that their application

is limited by their inefficient colonization of aspen roots. Our results indicate that functional imaging of

dynamic root-bacteria interactions in the RMI-chip requires careful matching between the host plant and

the bacterial root colonizer.

Introduction

The plant microbiome plays an important role in the rhizosphere [1-3]. Virtually all plant tissues host

microbes that can act as symbionts, commensals, or pathogens. Interactions between plant and microbes

can be beneficial, neutral, or harmful and directly influence plant growth and productivity [2, 4, 5]. Plant-

growth-promoting (PGP) rhizobacteria are bacteria that exert beneficial effects on plants through direct

or indirect interactions with the roots [6, 7]. PGP bacteria have the potential to increase the availability of

soil nutrients to the plant, produce metabolites such as plant hormones, elicit plant systemic defences,

and increase plant resistance to biotic and abiotic stresses [8, 9]. In return, the plant provides

photosynthetically-derived carbon, such as sugars and organic acids that are consumed by rhizosphere

microorganisms as well as a wide range of molecular compounds acting as environmental signals for the

root microbiota. Microbes attach to the root surface and form microcolonies that can eventually grow

into larger biofilms. The formation of biofilms at root surfaces was proposed to be part of the cellular PGP

activities of beneficial rhizobacteria [10].

Understanding these complex interactions between plant roots and microbes requires the ability to track

their dynamics at high spatial and temporal resolution. Real time monitoring of dynamic root-microbe

interactions at cellular resolution is now possible using microfluidics approaches coupled with advanced

live imaging microscopy. Microfluidic platforms provide a powerful approach to evaluate the responses

of growing plant cells to external perturbations (e.g., nutrients, media flow, temperature, hydrodynamics,

light, and stressors) at throughputs higher than with conventional methods using pots or plates, and in

precisely controlled environments. Multiple microfluidics devices such as "Plant on a chip" [11], RootChip

[12], RootArray [13], TipChip [14], and PlantChip [15] were developed to study various aspects of the cell

biology of *Arabidopsis thaliana*, including gene expression, cell biomechanics, cellular mechanism of

growth and cell division (reviewed in [16]). The PlantChip device enables the continuous monitoring of

phenotypic changes at the cellular level and also at the whole plant level, including seed germination and

root and shoot growth (hypocotyls, cotyledons, and leaves) [15].

Fewer studies have used microfluidic devices to visualize the interactions of *Arabidopsis* roots with

pathogenic or beneficial microorganisms. Using the plant-in-chip platform, visualization of the attack of

Arabidopsis roots by pathogenic nematodes and oomycetes motile spores revealed some physiological

changes taking place in the host plant and the pathogen during the attack [17]. Recently, a microfluidic

device to track root interactions system (TRIS) revealed a distinct chemotactic behaviour of the bacterium

Bacillus subtilis toward the root elongation zone and its rapid colonization, and allowed real-time

monitoring of bacterial preference between roots from various *Arabidopsis* genotypes [18]. To date,

studies of plant roots and root-microbe interactions (RMIs) using microfluidics have been focused on *A.*

thaliana, an annual herbaceous plant model that can complete is entire life cycle in 6 weeks, and grows a

single primary root that later produces smaller lateral roots. However, there is a need to study root

development and root-microbe interactions for other plants, including woody perennial plants such as

trees.

Here, we describe a microfluidic device, called RMI-chip, that enables the direct visualization of root-

microbe interactions taking place at early stages of tree seedling growth. We studied the interactions of

Populus tremuloides (trembling aspen tree) with the bacterium *Pseudomonas fluorescens.* These

interactions are biologically relevant in nature, as *P. fluorescens* is abundant in the rhizosphere of *Populus*

trees [1, 19-21], and exhibits functionality in laboratory assays. We have shown that *P. fluorescens*

promotes the growth of aspen seedlings [22], colonizes aspen seedling roots by forming dense and

dynamic biofilms [23], and modulates expression of anti-fungal defence response genes in roots of aspen

seedlings [24]. The RMI-chip device was designed to accommodate aspen seedling growth for periods up

to 5 weeks, and to enable direct observation of root growth and its dynamic colonisation by *P. fluorescens*

biofilms with high spatiotemporal resolution. The RMI-chip was also used to monitor the growth of rice

seedling roots and detect the production of reactive oxygen species (ROS) by the root using engineered

Bacillus subtilis strains as biosensors. We find that in the RMI-chip interactions between host plants and

bacterial species are specific, consistent with ecological observations and with colonization profiles

observed in other experimental systems, and that formation of bacterial biofilms on root surfaces is

needed for extended colonization.

Experimental

Materials and chemicals

Seeds of Populus tremuloides Michx. were obtained from the National Tree Seed Center, Natural

Resources Canada, Fredericton NB, Canada. The rice seeds were obtained from Baker Creek Heirloom

Seed Co. (Mansfield, MO, USA). Phytoblend was purchased from Caisson Laboratories, Inc. (Smithfield,

UT, USA). All other chemicals were purchased from Sigma-Aldrich (St. Louis, MO, USA).

RMI-chip and humidity chamber design

The chips were designed with AutoCAD software (Autodesk) and were fabricated via soft lithography at

the Scientific Device Laboratory (Des Plaines, IL, USA). Briefly, SU-8 2025 photoresist (MicroChem,

Westborough, MA, USA) was used to make moulds on a 4-inch silicon wafer. The two-part silicone

elastomer (SYLGARD 184, Thermo Fisher Scientific, Waltham, MA), the silane precursor and curing agent

were mixed 10:1, degassed, poured on silicon wafer mould, degassed and baked at 65 °C for 4 hours. The

PDMS elastomer was punched and bonded to a 48 x 65mm, thickness No. 1 cover glass (Ted Pella Inc.,

Redding, CA, USA). RMI-chip devices were used without Aquapel treatment. The humidity chambers and

microscope stage adapters were fabricated from PMMA thermoplastic via laser cutting (Ponoko, Oakland,

CA, USA) and assembled with acrylic adhesive (Weld-On, Compton, CA, USA). Tubing management and

other holder accessories were 3D printed using PLA. The AutoCAD files (ESI) of the RMI-chip design, the

SVG files used for laser cutting, and the SCAD and STL files used for 3D printing are available in the online

supporting material.

Fluorescent-labelled and biosensor bacterial strains

The mNeonGreen (mNG)- and dsRed-labelled *P. fluorescens* SBW25 strains harboured an environmentally

stable plasmid derivative that constitutively expresses the dsRed or mNG fluorescent proteins, as

described [25]. *B. subtilis* strains labelled with the mCherry fluorescent protein were constructed by

transferring the genetic construct P_{pen}-mCherry::*kan* from *B. subtilis* MMB1023 [26] into the NCIB3610

background by phage transduction. The resulting NCIB3610-mcherry strain, which expressed

constitutively mCherry, was used as a recipient for the transfer of sensory genetic elements. A xylose-

responsive genetic module was constructed by inserting a P_{xylA} BioBrick DNA block (PCR-amplified from

plasmid pBS1C3-PxylA) into plasmid pBS1C [27] to form pBS1C-PxylA, which was then used as recipient

for the insertion of the gfp_{sp} gene from pRD111 [28], using Spe1 and Pst1 restriction sites. In the final

plasmid pBS1C-P_{xylA}-gfp$_{sp}$, the gfp_{sp} gene is under control of the xylose-inducible P_{xylA} promoter, linked with

a chloramphenicol resistance gene (*cat*) and flanked by the right and left arms of the *B. subtilis amyE* gene.

The P_{xylA}-GFP::*cat* module was integrated in the *amyE* gene of *B. subtilis* strain 1A976 (SKC6) by

transformation of the competent cells. The module was then finally transferred in the NCIB3610-mCherry

strain by SPP1-mediated phage transduction. A ROS-responsive *B. subtilis* strain was constructed by SPP1

transduction of the genetic module P_{katA}-gfp$_{sp}$::*spc* from strain 1A1010 (Bacillus Genetic Stock Center,

http://www.bgsc.org), where P_{katA} is the promoter controlling the expression the catalase gene, in the

NCIB3610 strain for integration at *amyE locus*. Transductants were selected for resistance to

spectinomycin.

Aspen and rice cultivation

Populus tremuloides Michx. seeds were cold stratified in Milli-Q water at 4°C for 2 to 14 days. Seeds were

then surface sterilized by prewashing briefly in 1% Tween-20 and incubated in pH-reduced (10 mM HCl)

0.1M sodium hypochlorite. After 4 minutes of incubation the seeds were rinsed 8-10 times with sterile

Milli-Q water and incubated in sterile Milli-Q water overnight in the dark at room temperature. The

following day the seeds were spread onto 1% agar Johnson's plate (4 mM KNO_3, 2 mM $Ca(NO_3)_2$, 4 mM

NH_4NO_3, 0.5 mM KH_2PO_4, 1 mM $MgSO_4$, 440 µM KCl, 250 µM H_3BO_3, 20 µM $MnSO_4$, 20 µM $ZnSO_4$, 5

µM $CuSO_4$, 5 µM $NaMoO_4$, 5 µM $CoCl_2$, 200 µM Fe,Na-EDTA, pH 5.6, 1% Phytoblend). The seeds were

incubated under low to medium light on a 16hr light/8hr dark period, maintaining the temperature below

30°C. Once germinated, seedlings with hypocotyls shorter than 5 mm were transferred into cut 200 µL

pipette tips filled with 1% Johnson's agar to let the root elongate gravitropically. For a typical experiment

20-25 pipette tips were inserted into Johnson's agar (1%) plates at a ~45° angle and incubated under the

same conditions (Fig. S1). Root growth was checked periodically. Six seedling-containing tips with roots

grown close to the end of the tips were mounted into the RMI-chip, which was incubated at a ~45° angle

to facilitate root infiltration.

De-hulled rice seeds (*Oryza sativa* 'Diamante-INIA') were dehulled and prewashed briefly in 1% Tween-

20, followed by incubation in pH-reduced (40 mM HCl) 0.4M sodium hypochlorite solution on a rocker for

15-20 minutes. The seeds were rinsed 7 times with Milli-Q water, plated on 1% Johnson's agar, and

incubated at 30°C in the dark for 48-72 hours. When germinated and the radical protrusions reached 3-

6mm, the seedlings were planted directly into cut 200 µL pipette tips filled with Johnson's solution already

inserted into the RMI-chip.

Following hypochlorite sterilization, all seed and seedling manipulations were performed in a laminar flow

hood, including mounting in the RMI-chip. Syringes, tubing, and adaptors were autoclaved prior to

connection to the chip to maintain sterility during the extended continuous perfusion.

Bacterial inoculations of the RMI-chips

Bacterial species were cultivated overnight in LB at their respective optimal temperature (i.e. 28°C for *P.*

fluorescens, 37°C for *B. subtilis*) in the presence of the appropriate antibiotic selection. Overnight cultures

of *P. fluorescens* SBW25-mNG and -DsRed strains were centrifuged, rinsed in PBS, and re-suspended in

PBS to OD_{600}=4. A 10µl aliquot of cell suspension was used for inoculation of aspen roots. Overnight

cultures of *B. subtilis* biosensor strains were diluted 100-fold and grown in LB up to mid-exponential phase

(OD_{600}=0.4), rinsed with PBS and likewise adjusted to OD_{600}=4 for the inoculation of the roots of aspen

and rice seedlings. After overnight incubation without flow, excess bacteria were washed away with

Johnson's solution at a flow of 0.2 µL/min for 2 hours, followed by continuous perfusion at 0.02 µl/min.

The outlet ports were connected to 200 µL pipette tips during inoculation, excess liquid was removed,

and ports were connected to independent lines to prevent cross-communication between the 100 µm x

800 µm channels (Fig. 1a) and with the external Johnson's solution overlay during perfusion. Note that in

theory, the RMI-chip can be washed and autoclaved after first use making the second set of 6 independent

channels are available for a new experiment. In this study, RMI-chips were used only once.

Imaging experiments

An inverted Nikon C2+ laser-scanning confocal microscope was used for imaging experiments (Nikon,

Melville, NY, USA). An Eclipse Ti-E inverted microscope equipped with perfect focus system, an automated

stage, and with 10x, 20x, and 100x objective lenses (CFI Plan Fluor 10X, NA 0.3, WD 16 mm; CFI Plan

Apochromat Lambda 20x, NA 0.75, WD 1.00 mm; CFI Plan Apo Lambda 100x, NA 1.5, WD 130 um,

respectively) was used for single image, time series, and z-stack acquisitions. Laser illumination emission

at 488 nm coupled with a 525/50-nm excitation filter was used to capture mNeonGreen fluorescence, and

laser illumination at 561 nm coupled with a 595/50-nm excitation filter was used to capture dsRed (*P.*

fluorescens) or mCherry (*B. subtilis*) fluorescence. The transmitted light was also detected to image

bacterial colonization in the context of the root structure. A custom holder was designed to provide access

of the objective to the RMI-chip and imaging chamber.

Results and discussion

RMI-chip design optimized for aspen seedling growth

The RMI-chip was designed to observe the growth and rhizobacterial colonization of aspen seedling roots

over extended periods of time. The aspen root system consists of a taproot from which smaller branch

roots emerge. When a seed germinates, the first root to emerge is the primary root which develops into

the taproot. With seedlings grown into agar-filled pipette tips (Fig. S1a), we observed that the primary

root quickly branched after exiting the tip (Fig. S1b). Thus, seedlings were inserted into the RMI-chip

shortly before the primary root reached the end of the pipette tip. Further development of the primary

root, including formation of branch roots, was not hampered by the pipette tip (Fig. S2a). In early studies,

we observed that the primary root readily entered the RMI-chip circular chamber (Fig. 1a), but we found

that a substantial part of the root system di not continue into the linear growth channel, instead growing

in circles, which caused the seedling to wither quickly. This problem was resolved by growing the seedlings

semi-gravitropically in a chip tilted at a 45° angle, submerged in Johnson's solution for up to a week, until

the root tip reached the media inlets in the growth channel (Fig. S2c). Under these conditions, most

primary roots entered the growth channel, at which point the RMI-chip was placed horizontally in a

humidity chamber and continuous media flow was supplied (Fig. 2). Regular exchange of the nutrient

solution surrounding the RMI-chip and the maintenance of water-saturated paper used to provide the

required humidity in the growth chamber, were crucial to obtain healthy seedling growth.

The RMI-chip has 6 independent channels with one inlet for the seedling root, two dedicated inlets with

filters for media and bacterial inoculation, and one common outlet (Fig. 1 a). The features are generated

via soft lithography on a silicon wafer, which is then used as a mould to produce polydimethylsiloxane

(PDMS) slabs. The PDMS is bonded to a large thin microscope coverslip providing a means for imaging at

cellular resolution using a confocal microscope. Different channel depths were tested for aspen seedling

growth. Although aspen primary root can grow into an 80 μm x 80 μm channel in a submerged microfluidic

device (Fig. S2b), the root occupied the entire channel precluding flow experiments. Therefore, we

selected a channel width of 800 μm and fabricated channels with depths of 100 μm and 400 μm. While

aspen seedlings grew similarly in both devices, the 400 μm deep channel did not position the root close

to the glass coverslip, making it suboptimal for microscopic observations using objectives with standard

working distances (Fig. 2). Our final design has growth channels 100 μm deep, 800 μm wide and 36 mm

long, which can easily accommodate aspen primary roots and enable root growth under constant

perfusion. Flow characteristics of the 400 μm x 800 channel were measured with fluorescently labelled

styrene beads (Fig. 1b). Without perfusion, slight bulk material movement was detected. At 1 μL/min,

laminar flow was observed with a measured 50 μm/s bead velocity (Supplementary video 1).

It can be challenging for plant roots to grow under constant flow. For example, without flow in a growth

channel, root border cells and mucilage can be observed at the root tip (Fig. S2c, Fig. S3). These cells and

mucilage can be washed away at high flow rates (e.g., 2 μL/min). Nevertheless, nutrient flow is needed to

maintain seedling growth in the RMI-chip. Therefore, we determined a minimal flow rate of nutrient

solution that did not affect root growth, preserving root morphology, including root cap, root hairs and

mucilage, while keeping out air bubbles. After several iterations, each including at least 4 seedlings, the

minimal flow rate was found to be 0.02 μL/min. As each channel is connected to an individual 1 ml syringe,

this flow rate provides enough media for up to 5 weeks. Under these conditions, the calculated average

laminar flow is 4 µm/s and the media is replaced every ~14 minutes in each channel. Aspen seedlings

exhibited a high heterogeneity in primary root growth with an average growth rate of 1 ± 0.9 mm/d. This

heterogeneity may be due to the use of open pollinated seeds which have different genetic makeups. In

addition, only 4 or 5 seedlings out of 6 mounted in the RMI-chip typically sustained growth over a 5-week

period. On average, aspen primary roots grew much slower than Arabidopsis primary roots reported to

grow at 3.7 mm/d in the RootChip [12].

Continuous imaging of the RMI-chip

Confocal Laser Scanning Microscopy was used to perform spatial and temporal studies of the aspen root

growth. Because of the repeated microscopy observations, a flexible system was required which was

compatible with the transfer of the chip between the growth chamber and microscope stage. A modular

chamber was designed to overcome this technical challenge and to prevent water evaporation during

cultivation and imaging. An innermost imaging chamber was built to provide humidity during the repeated

1-hour imaging experiments, and a medium-size chamber allowed submersion of the RMI-chip in media

and its transport between incubation chamber and microscope. A six-channel syringe pump was

connected with Teflon™ FEP tubing to the chip and transferred with the chip during imaging experiments

in order to provide continuous flow of nutrients, even during long imaging experiments (Fig. S4). Finally,

a large closed chamber that can be filled with water was built to provide humidity during long-term

growth. In addition, a holder compatible with the imaging chamber was fabricated for the Nikon Ti-2

microscope (Fig. S4). The humidity chambers and microscope stage adapters were fabricated from PMMA

thermoplastic. 3D printed inserts were fabricated to manage the tubing during imaging experiments

(design files are available in the online supporting material). With this system, the RMI-chip was

continuously perfused with Johnson's solution flowing at 0.02 μL/min, including during the lengthy

observations under the microscope, thus minimizing the disturbance of seedling growth.

Growing rice seedlings in the RMI-chip

We tested the compatibility of the RMI-chip with other plant species such as rice. Rice seedling roots could

be introduced easily into the chamber without an agar tip by directly transplanting the seedlings within

days after germination. Healthy root growth was observed under continuous perfusion with Johnson's

solution at a 0.02 μL/min flow rate (Fig. S5). Under these conditions, we observed fast root growth where

the root tip reached the end of growth chamber after 4 days on average. As for aspen, variability between

seedlings was observed, some root tips reaching RMI-chip end after 3.5 to 5 days. Thus, rice root growth

rate is approximately 10 times faster than aspen root growth under the same conditions. Unfortunately,

the fast growth rate of rice seedlings limits observation time to 3-5 days. Once the root tip reaches the

outlet, it blocks the chamber and cuts off nutrient flow.

Colonization of aspen seedling root by *P. fluorescens*

Aspen seedlings grown for 8 days in the RMI-chip were inoculated with mNeonGreen-labelled *P.*

fluorescens SBW25 by injection of approximately 2×10^9 cells through an inlet port (Fig. S6a). After

incubation for 16 hours without flow, the flow was restored at 0.2 μL/min for 2 hours to wash away excess

bacterial cells, then set at 0.02 μl/min for continuous perfusion. Under these conditions, a small number

of SBW25 cells remained associated with the lower part of the root (Fig. S6b). Importantly, the only source

of carbon available for bacterial growth in the RMI-chip are the sugars and organic acids

photosynthetically produced by the plant. Five days after inoculation, actively dividing SBW25 cells were

colonizing the intercellular spaces between root epidermal cells of the root cortex, and, to a lesser extent,

the root surface (Fig. S7). Although this preferential colonization of crevices on the root surface may be

caused by the flow in the RMI-chip, it is consistent with our previous observations in a vertical plate

system, where SBW25 first colonizes the cell interstitial spaces along the aspen root cortex before forming

a variety of bacterial assemblies that range from microcolonies to highly structured biofilms [23].

Next, we observed aspen primary root colonization by a 1:1 mixture of two *P. fluorescens* SBW25

derivatives labelled with distinct fluorescent proteins, mNeonGreen and dsRed. Three days after

inoculation, we observed discrete patches of red and green cells with almost no area of mixed colours

(Fig. 3a). This finding indicates that bacterial cells divide and form expanding patches in the intercellular

spaces and on the root surface, ruling out that those patches are formed by random deposition and

aggregation of cells within the root crevices. Thirteen days after inoculation, bacterial patches were longer

and formed dense cell assemblies on the primary root surface (Fig. 3b). Close-up examination revealed

mature biofilms resulting from clonal growth of green and red bacterial populations. These biofilms

exhibited channel-like void spaces, which are a hallmark of mature biofilms [29]. Although cell density and

presence of small aggregates in the pure cultures prior to co-inoculation could play a role in determining

the colonisation patterns [30], the quasi-absence of intermixed biofilms is indicative of a competition

between the green and red strains. These strains have an identical genome but differ by the expressed

fluorescent protein, possibly creating differences in cell fitness and some competition.

Notably, we observed that a variety of SBW25 cell assemblies coexisted with biofilms on the root surface

(Fig. 4a). These assemblies included clusters of somewhat regularly spaced individual cells, clusters of cell

doublets which are presumably dividing, microcolonies and mature biofilms (Fig. S8). These assemblies

may reflect the different stages of biofilm formation, where cells attach to the root surface, divide, and

over time form more compact assemblies maturing into biofilms. Intriguingly, cells packed within a biofilm

appeared to have a round coccoid shape in contrast with the normal rod-like morphology of individual

SBW25 cells (Fig. 3a, Fig. 4a). Confocal 3D image reconstruction and side projection of the 3D volume (Fig.

4b) revealed that the coccoid shape was only apparent and resulted from the top view of vertically

arranged bacterial cells within the biofilm. This finding corroborates our previous observation of vertical

arrangements of SBW25 cells within biofilms and mucilage formed on the roots of aspen seedlings

growing in a vertical agar plate [23].

After 30 days in the RMI-chip, fluorescent SBW25 cells became rare on the root surface and were

preferentially found as micro-colonies located within the spaces between root epidermal cells (Fig. S9).

Overall, the dynamic of root colonization by SBW25, which includes active growth and colonization of the

root during the first two weeks, formation of mature biofilms associated with a variety of other cell

assemblies, and dispersal of these biofilms and assemblies after 5 weeks, appears to be remarkably

comparable in the RMI-chip and vertical plate assay, the latter being a very different set-up where seedling

roots grow on a nutrient agar surface [23]. This similarity suggests that the observed dynamics of cell

assemblies are not primarily shaped by the flow in the RMI--chip but rather reflect an intrinsic colonization

behaviour of aspen primary root by *P. fluorescens* SBW25. In addition, novel cell assemblies such as

clusters of somewhat regularly spaced individual cells and cell doublets were observed uniquely in the

RMI-chip.

RMI-chip and rhizobacterial biosensors to monitor dynamic root exudation

In the RMI-chip, the carbon sources available for bacterial growth are provided by root exudates. These

exudates are compositionally complex and contain ions (i.e. H+), inorganic acids, oxygen, water, and

multiple carbon-based compounds which include amino acids, organic acids, sugars, phenolics and an

array of secondary metabolites, and high-molecular weight compounds like mucilage and proteins [31].

Root exudates are key mediators of interactions with microbes in the rhizosphere and root exudation is a

regulated process which responds to biotic and abiotic stresses to the plant [31]. We assessed whether

biosensors, which are engineered rhizobacteria that can colonize root surfaces and express a fluorescent

protein in response to the presence of a specific metabolite or environmental stressor, could be used in

the RMI-chip to monitor dynamic changes in the composition of plant exudates.

We used *Bacillus subtilis,* which colonizes *Arabidospis* roots [18] and is used as biocontrol agent to colonize

and protect various herbaceous plants [32], to develop our biosensor strains. We selected two well

characterized promoters, one responding to the presence of xylose [33], a sugar abundant in root

exudates, and the other one responding to the presence of Reactive Oxygen Species (ROS) [34], which are

normal products of plant cellular metabolism and act as second messengers in plant responses to various

environmental stresses [35]. The biosensors expressed the green fluorescent protein (GFP) conditionally

in the presence of xylose or ROS and also expressed constitutively the red fluorescent protein mCherry,

which serves to label the cells (see Experimental methods).

Aspen primary roots were inoculated with *B. subtilis* biosensors in the RMI-chip, incubated overnight

without flow, and upon restoration of flow the root was observed using confocal laser microscopy. We

found that, as with *P. fluorescens*, most cells were rapidly washed away from the root surface. However,

biosensor cells were very sparse on the root after 3 days and could not be observed after 5 days. The use

of a low flow (0.02 μL/min) throughout inoculation of the RMI-chip did not enhance the number of

biosensor cells remaining attached to the primary root, reflecting an inability of *B. subtilis* to colonize

aspen primary root under these conditions. This is in contrast with our observations with *P. fluorescens*

and may reflect the fact that *B. subtilis* is not a natural colonizer of aspen roots. To circumvent this

problem, observations were done in a stopped-flow experiment limited to 5 days, after which the aspen

seedling starts to whither. Even In the absence of flow, *Bacillus* cells did not colonize but remained loosely

attached to the aspen root surface. Whereas biosensor cells did not express GFP after inoculation (Fig. 5

a), strong GFP signals were detected in biosensor cells remaining attached to the aspen root after 5 days

of incubation, indicating the production of xylose (Fig. 5b) or ROS (Fig. 5c) by the root. Exposure of the

ROS biosensor to a rice primary root under the same RMI-chip conditions (i.e., no flow) resulted in a robust

colonization of the root and in high levels of GFP after 3 days (Fig. 5d). This finding indicates that high

levels of ROS are produced by rice primary root. Together, these results provide a proof-of-concept that

bacterial biosensors can be used in RMI-chip as a way to investigate the dynamic chemical crosstalk

between root and rhizobacteria.

We also inoculated rice seedlings with *P. fluorescens* SBW25 to test for ability to colonize in the RMI-chip.

Repeated trials showed that SBW25 cells were rapidly washed away from the root surface and remained

barely detectable under nutrient flow after one day (data not shown), suggesting an inability of *P.*

fluorescens SBW25 to colonize rice primary root under these conditions. This result was unexpected as

rice was shown to host *Pseudomonas* endophytes [36]. These observations clearly show that biosensors

need to be developed from bacterial isolates that efficiently colonize the roots of the plant host of

interest. What may seem to be a caveat for the development of biosensors is actually grounded in ecology

and evolution, as root exudates from a particular plant species are known to maintain and support a highly

specific diversity of microbes in the rhizosphere of this plant [31].

Conclusions

In summary, a microfluidic device and accessories were fabricated enabling the long-term cultivation of

aspen seedlings under constant flow and the study of root-microbe interactions through repeated

functional imaging of the system. The device enabled high-resolution imaging through the various phases

of aspen colonization by *P. fluorescens* SBW25 at the single bacterial cell level under continuous nutrient

flow. Different biofilm morphologies and diverse bacterial assemblies were observed over time,

emphasizing the need for high-resolution imaging to understand colonization patterns and strategies over

time driven by local environments. *B. subtilis* biosensors provided a means for functional imaging of root

exudate uptake by the bacteria and the detection of an environmental stressor. However, the use of these

biosensors was limited by their lack of long term colonization of the host plant. The RMI-chip can be used

for the long-term observation of other slow-growing plants, or modified to study faster growing plants as

well.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was performed under the Small Worlds project and was supported by the Office of Biological

and Environmental Research (BER), Office of Science, United States Department of Energy (DOE), under

Contract DE-AC02-06CH11357.

Affiliation

a. Argonne National Laboratory, Biosciences Division, 9700 S. Cass Ave., Argonne, IL 60439 USA.

† Current address: Oil-Dri Innovation Center, 777 Forest Edge Rd., Vernon Hills, IL 60061 USA.

* Corresponding author.

Electronic Supplementary Information (ESI) available: [AutoCAD files of the RMI-chip, SCAD files

of humidity chamber and microscope stage adapter, vector graphics of laser cut accessories, video

files of flow in microfluidic channel].

Fig.1 RMI-chip design. (a) The CAD design of the RMI-chip. A single growth channel is 36 mm long, 100

µm high, and 800 µm wide (a). The media and inoculation inlets are equipped with filters to avoid

occlusions during week-long experiments. Design files are available in the online supporting materials. (b)

Flow dynamics. Fluorescently labelled latex beads were used to measure media flow in the empty RMI-

chip. The root entrance and the inoculation inlets were plugged with stopper pins to mimic operational

flow regime of the chip. The latex beads suspended in Johnson were injected and the chip images,

followed by perfusion with Johnson solution at 1µL/min. The latex beads were washed from the upstream

chambers and the velocity of the beads were averaged. The measured and calculated values agreed.

Laminar flow was observed under the microscope.

(a)

(b)

Fig. 2. Imaging setup. (a) The imaging setup with humidity chamber and 3D printed stage adaptor. The

RMI-chip has an inlet for media, a separate port for inoculation, and a wide channel to accommodate the

tree seedling root without interfering with constant media flow. (b) The continuous flow incubation

chamber enables the cultivation of six seedlings in parallel (left). Details of an aspen seedling (right).

Fig. 3. Spatial and temporal colonization of aspen primary roots by *P. fluorescens* in the RMI-chip: *P.*

fluorescens SBW25 strains expressing dsRed or mNeonGeen were co-inoculated on aspen roots growing

in the RMI-chip. (a) After 3 days, bacterial cells colonize aspen root surface by filling intercellular spaces

(left panel) and adhering to plant cell surfaces (right panel). (b) After 13 days, SBW25 cells formed spatially

segregated red and green long patches of cells that correspond to dense biofilm-like assemblies. Typical

of mature biofilms, some internal void spaces and channels are indicated by arrows.

Fig. 4: A gallery of bacterial assemblies on the root surface. (a) Image showing diverse assemblies of

green SBW25 cells on the root surface: 1, individual cells; 2, cell doublets; 3, microcolonies; and 4, biofilms.

(b) Projection of a reconstructed 3D volume with side views reveals a vertical alignment of cells within the

biofilm. Scale bar indicates 10 μm length.

Fig. 5. ***B. subtilis*** **biosensors exposed to aspen root (panels a-c) and rice root (panel d) in the RMI-chip.**

(a) Among the root hairs, the xylose biosensor cells appear red from constitutive mCherry expression with

a very faint GFP (green) signal detected in some cells. Similar results were obtained with the ROS biosensor

(not shown). (b) Five days after inoculation, rare attached biosensor cells overwhelmingly express GFP,

indicating the presence of xylose in root exudates. Note the plant tissue autofluorescence causing cell

outlines to be green and some plant cells to be entirely red. (c) After 5 days, most ROS biosensor cells

expressed GFP, indicating the presence of ROS produced by the root. Plant tissue autofluorescence is also

detected. (d) Three days after inoculation of a rice seedling, the *B. subtilis* ROS biosensor displayed robust

colonization of the root with a strong expression of GFP, indicating production of ROS by the rice root.

Scale bar indicates 10 µm length.

Notes and references

1.   Brown, S.D., et al., *Twenty-one genome sequences from Pseudomonas species and 19 genome sequences from diverse bacteria isolated from the rhizosphere and endosphere of Populus deltoides.* J Bacteriol, 2012. **194**(21): p. 5991-3.

2.   Compant, S., et al., *Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects.* Appl Environ Microbiol, 2005. **71**(9): p. 4951-9.

3.   Bulgarelli, D., et al., *Structure and functions of the bacterial microbiota of plants.* Annu Rev Plant Biol, 2013. **64**: p. 807-38.

4.   Newton, A.C., et al., *Pathogenesis, parasitism and mutualism in the trophic space of microbe-plant interactions.* Trends Microbiol, 2010. **18**(8): p. 365-73.

5.   Gaiero, J.R., et al., *Inside the root microbiome: bacterial root endophytes and plant growth promotion.* Am J Bot, 2013. **100**(9): p. 1738-50.

6.   Dupuy, L.X.a.S., W.K., *Mechanisms of Early Microbial Establishment on Growing Root Surfaces.* Vadose Zone J. | Advancing Critical Zone Science, 2016. **15**(2).

7.   Varivarn, K., et al., *Colonization strategies of Pseudomonas fluorescens Pf0-1: activation of soil-specific genes important for diverse and specific environments.* BMC Microbiol, 2013. **13**: p. 92.

8.   Barea, J.M., et al., *Microbial co-operation in the rhizosphere.* J Exp Bot, 2005. **56**(417): p. 1761-78.

9.   Rodriguez, H. and R. Fraga, *Phosphate solubilizing bacteria and their role in plant growth promotion.* Biotechnol Adv, 1999. **17**(4-5): p. 319-39.

10.   Seneviratne G., W.M.L.M.A.W., Seneviratne K.A.C.N., Zavahir J.S., Kecskés M.L., Kennedy I.R. , *Importance of Biofilm Formation in Plant Growth Promoting Rhizobacterial Action*, in *Plant Growth and Health Promoting Bacteria*, M. D., Editor. 2010, Springer, Berlin, Heidelberg: Berlin, Heidelberg.

11.   Meier, M., E.M. Lucchetta, and R.F. Ismagilov, *Chemical stimulation of the Arabidopsis thaliana*

root using multi-laminar flow on a microfluidic chip. Lab Chip, 2010. **10**(16): p. 2147-53.

12.   Grossmann, G., et al., *The RootChip: an integrated microfluidic chip for plant science.* Plant Cell,

2011. **23**(12): p. 4234-40.

13.   Busch, W., et al., *A microfluidic device and computational platform for high-throughput live*

imaging of gene expression. Nat Methods, 2012. **9**(11): p. 1101-6.

14.   Agudelo, C.G., et al., *TipChip: a modular, MEMS-based platform for experimentation and*

phenotyping of tip-growing cells. Plant J, 2013. **73**(6): p. 1057-68.

15.   Jiang, H., et al., *Plant chip for high-throughput phenotyping of Arabidopsis.* Lab Chip, 2014. **14**(7):

p. 1281-93.

16.   Sanati Nezhad, A., *Microfluidic platforms for plant cells studies.* Lab Chip, 2014. **14**(17): p. 3262-

74.

17.   Parashar, A. and S. Pandey, *Plant-in-chip: Microfluidic system for studying root growth and*

pathogenic interactions in Arabidopsis. Applied Physics Letters, 2011. **98**(26).

18.   Massalha, H., et al., *Live imaging of root-bacteria interactions in a microfluidics setup.* Proc Natl

Acad Sci U S A, 2017. **114**(17): p. 4549-4554.

19.   Gottel, N.R., et al., *Distinct microbial communities within the endosphere and rhizosphere of*

Populus deltoides roots across contrasting soil types. Appl Environ Microbiol, 2011. **77**(17): p. 5934-44.

20.   Timm, C.M., et al., *Metabolic functions of Pseudomonas fluorescens strains from Populus deltoides*

depend on rhizosphere or endosphere isolation compartment. Front Microbiol, 2015. **6**: p. 1118.

21.   Weston, D.J., et al., *Pseudomonas fluorescens induces strain-dependent and strain-independent*

host plant responses in defense networks, primary metabolism, photosynthesis, and fitness. Mol Plant

Microbe Interact, 2012. **25**(6): p. 765-78.

22.   Shinde, S., et al., *Pseudomonas fluorescens Transportome Is Linked to Strain-Specific Plant Growth*

Promotion in Aspen Seedlings under Nutrient Stress. Front Plant Sci, 2017. **8**: p. 348.

23.   Noirot-Gros, M.F., et al., *Dynamics of Aspen Roots Colonization by Pseudomonads Reveals Strain-*

Specific and Mycorrhizal-Specific Patterns of Biofilm Formation. Front Microbiol, 2018. **9**: p. 853.

24.   Shinde, S.Z., S.; Collart, RF.; Cumming, JR.; Noirot, P.; Larsen, P., *Pseudomonas fluorescens*

increases mycorrhization and modulates expression of anti-fungal defense response genes in roots of

aspen seedlings. BMC Plant Biology, in press.

25.   Wilton, R., et al., *A New Suite of Plasmid Vectors for Fluorescence-Based Imaging of Root*

Colonizing Pseudomonads. Front Plant Sci, 2017. **8**: p. 2242.

26.   Babic, A., et al., *Efficient gene transfer in bacterial cell chains.* MBio, 2011. **2**(2).

27.   Radeck, J., et al., *The Bacillus BioBrick Box: generation and evaluation of essential genetic building*

blocks for standardized work with Bacillus subtilis. Journal of Biological Engineering, 2013. **7**(1).

28.   Overkamp, W., et al., *Benchmarking Various Green Fluorescent Protein Variants in Bacillus subtilis,*

Streptococcus pneumoniae, and Lactococcus lactis for Live Cell Imaging. Applied and Environmental

Microbiology, 2013. **79**(20): p. 6481-6490.

29.   O'Toole, G., H.B. Kaplan, and R. Kolter, *Biofilm formation as microbial development.* Annu Rev

Microbiol, 2000. **54**: p. 49-79.

30.   Melaugh, G., et al., *Shaping the Growth Behaviour of Biofilms Initiated from Bacterial Aggregates.*

PLoS One, 2016. **11**(3): p. e0149683.

31.   Badri, D.V. and J.M. Vivanco, *Regulation and function of root exudates.* Plant Cell Environ, 2009.

32(6): p. 666-81.

32.   Allard-Massicotte, R., et al., *Bacillus subtilis Early Colonization of Arabidopsis thaliana Roots*

Involves Multiple Chemotaxis Receptors. MBio, 2016. **7**(6).

33.   Kim, L., A. Mogk, and W. Schumann, *A xylose-inducible Bacillus subtilis integration vector and its*

application. Gene, 1996. **181**(1-2): p. 71-6.

34.   Hoover, S.E., et al., *Changes in DnaA-dependent gene expression contribute to the transcriptional*

and developmental response of Bacillus subtilis to manganese limitation in Luria-Bertani medium. J

Bacteriol, 2010. **192**(15): p. 3915-24.

35.   Sharma, P.J., A.B.; Dubey, R.S.; Pessarakli, M, *Reactive Oxygen Species, Oxidative Damage, and*

Antioxidative Defense Mechanism in Plants under Stressful Conditions. Journal of Botany, 2012. **2012**: p.

26.

36.   Jha, Y. and R.B. Subramanian, *Characterization of root-associated bacteria from paddy and its*

growth-promotion efficacy. 3 Biotech, 2014. **4**(3): p. 325-330.

(a)

(b)

(a)

inlet 1

seedling

inlet 2

(b)

no flow

1 µl/min

100 µm