1
The effect of ocean warming on black sea bass (*Centropristis striata*) aerobic scope and hypoxia tolerance

3
Ocean warming and *C. striata* physiology

Emily Slesinger[1][*], Alyssa Andres[2], Rachael Young[1], Brad Seibel[2], Vincent Saba[3], Beth Phelan[4],
John Rosendale[4], Dan Wieczorek[4], Grace Saba[1]

[1] Center for Ocean Observing Leadership, Department of Marine and Coastal Sciences, School of
Environmental and Biological Sciences, Rutgers University, New Brunswick, NJ 08901, USA
[2] College of Marine Science, University of South Florida, St. Petersburg, FL 33701, USA
[3] NOAA Northeast Fisheries Science Center, Geophysical Fluid Dynamics Laboratory,
Princeton, NJ 08540
[4] NOAA Northeast Fisheries Science Center, James J. Howard Laboratory, Highlands, NJ 07732
[*]Corresponding Author: slesinger@marine.rutgers.edu, 848-932-3452

Key Words

Summary Statement

Laboratory-based physiological studies focused on aerobic scope and hypoxia tolerance of black

sea bass presented here support the recently observed poleward shift of their range on the U.S.

Northeast Shelf.

Abstract

Over the last decade, ocean temperature in the U.S. Northeast Continental Shelf (U.S. NES) has

warmed faster than the global average and is associated with observed distribution changes of the

northern stock of black sea bass (*Centropristis striata*). Mechanistic models based on

physiological responses to environmental conditions can improve future habitat suitability

projections. We measured maximum (MMR), resting metabolic rate (RMR), and hypoxia

tolerance (S_{crit}) of the northern adult black sea bass stock to assess performance across the known

temperature range of the species. A subset of individuals was held at 30˚C for one month

($30_{chronic}$°C) prior to experiments to test acclimation potential. MMR and absolute aerobic scope

(AAS = MMR-RMR) reached a maximum at 24.4°C (AS: 367.21 mgO_2 kg^{-1} hr^{-1}) while S_{crit}

continued to increase in proportion to RMR up to 30°C. The $30_{chronic}$°C group had a significant

decrease in MMR and AAS but RMR or S_{crit} were not affected. This suggests a decline in

performance of oxygen demand processes (e.g. muscle contraction) beyond 24°C despite

maintenance of oxygen supply. The Metabolic Index, calculated from S_{crit} as an estimate of

potential aerobic scope, closely matched the measured factorial aerobic scope (FAS =

MMR/RMR) and declined with increasing temperature to a minimum below 3. This may

represent a critical value for the species. Temperature in the U.S. NES is projected to increase

above 24°C in the southern portion of the northern stock's range. Therefore, these black sea bass

will likely continue to shift north as the ocean continues to warm.

Introduction

Marine environments are progressively warming as a consequence of climate change

(Belkin, 2009). Along the U.S. Northeast Shelf (U.S. NES), ocean temperature is rising faster

than the global average (Pershing *et al.*, 2015; Caesar *et al.*, 2018) resulting in a significant

temperature increase (Friedland and Hare, 2007; Kavanaugh *et al.*, 2017). Sea surface and

bottom temperatures in the U.S. NES are projected to rise an additional 4.1°C and 5.0°C,

respectively, along the U.S. NES (Saba *et al.*, 2016; Kleisner *et al.*, 2017). Contemporary ocean

warming in the U.S. NES has been associated with distribution shifts of many economically and

ecologically important fish species both in latitude and/or depth (Nye *et al.*, 2009; Bell *et al.*,

2015; Kleisner *et al.*, 2016, 2017; Morley *et al.*, 2018), associated with tracking local climate

velocities (Pinsky *et al.,* 2013). Understanding and projecting shifts in fish distribution will be

important for characterizing potential ecological and economic impacts and anticipating and

resolving fishery management conflicts (Pinsky *et al.*, 2018).

Temperature directly affects metabolic rates of marine ectotherms (Clarke and Johnston,

1999; Verberk *et al.*, 2016) and is believed to set the boundaries of species ranges (Pörtner and

Farrell, 2008; Deutsch *et al.*, 2015). One explanation for the effects of temperature on

ectothermic species, oxygen and capacity-limited thermal tolerance (OCLTT; Pörtner, 2010),

postulates that thermal limitation occurs due to a mismatch in oxygen demand and supply at sub-

optimal temperatures, which ultimately determines suitable thermal habitat (Pörtner and Knust,

2007). In this framework, the thermal optimum occurs where absolute aerobic scope (AAS), the

difference between maximum (MMR) and resting metabolic rate (RMR) (Schulte, 2015), is

highest. RMR is the cost of maintenance for an organism and increases with temperature (Clarke

and Johnston, 1999). MMR may be constrained differently across a temperature range by oxygen

uptake, transport or utilization. The drop in AAS beyond the thermal optimum is associated with

the failure of MMR to increase relative to the continuing rate of increase in RMR (Farrell, 2016).

Absolute AAS is thought to represent the capacity for oxygen uptake, beyond that supporting

maintenance metabolism, that can be utilized for activities that promote individual fitness (e.g.

growth, reproduction, predator avoidance; Pörtner & Peck 2010). Hence, the adaptive benefit of

living at suitable temperatures to maintain metabolic scope may provide a mechanistic

explanation for where fish may be distributed in their environment.

While the general distribution of fishes is broadly confined by thermal preferences,

oxygen availability can further constrain suitable habitat. The hypoxia tolerance of a fish can be

estimated as the critical oxygen saturation level (S_{crit}), the $\%O_2$ below which oxygen supply

cannot match the demands of maintenance metabolism. Further reductions in $\%O_2$ cause a

proportional decrease in RMR (Schurmann and Steffensen, 1997). Below the S_{crit}, a fish has

time-limited survival as ATP production progressively relies on unsustainable anaerobic

pathways and metabolic suppression (Nilsson and Renshaw, 2004; Seibel, 2011). Generally, a

fish with a low S_{crit} is tolerant of lower sustained oxygen levels (Claireaux and Chabot, 2016). As

ocean temperature increases, oxygen demand concomitantly increases (Del Toro-Silva *et al.*,

2008; Capossela *et al.*, 2012), potentially reducing hypoxia tolerance (Collins *et al.*, 2013;

McDonnell and Chapman, 2015). The S_{crit} further provides a means of calibrating the Metabolic

Index, which is a ratio of oxygen supply to demand that provides an estimate of sustained

factorial aerobic scope and metabolically suitable habitat (Deutsch et al., 2015). At S_{crit}, by

definition, supply exactly matches demand allowing statistical estimation of the equations'

physiological parameters.

The northern stock of black sea bass (*Centropristis striata*) on the U.S. NES extends from

Cape Hatteras to the Gulf of Maine and is centered in the Mid-Atlantic Bight (MAB; Roy *et al.*,

2012). These fish seasonally migrate from the continental shelf edge in cooler months to inshore

depths (5-50m) in warmer months (Musick and Mercer, 1977; Moser and Shepherd, 2008).

Seasonally migrating black sea bass thus experience a wide range of temperatures throughout the

year, ranging from 6°C during winter and up to 27°C during summer/early fall months (Steimle

et al., 1999). Off the coast of New Jersey, periodic hypoxic events can occur during the summer

as a result of high biological activity (Schofield *et al.*, 2012) fueled by upwelling of nutrient rich

waters (Glenn *et al.*, 2004). Therefore, during the warm summer months, black sea bass are

potentially subject to hypoxia in this region, contributing to the oxygen limitation that contains

suitable habitat.

The northern stock of black sea bass may already be exhibiting poleward shifts, likely

due to ocean warming (Hare *et al.*, 2016; Kleisner *et al.*, 2017). Evidence for current black sea

bass distribution shifts comes primarily from bottom trawl survey data (Kleisner *et al.*, 2017),

and is supported anecdotally through fishermen. Laboratory-based process studies focused on the

physiology of an organism provide detailed mechanistic relationships between the environment

and the animal (Wikelski and Cooke, 2006). The objectives of this study were to determine the

AAS and S_{crit} for the northern stock of adult black sea bass. We measured AAS and S_{crit} at a

range of temperatures similar to those experienced by black sea bass during their summer inshore

residency to compare thermal optima, if present, and metabolic index against the known

temperature range of the species. Results from these physiological studies may be useful for

modeling suitable habitat based on environmental parameters (Lefevre *et al.*, 2017) and model

black sea bass distributions (e.g., Manderson *et al.*, 2011; Deutsch *et al.*, 2015), and projecting

future distribution shifts in black sea bass with continued ocean warming.

Methods

Fish Collection and Husbandry

Adult black sea bass (*Centropristis striata*) from the northern stock (length = 221-

398mm; weight = 193.7-700.4g) were collected off the coast of New Jersey, USA at depths of

15-20m in early June from Sea Girt and Manasquan Reefs by fish traps (2016), and from local

reefs off Sandy Hook by hook-and-line (2017). Fish were housed in the NOAA James J. Howard

Marine Laboratory, held at ambient temperature (22 ± 1°C) and salinity (26ppt), maintained at a

natural photoperiod for New Jersey summer, and fed daily to satiation on a diet of sand lance and

silversides. Water temperature and salinity was monitored daily using a YSI (Pro-30; Yellow

Springs, Ohio, USA), and water chemistry remained at suitable levels (< 20 uM nitrate,

undetectable nitrite, < 0.05 uM ammonia, pH range of 7.98-8.04). Fish were acclimated to

captive conditions two weeks prior to the trials, after which all experimental fish ate regularly

and were in good condition. Any fish exhibiting apparent health issues (i.e. lack of appetite,

difficulties with buoyancy or orientation) were not used in experiments. After acclimation, fish

were measured for length (TL mm), weight (g), and tagged with individually numbered T-bar

Floy tags inserted underneath the dorsal rays. For each temperature treatment, fish were

acclimated at a rate of 2°C day^{-1} to reach experimental temperature, then held at the target

treatment temperature for at least 48hr prior to the start of experiments. Fish were starved 48hr

prior to the start of each experiment to eliminate effects of specific dynamic action (Chabot *et*

al., 2016a). At the end of each experiment, fish were euthanized with an overdose of MS-222

(250 mgL^{-1}). Fish were collected under New Jersey permits #1610 & #1717. Treatment of all

animals was held in accordance with Rutgers IACUC protocol 15-054.

Experimental Set-Up

Experimental tanks (1,200L) were filled with treated seawater from Sandy Hook Bay that

continuously circulated through a closed system. Circulating seawater was treated using filters

(sand and biological) and UV-light, and salinity was adjusted to mimic average summertime

inshore NJ bottom water (32±1). Experimental temperatures were achieved using in-line chillers

(Aqua Logic Delta Star; San Diego, California, USA) and/or titanium exchanger heaters

(Innovative Heat Concepts, Homestead, Florida, USA), and maintained at ±1°C from target

temperature.

Metabolic rates were measured using intermittent respirometry under the protocols

outlined in Clark *et al.*, (2013) and Svendsen *et al.*, (2016a). Flow-through respirometers (13.5

liter volume; 23[H]x26[W]x37[L] cm plexiglass) were placed into the two experimental tanks

(two respirometers per tank; four respirometers per trial). Flush pumps (Eheim Universal 600

l//h; Deizisau, Germany) connected to the respirometer were used to pull water from the

surrounding temperature bath to replenish dissolved oxygen and eliminate metabolic waste

buildup within the respirometer. The duration and timing of flushes set the intermittent cycles,

which were controlled through a pre-determined time sequence using a DAQ-M instrument

(Loligo Systems; Viborg, Denmark), and were set based on the trial temperature so that oxygen

saturation was never below 75% (Svendsen *et al.*, 2016b). For each closed measure period

(when flush pumps were off), the rate of decline in dissolved oxygen concentration within the

sealed chamber was used to calculate a mass specific rate of oxygen consumption, or metabolic

rate (MO_2: $mgO_2\,kg^{-1}\,hr^{-1}$). A closed recirculation loop connected with a smaller pump (Eheim

Universal 300 l/h; Deizisau, Germany) was also utilized to uniformly disperse dissolved oxygen

within the chamber and provide waterflow across the oxygen dipping probe optical mini sensor

(PreSens Pst3; Regensburg, Germany). Oxygen probes were calibrated in accordance with the

supplier's manual (Oxygen dipping probe PSt3, PreSens GmbH, Regensburg, Germany) and

checked with a YSI (ProSolo ODO; Yellow Springs, Ohio, USA) that was calibrated in 100%

and 0% oxygen saturation sample waters. Autoresp computer software (Loligo Systems; Viborg,

Denmark) and a Witrox-4 instrument (Loligo Systems; Viborg, Denmark) were used to

continuously monitor dissolved oxygen and temperature within the chamber over the course of

the experiment.

For hypoxia experiments, intermittent respirometry was also used to avoid a CO_2 and

metabolite build up (Rodgers et al. 2016). Each respirometer flush pump was connected to an

external water bath that was filled with the same system water. Within the external water bath, a

pump (Eheim Universal 1200 l/h; Deizisau, Germany) connected to a piece of Tygon tubing held

an oxygen optode to monitor source O_2 and served as a mixing device. Also within the external

water bath, four small microdiffusers were connected to a N_2 gas canister (Schurmann and

Steffensen, 1997) to allow for diffusion of nitrogen gas into the external bath and subsequent

displacement of O_2 within the external water bath and control of environmental $\%O_2$ within the

chambers over the course of the hypoxia experiment.

Background respiration was measured by taking background MO_2 (MO_{2br}) pre- and post-

trial in empty chambers for ~1.5hr. A linear regression between pre- and post-MO_{2br} was used to

apply a correction factor to each MO_2 value recorded throughout an experiment.

Experiments were conducted at a range of temperatures (12, 17, 22, 24, 27 and 30°C). An

additional subset of black sea bass were held at 30°C for one month to test acclimation potential

(Table 1). We used two different methods in an attempt to elicit maximum metabolic rate

(MMR): exhaustive-chase and swim-flume. For the chase method, individual black sea bass

(Table 1) were placed in a 4ft-diameter chase tank filled with water from the experimental tanks.

Fish were chased via tactile stimulation on the caudal tail and were determined exhausted when

unresponsive to further tactile stimulation and air exposure. Fish were then immediately

transferred to individual respirometers that were sealed within ~1 min from the end of the chase

and remained in the metabolic chambers for ~23hr allowing for resting metabolic rate (RMR)

measurement (Chabot *et al.*, 2016b). Once the experiment was finished, fish were either

exercised in a swim-flume (Loligo Systems 90L; Viborg, Denmark) after a 24hr rest period or

remained in the chamber for the hypoxia experiments. For the swim-flume, fish were tested

using a sprint protocol. Swimming speed was increased over a 5min period up to 0.95 BL s^{-1} with

a flush pump on as the fish adjusted to the flume. After an adjustment time (~10 minutes), the

speed was increased over a period of 5 minutes until the fish was sprinting (designated as >10

tail bursts during 30s intervals and an inability to maintain position in the working section

without burst swimming). Once a fish was sprinting, the flush pump was turned off and the

flume was sealed. Fish were held at their sprint speed for at least 10 minutes or until failure,

determined when the fish rested at the backgrate for >10s. Aerobic scope was calculated in

absolute (AAS = MMR-RMR) and factorial terms (FAS = MMR/RMR). An additional subset of

black sea bass was held at 30°C for one month to test acclimation potential. In 2016, fish were

only tested at 24, 27, and 30°C due to restrictions in maintaining temperatures. See Table 1 for

sample size at each temperature.

Critical $\%O_2$ determinations

231    Hypoxia (S_{crit}) experiments were conducted on the last 4 fish of each temperature

232 treatment trial. This allowed us to reliably use fish that were already acclimated to the

233 respirometers and had reached RMR overnight. Starting with 100% dissolved oxygen (DO)

234 saturation within the chambers, environmental %O_2 was incrementally decreased by 10%. Three

235 intermittent (flush, wait, measure) cycles were measured per DO level until S_{crit} was determined

236 to have been reached, indicated by a substantial decline in fish metabolic rate or loss of

237 equilibrium, and the experiment ended.

239 *Data Analysis*

240    Fish MO_2 was calculated via the AutoResp program from the slope of oxygen saturation

241 decline during each closed measurement period. Validation of each MO_2 value was conducted

242 using R_2 values from each measure period. MO_2 measurements with R_2 values < 0.9 were not

243 used.

244    We report our baseline metabolic rate as resting (RMR) instead of standard (SMR)

245 metabolic rate because the amount of time in the chamber was ~23 hours, which does not allow

246 for determination of full diel cycles (Chabot *et al.*, 2016b). RMR was calculated from a

247 truncated dataset without the first two hours of elevated MO_2 values following exercise and by

248 using the 20th quantile of the RMR data in the *calcSMR* package in R (Chabot *et al.*, 2016b).

249 Briefly, a frequency distribution of MO_2 values from the truncated data set was created and the

250 values at the 20th quantile were taken to calculate RMR. MMR in the chase protocol was defined

251 as the highest MO_2 value recorded during the trial and MMR was calculated for the duration of

252 the sprint interval in the swim-flume. AAS was taken as the difference between MMR and RMR.

253 There was a significant effect of mass on MO_2 ($F_{1,117}$ = 4.651; P < 0.05; Fig. 1). Therefore, the

254 effect of temperature on MO_2 was analyzed using a one-way ANCOVA with weight as a

255 covariate. A Tukey's HSD *post hoc* was used to determine significant pair-wise comparisons

256 between temperatures. MO_2 was adjusted for weight using the estimated marginal means from

257 the ANCOVA centered around a fish mean weight of 346.9g. Because the average weight of fish

258 in the 24°C treatment (253.9g) was lower than the mean weight of all other experimental fish, the

259 adjusted MO_2 for 24°C was slightly overestimated and had larger standard error for AAS and

260 MMR. Curves for aerobic scope were modeled using a 3rd degree polynomial fit and were used

to estimate a thermal optimum (temperature at the highest AS). All graphs and results report the

adjusted MO_2 (MO_{2adj}).

Q_{10} values were calculated for the adjusted MO_2 between temperature increments, and

between the range of temperatures using the formula:

$$Q_{10} = \frac{R_2}{R_1}^{10/(T_2-T_1)}$$

where Q_{10} is the temperature coefficient for MO_2, R_1 is the MO_2 at T_1 and R_2 is the MO_2 at T_2.

S_{crit} was determined by fitting two regression lines through the data: one through the region

where RMR was independent of $\%O_2$ and one through the portion where MO_2 decreased linearly

with a decrease in $\%O_2$. The intersection of the two regression lines is the critical point used for

S_{crit} (Yeager and Ultsch, 1989). This was analyzed using R code in the *calcO2crit* package from

Claireaux and Chabot (2016). Because we had a sample size of 4 fish per temperature trial, a

power analysis was run to determine the statistical power of this small sample size and four fish

provided enough statistical power (*Power* = 1, *n* = 4, *f* = 1.71, *sig. level* = 0.05). A one-way

ANOVA was used to assess the effect of temperature on S_{crit} and a Tukey's HSD *post hoc* test

was used to determine significant pair-wise comparisons between temperatures.

All statistical analyses were performed in R (Version 3.4.1). Data were checked for

assumptions of normality by the visual Q-Q norm plot and statistically with the Shapiro-Wilk

test where *P* > 0.05 indicate normally distributed data. Assumptions of homogeneity were

assessed using the Levene's test where a *P* > 0.05 indicates homogeneity. Data that did not fit

assumptions of normality were log-transformed prior to further statistical analysis. Data are

presented as mean ± SE and results from statistical analyses are defined as significant at *P* <

0.05.

Results

Metabolic rates and aerobic scope

RMR increased significantly with temperature (Figs 2a and 2b) and there was a

significant effect of weight and temperature*weight interaction on RMR (*P* < 0.05; Table 2).

While the results for the two MMR methods differed considerably, temperature, weight and

temperature*weight interaction all had a significant effect on MMR using either method (*P* <

0.05; Table 2). The chase MMR increased continuously with temperature, while flume MMR

increased with temperature until ~22°C (Fig 2a & 2b). The MMR values from the flume were

consistently higher across the temperature range than from the chase method, indicating that the

metabolic rate reached during the chase likely was not the maximum possible for this species.

While the chase method did not achieve MMR, it still provided an estimate of

submaximal exercise performance across a temperature range. The MMR using the chase

method increased continuously with temperature and reached a maximum adjusted value of

396.65 ± 11.48 mg O_2 kg^{-1} hr^{-1}at 30.0°C (the highest temperature measured; Table 3; Fig 1a).

The MMR measured using the flume reached a maximum of 497.96 ± 21.92 mgO_2 kg^{-1} hr^{-1} at

27°C (Table 3; Fig 1b). The absolute aerobic scope using the flume method reached a

maximum, typically referred to as "T_{opt}" at ~24.4°C (Fig. 2B). There was a significant effect of

temperature, weight, and the temperature*weight interaction on AAS ($P < 0.05$) using both

MMR methods (Table 2). Using different MMR methods resulted in differences in the shape of

the AAS curve and the estimated thermal optimum with consequences for its interpretation. All

RMR, MMR and AAS values are reported in Table 3 and Q_{10} values are reported in Table 4.

Critical %O_2

The critical %O_2 (S_{crit}) increased significantly with increasing temperature (Fig 4a; $F_{5,18} =$

14.023, $P < 0.05$), directly correlated with RMR (Fig 4b). There was no significant difference

between 12˚C (19.65 ± 1.72 %O_2), 17˚C (21.325 ± 1.75 %O_2) and 22˚C (21.80 ± 1.21 %O_2), but

S_{crit} increased significantly at 27˚C (31.60 ± 1.67 %O_2) and further at 30˚C (37.875 ± 3.39 %O_2).

However, non-significance between 12, 17 and 22°C could be due to low sample size.

Chronic high temperature exposure

The 30$_{chronic}$°C group AAS using both MMR methods significantly decreased when

compared to the 30°C treatment where fish were only held at this temperature for a week. Based

on Tukey *post hoc* differences, RMR did not change significantly between the 30$_{chronic}$°C and

30°C treatments but there was a significant decrease in MMR between the 30°C and 30$_{chronic}$°C

treatments. There was no significant difference in S_{crit} between 30°C and 30$_{chronic}$°C treatments.

Discussion

The primary objective of this study was to assess the use of physiological measurements

to determine habitat suitability for the northern stock of black sea bass at current and future

temperatures. We measured the oxygen consumption rate during two different exercise

protocols. The flume yielded much higher metabolic rates, indicating that the chase method did

not elicit MMR. Using the flume MMR, we found that AAS peaked at 24.4°C. S_{crit} increased

with increasing temperatures as is typical of most (but not all, Wishner et al., 2018) animals,

including fishes (Rogers et al. 2016). Chronic exposure to 30°C resulted in a significant drop in

AAS with no change in RMR or S_{crit}. That S_{crit} increased with temperature in proportion to RMR,

while MMR in the flume did not, suggests that chronic exposure to high temperature did not alter

the capacity for oxygen uptake and transport, but that the capacity to generate ATP was reduced,

perhaps due to a decrement in muscle function. The capacity for submaximal exercise (oxygen

consumption following a chase to exhaustion) also increased across the entire temperature range

further suggesting that the failure was not in the capacity for oxygen supply. Chronic exposure

to 30°C led to further reductions in MMR using both methods, but no loss of oxygen supply

capacity as estimated from S_{crit}, suggesting continued deterioration in muscle function with

longer exposure to warm temperatures.

Absolute AS typically increases with temperature up to a point, often termed "optimal",

and then declines at higher temperatures resulting in a roughly bell-shaped curve as has been

identified in fishes that include, but is not limited to, juvenile European sea bass *Dicentrarchus*

labrax (Claireaux *et al.*, 2006), turbot *Scophthalmus maximus* (Mallekh and Lagardere, 2002),

coho salmon *Oncorhynchus kisutch* (Raby *et al.*, 2016), and sockeye salmon *Oncorhynchus*

nerka (Farrell *et al.*, 2008). However, some studies have found left- or right-skewed curves (e.g.

Healy and Schulte 2012) while others find that AAS continues to increase up to the critical

temperature for the species (i.e. no temperature optimum for AAS is identifiable; e.g. Norin *et al.*

2014). In our study, the black sea bass AAS curve was more bell-shaped with an estimated

optimal temperature of 24.4°C. Bottom temperature in the southern portion of the black sea bass

range typically hovers around 24-26°C during the summer (Houghton *et al.* 1982; Castelao *et al.*

2008; from U.S. East Coast Regional ESPreSSO model, Wilkin and Hunter 2013), which would

suggest this area to be thermally optimal. However, if the loss in AAS at higher temperatures is

due to a failure in muscular performance rather than potential oxygen supply, then 24°C may

represent a maximum tolerable temperature rather than a temperature that allows optimal

performance. In support of this interpretation, the Metabolic Index (which closely matches the

factorial aerobic scope) declines with increasing temperature toward levels (~3 at 27°C in black

sea bass) known to limit the geographic range of some species (Deutsch et al., 2015). While the

average bottom temperature in the southern portion of the northern stock of black sea bass is near

24°C during the summer months, there has still been a consistent expansion of their range

northward into lower temperatures (NEFSC 2017) further suggesting that the temperature

eliciting maximum AAS is not, in fact, optimal. It is important to note that AAS is only a

measured capacity to supply oxygen under maximum sustained exercise (Farrell, 2016). The

required scope for other metabolic expenses (i.e. feeding, digestion; Holt and Jørgensen, 2015)

change with temperature in unknown ways and metabolic needs can change seasonally and with

ontogeny (Clark *et al.*, 2013). Thus, AAS may in this case be an inappropriate predictor of

fitness.

Black sea bass in the $30_{chronic}$°C treatment did not acclimate, indicated by no change in

RMR or S_{crit} and a significant decrease in their MMR and AS. Norin et al. (2014) similarly found

that MMR and AAS in juvenile barramundi decreased significantly following 5 weeks at the

highest study temperature (38°C).   However, unlike black sea bass in our study, the juvenile

barramundi RMR also decreased after the 5-week exposure. This same response has also been

found for short-horn sculpin (*Myoxocephalus scorpius*) whose RMR was restored after being

held at 16°C for 8 weeks to RMR values that were measured at 10°C (Sandblom *et al.*, 2014).

The decrease in RMR can be an acclimation response to lower their energetic costs at high

temperatures, but comes with its own caveats as this sometimes can reduce MMR. Importantly,

black sea bass in the $30_{chronic}$°C treatment may have suffered stress from long-term captivity,

which could also reduce AAS and time did not permit for a control chronic trial at a cooler

temperature (although all fish were held for at least 5 days). Understanding the acclimation

potential of black sea bass would benefit from future studies focusing on effects of a chronic

treatment at each temperature tested.

S_{crit} decreased as temperature increased, most likely caused by rising RMR with higher

temperatures, which has been shown in a majority of fish hypoxia studies (e.g. Schurmann and

Steffensen, 1997; although see Wishner et al., 2018). The $30_{chronic}$°C group did not have a

significant decrease in hypoxia tolerance compared to the 30°C group, which agrees with no

change in RMR between the two 30°C treatments. This suggests that the reduced MMR in

$30_{chronic}$°C animals resulted from reduced capacity to generate ATP, rather than to supply

oxygen. Black sea bass had lower S_{crit} than striped bass *Morone saxatilis* (Lapointe *et al.*, 2014)

and summer flounder *Paralichthys dentatus* (Capossela *et al.*, 2012), two important species

found throughout the MAB that periodically experience hypoxic water during the summer

months. However, when compared with fish that frequently experience hypoxia, such as

largemouth bass and crucian carp (Yamanaka *et al.*, 2007) and juvenile barramundi (Collins *et*

al., 2013), black sea bass were less hypoxia tolerant, especially in warmer water. Deutsch et al.

(2015) proposed a metabolic index (MI), as the ratio of oxygen supply to demand, which is

effectively an estimate of a species' time-averaged aerobic scope. By definition, the MI is equal

to 1 at the S_{crit}. A minimum MI of 2-5, indicating the capacity to supply oxygen at 2-5x the rate

required at rest, is supportive of a population and delineates the equatorward distribution limit in

the few species studied to date. Black sea bass factorial AS and MI both decreased with

increasing temperature (Fig. 6). During the summer months when bottom water temperature is

warmest along the coastal MAB, periodic hypoxic events occur after large phytoplankton blooms

in the surface waters. In the past, these hypoxic events decreased bottom water PO_2 below

~5.5kPa (26% saturation; 2.2 mg L^{-1} at 14°C Schofield et al. 2012), providing a metabolic index

of ~1.3 at those temperatures for black sea bass. Such environments can be tolerated for short

periods but are not likely supportive of a thriving population. At 30°C, even air-saturated water

provides a MI of only 2.6 which is near the physiological limits of many species (Deutsch *et al.*,

2015). Therefore, when determining the suitable habitat, both temperature and oxygen must be

taken into consideration as the interacting effects of these variables will effectively decrease

optimal thermal habitat.

The chase method did not elicit MMR in black sea bass since MMR from the flume

method was consistently higher. Which method, chase or flume, provides a more reliable

measure of MMR and AAS is actively debated (Norin and Clark, 2016; Killen *et al.*, 2017).

Whether a maximum rate of oxygen uptake is achieved by either method could depend on the

type of swimming the study fish species naturally exhibits in the wild. Norin et al (2014)

purposefully used a chase method for juvenile barramundi (*Lates calcarifer*), an ambush

predator, that typically swims in quick bursts. In other cases, a fish will exhibit marked post-

exercise oxygen consumption (EPOC; Plambech *et al.*, 2013), sometimes eliciting MMR minutes

to hours after the cessation of exercise (Reidy *et al.*, 1995). The swim flume method may be

more appropriate for endurance swimming exhibited by pelagic fish such as tunas (Killen *et al.*,

2017). Different MMR methods may promote a certain type of swimming which could exhaust a

fish before reaching MMR by depleting anaerobic stores, a noteworthy contributor to AAS

(Ejbye-Ernst *et al.*, 2016). For this study, we employed a sprint protocol for the swim-flume,

which prompted similar burst swimming as in the chase method. However, during the chase

protocol, black sea bass switched almost immediately to burst swimming accompanied with

quick turning/flipping movements, compared to a slower transition and continuously straight

burst swimming in the swim flume. The differences in MMR between the two methods could

have been related to different swimming types, durations and/or speeds which could recruit more

anaerobic resources (Svendsen *et al.*, 2010) in the chase method, leading to exhaustion before

reaching MMR.

In summary, the results from this study indicate that the northern stock of black sea bass

reach a peak in AAS at ~24°C, which is warmer than in the northern portion of their range in the

U.S. NES. The MI of 3.8 in air-saturated water, calculated from S_{crit} at 24°C, suggests relatively

limited scope for sustained activity at that temperature (Deutsch *et al.*, 2015). We suggest that,

rather than an optimal temperature, the peak in MMR and AAS indicates the maximum tolerable

temperature, beyond which black sea bass experience a failure in some subcellular or organ

systems that contribute to muscle performance. Our study only used individuals from the

northern stock that were collected during the summer off of the New Jersey coastline. Metabolic

research on the southern stock (south of Cape Hatteras, NC) and/or individuals from the northern

stock in waters outside of New Jersey could reveal variation in some of these physiological

metrics. However, the distribution of the northern stock of black sea bass has shifted northward

(Kleisner *et al.*, 2017) and this newly expanded habitat is almost 10°C colder than their apparent

thermal optimum for AAS. We believe the preference for cooler waters reflects physiological

limitation at higher temperatures, including possible limitation of oxygen supply relative to

demand for growth and reproduction (reduced Metabolic Index) despite maintenance of oxygen

supply capacity. However, many other factors, including food availability, additional energetic

costs (e.g., evading predators, mating), or lower optimal temperatures for other critical processes

may be important. This suggests AAS may not be the most appropriate predictor for habitat

suitability in this species. Additionally, the northern stock of black sea bass population size has

been increasing in the last decade (NEFSC 2017), and this increase in biomass could be pushing

part of the population northward. Regardless, the chronic exposure experiments presented here

suggest little capacity for physiological adjustment to future temperatures. Black sea bass

thermal habitat may shrink considerably in the southern region of the MAB as bottom water

temperatures reach >27°C and continue to expand into the northern region of the MAB as ocean

waters continue to warm, significantly impacting fisheries in these two regions.

Acknowledgements

We thank Doug Zemeckis and Captain Chad Hacker (R/V Tagged Fish) for helping us collect

black sea bass; Richard Brill and Andrij Horodysky for providing insight and suggestions for the

study design; and the students at the Marine and Science Technology Academy for their

assistance in animal husbandry. We also acknowledge the NOAA James J. Howard Laboratory

personnel for their support and help throughout the study.

Competing Interests

The authors declare no competing interests.

Funding

The research was supported by the NOAA Office of Oceanic and Atmospheric Research (OAR),

Coastal and Ocean Climate Applications (COCA) Program (NA15OAR4310119).

List of Abbreviations

AAS　　　　　Absolute aerobic scope

FAS　　　　　Factorial aerobic scope

MO_2　　　　Oxygen consumption rate

MO_{2adj}　　　Adjusted oxygen consumption rate

MO_{2br}　　　Background oxygen consumption rate

MMR　　　　Maximum metabolic rate

RMR　　　　Resting metabolic rate

S_{crit}　　　　Critical oxygen saturation

SMR　　　　Standard metabolic rate

$\%O_2$　　　　Oxygen saturation

Tables
477
478 Table 1. Number of black sea bass used in both the chase and flume MMR trials at each
479 temperature.
480

Temperature (°C)	Sample Sizes	
	Chase MMR	*Flume MMR*
12	16	10
17	16	12
22	16	12
24	17	0
27	30	12
30	17	10
30$_{chronic}$	9	4

482 Table 2. ANCOVA results for AS, MMR (both methods) and RMR. Bolded P-values are
483 significant.
484

Variable	Effect	DF	F-value	P-value
AAS (chase)	Temperature Weight Temperature*weight	6, 105 1, 105 6, 105	13.877 2.082 2.106	**< 0.001** > 0.05 0.0586
AAS(flume)	Temperature Weight Temperature*weight	5, 48 1, 48 5, 48	6.185 6.599 4.033	**< 0.001** **< 0.05** **< 0.01**
MMR (chase)	Temperature Weight Temperature*weight	6, 105 1, 105 6, 105	50.327 9.267 2.281	**< 0.001** **< 0.01** **< 0.05**
MMR (flume)	Temperature Weight Temperature*weight	5, 48 1, 48 5, 48	16.244 8.927 3.147	**< 0.001** **< 0.01** **< 0.05**
RMR	Temperature Weight Temperature*weight	6, 105 1, 105 6, 105	136.613 12.282 2.489	**< 0.001** **< 0.001** **< 0.05**

486　Table 3. The MO_{2adj} mean ± S.E. values for RMR, MMR (both methods), AS (both methods)
487　and S_{crit} for each temperature treatment. * = the adjusted MO_2 values for MMR and AAS at
488　24°C are overestimated due to the average weight of fish in the 24°C group to be smaller than
489　the average weight for all study fish combined.
490
491

Temperature (°C)	RMR	Chase MMR	Flume MMR	Chase AS	Flume AS	S_{crit} (% O_2 saturation)
12	46.44±1.94	169.12±11.78	286.57±24.02	117.67±7.55	242.67±24.24	19.65
17	65.27±3.27	215.05±14.15	369.27±24.47	143.63±11.07	303.55±24.70	21.33
22	95.69±4.51	266.03±13.32	462.30±22.01	167.18±12.13	363.73±22.21	21.80
24	106.61±11.03	342.06±29.20*	NA	230.39±36.65*	NA	NA
27	140.55±4.48	357.44±8.99	497.96±21.92	208.96±10.24	351.65±22.12	31.60
30	173.36±7.05	396.65±11.48	479.88±25.29	213.20±13.33	310.20±25.52	37.88
30$_{chronic}$	163.14±9.28	306.62±16.06	356.82±53.05	136.03±11.90	198.33±53.54	38.63

493   Table 4. Q_{10} values for AS, MMR (both methods) and RMR separated between each
494   temperature increment. * = the slightly overestimated adjusted MO_2 for the 24°C fish is
495   reflected in calculated Q_{10} values.
496

	12-17°C	17-22°C	22-24°C	22-27°C	24-27°C	27-30°C	27-30$_C$°C
AS_{chase}	1.49	1.35	4.97*	1.56	0.72*	1.07	0.24
AS_{flume}	1.56	1.44	NA	0.93	NA	0.66	0.15
MMR_{chase}	1.62	1.53	3.51*	1.81	1.16*	1.41	0.60
MMR_{flume}	1.66	1.58	NA	1.16	NA	0.88	0.33
RMR	1.96	2.15	1.72	2.16	2.51	2.01	1.64

497
498

    Figures
500
501   *Figure Captions*
502
503   **Figure 1. Temperature and body weight both affect resting metabolic rate in black sea**
504   **bass.** RMR (n=121) for each temperature treatment is plotted against body weight (g). A fitted
505   regression line demonstrates that in addition to the effect of temperature on RMR, body weight
506   also has an effect ($P<0.05$). 30c = $30_{chronic}°C$ treatment.
507
Figure 2A and 2B. Effect of temperature on resting metabolic rate and maximum
metabolic rate measured with a chase and a flume method. MMR (solid circles) and RMR
(open circles) presented as mean ± s.e. normalized to a mean weight of 350g for each
temperature treatment for chase method MMR (A) and flume method MMR (B). RMR is slightly
different between (A) and (B) based on which fish were used for the respective MMR method.
The $30_{chronic}°C$ group is denoted by triangles. Tukey *post hoc* significance between treatments is
shown by letters where data points with different letters indicate a significant difference
($P<0.05$). **2A**: n=9 for $30_{chronic}°C$; n=16 for 12, 17, and 22°C; n=17 for 24 and 30°C; n=30 for
27°C. **2B**: n=4 for $30_{chronic}°C$; n=10 for 12 and 30°C; n=12 for 17, 22, and 27°C.
517
Figure 3a and 3b. Effect of temperature on black sea bass aerobic scope. Aerobic scope
(mean ± s.e.) of black sea bass normalized around a mean weight of 350g at each temperature
treatment with the 30°C chronic group denoted by the black triangle. Letters indicate Tukey *post*
hoc significance between groups where data points sharing a letter are not significantly different
($P<0.05$). Aerobic scope curves were generated from a) the chase MMR treatment ($y = 180.17 +$
$89.15x – 15.40x^2 – 21.55x^3; R^2 = 0.878$) and b) flume MMR treatment ($y = 314.36 + 63.29x – 68.26x^2 –$
$19.65x^3; R^2 = 0.994$). **3A**: n=9 for $30_{chronic}°C$; n=16 for 12, 17, and 22°C; n=17 for 24 and 30°C;
n=30 for 27°C. **3B**: n=4 for $30_{chronic}°C$; n=10 for 12 and 30°C; n=12 for 17, 22, and 27°C.
526
527   **Figure 4. S_{crit} increases with increasing temperature.** S_{crit} presented as $\%O_2$ air saturation for
528   each temperature treatment. $30_{chronic}°C$ treatment is denoted by a triangle and there is no
529   significant difference between the $30_{chronic}°C$ and acute 30°C treatments. A linear-regression was
530   fitted for these data points ($R_2 = 0.793, P<0.001$) showing an increase in S_{crit} (e.g. a decrease in
531   hypoxia tolerance) with increasing temperature.
532
533   **Figure 5. S_{crit} dependence on resting metabolic rate.** S_{crit} is plotted against resting metabolic
534   rate measured during the hypoxia experiment. A linear-regression was fitted for these data points
535   ($R_2 = 0.823, P<0.001$) and shows an increase in S_{crit} as metabolic rates also rise.
536
537   **Figure 6. Factorial aerobic scope and metabolic index response to temperature.** Factorial
538   aerobic scope (FAS) and metabolic index (MI) plotted against temperature. Trends illustrate a
539   decreasing trend in both measures as temperature increases. Both FAS and MI are unitless
540   measures, but both measures scale similarly.

 Figures

543   Figure 1.

544
545
546   Figure 2.

556   Figure 3

557
558
559   Figure 4.

572
573   Figure 5
574

575
576
577   Figure 6

578
579

References

Belkin, I. M. 2009. Rapid warming of Large Marine Ecosystems. Progress in Oceanography, 81: 207–213.

Bell, R. J., Richardson, D. E., Hare, J. A., Lynch, P. D., and Fratantoni, P. S. 2015. Disentangling the effects of climate, abundance, and size on the distriubtion of marine fish: an example based on four stocks from the Northeast US shelf. ICES Journal of Marine Science, 72: 1311–1322.

Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V. 2018. Observed fingerprint of a weakening Atlantic Ocean overturning circulation. Nature, 556: 191–196.

Capossela, K. M., Brill, R. W., Fabrizio, M. C., and Bushnell, P. G. 2012. Metabolic and cardiorespiratory responses of summer flounder *Paralichthys dentatus* to hypoxia at two temperatures. Journal of Fish Biology, 81: 1043–1058.

Castelao, R., Glenn, S., Schofield, O., Chant, R., Wilkin, J., and Kohut, J. 2008. Seasonal evolution of hydrographic fields in the central Middle Atlantic Bight from glider observations. Geophysical Research Letters, 35: 6–11.

Chabot, D., McKenzie, D. J., and Craig, J. F. 2016a. Metabolic rate in fishes: definitions, methods and significance for conservation physiology. Journal of Fish Biology, 88: 1–9.

Chabot, D., Steffensen, J. F., and Farrell, A. P. 2016b. The determination of standard metabolic rate in fishes. Journal of Fish Biology, 88: 81–121.

Claireaux, G., Couturier, C., and Groison, A. 2006. Effect of temperature on maximum swimming speed and cost of transport in juvenile European sea bass (*Dicentrarchus labrax*). Journal of Experimental Biology, 209: 3420–3428.

Claireaux, G., and Chabot, D. 2016. Responses by fishes to environmental hypoxia: integration through Fry's concept of aerobic metabolic scope. Journal of Fish Biology, 88: 232–251.

Clark, T. D., Sandblom, E., and Jutfelt, F. 2013. Aerobic scope measurements of fishes in an era of climate change: respirometry, relevance and recommendations. Journal of Experimental Biology, 216: 2771–2782.

Clarke, A., and Johnston, N. 1999. Scaling of metabolic rate with body mass and temperature in teleost fish. Journal of Animal Ecology, 68: 893–905.

Collins, G. M., Clark, T. D., Rummer, J. L., and Carton, A. G. 2013. Hypoxia tolerance is conserved across genetically distinct sub-populations of an iconic, tropical Australian teleost (*Lates calcarifer*). Conservation Physiology, 1: doi:10.1093/conphys/cot29.

Del Toro-Silva, F. M., Miller, J. M., Taylor, J. C., and Ellis, T. A. 2008. Influence of oxygen and temperature on growth and metabolic performance of *Paralichthys lethostigma* (Pleuronectiformes: Paralichthyidae). Journal of Experimental Marine Biology and Ecology, 358: 113–123.

Deutsch, C., Ferrel, A., Seibel, B., Pörtner, H. O., and Huey, R. B. 2015. Climate change tightens a metabolic constraint on marine habitats. Science, 348: 1132–1136.

Ejbye-Ernst, R., Michaelsen, T. Y., Tirsgaard, B., Wilson, J. M., Jensen, L. F., Steffensen, J. F., Pertoldi, C., *et al.* 2016. Partitioning the metabolic scope: The importance of anaerobic metabolism and implications for the oxygen- and capacity-limited thermal tolerance (OCLTT) hypothesis. Conservation Physiology, 4: 1–13.

Fabrizio, M. C., Manderson, J. P., and Pessutti, J. P. 2013. Habitat associations and dispersal of black sea bass from a mid-Atlantic Bight reef. Marine Ecology Progress Series, 482: 241–253.

Farrell, A. P., Hinch, S. G., Cooke, S. J., Patterson, D. A., Crossin, G. T., Lapointe, M., and

Mathes, M. T. 2008. Pacific salmon in hot water: applying aerobic scope models and
biotelemetry to predict the success of spawning migrations. Physiological and Biochemical
Zoology, 81: 697–709.
Farrell, A. P. 2016. Pragmatic perspective on aerobic scope: peaking, plummeting, pejus and
apportioning. Journal of Fish Biology, 88: 322–343.
Friedland, K. D., and Hare, J. A. 2007. Long-term trends and regime shifts in sea surface
temperature on the continental shelf of the northeast United States. Continental Shelf
Research, 27: 2313–2328.
Glenn, S., Arnone, R., Bergmann, T., Bissett, W. P., Crowley, M., Cullen, J., Gryzmski, J., *et al.*
2004. Biogeochemical impact of summertime coastal upwelling on the New Jersey Shelf.
Journal of Geophysical Research C: Oceans, 109: 1–15.
Hare, J. A., Morrison, W. E., Nelson, M. W., Stachura, M. M., Teeters, E. J., Griffis, R. B.,
Alexander, M. A., *et al.* 2016. A vulnerability assessment of fish and invertebrates to
climate change on the northeast u.s. continental shelf. PLoS ONE, 11: e0146756.
Healy, T. M., and Schulte, P. M. 2012. Thermal acclimation is not necessary to maintain a wide
thermal breadth of aerobic scope in the common killifish (*Fundulus heteroclitus*).
Physiological and Biochemical Zoology, 85: 107–119.
Holt, R. E., and Jørgensen, C. 2015. Climate change in fish: effects of respiratory constraints on
optimal life history and behaviour. Biology Letters, 11: 20141032.
Houghton, R. W., Schlitz, R., Beardsley, R. C., Butman, B., and Chamberlin, J. L. 1982. The
Middle Atlantic Bight Cold Pool: Evolution of the Temperature Structure During Summer
1979. Journal of Physical Oceanography, 12: 1019–1029.
Kavanaugh, M. T., Rheuban, J. E., Luis, K. M. A., and Doney, S. C. 2017. Thirty-three years of
ocean benthic warming along the U.S. Northeast Continental shelf and slope: patterns,
drivers, and ecological consequences. Journal of Geophysical Research: Oceans: 1–16.
Killen, S. S., Norin, T., and Halsey, L. G. 2017. Do method and species lifestyle affect measures
of maximum metabolic rate in fishes? Journal of Fish Biology, 90: 1037–1046.
Kleisner, K. M., Fogarty, M. J., McGee, S., Barnett, A., Fratantoni, P., Greene, J., Hare, J. A., *et*
al. 2016. The effects of sub-regional climate velocity on the distribution and spatial extent
of marine species assemblages. PLoS ONE, 11: 1–21.
Kleisner, K. M., Fogarty, M. J., McGee, S., Hare, J. A., Moret, S., Perretti, C. T., and Saba, V. S.
2017. Marine species distribution shifts on the U.S. Northeast Continental Shelf under
continued ocean warming. Progress in Oceanography, 153: 24–36.
Lapointe, D., Vogelbein, W. K., Fabrizio, M. C., Gauthier, D. T., and Brill, R. W. 2014.
Temperature, hypoxia, and mycobacteriosis: effects on adult striped bass *Morone saxatilis*
metabolic performance. Diseases of Aquatic Organisms, 108: 113–127.
Lefevre, S., McKenzie, D. J., and Nilsson, G. E. 2017. Models projecting the fate of fish
populations under climate change need to be based on valid physiological mechanisms.
Global Change Biology, 23: 3449–3459.
Mallekh, R., and Lagardère, J. P. 2002. Effect of temperature and dissolved oxygen
concentration on the metabolic rate of the turbot and the relationship between metabolic
scope and feeding demand. Journal of Fish Biology, 60: 1105–1115.
Manderson, J., Palamara, L., Kohut, J., and Oliver, M. J. 2011. Ocean observatory data is useful
for regional habitat modeling of species with different vertical habitat preferences. Marine
Ecology Progress Series, 438: 1–17.
McDonnell, L. H., and Chapman, L. J. 2015. At the edge of the thermal window: effects of

elevated temperature on the resting metabolism, hypoxia tolerance and upper critical thermal limit of a widespread African cichlid. Conservation Physiology, 3: 1–13.

Morley, J. W., Selden, R. L., Latour, R. J., Frölicher, T. L., Seagraves, R. J., and Pinsky, M. L. 2018. Projecting shifts in thermal habitat for 686 species on the North American continental shelf. PLoS ONE, 13: 1–28.

Moser, J., and Shepherd, G. R. 2008. Seasonal distribution and movement of black sea bass (*Centropristis striata*) in the Northwest Atlantic as determined from a mark-recapture experiment. Journal of Northwest Atlantic Fishery Science, 40: 17–28.

Musick, J. A., and Mercer, L. P. 1977. Seasonal distribution of Black Sea Bass, *Centropristis striata*, in the Mid-Atlantic Bight with comments on ecology and fisheries of the species. Trans. Amer. Fish. Soc., 106: 12–25.

NEFSC. 2017. The 62nd northeast regional stock assessment workshop (62nd SAW). Ref. Doc. 17-03, NEFSC, Woods Hole, MA.

Nilsson, G. E., and Renshaw, G. M. C. 2004. Hypoxic survival strategies in two fishes: extreme anoxia tolerance in the North European crucian carp and natural hypoxic preconditioning in a coral-reef shark. The Journal of Experimental Biology, 207: 3131–3139.

Norin, T., Malte, H., and Clark, T. D. 2014. Aerobic scope does not predict the performance of a tropical eurythermal fish at elevated temperatures. The Journal of Experimental Biology, 217: 244–251.

Norin, T., and Clark, T. D. 2016. Measurement and relevance of maximum metabolic rate in fishes. Journal of Fish Biology, 88: 122–151.

Nye, J. A., Link, J. S., Hare, J. A., and Overholtz, W. J. 2009. Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf. Marine Ecology Progress Series, 393: 111–129.

Pershing, A. J., Alexander, M. A., Hernandez, C. M., Kerr, L. A., Le Bris, A., Mills, K. E., Nye, J. A., *et al.* 2015. Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. Science, 350: 809–812.

Pinsky, M. L., Worm, B., Fogarty, M. J., Sarmiento, J. L., and Levin, S. A. 2013. Marine taxa track local climate velocities. Science, 341: 1239–1242.

Pinsky, M. L., Reygondeau, G., Caddell, R., Palacios-Abrantes, J., Spijkers, J., and Cheung, W. W. L. 2018. Preparing ocean governance for species on the move. Science, 360: 1189–1192.

Plambech, M., Van Deurs, M., Steffensen, J. F., Tirsgaard, B., and Behrens, J. W. 2013. Excess post-hypoxic oxygen consumption in Atlantic cod *Gadus morhua*. Journal of Fish Biology, 83: 396–403.

Pörtner, H. O., and Knust, R. 2007. Climate change affects marine fishes through the oxygen limitation of thermal tolerance. Science, 315: 95–97.

Pörtner, H. O., and Farrell, A. P. 2008. Physiology and climate change. Science, 322: 690–692.

Pörtner, H. O. 2010. Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. Journal of Experimental Biology, 213: 881–893.

Pörtner, H. O., and Peck, M. A. 2010. Climate change effects on fishes and fisheries: towards a cause-and-effect understanding. Journal of Fish Biology, 77: 1745–1779.

R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Raby, G. D., Casselman, M. T., Cooke, S. J., Hinch, S. G., Farrell, A. P., and Clark, T. D. 2016.

Aerobic scope increases throughout an ecologically relevant temperature range in coho
salmon. The Journal of Experimental Biology, 219: 1922–1931.
Reidy, S. P., Nelson, J. A., Tang, Y., and Kerr, S. R. 1995. Post-exercise metabolic rate in
Atlantic cod and its dependence upon the methos of exhaustion. Journal of Fish Biology,
47: 377–386.
Rodgers, G. G., Tenzing, P., and Clark, T. D. 2016. Experimental methods in aquatic
respirometry: the importance of mixing devices and accounting for background respiration.
Journal of Fish Biology, 88: 65–80.
Rogers, N. J., Urbina, M. A., Reardon, E. E., Mckenzie, D. J., and Wilson, R. W. 2016. A new
analysis of hypoxia tolerance in fishes using a database of critical oxygen level (P_{crit}).
Conservation Physiology, 4: cow012.
Roy, E. M., Quattro, J. M., and Greig, T. W. 2012. Genetic management of black sea bass:
influence of biogeographic barriers on population structure. Marine and Coastal Fisheries,
4: 391–402.
Saba, V. S., Griffies, S. M., Anderson, W. G., Winton, M., Alexander, M. A., Delworth, T. L.,
Hare, J. A., *et al.* 2016. Enhanced warming of the Northwest Atlantic Ocean under climate
change. Journal of Geophysical Research: Oceans, 120: 1–15.
Sandblom, E., Gräns, A., Axelsson, M., and Seth, H. 2014. Temperature acclimation rate of
aerobic scope and feeding metabolism in fishes: implications in a thermally extreme future.
Proceedings of Royal Society of Biology, 281: 20141490.
Schofield, O., Roarty, H., Saba, G. K., Xu, Y., Kohut, J., Glenn, S., Manderson, J., *et al.* 2012.
Phytoplankton dynamics and bottom water oxygen during a large bloom in the summer of
2011. Oceans, 2012, IEEE: 1–6.
Schulte, P. M. 2015. The effects of temperature on aerobic metabolism: towards a mechanistic
understanding of the responses of ectotherms to a changing environment. The Journal of
Experimental Biology, 218: 1856–1866.
Schurmann, H., and Steffensen, J. F. 1997. Effects of temperature, hypoxia and activity on the
metabolism of juvenile Atlantic cod. Journal of Fish Biology, 50: 1166–1180.
Seibel, B. 2011. Critical oxygen levels and metabolic supression in oceanic oxygen minimum
zones. The Journal of Experimental Biology, 214: 326-336.
Steimle, F., Zetlin, C., Berrien, P., and Chang, S. 1999. Essential fish habitat source document:
Black sea bass, *Centropristis striata*, life history and habitat characteristics. NOAA Tech
Memo NMFS NE 143: 42.
Svendsen, J. C., Tudorache, C., Jordan, A. D., Steffensen, J. F., Aarestrup, K., and Domenici, P.
2010. Partition of aerobic and anaerobic swimming costs related to gait transitions in a
labriform swimmer. Journal of Experimental Biology, 213: 2177–2183.
Svendsen, M. B. S., Bushnell, P. G., and Steffensen, J. F. 2016a. Design and setup of
intermittent-flow respirometry system for aquatic organisms. Journal of Fish Biology, 88:
26–50.
Svendsen, M. B. S., Bushnell, P. G., Christensen, E. A. F., and Steffensen, J. F. 2016b. Sources
of variation in oxygen consumption of aquatic animals demonstrated by simulated constant
oxygen consumption and respirometers of different sizes. Journal of Fish Biology, 88: 51–
64.
Verberk, W. C. E. P., Bartolini, F., Marshall, D. J., Pörtner, H. O., Terblanche, J. S., White, C.
R., and Giomi, F. 2016. Can respiratory physiology predict thermal niches? Annals of the
New York Academy of Sciences, 1365: 73–88.

Wikelski, M., and Cooke, S. J. 2006. Conservation physiology. Trends in Ecology and
Evolution, 21: 38–46.
Wilkin, J. L., and Hunter, E. J. 2013. An assessment of the skill of real-time models of Mid-
Atlantic Bight continental shelf circulation. Journal of Geophysical Research: Oceans, 118:
2919–2933.
Wishner, K. F., Seibel, B. A., Roman, C., Deutsch, C., Outram, D., Shaw, C. T., Birk, M. A., *et*
al. 2018. Ocean deoxygenation and zooplankton: very small oxygen differences matter.
Science Advances, 4: eaau5180.
Yamanaka, H., Kohmatsu, Y., and Yuma, M. 2007. Difference in the hypoxia tolerance of the
round crucian carp and largemouth bass: Implications for physiological refugia in the
macrophyte zone. Ichthyological Research, 54: 308–312.
Yeager, G. R., and Ultsch, D. P. 1989. Physiological regulation and conformation : A BASIC
program for the determination of critical points. Physiological Zoology, 62: 888–907.