1 TIGAR: An Improved Bayesian Tool for Transcriptomic Data Imputation Enhances Gene
2 Mapping of Complex Traits
3

4 Sini Nagpal[1], Xiaoran Meng[2,3], Michael P. Epstein[2,3], Lam C. Tsoi[4], Matthew Patrick[5], Greg
5 Gibson[1], Philip L. De Jager[6], David A. Bennett[7], Aliza P. Wingo[8,9], Thomas S. Wingo[3,10], Jingjing
6 Yang[3*]
7
8   1. School of Biology, Georgia Institute of Technology, Atlanta, GA, 30332, USA
9   2. Department of Biostatistics and Bioinformatics, Emory University School of Public Health,
10   Atlanta, GA, 30322, USA
11   3. Center for Computational and Quantitative Genetics, Department of Human Genetics, Emory
12   University School of Medicine, Atlanta, GA, 30322, USA
13   4. Department of Dermatology; Department of Computational Medicine & Bioinformatics;
14   Department of Biostatistics, University of Michigan, Ann Arbor, MI, 48109, USA
15   5. Department of Dermatology, University of Michigan Medical School, Ann Arbor, MI, 48109,
16   USA
17   6. Medical Center Neurological Institute, Columbia University, New York, NY, 10032,
18   USA
19   7. Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, 60612, USA
20   8. Division of Mental Health, Atlanta VA Medical Center, Decatur, GA, USA
21   9. Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine,
22   Atlanta, GA, 30322, USA
23   10. Department of Neurology, Emory University School of Medicine, Atlanta, GA, 30322, USA

24   *Correspondence author: jyang51@emory.edu
25
26

Abstract

The transcriptome-wide association studies (TWAS) that test for association between the study

trait and the imputed gene expression levels from cis-acting expression quantitative trait loci

(cis-eQTL) genotypes have successfully enhanced the discovery of genetic risk loci for complex

traits. By using the gene expression imputation models fitted from reference datasets that have

both genetic and transcriptomic data, TWAS facilitates gene-based tests with GWAS data while

accounting for the reference transcriptomic data. The existing TWAS tools like PrediXcan and

FUSION use parametric imputation models that have limitations for modeling the complex

genetic architecture of transcriptomic data. Therefore, we propose an improved Bayesian

method that assumes a data-driven nonparametric prior to impute gene expression. Our method

is general and flexible and includes both the parametric imputation models used by PrediXcan

and FUSION as special cases. Our simulation studies showed that the nonparametric Bayesian

model improved both imputation R^2 for transcriptomic data and the TWAS power over

PrediXcan. In real applications, our nonparametric Bayesian method fitted transcriptomic

imputation models for 2X number of genes with 1.7X average regression R^2 over PrediXcan,

thus improving the power of follow-up TWAS. Hence, the nonparametric Bayesian model is

preferred for modeling the complex genetic architecture of transcriptomes and is expected to

enhance transcriptome-integrated genetic association studies. We implement our Bayesian

approach in a convenient software tool "**TIGAR**" (**T**ranscriptome-**I**ntegrated **G**enetic **A**ssociation

Resource), which imputes transcriptomic data and performs subsequent TWAS using individual-

level or summary-level GWAS data.

Introduction

Genome-wide association studies (GWAS) have successfully identified thousands of

genetic risk loci for complex traits. However, the majority of these loci are located within

noncoding regions whose molecular mechanisms remain unknown[1-3]. Recent studies have

shown that these associated regions were enriched for regulatory elements such as enhancers

(H3K27ac marks)[4; 5] and expression of quantitative trait loci (eQTL)[6; 7], suggesting that the

genetically regulated gene expression might play a key role in the biological mechanisms of

complex traits. Multiple studies have recently generated rich transcriptomic datasets for diverse

tissues of the human body, e.g., the Genotype-Tissue Expression (GTEx) project for 44 human

tissues[6], Genetic European Variation in Health and Disease (GEUVADIS) for lymphoblastoid

cell lines[8], Depression Genes and Networks (DGN) for whole-blood samples[9], and the North

American Brain Expression Consortium (NABEC) for cortex tissues[10]. Previous studies[11-16] have

also shown that integrating transcriptomic data in GWAS can help identify novel functional loci.

The majority of GWAS projects do not possess transcriptomic data and thus cannot

directly conduct integrative analysis. Existing studies[11; 12] have shown that one can impute the

genetically regulated gene expression (GReX) within such GWAS projects by using reference

datasets like GTEx[6] and GEUVADIS[8], and then test for association between imputed GReX for

GWAS samples and the trait of interest — referred to as transcriptome-wide association studies

(TWAS)[11; 12]. Specifically, the gene expression imputation models are fitted by regressing

assayed gene-expression levels on cis-eQTL genotypes with reference dataset. For examples,

the PrediXcan[11] method uses an Elastic-Net[17] variable selection model on reference dataset to

estimate the cis-eQTL effect-sizes that are used to impute the GReX for GWAS samples, while

the FUSION[12] tool implements a Bayesian sparse linear mixed model (BSLMM)[18] to estimate

the cis-eQTL effect-sizes.

Basically, the Elastic-Net[17] model used by PrediXcan[11] assumes a combination of

LASSO[19] (L_1) and Ridge[20] (L_2) penalties on the cis-eQTL effect-sizes, which is equivalent to a

Bayesian model with a mixture Gaussian and Laplace prior[21]. Whereas, the BSLMM[18] used by

FUSION[12] is a combination of Bayesian variable selection model (BVSR)[22] and linear mixed

model (LMM)[23] by assuming a normal mixture prior. However, a parametric prior is assumed for

the cis-eQTL effect-sizes by both Elastic-Net and BSLMM, which restricts the capability of

PrediXcan and FUSION for handling the underlying complex genetic architecture of

transcriptomes. Existing studies[11; 12] have also shown that both PrediXcan[11] and FUSION[12]

estimated the average regression R^2 (i.e., the percentage of gene expression variation that can

be explained by cis- genotypes) as ~5% for human whole blood transcriptome, while the

average genome-wide heritability of gene expression in human whole blood transcriptome is

estimated to be more than double that quantity[24; 25]. As shown in the result section of this paper,

similar underestimation of regression R^2 by the Elastic-Net model (used by PrediXcan[11]) was

observed in both simulated and real datasets.

Therefore, to flexibly model cis-eQTL distributions, we propose to use a nonparametric

Bayesian method, where the prior for effect-sizes is nonparametric and can be estimated from

the data by assuming a Dirichlet process prior on effect-size variance. This Bayesian model is

also known as latent Dirichlet process regression (DPR) model[26], which can flexibly model the

underlying complex genetic architecture of transcriptomes. Thus, DPR is a more generalized

model that includes Elastic-Net (implemented in PrediXcan[11]) and BSLMM (implemented in

FUSION[12]) as special cases. Consequently, DPR can robustly estimate cis-eQTLs and then

improve the estimation of imputation R^2 (the squared correlation between the observed and

imputed values on test samples). Moreover, we will use a variational Bayesian algorithm[26-28] as

an alternative of Monte Carlo Markov Chain (MCMC)[29] to efficiently fit the Bayesian model.

Similar to PrediXcan[11] and FUSION[12] methods, we can use DPR to estimate cis-eQTLs

effect-sizes from a reference dataset, which can then be used for downstream TWAS using

either individual-level or summary-level GWAS data. In subsequent sections, we first describe

the DPR approach for estimating cis-eQTL effect-sizes from a reference dataset and how we

can then use these effect-sizes for downstream TWAS in a target GWAS using either individual-

level or summary-level data. We then compare the performance of DPR with PrediXcan using

both simulated data and real SNP and transcriptome data from the Religious Orders Study and

Rush Memory Aging Project (ROS/MAP)[30-33] for studying Alzheimer's Disease (AD).

Our in-depth simulation studies demonstrated that the DPR method obtained higher

imputation R^2 on test samples, when the true expression heritability is <10% or when >1%

variants are true causal cis-eQTL. Consequently, better imputation R^2 resulted in improved

power for follow-up association studies. Meanwhile, application of DPR to the ROS/MAP study

imputed GReX for twice the number of genes as PrediXcan with 70% higher regression R^2 on

average. Using DPR, we also found a potentially associated gene *TRAPPC6A* for AD pathology

indices that was missed by PrediXcan. Further, by using the transcriptomic imputation models

fitted from ROS/MAP data and summary-level GWAS data generated from the International

Genomics of Alzheimer's Project (IGAP)[34], we identified 7 out of 26 known AD loci[34-38] that

potentially affect the late-onsite AD risk through transcript abundance. We conclude with a

discussion of future topics and further describe our novel software tool TIGAR (**T**ranscriptome-

Integrated **G**enetic **A**ssociation **R**esource) implementing our nonparametric Bayesian method

for public use.

<u>Materials and Methods</u>

We consider the following additive linear regression model for estimating the cis-eQTL

effect-sizes from a reference study that has both genetic and transcriptomic data available,

$$E_g = Xw + \epsilon, \quad \epsilon \sim N(0, \sigma_\epsilon^2 I); \qquad \textit{Equation 1}$$

where E_g denotes the gene expression levels (after corrections for confounding

covariates such as age, sex, and principal components (PCs)) for gene g, X denotes the

genotype matrix for all cis- genotypes (encoded as the number of minor alleles), w denotes the

corresponding cis-eQTL effect-size vector, and ϵ denotes the error term. The intercept term is

dropped in *Equation 1* for assuming both E_g and X are centered at 0. Generally, SNPs within

1MB of the flanking 5' and 3' ends (cis-SNPs) are included in this regression model and non-

zero $\hat{w}$ will be used for follow-up analysis. In particular, the GReX will be imputed by

$$\widehat{GReX} = X_{new}\hat{w},$$

with cis-SNP data X_{new} for GWAS samples.

<u>Nonparametric Bayesian Method</u>

Following the latent Dirichlet process regression (DPR) model[26], we assume a normal

prior $N(0, \sigma_w^2)$ for the cis-eQTL effect-sizes $(w_i, i = 1, \dots, p)$ and a Dirichlet process (DP) prior[39]

for the effect-size variance σ_w^2 (as in *Equation 1*):

$$w_i \sim N(0, \sigma_w^2), \ \sigma_w^2 \sim D, \ D \sim DP\ (IG(a, b), \xi). \quad \textit{Equation 2}$$

The prior distribution D deviates from the DP with base distribution as an Inverse Gamma (IG)

distribution and concentration parameter ξ. Note that σ_w^2 can be viewed as a latent variable and

integrating out σ_w^2 will induce a nonparametric prior distribution on w_i, which is equivalent to a

DP normal mixture model[26-28],

$\quad w_i \sim \sum_{k=0}^{+\infty} \pi_k N(0, \sigma_k^2), \ \sigma_k^2 \sim IG(a_k, b_k), \ \pi_k = v_k \prod_{l=0}^{k-1}(1 - v_l), \ v_k \sim Beta(1, \ \xi).$ *Equation 3*

Here, the nonparametric prior distribution on w_i is equivalently represented by a mixture normal

prior that is a weighted sum of an infinitely number of normal distributions $\left(N(0, \sigma_k^2), k = \right.$

$\left. 0, \dots, +\infty\right)$, corresponding weight π_k is determined by $(v_l, l = 0, \dots, k)$ with a Beta prior, and ξ in

the Beta prior (the same concentration parameter as in Equation 2) determines the number of

components with non-zero weights in the mixture normal prior. Conjugate hyper priors

$\xi \sim Gamma(a_\xi, b_\xi)$ and $\sigma_\epsilon^2 \sim IG(a_\epsilon, b_\epsilon)$ are assumed.

Generally, the hyper parameters $a_k, b_k, a_\epsilon, b_\epsilon$ in the *Inverse Gamma* distributions can be

set as 0.1 and (a_ξ, b_ξ) in the *Gamma* distribution can be set as (1, 0.1) to induce non-informative

priors for $(\sigma_k^2, \sigma_\epsilon^2, \xi)$. That is, the parameters $(\sigma_k^2, \sigma_\epsilon^2, \xi)$ will be adaptively estimated from the

data and the nonparametric prior on w_i will be data-driven. The posterior estimates for $\boldsymbol{w}$ can be

obtained by the MCMC[29] or variational Bayesian algorithm[28; 40], from the following joint

conditional posterior distribution

$$P\left(\boldsymbol{w}, \boldsymbol{\pi}, \boldsymbol{v}, \xi, \sigma_\epsilon^2 | \boldsymbol{E_g}, \boldsymbol{X}\right) \propto$$

$$P\left(\boldsymbol{E_g} | \boldsymbol{w}, \boldsymbol{X}, \sigma_\epsilon^2\right) P\left(\boldsymbol{w} | \boldsymbol{\pi}, \sigma_1^2, \dots, \sigma_k^2, \dots\right) \left(\prod_{k=0}^{+\infty} P\left(\sigma_k^2 | a_k, b_k\right)\right) P(\boldsymbol{\pi} | \boldsymbol{v}) P(\boldsymbol{v} | \xi) P\left(\xi | a_\xi, b_\xi\right) P(\sigma_\epsilon^2 | a_\epsilon, b_\epsilon).$$

Particularly, the variational Bayesian algorithm[28; 40] is an approximation for the MCMC[29] with

greatly improved computational efficiency, which is also used in our tool. Please refer to the

supplementary note for technical details of both MCMC sampling and variational inference

algorithms for obtaining the Bayesian posterior estimates for the cis-eQTL effect-sizes.

Elastic-Net and BSLMM Methods

The Elastic-Net model[17] (used by PrediXcan[11]) estimates the cis-eQTL effect-sizes $\hat{w}$ in

Equation 1 with a combination of L_1 (LASSO)[19] and L_2 (Ridge)[20] penalties by

$$\hat{w} = \underset{w}{\mathrm{argmin}} \left(\left\| E_g - Xw \right\|_2^2 + \lambda \left(\alpha \|w\|_1 + \frac{1}{2}(1-\alpha)\|w\|_2^2 \right) \right),$$

where $\|\cdot\|_2$ denotes L_2 norm, $\|\cdot\|_1$ denotes L_1 norm, $\alpha \in [0,1]$ denotes the proportion of L_1

penalty, and λ denotes the penalty parameter. Particularly, PrediXcan[11] takes $\alpha = 0.5$ and tunes

the penalty parameter λ by a 5-fold cross validation.

As pointed out by previous studies[17; 21], the Elastic-Net model is equivalent to a Bayesian

model with a mixture Gaussian and Laplace (mixture normal) prior for w, that is, $p(w) \propto$

$\exp\left(-\lambda \left(\alpha\|w\|_1 + \frac{1}{2}(1-\alpha)\|w\|_2^2 \right) \right)$. Whereas, the BSLMM[18] assumes a mixture of two normal

as the prior for cis-eQTL effect-sizes, $w_i \sim \pi N(0, (\sigma_1^2 + \sigma_2^2)) + (1-\pi)N(0, \sigma_2^2)$. That is, the

BSLMM[18] assumes all cis-SNPs have at least a small effect, which are normally distributed with

variance σ_2^2, and some proportion (π) of cis-SNPs have an additional effect, normally distributed

with variance σ_1^2. Particularly, with $\sigma_2^2 = 0$, BSLMM becomes BVSR[22], and with $\pi = 0$, the

BSLMM becomes the LMM[23]. Therefore, the DP normal mixture[26-28] as assumed by the DPR

method includes the parametric (mixture normal) priors for Bayesian Elastic-Net[21] and BSLMM[18]

as special cases, which is the main reason why DPR is a more generalized model including

Elastic-Net and BSLMM as special cases. Thus, we believe the DPR method can robustly

model complex genetic architecture and then improve the imputation R^2.

<u>Association Study with Univariate Phenotype</u>

Given individual-level GWAS data (genotype data X_{new}, phenotype Y, and covariant

matrix C) and cis-eQTL effect-size estimates $\hat{w}$, the follow-up TWAS (using a burden type gene-

based test[41]) is to test for association between $\widehat{GReX} = X_{new}\hat{w}$ and Y based on the following

generalized linear regression model

183
$$f(E[Y|X, C]) = \eta C + \beta \widehat{GReX} .$$ *Equation 4*

Here, $f(\cdot)$ is a pre-specified link function, which can be set as identify function for quantitative

phenotype or set as logit function for dichotomous phenotype. The gene-based association test

is equivalent to test $H_0: \beta = 0$ in Equation 4.

If only summary-level data are available, we take the same approach as implemented by

the FUSION[12] method. Let Z denote the vector of Z-scores by single variant tests (Wald,

likelihood ratio, score tests, etc.) for all cis-SNPs. The burden Z-score for gene-based

association test is defined as

191
$$\tilde{Z} = \frac{Z\widehat{w}}{\sqrt{Z\widehat{w}}} = \frac{Z\widehat{w}}{\sqrt{\widehat{w}'V\widehat{w}}} ,$$ *Equation 5*

where V denotes the covariance matrix across study SNPs that can be estimated from training

data or reference panels such as 1000 Genome Project[42].

<u>Association Study with Multivariate Phenotype</u>

To test for association between multivariate phenotypes and imputed GReX of the focal

gene, we take a similar approach as the MultiPhen method[43] using individual-level data. For

example, consider two phenotypes (Y_1, Y_2) and a covariate matrix C, we first adjust for the

covariates by taking the residuals $(\widetilde{Y_1}, \widetilde{Y_2})$ respectively from the linear regression models

$Y_j = \eta C + \epsilon, \; j = 1,2$. Then we test if the regression R^2 is significantly greater than zero

$(H_0: R^2 = 0)$ for the following regression model

202
$$\widehat{GReX}_g = \beta_1 \widetilde{Y_1} + \beta_2 \widetilde{Y_2} + \epsilon .$$ *Equation 6*

That is, we test if the multivariate phenotypes can jointly explain a non-zero percentage of

variance in the imputed GReX. The p-value can be calculated by using the F-statistic for the

regression R^2 in *Equation 6.*

Even when only summary-level GWAS data are available, we can first obtain a burden

Z-score per phenotype from *Equation 5*, i.e., $\tilde{\mathbf{Z}} = (\widetilde{Z_1}, \widetilde{Z_2})$ with two phenotypes. Then a similar

burden approach can be used to obtain a joint Z-score for multi-phenotype test,

$$\tilde{Z}_{joint} = \frac{\tilde{\mathbf{Z}}\mathbf{J}}{\sqrt{\tilde{\mathbf{Z}}\mathbf{J}}} = \frac{\tilde{\mathbf{Z}}\mathbf{J}}{\sqrt{\mathbf{J}'\mathbf{V}_Y\mathbf{J}}}, \quad \mathbf{J} = (1, \ldots, 1)',$$

where $\mathbf{V}_Y$ is the covariance matrix among study traits.

<u>Simulation Study Design</u>

We conducted in-depth simulation studies to compare the performance of both

PrediXcan and DPR methods with respect to imputation R^2 in the test data and the power of

association studies. Specifically, we used 499 ROS/MAP participants[44] with both RNA-

sequencing and genotype data as training data, and genotype data from an additional 1,200

ROS/MAP participants[44] as test data. The test sample size (1,200) was chosen arbitrarily

(randomly selected from the ROS/MAP study) to be comparable with the sample size (1,164) in

the real association study of AD pathology indices. The genotyped and imputed genetic data for

2,799 cis-SNPs (with minor allele frequency (MAF) > 5% and Hardy-Weinberg p-value $> 10^{-5}$)

of gene *ABCA7* were used to simulate gene expression levels.

We performed comprehensive scenarios that varied the proportion of causal SNPs (out

of 2,799 SNPs, influenced gene expression) among values in the vector

$p_{causal} = (1\%, 5\%, 10\%, 20\%)$. We varied the proportion of gene expression variance explained

by causal SNPs (i.e., expression heritability), along with the proportion of phenotypic variance

explained by simulated gene expression levels (i.e., phenotypic heritability), among values in

the vector $\left(h_e^2, h_p^2\right) = ((5\%, 80\%), (10\%, 50\%), (20\%, 25\%), (50\%, 10\%))$. The phenotypic

heritability was selected arbitrarily with respect to expression heritability such that the follow-up

association study power fell within the range of $(25\%, 85\%)$. With genotype matrix $\mathbf{X}_g$ of the

randomly selected causal SNPs (according to p_{causal}), we generated effect-sizes w_i from

$N(0,1)$ and then re-scaled the effect-sizes to ensure the targeted h_e^2. Gene expression levels

were generated by $E_g = X_g w + \epsilon$, with $\epsilon \sim N(0,(1-h_e^2))$. Then the phenotype values were

generated by $Y = \beta E_g + \epsilon$, where β was selected with respect to h_p^2 and $\epsilon \sim N(0,(1-h_p^2))$.

For each scenario, we repeated simulations for 1,000 times, where we applied both

PrediXcan[11] and DPR methods to obtain imputation models with training samples, impute the

GReX for test samples, and then conduct follow-up association studies using the imputed GReX.

We did not compare with FUSION[12] using BSLMM because of the computational burden of

estimating cis-eQTL effect-sizes by MCMC (~2 hours per gene). The association study power

was calculated as the proportion of 1,000 repeated simulations with p-value $< 2.5 \times 10^{-6}$

(genome-wide significance threshold adjusting for testing 20K independent genes).

<u>ROS/MAP Data</u>

Samples in the ROS/MAP data were collected from participants of the Religious Orders

Study (ROS) and the Rush Memory and Aging Project (MAP), which are prospective cohort

studies of studying aging and dementia[30; 31; 33]. The ROS/MAP study recruited senior adults

without known dementia at enrollment, who underwent annual clinical evaluation. Brain autopsy

was done at the time of death for each participant. All participants signed an informed consent

and Anatomic Gift Act, and the studies were approved by the Institutional Review Board of Rush

University Medical Center, Chicago, IL. Specifically, microarray genotype data generated for

2,093 European-decent participants[44], were further imputed to the 1000 Genome Project Phase

$3^{[42]}$ in our analysis. The post-mortem brain samples (gray matter of the dorsolateral prefrontal

cortex) from ~30% these participants were profiled for transcriptomic data by next-generation

RNA-seqencing[45]. In this paper, we conducted gene-based association studies for two important

indices of AD pathology that were quantified with β-antibody specific immunostains[30; 31; 33]:

neurofibrillary tangle density (tangles) with stereology and β-amyloid load (amyloid) with image

analysis. The neurofibrillary tangle density quantifies the average Tau tangle density within two

or more 20μm sections from eight brain regions — hippocampus, entorhinal cortex, midfrontal

cortex, inferior temporal, angular gyrus, calcarine cortex, anterior cingulate cortex, and superior

frontal cortex. The β-amyloid load quantifies the average percent area of cortex occupied by β-

amyloid protein in adjacent sections from the same eight brain regions.

Results

Simulation Studies

In the simulation studies, we observed that the DPR method performed robustly with

respect to different causal proportions and gene expression heritability. Especially, when

$p_{causal} > 1\%$, DPR out-performed PrediXcan across all expression heritability values, giving

higher imputation R^2 in test data (Figure 1). For example, when $p_{causal} = 20\%$, the average

imputation R^2 of 1,000 simulations was estimated as 4.54% by using DPR versus 2.64% by

using PrediXcan with $h_e^2 = 10\%$, while the average imputation R^2 was estimated as 12.02% by

using DPR versus 9.13% by using PrediXcan with $h_e^2 = 20\%$ (Table 1). When $p_{causal} = 1\%$,

DPR performed similarly as PrediXcan with $h_e^2 = 20\%, 50\%$ but slightly out-performed

PrediXcan with $h_e^2 = 5\%, 10\%$ (Table 1, Figure 1).

Consequently, when $p_{causal} > 1\%$ and $h_e^2 < 50\%$, or $p_{causal} = 1\%$ and $h_e^2 < 20\%$, the

power of association studies was higher by using DPR versus using PrediXcan imputation

models (Figure 2). When $h_e^2 = 50\%$, using both imputation models led to comparable power for

association studies (Figure 2). Interestingly, even when both methods had similar training

regression R^2 (Figure S1), the DPR method still resulted in higher imputation R^2 for test data

(Figures 1; Table 1) and higher power for association studies (Figure 2). Overall, these results

demonstrated the advantages of the DPR method for modeling the complex genetic architecture

of transcriptomes, especially when the causal proportions $> 1\%$ or the expression heritability

$< 10\%$.

<u>Real Applications to ROS/MAP Data</u>

To illustrate the performance of the DPR method in real studies, we applied both DPR

and PrediXcan on the ROS/MAP data (see Methods). We trained the gene expression

imputation models using 499 samples that have both transcriptomic data for frontal cortex

tissues and genotype data (imputed to 1000 Genome Phase 3, with MAF > 5%, Hardy-

Weinberg p-value $> 10^{-5}$, and imputation R^2 > 0.3). A total of 15,583 genes had gene

expression levels after standard RNA-sequencing quality control. The gene expression levels

were first adjusted for age at death, sex, postmortem interval, study (ROS or MAP), batch

effects, RNA integrity number scores, and cell type proportions (with respect to

oligodendrocytes, astrocytes, microglia, neurons) by linear regression models. For each gene,

cis- SNPs within the 1MB of the flanking 5' and 3' ends were used in the imputation models as

predictors.

In Figure 3, we compared transcriptome-wide regression R^2 estimated by using both

DPR and PrediXcan methods. Specifically, DPR fitted imputation models for 14,214 genes with

non-zero regression R^2 (average 10.3%), versus 6,549 genes with non-zero regression R^2

(average 6.05%) by PrediXcan. That is, DPR fitted imputation models for 2X genes with 1.7X

average regression R^2 over PrediXcan. Thus, the DPR method featuring data-driven

nonparametric prior for the cis-eQTL is indeed preferred in real studies.

Next, we imputed the GReX for all GWAS samples using the trained transcriptomic

imputation models by both DPR and PrediXcan, and then conducted gene-based association

studies (Methods) using all GWAS samples (N = 1,164) that have the AD pathology indices

(neurofibrillary tangle density and β-amyloid load, with Pearson correlation 0.48) quantified.

Possible confounding covariates including age at death, sex, study (ROS or MAP), smoking,

education, and first 3 genotype principle components were adjusted in the association studies.

Interestingly, the association studies for both AD pathology indices using the DPR imputation

models identified the same top significant gene *TRAPPC6A* (within the 2MB region from the

major risk gene *APOE*, encoding apolipoprotein E, but independent of *APOE*) with p-

values 1.64×10^{-5} and 5.35×10^{-5} (Figures S2 and S3). Moreover, the multivariate phenotype

association studies (Methods) for both AD pathology indices also identified *TRAPPC6A* as the

most significant gene with p-value 5.81×10^{-6} and FDR 0.08 (Figure 4). On the other hand, the

PrediXcan failed to obtain a transcriptomic imputation model for *TRAPPC6A* (Figures S4–S6).

In addition, for 26 known common and rare loci of late-onset AD[34-38], we conducted

association studies using transcriptomic imputation models (DPR and PrediXcan) fitted from

ROS/MAP data and standard GWAS results from IGAP[34]. We identified 7 significant loci (FDR

< 0.05; Figure 5; Figure S7) — *ABCA1*, *ADAM10, CD2AP, HLA_DRB5, MAPT, MS4A6A*, and

TREM2 — that potentially affect late-onsite AD risk through transcript abundance. Specifically,

5 out of 7 significant loci were uniquely identified by using the DPR imputation models, 1 locus

was uniquely identified by using the PrediXcan imputation model, while 1 locus was identified by

using both imputation models (Figure 5). From the individual-level association studies with both

AD pathology indices (ROS/MAP data) for these 26 loci (Figure S8), we see that the PrediXcan

method only imputed GReX for 9 out of 26 loci with average regression $R^2 = 10.3\%$, while the

DPR method imputed GReX for all 26 genes with average regression $R^2 = 16.7\%$. In summary,

these results show that the DPR method has superior power for follow-up TWAS.

Discussion

In this paper, we propose an improved nonparametric Bayesian method for

transcriptomic data imputation, accompanied with an integrated tool (freely available on

GITHUB) that is referred as **T**ranscriptome-**I**ntegrated **G**enetic **A**ssociation **R**esource (**TIGAR**).

TIGAR integrates both nonparametric Bayesian (DPR) and Elastic-Net models as two options

for transcriptomic data imputation, along with TWAS options using individual-level and

summary-level GWAS data for univariate and multi-variate phenotypes. We show the

usefulness of the DPR method by both in-depth simulations and real applications using

individual-level ROS/MAP[30-33] and summary-level IGAP[34] data. We demonstrate that DPR is

preferred over PrediXcan because of the flexible nonparametric modeling of cis-eQTL effect-

sizes that results in improved imputation R^2 for gene expression levels and higher power for

TWAS.

With respect to user-friendly interface and computational efficiency, TIGAR can: (i) Take

standard input files such as genotype files in VCF and dosage formats, phenotype files in PED

format, and a combined file (tab delimited) for gene annotations and expression levels; (ii) Load

input data per gene by TABIX for memory efficiency; (iii) Filter SNPs by given MAF and Hardy-

Weinberg p-value thresholds; (iv) Provide options of training both Elastic-Net (using Python

scripts) and DPR (using the executable tool developed with C++[26]) imputation models with

unified output format; (v) Implement multi-threaded computation to take full advantage of multi-

core clusters. These features make TIGAR a preferred tool for saving tedious data preparation

and computation time. For example, TIGAR can complete training imputation models for ~20K

genes and ~1K samples within ~20 hours and TWAS within ~1hour with a 2.4GHz 16-core CPU.

The computation time is expected to increase linearly with respect to the sample size.

However, as observed in the real ROS/MAP studies, there remains a large gap between

the regression R^2 estimated by DPR using cis-eQTL predictors (10.3%) and the average

genome-wide heritability of gene expression levels (21.8% estimated by GCTA[46] based on a

LMM). This is likely due to the large trans-acting contribution to transcript abundance

documented for most genes. Thus, we hypothesize that it should be possible to further improve

the regression R^2 and imputation R^2 by fitting transcriptomic imputation models with genome-

wide variants as predictors. To resolve the computational bottleneck of using genome-wide

variants, the current DPR model might be extended using a scalable Bayesian inference

technique such as an Expectation Maximization MCMC (EM-MCMC) algorithm[47]. To provide

more flexibility for the DPR model, the cis- and trans-acting genotype predictors could be

modeled as two different categories with different nonparametric DP priors. As a result, we

expect to further improve the prediction accuracy for GReX and then enhance the power of

follow-up TWAS.

Besides the follow-up gene-based association studies (i.e., TWAS) described in this

paper, the transcriptomic imputation models can be further extended by incorporating

environmental contributions. The imputed transcript abundance levels can then be used for

gene network analysis, differential gene expression analysis, and transcriptome mediation

analysis with GWAS data. Validation of transcriptomic prediction accuracy in independent

datasets will be critical in this regard, but unfortunately multiple large and similar datasets are

not yet generally available for tissues other than peripheral blood.

In conclusion, we expect our work will provide a convenient and improved tool for

transcriptomic imputation using the currently available rich reference datasets, as well as

enhanced gene mapping for better understanding the genetic etiology of complex traits.

Supplemental Data

Supplementary Text and Figures

Acknowledgments

J.Y. was supported by the startup funding from Emory University Department of Human

Genetics. A.P.W. and T.S.W. were supported by National Institutes of Health (NIH)

R01AG056533. M.P.E. was supported by NIH R01GM11796. L.C.T. was supported by the

Dermatology Foundation, the Arthritis National Research Foundation, the National

Psoriasis Foundation, and NIH K01AR072129. ROS/MAP study data were provided by the

Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL. Data collection

was supported through funding by NIA grants P30AG10161, R01AG15819, R01AG17917,

R01AG30146, R01AG36836, U01AG32984, U01AG46152, the Illinois Department of Public

Health, and the Translational Genomics Research Institute. In addition, we thank Thanneer

Perumal and Benjamin Logsdon for performing quality control of the ROS/MAP RNA-

sequencing data and for creating the brain cell type proportions.

Web Resources

TIGAR, https://github.com/xmeng34/TIGAR

PrediXcan, https://github.com/hakyim/PrediXcan

FUSION, http://gusevlab.org/projects/fusion/

ROS/MAP data, https://www.synapse.org/#!Synapse:syn3219045; http://www.radc.rush.edu/

IGAP data, http://web.pasteur-lille.fr/en/recherche/u744/igap/igap_download.php

References

1. Visscher, P.M., Brown, M.A., McCarthy, M.I., and Yang, J. (2012). Five years of GWAS
discovery. American journal of human genetics 90, 7-24.
2. McCarthy, M.I., Abecasis, G.R., Cardon, L.R., Goldstein, D.B., Little, J., Ioannidis, J.P., and
Hirschhorn, J.N. (2008). Genome-wide association studies for complex traits: consensus,
uncertainty and challenges. Nature reviews Genetics 9, 356-369.
3. Huang, Q. (2015). Genetic study of complex diseases in the post-GWAS era. J Genet
Genomics 42, 87-98.
4. Farh, K.K., Marson, A., Zhu, J., Kleinewietfeld, M., Housley, W.J., Beik, S., Shoresh, N., Whitton,
H., Ryan, R.J., Shishkin, A.A., et al. (2015). Genetic and epigenetic fine mapping of causal
autoimmune disease variants. Nature 518, 337-343.
5. Tsoi, L.C., Stuart, P.E., Tian, C., Gudjonsson, J.E., Das, S., Zawistowski, M., Ellinghaus, E.,
Barker, J.N., Chandran, V., Dand, N., et al. (2017). Large scale meta-analysis
characterizes genetic architecture for common psoriasis associated variants. Nat
Commun 8, 15382.

6. Consortium, G.T., Laboratory, D.A., Coordinating Center -Analysis Working, G., Statistical
Methods groups-Analysis Working, G., Enhancing, G.g., Fund, N.I.H.C., Nih/Nci,
Nih/Nhgri, Nih/Nimh, Nih/Nida, et al. (2017). Genetic effects on gene expression across
human tissues. Nature 550, 204-213.
7. Nicolae, D.L., Gamazon, E., Zhang, W., Duan, S., Dolan, M.E., and Cox, N.J. (2010). Trait-
associated SNPs are more likely to be eQTLs: annotation to enhance discovery from
GWAS. PLoS genetics 6, e1000888.
8. Lappalainen, T., Sammeth, M., Friedlander, M.R., t Hoen, P.A., Monlong, J., Rivas, M.A.,
Gonzalez-Porta, M., Kurbatova, N., Griebel, T., Ferreira, P.G., et al. (2013).
Transcriptome and genome sequencing uncovers functional variation in humans. Nature
501, 506-511.
9. Battle, A., Mostafavi, S., Zhu, X., Potash, J.B., Weissman, M.M., McCormick, C., Haudenschild,
C.D., Beckman, K.B., Shi, J., Mei, R., et al. (2014). Characterizing the genetic basis of
transcriptome diversity through RNA-sequencing of 922 individuals. Genome research
24, 14-24.
10. Gibbs, J.R., van der Brug, M.P., Hernandez, D.G., Traynor, B.J., Nalls, M.A., Lai, S.L., Arepalli,
S., Dillman, A., Rafferty, I.P., Troncoso, J., et al. (2010). Abundant quantitative trait loci
exist for DNA methylation and gene expression in human brain. PLoS genetics 6,
e1000952.
11. Gamazon, E.R., Wheeler, H.E., Shah, K.P., Mozaffari, S.V., Aquino-Michaels, K., Carroll, R.J.,
Eyler, A.E., Denny, J.C., Consortium, G.T., Nicolae, D.L., et al. (2015). A gene-based
association method for mapping traits using reference transcriptome data. Nature
genetics 47, 1091-1098.
12. Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W., Jansen, R., de Geus, E.J.,
Boomsma, D.I., Wright, F.A., et al. (2016). Integrative approaches for large-scale
transcriptome-wide association studies. Nature genetics 48, 245-252.
13. Zhu, Z., Zhang, F., Hu, H., Bakshi, A., Robinson, M.R., Powell, J.E., Montgomery, G.W.,
Goddard, M.E., Wray, N.R., Visscher, P.M., et al. (2016). Integration of summary data
from GWAS and eQTL studies predicts complex trait gene targets. Nature genetics 48,
481-487.
14. Mancuso, N., Shi, H., Goddard, P., Kichaev, G., Gusev, A., and Pasaniuc, B. (2017).
Integrating Gene Expression with Summary Association Statistics to Identify Genes
Associated with 30 Complex Traits. American journal of human genetics 100, 473-487.
15. Su, Y.R., Di, C., Bien, S., Huang, L., Dong, X., Abecasis, G., Berndt, S., Bezieau, S., Brenner, H.,
Caan, B., et al. (2018). A Mixed-Effects Model for Powerful Association Tests in
Integrative Functional Genomics. American journal of human genetics 102, 904-919.
16. Hu, Y., Li, M., Lu, Q., Weng, H., Wang, J., Zekavat, S.M., Yu, Z., Li, B., Muchnik, S., Shi, Y., et al.
(2018). A statistical framework for cross-tissue transcriptome-wide association analysis.
bioRxiv.
17. Zou, H., and Hastie, T. (2005). Regularization and variable selection via the elastic net.
Journal of the Royal Statistical Society: Series B (Statistical Methodology) 67, 301-320.
18. Zhou, X., Carbonetto, P., and Stephens, M. (2013). Polygenic modeling with bayesian sparse
linear mixed models. PLoS genetics 9, e1003264.

19. Tibshirani, R. (1996). Regression Shrinkage and Selection via the Lasso. Journal of the Royal
Statistical Society Series B (Methodological) 58, 267-288.
20. Hoerl, A.E., and Kennard, R.W. (2000). Ridge Regression: Biased Estimation for
Nonorthogonal Problems. Technometrics 42, 80-86.
21. Li, Q., and Lin, N. (2010). The Bayesian elastic net. Bayesian Anal 5, 151-170.
22. Guan, Y.T., and Stephens, M. (2011). Bayesian Variable Selection Regression for Genome-
Wide Association Studies and Other Large-Scale Problems. Annals of Applied Statistics 5,
1780-1815.
23. Yu, J., Pressoir, G., Briggs, W.H., Vroh Bi, I., Yamasaki, M., Doebley, J.F., McMullen, M.D.,
Gaut, B.S., Nielsen, D.M., Holland, J.B., et al. (2005). A unified mixed-model method for
association mapping that accounts for multiple levels of relatedness. Nature genetics 38,
203.
24. Huan, T., Liu, C., Joehanes, R., Zhang, X., Chen, B.H., Johnson, A.D., Yao, C., Courchesne, P.,
O'Donnell, C.J., Munson, P.J., et al. (2015). A systematic heritability analysis of the
human whole blood transcriptome. Human genetics 134, 343-358.
25. Lloyd-Jones, L.R., Holloway, A., McRae, A., Yang, J., Small, K., Zhao, J., Zeng, B., Bakshi, A.,
Metspalu, A., Dermitzakis, M., et al. (2017). The Genetic Architecture of Gene Expression
in Peripheral Blood. American journal of human genetics 100, 371.
26. Zeng, P., and Zhou, X. (2017). Non-parametric genetic prediction of complex traits with
latent Dirichlet process regression models. Nat Commun 8, 456.
27. Blei, D.M., and Jordan, M.I. (2006). Variational inference for Dirichlet process mixtures.
Bayesian Anal 1, 121-143.
28. Blei, D.M., Kucukelbir, A., and McAuliffe, J.D. (2017). Variational Inference: A Review for
Statisticians. Journal of the American Statistical Association 112, 859-877.
29. Casella, G. (2001). Empirical Bayes Gibbs sampling. Biostatistics 2, 485-500.
30. Bennett, D.A., Schneider, J.A., Arvanitakis, Z., and Wilson, R.S. (2012). Overview and findings
from the religious orders study. Curr Alzheimer Res 9, 628-645.
31. Bennett, D.A., Schneider, J.A., Buchman, A.S., Barnes, L.L., Boyle, P.A., and Wilson, R.S.
(2012). Overview and findings from the rush Memory and Aging Project. Curr Alzheimer
Res 9, 646-663.
32. Ng, B., White, C.C., Klein, H.U., Sieberts, S.K., McCabe, C., Patrick, E., Xu, J., Yu, L., Gaiteri, C.,
Bennett, D.A., et al. (2017). An xQTL map integrates the genetic architecture of the
human brain's transcriptome and epigenome. Nat Neurosci 20, 1418-1426.
33. Bennett, D.A., Buchman, A.S., Boyle, P.A., Barnes, L.L., Wilson, R.S., and Schneider, J.A.
(2018). Religious Orders Study and Rush Memory and Aging Project. J Alzheimers Dis 64,
S161-S189.
34. Lambert, J.C., Ibrahim-Verbaas, C.A., Harold, D., Naj, A.C., Sims, R., Bellenguez, C.,
DeStafano, A.L., Bis, J.C., Beecham, G.W., Grenier-Boley, B., et al. (2013). Meta-analysis
of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. Nature
genetics 45, 1452-1458.
35. Reitz, C. (2014). Genetic loci associated with Alzheimer's disease. Future Neurol 9, 119-122.
36. Reitz, C. (2015). Novel susceptibility loci for Alzheimer's disease. Future Neurol 10, 547-558.
37. Sims, R., van der Lee, S.J., Naj, A.C., Bellenguez, C., Badarinarayan, N., Jakobsdottir, J.,
Kunkle, B.W., Boland, A., Raybould, R., Bis, J.C., et al. (2017). Rare coding variants in

PLCG2, ABI3, and TREM2 implicate microglial-mediated innate immunity in Alzheimer's
disease. Nature genetics 49, 1373-1384.
38. Yuan, X.Z., Sun, S., Tan, C.C., Yu, J.T., and Tan, L. (2017). The Role of ADAM10 in Alzheimer's
Disease. J Alzheimers Dis 58, 303-322.
39. Muller, P., and Mitra, R. (2013). Bayesian Nonparametric Inference - Why and How.
Bayesian Anal 8.
40. Carbonetto, P., and Stephens, M. (2012). Scalable Variational Inference for Bayesian
Variable Selection in Regression, and Its Accuracy in Genetic Association Studies.
Bayesian Analysis 7, 73-107.
41. Li, B., and Leal, S.M. (2008). Methods for detecting associations with rare variants for
common diseases: application to analysis of sequence data. American journal of human
genetics 83, 311-321.
42. Genomes Project, C., Abecasis, G.R., Auton, A., Brooks, L.D., DePristo, M.A., Durbin, R.M.,
Handsaker, R.E., Kang, H.M., Marth, G.T., and McVean, G.A. (2012). An integrated map
of genetic variation from 1,092 human genomes. Nature 491, 56-65.
43. O'Reilly, P.F., Hoggart, C.J., Pomyen, Y., Calboli, F.C., Elliott, P., Jarvelin, M.R., and Coin, L.J.
(2012). MultiPhen: joint model of multiple phenotypes can increase discovery in GWAS.
PloS one 7, e34861.
44. De Jager, P.L., Shulman, J.M., Chibnik, L.B., Keenan, B.T., Raj, T., Wilson, R.S., Yu, L., Leurgans,
S.E., Tran, D., Aubin, C., et al. (2012). A genome-wide scan for common variants
affecting the rate of age-related cognitive decline. Neurobiol Aging 33, 1017 e1011-1015.
45. De Jager, P.L., Srivastava, G., Lunnon, K., Burgess, J., Schalkwyk, L.C., Yu, L., Eaton, M.L.,
Keenan, B.T., Ernst, J., McCabe, C., et al. (2014). Alzheimer's disease: early alterations in
brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. Nat Neurosci 17, 1156-
1163.
46. Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a tool for genome-wide
complex trait analysis. American journal of human genetics 88, 76-82.
47. Yang, J., Fritsche, L.G., Zhou, X., Abecasis, G., and International Age-Related Macular
Degeneration Genomics, C. (2017). A Scalable Bayesian Method for Integrating
Functional Information in Genome-wide Association Studies. American journal of human
genetics 101, 404-416.
529

Table 1. Average regression R^2 in the training data and imputation R^2 in the test data under various simulation scenarios, when the proportion of true causal SNPs is 1% and 20% with expression heritability $h_e^2 = (5\%, 10\%, 20\%, 50\%)$. The best imputation R^2 in the test samples are bold.

h_e^2		Causal Proportion 1%		Causal Proportion 20%	
		DPR	PrediXcan	DPR	PrediXcan
5%	Training R^2	17.60%	12.38%	17.60%	12.54%
	Test R^2	**1.60%**	1.12%	**1.54%**	0.76%
10%	Training R^2	21.57%	17.48%	21.33%	18.24%
	Test R^2	**4.44%**	4.13%	**4.54%**	2.64%
20%	Training R^2	31.80%	30.27%	31.48%	31.30%
	Test R^2	12.26%	**12.29%**	**12.02%**	9.13%
50%	Training R^2	60.50%	58.40%	60.71%	61.91%
	Test R^2	39.30%	**42.05%**	**39.32%**	36.04%

Figure 1. Boxplots of average imputation R^2 in test samples by DPR and PrediXcan under various simulation scenarios with various proportions of true causal SNPs (1%, 5%, 10%, 20%) and true expression heritability $h_e^2 = (5\%, 10\%, 20\%, 50\%)$.

Figure 2. Power of genetic association studies by using DPR and PrediXcan imputation models, under various simulation scenarios with proportions of true causal SNPs (1%, 5%, 10%, 20%), expression and phenotype heritability $\left(h_e^2, h_p^2\right) = \left((5\%, 80\%), (10\%, 50\%), (20\%, 25\%), (50\%, 10\%)\right)$.

Figure 3. Histogram plots of transcriptome-wide regression R^2 by DPR and PrediXcan.

Figure 4. Manhattan plot for the multivariate phenotype associations studies using imputed GReX by DPR and two AD pathologic indices, neurofibrillary tangle density (Tangles) and β-amyloid load (Amyloid), from individual-level ROS/MAP data.

Figure 5. Bar plot of -log10(FDR) of gene-based association studies for known AD loci using summary-level GWAS statistics from IGAP and the transcriptomic imputation models by DPR and PrediXcan from the ROS/MAP data. The horizontal line shows the significance level with FDR=0.05. The missing values are due to either zero regression R^2 by PrediXcan or large FDR values (~1) using the DPR imputation models.