TIGAR: An Improved Bayesian Tool for Transcriptomic Data Imputation Enhances Gene
Mapping of Complex Traits

Sini Nagpal[1], Xiaoran Meng[2,3], Michael P. Epstein[2,3], Lam C. Tsoi[4], Matthew Patrick[5], Greg
Gibson[1], Philip L. De Jager[6], David A. Bennett[7], Aliza P. Wingo[8,9], Thomas S. Wingo[3,10], Jingjing
Yang[3*]

1. School of Biology, Georgia Institute of Technology, Atlanta, GA, 30332, USA
2. Department of Biostatistics and Bioinformatics, Emory University School of Public Health, Atlanta, GA, 30322, USA
3. Center for Computational and Quantitative Genetics, Department of Human Genetics, Emory University School of Medicine, Atlanta, GA, 30322, USA
4. Department of Dermatology; Department of Computational Medicine & Bioinformatics; Department of Biostatistics, University of Michigan, Ann Arbor, MI, 48109, USA
5. Department of Dermatology, University of Michigan Medical School, Ann Arbor, MI, 48109, USA
6. Medical Center Neurological Institute, Columbia University, New York, NY, 10032, USA
7. Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, 60612, USA
8. Division of Mental Health, Atlanta VA Medical Center, Decatur, GA, USA
9. Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, 30322, USA
10. Department of Neurology, Emory University School of Medicine, Atlanta, GA, 30322, USA

*Correspondence author: jingjing.yang@emory.edu

Abstract

The transcriptome-wide association studies (TWAS) that test for association between the study trait and the imputed gene expression levels from cis-acting expression quantitative trait loci (cis-eQTL) genotypes have successfully enhanced the discovery of genetic risk loci for complex traits. By using the gene expression imputation models fitted from reference datasets that have both genetic and transcriptomic data, TWAS facilitates gene-based tests with GWAS data while accounting for the reference transcriptomic data. The existing TWAS tools like PrediXcan and FUSION use parametric imputation models that have limitations for modeling the complex genetic architecture of transcriptomic data. Therefore, to improve on this, we propose to use a Bayesian method that assumes a data-driven nonparametric prior to impute gene expression. The nonparametric Bayesian method is flexible and general because it includes both of the parametric imputation models used by PrediXcan and FUSION as special cases. Our simulation studies

showed that the nonparametric Bayesian model improved both imputation R^2 for transcriptomic

data and the TWAS power over PrediXcan. In real applications, our nonparametric Bayesian

method fitted transcriptomic imputation models for 57.6% more genes over PrediXcan, thus

improving the power of follow-up TWAS. Hence, the nonparametric Bayesian model is preferred

for modeling the complex genetic architecture of transcriptomes and is expected to enhance

transcriptome-integrated genetic association studies. We implement our Bayesian approach in a

convenient software tool "**TIGAR**" (**T**ranscriptome-**I**ntegrated **G**enetic **A**ssociation **R**esource),

which imputes transcriptomic data and performs subsequent TWAS using individual-level or

summary-level GWAS data.

Introduction

Genome-wide association studies (GWAS) have successfully identified thousands of

genetic risk loci for complex traits. However, the majority of these loci are located within noncoding

regions whose molecular mechanisms remain unknown[1-3]. Recent studies have shown that these

associated regions were enriched for regulatory elements such as enhancers (H3K27ac marks)[4;

5] and expression of quantitative trait loci (eQTL)[6; 7], suggesting that the genetically regulated gene

expression might play a key role in the biological mechanisms of complex traits. Multiple studies

have recently generated rich transcriptomic datasets for diverse tissues of the human body, e.g.,

the Genotype-Tissue Expression (GTEx) project for 44 human tissues[6], Genetic European

Variation in Health and Disease (GEUVADIS) for lymphoblastoid cell lines[8], Depression Genes

and Networks (DGN) for whole-blood samples[9], and the North American Brain Expression

Consortium (NABEC) for cortex tissues[10]. Previous studies[11-16] have also shown that integrating

transcriptomic data in GWAS can help identify novel functional loci.

The majority of GWAS projects do not possess transcriptomic data and thus cannot

directly conduct integrative analysis. Existing studies[11; 12] have shown that one can impute the

genetically regulated gene expression (GReX) within such GWAS projects by using reference

datasets like GTEx[6] and GEUVADIS[8], and then test for association between imputed GReX for

GWAS samples and the trait of interest — referred to as transcriptome-wide association studies

(TWAS)[11; 12]. Specifically, the gene expression imputation models are fitted by regressing assayed

gene-expression levels on cis-eQTL genotypes with reference dataset. For examples, the

PrediXcan[11] method uses an Elastic-Net[17] variable selection model on reference dataset to

estimate the cis-eQTL effect-sizes that are used to impute the GReX for GWAS samples, while

the FUSION[12] tool implements a Bayesian sparse linear mixed model (BSLMM)[18] to estimate the

cis-eQTL effect-sizes.

73   Basically, the Elastic-Net[17] model used by PrediXcan[11] assumes a combination of

LASSO[19] (L_1) and Ridge[20] (L_2) penalties on the cis-eQTL effect-sizes, which is equivalent to a

Bayesian model with a mixture Gaussian and Laplace prior[21]. Whereas, the BSLMM[18] used by

FUSION[12] is a combination of Bayesian variable selection model (BVSR)[22] and linear mixed

model (LMM)[23] by assuming a normal mixture prior. However, a parametric prior is assumed for

the cis-eQTL effect-sizes by both Elastic-Net and BSLMM, which restricts the capability of

PrediXcan and FUSION for handling the underlying complex genetic architecture of

transcriptomes. Existing studies[11; 12] have also shown that both PrediXcan[11] and FUSION[12]

estimated the average regression R^2 (i.e., the percentage of gene expression variation that can

be explained by cis- genotypes) as ~5% for human whole blood transcriptome, while the average

genome-wide heritability of gene expression in human whole blood transcriptome is estimated to

be more than double that quantity[24; 25]. As shown in the result section of this paper, similar

underestimation of regression R^2 by the Elastic-Net model (used by PrediXcan[11]) was observed

in both simulated and real datasets.

87   Therefore, to flexibly model cis-eQTL distributions, we propose to use a nonparametric

Bayesian method, where the prior for effect-sizes is nonparametric and can be estimated from

the data by assuming a Dirichlet process prior on effect-size variance. This Bayesian model is

also known as latent Dirichlet process regression (DPR) model[26], which can flexibly model the

underlying complex genetic architecture of transcriptomes. Thus, DPR is a more generalized

model that includes Elastic-Net (implemented in PrediXcan[11]) and BSLMM (implemented in

FUSION[12]) as special cases. Consequently, DPR can robustly estimate cis-eQTLs and then

improve imputation R^2 (the squared Pearson correlation between the observed and imputed

values on test samples). Moreover, we will use a variational Bayesian algorithm[26-28] as an

alternative of Monte Carlo Markov Chain (MCMC)[29] to efficiently fit the Bayesian model.

Similar to PrediXcan[11] and FUSION[12] methods, we can use DPR to estimate cis-eQTLs

effect-sizes from a reference dataset, which can then be used for downstream TWAS using either

individual-level or summary-level GWAS data. In subsequent sections, we first describe the DPR

approach for estimating cis-eQTL effect-sizes from a reference dataset and how we can then use

these effect-sizes for downstream TWAS using either individual-level or summary-level GWAS

data. We then compare the performance of DPR with PrediXcan using both simulated data and

real SNP and transcriptome data from the Religious Orders Study and Rush Memory Aging

Project (ROS/MAP)[30-33] for studying Alzheimer's Disease (AD).

Our in-depth simulation studies demonstrated that the DPR method obtained higher

imputation R^2 on test samples, when the true expression heritability is <10% or when >1%

variants are true causal cis-eQTL. Consequently, better imputation R^2 resulted in improved power

for follow-up association studies. Meanwhile, application of DPR to the ROS/MAP study imputed

GReX for 57.6% more genes than PrediXcan. Using DPR, we also found a potentially associated

gene *TRAPPC6A* for AD pathology indices that was missed by PrediXcan. Further, by using the

transcriptomic imputation models fitted from ROS/MAP data and summary-level GWAS data

generated from the International Genomics of Alzheimer's Project (IGAP)[34], we identified 3 known

AD loci[34-38] that potentially affect the late-onsite AD risk through transcript abundance. We

conclude with a discussion of future topics and further describe our novel software tool TIGAR

(**T**ranscriptome-**I**ntegrated **G**enetic **A**ssociation **R**esource) implementing the nonparametric

Bayesian method for public use.

Materials and Methods

We consider the following additive linear regression model for estimating the cis-eQTL

effect-sizes from a reference study that has both genetic and transcriptomic data available,

$$E_g = Xw + \epsilon, \quad \epsilon \sim N(0, \sigma_\epsilon^2 I) \,; \qquad \textit{Equation 1}$$

where E_g denotes the gene expression levels (after corrections for confounding

covariates such as age, sex, and principal components (PCs)) for gene g, X denotes the

genotype matrix for all cis- genotypes (encoded as the number of minor alleles), w denotes the

corresponding cis-eQTL effect-size vector, and ϵ denotes the error term. The intercept term is

dropped in *Equation 1* for assuming both E_g and X are centered at 0. Generally, SNPs within

1MB of the flanking 5' and 3' ends (cis-SNPs) are included in this regression model and non-zero

$\widehat{w}$ will be used for follow-up analysis. In particular, the GReX will be imputed by

$$\widehat{GReX} = X_{new}\widehat{w},$$

with cis-SNP data X_{new} for GWAS samples.

Nonparametric Bayesian Method

Following the latent Dirichlet process regression (DPR) model proposed in previous

studies for predicting complex phenotypes[26], we assume a normal prior $N(0, \sigma_w^2)$ for the cis-eQTL

effect-sizes $(w_i, i = 1, \dots, p)$ and a Dirichlet process (DP) prior[39] for the effect-size variance σ_w^2 (as

in *Equation 1*):

$$w_i \sim N(0, \sigma_w^2), \ \sigma_w^2 \sim D, \ D \sim DP\,(IG(a,b), \xi). \quad \textit{Equation 2}$$

The prior distribution D deviates from the DP with base distribution as an Inverse Gamma (IG)

distribution and concentration parameter ξ. Note that σ_w^2 can be viewed as a latent variable and

integrating out σ_w^2 will induce a nonparametric prior distribution on w_i, which is equivalent to a DP

normal mixture model[26-28],

$\quad w_i \sim \sum_{k=0}^{+\infty} \pi_k N(0, \sigma_k^2), \ \sigma_k^2 \sim IG(a_k, b_k), \ \pi_k = v_k \prod_{l=0}^{k-1}(1 - v_l), \ v_k \sim Beta(1, \ \xi).$ *Equation 3*

Here, the nonparametric prior distribution on w_i is equivalently represented by a mixture normal

prior that is a weighted sum of an infinitely number of normal distributions $\left(N(0, \sigma_k^2), k = 0, \ldots, +\infty\right)$,

corresponding weight π_k is determined by $(v_l, l = 0, \ldots, k)$ with a Beta prior, and ξ in the Beta prior

(the same concentration parameter as in Equation 2) determines the number of components with

non-zero weights in the mixture normal prior. Conjugate hyper priors $\xi \sim Gamma(a_\xi, b_\xi)$ and

$\sigma_\epsilon^2 \sim IG(a_\epsilon, b_\epsilon)$ are assumed.

Generally, the hyper parameters $a_k, b_k, a_\epsilon, b_\epsilon$ in the *Inverse Gamma* distributions can be

set as 0.1 and (a_ξ, b_ξ) in the *Gamma* distribution can be set as (1, 0.1) to induce non-informative

priors for $(\sigma_k^2, \sigma_\epsilon^2, \xi)$. That is, the parameters $(\sigma_k^2, \sigma_\epsilon^2, \xi)$ will be adaptively estimated from the data

and the nonparametric prior on w_i will be data-driven. The posterior estimates for $\boldsymbol{w}$ can be

obtained by the MCMC[29] or variational Bayesian algorithm[28; 40], from the following joint conditional

posterior distribution

$$P\left(\boldsymbol{w}, \boldsymbol{\pi}, \boldsymbol{v}, \xi, \sigma_\epsilon^2 | \boldsymbol{E_g}, \boldsymbol{X}\right) \propto$$

$\quad P\left(\boldsymbol{E_g} | \boldsymbol{w}, \boldsymbol{X}, \sigma_\epsilon^2\right) P\left(\boldsymbol{w} | \boldsymbol{\pi}, \sigma_1^2, \ldots, \sigma_k^2, \ldots\right) \left(\prod_{k=0}^{+\infty} P(\sigma_k^2 | a_k, b_k)\right) P(\boldsymbol{\pi} | \boldsymbol{v}) P(\boldsymbol{v} | \xi) P(\xi | a_\xi, b_\xi) P(\sigma_\epsilon^2 | a_\epsilon, b_\epsilon).$

Particularly, the variational Bayesian algorithm[28; 40] is an approximation for the MCMC[29] with

greatly improved computational efficiency, which is also used in our tool. Please refer to the

supplementary note for technical details of both MCMC sampling and variational inference

algorithms for obtaining the Bayesian posterior estimates for the cis-eQTL effect-sizes.

162 <u>Elastic-Net and BSLMM Methods</u>

163  The Elastic-Net model[17] (used by PrediXcan[11]) estimates the cis-eQTL effect-sizes $\hat{w}$ in

164 *Equation 1* with a combination of L_1 (LASSO)[19] and L_2 (Ridge)[20] penalties by

165 $$\hat{w} = \underset{w}{\operatorname{argmin}} \left(\left\| E_g - Xw \right\|_2^2 + \lambda \left(\alpha \|w\|_1 + \frac{1}{2}(1-\alpha)\|w\|_2^2 \right) \right),$$

166 where $\|\cdot\|_2$ denotes L_2 norm, $\|\cdot\|_1$ denotes L_1 norm, $\alpha \in [0,1]$ denotes the proportion of L_1

167 penalty, and λ denotes the penalty parameter. Particularly, PrediXcan[11] takes $\alpha = 0.5$ and tunes

168 the penalty parameter λ by a 5-fold cross validation.

169  As pointed out by previous studies[17; 21], the Elastic-Net model is equivalent to a Bayesian

170 model with a mixture Gaussian and Laplace (mixture normal) prior for w, that is, $p(w) \propto$

171 $\exp\left(-\lambda \left(\alpha\|w\|_1 + \frac{1}{2}(1-\alpha)\|w\|_2^2 \right) \right)$. Whereas, the BSLMM[18] assumes a mixture of two normal as

172 the prior for cis-eQTL effect-sizes, $w_i \sim \pi N(0, (\sigma_1^2 + \sigma_2^2)) + (1-\pi)N(0, \sigma_2^2)$. That is, the BSLMM[18]

173 assumes all cis-SNPs have at least a small effect, which are normally distributed with variance

174 σ_2^2, and some proportion (π) of cis-SNPs have an additional effect, normally distributed with

175 variance σ_1^2. Particularly, with $\sigma_2^2 = 0$, BSLMM becomes BVSR[22], and with $\pi = 0$, the BSLMM

176 becomes the LMM[23]. Therefore, the DP normal mixture[26-28] as assumed by the DPR method

177 includes the parametric (mixture normal) priors for Bayesian Elastic-Net[21] and BSLMM[18] as

178 special cases, which is the main reason why DPR is a more generalized model including Elastic-

179 Net and BSLMM as special cases. Thus, we believe the DPR method can robustly model complex

180 genetic architecture and then improve the imputation R^2.

182 <u>Association Study with Univariate Phenotype</u>

183  Given individual-level GWAS data (genotype data X_{new}, phenotype Y, and covariant

184 matrix C) and cis-eQTL effect-size estimates $\hat{w}$, the follow-up TWAS (using a burden type gene-

185 based test[41]) is to test for association between $\widehat{GReX} = X_{new}\widehat{w}$ and Y based on the following

186 generalized linear regression model

$$f(E[Y|X,C]) = \eta C + \beta \widehat{GReX}. \qquad \textit{Equation 4}$$

Here, $f(\cdot)$ is a pre-specified link function, which can be set as identify function for quantitative

phenotype or set as logit function for dichotomous phenotype. The gene-based association test

is equivalent to test $H_0: \beta = 0$ in Equation 4.

If only summary-level data are available, we take the same approach as implemented by

the FUSION[12] method. Let Z denote the vector of Z-scores by single variant tests (Wald, likelihood

ratio, score tests, etc.) for all cis-SNPs. The burden Z-score for gene-based association test is

defined as

$$\tilde{Z} = \frac{Z\widehat{w}}{\sqrt{Z\widehat{w}}} = \frac{Z\widehat{w}}{\sqrt{\widehat{w}'V\widehat{w}}}, \qquad \textit{Equation 5}$$

where V denotes the covariance matrix across study SNPs that can be estimated from training

data or reference panels such as 1000 Genome Project[42].

<u>Association Study with Multivariate Phenotype</u>

To test for association between multivariate phenotypes and imputed GReX of the focal

gene, we take a similar approach as the MultiPhen method[43] using individual-level data. For

example, consider two phenotypes (Y_1, Y_2) and a covariate matrix C, we first adjust for the

covariates by taking the residuals $(\widetilde{Y_1}, \widetilde{Y_2})$ respectively from the linear regression models $Y_j =$

$\eta C + \epsilon$, $j = 1,2$. Then we test if the regression R^2 is significantly greater than zero ($H_0: R^2 = 0$) for

the following regression model

$$\widehat{GReX}_g = \beta_1 \widetilde{Y_1} + \beta_2 \widetilde{Y_2} + \epsilon. \qquad \textit{Equation 6}$$

That is, we test if the multivariate phenotypes can jointly explain a non-zero percentage of

variance in the imputed GReX. The p-value can be calculated by using the F-statistic for the

regression R^2 in *Equation 6*.

Even when only summary-level GWAS data are available, we can first obtain a burden Z-

score per phenotype from *Equation 5*, i.e., $\widetilde{\mathbf{Z}} = (\widetilde{Z_1}, \widetilde{Z_2})$ with two phenotypes. Then a similar

burden approach can be used to obtain a joint Z-score for multi-phenotype test,

$$\tilde{Z}_{joint} = \frac{\widetilde{\mathbf{Z}}\mathbf{J}}{\sqrt{\widetilde{\mathbf{Z}}\mathbf{J}}} = \frac{\widetilde{\mathbf{Z}}\mathbf{J}}{\sqrt{\mathbf{J}'\mathbf{V_Y}\mathbf{J}}}, \ \mathbf{J} = (1, \dots, 1)',$$

where $\mathbf{V_Y}$ is the covariance matrix among study traits.

<u>Simulation Study Design</u>

We conducted in-depth simulation studies to compare the performance of both PrediXcan

and DPR methods with respect to imputation R^2 in the test data and the power of association

studies. Specifically, we used 499 ROS/MAP participants[44] with both RNA-sequencing and

genotype data as training data, and genotype data from an additional 1,200 ROS/MAP

participants[44] as test data. The test sample size (1,200) was chosen arbitrarily (randomly selected

from the ROS/MAP study) to be comparable with the sample size (1,164) in the real association

study of AD pathology indices. The genotyped and imputed genetic data for 2,799 cis-SNPs (with

minor allele frequency (MAF) > 5% and Hardy-Weinberg p-value $> 10^{-5}$) of the arbitrarily chosen

gene *ABCA7* (see Figure S1 for the LD block structure) were used to simulate gene expression

levels.

We performed comprehensive scenarios that varied the proportion of causal SNPs (out of

2,799 SNPs, influenced gene expression) among values in the vector $p_{causal} =$

$(0.01, 0.05, 0.1, 0.2)$. We varied the proportion of gene expression variance explained by causal

SNPs (i.e., expression heritability), along with the proportion of phenotypic variance explained by

simulated gene expression levels (i.e., phenotypic heritability), among values in the vector

$\left(h_e^2, h_p^2\right) = \left((0.05, 0.8), (0.1, 0.5), (0.2, 0.25), (0.5, 0.1)\right)$. The phenotypic heritability was selected

arbitrarily with respect to expression heritability such that the follow-up association study power

fell within the range of $(25\%, 85\%)$. We also considered various training sample sizes (100, 300,

499) for simulation scenario with $p_{causal} = 0.2$ and $\left(h_e^2, h_p^2\right) = (0.2, 0.25)$.

With genotype matrix $\boldsymbol{X_g}$ of the randomly selected causal SNPs (according to p_{causal}), we

generated effect-sizes w_i from $N(0, 1)$ and then re-scaled the effect-sizes to ensure the targeted

h_e^2. Gene expression levels were generated by $\boldsymbol{E_g} = \boldsymbol{X_g w} + \epsilon$, with $\epsilon \sim N(0, (1 - h_e^2))$. Then the

phenotype values were generated by $\boldsymbol{Y} = \beta \boldsymbol{E_g} + \epsilon$, where β was selected with respect to h_p^2 and

$\epsilon \sim N(0, (1 - h_p^2))$.

For each scenario, we repeated simulations for 1,000 times, where we applied both

PrediXcan[11] and DPR methods to obtain imputation models with training samples, impute the

GReX for test samples, and then conduct follow-up association studies using the imputed GReX.

We did not compare with FUSION[12] using BSLMM because of the computational burden of

estimating cis-eQTL effect-sizes by MCMC (~2 hours per gene). The association study power

was calculated as the proportion of 1,000 repeated simulations with p-value $< 2.5 \times 10^{-6}$

(genome-wide significance threshold adjusting for testing 20K independent genes).

<u>ROS/MAP Data</u>

Samples in the ROS/MAP data were collected from participants of the Religious Orders

Study (ROS) and the Rush Memory and Aging Project (MAP), which are prospective cohort

studies of studying aging and dementia[30; 31; 33]. The ROS/MAP study recruited senior adults

without known dementia at enrollment, who underwent annual clinical evaluation. Brain autopsy

was done at the time of death for each participant. All participants signed an informed consent

and Anatomic Gift Act, and the studies were approved by the Institutional Review Board of Rush

University Medical Center, Chicago, IL. Specifically, microarray genotype data generated for

2,093 European-decent participants[44], were further imputed to the 1000 Genome Project Phase

3[42] in our analysis. The post-mortem brain samples (gray matter of the dorsolateral prefrontal

cortex) from ~30% these participants were profiled for transcriptomic data by next-generation

RNA-seqencing[45]. In this paper, we conducted gene-based association studies for two important

indices of AD pathology that were quantified with β-antibody specific immunostains[30; 31; 33]:

neurofibrillary tangle density (tangles) with stereology and β-amyloid load (amyloid) with image

analysis. The neurofibrillary tangle density quantifies the average Tau tangle density within two

or more 20μm sections from eight brain regions — hippocampus, entorhinal cortex, midfrontal

cortex, inferior temporal, angular gyrus, calcarine cortex, anterior cingulate cortex, and superior

frontal cortex. The β-amyloid load quantifies the average percent area of cortex occupied by β-

amyloid protein in adjacent sections from the same eight brain regions.

Results

Simulation Studies

In the simulation studies, we observed that the DPR method performed robustly with

respect to different causal proportions and gene expression heritability. Especially, when

$p_{causal} > 0.01$ DPR outperformed PrediXcan across all expression heritability values, giving

higher imputation R^2 in test data (Figure 1A). For example, when $p_{causal} = 0.2$, the average

imputation R^2 of 1,000 simulations was estimated as 4.54% by using DPR versus 2.64% by using

PrediXcan with $h_e^2 = 0.1$, while the average imputation R^2 was estimated as 12.02% by using

DPR versus 9.13% by using PrediXcan with $h_e^2 = 0.2$ (Table 1). When $p_{causal} = 0.01$, DPR

performed similarly as PrediXcan with $h_e^2 = (0.2, 0.5)$ but slightly out-performed PrediXcan with

$h_e^2 = (0.05, 0.1)$ (Table 1, Figure 1).

Consequently, when $p_{causal} > 0.01$ and $h_e^2 < 0.5$, or $p_{causal} = 0.01$ and $h_e^2 < 0.2$, the

power of association studies was higher by using DPR versus using PrediXcan imputation models

(Figure 1B). When $h_e^2 = 0.5$, using both imputation models led to comparable power for

association studies (Figure 1B). Even though both methods had similar over-estimated training

R^2 (Figure S2), the DPR method still resulted in higher imputation R^2 for test data (Table 1;

Figures 1A) and higher power for association studies (Figure 1B). In addition, from the simulation

studies with various training sample sizes (100, 300, 499) and $p_{causal} = 0.2$ and $\left(h_e^2, h_p^2\right) =$

$(0.2, 0.25)$, the imputation R^2 and TWAS power increases as sample size increases while the DPR

method consistently outperforms PrediXcan (Figure 2). Overall, these results demonstrated the

advantages of the DPR method for modeling the complex genetic architecture of transcriptomes,

especially when the causal proportions > 0.01 or the expression heritability < 0.2.

292
<u>Real Applications to ROS/MAP Data</u>

To illustrate the performance of the DPR method in real studies, we applied both DPR and

PrediXcan on the ROS/MAP data (see Methods). We trained the gene expression imputation

models using 499 samples that have both transcriptomic data for frontal cortex tissues and

genotype data (imputed to 1000 Genome Phase 3, with MAF > 5%, Hardy-Weinberg p-value >

10^{-5}, and genotype imputation $R^2 > 0.3$). A total of 15,583 genes had gene expression levels

after standard RNA-sequencing quality control. The gene expression levels were first adjusted for

age at death, sex, postmortem interval, study (ROS or MAP), batch effects, RNA integrity number

scores, and cell type proportions (with respect to oligodendrocytes, astrocytes, microglia, neurons)

by linear regression models. For each gene, cis- SNPs within the 1MB of the flanking 5' and 3'

ends were used in the imputation models as predictors.

In this real study, we compared transcriptome-wide 5-fold cross validation (CV) regression

R^2 estimated by using both DPR and PrediXcan methods. Specifically, we randomly split 499

training samples into 5 folds, where the imputation R^2 of each fold was calculated using the model

trained with the other 4 folds samples. If the training model is null, we take the imputation R^2 as

0 and take the average imputation R^2 across all 5-fold test samples as 5-fold CV R^2. The

transcriptome-wide median of 5-fold CV R^2 is 0.013 by DPR versus 0.005 by PrediXcan. The 5-

fold CV R^2 was used as the criterion for selecting significant imputation models ($R^2 > 0.01$ as

used by previous studies[11; 46]). From Figure 3A, we can see that the DPR method obtained more

imputation models and higher imputation R^2 when 5-fold CV R^2 is in the range of (0.01, 0.05),

which is also consistent with our simulation studies. Overall, the DPR method obtained significant

imputation models for 8,742 genes versus 5,547 genes by PrediXcan (with 57.6% increases).

Thus, the DPR method featuring data-driven nonparametric prior for the cis-eQTL is preferred in

real studies for identifying more genes with imputable expression levels.

Next, we used all 499 training samples to fit imputation models for genes with respective

5-fold CV $R^2 > 0.01$ by both DPR and PrediXcan, and then used these models to impute the

GReX for all GWAS samples. We conducted univariate phenotype association studies (Methods)

using all GWAS samples (N = 1,164) that have the AD pathology indices (neurofibrillary tangle

density and β-amyloid load, with Pearson correlation 0.48) quantified. Possible confounding

covariates including age at death, sex, study (ROS or MAP), smoking, education, and first 3

genotype principle components were adjusted in the association studies. Interestingly, the

association studies for both AD pathology indices using the DPR imputation models identified the

same top significant gene *TRAPPC6A* (within the 2MB region from the major risk gene *APOE*,

encoding apolipoprotein E, but independent of *APOE*) with p-values 1.64×10^{-5} and 5.35×10^{-5}

(Figure S3A and Figure S4A). Moreover, the multivariate phenotype association studies (Methods)

for both AD pathology indices also identified *TRAPPC6A* as the most significant gene with p-value

5.81×10^{-6} and FDR 0.08 (Figure 3C). On the other hand, the PrediXcan failed to obtain a

transcriptomic imputation model for *TRAPPC6A* (Figure S3B, Figure S4B, and Figure S6).

Quantile-Quantile plots for these TWAS p-values were presented in Figure S5.

In addition, for 14 known common and rare loci of late-onset AD[34-38] with significant

imputation models, we conducted association studies using transcriptomic imputation models

(DPR and PrediXcan) fitted from ROS/MAP data and standard GWAS results from IGAP[34]. Using

the imputation models fit by DPR, we identified 3 significant loci with FDR < 0.05 (Figure 3B) —

ADAM10, CD2AP, and *TREM2* — that potentially affect late-onsite AD risk through transcript

abundance. Here, *TREM2* was also identified by using the PrediXcan imputation model (Figure

3B). Particularly, the PrediXcan method only imputed GReX for 5 out of these 14 loci. In summary,

these results show that the DPR method has superior power for follow-up TWAS.

Discussion

In this paper, we propose an improved nonparametric Bayesian method for transcriptomic

data imputation, accompanied with an integrated tool (freely available on GITHUB) that is referred

as **T**ranscriptome-**I**ntegrated **G**enetic **A**ssociation **R**esource (**TIGAR**). TIGAR integrates both

nonparametric Bayesian (DPR) and Elastic-Net models as two options for transcriptomic data

imputation, along with TWAS options using individual-level and summary-level GWAS data for

univariate and multi-variate phenotypes. We show the usefulness of the DPR method by both in-

depth simulations and real applications using individual-level ROS/MAP[30-33] and summary-level

IGAP[34] data. We demonstrate that DPR is preferred over PrediXcan because of the flexible

nonparametric modeling of cis-eQTL effect-sizes that results in improved imputation R^2 for gene

expression levels and higher power for TWAS.

With respect to user-friendly interface and computational efficiency, TIGAR can: (i) Take

standard input files such as genotype files in VCF and dosage formats, phenotype files in PED

format, and a combined file (tab delimited) for gene annotations and expression levels; (ii) Load

input data per gene by TABIX for memory efficiency; (iii) Filter SNPs by given MAF and Hardy-

Weinberg p-value thresholds; (iv) Provide options of training both Elastic-Net (using Python

scripts) and DPR (using the executable tool developed with C++[26]) imputation models with unified

output format; (v) Implement multi-threaded computation to take full advantage of multi-core

clusters. These features make TIGAR a preferred tool for saving tedious data preparation and

computation time. For example, TIGAR can complete training imputation models for ~20K genes

and ~1K samples within ~20 hours and TWAS within ~1hour with a 2.4GHz 16-core CPU. The

computation time is expected to increase linearly with respect to the sample size.

It is important to notice that imputing GReX with cis-eQTL effect-sizes estimated from a

training dataset is analogous to the idea of estimating polygenic risk scores (PRS)[47]. Even though

studies of population heterogeneity are lacked for imputing GReX, the same philosophy of

estimating PRS still applies because of the same underlying statistical models. That is, given both

genetic and transcriptomic heterogeneities across different populations, one need to be cautious

not using training dataset of a different ethnicity for TWAS[47].

As observed in the real ROS/MAP studies, there remains a large gap between the 5-fold

CV R^2 using cis-eQTL predictors (~5%) and the average genome-wide heritability of gene

expression levels (21.8% estimated by GCTA[48] based on a LMM). This is likely due to the large

trans-acting contribution to transcript abundance documented for most genes. Thus, we

hypothesize that it is promising to further improve the imputation R^2 by fitting transcriptomic

imputation models with genome-wide variants as predictors. Scalable Bayesian inference

techniques such as the Expectation Maximization MCMC (EM-MCMC) algorithm[49] are required

for incorporating genome-wide variants.

Another limitation of existing TWAS methods is that the uncertainty of cis-eQTL effect-size

estimates has not been taken into accounted in the association studies. A Bayesian framework

can also be derived by taking the standard errors of these cis-eQTL effect-size estimates as prior

standard deviations, which is part of our continuing research.

Besides the follow-up gene-based association studies (i.e., TWAS) described in this paper,

the transcriptomic imputation models can be further extended by incorporating environmental

contributions. The imputed transcript abundance levels can then be used for gene network

analysis, differential gene expression analysis, and transcriptome mediation analysis with GWAS

data. Validation of transcriptomic prediction accuracy in independent datasets will be critical in

this regard, but unfortunately multiple large and similar datasets are not yet generally available

for tissues other than peripheral blood.

In conclusion, we expect our work will provide a convenient and improved tool for

transcriptomic imputation using the currently available rich reference datasets, as well as

enhanced gene mapping for better understanding the genetic etiology of complex traits.

Supplemental Data

Supplementary Text and Figures

Acknowledgments

J.Y. was supported by the startup funding from Emory University Department of Human Genetics.

A.P.W. and T.S.W. were supported by National Institutes of Health (NIH) R01AG056533.

M.P.E. was supported by NIH R01GM11796. L.C.T. was supported by the Dermatology

Foundation, the Arthritis National Research Foundation, the National Psoriasis

Foundation, and NIH K01AR072129. ROS/MAP study data were provided by the Rush

Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL. Data collection was

supported through funding by NIA grants P30AG10161, R01AG15819, R01AG17917,

R01AG30146, R01AG36836, U01AG32984, U01AG46152, the Illinois Department of Public

Health, and the Translational Genomics Research Institute. In addition, we thank Thanneer

Perumal and Benjamin Logsdon for performing quality control of the ROS/MAP RNA-sequencing

data and for creating the brain cell type proportions.

Web Resources

409   TIGAR, https://github.com/yanglab-emory/TIGAR

410   PrediXcan, https://github.com/hakyim/PrediXcan

411   FUSION, http://gusevlab.org/projects/fusion/

412   ROS/MAP data, https://www.synapse.org/#!Synapse:syn3219045; http://www.radc.rush.edu/

413   IGAP data, http://web.pasteur-lille.fr/en/recherche/u744/igap/igap_download.php

References

416   1. Visscher, P.M., Brown, M.A., McCarthy, M.I., and Yang, J. (2012). Five years of GWAS discovery.
417   American journal of human genetics 90, 7-24.
418   2. McCarthy, M.I., Abecasis, G.R., Cardon, L.R., Goldstein, D.B., Little, J., Ioannidis, J.P., and
419   Hirschhorn, J.N. (2008). Genome-wide association studies for complex traits: consensus,
420   uncertainty and challenges. Nature reviews Genetics 9, 356-369.
421   3. Huang, Q. (2015). Genetic study of complex diseases in the post-GWAS era. J Genet Genomics
422   42, 87-98.
423   4. Farh, K.K., Marson, A., Zhu, J., Kleinewietfeld, M., Housley, W.J., Beik, S., Shoresh, N., Whitton,
424   H., Ryan, R.J., Shishkin, A.A., et al. (2015). Genetic and epigenetic fine mapping of causal
425   autoimmune disease variants. Nature 518, 337-343.
426   5. Tsoi, L.C., Stuart, P.E., Tian, C., Gudjonsson, J.E., Das, S., Zawistowski, M., Ellinghaus, E., Barker,
427   J.N., Chandran, V., Dand, N., et al. (2017). Large scale meta-analysis characterizes genetic
428   architecture for common psoriasis associated variants. Nat Commun 8, 15382.
429   6. GTEx Consortium. (2017). Genetic effects on gene expression across human tissues. Nature
430   550, 204-213.
431   7. Nicolae, D.L., Gamazon, E., Zhang, W., Duan, S., Dolan, M.E., and Cox, N.J. (2010). Trait-
432   associated SNPs are more likely to be eQTLs: annotation to enhance discovery from GWAS.
433   PLoS genetics 6, e1000888.

8. Lappalainen, T., Sammeth, M., Friedlander, M.R., t Hoen, P.A., Monlong, J., Rivas, M.A.,
Gonzalez-Porta, M., Kurbatova, N., Griebel, T., Ferreira, P.G., et al. (2013). Transcriptome
and genome sequencing uncovers functional variation in humans. Nature 501, 506-511.
9. Battle, A., Mostafavi, S., Zhu, X., Potash, J.B., Weissman, M.M., McCormick, C., Haudenschild,
C.D., Beckman, K.B., Shi, J., Mei, R., et al. (2014). Characterizing the genetic basis of
transcriptome diversity through RNA-sequencing of 922 individuals. Genome research 24,
14-24.
10. Gibbs, J.R., van der Brug, M.P., Hernandez, D.G., Traynor, B.J., Nalls, M.A., Lai, S.L., Arepalli,
S., Dillman, A., Rafferty, I.P., Troncoso, J., et al. (2010). Abundant quantitative trait loci
exist for DNA methylation and gene expression in human brain. PLoS genetics 6, e1000952.
11. Gamazon, E.R., Wheeler, H.E., Shah, K.P., Mozaffari, S.V., Aquino-Michaels, K., Carroll, R.J.,
Eyler, A.E., Denny, J.C., GTEx Consortium, Nicolae, D.L., et al. (2015). A gene-based
association method for mapping traits using reference transcriptome data. Nature
genetics 47, 1091-1098.
12. Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W., Jansen, R., de Geus, E.J.,
Boomsma, D.I., Wright, F.A., et al. (2016). Integrative approaches for large-scale
transcriptome-wide association studies. Nature genetics 48, 245-252.
13. Zhu, Z., Zhang, F., Hu, H., Bakshi, A., Robinson, M.R., Powell, J.E., Montgomery, G.W., Goddard,
M.E., Wray, N.R., Visscher, P.M., et al. (2016). Integration of summary data from GWAS
and eQTL studies predicts complex trait gene targets. Nature genetics 48, 481-487.
14. Mancuso, N., Shi, H., Goddard, P., Kichaev, G., Gusev, A., and Pasaniuc, B. (2017). Integrating
Gene Expression with Summary Association Statistics to Identify Genes Associated with
30 Complex Traits. American journal of human genetics 100, 473-487.
15. Su, Y.R., Di, C., Bien, S., Huang, L., Dong, X., Abecasis, G., Berndt, S., Bezieau, S., Brenner, H.,
Caan, B., et al. (2018). A Mixed-Effects Model for Powerful Association Tests in Integrative
Functional Genomics. American journal of human genetics 102, 904-919.
16. Hu, Y., Li, M., Lu, Q., Weng, H., Wang, J., Zekavat, S.M., Yu, Z., Li, B., Muchnik, S., Shi, Y., et al.
(2018). A statistical framework for cross-tissue transcriptome-wide association analysis.
bioRxiv.
17. Zou, H., and Hastie, T. (2005). Regularization and variable selection via the elastic net. Journal
of the Royal Statistical Society: Series B (Statistical Methodology) 67, 301-320.
18. Zhou, X., Carbonetto, P., and Stephens, M. (2013). Polygenic modeling with bayesian sparse
linear mixed models. PLoS genetics 9, e1003264.
19. Tibshirani, R. (1996). Regression Shrinkage and Selection via the Lasso. Journal of the Royal
Statistical Society Series B (Methodological) 58, 267-288.
20. Hoerl, A.E., and Kennard, R.W. (2000). Ridge Regression: Biased Estimation for Nonorthogonal
Problems. Technometrics 42, 80-86.
21. Li, Q., and Lin, N. (2010). The Bayesian elastic net. Bayesian Anal 5, 151-170.
22. Guan, Y.T., and Stephens, M. (2011). Bayesian Variable Selection Regression for Genome-
Wide Association Studies and Other Large-Scale Problems. Annals of Applied Statistics 5,
1780-1815.
23. Yu, J., Pressoir, G., Briggs, W.H., Vroh Bi, I., Yamasaki, M., Doebley, J.F., McMullen, M.D., Gaut,
B.S., Nielsen, D.M., Holland, J.B., et al. (2005). A unified mixed-model method for

association mapping that accounts for multiple levels of relatedness. Nature genetics 38,
203.
24. Huan, T., Liu, C., Joehanes, R., Zhang, X., Chen, B.H., Johnson, A.D., Yao, C., Courchesne, P.,
O'Donnell, C.J., Munson, P.J., et al. (2015). A systematic heritability analysis of the human
whole blood transcriptome. Human genetics 134, 343-358.
25. Lloyd-Jones, L.R., Holloway, A., McRae, A., Yang, J., Small, K., Zhao, J., Zeng, B., Bakshi, A.,
Metspalu, A., Dermitzakis, M., et al. (2017). The Genetic Architecture of Gene Expression
in Peripheral Blood. American journal of human genetics 100, 371.
26. Zeng, P., and Zhou, X. (2017). Non-parametric genetic prediction of complex traits with latent
Dirichlet process regression models. Nat Commun 8, 456.
27. Blei, D.M., and Jordan, M.I. (2006). Variational inference for Dirichlet process mixtures.
Bayesian Anal 1, 121-143.
28. Blei, D.M., Kucukelbir, A., and McAuliffe, J.D. (2017). Variational Inference: A Review for
Statisticians. Journal of the American Statistical Association 112, 859-877.
29. Casella, G. (2001). Empirical Bayes Gibbs sampling. Biostatistics 2, 485-500.
30. Bennett, D.A., Schneider, J.A., Arvanitakis, Z., and Wilson, R.S. (2012). Overview and findings
from the religious orders study. Curr Alzheimer Res 9, 628-645.
31. Bennett, D.A., Schneider, J.A., Buchman, A.S., Barnes, L.L., Boyle, P.A., and Wilson, R.S. (2012).
Overview and findings from the rush Memory and Aging Project. Curr Alzheimer Res 9,
646-663.
32. Ng, B., White, C.C., Klein, H.U., Sieberts, S.K., McCabe, C., Patrick, E., Xu, J., Yu, L., Gaiteri, C.,
Bennett, D.A., et al. (2017). An xQTL map integrates the genetic architecture of the human
brain's transcriptome and epigenome. Nat Neurosci 20, 1418-1426.
33. Bennett, D.A., Buchman, A.S., Boyle, P.A., Barnes, L.L., Wilson, R.S., and Schneider, J.A. (2018).
Religious Orders Study and Rush Memory and Aging Project. J Alzheimers Dis 64, S161-
S189.
34. Lambert, J.C., Ibrahim-Verbaas, C.A., Harold, D., Naj, A.C., Sims, R., Bellenguez, C., DeStafano,
A.L., Bis, J.C., Beecham, G.W., Grenier-Boley, B., et al. (2013). Meta-analysis of 74,046
individuals identifies 11 new susceptibility loci for Alzheimer's disease. Nature genetics
45, 1452-1458.
35. Reitz, C. (2014). Genetic loci associated with Alzheimer's disease. Future Neurol 9, 119-122.
36. Reitz, C. (2015). Novel susceptibility loci for Alzheimer's disease. Future Neurol 10, 547-558.
37. Sims, R., van der Lee, S.J., Naj, A.C., Bellenguez, C., Badarinarayan, N., Jakobsdottir, J., Kunkle,
B.W., Boland, A., Raybould, R., Bis, J.C., et al. (2017). Rare coding variants in PLCG2, ABI3,
and TREM2 implicate microglial-mediated innate immunity in Alzheimer's disease. Nature
genetics 49, 1373-1384.
38. Yuan, X.Z., Sun, S., Tan, C.C., Yu, J.T., and Tan, L. (2017). The Role of ADAM10 in Alzheimer's
Disease. J Alzheimers Dis 58, 303-322.
39. Muller, P., and Mitra, R. (2013). Bayesian Nonparametric Inference - Why and How. Bayesian
Anal 8.
40. Carbonetto, P., and Stephens, M. (2012). Scalable Variational Inference for Bayesian Variable
Selection in Regression, and Its Accuracy in Genetic Association Studies. Bayesian Analysis
7, 73-107.

41. Li, B., and Leal, S.M. (2008). Methods for detecting associations with rare variants for common
diseases: application to analysis of sequence data. American journal of human genetics
83, 311-321.
42. 1000 Genomes Project Consortium. (2012). An integrated map of genetic variation from 1,092
human genomes. Nature 491, 56-65.
43. O'Reilly, P.F., Hoggart, C.J., Pomyen, Y., Calboli, F.C., Elliott, P., Jarvelin, M.R., and Coin, L.J.
(2012). MultiPhen: joint model of multiple phenotypes can increase discovery in GWAS.
PloS one 7, e34861.
44. De Jager, P.L., Shulman, J.M., Chibnik, L.B., Keenan, B.T., Raj, T., Wilson, R.S., Yu, L., Leurgans,
S.E., Tran, D., Aubin, C., et al. (2012). A genome-wide scan for common variants affecting
the rate of age-related cognitive decline. Neurobiol Aging 33, 1017 e1011-1015.
45. De Jager, P.L., Srivastava, G., Lunnon, K., Burgess, J., Schalkwyk, L.C., Yu, L., Eaton, M.L.,
Keenan, B.T., Ernst, J., McCabe, C., et al. (2014). Alzheimer's disease: early alterations in
brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. Nat Neurosci 17, 1156-
1163.
46. Wu, L., Shi, W., Long, J., Guo, X., Michailidou, K., Beesley, J., Bolla, M.K., Shu, X.O., Lu, Y., Cai,
Q., et al. (2018). A transcriptome-wide association study of 229,000 women identifies new
candidate susceptibility genes for breast cancer. Nature genetics 50, 968-978.
47. Martin, A.R., Gignoux, C.R., Walters, R.K., Wojcik, G.L., Neale, B.M., Gravel, S., Daly, M.J.,
Bustamante, C.D., and Kenny, E.E. (2017). Human Demographic History Impacts Genetic
Risk Prediction across Diverse Populations. American journal of human genetics 100, 635-
649.
48. Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a tool for genome-wide
complex trait analysis. American journal of human genetics 88, 76-82.
49. Yang, J., Fritsche, L.G., Zhou, X., Abecasis, G., and International Age-Related Macular
Degeneration Genomics, C. (2017). A Scalable Bayesian Method for Integrating Functional
Information in Genome-wide Association Studies. American journal of human genetics
101, 404-416.
548

Table 1. *Average imputation R^2 in the test data under various simulation scenarios, when the proportion of true causal SNPs $p_{causal} = (0.01, 0.2)$ and expression heritability $h_e^2 = (0.05, 0.1, 0.2, 0.5)$. The best imputation R^2 in the test samples are bold.*

h_e^2	Causal Proportion 0.01		Causal Proportion 0.2	
	DPR	PrediXcan	DPR	PrediXcan
0.05	**1.60%**	1.12%	**1.54%**	0.76%
0.1	**4.44%**	4.13%	**4.54%**	2.64%
0.2	12.26%	**12.29%**	**12.02%**	9.13%
0.5	39.30%	**42.05%**	**39.32%**	36.04%

Figure 1. *Plots of average imputation R^2 (**A**) and TWAS power (**B**) in test samples by DPR and PrediXcan, with various proportions of true causal SNPs $p_{causal} = (0.01, 0.05, 0.1, 0.2)$ and true expression heritability $h_e^2 = (0.05, 0.1, 0.2, 0.5)$. TWAS Power was evaluated with paired expression and phenotype heritability $\left(h_e^2, h_p^2\right) = \left((0.05,0.8), (0.1, 0.5), (0.2,0.25), (0.5,0.1)\right)$.*

Figure 2. Test R^2 (**A**) and TWAS power (**B**) from simulation studies with causal proportion $p_{causal} = 0.2$, expression heritability and phenotype heritability $(h_e^2, h_p^2) = (0.2, 0.25)$, and various training sample sizes $(100, 300, 499)$.

Figure 3. *Transcriptome-wide 5-fold cross validation R² (**A**) by PrediXcan and DPR with 499 ROS/MAP training samples, with different colors denoting whether the imputation $R^2 > 0.01$ by DPR, PrediXcan, or both methods (genes with $R^2 < 0.01$ by both DPR and PrediXcan were excluded from the plot). TWAS results (**B**) at known AD loci using GWAS summary-level statistics from IGAP and imputation models fitted from ROS/MAP data, where missing values are due to NULL imputation models by PrediXcan. Manhattan plot (**C**) for the multiphenotype TWAS (with neurofibrillary tangle density and β-amyloid load), using individual-level ROS/MAP data.*