Title: Mitf links neuronal activity and long-term homeostatic intrinsic plasticity

Abbreviated title: Mitf regulates intrinsic plasticity

Diahann A. M. Atacho[1,3], Hallur Reynisson[2], Anna Þóra Pétursdóttir[1,3], Thor Eysteinsson[2], Eiríkur Steingrímsson[3], Pétur Henry Petersen[1,4*]

[1]Department of Anatomy; [2]Department of Physiology; [3]Department of Biochemistry and Molecular Biology, [4]Department of Pharmacology, BioMedical Center, Faculty of Medicine, University of Iceland, 101 Reykjavik, Iceland.

*Corresponding author: phenry@hi.is, tel: +354 845 53 46, address: Vatnsmyrarvegur 16, 101 Reykjavik, Iceland.

Number of pages: 29 + SI

Number of figures: 11

Number of tables: 3

Number of words abstract: 230

Number of words introduction: 423

Number of words discussion: 1016

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: This research was supported by grants of the Icelandic Research Fund, Rannís grant numbers: [152715-053 and 163068-051]. We thank our colleagues from the BioMedical Center at the University of Iceland, Erna Magnúsdóttir PhD, Ragnhildur Káradóttir PhD, Margrét Helga Ögmundsdóttir PhD, Ramile Dilshat, Kimberley Anderson, Alba Sabate and Unnur Diljá Teitsdóttir for their help.

Abstract

Neuroplasticity forms the basis of neuronal circuit functional complexity and can determine differences between otherwise similar neuronal circuits. Although synaptic plasticity is fairly well characterized, much less is known about the molecular mechanisms underlying intrinsic plasticity, especially its transcriptional regulation. We show here that the Microphthalmia-associated transcription factor (*Mitf*), best known as the master regulator of melanocytic cell fate and differentiation, plays a central role in homeostatic intrinsic plasticity of olfactory bulb (OB) projection neurons. Mitral and tufted (M/T) neurons from *Mitf* mutant mice are hyperexcitable due to reduced Type-A potassium current (I_A) and they exhibit reduced expression of *Kcnd3* (Kv4.3)*, which encodes a potassium voltage-gated channel subunit important for the generation of the I_A. Furthermore, *Mitf* gene expression is activity-dependent in the OB and its absence leads to the loss of activity-dependent transcription of *Kcnd3* in projection neurons. The MITF protein binds to and activates expression from *Kcnd3* regulatory elements. Activity can therefore affect *Kcnd3* expression directly via MITF. Moreover, *Mitf* mutant mice have increased olfactory discriminatory ability and, unlike wild type mice, have reduced ability to detect odour following long-term odour exposure, indicating that regulation of *Kcnd3* is pivotal for the ability to recover from prolonged odour exposure. Our findings show that *Mitf* acts as a direct regulator of intrinsic homeostatic feedback, plays a key role in olfactory adaptation and links neuronal activity, transcriptional changes and neuronal function.

Significance statement

A direct, *Mitf*-dependent link between neuronal activity and homeostatic changes in the expression of a key potassium channel subunit is demonstrated in projection neurons of the mouse OB. This is one of the first studies that directly link activity and genetically defined changes in intrinsic plasticity, leading to changes in neuronal response. These findings broaden the general understanding of transcriptional regulation of homeostatic intrinsic plasticity in learning and memory. The results are also important for understanding the role of *Mitf* in other cell types. Regulation of intrinsic plasticity has wide-ranging implications and fundamental importance for neurological diseases such as neurodegeneration, autism and epilepsy.

Introduction

Activity-dependent homeostatic mechanisms lead to neurons with high activity becoming less sensitive and neurons with low activity more sensitive, thereby reestablishing a set point for neuronal activity [1-4]. Neuronal plasticity in general, is comprised of activity-dependent changes that alter synaptic or intrinsic excitability of the neuron [5]. Synaptic plasticity and intrinsic plasticity, are thought to work at different time-scales and to interact and complement each other [6, 7]. Although synaptic plasticity is fairly well characterized, much less is known about the molecular mechanisms underlying intrinsic plasticity and its transcriptional regulation [8, 9].

The olfactory system is an attractive system to study plasticity, due to its clear flow of information, well-defined neuronal subtypes, ease of activation with odourants and delimited neuronal circuits. The olfactory bulb (OB) is the first relay of peripheral olfactory information within the CNS. Each olfactory bulb (OB) in the mouse contains around 2000 glomeruli, spherical neuropil structures connecting peripheral olfactory sensory input and cortical structures. In the OB, two types of projection neurons are designated to each glomerulus and synapse with the OSN and target other CNS regions. These are the mitral cells (MC) [10, 11] located in the mitral cell layer (MCL) and the tufted neurons located in the external plexiform layer (EPL). Projection neurons of the OB show remarkable capacity in normalizing sensory input, by amplifying low signals and reducing high signals [12]. Odour exposure also reduces sensitivity of mitral and tufted cells (M/T) [13, 14]. In addition, it has been shown that there is high inter-glomerular functional diversity [15], showing that each glomerulus is, to a large degree, regulated independently by its activity. The OB therefore shows clear changes in gain-control and homeostasic plasticity and it has been suggested that intrinsic changes are fundamental to sensory adaptation in the OB [16].

The transcription factor MITF, best known as a master regulator of melanocytes [17], is specifically expressed in the M/T neurons of the OB [18]. As the M/T neurons are the key output neurons in the OB circuitry, Mitf might play a role in olfaction [18]. Mitf is also expressed in a subset of tufted cells, the external tufted cells (ETC), which are excitatory and synapse with MC, but do not project out of the OB [19-21]. Here we show that Mitf links neuronal activity and intrinsic activity-dependent changes in the OB by regulating the expression of the potassium channel subunit *Kcnd3*, leading to a homeostatic response in OB´s projection neurons. Therefore, we suggest that Mitf plays a key role in olfactory adaptation and intrinsic homeostatic plasticity.

Results

***Mitf^mi-vga9/mi-vga9* mice have increased numbers of inhibitory and excitatory neuronal subtypes in the glomerular layer.**

In order to characterize *Mitf* expression in the OB, we used multiplex fluorescent RNA *in situ* hybridization (mFISH) which generates a single fluorescent dot per transcript detected [22], allowing for single-cell analysis of gene expression. Consistent with previous observations [18], this showed that in wild type mice *Mitf* is expressed in M/T neurons, including in external tufted cells (ETC) (Figure 1a, b) and at low levels in the granule cell layer (GCL) (Figure 1a, arrowheads). Interestingly, there is significantly more *Mitf* is expressed in the ETCs compared to MCs ($F_{(1,15)}=8.94$, p=0.0092, Two-way ANOVA). Mice homozygous for the *Mitf^mi-vga9* mutation are white and microphthalmic due to a transgene insertion mutation that affects expression of *Mitf* [23]. As expected, *Mitf* expression was decreased in the ETCs ($t(14)=7.923$, p<0.0001, Sidak multiple comparison test) and MCs ($t(14)=2.969$, p=0.0202, Sidak multiple comparison) in OBs of *Mitf^mi-vga9/mi-vga9* mice.

Histological analysis showed no detectable defects in the cellular architecture of OBs from P75-P95 *Mitf^mi-vga9/mi-vga9* mice (Figure 1-1 a). Expression of tyrosine hydroxylase (TH) is reduced upon loss of activity in the OB [24] but was not affected in the OBs of *Mitf^mi-vga9/mi-vga9* mice (Figure 1-1 b, c). This also suggests normal OB function. Importantly, *Tbr2*, a marker for M/T cells [25, 26], is expressed in *Mitf^mi-vga9/mi-vga9* OBs (Figure 1 a) showing normal OB ontogeny. To assess changes in neuronal circuits of the mutant OB, the number of excitatory neurons and inhibitory interneurons was determined (Figure 1 d-i). Counting the number of excitatory and inhibitory interneurons showed an increase in glomerular Tbr2+ excitatory neurons ($t(20)=2.914$, p=0.0171, Sidak multiple comparison), Calbindin+ interneurons ($t(14)=2.469$, p=0.0489, Sidak multiple comparison) and Parvalbumin+ interneurons in the *Mitf* mutant OBs

as compared to wild type (wild-type vs $Mitf^{mi\text{-}vga9/+}$: t(30)=3.12, p=0.0236; wild-type vs $Mitf^{mi\text{-}vga9/mi\text{-}vga9}$: t(30)=4.407, p=0.0007, Sidak multiple comparison). No change was observed in Calretinin+ interneurons (Figure 1-2 a, b) or TH+ neurons (Figure1-1 b, c). Additionally, the ratio between inhibitory and excitatory neurons was not changed in the mutant mouse (SI Table 2). Interestingly, analysis of apoptosis showed a twofold increase in cell death in periglomerular cells of $Mitf^{mi\text{-}vga9/mi\text{-}vga9}$ mice (Figure 1-3; t(16)=2.724, p=0.015, Two-tailed t-test) suggesting increased glomerular neuronal turnover in the mutant mice.

$Mitf^{mi\text{-}vgag/mi\text{-}vga9}$ mice have reduced I_A and a concomittant increase in mitral cell activity.

To examine if the $Mitf^{mi\text{-}vga9}$ mutation resulted in altered M/T neuronal function, we performed whole-cell patch clamp analysis on cultured primary M/T neurons from wild-type and $Mitf^{mi\text{-}vga9/mi\text{-}vga9}$ mice both under voltage clamp and current clamp conditions, bypassing possible secondary effects due to cortical feedback mechanisms. To establish the identity of the cells in culture, several biophysical properties were determined (see methods). The likelihood of an action potential is decided by the balance between sodium influx and potassium efflux. The initial potassium efflux activated near the spiking threshold is termed the A-type potassium current (I_A) and is a major determinant of the likelihood of an action potential. Patch clamp recordings under voltage clamp showed a reduction in the I_A K⁺-current in M/T neurons from $Mitf^{mi\text{-}vga9/mi\text{-}vga9}$ mice. (Figure 2a-d; t(50)=2.313, p=0.0491, Sidak multiple comparison). No significant reduction was observed in the I_{DR} current (Figure 2d). The I/V relationship denotes the relationship between the K⁺-current density (pA/pF) across the membrane and the membrane potential, and thus takes size of the cells into account based on their measured membrane capacitance. The I/V curves shows that the mutant neurons have an altered voltage dependence of activation as the threshold for activation of the outward current is shifted approximately +5mV in the positive direction (Figure 2e), and the current density reduced at depolarized membrane potentials. However, the channel kinetics, i.e. activation and

inactivation of the channels mediating the I_A, remained the same in cells from wild-type and mutant mice (Figure 2-1). Reduction in the I_A leads to increased likelihood of action potentials and hyperexcitability [27-29]. Accordingly, increased spiking was observed in the mutant neurons (Figure 2f). The mean firing frequency of spikes in wild-type M/T neurons was 0.79 ± 0.17 Hz, while 1.75 ± 0.15 Hz in mutant M/T neurons (t(11)=1.61, p=0.007, Two-tailed t-test). The mean amplitude of action potentials was similar, or 62 ± 4 mV in wild-type M/T neurons and 54 ± 2.6 mV in mutant M/T neurons (t(11)=1.06, p=0.31, Two-tailed t-test). In addition to spikes, miniature excitatory postsynaptic potentials (EPSP) could be observed in the current clamp recordings. The mean frequency of miniature EPSPs in the wild-type M/T neurons was 0.27 ± 0.04 Hz and i 0.47 ± 0.21 Hz (t(11)=-1.55, p=0.148, Two-tailed t-test) in mutant M/T neurons. Additionally, expression of *c-Fos*, an activity-dependent gene frequently used to monitor neuronal acivity, was increased in the *Mitf*$^{mi\text{-}vga9/mi\text{-}vga9}$ MCs (Fig. 2g, h; t(9)=3.093, p=0.0386, Sidak multiple comparison), indicating increased neuronal activity *in vivo*. A second cell population in the GCL also showed clear expression of *c-Fos* in the mutant (Fig. 2g, arrowheads). These are likely the deep short-axon-cells, which inhibit ETC neurons [30].

Expression of *Kcnd3* and *Mitf* is activity-dependent in the OB.

Increased neuronal activity due to the loss of I_A led us to examine the expression of key potassium channel subunits in the OB. As displayed in SI Table 1, many potassium channel subunits [31] are expressed in the OB [32-34]. We focused our analysis on the I_A-current regulating Kv4.3/*Kcnd3* potassium channel subunit, as according to ChIPseq data, MITF binds to its regulatory region (SI Table 1). RNA *in situ* analysis showed reduced expression of *Kcnd3* in ETCs (t(12)=3.297, p=0.0190, Sidak multiple comparison) and MCs (t(12)=3.249, p=0.0208, Sidak multiple comparison) of *Mitf* $^{mi\text{-}vga9/mi\text{-}vga9}$ OBs, whereas the levels remained similar in cells of the GCL (Fig. 3a, b). Thus, the effects of *Mitf* on *Kcnd3* expression are cell autonomous,

consistent with the possibility of *Mitf* regulating *Kcnd3* in M/T neurons. The I_A-regulating potassium channel subunit *Kcnd2* did not show MITF binding to a regulatory region according to ChIPseq data (SI Table 1). However, it did show a cell-specific increase in expression in the mutant mouse OB (t(6)=5.051, p=0.0047, Sidak multiple comparison), possibly as a compensatory mechanism for the reduction in *Kcnd3* (Figure 3-1).

Neuronal activity is a major determinant of OB circuit plasticity. It leads to an increase in cAMP levels, a known regulator of MITF signaling [35]. We therefore explored the effects of OSN activity on the expression of *Mitf* and *Kcnd3* in the OB. Amyl-acetate is known to activate a multitude of glomeruli [36]. Amyl acetate treatment of the wild type mice (Figure 3c) resulted in an immediate increase in the global transcription of *c-Fos* (Figure 3-2, Figure 3e; q(40)=4.875, p<0.0001, Dunnett's multiple comparison). Similarly RNA *in situ* hybridization showed significant cell-specific increase of *c-Fos* in the ETCs (q(57)=2.687, p=0.0241, Dunnett's multiple comparison) and GCs (q(57)=3.359, p=0.0036, Dunnett's multiple comparison) upon treatment. An increase was also observed in *c-Fos* expression in MCs, but was not significant (q(57)=2.365, p=0.0561). Note, however that not all glomeruli are affected by amyl acetate[36]. The increase in *c-Fos* expression was followed by an increase in *Mitf* expression 30 minutes after exposure to amyl acetate both in the ETCs (Figure 3-2, Figure 3f) (q(45)=3.325, p=0.005, Dunnett's multiple comparison) and the MCs (q(45)=5.031, p=0.0001, Dunnett's multiple comparison). Expression of *Kcnd3* also increased 30 minutes after exposure, but only in the MCs (Figure 3-3, Figure 3g; q(30)=3.271, p=0.0075, Dunnet's multiple comparison) and not in the ETCs. This showed that both *Mitf* and *Kcnd3* expression are activity-dependent in the MC projection neurons of the OB.

Activity-dependent increase in *Kcnd3* expression requires MITF enhancer activity.
Induction of large-scale OB activity with amyl acetate was also performed on *Mitf* mutant mice. The increase in *Kcnd3* expression observed upon inducing neuronal activity in wild type OBs

was not observed in *Mitf* mutant OBs (Figure 4a, b). Hence, *Mitf* and *Kcnd3* are both regulated

in an activity-dependent manner in the MC and the expression of *Kcnd3* depends on *Mitf* in

both MC and ETC. Analysis of two MITF ChIPseq data sets [32, 34] showed that MITF binds

to intronic regions of *KCND3* in 501mel melanoma cells. The two MITF ChIPseq datasets show

the same overlapping binding peak A (Figure 4c, yellow). A second MITF binding peak, peak

B, was observed only in the Laurette dataset and overlaps with H3K27ac and H3K4me1

ChIPseq peaks in 501mel melanoma cells [37] indicative of an active enhancer site (Figure 4c,

orange). Using wild-type MITF and a transcriptionally inactive mutant (Figure 4d),

transcription activation analysis showed that MITF activates *Kcnd3* transcription from the

region containing peak B in both HEK293T cells (Figure 4e; MITF-pCMV vs pCMV:

q(12)=22.58, p=0.0001; MITF-pCMV vs MITFB4RA-pCMV: q(12)=20.78, p=0.0001,

Dunnett's multiple comparison) and N2A cells (Figure 4f; MITF-pCMV vs pCMV:

q(12)=15.39, p=0.0001; MITF-pCMV vs MITFB4RA-pCMV: q(12)=15.18, p=0.0001,

Dunnett's multiple comparison), whereas transcriptionally inactive MITF does not (Fig. 4 e,f).

Tyr was used as a positive control and was significantly activated in both cell types. No

activation was observed in either cell type from a fragment containing peak A. As the region

around peak B has active enhancer marks, we conclude that MITF signaling is induced by

activity and regulates an enhancer region in an intron of *Kcnd3* leading to activity-dependent

increase in *Kcnd3* expression in projection neurons of the OB.

***Mitf* regulates olfactory discriminatory ability and odor fatigue recovery.**

Hyperactive MCs are likely to affect olfaction. Changes in the olfactory neuronal circuits

frequently lead to reduced ability to detect odorants or to differentiate between them. Both are

critical functions of the olfactory system. Olfactory testing, however, showed no difference in

the ability of *Mitf*^mi-vga9/mi-vga9^ and wild-type mice to detect or avoid odours (Figure 5a, b, Figure

5- SI Table 1). Next, we tested the ability of *Mitf*^mi-vga9/mi-vga9^ mice to differentiate between

odourants. Repeated introduction of the same odourant leads to reduced interest (habituation) and when followed by a novel odourant, renewed interest (dishabituation) (Figure 5c) - an established measurement of olfactory discriminatory ability [38]. In this assay, the $Mitf^{mi\text{-}vga9/mi\text{-}vga9}$ mice showed increased dishabituation when exposed to the chemically similar odourant pair vanilla-almond (Figure 5d; t(84)=5.223, p<0.0001, Dunnett's multiple comparison). This was also observed when the odourants were introduced in the reverse order (Figure 5e; t(126)=5.838, p=0.0001, Dunnett's multiple comparison) and when the less chemically similar odourant pair lemon-vanilla was used (Figure 5f; t(147)=2.708, p=0.0144, Dunnett's multiple comparison). This demonstrates increased olfactory discriminatory ability or sensitivity in the $Mitf$ mutant mice. Importantly, $Mitf^{mi\text{-}vga9/+}$ heterozygotes, which are phenotypically similar to wild-type mice, show an intermediate phenotype (Figure 5d-f). This demonstrates that the increased olfactory discrimination of the $Mitf^{mi\text{-}vga9/mi\text{-}vga9}$ mice is not due to larger cortical area devoted to olfactory processing caused by their blindness and deafness. Odour exposure reduces sensitivity of M/T neurons [14] and the appropriate long-term adaptation to sensory input is fundamental to olfaction. Our results suggested that the mutant mice might not adapt well on a longer time-scale, as $Mitf$ dependent transcription would be required. We therefore tested the effects of a long-term exposure of an odourant on olfactory ability, also termed odour fatigue. Under normal circumstances this leads to a reduction in ability to detect the odorant, followed by recovery once the odorant is removed. However, in the mutant mice, there was a stark reduction in of the ability to detect the odour as 6 out of 8 mutant mice tested did not detect the odor when reintroduced (Figure 5g; t(84)=3, p=0.0246, Sidak's multiple comparison).

We propose a model for a functional role of $Mitf$ in projection neurons of the olfactory bulb, where it regulates the intrinsic excitability of the neurons following activity induction, by reducing the neuronal firing rate of neurons through its regulation of $Kcnd3$ expression. In our

model, the activity of OSNs leads to an increase in *Mitf* expression, and an *Mitf*-dependent increase in the transcription of *Kcnd3* in the output neurons of the OB. An increased I_A-current, in turn, decreases the likelihood of the generation of action potential and reduces activity leading to decreased sensitivity of the glomeruli in question (Figure 6). Conversely, lack of activity in a glomerulus would lead to the opposite mechanism and increased sensitivity [16].

Discussion

The transcription factor MITF is expressed specifically in the excitatory neurons of the OB [18] and here we show that it plays a key role in activity-dependent changes in neuronal excitability. Neuronal activity sculpts the nervous system and is required for adaptive changes in neuronal function. To do so, neuronal activity must functionally change the neuron, by impacting its firing frequency or the strength of individual synapses. However, neurons also need to tightly regulate their own activity: too much neuronal activity is costly and can lead to excitotoxicity or uncontrolled firing, whereas the loss of neuronal activity leads to loss of information and neurodegeneration. To avoid this, mechanisms are in place to maintain neuronal activity within a set window, and the ability of a neuron to be plastic has to be within the confines of such neuronal homeostasis [3, 39, 40]. Our results show how sensory activity can affect firing rate of the olfactory bulb projection neurons in an *Mitf* dependent manner due to the effect on A-type currents via *Kcnd3* expression. This mechanism can maintain neuronal homeostasis or it can function to determine the sensitivity or neuronal gain-control of the projection neurons on a longer time scale.

The fact that the expression of both *Mitf* and *Kcnd3* in M/T are activity-dependent in the wild-type mouse suggested a direct link between MITF and *Kcnd3*, especially as *Kcnd3* expression following activity induction was not increased in the *Mitf* mutant mice. This suggested to us that *Kcnd3* might be a direct MITF target gene. Previously published ChIPseq data shows that MITF can bind to the *Kcnd3* locus. Furthermore, transcription activation experiments showed that MITF transcriptionally activates expression from an intronic enhancer element of *Kcnd3*. This implies that while MITF may not be required for *Kcnd3* expression *per se*, it is required for the activity-dependent increase in *Kcnd3* expression upon activity induction in mitral neurons. In accordance with this, stimulus dependent enhancer activity has previously been

described in mouse cortical neurons, possibly as a transcription activation mechanism upon induction of neuronal activity[9, 41-43].

ETCs drive olfactory output [44] and link OSN with M/T neurons [45], while M/T neurons are the projection neurons of the OB. Expression of *C-Fos*, a marker of neuronal activity, was increased in MCs of *Mitf^{mi-vga9}* mutant OBs but not in ETCs under baseline conditions. Interestingly, while we did observe an increase in *Mitf* and *Kcnd3* expression in the MCs, and the expression of both *c-Fos* and *Mitf* were increased upon activity in the ETC of wild type mice, *Kcnd3* was not. Additionally, increased activity of putative ETC inhibiting cells was also detected. This demonstrates that complex inhibitory and compensatory mechanisms are at work within the glomeruli. This also demonstrates that these two cell types, which have different functions in the olfactory neuronal circuit, also have a different intrinsic response to activity.

As MITF might play a role in the differentiation of M/T neurons we examined the number of excitatory neurons in the adult *Mitf* mutant OB and showed that the number of glomerular excitatory neurons was increased by around 30% in the *Mitf* mutant mice OB. The same was true for glomerular inhibitory neurons. Hence M/T neurons are generated in the *Mitf* mutant mouse and express the appropriate neuronal markers. The cause of the increase in glomerular excitatory neuron number is most likely due to the hyperactivity of the M/T neurons. which should have similar effects on the ontogeny of the OB as a very diverse olfactory environment would [46, 47]. It is unlikely that *Mitf* affects the number of inhibitory neurons directly as the lineage of excitatory neurons and inhibitory neurons in the OB are separate [48]. Calbindin+ and PV+ neurons showed a similar increase in numbers, whereas Calretinin+ interneurons did not. Interestingly, Calretinin+ interneurons appear to be less affected by activity within the

glomerular circuit [49], offering an explanation as to why this neuronal subtype was not increased in numbers. While many studies have shown that activity affects the number of excitatory neurons [47, 50, 51], and that at the synaptic level the balance between excitatory and inhibitory neurons is pivotal for OB function [52-54], our study is the first to demonstrate that the number of excitatory and inhibitory cells are balanced in the OB upon increase in neuronal activity of excitatory neurons.

The olfactory phenotype of the *Mitf* mutant mice is interesting – they show increased ability to discriminate between odours or increased olfactory sensitivity. This is a rarely reported characteristic of mutant mice [55], as the expectation is that olfaction works at maximum discriminatory power, since this is most beneficial for the organism. One explanation, at least in part, for this change in discriminatory ability could be the increased number of glomerular excitatory neurons and inhibitory interneurons of the OB. Apart from that, the nature of the olfactory phenotype in the *Mitf*$^{mi-vga9/mi-vga9}$ mouse is likely due to increased levels of lateral inhibition due to increased activity [56, 57] and/or more glomeruli being activated by each odourant, leading to reduced cross-habituation [58]. Further studies will determine this. Importantly, while the *Mitf* mutant mouse has an increase in the ability to differentiate between odours, long-term odour exposure leads to a reduction in the ability to detect that odour. This shows that *Mitf* is required for normal long-term adaptation in the OB and that the increase in olfactory discriminatory ability observed in the mutant mice comes at a cost.

Sensory systems focus on detecting novelty or change. In olfaction, this translates into the ability to quickly distinguish rare but important odours from background odours [14, 16]. In the OB, *Mitf* is pivotal to the adaptation of individual glomeruli through regulating intrinsic homeostatic plasticity in an activity-dependent manner. Most glomeruli are activated rarely. However, they play important roles in detecting rare threats or food sources, both of high

survival value. Thus, regulation of intrinsic plasticity, as described here, provides a mechanism through which olfaction can be insensitive to background odours but sensitive to rare odours. Keeping these glomeruli sensitive is therefore of high adaptive value and our model provides a mechanism through which this can occur.

Materials & Methods

For full details, see SI Appendix.

References:

1.   Davis, G.W., *Homeostatic control of neural activity: From phenomenology to molecular design.* Annual Review of Neuroscience, 2006. **29**: p. 307-323.
2.   Turrigiano, G., *Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement.* Annu Rev Neurosci, 2011. **34**: p. 89-103.
3.   Turrigiano, G.G., *The self-tuning neuron: synaptic scaling of excitatory synapses.* Cell, 2008. **135**(3): p. 422-35.
4.   Lemasson, G., E. Marder, and L.F. Abbott, *Activity-Dependent Regulation of Conductances in Model Neurons.* Science, 1993. **259**(5103): p. 1915-1917.
5.   Sweatt, J.D., *Neural plasticity and behavior - sixty years of conceptual advances.* J Neurochem, 2016. **139 Suppl 2**: p. 179-199.
6.   Lisman, J., et al., *Memory formation depends on both synapse-specific modifications of synaptic strength and cell-specific increases in excitability.* Nature Neuroscience, 2018. **21**(3): p. 309-314.
7.   Kourrich, S., D.J. Calu, and A. Bonci, *Intrinsic plasticity: an emerging player in addiction.* Nat Rev Neurosci, 2015. **16**(3): p. 173-84.
8.   Hrvatin, S., et al., *Single-cell analysis of experience-dependent transcriptomic states in the mouse visual cortex.* Nature Neuroscience, 2018. **21**(1): p. 120-+.
9.   Guzman-Karlsson, M.C., et al., *Transcriptional and epigenetic regulation of Hebbian and non-Hebbian plasticity.* Neuropharmacology, 2014. **80**: p. 3-17.
10.   Kikuta, S., et al., *Odorant response properties of individual neurons in an olfactory glomerular module.* Neuron, 2013. **77**(6): p. 1122-35.
11.   Tan, J., et al., *Odor information processing by the olfactory bulb analyzed in gene-targeted mice.* Neuron, 2010. **65**(6): p. 912-26.
12.   Roland, B., et al., *Massive normalization of olfactory bulb output in mice with a 'monoclonal nose'.* Elife, 2016. **5**.
13.   Chaudhury, D., et al., *Olfactory bulb habituation to odor stimuli.* Behav Neurosci, 2010. **124**(4): p. 490-9.

14.   Kato, H.K., et al., *Dynamic sensory representations in the olfactory bulb: modulation by wakefulness and experience.* Neuron, 2012. **76**(5): p. 962-75.

15.   Angelo, K., et al., *A biophysical signature of network affiliation and sensory processing in mitral cells.* Nature, 2012. **488**(7411): p. 375-8.

16.   Tyler, W.J., et al., *Experience-dependent modification of primary sensory synapses in the mammalian olfactory bulb.* J Neurosci, 2007. **27**(35): p. 9427-38.

17.   Tachibana, M., et al., *Ectopic expression of MITF, a gene for Waardenburg syndrome type 2, converts fibroblasts to cells with melanocyte characteristics.* Nat Genet, 1996. **14**(1): p. 50-4.

18.   Ohba, K., et al., *Microphthalmia-associated transcription factor is expressed in projection neurons of the mouse olfactory bulb.* Genes to Cells, 2015. **20**(12): p. 1088-1102.

19.   De Saint Jan, D., et al., *External tufted cells drive the output of olfactory bulb glomeruli.* J Neurosci, 2009. **29**(7): p. 2043-52.

20.   Hayar, A., et al., *Olfactory bulb glomeruli: External tufted cells intrinsically burst at theta frequency and are entrained by patterned olfactory input.* Journal of Neuroscience, 2004. **24**(5): p. 1190-1199.

21.   Whitesell, J.D., et al., *Interglomerular lateral inhibition targeted on external tufted cells in the olfactory bulb.* J Neurosci, 2013. **33**(4): p. 1552-63.

22.   Wang, F., et al., *RNAscope: a novel in situ RNA analysis platform for formalin-fixed, paraffin-embedded tissues.* J Mol Diagn, 2012. **14**(1): p. 22-9.

23.   Hodgkinson, C.A., et al., *Mutations at the Mouse Microphthalmia Locus Are Associated with Defects in a Gene Encoding a Novel Basic-Helix-Loop-Helix-Zipper Protein.* Cell, 1993. **74**(2): p. 395-404.

24.   Liu, N., et al., *Unique regulation of immediate early gene and tyrosine hydroxylase expression in the odor-deprived mouse olfactory bulb.* J Biol Chem, 1999. **274**(5): p. 3042-7.

25.   Imamura, F. and C.A. Greer, *Pax6 regulates Tbr1 and Tbr2 expressions in olfactory bulb mitral cells.* Molecular and Cellular Neuroscience, 2013. **54**: p. 58-70.

26.   Mizuguchi, R., et al., *Tbr2 Deficiency in Mitral and Tufted Cells Disrupts Excitatory-Inhibitory Balance of Neural Circuitry in the Mouse Olfactory Bulb (June 27, 8831, 2012).* Journal of Neuroscience, 2012. **32**(39): p. 13639-13639.

27.   Fransen, E. and J. Tigerholm, *Role of A-type potassium currents in excitability, network synchronicity, and epilepsy.* Hippocampus, 2010. **20**(7): p. 877-87.

28.   Yuan, W., A. Burkhalter, and J.M. Nerbonne, *Functional role of the fast transient outward K+ current IA in pyramidal neurons in (rat) primary visual cortex.* J Neurosci, 2005. **25**(40): p. 9185-94.

29.   Kim, J., et al., *Regulation of dendritic excitability by activity-dependent trafficking of the A-type K+ channel subunit Kv4.2 in hippocampal neurons.* Neuron, 2007. **54**(6): p. 933-47.

30.   Burton, S.D., et al., *Olfactory Bulb Deep Short-Axon Cells Mediate Widespread Inhibition of Tufted Cell Apical Dendrites.* Journal of Neuroscience, 2017. **37**(5): p. 1117-1138.

31.   Gutman, G.A., et al., *International Union of Pharmacology. LIII. Nomenclature and molecular relationships of voltage-gated potassium channels.* Pharmacol Rev, 2005. **57**(4): p. 473-508.

32.   Laurette, P., et al., *Transcription factor MITF and remodeller BRG1 define chromatin organisation at regulatory elements in melanoma cells.* Elife, 2015. **4**.

33.   Calero-Nieto, F.J., et al., *Key regulators control distinct transcriptional programmes in blood progenitor and mast cells.* EMBO J, 2014. **33**(11): p. 1212-26.

34.   Strub, T., et al., *Essential role of microphthalmia transcription factor for DNA replication, mitosis and genomic stability in melanoma.* Oncogene, 2011. **30**(20): p. 2319-32.

35.   Shoag, J., et al., *PGC-1 coactivators regulate MITF and the tanning response.* Mol Cell, 2013. **49**(1): p. 145-57.

36.   Bepari, A.K., et al., *Visualization of odor-induced neuronal activity by immediate early gene expression.* Bmc Neuroscience, 2012. **13**.

37.   Fontanals-Cirera, B., et al., *Harnessing BET Inhibitor Sensitivity Reveals AMIGO2 as a Melanoma Survival Gene.* Molecular Cell, 2017. **68**(4): p. 731-+.

38.   Yang, M. and J.N. Crawley, *Simple behavioral assessment of mouse olfaction.* Curr Protoc Neurosci, 2009. **Chapter 8**: p. Unit 8 24.

39.   Titley, H.K., N. Brunel, and C. Hansel, *Toward a Neurocentric View of Learning.* Neuron, 2017. **95**(1): p. 19-32.

40.   Pozo, K. and Y. Goda, *Unraveling mechanisms of homeostatic synaptic plasticity.* Neuron, 2010. **66**(3): p. 337-51.

41.   Walke, W., G. Xiao, and D. Goldman, *Identification and characterization of a 47 base pair activity-dependent enhancer of the rat nicotinic acetylcholine receptor delta-subunit promoter.* J Neurosci, 1996. **16**(11): p. 3641-51.

42.   Kim, T.K., et al., *Widespread transcription at neuronal activity-regulated enhancers.* Nature, 2010. **465**(7295): p. 182-U65.

43.   Malik, A.N., et al., *Genome-wide identification and characterization of functional neuronal activity-dependent enhancers.* Nature Neuroscience, 2014. **17**(10): p. 1330-1339.

44.   Hayar, A., et al., *External tufted cells: a major excitatory element that coordinates glomerular activity.* J Neurosci, 2004. **24**(30): p. 6676-85.

45.   Gire, D.H., et al., *Mitral Cells in the Olfactory Bulb Are Mainly Excited through a Multistep Signaling Path.* Journal of Neuroscience, 2012. **32**(9): p. 2964-2975.

46.   Johnson, M.C., et al., *Odor enrichment sculpts the abundance of olfactory bulb mitral cells.* Neurosci Lett, 2013. **541**: p. 173-8.

47.   Liu, A., S. Savya, and N.N. Urban, *Early Odorant Exposure Increases the Number of Mitral and Tufted Cells Associated with a Single Glomerulus.* J Neurosci, 2016. **36**(46): p. 11646-11653.

48.   Wichterle, H., et al., *In utero fate mapping reveals distinct migratory pathways and fates of neurons born in the mammalian basal forebrain.* Development, 2001. **128**(19): p. 3759-3771.

49.   Fogli Iseppe, A., A. Pignatelli, and O. Belluzzi, *Calretinin-Periglomerular Interneurons in Mice Olfactory Bulb: Cells of Few Words.* Front Cell Neurosci, 2016. **10**: p. 231.

50.   Corotto, F.S., J.R. Henegar, and J.A. Maruniak, *Odor deprivation leads to reduced neurogenesis and reduced neuronal survival in the olfactory bulb of the adult mouse.* Neuroscience, 1994. **61**(4): p. 739-44.

51.   Mandairon, N., et al., *Long-term fate and distribution of newborn cells in the adult mouse olfactory bulb: Influences of olfactory deprivation.* Neuroscience, 2006. **141**(1): p. 443-51.

52.   Abraham, N.M., et al., *Synaptic inhibition in the olfactory bulb accelerates odor discrimination in mice.* Neuron, 2010. **65**(3): p. 399-411.

53.   Takahashi, H., et al., *A Subtype of Olfactory Bulb Interneurons Is Required for Odor Detection and Discrimination Behaviors.* J Neurosci, 2016. **36**(31): p. 8210-27.

54.   Zou, J., et al., *Targeted deletion of ERK5 MAP kinase in the developing nervous system impairs development of GABAergic interneurons in the main olfactory bulb and behavioral discrimination between structurally similar odorants.* J Neurosci, 2012. **32**(12): p. 4118-32.

55.   Fadool, D.A., et al., *Kv1.3 channel gene-targeted deletion produces "Super-Smeller Mice" with altered glomeruli, interacting scaffolding proteins, and biophysics.* Neuron, 2004. **41**(3): p. 389-404.

56.   Nunez-Parra, A., et al., *Disruption of centrifugal inhibition to olfactory bulb granule cells impairs olfactory discrimination.* Proceedings of the National Academy of Sciences of the United States of America, 2013. **110**(36): p. 14777-14782.

57.   Nunes, D. and T. Kuner, *Disinhibition of olfactory bulb granule cells accelerates odour discrimination in mice.* Nature Communications, 2015. **6**.

58.   Mombaerts, P., et al., *Visualizing an olfactory sensory map.* Cell, 1996. **87**(4): p. 675-86.

Figure legends

Figure 1: *Mitf^{mi-vga9/mi-vga9}* mice have an increase in inhibitory and excititatory neuronal subtypes in the glomerular layer. a. RNA *in situ* hybridization of *Mitf* (red) and *Tbr2* (green) in glomeruli and MC/GCL of wild-type and *Mitf^{mi-vga9/mi-vga9}* mice. Scale bars are 50μm. **b.** *Mitf* count per *Tbr2*-positive cells below glomeruli and MCL. N=9 per genotype. **c**. *Mitf* mRNA expression, determined by RT-qPCR. N=6 per genotype. **d.** Immunofluorescent staining of Tbr2/Eomes (Tbr2+) neurons (green) in the OBs of wild-type, *Mitf^{mi-vga9/+}* and *Mitf^{mi-vga9/mi-vga9}* mice. **e**. Cell count of Tbr2/Eomes (Tbr2+) cells in the glomeruli and MCL of wild-type, *Mitf^{mi-vga9/+}* and *Mitf^{mi-vga9/mi-vga9}* mice. N=6 per genotype. **f**. Immunofluorescent staining of Calbindin (Calb+) neurons (green) in the OBs of wild-type *Mitf^{mi-vga9/+}* and *Mitf^{mi-vga9/mi-vga9}* mice **g.** Cell count of Calbindin (Calb+) cells in the glomeruli of wild-type, *Mitf^{mi-vga9/+}* and *Mitf^{mi-vga9/mi-vga9}* mice. N=6 per genotype. **h**. Immunofluorescent staining of Parvalbumin (Pv+) neurons (green) in the OBs of wild-type, *Mitf^{mi-vga9/+}* and *Mitf^{mi-vga9/mi-vga9}* mice. **i**. Cell count of Parvalbumin (Pv+) cells in the glomeruli, EPL and MC/GCL of the medial OBs of wild-type, *Mitf^{mi-vga9/+}* and *Mitf^{mi-vga9/mi-vga9}* mice. N=6 per genotype. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bars are 100μm unless otherwise indicated. P-values were calculated using two-way ANOVA (**b,e,i**), two-tailed unpaired student's t-test (**c**) or one-way ANOVA (**g**). *= $P<0.05$, ***$P<0.001$. MC= mitral cells, MCL=mitral cell layer, EPL= external plexiform layer.

Figure 2: *Mitf^{mi-vgag/mi-vga9}* mice have a decrease in I_A and a concomittant increase in mitral cell activity. a. Representative images of voltage clamp recordings. **b**. Representative images of voltage clamp recordings, where the I_A was inactivated by a -40mV pre-pulse. **c**. The isolated I_A recorded from wild-type and *Mitf^{mi-vga9/mi-vga9}* M/T neurons. **d**. $I_{A\ max}$ and $I_{DR\ max}$ in wild-type and *Mitf^{mi-vga9/mi-vga9}*. N=12 per genotype. **e**. The relation between current density (pA/pF) and

membrane voltage (mV) of wild-type and *Mitf^mi-vga9/mi-vga9* M/T neurons. N=12 per genotype. **f**. Representative recordings of spontaneous action potentials observed in wild-type (N=5) and *Mitf^mi-vga9/mi-vga9* M/T neurons (N=3) by current clamp. **g**. RNA *in situ* hybridization of *c-Fos* (green) and *Mitf* (red) in wild-type and *Mitf^mi-vga9/mi-vga9* OBs. **h**. *c-Fos* dots per cell in the ETC, MC and GC. N=4 per genotype. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bars are 50µm. P-values were calculated using two-tailed unpaired student's t-test (**d**) and two-way ANOVA (**h**) and. *P<0.05. ETC= external tufted cells, MC= mitral cells, GC= granule cells.

Figure 3: *Kcnd3* and *Mitf* expression are activity-dependent in the OB. a. RNA *in situ* hybridization of *Tbr2* (green), *Mitf* (red) and *Kcnd3* (magenta) performed on wild-type and *Mitf^mi-vga9/mi-vga9* OBs. **b**. *Kcnd3* count per cell. N=5 per genotype. **c.** Schematic representation of the amyl acetate (AA) experiment where mice were habituated to an odourless cage for 60 minutes. Upon exposure to AA for 30 minutes (red block), they were sacrificed at different time points (black vertical lines) **d.** *Mitf* and *c-Fos* mRNA expression in wild-type OB following AA, determined by RT-qPCR. N=6 per time point. **e.** *c-Fos* dots per cell in wild-type OB mice following AA. N=6 per time point. **f.** *Mitf* dots per cell in wild-type OB following AA. N=6 per time point. **g.** *Kcnd3* dots per cell in wild-type OB following AA. N=6 per time point. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bars are 50µm. P-values were calculated using two-way ANOVA (**b,d,f,g,i**). *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. PG= periglomerular.

Figure 4 Activity-dependent increase in *Kcnd3* expression requires MITF enhancer activity.a. RNA *in situ* hybridization of *c-Fos* (red) and *Kcnd3* (green) performed on *Mitf^mi-vga9/mi-vga9* OB following AA. **b**. *Kcnd3* dots per cell in *Mitf^mi-vga9/mi-vga9* OB following AA. N=3-5 per time point. **c.** ChIPseq peaks of MITF, H3K27ac and H3K4me1 on *KCND3* gene in

501mel cells. The MITF peaks are labeled A (orange) and B (yellow). Yellow shows overlapping peak of MITF binding in Strub and Laurette datasets (B) whereas Orange indicates MITF peaks that overlap with H3K27ac and H3K4me1 peaks (A). **d.** Sequence of the basic region of wild-type (MITF-M) and transcriptionally inactive MITF with four argenines mutated to alanines (MITF-M B4RA). **e.** Transcription activation assays performed in HEK293T cells co-transfected with constructs containing the hTYR, hKCND3_pA, and hKCND3_pB regulatory regions fused to luciferase, together with empty vector or wild-type or transcriptionally inactive (B4RA) MITF constructs. N=3. **f**. Transcription activation assays performed in N2A cells co-transfected with constructs containing the hTYR, hKCND3_pA, and hKCND3_pB regulatory regions fused to luciferase, together with empty vector or wild-type or transcriptionally inactive (B4RA) MITF constructs. N=3. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bars are 50μm. P-values were calculated using two-way ANOVA (**b, e, f**) *P<0.05, ****P<0.0001.

Figure 5: *Mitf* **regulates olfactory discriminatory ability and odor fatigue recovery. a.** Quantification of the results of the hidden cereal odour detection-assay. N=8 per genotype **b.** Avoidance assay where a cage was sectioned in 3 equal areas and mice were exposed to water or propionic acid (PA) in section 1. Quantification shows time spent in each section. SI Table 3. N=6-8 per genotype. **c**. Schematic overview of the dishabituation-habituation assay, where mice are exposed to odour A (red) for 30 seconds 6 times with 5-minute intervals, after which they are exposed to odour B (green) for 30 seconds. **d** Dishabituation-habituation, mice were exposed to vanilla as odour A and almond as odour B. N=8 per genotype. **e**. Dishabituation-habituation, where mice were exposed to almond as odour A and vanilla as odour B. N=6-8 per genotype. **f**. Dishabituation-habituation, mice were exposed to almond as odour A and lime as odour B. N=8 per genotype. **g.** Dishabituation-habituation assay where the mice were exposed

to vanilla for two hours, followed by almond as odour A and vanilla as odour B. N=15 per genotype. The values on the graphs are mean $\pm$ SEM. P-values were calculated using one-way ANOVA (**a**) or two-way ANOVA (**b, d-g**). *P<0.05, **P<0.01****P<0.0001.

Figure 6: Model of MITF as a regulator of homeostatic intrinsic plasticity. Upon neuronal activity in the OB, MITF is required for an increase in *Kcnd3* expression. More Kv4.3 protein results in an increase in the I_A potassium current and reduced likelihood of action potentials, subsequently leading to decreased activity of M/T cells.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Information

Animals

All animal procedures were approved by the Committee on Experimental Animals, according to regulation 460/2017 and European Union Directive 2010/63 (license number 2013-03-01). Mice of the genotypes *C57BL/6J* (RRID:IMSR_JAX:000664, wild-type), *C57BL/6J-Mitf^{mi-vga9/mi-vga9}* (RRID:MGI:4455017, *Mitf^{mi-vga9/mi-vga9}*) and *C57BL/6J-Mitf^{mi-vga9/+}* (*Mitf^{mi-vga9/+}*) were maintained at the mouse facility at the University of Iceland. Unless otherwise specified, mice were housed two to five per per cage in a temperature-controlled environment (21-22°C). Unless otherwise specified, mice consumed water and food *ad libitum*. Mice of P75-P95 of both genders were used for all experiments.

Electrophysiology

Primary OB neuronal cultures from P0-P2 mice were obtained as described by Lutz et al., (2007) with slight modifications. Briefly, mice were decapitated and submerged in ethanol, upon which the head was placed in Ca^{2+} and Mg^{2+}-free HBSS (Gibco) containing 1% Penicillin/Streptomycin (Gibco) and 1% Amphoterecin B (Gibco) under a dissecting microscope. OBs were dissected from the heads, the meninges removed, then trypsinized for 7-10 minutes at 37°C. The cells were washed with 0.05% trypsin-EDTA prior to incubation in the trypsin solution. The neurons were then gently resuspended in Neurobasal-A Medium (Gibco) supplemented with 1% GlutaMAX (Gibco) 1% Penicillin/Streptomycin, 1% Amphoterecin B, 2% 50x serum-free B27 Supplement (Gibco) and 10% FBS. The cells were diluted to a density of $1x10^6$cells·mL^{-1} and plated at $1.25x10^5$cells·cm^2 on 0.1mg/mL Poly-D-Lysine hydrobromide (Sigma-Aldrich) and 0.075mg·mL^{-1} PureCol Purfied bovine Collagen

Solution, Type 1 (Advanced BioMatrix) pre-coated coverslips in Nunclon Delta Surface 24-well plates (Thermo Scientific). Neurons were cultured in Neurobasal-A Medium supplemented with 1% GlutaMAX 1% Penicillin/Streptomycin, 1% Gibco Amphoterecin B and 2% 50x serum-free B27 Supplement for 2 days, upon which they were placed in GlutaMAX-free Medium, which was replenished every 3 days.

The mean DIV was 12 +/- 3 days. The cultured cells were maintained in a solution with NBM-B27, 1% GlutaMAX until DIV = 2, and subsequently in NBM-B27 without GlutaMAX. The medium was replaced every 2-3 days. The osmolarity of the solutions was measured with an osmometer and the pipette solution was 319 mOsm, while the extracellular Krebs solution was adjusted to 312 mOsm. Isolated cells on coverslips were placed in a recording chamber, and kept at room temperature. The cells were continuously superfused during recordings with an oxygenated extracellular solution. The extracellular mammalian Krebs solution contained the following (all in mM): 150 NaCl, 5 KCl, 2.6 $CaCl_2$, 2 $MgCl_2$, 10 HEPES, 5 glucose, at pH 7.3. The solution was filtered with a 0.2 µm filter prior to use. Pipettes were pulled out of 1.5mm O.D. borosilicate glass capillary tubings (Science Products GmbH) with a two-stage Narishige PP83 horizontal puller (Narishige Co), and then fire-polished to about 1 µm tip diameter in a Narishige MF-83 microforge. Pipette resistances were 2-6MΩ when placed in the Krebs solution while filled with the internal solution. The Internal solution contained the following (all in mM): 145 KCl, 2 $MgCl_2$, 10 EGTA, 10 HEPES, 0.2 NaATP, pH 7.3. The reference Ag/AgCl electrode was placed in a compartment separated from the recording chamber, and electrically linked to the superfused compartment containing the cells with a glass tube bridge filled with 2M KCl/Agar solution. Recordings were made using the whole-cell configuration of the patch-clamp technique (Hamill, et al., 1981) on M/T cells with large nuclei, with an Axopatch 1D patch clamp amplifier (Axon Instruments Inc) in voltage or current clamp mode. The mean membrane resistance of wild-type M/T neurons was 104±20.3 MOhm, and 98±17

MOhm (t(25)= 0.234, p=0.82, Two-tailed test) in *Mitf^{mi-vga9/mi-vga9}* neurons. The mean series resistance of wild-type M/T neurons was 13±2.3 MOhm and 17±1.6 MOhm (t(25)= -1.415, p=0.17, Two-tailed test) in *Mitf^{mi-vga9/mi-vga9}* neurons. The mean cell capacitance of wild-type neurons was 46±7.3 pF and that of *Mitf^{mi-vga9/mi-vga9}* neurons was 56±11 pF (t(25)= -0.7308, p=0.47, Two-tailed t-test). The mean resting membrane potentials of wild-type M/T neurons were -58±2 mV and of *Mitf^{mi-vga9/mi-vga9}* neurons-57±2 mV, which is not signficantly different (t(11)=-0.102, p=0.92, Two-tailed t-test). The measured values of all these biophysical parameters of the cultured cells are comparable to previously published values measured from cultured mouse olfactory bulb M/T cells [28, 29]. The output of the amplifier was low-pass filtered at 2KHz and digitized at 10KHz with a Digidata 1440A (Molecular Devices) 16-bit A/D-D/A converter. Acquisition of recordings and generation of voltage or current pulses were performed with the pClamp 10.0 software (Molecular Devices) in conjunction with the Axopatch 1D amplifier. No corrections were made for liquid junction potentials or leak currents. After a break-in under voltage clamp, the cell was clamped at a holding potential of -70 mV, and then a 5 mV hyperpolarizing voltage step of 50 msec in duration was applied to evoke capacitance transients in the whole-cell current, to estimate the series resistance (Rs), input resistance (Ri), membrane resistance (Rm) and capacitance (Cm) of the cell. Subsequently, the cultured cells were then first voltage clamped at a holding potential of -70mV, and then stepped to -90mV for 80msec. Whole-cell outward currents were then evoked by a protocol using successive depolarizing voltage pulses from -90mV, 300msec in duration at +5 mV steps. To separate the A-type current from other outward currents another protocol with a series of voltage pulses 300 msec in duration was then applied to the cell, except that before the depolarizing pulses were delivered, the cell was held at -40mV for 80msec, after having been kept hyperpolarized for 55msec at -90mV. In order to isolate the A-type current from the total outward current, the current traces evoked by the second protocol, which

primarily involves delayed rectifier K^+ currents, were subtracted from the corresponding traces evoked by the first protocol. The resting membrane potential was recorded immediately after break in and turning to whole-cell current clamp mode with current at zero. Spontaneous membrane potential fluctuations and spiking activity of projection neurons were measured in whole-cell current clamp mode, while holding the cell near the resting potential (-65±5mV).

Immunofluorescence

Mice were transcardially perfused (Experimental license number 2014-07-02) with 1x Phosphate buffer solution (PBS; Dulbecco) followed by 4% paraformaldehyde (PFA; Sigma Aldrich) in 1xPBS. Following dissection, the brain was post-fixed for 2 hours in 4% PFA at room temperature. The OBs were sectioned at -20°C, in 20µm thin sections and kept at 4°C until used for immunofluorescence. Sections were placed in a blocking buffer of 0.1% Triton X-114 (Sigma Aldrich) and 5% Normal Goat Serum (NGS; Gibco) in PBS for 1 hour at room temperature. Blocking was followed by incubation with primary antibody (diluted 1:500) in blocking buffer overnight at 4°C. The antibodies used were TBR2/Eomes (Abcam, Ab23345, RRID:AB_778267), Parvalbumin (Abcam, Ab11427, RRID:AB_298032), Calbindin (Millipore, Ab1778, RRID:AB_2068336), Calretinin (Abcam, Ab702, RRID:AB_305702) and Tyrosine hydroxylase (Millipore, MAB318, RRID:AB_2313764). Sections were subsequently incubated for 1 hour at room temperature in blocking buffer containing the Alexa Fluor goat anti-mouse 488 or Alexa Fluor goat anti-rabbit 546 (Life Technologies) secondary antibodies (each at 1:1000 dilution) and DAPI (1:1000; Sigma Aldrich). The sections were then washed and subsequently kept at 4°C until imaging at 20x magnification using an Olympus FV10-MCPSU confocal microscope; 2-5 images were obtained from the medial OB of each sample. Cells were counted manually. Quantification of the samples was performed under blind

condition. The average of cells per sample was calculated and plotted using GraphPad Prism 7.

TUNEL

Fixed frozen sections were obtained as described above. For TUNEL staining of the fixed frozen sections, In Situ Cell Death Detection Kit, Fluorescin Version 17 (Roche) was used. Images were obtained at 20x magnification using Z-stack 3D images. From each sample, 2-3 images were obtained from the medial OB. Using the Fiji software, images were stacked and positively stained cells were counted manually. The average number of positive cells per image was calculated using GraphPad Prism 7.

Histological analysis

Mice were sacrificed by cervical dislocation and the brain was post-fixed for 24 hours in 4% Formaldehyde (Sigma Aldrich). Following fixation, the OB was embedded in paraffin and sectioned in 4µm thin sections. The sections were stained for hematoxylin and eosin (H&E).

Plasmids

The MITF-M cDNA was cloned into the p3xFlag-CMV14 vector using EcoRI and BamHI restriction sites. The 4 arginines in the basic domain were mutated to alanines by in situ mutagenesis in order to generate the construct MITF-B4RA [30]. The human TYR promoter in pGL3 Basic Luciferase Reporter vector was used for co-transfection assays.

MITF binding sites in the intronic region of KCND3 were identified using the ChIPseq data from Laurette et al., 2015 (GSM1517751) and [31] (GSE64137). The data were imported into IGV Tools and binding peaks near KCND3 identified. Similarly, ChIPseq data for H3K4me1 (GSM2476344) and H3K27ac modifications in melanoma cells (GSM2476350) was was analyzed [32]. Two MITF peaks termed p2 and p4 were identified. In order to clone the KCND3_p2 region, a 655bp fragment was amplified from genomic DNA of the human 501mel melanoma cells using forward primer 5'-TTGTGAGAGTAGCAGAGTGCTTTGC-3' and reverse primer 5'-GAGCAGATTCAGAGATCAGAAATCAATGG-3'; the primers also contained restriction sites for KpnI and XhoI. In order to clone the KCND3-p4 region, a 469bp fragment was amplified from genomic DNA of 501mel melanoma cells using forward primer 5'-GCTTCTGGAAGGTGAGAGAAGGA-3' and reverse primer 5'-AGTGTCCTGATAGCCACATTAGGTC-3'; the primers also contained restriction sites for NheI and XhoI. Both fragments were subsequently cloned into the pGL3 Promoter reporter vector.

Cell culture

HEK293T (RRID:CVCL_0063) and N2A (RRID:CVCL_0470) neuronal cells were obtained from the Leibniz Institute DSMZ-Deutsche Sammlung von Mikroorganism un Zellkuturen GmbH. HEK293T cells were cultured in DMEM-Glutamax media (Gibco) supplemented with 10% Fetal bovine serum (FBS; Gibco), N2A cells were cultured in DMEM-Glutamax media supplemented with 10% FBS and 1% Non-Essential Amino Acids (Gibco). Cells were cultured in 75cm^2 Nunc EasYflask with Nunclon Delta surface (Thermo Scientific) in HERAcell 240i incubator (Thermo Scientific) at 37°C and 5% CO_2. HEK293T were passaged once 90%

confluency was reached, by gently trypsinizing the cells, resuspending them in media and diluting the cells 1:10 and replating them. N2A cells were passaged once 70% confluency was reached, by gently trypsinizing the cells, resuspending them in media and diluting the cells 1:5 and replating them.

Luciferase

N2A and HEK293T cells were plated in Falcon 96-well cell culture plates at 13.800 and 20.000 cells per well respectively. After 24 hours, the N2A and HEK293T cells were transfected with 0.1 µg of DNA mixture at 7:1 and 2.8:1 Fugene HD to DNA ratio, respectively. This mixture contained equal amounts of the luciferase reporter plasmid, renilla reporter plasmid and MITF-FlagpCMV, MITF-B4RAFlagpCMV, or FlagpCMV. Each combination was plated and transfected in triplicate. Cells were harvested for luciferase assays after 24 hours of incubation using Dual-Glo Luciferase Assay System (Promega Corporation) as described by their protocol and the luciferase and renilla luminescence measured separately on MODULUS II TURNER (Biosystems). To quantify the normalized luminescence, the luciferase and renilla luminescence values were divided by one another for each well. Subsequently an average was obtained for the triplicate. These values were then plotted using GraphPad Prism 7.

Odour exposure

Mice were placed in empty cages, devoid of bedding, food and water for one hour in an odour free room. After one hour, mice were moved to a different room where an open eppendorf tube containing 60µL of amyl acetate (Sigma-Aldrich) was taped inside their cage at nose-level. After 30 minutes, the mice were placed in a new cage without amyl acetate and moved back to the odour free room. The mice were sacrificed by cervical dislocation at different intervals:

before amyl acetate exposure, immediately after exposure and 30, 90, and 210 minutes after exposure to amyl acetate. The OBs were either flash frozen in liquid nitrogen for quantitative real time PCR or placed in OCT medium (Sakura) in a 15x15x15mm Tissue-Tek Cryomold (Sakura) and subsequently flash frozen for sectioning and RNA *in situ* hybridization.

RNA mFISH

Mice were sacrificed by cervical dislocation. The OB was dissected out and placed in OCT compound in a 15x15x15mm Tissue-Tek Cryomold and flash frozen with liquid nitrogen. The caudal OB was sequentially sectioned unto 11 slides, with 2 sections per slide, at a thickness of 20µm. The slides were stored in a -80°C freezer in an airtight bag until use. The samples were pre-treated using the protocol "RNAscope Sample Preparation and Pretreatment Guide for Fresh Frozen Tissue (Manual RNAScope assay)" (Advanced Cell Diagnostics). mFISH was subsequently performed according to the manufacturer's instruction for the RNAscope Fluorescent Multiplex Kit (Advanced Cell Diagnostics) with probes 422501 (*Mitf*), 429641-C2 (*Tbr2/Eomes*), 452598-C3 (*Kcnd3*) and 316921-C2 (*c-Fos*). Sections were kept at 4°C until imaging using Z-stack 3D images with the Olympus FluoView FV10 confocal microscope at 30x magnification. Two images were obtained from the medial OB per sample - one image of the glomerular region of the OB and a separate image of the mitral and granule cell layer. Using the Fiji software, fluorescent dots were counted. *Tbr2+* cells located directly below the glomeruli, were considered external tufted cells. Similarly, *Tbr2+* cells located in the mitral cell layer, and not in the external plexiform layer, were considered as mitral cells. In the cases where *Tbr2* coexpression was not determined, *Mitf* expression, localization and nuclear morphology was used to determine cell type. Average dots per cell was calculated for each sample and plotted using GraphPad Prism 7.

Quantitative PCR

Mice were sacrificed by cervical dislocation and OBs flash frozen in liquid nitrogen. The samples were kept at -80°C until RNA was isolated using NucleoSpin RNA (Machery Nagel). In order to generate cDNA, the SuperScript II Reverse Transcriptase kit (Invitrogen) was used, using 1 ug of RNA and oligo(T) primers. Mouse specific exon-spanning primers were designed using Primer-blast. The primers-pairs used were the following: *Gapdh* forward 5'-ATGACATCAAGAAGGTGGTG-3', reverse 5'-CATACCAGGAAATGAGCTTG-3'; *Actin* forward 5'-CACTGTCGAGTCGCGTCC-3', reverse 5'-TCATCCATGGCGAACTGGTG-3'; *Mitf* forward 5'-AGCAAGAGCATTGGCTAAAGA-3', reverse 5'-GCATGTCTGGATCATTTGACT-3' and *c-Fos* forward 5'-TTTCAACGCCGACTACGAGG-3', reverse 5'-TCTGCGCAAAAGTCCTGTGT-3'. Quantitative qPCR was performed using Power SYBR Green PCR Master Mix (ThermoFisher Scientific). The $2^{-\Delta\Delta Ct}$ method was used to normalize expression to *C57BL/6J,* relative to *Gapdh* and *Actin* as described previously [33].

Behavior

Prior to performing any behavior experiments, individual mice were placed in separate cages for at least 24 hours. Mice were used for a maximum of two behavioral experiments but only once for the same behavioral test. Each mouse was only used once for each habituation/dishabituation experiment. All behavior experiments were performed at 9pm, 1 hour after the lights were turned off. All behavior was filmed, and the behavior of interest measured as a function of time and plotted using GraphPad Prism 7.

Hidden food assay

The food pellets were replaced by sweetened cereal – Cocoa Puffs (Nesquik General Mills) for 12 hours. After 12 hours, all Cocoa Puffs pellets were removed from the food section of the cage and one Cocoa Puffs pellet was placed into the cage. Following overnight starvation (license number 2016-05-01), a cage was filled with 4cm of bedding and a Cocoa Puffs pellet was hidden in one corner, under the bedding. The mouse was placed in the posterior section of the cage, opposite to the pellet, and the time it took the mouse to find the pellet was determined. Mice which showed obvious distress (tremors, unsteady gait or unwillingness to explore) and could not find the pellet in under 180 seconds were removed from the experiment.

Avoidance assay

A 41.5 x 24 x 18.5cm cage was sectioned into three equal sections numbered 1-3 and the bottom covered with 0.5cm of bedding. Prior to the experiment, 30µL of water was placed onto a 5 x 5 cm piece of Whatman paper which was then taped onto section 1, at nose height. The mouse was gently placed in section 3 and allowed to roam for 120 seconds in the cage. After 120 seconds, the mouse was removed from the cage and the 5 x 5cm Whatman paper containing 30µL of water was replaced with a 5 x 5cm paper of Whatman paper containing 30µL of propionic acid (Sigma Aldrich). The mouse was gently placed in section 3 and allowed to roam for 120 seconds in the cage. After 120 seconds, the mouse was removed from the cage. The time the mouse spent in each of the 3 sections, using the nose as the reference point, was measured for both conditions.

Habituation-Dishabituation

In order to perform the habituation-dishabituation experiment, we used the protocol of Lehmkuhl and associates [34]. Mice were kept in individual cages for 24 hours. Food and water

were subsequently removed, and the mice were kept in an odour free room for 1 hour prior to the experiment. The mice were habituated for 30 seconds every 5 minutes by placing 5µL of a particular odour in the cage, a total of 6 times, and subsequently exposed to 5µL of another odour for 30 seconds. Time spent sniffing the object, or in the direction of the object was determined. Odourants used were almond, lime (brand name Dr Oetker) and vanilla extracts (brand name Katla). For long-term exposure to odour, the mice were exposed to vanilla odourant for 2 hours prior to habituation-dishabituation with almond-vanilla odour pairs.

Statistical analysis

No statistical methods were used to predetermine sample size. Grouped analyses were used for the *in vivo* experiments as at least three mice per genotype were used. Quantitative results were analysed by one- or two-way analyses of variance (ANOVA), and two-sided unpaired or paired Student's t-tests and Chi-square tests using GraphPad Prism 7. To obtain p-values, multiple comaparisons for ANOVA tests were performed with Livak's correction or Dunnett's correction in cases where there was a comparison to a control time point. All numerical results are presented as mean and SEM unless stated otherwise. Degrees of freedom are indicated between brackets.

Author Contributions:

D.A.M.A. took part in designing experiments with P.H.P. and E.S., performed all experiments and analyzed the data, aside from the electrophysiology experiments. A.P provided data. T.E and H.R. performed the electrophysiology experiments. P.H.P., E.S. and D.A.M.A wrote the manuscript and T.E. commented on the manuscript at all stages.

Supplementary Figures

Figure 1-1: *Mitf^(mi-vga9/mi-vga9)* **mice have normal olfactory bulbs. a**. Representative images of the H&E histological analysis of coronal sections of the medial olfactory bulbs of the indicated genotypes. **b.** Immunofluorescent staining of Tyrosine hydroxylase (TH) showing the localization of TH in the glomeruli of the indicated genotypes. **c.** Cell count of TH+ cells of the OB of wild-type, *Mitf^(mi-vga9/+)* and *Mitf^(mi-vga9/mi-vga9)* mice. N=6 per genotype. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bar indicates 100µm**.** P-values were calculated using one-way ANOVA (**c**).

Figure 1-2: CALR staining of wild-type and *Mitf* mutant OBs. a. Immunofluorescent staining of calretinin (CALR) showing the localization of CALR in the glomeruli, external plexiform (EPL) and in the mitral and granule cell layers (MC/GC) in OB. Scale bar indicates 100µm. **b.** Number of CALR+ cells OBs of the indicated genotypes. N=6 per genotype. The values on the graphs are mean ± SEM. P-values were calculated using two-way ANOVA (**b**).

Figure 1-3: Increase in apoptosis in OBs of *Mitf^(mi-vga9/mi-vga9)* mice. a. TUNEL staining of sections of OBs of the indicated genotypes, showing localization of apoptotic cells. **b.** The number of apoptotic cells of OB of wild-type and *Mitf^(mi-vga9/mi-vga9)* mice. N=9 per genotype. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bars are 50µm. P-values were calculated using two-tailed unpaired student's t-test. *P<0.05.

Figure 2-1: Gating kinetics of the A-type current (I$_A$). a. Representative recordings obtained to assess the gating kinetics of the I$_A$ current mediating channels. The projection neurons were first voltage clamped at -30mV and subsequently depolarized to 20mV utilizing a pre-hyperpolarizing pulse in between to recover the I$_A$ current from inactivation. This was done by increasing the time, Δt, spent in the hyperpolarized state, as indicated by the voltage trace on the left. To gauge its activation kinetics, the neurons were held at -90mV, as shown in the voltage trace on the right, and subsequently depolarized to 20mV, with a pre-depolarization pulse to -30mV applied in between. The A-type current gets inactivated by increasing the duration of the pre-pulse, Δt. **b**. Normalized mean currents (±SEM) plotted separately during recovery of the I$_A$ current from inactivation and during deactivation, in cells from WILD-TYPE and *Mitf$^{mi-vga9/mi-vga9}$* mice. The functions fitted to the data yield half-activation and half-recovery times of 15.2±4.7msec and 42.6±7.2msec for the wild-type (N=5, P=0.53) and 18.2±1.5msec and 51.3±9.3msec for the Mitf$^{mi-vga9/mi-vga9}$ (N=6, P=0.49) M/T

Figure 3-1 : *Kcnd2* expression is increased in the *Mitf* mutant olfactory bulb. a. RNA *in situ* hybridization of *Tbr2* (green), *Mitf* (red) and *Kcnd2* (magenta) performed on wild-type and *Mitf$^{mi-vga9/mi-vga9}$* OBs. **b**. *Kcnd3* count per cell. N=4 per genotype. The values on the graphs are mean ± SEM. DAPI nuclear staining is shown in blue. Scale bars are 50μm. P-values were calculated using two-way ANOVA (**b**) *P<0.05.

Figure 3-2: *c-Fos* and *Mitf* expression is increased upon activity induction in the olfactory bulb. RNA *in situ* hybridization of *c-Fos* (red) and *Mitf* (green) performed on wild-type OBs following AA. DAPI nuclear staining is shown in blue. Scale bars are 50μm.

Figure 3-3: Kcnd3 expression is increased upon activity induction in the olfactory bulb.

RNA *in situ* hybridization of *Mitf* (red) and *Kcnd3* (green) performed on wild-type OBs

following AA. DAPI nuclear staining is shown in blue. Scale bars are 50µm.

Figure 1-1

Figure 1-2

Figure 1-3

Figure 2-1

a

b

C57Bl/6J

Mitf^{mi-vga9/mi-vga9}

Figure 3-1

Figure 3-2

Figure 3-3

Supplementary Tables

Table 1: Potassium channel subunits in the OB.

Gene	K[+]-channel subunit	ChIPseq peak[1]				ABA expression in OB[2]	Current[3] Gutman 2005
		Strub[4] 2011	Laurette[5] 2015	Young[6] 2014	Calero-Nieto[7] 2014		
Kcna1	Kv1.1	No	No	No	No	Low expression in GCL	I_{DR}
Kcna2	Kv1.2	Yes	Yes	No	No	Expressed in MCL and ETC	I_{DR}
Kcna3	Kv1.3	No	Yes	No	No	Expressed in MCL and ETC	I_{DR}
Kcna4	Kv1.4	No	No	No	No	Expressed in MCL, ETC and GCL	I_A
Kcna5	Kv1.5	No	No	No	No	Low expression in ETC	I_A with β-

							subunit KvB3
Kcnc1	Kv3.1	No	No	No	No	Expressed in MCL and ETC	I_{DR}
Kcnc2	Kv3.2	No	No	No	No	Expressed in PGC, ETC, MCL	I_{DR}
Kcnc3	Kv3.3	No	No	No	No	Expressed in MCL and ETC	I_A
Kcnc4	Kv3.4	No	No	No	No	Low expression in MCL	I_A
Kcnd1	Kv4.1	No	No	No	No	Expressed throughout the OB	I_A
Kcnd2	Kv4.2	No	No	No	No	Expressed in MCL and GCL	I_A
Kcnd3	Kv4.3	Yes	Yes	No	No	Expressed throughout the OB	I_A

Available ChIPseq data, Allen Brain Atlas expression of potassium channel subunits in the olfactory bulb (OB) and published data on the type of current these subunits are regulating.

[1]"No" indicates that no peak observed was higher than negative control, whereas "yes" indicates that the peak observed was higher than the negative control. Region screened encompassed whole intronic regions and approximately 2kbp up and downstream of transcription start site.

[2]GCL= granule cell layer, MCL= mitral cell layer, ETC= external tufted cells, PGC= periglomerular cells

[3]I_{DR} = delayed rectifying current, I_A= A-type current.

[4] GEO Accession number GSE64137

[5] GEO Accession number GSM1517751

[6] GEO Accession number GSM1567047

[7] GEO Accession number GSM1167584

Table 2: Ratio of inhibitory to excitatory cells in the glomeruli.

Genotype	Ratio inhibitory/excitatory ±SEM	P-value vs C57Bl/6J	N
C57Bl/6J	2.071 ± 0.3292	-	6
$Mitf^{mi-vga9/+}$	2.313 ± 0.2668	0.7738	6
$Mitf^{mi-vga9/mi-vga9}$	2.012 ± 0.1278	0.9805	6

The ratio of inhibitory to excitatory cells was calculated by adding the number of CALB and PV positive cells from each individual mouse, and dividing it by the number of TBR2 positive cells in the same mouse. P-value was calculated using One-way ANOVA.

Table 3: Two-tailed paired student's t-test performed on avoidance behavior.

	C57Bl/6J	$Mitf^{mi-vga9/+}$	$Mitf^{mi-vga9/mi-vga9}$
Section 1	0.0016*	0.0009*	0.0004*
Section 2	0.6943	0.9325	0.3859
Section 3	0.0408*	0.0138*	0.0267*

Two-tailed paired student's t-test performed on the time mice spent in each section, comparing water and propionic acid conditions. *P<0.05. N=6-8.