The complete structure of the human TFIIH core complex

Basil J. Greber[1,2], Daniel B. Toso[1], Jie Fang[3], Eva Nogales[1,2,3,4,*]

[1] California Institute for Quantitative Biosciences (QB3), University of California,

Berkeley, California 94720, USA

[2] Molecular Biophysics and Integrative Bio-Imaging Division, Lawrence Berkeley

National Laboratory, Berkeley, California 94720, USA

[3] Howard Hughes Medical Institute, University of California, Berkeley, California

94720, USA

[4] Department of Molecular and Cell Biology, University of California, Berkeley,

California 94720, USA

[*] **For correspondence:** Enogales@lbl.gov (E.N.)

Competing interests: The authors declare that no competing interests exist.

17
18

ABSTRACT

Transcription factor IIH (TFIIH) is a heterodecameric protein complex critical for transcription initiation by RNA polymerase II and nucleotide excision DNA repair. The TFIIH core complex is sufficient for its repair functions and harbors the XPB and XPD DNA-dependent ATPase/helicase subunits, which are affected by human disease mutations. Transcription initiation additionally requires the CdK activating kinase subcomplex. Previous structural work has provided only partial insight into the architecture of TFIIH and its interactions within transcription pre-initiation complexes. Here, we present the complete structure of the human TFIIH core complex, determined by phase-plate cryo-electron microscopy at 3.7 Å resolution. The structure uncovers the molecular basis of TFIIH assembly, revealing how the recruitment of XPB by p52 depends on a pseudo-symmetric dimer of homologous domains in these two proteins. The structure also suggests a function for p62 in the regulation of XPD, and allows the mapping of previously unresolved human disease mutations.

INTRODUCTION

Transcription factor IIH (TFIIH) is a 10-subunit protein complex with a total molecular weight of 0.5 MDa that serves a dual role as a general transcription factor for transcription initiation by eukaryotic RNA polymerase II (Pol II), and as a DNA helicase complex in nucleotide excision DNA repair (NER) (Compe & Egly, 2016; Sainsbury et al., 2015). Mutations in TFIIH subunits that cause the inherited autosomal recessive disorders xeroderma pigmentosum (XP), trichothiodystrophy (TTD), and Cockayne syndrome (CS) are characterized by high incidence of cancer or premature ageing (Cleaver et al., 1999; Rapin, 2013). Furthermore, TFIIH is a possible target for anti-cancer compounds (Berico & Coin, 2017) and therefore of great importance for human health and disease.

The TFIIH core complex is composed of the seven subunits XPB, XPD, p62, p52, p44, p34, and p8, and is the form of TFIIH active in DNA repair (Svejstrup et al., 1995), where TFIIH serves as a DNA damage verification factor (Li et al., 2015;

Mathieu et al., 2013) and is responsible for opening a repair bubble around
damaged nucleotides. This activity depends on both the SF2-family DNA-
dependent ATPase XPB, and the DNA helicase activity of XPD (Coin et al., 2007;
Evans et al., 1997; Kuper et al., 2014). TFIIH function in transcription initiation
requires the double-stranded DNA translocase activity of XPB to regulate
opening of the transcription bubble (Alekseev et al., 2017; Fishburn et al., 2015;
Grünberg et al., 2012), and additionally the CdK activating kinase (CAK) complex,
which harbors the kinase activity of CDK7 as well as the Cyclin H and MAT1
subunits (Devault et al., 1995; Fisher et al., 1995; Fisher & Morgan, 1994;
Shiekhattar et al., 1995; Svejstrup et al., 1995). Targets of human CDK7 include
the C-terminal heptapeptide repeat domain of the largest subunit of Pol II, as
well as cell-cycle regulating CDKs (Fisher & Morgan, 1994; Shiekhattar et al.,
1995). MAT1 serves as a bridging subunit that promotes CAK subcomplex
formation by interacting with Cyclin H and CDK7 (Devault et al., 1995; Fisher et
al., 1995), recruits the CAK to the core complex by interactions with XPD and XPB
(Abdulrahman et al., 2013; Busso et al., 2000; Greber et al., 2017; Rossignol et al.,
1997), and also aids in Pol II-PIC formation by establishing interactions with the
core PIC (He et al., 2013; He et al., 2016; Schilbach et al., 2017). The presence of
MAT1 inhibits the helicase activity of XPD (Abdulrahman et al., 2013; Sandrock &
Egly, 2001), but the mechanism of this inhibition is not fully understood. While
the enzymatic activity of XPD is not required for transcription initiation, it is
critical for the DNA repair function of TFIIH (Dubaele et al., 2003; Evans et al.,
1997; Kuper et al., 2014). Therefore, NER requires the release of the CAK
subcomplex from the core complex (Coin et al., 2008). The activities of both XPB
and XPD are regulated by interactions with additional TFIIH components,
including that of p44 with XPD (Coin et al., 1998; Dubaele et al., 2003; Kim et al.,
2015), and those of the p52-p8 module with XPB (Coin et al., 2007; Coin et al.,
2006; Jawhari et al., 2002; Kainov et al., 2008). These interactions are likely to be
crucial for TFIIH function, as some are affected by disease mutations (Cleaver et
al., 1999), but they have been only partially characterized mechanistically.
82
Our previous structure of the TFIIH core-MAT1 complex at 4.4 Å resolution
(Greber et al., 2017) allowed modeling of TFIIH in the best-resolved parts of the

density map, but several functionally important regions remained unassigned or
only partially interpreted because reliable *de novo* tracing of entire domains in
the absence of existing structural models was not possible. Here, we present the
complete structure of the human TFIIH core complex in association with the CAK
subunit MAT1, determined by phase plate cryo-electron microscopy (cryo-EM)
at 3.7 Å resolution. Our structure reveals the complete architecture of the TFIIH
core complex and provides detailed insight into the interactions that govern its
assembly. Additionally, our cryo-EM maps define the molecular contacts that
control the regulation of the XPB and XPD subunits of TFIIH, including the critical
p52-XBP interaction, and an extensive regulatory network around XPD, formed
by XPB, p62, p44, and MAT1.

RESULTS

Structure determination of TFIIH

To determine the complete structure of the human TFIIH core complex, we
collected several large cryo-EM datasets (Supplementary Table 1) of TFIIH
immuno-purified from HeLa cells using an electron microscope equipped with a
Volta phase plate (VPP) (Danev & Baumeister, 2017) and a direct electron
detector camera mounted behind an energy filter. From a homogeneous subset
of approximately 140,000 TFIIH particle images identified by 3D classification
(Scheres, 2010), we reconstructed a 3D cryo-EM density map at 3.7 Å resolution
(Figure 1 – figure supplements 1, 2A-C). This VPP-based cryo-EM map was
substantially improved compared to our previous maps obtained without phase
plate, both in resolution and interpretability (Figure 1 – figure supplement 2D-
G), and enabled building, refinement, and full validation of an atomic model of
the TFIIH core complex and the MAT1 subunit of the CAK subcomplex (Figure
1A-C, Figure 1 – figure supplement 2B, C, Supplementary Tables 2, 3), while the
remainder of the CAK is invisible in our map due to extreme motion. Tracing and
sequence register assignment of protein components modeled *de novo* was
facilitated by density maps obtained from focused classification and multibody
refinement (Figure 1 – figure supplements 3-5) (Bai et al., 2015; Nakane et al.,

2018), which resulted in density maps for sub-volumes of the complex at up to 3.6 Å resolution. Both the overall and multibody-refined maps showed clear side chain information (Figure 1 – figure supplement 6A-D). Furthermore, our model was corroborated by existing chemical crosslinking-mass spectrometry (CX-MS) data of human TFIIH (Luo et al., 2015) and site-specific crosslinks from yeast TFIIH (Warfield et al., 2016) (Figure 1 – figure supplement 6E-I).

Figure 1. Structure of the TFIIH core complex. (**A**, **B**, **C**) Three views of the structure of the TFIIH core complex and MAT1. Subunits are color-coded and labeled (in color); individual domains are labeled (in black) and circled if needed for clarity. (**D**) Domain-level interaction network between the components of the TFIIH core complex and MAT1 derived from our structure. Proteins are shown with the same colors as in A and major unmodeled regions are shown in grey. Abbreviations: CTD: C-terminal domain; DRD: DNA damage recognition domain; FeS: iron sulfur cluster domain; NTD: N-terminal domain; vWFA: von Willebrand Factor A.

Figure supplement 1. Data processing scheme for global reconstruction.

Figure supplement 2. Resolution estimation, validation statistics, and quality of the density.

Figure supplement 3. Focused classification of the p62 BSD2 domain.

Detailed architecture of TFIIH and structure of p62

Our structure of the TFIIH core complex shows its horseshoe-like overall shape

(Figure 1A-C), as observed in previous lower-resolution reconstructions of free

and PIC-bound TFIIH (Gibbons et al., 2012; Greber et al., 2017; He et al., 2016;

Murakami et al., 2015; Schilbach et al., 2017), and allows us to define the

complete set of inter-subunit interactions that lead to the formation of the TFIIH

core complex (Figure 1D).

The largest subunits of the complex, the SF2-family DNA-dependent ATPases

XPB and XPD, both containing two RecA-like domains (RecA1 and RecA2),

interact directly (Greber et al., 2017) and are on one side of the complex, and are

additionally bridged by MAT1 (Figure 1B), which has been shown to interact

with either ATPase in isolation (Busso et al., 2000). On the side facing away from

MAT1, XPD interacts with the von Willebrand Factor A (vWFA) domain of p44

(Coin et al., 1998; Dubaele et al., 2003; He et al., 2016; Kim et al., 2015), which in

turn forms a tight interaction with p34 via interlocking eZnF domains (Schilbach

et al., 2017) and a p44 RING domain interaction (Radu et al., 2017) (Figure 1B-D,

Figure 1 – figure supplement 6J, K), consistent with the formation of a

multivalent interaction network between p34 and p44 (Radu et al., 2017). The

vWFA domain of p34 recruits p52 by a three-way interaction that involves the

most N-terminal winged helix domain in p52 and a helical segment of p62

(Schilbach et al., 2017) (Figure 1 – figure supplement 6L). The p52 C-terminal

region comprises two domains; first, the 'clutch' that interacts with XPB (Jawhari

et al., 2002) and second, a dimerization module that binds p8 (Kainov et al.,

2008), thereby recruiting XPB to TFIIH and cradling XPB RecA2 (see below). In

addition to this structural framework that is formed by folded domains, our cryo-

EM map reveals several interactions involving extended protein segments,

including several interactions formed by p62 (Figure 2), and an interaction
between the p44 N-terminal extension (NTE) and the N-terminal domain (NTD)
of XPB (Figure 1B). To form this interaction, approx. 15 residues of p44 span the
distance between the p44 vWFA domain and the XPB NTD, where a small helical
motif in p44 contacts XPB residues 72-75, 95-102, and 139-143, in agreement
with CX-MS data (Luo et al., 2015) (Figure 1 – figure supplement 6E). Partial
deletion of the p44 NTE in yeast causes a slow-growth phenotype, suggesting a
functional role for this p44-XPB interaction (Warfield et al., 2016).

Figure 2. The structure of p62. (**A**) View of the top surface of the TFIIH core
complex; p62 is color-coded by structural region. (**B**) The BSD2 (blue) and XPD
anchor segments (teal) of p62 (surface) interact with the region around the XPD
substrate-binding cavity. (**C**) Residues 346-365 of p62 (yellow) approach the

nucleotide-binding pocket of XPD. ADP superposed from the structure of the

Brr2 helicase (Nguyen et al., 2013). (**D**) Superposition of DNA from the structure

of the substrate-bound helicase Hel308 (Büttner et al., 2007) shows that the

positions of p62 and RecA1-bound DNA overlap.

Figure supplement 1. Analysis of the structure of p62.

The p62 subunit is almost completely resolved in our structure and exhibits a

complex beads-on-a-string-like topology. It fully encircles the top surface of

TFIIH (Figures 1C, 2A, Figure 2 – figure supplement 1), interacting with XPD,

p52, p44, and p34, in agreement with previous structural findings (Greber et al.,

2017; Schilbach et al., 2017). Based on these interactions, p62 can be subdivided

into three functional regions: (i) the N-terminal PH-domain, disordered in our

structure, is responsible for mediating interactions with components of the core

transcriptional machinery (Di Lello et al., 2008; He et al., 2016; Schilbach et al.,

2017), transcriptional regulators (Di Lello et al., 2006), and DNA repair pathways

(Gervais et al., 2004; Lafrance-Vanasse et al., 2013; Okuda et al., 2017); (ii)

residues 108-148 and 454-548 of p62, including the first BSD (BTF2-like,

synapse-associated, DOS2-like) domain (BSD1) and the C-terminal 3-helix

bundle, play an architectural role by binding to p34 and the extended zinc finger

(eZnF) domain of p44 (Figure 2A, C, Figure 1 – figure supplement 6J-L, Figure 2 –

figure supplement 1B, C); and (iii) residues 160-365, including the BSD2 domain,

are responsible for interactions with and regulation of XPD (Figure 2B-D, Figure

2 – figure supplement 1D-F).

Specifically, p62 residues 160-365 form three structural elements that interact

with XPD (Figure 2B), in agreement with previous biochemical, structural, and

CX-MS data (Figure 1 – figure supplement 6G) (Jawhari et al., 2004; Luo et al.,

2015; Schilbach et al., 2017). First, an α-helix formed by p62 residues 295-318

binds directly to XPD RecA2 and thereby recruits residues 160-258 of p62,

comprising the BSD2 domain and adjacent sequence elements, to this surface of

XPD RecA2 (Figures 2A, B, Figure 2 – figure supplement 1D). Second, p62

residues 266-287 are inserted into the DNA-binding cavity of XPD (Figures 2B,

D), in agreement with previous observations (Schilbach et al., 2017). This inserted p62 segment directly blocks a DNA-binding site on XPD RecA1 (Figure 2D) and localizes near the access path to a pore-like structure between the XPD FeS and ARCH domains. While p62 does not directly contact the DNA-binding surface on XPD RecA2, it may still sterically interfere with DNA binding to the helicase elements of XPD in this region (Figure 2 – figure supplement 1E). Therefore, this segment of p62 may need to move away when XPD binds and unwinds DNA. Third, p62 residues 350-358 form a short α-helix that binds in a cleft between the two RecA-like domains of XPD (Figure 2C), so that it not only closes the entrance to the nucleotide binding pocket in XPD RecA1 (Figure 2 – figure supplement 1F), but also partially overlaps with the predicted location of the nucleotide itself (Figure 2C), strongly suggesting a role for this p62 sequence element in XPD regulation. The density for these structural elements of p62 (residues 260-300 and 346-365) in our cryo-EM map is weaker than for the remainder of the complex, suggesting a dynamic interaction with XPD that enables them to modulate the access to the nucleotide-binding pocket, the DNA-binding cavity, and the DNA-translocating pore of XPD, depending on the functional state of TFIIH. 3D reconstructions of TFIIH classified for these regions of p62 (Figure 1 – figure supplement 3) show globally intact TFIIH, both in the presence and absence of the p62 segments at these XPD sites (Figure 2 – figure supplement 1G-J), supporting our hypothesis of dynamic regulation, rather than the alternative hypothesis of p62 binding to XPD as a requirement for TFIIH stability (Luo et al., 2015).

Molecular basis of XPB recruitment by p52

Our structure of TFIIH resolves the structure and interactions of all four folded domains of human XPB – two RecA-like domains that form the SF2-family type helicase cassette, a DNA damage recognition domain (DRD)-like domain, and an N-terminal domain (NTD) (Figure 3A, Figure 3 – figure supplement 1) – and reveals the molecular basis of XPB recruitment by p52. The XPB NTD encompasses residues 1-165, with the first 55 residues forming an N-terminal extension (NTE), and the remainder assuming a mixed α/β-fold with four α-

helices and five β-strands (Figure 3A). The side chain densities in the cryo-EM

map (Figure 1 – figure supplement 6A) and CX-MS data (Luo et al., 2015) (Figure

1 – figure supplement 6F) both confirm our assignment of this domain. Existing

biochemical data show that the XPB NTD is required for integration of XPB into

TFIIH (Jawhari et al., 2002) by forming an interaction with p52 that has been

referred to as the 'clutch' (Schilbach et al., 2017). In our structure, the p52

contribution to the clutch encompasses p52 residues 306-399, which, strikingly,

assume the same overall fold as the XPD NTD (Figure 3B), as hypothesized

previously (He et al., 2016; Luo et al., 2015), thereby forming a pseudo-

symmetric dimer of structurally homologous domains. The two domains interact

through their β-sheets, via both hydrophobic and charged interactions (Figure 3

– figure supplement 2A-C), and with the most N-terminal β-strand emanating

from the XPB NTD extending the p52 β-sheet by additional lateral interactions

(Figure 3A, C).

Our structural findings rationalize biochemical data that show that deletion of

XPB residues 1-207, but not deletion of residues 1-44, impairs the p52-XPB

interaction (Jawhari et al., 2002) (Figure 3C). Our structure is also consistent

with data indicating that p52 residues 304-381 are critical for the XPB-p52

interaction (Coin et al., 2007; Jawhari et al., 2002), but does not show any

contacts that could explain that reported binding of XPB to p52 residues 1-135

or 1-304 (Jawhari et al., 2002) (Figure 3 – figure supplement 2D, E). The

interaction between p52 and XPB not only recruits XPB to TFIIH, but also

stimulates its ATPase activity *in vitro* (Coin et al., 2007). Because our structure

does not shown any elements of p52 approaching the XPB nucleotide-binding

pocket, we propose that this effect is likely induced by the interactions of p52

with the XPB NTD and RecA2, which may, together with p8 (Coin et al., 2006),

properly arrange the XPB helicase cassette to bind and hydrolyze ATP (Figure

3D) (Grünberg et al., 2012).

Figure 3. Structure and interactions of XPB. (**A**) Bottom lobe of TFIIH. XPB

RecA1/2 teal, DRD blue, NTD dark blue, p52 yellow, p8 green, p44 NTE red. (**B**)

Superposition of the XPB NTD and the p52 clutch domain. (**C**) Mapping of

mutations on the XPB NTD and the p52 clutch domain; mutated regions are

color-coded or shown as spheres (see text for details). (**D**) The combined

interactions of the p52 clutch, the p8-p52 CTD dimer, and the XPB NTD with XPB

RecA2 may restrict the conformational flexibility of XPB RecA2 to optimize XPB

activity. (**E**) An extension of the DRD (blue) contacts XPD (green). The sequence

for which formation of an α-helix is predicted (Kelley et al., 2015) is indicated.

(**F**) The DRD extension overlaps with the substrate-binding site on XPD RecA2.

Substrate DNA modeled from PDB ID 5HW8 (Constantinescu-Aruxandei et al.,

2016).

Figure supplement 1. Domain organization of XPB and sequence alignment of

the N-terminal regions of XPB and XPB-like enzymes from eukaryotes, archaea,

and bacteria.

Figure supplement 2. Structure and interactions of the XPB DRD and NTD.

The XPB NTD is the site of the two human disease mutations F99S and T119P, which cause XP and TTD, respectively (Cleaver et al., 1999). Or structure shows that neither of these residues is in direct contact with p52 or the RecA-like domains of XPB, suggesting that the F99S and T119P mutations exert their detrimental effects through structural perturbation of the XPB NTD (Figure 3 – figure supplements 1B, 2F-I). Specifically, T119 is located near a turn at the end of a β-strand (Figure 3 – figure supplement 2G), where its side chain points towards the solvent. Nevertheless, this residue is highly conserved in eukaryotic XPB from ciliates to humans, in some archaeal and bacterial XPB homologs (Figure 3 – figure supplement 1B), and in the structurally homologous clutch domain in p52 (Figure 3 – figure supplement 2F-H). This conservation suggests that a threonine at this position is important for the efficient folding of this domain in general, and that the T119P mutation may cause its destabilization, resulting in lower levels of active enzyme in TTD patients. Lower overall levels of properly assembled TFIIH have been shown to be a hallmark of TTD (Botta et al., 2002; Dubaele et al., 2003; Giglia-Mari et al., 2004) and could explain the disease-causing effect of T119P *in vivo* even though recombinant TFIIH carrying this mutation retains some activity in both transcription initiation and NER (Coin et al., 2007). A less likely alternative, given the conservation of the equivalent residue in the p52 clutch, is that T119 is involved in an interaction with a factor that is critical for cellular function, e.g. in NER.

The F99S mutation affects a residue that is conserved throughout eukaryotic XPB (Figure 3 – figure supplement 1B), is inserted into a conserved hydrophobic pocket, and localizes to an α-helix at the XPB contact site with the p44 N-terminal extension (Figure 3 – figure supplement 2I). This mutation is likely to impair the stability and folding of the XPB NTD. Unlike T119P, this mutation leads to impaired DNA opening in NER assays, reduced interaction with p52, reduced ATPase activity (Coin et al., 2007), and strong impairment in DNA damage repair (Riou et al., 1999), suggesting a severe effect on the structure of the XPB NTD.

Natural and synthetic mutations in the *Drosophila melanogaster* homolog of p52 that lead to disease-like phenotypes in flies and have similar defects when

335   introduced into human cells (Fregoso et al., 2007) map directly to the p52-XPB
336   interface, explaining their detrimental phenotypes (Figure 3C, Figure 3 – figure
337   supplement 2A).
338
Our structure assigns XPB residues 165-300 to a DRD-like domain that connects
the NTD to the RecA-like domain (Figure 3A, Figure 3 – figure supplement 1A),
the deletion of which is lethal in yeast (Warfield et al., 2016). The DRD is a DNA-
binding domain found in DNA repair enzymes and chromatin remodelers (Mason
et al., 2014; Obmolova et al., 2000) and has been implicated in DNA damage
recognition in archaeal XPB (Fan et al., 2006; Rouillon & White, 2010). Our 3.7 Å-
resolution map of TFIIH reveals that in eukaryotic XPB, one β-strand of the DRD
of archaeal XPB is replaced by an insertion of approximately 70 residues that
exhibits relatively low sequence conservation (Figure 3E, Figure 3 – figure
supplement 1B) and shifts the domain boundaries of the human XPB DRD-like
domain with respect to previous sequence alignments (Fan et al., 2006;
Oksenych et al., 2009). The part of this insertion resolved in our map consists of
a negatively charged linker and an α-helical element that contacts XPD directly
(Figure 3E, F). The surface on XPD involved in this interaction has been
implicated in the initial step of DNA substrate binding by XPD (Constantinescu-
Aruxandei et al., 2016; Kuper et al., 2012). Density features and secondary
structure prediction indicate the presence of several aromatic side chains of XPB
near the interface (Figure 3E), where they might form contacts resembling those
of nucleoside bases of XPD-bound DNA substrates (Figure 3F). Thus, XPB may
modulate substrate binding by XPD, further reinforcing the idea that XPD activity
is regulated by several other components of TFIIH.

Figure 4. Conformational dynamics of TFIIH. (**A**) Results of multibody analysis. Several major modes of motion involve the enzymatic subunits of the TFIIH core complex or their domains. (**B**) Motion of the p52 clutch domain (closed conformation gold, open conformation light yellow) and associated XPB NTD (closed conformation blue, open conformation light blue) relative to the remainder of p52 (brown), based on comparison of free and PIC-bound TFIIH structures. (**C**) XPB motions from the closed conformation (free TFIIH; darker blue hues) and the open conformation (TFIIH-PIC; lighter hues). The MAT1-XPB contact dissociates during this rearrangement. (**D**) Schematic model for the conformational transitions in MAT1 and repositioning of the CAK kinase module during Pol II-PIC entry of TFIIH.

Figure supplement 1. Analysis of the conformational variance of TFIIH.

Figure supplement 2. Conformational dynamics of TFIIH and comparison with the PIC-bound TFIIH.

Conformational dynamics of the TFIIH core complex

In order to investigate the dynamics of TFIIH, we analyzed the conformational landscape of the particles in our cryo-EM dataset (Figure 4A, Figure 4 – figure supplement 1) (Nakane et al., 2018). This analysis revealed the relative motions

of the two ATPases and their domains (Figure 4A). The major mode of motion,
which involves the breaking of the interaction between XPB and XPD, resembles
the conformational transition of TFIIH when it enters the Pol II-PIC and binds to
DNA (Greber et al., 2017; He et al., 2016; Schilbach et al., 2017) (Figure 4 – figure
supplement 2A-C). Analysis of our TFIIH structure and comparison with that of
the complex within the Pol II-PIC maps (He et al., 2016; Schilbach et al., 2017)
allowed us to identify a specific rearrangement at the interface between the
clutch and adjacent winged helix domain in p52 (Figure 4B) as the basis of this
conformational change. A structural unit composed of XPB, p8 and the clutch
domain of p52 undergoes a downward motion upon DNA-binding within the Pol
II-PIC (Figure 4B, Figure 4 – figure supplement 2B, C). This conformational
change in TFIIH upon PIC entry also appears to break the interaction between
MAT1 and the XPB DRD-like domain (Figure 4C), which in turn might serve to
enable positioning of the CDK7-cyclin H dimer within the CAK subcomplex at the
appropriate location for Pol II-CTD phosphorylation in the mediator-bound Pol
II-PIC (Figure 4D) (Schilbach et al., 2017). Our structural comparison also reveals
that a TFIIE-XPB interaction that has been implicated in XPB regulation
(Schilbach et al., 2017) may depend on the existence of the open conformation of
TFIIH, as there would be steric hindrance in a complex involving the closed
conformation of TFIIH (Figure 4 – figure supplement 2D-F).
402
403 **Structure of XPD**
404
The structure of XPD shows the conserved domain arrangement of two RecA-like
domains (RecA1 and RecA2), with the FeS and ARCH domain insertions in RecA1
(Constantinescu-Aruxandei et al., 2016; Fan et al., 2008; Kuper et al., 2012). The
quality of the map allowed us to interpret the density for the N- and C-termini of
XPD, which closely approach each other near the nucleotide-binding site within
RecA1 (Figure 5A). The N-terminus of XPD forms a short two-stranded β-sheet
near the nucleotide-binding site in XPD RecA1 (which is empty in our structure)
and may contribute to the stabilization of the bound nucleotide via the aromatic
side chains of Y14 and Y18, the latter being affected by the Y18H mutation in an
XP/TTD patient (Kralund et al., 2013) (Figure 5 – figure supplement 1A-C). The

XPD C-terminal segment runs along the side of XPD RecA2 and interacts with the linker between RecA1 and RecA2 (Figure 5 – figure supplement 1D). The C-terminal segment includes the site of the K751Q polymorphism (Figure 5 – figure supplement 1D), and deletion of this terminal segment causes XP in human patients (Cleaver et al., 1999).

Figure 5. Structure and regulation of XPD. (**A**) Structure of XPD colored by domain. N- and C-terminal segments (blue and purple, respectively) of XPD are indicated. An ADP molecule (superposed from (Nguyen et al., 2013)) denotes the nucleotide-binding pocket in XPD RecA1, which is empty in our structure. (**B**) Structure of the FeS domain. Residues critical for XPD enzymatic activity (blue) and damage verification (teal) are indicated. The R112H mutation causes TTD in human patients. ARCH domain insertion brown. DNA superposed from (Büttner et al., 2007). (**C**) Interaction network of XPD with surrounding TFIIH subunits (interacting regions colored, remainder grey). (**D**) Cartoon model for repression and de-repression of XPD by MAT1, XPB, and p62. (**E**) XPD-p44 interacting regions (defined as residues within < 4 Å of the neighboring protein) are colored in dark green (XPD) and dark red (p44). Residues discussed in the text are shown as sticks; those with mutation data (natural variants or experimental

constructs) are colored yellow on XPD, teal on p44. The remainder of the β4-α5

loop harboring the synthetic p44 mutations is colored teal as well.

Figure supplement 1. Structure of XPD.

Figure supplement 2. XPD-p44 interaction.

Before XPD-bound DNA reaches the helicase motifs in the RecA like domains, it

passes through a pore-like structure next to the 4FeS cluster at the interface

between the FeS and ARCH domains (Figure 5 – figure supplement 1E, F) (Cheng

& Wigley, 2018; Constantinescu-Aruxandei et al., 2016; Kuper et al., 2012; Liu et

al., 2008; Wolski et al., 2008). This region was poorly defined in previous TFIIH

reconstructions, but our cryo-EM map now shows side-chain densities for the

aromatic residues Y158, F161, and F193, which are critical for the DNA-binding,

ATPase, and helicase activities of XPD (Kuper et al., 2014), as well as for residues

Y192 and R196, which form part of a DNA lesion recognition pocket (Mathieu et

al., 2013) (Figure 5B). This functionally important region is only partially

conserved in archaeal XPD homologs (Figure 5 – figure supplement 1G-I) (Fan et

al., 2008; Kuper et al., 2012; Wolski et al., 2008). A eukaryotic-specific loop

insertion in the XPD ARCH domain (Greber et al., 2017; Schilbach et al., 2017)

closely approaches this binding pocket (Figure 5B) and may serve to regulate the

binding of DNA in the lesion recognition pocket such as to prevent untimely

access of substrates to the XPD pore.

Interactions and regulation of XPD

Our structure of TFIIH shows that XPD forms architectural and regulatory

interactions with four other TFIIH subunits: XPB, p62, p44, and MAT1, which

together form a cradle-like structure around XPD (Figure 5C). We described

above two interactions that could potentially regulate XPD activity: the newly

defined interaction of an insertion element in the XPB DRD with a DNA-binding

site in XPD (Figure 3E, F); and XPD-p62 interactions involving the XPD

nucleotide binding pocked and DNA binding cavity (Figure 2, Figure 2 – figure

supplement 1D-F) that implicate p62, as well as XPB, in XPD regulation.

Additionally, it is known that the helicase activity of XPD is inhibited by the CAK

subcomplex (Araújo et al., 2000; Sandrock & Egly, 2001). The contacts we see

between MAT1 and XPD localize to the ARCH domain of XPD and the N-terminal

RING domain and helical bundle of MAT1 (residues 1-210) (Figure 5C), in

agreement with previous structural (Greber et al., 2017; Schilbach et al., 2017)

and biochemical analysis (Abdulrahman et al., 2013; Luo et al., 2015; Warfield et

al., 2016). The interaction between the XPD ARCH domain and the MAT1 helical

bundle is characterized by charge complementarity (Figure 5 – figure

supplement 1J-M). This interface is highly dynamic, enabling the release of MAT1

and the entire CAK subcomplex from TFIIH during NER, as well as its subsequent

re-association to regenerate a transcription-competent TFIIH (Coin et al., 2008).

Insertion of substrate DNA into the pore between the XPD ARCH and FeS

domains requires the flexibility of the XPD ARCH domain (Constantinescu-

Aruxandei et al., 2016), and large domain motions have been observed in the

structure of the DNA-bound homologous helicase DinG upon nucleotide binding

(Cheng & Wigley, 2018). This suggests a role for the mobility of the ARCH

domain in both DNA loading and DNA translocation by the XPD helicase. Our

structure suggests that binding of the MAT1 helical bundle and RING domain to

the ARCH domain may prevent such motion and therefore the subsequent

substrate loading and XPD helicase activity (Figure 5D), in agreement with

biochemical data that show XPD inhibition upon MAT1 binding (Sandrock &

Egly, 2001), as well as reduced single-stranded DNA affinity of TFIIH in the

presence of the CAK (Li et al., 2015). Conversely, release of MAT1 from XPD

might allow the ARCH domain to move more freely, thereby de-repressing XPD.

Furthermore, displacement of the MAT1 α-helix that connects XPD to XPB may

allow XPB to move away from XPD, thereby unmasking the substrate-binding

site on XPD RecA2 that is otherwise occluded by the DRD insertion element

(Figure 5D). This latter conformational change would be similar, overall, to that

seen for TFIIH upon incorporation into the Pol II-PIC, where XPD and XPB move

apart and density for the MAT1 helix is missing (Figure 4C) (He et al., 2016;

Schilbach et al., 2017). We propose that the combined unmasking of the XPD

substrate binding site and the enhanced flexibility of the XPD ARCH domain may

both contribute to de-repression of the XPD helicase upon release of MAT1. This mechanism of XPD inhibition by MAT1 does not exclude the possibility of additional repression of NER activity by the CAK subcomplex through phosphorylation of NER pathway components (Araújo et al., 2000).

Figure 6. Disease mutations in XPD. (**A**) Residues affected by disease mutations are shown as spheres (XP purple; TTD yellow; CS-XP orange). Conserved helicase elements for DNA binding are shown in blue, for nucleotide binding and hydrolysis in green, and for coupling of nucleotide hydrolysis and DNA translocation in brown. DNA superposed from (Büttner et al., 2007). (**B**) Salt bridge between R658 (RecA2) and D240 (RecA1) visualized in our structure that is affected by the temperature sensitive TTD mutation R658C (Vermeulen et al., 2001).

Figure supplement 1. Mapping of XPD disease mutations on the TFIIH structure.

Our structure also resolves in detail the XPD-p44 interaction, a known regulatory interface (Dubaele et al., 2003; Kim et al., 2015; Kuper et al., 2014) affected by numerous disease mutations (Cleaver et al., 1999; Greber et al., 2017; Kuper et al., 2014) (Figure 5E). The relatively small interaction surface between p44 and XPD, of just 1010 Å² (Figure 5 – figure supplement 2A), contrasts with the much larger buried surface of 3370 Å² between XPD and p62, or 1570 Å² for the p52-

XPB interaction. This smaller interaction surface may result in higher sensitivity
to mutations that localize at the XPD-p44 interface. Our structure, thus,
rationalizes the deleterious effect of a number of natural and synthetic mutations
in this interface (Supplementary Note 1, Figure 5 – figure supplement 2B-E),
including mutations L174W and T175R in the β4-α5 loop of p44 (Figure 5E)
(Kim et al., 2015; Seroz et al., 2000), which may lead to steric clashes in the
densely packed interface (Figure 5 – figure supplement 2B), and the XPD R722W
mutation (Kuper et al., 2014), which disrupts the salt bridge with D75 in p44 and
may additionally cause steric clashes with neighboring p44 residues due to the
bulky tryptophan side chain (Figure 5 – figure supplement 2C). Our structure
also shows that, in contrast to a previously proposed model (Luo et al., 2015),
the XPD R616P, D673G, and G675R disease mutations act either via disruption of
the XPD structure or the XPD-p44 interface, but not via disruption of the
interaction with p62 (see Supplementary Note 1, Figure 5 – figure supplement
2D). Notably, the p44-dependet stimulation of XPD activity does not depend on
the presence of p62 (Dubaele et al., 2003; Kim et al., 2015; Kuper et al., 2014).
We were also able to map a number of disease mutations onto our XPD structure
(Figure 6A) and analyze in detail the interactions involving the affected residues
(example shown in Figure 6B). Our analysis confirms that XP mutations mostly
localize near the helicase substrate-binding or active sites, while TTD mutations
predominantly localize to the periphery of XPD (Figure 6, Figure 6 – figure
supplement 1) (Fan et al., 2008; Liu et al., 2008), where they disrupt TFIIH
assembly and cause the transcription defects that are a hallmark of this disease
(Dubaele et al., 2003) (Supplementary Note 2).
549
550 **DISCUSSION**
551
Our study reveals the complete structure of the TFIIH core complex and provides
mechanistic insights into the regulation of its two component helicases.
Specifically, it shows XPD wrapped by numerous interactions with XPB, p62, p44,
and MAT1 (Figure 5C, D), indicating how its activity can be tightly controlled and
de-repressed only when its enzymatic function is needed. XPD activity is not
needed and most likely inhibited during transcription initiation, but it may also

be tightly controlled during NER, when repair bubble opening and lesion verification need to be coordinated with the recruitment and activation of the damage recognition and processing machinery (Figure 7 and Discussion). While the regulation of XPD by MAT1 and p44 has been studied in some detail, and the domain motions in TFIIH suggest a straightforward mechanism for liberating the substrate-binding site on XPD RecA2, less was known about the interplay between XPD and p62. Our structure now shows how p62 is able to impede both substrate and nucleotide binding in XPD RecA1, and hints at dynamic structural changes of p62 during de-repression and enzymatic activity of XPD, possibly regulated by other components of the transcription or NER pathways.

Figure 7. Implications for assembly of the repair bubble during NER. (**A**) Schematic of DNA-bound TFIIH (DNA damage verification pocket in XPD and DNA 5'-phosphates indicated by purple and orange spheres, respectively) Binding of both XPB and XPD to the damaged strand would lead to extrusion of a bubble when XPD scans in the 5'-3' direction, while XPB may be stationary or contribute to bubble extrusion if translocating in the 3'-5' direction (DNA superposed from PDB IDs 5H8W, 2P6R, 5OQJ (Büttner et al., 2007; Constantinescu-Aruxandei et al., 2016; Schilbach et al., 2017)). (**B**) Binding of XPB to the undamaged strand would enable the entire complex to scan in 5'-3' direction, given the opposing polarities of the two ATPases/helicases involved. (**C**) Model for the assembled repair bubble. Positions of NER factors are approximate. See Discussion for details.

Our results allow us to put extensive biochemical data on the NER pathway into a

structural context (Figure 7). Depending on whether XPB binds to the damaged

(Figure 7A) or undamaged (Figure 7B) strand, the combined action of XPD and

XPB could lead to the extrusion of a DNA bubble (Figure 7A) or to the tracking of

the entire complex towards the lesion (Figure 7B), which is initially located 3' of

the TFIIH binding site (Sugasawa et al., 2009). The latter hypothesis is attractive

in the context of biochemical data that show that XPD tracks along the damaged

strand in the 5' to 3' direction until it encounters the DNA lesion in order to

verify the presence of a bona fide NER substrate (Buechner et al., 2014; Li et al.,

2015; Mathieu et al., 2013; Naegeli et al., 1993; Sugasawa et al., 2009; Wirth et

al., 2016). It is worth noting that the length of DNA fragments excised during

NER is approx. 29 nt, with 22 nt located 5' and 5 nt located 3' of a thymine dimer

lesion (Huang et al., 1992). According to our structure, the 22 nt 5'-fragment

corresponds well to the estimated 20 nt of DNA that are required to span the

distance from the DNA damage verification pocket in XPD (Mathieu et al., 2013)

to the helicase elements of XPB. This proposal is compatible with a model in

which TFIIH sitting on the open repair bubble might track towards the lesion,

where it would stop due to inhibition of XPD (Li et al., 2015; Mathieu et al., 2013;

Naegeli et al., 1993), at which point double incision could be initiated. However,

this model (Figure 7B) would require strong DNA bending before both XPB and

XPD could be loaded. Additionally, it has not been fully resolved whether XPB

participates in DNA translocation or unwinding during TFIIH activity in NER (Li

et al., 2015), which would be required in the tracking model (Li et al., 2015), or

whether it exclusively acts to anchor the complex in the vicinity of the DNA

lesion (Coin et al., 2007; Oksenych et al., 2009). Independently of the orientation

of the repair bubble, our structural data are compatible with literature data

introduced above and a model that localizes XPG near XPD and p62 (site of 3'-

incision), XPF-ERCC1 near XPB (site of 5'-incision), and with RPA binding the

non-damaged strand (Figure 7C). We have currently not included XPA in this

model because its interactions with distinct partners or participation in various

processes, such as involvement in CAK release (Coin et al., 2008), binding to p8

(Ziani et al., 2014), and participation in helicase stalling after lesion recognition

(Li et al., 2015), suggest its localization to various, often distant sites on TFIIH, or

the repair bubble in general (Sugitani et al., 2016).

In summary, our structure of the human TFIIH core complex reveals the

interactions that govern the architecture and function of this molecular machine,

provides new insights into the regulation of its enzymatic subunits, and thus

constitutes an excellent framework for further mechanistic studies of TFIIH in

the context of larger DNA repair and transcription assemblies.

METHODS

TFIIH purification, cryo-EM specimen preparation, and data collection

TFIIH was purified and cryo-EM grids were prepared on carbon-coated C-flat CF

4/2 holey carbon grids (Protochips) using a Thermo Fisher Scientific Vitrobot

Mk. IV, as previously described (Greber et al., 2017). To improve on our previous

4.4 Å cryo-EM map of human TFIIH (Greber et al., 2017), which was based on 4

cryo-EM datasets (3 of which were retained in the 4.4 Å reconstruction, datasets

8-10 in Supplementary Table 1) from a Titan microscope (Thermo Fisher

Scientific) equipped with a side-entry holder (Gatan) and a K2 Summit direct

electron detector (Gatan), we collected new data (dataset 7 in Supplementary

Table 1) on a Titan KRIOS microscope (Thermo Fisher Scientific) operated at 300

kV extraction voltage and equipped with a C_S-corrector, a K2 Summit direct

electron detector (Gatan) operated in super-resolution counting mode, and a

Quantum energy filter (Gatan). This dataset was collected under the same

imaging conditions as our previous data (i.e. 37,879 x magnification resulting in

1.32 Å pixel size, and at a total exposure of 40 e$^-$ Å^{-2}), except for the change of

microscope. Datasets 7-10 could be combined to yield a cryo-EM map at 4.3 Å

resolution (not shown), however, this did not lead to a substantial improvement

in map quality, suggesting that particle alignment quality was limiting. We

therefore opted to collect further data on a Titan KRIOS electron microscope

(Thermo Fisher Scientific) operated at 300 kV acceleration voltage and equipped

with a Volta Phase Plate (VPP), a Gatan Quantum energy filter (operated at 20 eV

slit width), and a Gatan K2 Summit direct electron detector (operated in super-resolution counting mode). VPP data (datasets 1-6 in Supplementary Table 1) were collected according to the defocus acquisition technique (Danev & Baumeister, 2017; Khoshouei et al., 2017) at 43,478 x magnification, resulting in a physical pixel size of 1.15 Å on the object scale, with a total electron exposure of 50 e⁻ Å^{-2} at an exposure rate of 6.1 e⁻ Å^{-2} s^{-1} during an exposure time of 8.25 s, dose fractionated into 33 movie frames (50 frames for dataset 6). Data collection was monitored on-the-fly using FOCUS (Biyani et al., 2017) to ensure proper evolution of the VPP-induced phase shift.

Cryo-EM data processing

Initially, we used data collected in 10 microscopy sessions, 6 sessions using the VPP and 4 sessions without the VPP, resulting in > 30'000 total micrographs, of which approx. 15,000 were retained after inspection of the quality of Thon rings and CTF fitting (for details, see Supplementary Table 1). Movie stacks were aligned and dose weighed using MOTIONCOR2 (Zheng et al., 2017). The aligned, dose weighed sums from the datasets collected at 1.32 Å pixel size (datasets 7-10) were up-sampled to 1.15 Å per pixel to match the scale of the micrographs collected using the VPP (datasets 1-6) after calibrating the two magnifications to each other based on 3D reconstructions computed from the two types of data. CTF parameters were estimated using GCTF (Zhang, 2016) and particles were picked using GAUTOMATCH (Kai Zhang, MRC Laboratory of Molecular Biology, Cambridge UK) or RELION (Scheres, 2015) using templates generated from a preliminary run without reference templates. All subsequent data processing was performed in RELION 2 (Kimanius et al., 2016; Scheres, 2012) or RELION 3 (Nakane et al., 2018).

To remove false positive particle picks and broken particles, initial 3D classification was performed on each dataset individually (datasets 3, 4, 5), or on a few pooled datasets if appropriate (datasets 1 and 2 were pooled as they used the same batch of specimen; the non-VPP datasets 7-10 were joined because only few micrographs were retained due to more stringent quality criteria compared

to our previous study; and dataset 6 was initially classified together with
particles from dataset 4 to compensate for particle orientation bias, see Figure 1
– figure supplement 1). In summary, a total of > 2,000,000 initial particle picks
were subjected to this initial 3D classification, identifying approx. 820,000 intact
particles that were subjected to further processing. After 3D auto-refinement and
another round of 3D classification, performed separately for the VPP and non-
VPP data because the two data types were spuriously separated into distinct
classes in combined RELION 3D classification runs, the best classes (one from
VPP and non-VPP data each) resulting from the high-resolution 3D classifications
were refined according to the gold-standard refinement procedure (fully
independent half-sets), resulting in a 3.9 Å-resolution reconstruction according
to the FSC = 0.143 criterion (Rosenthal & Henderson, 2003; Scheres & Chen,
2012). Beam tilt refinement in RELION 3 improved the map computed from the
VPP data to 3.7 Å resolution. The non-VPP data no longer improved the
reconstruction after beam tilt correction and was therefore discarded at this
point. The final map was post-processed by application of b-factor of -175 A^2 and
low-pass filtration to the nominal 3.7 Å resolution for visualization and later
coordinate refinement.
700
To facilitate the interpretation of less ordered or only partially occupied parts of
the structure, including the p62 BSD2 domain, the MAT1 RING domain, the MAT1
three-helix bundle at the XPD arch domain, and the N-terminus of XPD, we used
signal subtracted classification (Bai et al., 2015; Nguyen et al., 2015) (p62 BSD2
domain, Figure 1 – figure supplement 3), focused classification (MAT1 RING
domain, Figure 1 – figure supplement 4), and multibody refinement (Nakane et
al., 2018) (MAT1 three-helix bundle and XPD N-terminus, Figure 1 – figure
supplement 5). For these classification procedures, we used only the VPP data
because 3D classification separated VPP and conventional cryo-EM data into
distinct classes, rendering combined classification ineffective. Multibody
refinement led to only a slight improvement in resolution (to 3.6 Å) relative to
the overall refined best map, and only during the first two iterations, likely due
to the relatively small size of the individual bodies and the resulting limited
signal for alignment. However, the above-mentioned structural elements showed

improved density features (Figure 1 – figure supplement 1B) and could be more

reliably interpreted in the multibody-refined XPD-MAT1 map (green in Figure 1

– figure supplement 5; also used in Figure 1 – figure supplement 2E, F). Overall,

the use of VPP data in this work resulted in substantial improvements both in

nominal resolution (Figure 1 – figure supplement 2D) and map quality (Figure 1

– figure supplement 2E-G) compared to our previous 4.4 Å-resolution structure.

Model building and refinement

The previous structure of the human TFIIH core complex (Greber et al., 2017)

and of yeast TFIIH in the Pol II-PIC (Schilbach et al., 2017) were docked into the

cryo-EM map and used as the basis for atomic modeling in O (Jones et al., 1991)

and COOT (Emsley et al., 2010). In addition to these models, the structure of the

human p34 VWFA-p44 RING domain complex (Radu et al., 2017), the N-terminal

RING domain of MAT1 (Gervais et al., 2001), the C-terminal RecA-like domain of

human XPB (Hilario et al., 2013) and several homology models for the p52

winged-helix domains generated using the PHYRE2 web server (Kelley et al.,

2015) based on templates PDB IDs 3F6O and 1STZ (Liu et al., 2005) were used

for model building.

Building of 52: The structure of p52 was traced and built completely *de novo*,

with the exception of the winged-helix domains and the very C-terminal domain,

where a homology model was placed into the cryo-EM map together with the p8

structure (Kainov et al., 2008; Vitorino et al., 2007) and adjusted to the density.

XPB: The structure of the XPB NTD was also built *de novo*, the structure of the

DRD was extensively re-built, and the RecA-like domains were rebuilt to improve

the fit to the density.

XPD: The improved cryo-EM map enabled detailed re-building of XPD, including

correction of register shifts in the more poorly ordered regions of the protein

and extension of the termini.

MAT1: The N-terminal MAT1 RING domain (Gervais et al., 2001) was first docked into the focused classified density and combined with the rest of the TFIIH model, were it helped guide the assignment of the sequence register to the MAT1 model, in combination with density features of large side chains in the helical regions.

Building of p34: The human p34 structure (Radu et al., 2017) was docked into the map as is, extended near the interaction site with p52, and combined with a completely re-built model of the C-terminal eZnF domain (Schilbach et al., 2017).

Building of p44: The p44 VWFA fold needed only minor rebuilding and was combined with the eZnF domain and the C-terminal human RING domain model (Radu et al., 2017; Schilbach et al., 2017). The p44 NTE was built according to the density at the contact site with XPB and guided by CX-MS data (Luo et al., 2015). Both the features of the cryo-EM map and crosslinks of the p44 NTE to p34, p52, and XPB (Figure 1 – figure supplement 6E) unambiguously confirm the tracing of this segment towards the XPB NTD, rather than alternative tracing towards XPD (this density is now assigned to p62, in agreement with p62-XPD crosslinks; see below and Figure 1 – figure supplement 6G).

Building of p62: The p62 protein was modeled based on the placement of the BSD domains (PDB ID 2DII), secondary structure prediction, extension of docked coordinates (Greber et al., 2017; Schilbach et al., 2017), and new tracing of the protein chain. Placement of the regions near XPD, where density is weak overall, was guided by matching the succession of secondary structure elements along the p62 sequence with helical densities in the cryo-EM map (Figure 2 – figure supplement 1A), and corroborated by CX-MS data (Luo et al., 2015), which showed excellent agreement of p62-XPD crosslinking data with the structure (Figure 1 – figure supplement 6G). Crosslinks between p62 and p44 showed a relatively large proportion of outliers (Supplementary Data Table 1), which may be due to the fact that the sequence register of the p62 segments to which these crosslinks map is not well constrained. These segments were modeled as poly-alanines and deposited without sequence assignment (UNK; Supplementary

Table 2). Maps low-pass filtered to 6 Å and sharpened by a b-factor of only -100

Å^2 were used to guide docking of domains and assess the continuity of the

density in poorly ordered regions of the protein (Figure 2 – figure supplement

1B, C).

The resulting coordinate model (Supplementary Table 2) was refined using the

real space refinement program in PHENIX (Adams et al., 2010; Afonine et al.,

2018) and validated using the MTRIAGE program in PHENIX and the

MOLPROBITY web server (Chen et al., 2010). Ramachandran, C$_\beta$, rotamer, and

secondary structure restraints were used throughout the refinement to ensure

good model geometry at the given resolution. Data used in the refinement

excluded spatial frequencies beyond the nominal 3.7 Å resolution of the cryo-EM

map to prevent over-fitting. The N-terminal RING domain of MAT1 and the BSD1

domain of p62, for which only poorly resolved density is present in the final

cryo-EM map, were additionally restrained by reference restraints (Headd et al.,

2012) using the NMR structures of the corresponding domains (PDB ID 1G25

and 2DII, respectively) (Gervais et al., 2001). The side chains of these two

domains (with the exception of residues involved in zinc finger formation and of

prolines) were truncated at the Cβ position to reflect the lower resolution of the

corresponding densities. The FSC curve between the refined coordinate model

and the cryo-EM map extends to 3.9 Å and the distribution of b-factors in the

refined coordinate model (Figure 1 – figure supplement 2C) mirror the local

resolution of the cryo-EM map (Figure 1 – figure supplement 2D), as expected.

Refinement statistics are given in Supplementary Table 3 and are typical for

structures in this resolution range (97[th] and 99[th] percentiles for MOLPROBITY

clash score and overall score (Chen et al., 2010)).

Flexibility analysis

For the analysis of conformational dynamics of TFIIH, VPP datasets 1 and 2 were

subjected to multibody refinement in RELION 3 (Nakane et al., 2018) using 6

masks (Figure 4 – figure supplement 1). The volume series for the first 16

principal components were reconstructed and difference densities (green and

purple in Figure 4 – figure supplement 1) were computed between the most extreme states in each series (blue and yellow in Figure 4 – figure supplement 1). Subsequently, roughly 20,000 particles corresponding to both ends of the distribution were used for selected principal components and subjected to 3D refinement, resulting in maps of approx. 10 Å resolution. The refined atomic model of the TFIIH core complex, subdivided into suitable rigid bodies, was then rigid-body refined into these volumes using PHENIX real space refinement (Afonine et al., 2018) and coordinate displacement between the two resulting models for each principal component was plotted to obtain an initial assessment of the modes of motion present in the TFIIH dataset (Figure 4A). For actual structural interpretation (Figure 4B, C), the final cryo-EM maps of the TFIIH core complex (this work) and TFIIH in the context of the Pol II-PIC (Schilbach et al., 2017) were used.

Other

Figures were created using PyMol (The PyMOL Molecular Graphics System, Version 1.8 Schrödinger, LLC.) and the UCSF Chimera package from the Computer Graphics Laboratory, University of California, San Francisco (supported by NIH P41 RR-01081) (Pettersen et al., 2004). Protein-protein interface statistics were determined using PISA (Krissinel & Henrick, 2007).

ACKNOWLEDGMENTS

We thank D. Wigley for sharing of structural data, S. Scheres for sharing computer code before publication, R. Tjian, S. Zheng, and D. King for supplying XPB mAB and peptides for TFIIH purification, P. Grob and A. Chintangal for support with microscopy and computing, respectively, and Z. Yu and C. Hong for data collection at the Janelia farm cryo-EM facility. We acknowledge the use of the resources of the National Energy Research Scientific Computing Center (NERSC), a DOE Office of Science user facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

ADDITIONAL INFORMATION

Funding

This work was funded through NIGMS grants R01-GM63072 and P01-GM063210

to E.N.. B.J.G. was supported by fellowships from the Swiss National Science

Foundation (projects P300PA_160983, P300PA_174355). E.N. is a Howard

Hughes medical investigator.

Author contributions

E.N. directed the study. J.F. performed HeLa cell culture and purified TFIIH. B.J.G.

performed cryo-EM specimen preparation, data processing, atomic model

building, and coordinate refinement. D.B.T and B.J.G collected the cryo-EM data.

B.J.G. wrote the initial draft of the manuscript and all authors contributed to the

final version.

Supplementary Data Table 1 (separate file)

Result summary of mapping of CX-MS data (Luo et al., 2015) onto the TFIIH

structure.

REFERENCES

Abdulrahman W., Iltis I., Radu L., Braun C., Maglott-Roth A., Giraudon C., Egly J.-
M., Poterszman A. (2013) ARCH domain of XPD, an anchoring platform for CAK
that conditions TFIIH DNA repair and transcription activities. *Proc Natl Acad Sci*
USA **110:** E633-642

Adams P. D., Afonine P. V., Bunkóczi G., Chen V. B., Davis I. W., Echols N., Headd J.
J., Hung L. W., Kapral G. J., Grosse-Kunstleve R. W. et al (2010) PHENIX: a
comprehensive Python-based system for macromolecular structure solution.
Acta Crystallogr D Biol Crystallogr **66:** 213-221

Afonine P. V., Poon B. K., Read R. J., Sobolev O. V., Terwilliger T. C., Urzhumtsev A.,
Adams P. D. (2018) Real-space refinement in PHENIX for cryo-EM and
crystallography. *Acta Crystallogr D Struct Biol* **74:** 531-544

Alekseev S., Nagy Z., Sandoz J., Weiss A., Egly J.-M., Le May N., Coin F. (2017)
Transcription without XPB Establishes a Unified Helicase-Independent
Mechanism of Promoter Opening in Eukaryotic Gene Expression. *Mol Cell* **65:**
504-513.e505

Araújo S. J., Tirode F., Coin F., Pospiech H., Syväoja J. E., Stucki M., Hübscher U.,
Egly J. M., Wood R. D. (2000) Nucleotide excision repair of DNA with
recombinant human proteins: definition of the minimal set of factors, active
forms of TFIIH, and modulation by CAK. *Genes & Dev* **14:** 349-359

Bai X.-C., Rajendra E., Yang G., Shi Y., Scheres S. H. (2015) Sampling the
conformational space of the catalytic subunit of human γ -secretase. *Elife* **4:**
e11182

Balasingham S. V., Zegeye E. D., Homberset H., Rossi M. L., Laerdahl J. K., Bohr V.
A., Tønjum T. (2012) Enzymatic activities and DNA substrate specificity of
Mycobacterium tuberculosis DNA helicase XPB. *PLoS ONE* **7:** e36960

Berico P., Coin F. (2017) Is TFIIH the new Achilles heel of cancer cells?
Transcription **9:** 47-51

Bienstock R. J., Skorvaga M., Mandavilli B. S., Van Houten B. (2003) Structural and
functional characterization of the human DNA repair helicase XPD by
comparative molecular modeling and site-directed mutagenesis of the bacterial
repair protein UvrB. *J Biol Chem* **278:** 5309-5316

Biyani N., Righetto R. D., McLeod R., Caujolle-Bert D., Castaño-Diez D., Goldie K.
N., Stahlberg H. (2017) Focus: The interface between data collection and data
processing in cryo-EM. *J Struct Biol* **198:** 124-133

Botta E., Nardo T., Lehmann A. R., Egly J.-M., Pedrini A. M., Stefanini M. (2002)
Reduced level of the repair/transcription factor TFIIH in trichothiodystrophy.
Hum Mol Genet **11:** 2919-2928

Buechner C. N., Heil K., Michels G., Carell T., Kisker C., Tessmer I. (2014) Strand-
specific recognition of DNA damages by XPD provides insights into nucleotide
excision repair substrate versatility. *J Biol Chem* **289:** 3613-3624

Busso D., Keriel A., Sandrock B., Poterszman A., Gileadi O., Egly J. M. (2000)
Distinct regions of MAT1 regulate cdk7 kinase and TFIIH transcription activities.
J Biol Chem **275:** 22815-22823

Büttner K., Nehring S., Hopfner K.-P. (2007) Structural basis for DNA duplex
separation by a superfamily-2 helicase. *Nat Struct Mol Biol* **14:** 647-652

Chen V. B., Arendall W. B., Headd J. J., Keedy D. A., Immormino R. M., Kapral G. J.,
Murray L. W., Richardson J. S., Richardson D. C. (2010) MolProbity: all-atom
structure validation for macromolecular crystallography. *Acta Crystallogr D Biol
Crystallogr* **66:** 12-21

Cheng K., Wigley D. B. (2018) DNA translocation mechanism of an XPD family
helicase. *Elife* **7**

Clarkson S. G., Wood R. D. (2005) Polymorphisms in the human XPD (ERCC2)
gene, DNA repair capacity and cancer susceptibility: an appraisal. *DNA Repair*
(Amst) **4:** 1068-1074

Cleaver J. E., Thompson L. H., Richardson A. S., States J. C. (1999) A summary of
mutations in the UV-sensitive disorders: xeroderma pigmentosum, Cockayne
syndrome, and trichothiodystrophy. *Hum Mutat* **14:** 9-22

Coin F., Marinoni J. C., Rodolfo C., Fribourg S., Pedrini A. M., Egly J. M. (1998)
Mutations in the XPD helicase gene result in XP and TTD phenotypes, preventing
interaction between XPD and the p44 subunit of TFIIH. *Nat Genet* **20:** 184-188

Coin F., Oksenych V., Egly J.-M. (2007) Distinct roles for the XPB/p52 and
XPD/p44 subcomplexes of TFIIH in damaged DNA opening during nucleotide
excision repair. *Mol Cell* **26:** 245-256

Coin F., Oksenych V., Mocquet V., Groh S., Blattner C., Egly J.-M. (2008) Nucleotide
excision repair driven by the dissociation of CAK from TFIIH. *Mol Cell* **31:** 9-20

Coin F., Proietti De Santis L., Nardo T., Zlobinskaya O., Stefanini M., Egly J.-M.
(2006) p8/TTD-A as a repair-specific TFIIH subunit. *Mol Cell* **21:** 215-226

Compe E., Egly J.-M. (2016) Nucleotide Excision Repair and Transcriptional
Regulation: TFIIH and Beyond. *Annu Rev Biochem* **85:** 265-290

Constantinescu-Aruxandei D., Petrovic-Stojanovska B., Penedo J. C., White M. F.,
Naismith J. H. (2016) Mechanism of DNA loading by the DNA repair helicase XPD.
Nucleic Acids Res **44:** 2806-2815

Danev R., Baumeister W. (2017) Expanding the boundaries of cryo-EM with
phase plates. *Curr Opin Struct Biol* **46:** 87-94

Devault A., Martinez A. M., Fesquet D., Labbé J. C., Morin N., Tassan J. P., Nigg E. A.,
Cavadore J. C., Dorée M. (1995) MAT1 ('menage à trois') a new RING finger
protein subunit stabilizing cyclin H-cdk7 complexes in starfish and Xenopus CAK.
EMBO J **14:** 5027-5036

Di Lello P., Jenkins L. M. M., Jones T. N., Nguyen B. D., Hara T., Yamaguchi H.,
Dikeakos J. D., Appella E., Legault P., Omichinski J. G. (2006) Structure of the
Tfb1/p53 complex: Insights into the interaction between the p62/Tfb1 subunit
of TFIIH and the activation domain of p53. *Mol Cell* **22:** 731-740

Di Lello P., Miller Jenkins L. M., Mas C., Langlois C., Malitskaya E., Fradet-Turcotte
A., Archambault J., Legault P., Omichinski J. G. (2008) p53 and TFIIEalpha share a
common binding site on the Tfb1/p62 subunit of TFIIH. *Proc Natl Acad Sci USA*
105: 106-111

Drapkin R., Le Roy G., Cho H., Akoulitchev S., Reinberg D. (1996) Human cyclin-
dependent kinase-activating kinase exists in three distinct complexes. *Proc Natl*
Acad Sci USA **93:** 6488-6493

Dubaele S., Proietti De Santis L., Bienstock R. J., Keriel A., Stefanini M., Van Houten
B., Egly J.-M. (2003) Basal transcription defect discriminates between xeroderma
pigmentosum and trichothiodystrophy in XPD patients. *Mol Cell* **11:** 1635-1646

Emsley P., Lohkamp B., Scott W. G., Cowtan K. (2010) Features and development
of Coot. *Acta Crystallogr D Biol Crystallogr* **66:** 486-501

Evans E., Moggs J. G., Hwang J. R., Egly J. M., Wood R. D. (1997) Mechanism of
open complex and dual incision formation by human nucleotide excision repair
factors. *EMBO J* **16:** 6559-6573

Fairman-Williams M. E., Guenther U.-P., Jankowsky E. (2010) SF1 and SF2
helicases: family matters. *Curr Opin Struct Biol* **20:** 313-324

Fan L., Arvai A. S., Cooper P. K., Iwai S., Hanaoka F., Tainer J. A. (2006) Conserved
XPB core structure and motifs for DNA unwinding: implications for pathway
selection of transcription or excision repair. *Mol Cell* **22:** 27-37

Fan L., Fuss J. O., Cheng Q. J., Arvai A. S., Hammel M., Roberts V. A., Cooper P. K.,
Tainer J. A. (2008) XPD helicase structures and activities: insights into the cancer
and aging phenotypes from XPD mutations. *Cell* **133:** 789-800

Fishburn J., Tomko E., Galburt E., Hahn S. (2015) Double-stranded DNA
translocase activity of transcription factor TFIIH and the mechanism of RNA
polymerase II open complex formation. *Proc Natl Acad Sci USA* **112:** 3961-3966

Fisher R. P., Jin P., Chamberlin H. M., Morgan D. O. (1995) Alternative
mechanisms of CAK assembly require an assembly factor or an activating kinase.
Cell **83:** 47-57

Fisher R. P., Morgan D. O. (1994) A novel cyclin associates with MO15/CDK7 to
form the CDK-activating kinase. *Cell* **78:** 713-724

Fregoso M., Lainé J.-P., Aguilar-Fuentes J., Mocquet V., Reynaud E., Coin F., Egly J.-
M., Zurita M. (2007) DNA repair and transcriptional deficiencies caused by
mutations in the Drosophila p52 subunit of TFIIH generate developmental
defects and chromosome fragility. *Mol Cell Biol* **27:** 3640-3650

Gervais V., Busso D., Wasielewski E., Poterszman A., Egly J. M., Thierry J. C.,
Kieffer B. (2001) Solution structure of the N-terminal domain of the human
TFIIH MAT1 subunit: new insights into the RING finger family. *J Biol Chem* **276:**
7457-7464

Gervais V., Lamour V., Jawhari A., Frindel F., Wasielewski E., Dubaele S., Egly J.-M.,
Thierry J.-C., Kieffer B., Poterszman A. (2004) TFIIH contains a PH domain
involved in DNA nucleotide excision repair. *Nat Struct Mol Biol* **11:** 616-622

Gibbons B. J., Brignole E. J., Azubel M., Murakami K., Voss N. R., Bushnell D. A.,
Asturias F. J., Kornberg R. D. (2012) Subunit architecture of general transcription
factor TFIIH. *Proc Natl Acad Sci USA* **109:** 1949-1954

Giglia-Mari G., Coin F., Ranish J. A., Hoogstraten D., Theil A., Wijgers N., Jaspers N.
G. J., Raams A., Argentini M., van der Spek P. J. et al (2004) A new, tenth subunit of
TFIIH is responsible for the DNA repair syndrome trichothiodystrophy group A.
Nat Genet **36:** 714-719

Greber B. J., Nguyen T. H. D., Fang J., Afonine P. V., Adams P. D., Nogales E. (2017)
The cryo-electron microscopy structure of human transcription factor IIH.
Nature **549:** 414-417

Grünberg S., Warfield L., Hahn S. (2012) Architecture of the RNA polymerase II
preinitiation complex and mechanism of ATP-dependent promoter opening. *Nat*
Struct Mol Biol **19:** 788-796

Guzder S. N., Qiu H., Sommers C. H., Sung P., Prakash L., Prakash S. (1994) DNA
repair gene RAD3 of S. cerevisiae is essential for transcription by RNA
polymerase II. *Nature* **367:** 91-94

He Y., Fang J., Taatjes D. J., Nogales E. (2013) Structural visualization of key steps
in human transcription initiation. *Nature* **495:** 481-486

He Y., Yan C., Fang J., Inouye C., Tjian R., Ivanov I., Nogales E. (2016) Near-atomic
resolution visualization of human transcription promoter opening. *Nature* **533:**
359-365

Headd J. J., Echols N., Afonine P. V., Grosse-Kunstleve R. W., Chen V. B., Moriarty
N. W., Richardson D. C., Richardson J. S., Adams P. D. (2012) Use of knowledge-
based restraints in phenix.refine to improve macromolecular refinement at low
resolution. *Acta Crystallogr D Biol Crystallogr* **68:** 381-390

Herrera-Moyano E., Moriel-Carretero M., Montelone B. A., Aguilera A. (2014) The
rem mutations in the ATP-binding groove of the Rad3/XPD helicase lead to
Xeroderma pigmentosum-Cockayne syndrome-like phenotypes. *PLoS Genet* **10:**
e1004859

Hilario E., Li Y., Nobumori Y., Liu X., Fan L. (2013) Structure of the C-terminal half
of human XPB helicase and the impact of the disease-causing mutation XP11BE.
Acta Crystallogr D Biol Crystallogr **69:** 237-246

Huang J. C., Svoboda D. L., Reardon J. T., Sancar A. (1992) Human nucleotide
excision nuclease removes thymine dimers from DNA by incising the 22nd
phosphodiester bond 5' and the 6th phosphodiester bond 3' to the photodimer.
Proc Natl Acad Sci USA **89:** 3664-3668

Jawhari A., Boussert S., Lamour V., Atkinson R. A., Kieffer B., Poch O., Potier N.,
Van Dorsselaer A., Moras D., Poterszman A. (2004) Domain architecture of the
p62 subunit from the human transcription/repair factor TFIIH deduced by

limited proteolysis and mass spectrometry analysis. *Biochemistry* **43:** 14420-
14430

Jawhari A., Lainé J.-P., Dubaele S., Lamour V., Poterszman A., Coin F., Moras D.,
Egly J.-M. (2002) p52 Mediates XPB function within the transcription/repair
factor TFIIH. *J Biol Chem* **277:** 31761-31767

Jones T. A., Zou J. Y., Cowan S. W., Kjeldgaard M. (1991) Improved methods for
building protein models in electron density maps and the location of errors in
these models. *Acta Crystallogr A Found Crystallogr* **47 (Pt. 2):** 110-119

Kainov D. E., Vitorino M., Cavarelli J., Poterszman A., Egly J.-M. (2008) Structural
basis for group A trichothiodystrophy. *Nat Struct Mol Biol* **15:** 980-984

Kelley L. A., Mezulis S., Yates C. M., Wass M. N., Sternberg M. J. E. (2015) The
Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc* **10:**
845-858

Khoshouei M., Radjainia M., Baumeister W., Danev R. (2017) Cryo-EM structure
of haemoglobin at 3.2 Å determined with the Volta phase plate. *Nat Commun* **8:**
16099

Kim J. S., Saint-André C., Lim H. S., Hwang C.-S., Egly J.-M., Cho Y. (2015) Crystal
structure of the Rad3/XPD regulatory domain of Ssl1/p44. *J Biol Chem* **290:**
8321-8330

Kimanius D., Forsberg B. O., Scheres S. H., Lindahl E. (2016) Accelerated cryo-EM
structure determination with parallelisation using GPUs in RELION-2. *Elife* **5:**
e18722

Kralund H. H., Ousager L., Jaspers N. G., Raams A., Pedersen E. B., Gade E., Bygum
A. (2013) Xeroderma Pigmentosum-Trichothiodystrophy overlap patient with
novel XPD/ERCC2 mutation. *Rare Dis* **1:** e24932

Krissinel E., Henrick K. (2007) Inference of macromolecular assemblies from
crystalline state. *J Mol Biol* **372:** 774-797

Kuper J., Braun C., Elias A., Michels G., Sauer F., Schmitt D. R., Poterszman A., Egly
J.-M., Kisker C. (2014) In TFIIH, XPD helicase is exclusively devoted to DNA
repair. *PLoS Biol* **12:** e1001954

Kuper J., Wolski S. C., Michels G., Kisker C. (2012) Functional and structural
studies of the nucleotide excision repair helicase XPD suggest a polarity for DNA
translocation. *EMBO J* **31:** 494-502

Lafrance-Vanasse J., Arseneault G., Cappadocia L., Legault P., Omichinski J. G.
(2013) Structural and functional evidence that Rad4 competes with Rad2 for
binding to the Tfb1 subunit of TFIIH in NER. *Nucleic Acids Res* **41:** 2736-2745

Li C.-L., Golebiowski F. M., Onishi Y., Samara N. L., Sugasawa K., Yang W. (2015)
Tripartite DNA Lesion Recognition and Verification by XPC, TFIIH, and XPA in
Nucleotide Excision Repair. *Mol Cell* **59:** 1025-1034

Liu H., Rudolf J., Johnson K. A., McMahon S. A., Oke M., Carter L., McRobbie A.-M.,
Brown S. E., Naismith J. H., White M. F. (2008) Structure of the DNA Repair
Helicase XPD. *Cell* **133:** 801-812

Liu J., Huang C., Shin D.-H., Yokota H., Jancarik J., Kim J.-S., Adams P. D., Kim R.,
Kim S.-H. (2005) Crystal structure of a heat-inducible transcriptional repressor
HrcA from Thermotoga maritima: structural insight into DNA binding and
dimerization. *J Mol Biol* **350:** 987-996

Luo J., Cimermančič P., Viswanath S., Ebmeier C. C., Kim B., Dehecq M., Raman V.,
Greenberg C. H., Pellarin R., Sali A. et al (2015) Architecture of the Human and
Yeast General Transcription and DNA Repair Factor TFIIH. *Mol Cell* **59:** 794-806

Mason A. C., Rambo R. P., Greer B., Pritchett M., Tainer J. A., Cortez D., Eichman B.
F. (2014) A structure-specific nucleic acid-binding domain conserved among
DNA repair proteins. *Proc Natl Acad Sci USA* **111:** 7618-7623

Mathieu N., Kaczmarek N., Rüthemann P., Luch A., Naegeli H. (2013) DNA quality
control by a lesion sensor pocket of the xeroderma pigmentosum group D
helicase subunit of TFIIH. *Curr Biol* **23:** 204-212

Mlak R., Krawczyk P., Homa-Mlak I., Powrózek T., Ciesielka M., Kozioł P.,
Milanowski J., Małecka-Massalska T. (2018) Predictive Value of Single Nucleotide
Polymorphisms of ERCC1, XPA, XPC, XPD and XPG Genes, Involved in NER
Mechanism in Patients with Advanced NSCLC Treated with Cisplatin and
Gemcitabine. *Pathol Oncol Res***:** 1-11

Moriel-Carretero M., Aguilera A. (2010) A postincision-deficient TFIIH causes
replication fork breakage and uncovers alternative Rad51- or Pol32-mediated
restart mechanisms. *Mol Cell* **37:** 690-701

Moriel-Carretero M., Herrera-Moyano E., Aguilera A. (2015) A unified model for
the molecular basis of Xeroderma pigmentosum-Cockayne Syndrome. *Rare Dis* **3:**
e1079362

Murakami K., Tsai K.-L., Kalisman N., Bushnell D. A., Asturias F. J., Kornberg R. D.
(2015) Structure of an RNA polymerase II preinitiation complex. *Proc Natl Acad*
Sci USA **112:** 13543-13548

Naegeli H., Bardwell L., Friedberg E. C. (1993) Inhibition of Rad3 DNA helicase
activity by DNA adducts and abasic sites: implications for the role of a DNA
helicase in damage-specific incision of DNA. *Biochemistry* **32:** 613-621

Nakane T., Kimanius D., Lindahl E., Scheres S. H. (2018) Characterisation of
molecular motions in cryo-EM single-particle data by multi-body refinement in
RELION. *Elife* **7:** e36861

Nguyen T. H. D., Galej W. P., Bai X.-C., Oubridge C., Newman A. J., Scheres S. H. W.,
Nagai K. (2016) Cryo-EM structure of the yeast U4/U6.U5 tri-snRNP at 3.7 Å
resolution. *Nature* **530:** 298-302

Nguyen T. H. D., Galej W. P., Bai X.-C., Savva C. G., Newman A. J., Scheres S. H. W.,
Nagai K. (2015) The architecture of the spliceosomal U4/U6.U5 tri-snRNP.
Nature **523:** 47-52

Nguyen T. H. D., Li J., Galej W. P., Oshikane H., Newman A. J., Nagai K. (2013)
Structural basis of Brr2-Prp8 interactions and implications for U5 snRNP
biogenesis and the spliceosome active site. *Structure* **21:** 910-919

Obmolova G., Ban C., Hsieh P., Yang W. (2000) Crystal structures of mismatch
repair protein MutS and its complex with a substrate DNA. *Nature* **407:** 703-710

Oksenych V., Bernardes de Jesus B., Zhovmer A., Egly J.-M., Coin F. (2009)
Molecular insights into the recruitment of TFIIH to sites of DNA damage. *EMBO J*
28: 2971-2980

Okuda M., Nakazawa Y., Guo C., Ogi T., Nishimura Y. (2017) Common TFIIH
recruitment mechanism in global genome and transcription-coupled repair
subpathways. *Nucleic Acids Res* **45:** 13043-13055

Peters G. J., Avan A., Ruiz M. G., Orsini V., Avan A., Giovannetti E., Smit E. F. (2014)
Predictive role of repair enzymes in the efficacy of Cisplatin combinations in
pancreatic and lung cancer. *Anticancer Res* **34:** 435-442

Pettersen E. F., Goddard T. D., Huang C. C., Couch G. S., Greenblatt D. M., Meng E.
C., Ferrin T. E. (2004) UCSF Chimera--a visualization system for exploratory
research and analysis. *J Comput Chem* **25:** 1605-1612

Poterszman A., Lamour V., Egly J. M., Moras D., Thierry J. C., Poch O. (1997) A
eukaryotic XPB/ERCC3-like helicase in Mycobacterium leprae? *Trends Biochem*
Sci **22:** 418-419

Radu L., Schoenwetter E., Braun C., Marcoux J., Koelmel W., Schmitt D. R., Kuper J.,
Cianférani S., Egly J. M., Poterszman A. et al (2017) The intricate network
between the p34 and p44 subunits is central to the activity of the
transcription/DNA repair factor TFIIH. *Nucleic Acids Res* **45:** 10872-10883

Rapin I. (2013) Disorders of nucleotide excision repair. *Handb Clin Neurol* **113:**
1637-1650

Riou L., Zeng L., Chevallier-Lagente O., Stary A., Nikaido O., Taïeb A., Weeda G.,
Mezzina M., Sarasin A. (1999) The relative expression of mutated XPB genes
results in xeroderma pigmentosum/Cockayne's syndrome or
trichothiodystrophy cellular phenotypes. *Hum Mol Genet* **8:** 1125-1133

Robert X., Gouet P. (2014) Deciphering key features in protein structures with
the new ENDscript server. *Nucleic Acids Res* **42:** W320-324

Robinson P. J., Trnka M. J., Bushnell D. A., Davis R. E., Mattei P.-J., Burlingame A. L.,
Kornberg R. D. (2016) Structure of a Complete Mediator-RNA Polymerase II Pre-
Initiation Complex. *Cell* **166:** 1411-1422.e1416

Rosenthal P. B., Henderson R. (2003) Optimal determination of particle
orientation, absolute hand, and contrast loss in single-particle electron
cryomicroscopy. *J Mol Biol* **333:** 721-745

Rossignol M., Kolb-Cheynel I., Egly J. M. (1997) Substrate specificity of the cdk-
activating kinase (CAK) is altered upon association with TFIIH. *EMBO J* **16:** 1628-
1637

Rouillon C., White M. F. (2010) The XBP-Bax1 helicase-nuclease complex
unwinds and cleaves DNA: implications for eukaryal and archaeal nucleotide
excision repair. *J Biol Chem* **285:** 11013-11022

Rudolf J., Makrantoni V., Ingledew W. J., Stark M. J. R., White M. F. (2006) The
DNA repair helicases XPD and FancJ have essential iron-sulfur domains. *Mol Cell*
23: 801-808

Sainsbury S., Bernecky C., Cramer P. (2015) Structural basis of transcription
initiation by RNA polymerase II. *Nat Rev Mol Cell Biol* **16:** 129-143

Sandrock B., Egly J. M. (2001) A yeast four-hybrid system identifies Cdk-
activating kinase as a regulator of the XPD helicase, a subunit of transcription
factor IIH. *J Biol Chem* **276:** 35328-35333

Scheres S. H. W. (2010) Classification of Structural Heterogeneity by Maximum-
Likelihood Methods. *Meth Enz* **482:** 295-320

Scheres S. H. W. (2012) RELION: implementation of a Bayesian approach to cryo-
EM structure determination. *J Struct Biol* **180:** 519-530

Scheres S. H. W. (2015) Semi-automated selection of cryo-EM particles in
RELION-1.3. *J Struct Biol* **189:** 114-122

Scheres S. H. W., Chen S. (2012) Prevention of overfitting in cryo-EM structure
determination. *Nat Meth* **9:** 853-854

Schilbach S., Hantsche M., Tegunov D., Dienemann C., Wigge C., Urlaub H., Cramer
P. (2017) Structures of transcription pre-initiation complex with TFIIH and
Mediator. *Nature* **551:** 204-209

Seroz T., Perez C., Bergmann E., Bradsher J., Egly J. M. (2000) p44/SSL1, the
regulatory subunit of the XPD/RAD3 helicase, plays a crucial role in the
transcriptional activity of TFIIH. *J Biol Chem* **275:** 33260-33266

Shiekhattar R., Mermelstein F., Fisher R. P., Drapkin R., Dynlacht B., Wessling H.
C., Morgan D. O., Reinberg D. (1995) Cdk-activating kinase complex is a
component of human transcription factor TFIIH. *Nature* **374:** 283-287

Sugasawa K., Akagi J.-i., Nishi R., Iwai S., Hanaoka F. (2009) Two-step recognition
of DNA damage for mammalian nucleotide excision repair: Directional binding of
the XPC complex and DNA strand scanning. *Mol Cell* **36:** 642-653

Sugitani N., Sivley R. M., Perry K. E., Capra J. A., Chazin W. J. (2016) XPA: A key
scaffold for human nucleotide excision repair. *DNA Repair (Amst)* **44:** 123-135

Svejstrup J. Q., Wang Z., Feaver W. J., Wu X., Bushnell D. A., Donahue T. F.,
Friedberg E. C., Kornberg R. D. (1995) Different forms of TFIIH for transcription
and DNA repair: holo-TFIIH and a nucleotide excision repairosome. *Cell* **80:** 21-
28

Theron T., Fousteri M. I., Volker M., Harries L. W., Botta E., Stefanini M., Fujimoto
M., Andressoo J.-O., Mitchell J., Jaspers N. G. J. et al (2005) Transcription-
associated breaks in xeroderma pigmentosum group D cells from patients with
combined features of xeroderma pigmentosum and Cockayne syndrome. *Mol Cell*
Biol **25:** 8368-8378

Tirode F., Busso D., Coin F., Egly J. M. (1999) Reconstitution of the transcription
factor TFIIH: assignment of functions for the three enzymatic subunits, XPB, XPD,
and cdk7. *Mol Cell* **3:** 87-95

Tremeau-Bravard A., Perez C., Egly J. M. (2001) A role of the C-terminal part of
p44 in the promoter escape activity of transcription factor IIH. *J Biol Chem* **276:**
27693-27697

Tsai K.-L., Yu X., Gopalan S., Chao T.-C., Zhang Y., Florens L., Washburn M. P.,
Murakami K., Conaway R. C., Conaway J. W. et al (2017) Mediator structure and
rearrangements required for holoenzyme formation. *Nature Publishing Group***:** 1-
19

Vermeulen W., Rademakers S., Jaspers N. G., Appeldoorn E., Raams A., Klein B.,
Kleijer W. J., Hansen L. K., Hoeijmakers J. H. (2001) A temperature-sensitive
disorder in basal transcription and DNA repair in humans. *Nat Genet* **27:** 299-
303

Vitorino M., Coin F., Zlobinskaya O., Atkinson R. A., Moras D., Egly J.-M.,
Poterszman A., Kieffer B. (2007) Solution structure and self-association
properties of the p8 TFIIH subunit responsible for trichothiodystrophy. *J Mol Biol*
368: 473-480

Warfield L., Luo J., Ranish J., Hahn S. (2016) Function of conserved topological
regions within the S. cerevisiae basal transcription factor TFIIH. *Mol Cell Biol* **36:**
2464-2475

Winkler G. S., Araújo S. J., Fiedler U., Vermeulen W., Coin F., Egly J. M.,
Hoeijmakers J. H., Wood R. D., Timmers H. T., Weeda G. (2000) TFIIH with
inactive XPD helicase functions in transcription initiation but is defective in DNA
repair. *J Biol Chem* **275:** 4258-4266

Wirth N., Gross J., Roth H. M., Buechner C. N., Kisker C., Tessmer I. (2016)
Conservation and Divergence in Nucleotide Excision Repair Lesion Recognition. *J*
Biol Chem **291:** 18932-18946

Wolski S. C., Kuper J., Hänzelmann P., Truglio J. J., Croteau D. L., Van Houten B.,
Kisker C. (2008) Crystal structure of the FeS cluster-containing nucleotide
excision repair helicase XPD. *PLoS Biol* **6:** e149

Zhang K. (2016) Gctf: Real-time CTF determination and correction. *J Struct Biol*
193: 1-12

Zheng S. Q., Palovcak E., Armache J.-P., Verba K. A., Cheng Y., Agard D. A. (2017)
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. *Nat Meth* **14:** 331-332

Ziani S., Nagy Z., Alekseev S., Soutoglou E., Egly J.-M., Coin F. (2014) Sequential
and ordered assembly of a large DNA repair complex on undamaged chromatin. *J*
Cell Biol **206:** 589-598
1237

1238

1239

Figure supplements

Figure 1 - figure supplement 1. Data processing scheme for global reconstruction. The datasets (DS) were initially processed individually, with the exception of DS6, where particles from DS4 were used to compensate for orientation bias, and DS1/DS2, which were joined for processing. Subsequently, the datasets were pooled and sub-classified. The best class of was used for the final refinement to obtain a globally refined map at 3.7 Å resolution (see Figure 1 – figure supplement 2). The classified VPP dataset was also used for further processing and classification of sub-regions (Figure 1 – figure supplements 3-5).

1253

Figure 1 – figure supplement 2. Resolution estimation, validation statistics,

and quality of the density. (**A**) Three views of the refined, sharpened, and low-

pass filtered cryo-EM map of the TFIIH core complex. Protein subunits are color-

coded and labeled. (**B**) Fourier shell correlation (FSC) curves. Black: FSC curve

between half-maps. Red: FSC curve between map and refined model. The

resolution at the appropriate thresholds (Rosenthal & Henderson, 2003; Scheres

& Chen, 2012) is indicated. (**C**) B-factor distribution on the refined TFIIH

coordinate model. (**D**) Local resolution estimation, computed by windowed FSC

in RELION 2. (**E**, **F**, **G**) Comparison of the 3.6 Å multibody-refined cryo-EM map of

the TFIIH core complex (E, F) with our previous 4.4 Å cryo-EM map (G) shows

substantial improvements in map quality in the more peripheral and flexible

regions of the complex (an α-helix from the XPD ARCH domain is shown).

1266

1267

Figure 1 – figure supplement 3. Focused classification of the p62 BSD2
domain. (A) Focused classification for the p62 BSD2 domain and adjacent
structural elements using the classified VPP dataset (indicated in Figure 1 –

figure supplement 1). Partial signal subtraction (Bai et al., 2015) and masking

were used to focus the classification on the region of interest (mask shown in

transparent yellow). Selected classes were refined using the selected but un-

subtracted particle images because the region of interest is too small for efficient

image alignment (Nguyen et al., 2016). (**B**) FSC curves for the three refined maps

shown in (A), indicating resolution of 3.9-4.3 Å according to the FSC = 0.143

criterion (Rosenthal & Henderson, 2003).

1279

1280

Figure 1 – figure supplement 4. Focused classification and interpretation of the MAT1 RING domain density. (**A**) The entire pool of pre-classified TFIIH

particles from the VPP datasets was subjected to alignment-free focused 3D
classification using a mask that covers the MAT1 RING domain and helical
bundle. Signal subtraction was not required due to the peripheral location of the
region in the complex and the orientation distribution of the particle images.
After one round of additional global classification (to select for overall particle
quality and conformation), the class showing the most highly occupied density
for the MAT1 RING domain was refined to 4.1 Å resolution (using beam tilt
values imported from the best-resolution refinement, Figure 1 – figure
supplement 1). (**B**) 3D classes differed not only in occupancy (i.e. classes 2 and 3
showed low occupancy and fragmented density for the MAT1 RING domain) but
also in the orientation of the MAT1 RING domain, which is flexibly attached to
the complex. Classes 1 and 4 are shown in different colors individually and
superposed to highlight the different orientations of the MAT1 RING domain. (**C**)
The FSC curve estimates a resolution of 4.1 Å for the refined map of TFIIH with
high occupancy of the MAT1 RING domain. (**D**) Best fit of the structure of the
MAT1 RING domain (Gervais et al., 2001) into the cryo-EM map, obtained using
the FITMAP command in UCSF CHIMERA (Pettersen et al., 2004) with 100
independent initial placements before local optimization at 5 Å resolution. This
fit shows the best correlation between model and map and additionally satisfies
geometrical constraints imposed by the continuity of the peptide chain between
the RING domain and the MAT1 helical bundle.
1305

Figure 1 – figure supplement 5. Multibody refinement. (**A**) Schematic for multibody refinement (Nakane et al., 2018) of the classified VPP dataset (indicated in Figure 1 – figure supplement 1). Three bodies were used and subjected to multibody refinement using the corresponding masks. (**B**) Example of the improvement of the cryo-EM density for the XPD N-terminus after multibody refinement (top) relative to the global refinement (bottom). (**C**) FSC curves for the refined bodies, indicating resolutions of 3.6 Å according to the FSC = 0.143 criterion (Rosenthal & Henderson, 2003).

1316

Figure 1 – figure supplement 6. Detailed architecture of the TFIIH core

complex. (**A-D**) The refined coordinate model is shown in the 3.7 Å-resolution

cryo-EM map. Large side chains are indicated. (**E-H**) Mapping of CX-MS data (Luo

et al., 2015) onto the atomic coordinate model. Crosslinked residues are shown

as spheres, crosslinks as black lines (crosslinks > 30 Å apart as yellow lines). (**E**)

Crosslinks between the p44 NTE and XPB, p52, and p34 confirm the tracing of

the p44 NTE towards the XPB NTD. (**F**) Intramolecular crosslinks in XPB validate

the tracing of the XPB NTD. (**G**) Crosslinks between p62 and XPD. Crosslinks

between p62 and p44 are not shown (see Methods and Supplementary Data

Table 1). (**H**) Crosslinks between MAT1 and XPB and XPD. One strong outlier

connecting XPB K44 to the MAT1 helical bundle (>100 Å distance) is a likely false

positive and is not shown. (**I**) Location of site-specific crosslinks between RAD3,

the yeast XPD homolog (RAD3 residues carrying the crosslinker mapped onto

the XPD structure and shown as purple spheres) and TFB3, the yeast MAT1

homolog (Warfield et al., 2016). (**J**) Interlocking zinc-binding domains of p44 and

p34 mediate interactions between these proteins in the hinge region. The p62 C-

terminal 3-helix bundle binds to the p44 eZnF domain. (**K**) Residues 378-388 of

p62 are sandwiched between the zinc-binding domains of p34 and p44. (**L**) View

of the p62-p52-p34 interaction region.

1336

1337

1338

Figure 2 – figure supplement 1. Analysis of the structure of p62. (**A**) p62

architecture and sequence assignment. Fitting of domains (see **B**), secondary

structure prediction for α-helices (Kelley et al., 2015), helical segments of the

cryo-EM map, and density for large side chains were used for initial assignment

of the trace of p62. The final model shows good agreement of observed helical

segments (residue numbers indicated) and secondary structure prediction for α-

helices (residue numbers given in brackets). (**B**) Fitting of p62 BSD1 (PDB ID
2DII; blue) into the TFIIH cryo-EM map. Due to flexibility, the density in this
region is weak and was low-pass filtered to 6 Å for interpretation and fitting. In
agreement with previous results (Schilbach et al., 2017), this domain interacts
with the p62 C-terminal three-helix bundle (purple), which in turn sits on the
eZnF domain of p44 (Figure 1 – figure supplement 6J), thereby explaining the
importance of the latter region for functional TFIIH (Tremeau-Bravard et al.,
2001). (**C**) Extended segments of the p62 chain were interpreted using the low-
pass filtered map as described in (B) to maintain continuity of the cryo-EM
density. (**D**) The p62 helix 295-317 is sandwiched between the BSD2 region of
p62 and XPD RecA2, thereby bridging these two elements of TFIIH. (**E**) XPD-
bound DNA was modeled by domain-wise superposition of the structures of a
XPD-DNA complex (Constantinescu-Aruxandei et al., 2016) and the structure of
DNA-bound Hel308 (Büttner et al., 2007) onto TFIIH. The presence of the p62
anchor near XPD RecA2 leads to steric hindrance with the DNA that would
prevent binding. (**F**) Residues 346-361 of p62 block access to the XPD
nucleotide-binding pocket. (**G-J**) Analysis of p62 (teal) conformational dynamics.
(**G**) Particle images were sorted for p62 elements near XPD using signal-
subtracted focused classification (mask shown in yellow; see Figure 1 – figure
supplement 3). (**H**) Refined class with p62 density near XPD. (**I**) Refined class
without p62 density near XPD. (**J**) Difference density (red) between the densities
shown in H, I. Both classes show intact particles, even in the absence of the p62
segment inserted into the XPD substrate binding cavity.

1368

1369

1370

Figure 3 – figure supplement 1. Domain organization of XPB and sequence alignment of the N-terminal regions of XPB or XPB-like enzymes from eukaryotes, archaea, and bacteria. (**A**) Domain organization of human XPB, color-coded as in Figure 2C. (**B**) Sequence alignments. Sequences used (species and UNIPROT or GENBANK accession code indicated): Human (*Homo sapiens*, P19447), mouse (*Mus musculus*, P49135), fruit fly (*Drosophila melanogaster*, Q02870), Choanoflagellate (*Monosiga brevicollis*, A9V0A0), fungus

(*Saccharomyces cerevisiae*, Q00578), ciliate (*Paramecium tetraurelia*,

XP_001448651.1), plant (*Arabidopsis thaliana*, Q38861), archaea (*Ferroplasma*,

EQB73227.1, *Thermoplasmatales*, EQB71900.1), bacteria (*Mycobacterium leprae*,

Q9CBE0) (Balasingham et al., 2012; Poterszman et al., 1997). Some archaeal

XPBs, including the *Archaeoglobus fulgidus* homolog that served for structure

determination (Fan et al., 2006), contain only the DRD, lacking the NTD/NTE,

and were not included. Residues are numbered according to the human

sequence. Identical residues are printed white on red background, similar

residues in red on white background. Secondary structure elements, residues

mutated in human disease (F99S, T119S), and the approx. 70-residue insertion

in the DRD are indicated. Figure generated using the ESPript web server (Robert

& Gouet, 2014).

1390

1391

1392

1393

Figure 3 – figure supplement 2. Structure and interactions of the XPB DRD

and NTD. (**A**) Interface between the XPB NTD and the p52-clutch domains.

Contacts include several large hydrophobic residues, as well as a salt bridge

between p52 R314 and XPB E115 (shown as sticks). Mutations of residues

equivalent to R314 and E310 in *D. melanogaster* p52 lead to disease-like

phenotypes (Fregoso et al., 2007). (**B**, **C**) Electrostatic surface potential of the

interacting surfaces of the XPB NTD (**B**) and the p52 clutch domain (**C**) show

charge complementarity (strongest complementary charge peak indicated by a

circle). (**D**) Front and back views of the domain organization of p52. Residues 1-

130 (red) and 131-304 (orange) are not in contact with the XPB NTD. Residues

304-400 of p52 form the p52 clutch. Protein subunits or domains interacting

with p52 are shown in grey. (**E**) Same as **D**, but p52 is colored using a gradient

from blue (N-terminus) to red (C-terminus). (**F**) Sequence alignment of human

and yeast p52 clutch and XPB NT domains highlights several highly conserved

residues, including XPB T119, which is affected by a human disease mutation.

Generated using the ESPript web server (Robert & Gouet, 2014). (**G**) Location of

XPB T119, affected by the TTD mutation T119P, at a junction between a β-strand

and an α-helix near the interface between XPB NTD, NTE, and RecA1 domains.

(**H**) Superposition of the XPB NTD and the p52 clutch shows that a threonine at

the position of XPB T119 also occurs in the p52 clutch. (**I**) The XP/CS mutation

F99S affects a XPB NTD residue (orange) situated in a hydrophobic pocket

formed by aliphatic residues (cyan) that are conserved across yeast and human

XPB and p52 (**F**).

1417

1418

Datasets 1 and 2
(after initial 3D
classification, Fig. S1)

3D refinement

XPD RecA1, ARCH, FeS
MAT1 RING, helical bundle

XPD RecA2
p44 VWFA domain
p62 BSD2

p34 (full protein)
p44 zinc binding domains
p62 3-helix bundle, BSD1

Mask
creation

p52 winged-helix domains

XPB NTD, DRD, RecA1
p52 clutch

XPB RecA2
p52 C-terminal domain
p8

Multibody refinement
with 6 masks

Principal component
analysis

First and last frame of
eigenvector movies

Difference densities:
first frame minus last
last frame minus first

Component 1

Component 2

Component 3

Component 4

Component 5

Component 6

Component 7

Component 8

Component 9

Component 10

Component 11

Component 12

Component 13

Component 14

Component 15

Component 16

Figure 4 – figure supplement 1. Analysis of conformational variance of TFIIH. VPP datasets 1 and 2 were refined and subjected to multibody refinement using 6 masks followed by principal component analysis. The first and last

1419
1420
1421
1422

1423 frames of the resulting eigenvector movies of the first 16 principal components

1424 are shown (in light blue and light yellow), along with difference densities

1425 obtained by subtracting the last volume from the first (green) and vice versa

1426 (purple). For the boxed components, the particles with the highest and lowest

1427 eigenvalues were extracted and refined (see Figure 4A).

1428

1429

1430
Figure 4 – figure supplement 2. Conformational dynamics of TFIIH and
comparison with the PIC-bound TFIIH. (**A**) Rendering of the yeast TFIIH-
containing Pol II-PIC (Schilbach et al., 2017). TFIIH subunits are colored and
labeled with their yeast names, with human names in brackets. (**B**)
Superposition of TFIIH from the yeast PIC (as shown in A, light colors) (Schilbach
et al., 2017) and the complete structure of the human TFIIH core complex (this
work; superposed on p44, bright colors) as single bodies. The top lobe of TFIIH
(MAT1 helical bundle, XPB, p44, p34, p52 winged-helix domains; p62 not shown
for clarity) aligns well, while the XPB-p52 (clutch/C-terminal domain)-p8
module is in a different position for the free and PIC-bound TFIIH (indicated). (**C**)
Same as **B**, but with the human XPB-p52 (clutch/C-terminal domain)-p8 module
superposed on the yeast structure as a rigid body. (**D**) View of DNA-bound Ssl2
(XPB) and neighboring subunits in the yeast Pol II-PIC (Schilbach et al., 2017).
(**E**) View of human XPB, superposed on the yeast structure as in **B**. XPB could not
bind DNA in this conformation, and the Tfa1 (TFIIE α-subunit) bridge helix, as
observed in the yeast Pol II-PIC, clashes with XPB. (**F**) Panels **D** and **E** combined,
visualizing the conformational change of TFIIH upon entry into the Pol II-PIC.
1448
1449

1450

Figure 5 – figure supplement 1. Structure of XPD. (**A**) Close-up view of the XPD N- and C-terminal segments near the nucleotide binding pocket of the enzyme. Y12 and Y18 may stabilize nucleotides when bound (not present in our structure). (**B**) View of the nucleotide binding site of XPD with superposed ADP from PDB ID 4BGD (Nguyen et al., 2013). Residues mutated in human disease are shown as sticks and colored (purple: XP, orange: XP/CS, dark green, TTD). These include Y18, which we propose could form a stacking interaction with the base of a bound nucleotide, as well as residues in conserved helicase motifs (see **C**). (**C**) Same view as **B**, but with conserved helicase motifs color-coded. Motifs Q, I, II, and IV are involved in nucleotide binding and hydrolysis. (**D**) View of the back side of XPD, with the N- and C-terminal segments indicated. The C-terminal segment interacts with the linker between XPD RecA1 and RecA2 (residues 476-

486). CX-MS crosslinks support the assignment of the C-terminal segment. The K751Q polymorphism has been associated with cancer therapy outcomes (Mlak et al., 2018; Peters et al., 2014), though functional differences between the variants has been debated (Clarkson & Wood, 2005). (**E**, **F**) Electrostatic surface (E) and cartoon (F) representations of the XPD structure, highlighting the pore between the XPD ARCH an FeS domains through which DNA (superposed from PDB IDs 5H8W, 2P6R (Büttner et al., 2007; Constantinescu-Aruxandei et al., 2016)) passes while being translocated by XPD. (**G**, **H**) Comparison of the structures of the human and archaeal FeS domains (**F**, *Thermoplasma*, PDB ID 4A15 (Kuper et al., 2012), **G**, *Sulfolobus*, PDB ID 3CRV (Fan et al., 2008)). There are differences at the secondary structure level, and functionally important residues are only partially conserved. (**I**) Sequence alignment of FeS domains from human, *Sulfolobus*, and *Thermoplasma* XPD. With the exception of the conserved cysteine residues that bind the FeS-cluster, sequence conservation in this domain is relatively low. Residues discussed in the text are indicated. (**J**-**M**) Charge complementary of regions of negative and positive electrostatic potential near the MAT1 3-helix bundle and the XPD ARCH domain interface (circled).

1482

Figure 5 – figure supplement 2. XPD-p44 interaction. (**A**) Top: XPD-p44 (XPD

green, p44 red) interaction surfaces are shown in darker shade (interacting

residues in p44 β4-α5 loop are shown in teal). Bottom: Charge complementarity

of the interface (two major complementary regions numbered 1 and 2). (**B**)

Detailed view of the β4-α5 loop (teal) with mutations discussed in the text. XPD-

p44 interacting regions (defined as residues within < 4 Å of neighboring protein)

are colored in dark green (XPD) and dark red (p44). Residues discussed in the

text are shown as sticks; those with mutation data (natural variants or

experimental constructs) are colored yellow on XPD, teal on p44. (**C**) Detailed

view of the interactions formed by XPD R722, which is mutated in TTD. (**D**) View

of XPD residues R616, D673, and G675 in the context of p62 and p44. (**E**) XPD-

p44-p62 interaction region. p62 segments interacting with both XPD and p44 are

colored dark blue and yellow.

1496

1497

1498

1499

**Figure 6 – figure supplement 1. Mapping of XPD disease mutations on the
TFIIH structure.** (**A**) Several residues affected by XP mutations localize near the
DNA binding surfaces of XPD. This figure uses the same color scheme as Figure 6
(XP mutations purple; TTD yellow; CS-XP orange; helicase elements for DNA
binding in blue, nucleotide binding and hydrolysis in green, coupling of
nucleotide hydrolysis to DNA translocation in brown; DNA superposed from
(Büttner et al., 2007)). (**B**) Mapping of mutations near the catalytic site of XPD.
(**C**) Mutations near the interface with p44. Note the predominance of TTD
mutations in this peripheral region of XPD. Also see Figure 5E, Figure 5 – figure
supplement 2, and text for details. (**D**) XPD R592, affected by a TTD mutation is
in a key location where p44, p62, and XPD interact. (**E**) Location of C259, affected
by the C259Y mutation, which likely causes disruption of ARCH domain integrity
and leads to defects in CAK subcomplex recruitment. (**F**) Mapping of yeast Rad3
rem mutations (green) onto our XPD structure. E235 and A236 are part of the
conserved DEAH sequence motif (helicase motif II), as is the immediately
adjacent XP mutation D234. H659 localizes near DNA (according to the
superposed model) and may be involved in ordering R601, which is in direct
contact with the modeled DNA and affected by an XP mutation.

The complete structure of the human TFIIH core complex

Basil J. Greber, Daniel B. Toso, Jie Fang, Eva Nogales[*]

[*] Correspondence should be addressed to E.N. (ENogales@lbl.gov).

This PDF files includes:

Supplementary Tables 1-3
Supplementary Notes 1-2

Other Supplementary Information for this manuscript includes the following:

Supplementary Data Table S1

1549

1550 **Supplementary Table 1. Data collection statistics.** All datasets were acquired

1551 on Gatan K2 Summit direct electron detectors mounted in 300 kV-electron

1552 microscopes with three-condenser type electron optics. Abbreviations: TEM,

1553 transmission electron microscope; VPP, volta phase plate; Σ, sum.

Data set #	TEM	VPP	Pixel size (Å)	Total dose (e⁻/Å²)	Frames	K2 acquisition mode	Total movies	Movies retained	Particles after initial 3D sorting	Particles in final map	Data reported
1	Titan KRIOS	Yes	1.15	50	33	Super-resolution	3,270	2,047	154,298	27,165	This study
2	Titan KRIOS	Yes	1.15	50	33	Super-resolution	3,190	1,804	134,654	23,034	This study
3	Titan KRIOS	Yes	1.15	50	33	Super-resolution	4,804	3,269	191,830	18,365	This study
4	Titan KRIOS	Yes	1.15	50	33	Super-resolution	3,283	2,709	107,390	32,008	This study
5	Titan KRIOS	Yes	1.15	50	33	Super-resolution	4,958	2,857	85,582	22,717	This study
6	Titan KRIOS	Yes	1.15	50	50	Super-resolution	1,932	1,631	113,001	15,370	This study
7	Titan KRIOS	No	1.32	40	30	Super-resolution	2,719	476	12,788	-	This study
8	Titan	No	1.32	40	30	Counting	1,734	215	3,942	-	Greber et al. 2017
9	Titan	No	1.32	40	30	Counting	2,694	262	4,617	-	Greber et al. 2017
10	Titan	No	1.32	40	30	Counting	2,593	653	19,642	-	Greber et al. 2017
Σ							31,177	15,923	827,744	138,659	

1554

1555

Supplementary Table 2. Components of the TFIIH structure.

Protein	Chain ID	Size* (aa)	Modeled residues	Sequence accession code	Alternative names	Comments
XPB	A	782	34-204, 247-730	P19447	ERCC3	-
XPD	B	760	1-760	P18074	ERCC2	4Fe4S-cluster: SF4, residue 1000.
p62	C	548	107-173, 183-323, 345-547	P32780	GTF2H1	BSD1 domain docked and refined with reference restraints; residues 233-295 built as poly-alanines and deposited as UNK.
p52	D	462	7-458	Q92759	GTF2H4	
p44	E	395	15-267, 282-387	Q13888	GTF2H2	Zn^{2+}, residues 401, 402, 403
p34	F	308	8-72, 95-292	Q13889	GTF2H3	Zn^{2+}, residues 401
p8	G	71	2-67	Q6ZYL4	GTF2H5, TTDA	-
CDK7	-	346	-	P50613	MO15	Not visualized.
Cyclin H	-	323	-	P51946	CCNH	Not visualized.
MAT1	H	309	1-210	P51948	MNAT1	N-terminal RING domain docked and refined with tight reference restraints. Zn^{2+}, residues 400, 401

 * According to Uniprot (http://www.uniprot.org).

 † Unassigned sequences were initially modeled as poly-alanine and deposited as UNK.

1561 **Supplementary Table 3. Data collection, map and model refinement, model**

1562 **validation.**

Data collection	Datasets 1-6
Microscope	Titan KRIOS
Stage type	Autoloader
Voltage (kV)	300
Detector	K2 Summit
Energy filter	GIF Quantum
Other	Volta phase plate
Pixel size (Å)	1.15
Defocus range (μm)	0.3-1.2
Electron dose (e^-/A^2)	50
Reconstruction	**EMD-XXXX**
Software	RELION 3
Particles	138,659
Box size (pixels)	256 x 256 x 256
Accuracy rotations (°)	1.4
Accuracy translations (pixels)	0.8
Map resolution (Å)	3.7
Map sharpening b-factor ($Å^2$)	-142
Coordinate refinement	
Software	PHENIX
Algorithm	REAL SPACE REFINE
Box size (Å)	294.4 x 294.4 x 294.4
Resolution cutoff (Å)	3.7
$FSC_{model-vs-map}$=0.5 (Å)	3.9
Model	**PDB-XXXX**
Number of residues	3203
Protein	3196
Ligand (Fe_4S_4, Zn^{2+})	7
B-factors overall	74.5
Protein	74.5
Ligand (Fe_4S_4)	63.4
Ligand (Zn^{2+})	133.8
R.m.s. deviations	
Bond lengths (Å)	0.009
Bond angles (°)	1.190
Validation	
Molprobity score (percentile)	2.0 (99th)
Molprobity clashscore (percentile)	6.7 (97th)
Rotamer outliers (%)	0.3
C_β deviations (%)	0
Ramachandran plot	
Favored (%)	87.8
Allowed (%)	11.9
Outliers (%)	0.3

1563

1564

1565

1566

Supplementary Discussion

Supplementary Note 1: The XPD-p44 interface. XPD serves a dual role within TFIIH: Its helicase activity is essential for TFIIH function in NER (Kuper et al., 2014; Winkler et al., 2000), and it serves as an architectural element in transcription, enabling the proper placement of the CAK subcomplex within the Pol II PIC (Abdulrahman et al., 2013; Drapkin et al., 1996; Dubaele et al., 2003; Greber et al., 2017; Guzder et al., 1994; He et al., 2016; Murakami et al., 2015; Robinson et al., 2016; Rossignol et al., 1997; Schilbach et al., 2017; Tirode et al., 1999; Tsai et al., 2017). The XPD-p44 interaction is critical both for the enzymatic and architectural roles of XPD (Coin et al., 1998; Dubaele et al., 2003; Kim et al., 2015; Kuper et al., 2014). Therefore, disruption of the XPD-p44 interface can simultaneously cause NER defects and impair transcription initiation, which leads to a TTD phenotype in human patients (Dubaele et al., 2003). Our structure provides a detailed view of this interaction interface, which encompasses the p44 loops β1-α1, β4-α5, β5-α6, and β6-α7 connecting β-strands and α-helices in the p44 VWA domain (Figure 5E, Figure 5 – figure supplement 2). Mutations L174W, T175R, and D178A in the p44 β4-α5 loop have been shown to abrogate the p44-XPD interaction and to reduce the helicase activity of XPD, implicating this region in XPD activation (Kim et al., 2015; Seroz et al., 2000). Our structure suggests that the insertion of large residues (L174W, T175R) into the tightly packed p44-XPD interface (Figure 5 – figure supplement 2B) is likely to lead to steric clashes, while the D178A mutation would break a likely salt bridge with XPD R592 (Figure 5C). Strikingly, the R592P mutation in XPD, the side chain of which packs against the p44 β4-α5 loop in wild type XPD, causes TTD in human patients (Cleaver et al., 1999), underscoring the importance of this interaction.

In XPD, RecA2 loop residues 531-533, 563-565, 589-592, 612-616, and the helical segment 718-725 form the p44 interaction interface (Figure 5E). Using recombinant human and *Chaetomium thermophilum* proteins, it has been shown that the XPD R722W mutation impairs the interaction between p44 and XPD, thereby interfering with both the p44-dependent stimulation of the XPD ATPase

and helicase activities in NER, and architectural role of p44 in transcription
initiation (Kuper et al., 2014). Our structure shows that the R722W mutation
disrupts a salt bridge to p44 D75, and the bulky tryptophan side chain may
additionally cause steric clashes with neighboring p44 residues, explaining the
deleterious effect of the mutation (Figure 5E, Figure 5 – figure supplement 2C). It
has been proposed that the R616P, D673G, and G675R disease mutations, which
impair the p44-XPD interaction and the XPD helicase activity (Dubaele et al.,
2003), exert their deleterious effect by impairing the interaction between XPD
and p62 (Luo et al., 2015). However, our structure shows that G675 is buried in
the enzymatic core (Figure 5 – figure supplement 2D), where accommodation of
the large arginine side chain in G675R will disrupt the structure of the protein.
XPD D673 is located in a β-strand in the vicinity of the XPD-p62 interface (Figure
5 – figure supplement 2D); however, no direct interactions of this residue with
p62 are observed. XPD R616 is located at the p44-XPD interface, where its side
chain is packed against neighboring residues and may engage in a salt bridge
with p44 D204 (Figure 5 – figure supplement 2D). These interactions would be
disrupted in the R616P mutant. Altogether, these observations suggest that these
mutations act either via disruption of XPD structure or the XPD-p44 interface,
not via p62. Nevertheless, a contribution of p62 to the stabilization of the p44-
XPD interface cannot be excluded because several p62 elements are observed
near the interface (Figure 5 – figure supplement 2E). As discussed in the main
text, some of these interactions are dynamic (Figure 4 – figure supplement 2G-J)
and unlikely to be strictly required for the stability of the p44-XPD interface.
1623
Supplementary Note 2: Mapping of XPD mutations. In addition to our analysis
of the XPD-p44 interface and the mutations located in its vicinity, we were also
able to map numerous other human disease mutations onto our structure
(Figure 6). Our results are in good agreement with the hypotheses derived from
comparative modeling based on homologous bacterial DNA repair enzymes
(Bienstock et al., 2003), archaeal XPD homologs (Fan et al., 2008; Liu et al., 2008;
Wolski et al., 2008), and our previous lower-resolution TFIIH structure (Greber
et al., 2017). Several XP mutations map in the regions that interact with bound
DNA, as confirmed by superposing homologous DNA-bound helicase structures

(Figure 6 – figure supplement 1A), or in the conserved motifs near the ATPase

active site (Figure 6 – figure supplement 1B). These mutations impair XPD

helicase activity (Dubaele et al., 2003; Fan et al., 2008; Liu et al., 2008), causing

the NER defects that manifest in the XP disease symptoms. As explained above

(Supplementary Note 1), TTD mutations tend to map to the periphery of XPD,

including the XPD-p44 interface (Supplementary Note 1, Figure 5E, Figure 5 –

figure supplement 2, Figure 6 – figure supplement 1C, D), the ARCH domain

(Figure 6 – figure supplement 1E), or the FeS domain (Figures 5B, 6). The C259Y

mutation in the ARCH domain likely impairs the proper folding of this domain

due to steric hindrance introduced by the large tyrosine side chain, which in turn

impairs association of the CAK subcomplex through disruption of MAT1

association, as discussed previously (Greber et al., 2017; Liu et al., 2008; Wolski

et al., 2008). Consistent with its observed location in our structure (Figure 5B),

the R112H mutation in the FeS domain leads to destabilization of the 4FeS

cluster (Rudolf et al., 2006), which in turn leads to the disruption of folding of

this entire domain (Fan et al., 2008) and lower TFIIH levels overall (Botta et al.,

2002). Another TTD mutation that does not localize to the periphery of XPD is

the particularly interesting temperature-sensitive mutation R658C, for which

patients show exacerbated symptoms, including complete hair loss, when they

develop a fever (Vermeulen et al., 2001). This residue is located at the interface

between the XPD RecA1 and RecA2 domains, where it forms a salt bridge across

the domain interface (Figure 6B). This interface would be partially destabilized

by the mutation at normal temperature and more severely disrupted at elevated

temperature, leading to stronger disease symptoms.

1657

We also mapped onto our structure the location of the so-called *rem* mutations in

Rad3, the yeast homolog of XPD. These mutations increase the affinity of Rad3

for DNA, which leads to retention of Rad3/XPD on DNA lesions, incomplete NER

reactions, and subsequent DNA breakage upon arrival of a DNA replication fork

(Herrera-Moyano et al., 2014; Moriel-Carretero & Aguilera, 2010). We mapped

the *rem* mutations Rad3-E236G (XPD E235), Rad3-A237T (XPD A236), and

Rad3-H661Y (XPD-H659Y), as well as the human XP-CS mutation XPD-G675R,

which also exhibits elevated DNA affinity but reduced helicase activity in an

archaeal model system (Fan et al., 2008) and DNA breakage in patient cells (Theron et al., 2005), onto our structure (Figure 6 – figure supplement 1F). XPD E235 and A236 are part of the conserved helicase motif II (signature sequence DEAH, Figure 5 – figure supplement 1C, Figure 6 – figure supplement 1F) (Fairman-Williams et al., 2010), and as such critical for nucleotide hydrolysis. Notably, the neighboring XPD D234, also part of motif II (Figure 5 – figure supplement 1C, Figure 6 – figure supplement 1C, F), is a human XP mutation (Cleaver et al., 1999). XPD H659 is located nearby, and contacts R601, which locates to the immediate vicinity of the DNA based on superposition with structures of DNA-bound helicases (Büttner et al., 2007; Cheng & Wigley, 2018) (Figure 6 – figure supplement 1F) and is affected by an XP mutation (Cleaver et al., 1999). In agreement with previous proposals, the *rem* mutations are therefore localized near regions responsible for ATP hydrolysis and DNA binding, consistent with the idea that they lead to stalling of NER complexes at DNA lesions, which may ultimately lead to incomplete NER, DNA breaks, failure to re-start transcription, and the XP-CS phenotype (Moriel-Carretero et al., 2015).

In summary, our structure pinpoints the locations of XPD mutations that provide insight into human disease mechanisms, and in most cases resolves the density for the side chains of the affected residues, thereby providing a structural framework for the detailed analysis of the interactions of the affected residues and the effects of these mutations.

Analysis of CX-MS data for human TFIIH

Chemical crosslinking-mass spectrometry data taken from Luo et al., Mol. Cell 59 (5): 794-806 (2015)
and mapped onto the cryo-EM structure of human TFIIH.

Residues not built in in the cryo-EM structure of human TFIIH are printed red.
Crosslinking distances refer to Cα-Cα distance of crosslinked lysine residues.
The expected maxium crosslink distance for BS3-induced crosslinks is 30 Å.

Color coding for crosslinking distances:

Agreement	Borderline	Incompatible
(< 30 Å)	(30-35 Å)	(> 35 Å)

Intramolecular crosslinks in human TFIIH subunits

Protein	Residue 1	Residue 2	Distance (Å)	Comments
p34	100	56	18.0	
p34	149	56	14.0	
p34	97	56	23.2	
p34	74	56	-	
p34	152	56	14.6	
p34	125	56	35.2	
p34	125	74	-	
MAT1	127	137	16.8	
MAT1	167	180	19.6	
MAT1	176	202	11.9	
MAT1	55	70	16.0	
MAT1	137	55	27.3	
MAT1	70	76	10.0	
MAT1	70	77	10.7	
XPD	268	274	9.7	
XPD	128	603	20.5	
XPD	692	751	28.6	
XPD	223	228	9.3	
XPD	223	69	13.0	
XPD	223	445	14.0	
XPD	223	35	16.8	
XPD	133	181	18.4	
XPD	35	751	26.8	
XPD	603	82	17.5	
XPD	228	35	10.4	
XPD	682	689	10.1	
XPD	113	603	18.6	
XPD	113	128	9.3	
XPD	113	181	11.3	
XPD	507	634	19.5	
XPD	507	682	9.8	
CDK7	32	342	-	CKD7 is not modeled in our structure.
CDK7	44	52	-	
CDK7	328	42	-	
CDK7	14	42	-	
CDK7	14	32	-	
CDK7	14	44	-	
CDK7	330	343	-	
XPB	167	222	-	Crosslink of the DRD loop insertion
XPB	167	326	16.3	
XPB	569	715	12.4	
XPB	171	311	29.2	
XPB	171	217	-	Crosslinks of the DRD loop insertion
XPB	171	222	-	that is not visible in the cryo-EM map
XPB	44	778	-	
XPB	44	68	28.1	
XPB	322	326	6.2	
XPB	157	44	13.1	
XPB	549	577	12.0	
XPB	282	463	18.0	
XPB	282	476	21.5	
XPB	577	601	11.3	
XPB	142	222	-	Crosslinks of the DRD loop insertion
XPB	142	241	-	that is not visible in the cryo-EM map
XPB	142	68	12.1	
XPB	311	379	10.0	
XPB	311	326	23.2	
XPB	217	326	-	Crosslinks of the DRD loop insertion
XPB	222	326	-	that is not visible in the cryo-EM map
XPB	222	257	-	
p62	129	504	15.3	
p62	18	38	-	Crosslinks of the N-terminal PH-
p62	104	106	-	domain of p62 and other N-terminal
p62	19	38	-	segments, which are not modeled
p62	161	38	-	in our structure.
p62	102	106	-	
p62	102	109	-	
Cyclin H	253	260	-	Cyclin H is not modeled in our
Cyclin H	274	62	-	structure.
Cyclin H	293	295	-	
Cyclin H	189	72	-	
Cyclin H	62	72	-	
Cyclin H	292	295	-	
p44	116	126	13.3	
p44	116	306	20.1	
p44	116	157	15.6	
p44	23	40	18.2	
p44	126	23	55.5	
p44	126	306	16.9	
p44	126	133	10.4	
p44	126	268	28.6	
p44	126	297	20.4	
p52	76	81	10.1	
p52	452	459	-	
p52	452	456	6.7	
p52	378	53	31.8	
p52	127	53	24.6	
p8	2	31	24.5	
p8	6	31	17.0	

Intermolecular crosslinks in human TFIIH subunits

Protein 1	Residue 1	Protein 2	Residue 2	Distance (Å)	Comments
p34	149	p52	378	15.3	
p34	152	p52	378	10.2	
p34	97	p52	127	8.7	
XPB	379	XPD	682	29.7	
XPB	222	XPD	682	-	
XPB	326	XPD	682	18.4	
XPB	171	XPD	682	19.6	
XPB	171	XPD	507	19.1	
p62	378	p34	125	25.7	
p62	38	p34	125	-	
p62	38	XPB	549	-	
p62	353	p44	126	50.0	p62-p44 crosslinks were discarded
p62	38	p44	126	-	in modeling due to high incidence
p62	252	p44	31	73.1	of distance violations and excellent
p62	161	p44	116	22.3	agreement of p62 model with
p62	161	p44	126	34.0	p62-XPD crosslinks.
p62	350	XPD	82	10.2	
p62	350	XPD	113	17.0	
p62	365	XPD	96	16.9	
p62	353	XPD	603	19.6	
p62	353	XPD	113	20.8	
p62	353	XPD	82	9.4	
p62	275	XPD	603	13.6	
p62	293	XPD	603	23.0	
p62	293	XPD	113	32.1	
p62	296	XPD	603	25.7	
Cyclin H	239	MAT1	55	-	Cyclin H is not modeled.
MAT1	121	XPB	44	116.7	
MAT1	180	XPB	171	17.7	
MAT1	202	XPB	142	37.5	
MAT1	176	XPB	171	19.3	
MAT1	167	XPB	217	-	Crosslinks of the DRD loop insertion
MAT1	167	XPB	222	-	that is not visible in the cryo-EM map
MAT1	167	XPB	257	19.1	
MAT1	167	XPD	689	26.9	
MAT1	167	XPD	634	19.7	
p44	40	p34	149	21.0	
p44	126	p34	149	34.1	
p44	42	p34	149	22.2	
p44	229	p34	149	23.0	
p44	229	p34	152	26.7	
p44	23	XPB	142	7.7	
p44	31	XPB	142	8.7	
p44	23	p52	378	32.7	
p44	40	p52	378	23.2	
p44	268	p52	77	30.0	
p44	77	p52	378	20.8	
p52	438	XPB	526	33.9	
p52	438	XPB	706	20.4	
p52	378	XPB	526	51.6	
p52	378	XPB	241	-	
p52	378	XPB	257	63.0	
p52	140	XPD	29	92.4	Likely false positives given the reliable
p52	277	XPD	101	116.0	assignment of p52 and XPD.
p52	53	p8	31	59.3	
p52	452	p8	2	9.8	
p52	452	p8	6	12.5	
p52	77	p8	71	-	
p52	456	p8	2	11.8	
p52	456	p8	6	18.5	
CDK7	52	Cyclin H	293	-	CDK7 and Cyclin H are not modeled
CDK7	293	XPD	634	-	CDK7 is not modeled.
p8	71	XPB	767	-	Crosslinks of p8 with the XPB RecA2
p8	71	XPB	526	-	domain are compatible with the
p8	71	XPB	59	-	architecture of TFIIH overall.
p8	71	XPB	476	-	
p8	6	XPB	167	88.0	