High throughput genotyping of structural variations in a complex plant genome using an original Affymetrix® Axiom® array

Clément Mabire[2*], Jorge Duarte[1*], Aude Darracq[2], Ali Pirani[3], Hélène Rimbert[1,4], Delphine Madur[2], Valérie Combes[2], Clémentine Vitte[2], Sébastien Praud[1], Nathalie Rivière[1], Johann Joets[2], Jean-Philippe Pichon[1], Stéphane D. Nicolas[2]

*Two authors contributed equally to work

Authors Affiliation:

1 Biogemma - Centre de Recherche de Chappes, CS 90126, Chappes 63720, France

2 GQE – Le Moulon, INRA, Univ. Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

3 Thermo Fisher Scientific, 3450 Central Expressway, Santa Clara, CA, 95051, USA

4 Present address: GDEC, INRA, Université Clermont Auvergne, 63000 Clermont-Ferrand, France

Corresponding authors: stephane.nicolas@inra.fr

Abstract

Background

Insertions/deletions (InDels) and more specifically presence/absence variations (PAVs) are pervasive in several species and have strong functional and phenotypic effect by removing or drastically modifying genes. Genotyping of such variants on large panels remains poorly addressed, while necessary for approaches such as association mapping or genomic selection.

Results

We have developed, as a proof of concept, a new high-throughput and affordable approach to genotype InDels. We first identified 141,000 InDels by aligning reads from the B73 line against the genome of three temperate maize inbred lines (F2, PH207, and C103) and reciprocally. Next, we designed an Affymetrix® Axiom® array to target these InDels, with a combination of probes selected at breakpoint sites (13%) or within the InDel sequence, either at polymorphic (25%) or non-polymorphic sites (63%) sites. The final array design is composed of 662,772 probes and targets 105,927 InDels, including PAVs ranging from 35bp to 129kbp. After Affymetrix® quality control, we successfully genotyped 86,648 polymorphic InDels (82% of all InDels interrogated by the array) on 445 maize DNA samples with 422,369 probes. Genotyping InDels using this approach produced a highly reliable dataset, with low genotyping error (~3%), high call rate (~98%), and high reproducibility (>95%). This reliability can be further increased by combining genotyping of several probes calling the same InDels (<0.1% error rate and >99.9% of call rate for 5 probes). This "proof of concept" tool was used to estimate the kinship matrix between 362 maize lines with 57,824 polymorphic InDels. This InDels kinship matrix was highly correlated with kinship estimated using SNPs from Illumina 50K SNP arrays.

Conclusions

We efficiently genotyped thousands of small to large InDels on a sizeable number of individuals using a new Affymetrix® Axiom® array. This powerful approach opens the way to studying the contribution of InDels to trait variation and heterosis in maize. The approach is easily extendable to other species and should contribute to decipher the biological impact of InDels at a larger scale.

Keywords

Present Absent Variation, Copy Number Variation, Structural Variation, genotyping, array, Zea mays, Genome assembly, Breakpoint, Chromosomal rearrangements

Background

In the past decade, there has been growing evidence that structural variations (SVs) are pervasive within plant genomes [1–9]. Insertion/deletions (InDels) are one class of SVs of particular interest, since they lead to the presence or absence of, sometimes large, genomic regions at a given locus, among individuals from the same species. The content of these InDels can be present elsewhere in the genome, but they can also be completely absent from the genome, in which case they are referred to as presence/absence variants (PAVs). Some InDels carry entire genes or affect gene regulatory elements and are thus likely to have a functional and phenotypic impact [10–12, 7, 13]. Hundreds to thousands of SVs, including PAVs and copy number variations (CNVs), have been discovered in several plant species, including wheat [14], rice [15], *Arabidopsis thaliana* [13], potato [16] , pigeon peas [17], and sorghum [18]. These results support the idea that one single reference genome cannot properly represent the complete gene set of a given species. There has been an increasing interest for building new individual genomes in complement to the reference genome, in order to better describe the genetic diversity within a plant species [3, 19–25] .

In maize, BAC sequence comparison first revealed that gene and transposable element content greatly vary between inbred lines [26, 27]. Whole genome sequencing of the B73 inbred line then provided the opportunity to explore the extent of SVs across the entire maize genome [28] by designing Comparative Genomic Hybridization (CGH) technology [29] . Several CGH studies found multiple CNVs between the B73 reference genome and other maize inbred lines or teosintes [2, 8, 9]. These studies demonstrated the large extent of SVs among maize inbred lines, including presence/absence variations of low copy sequences, such as genes. This was well illustrated by the discovery of a large 2 Mbp presence/absence region between Mo17 and B73 carrying several genes [2, 9, 20, 21]. However, CGH array technology shows several major drawbacks since (i) it does not allow the discovery of sequences that are not present in the reference genome used for designing probes of the arrays, (ii) it has a limited resolution which does not allow detection of InDels smaller than 1kb, and (iii) it is costly and labor-intensive, and therefore not adapted for genotyping several hundreds of individuals.

Methods based on SNP array experiments have been developed to detect CNVs and were shown to be more affordable and with higher throughput than CGH arrays [32–35] . Didion et al. (2012) identified atypical patterns of reduced hybridization intensities that were highly reproducible, so called "off-target variants" (OTVs) [36]. OTV patterns could originate either from the absence of the sequence due to a PAV polymorphism, or to a single nucleotide polymorphism within the probe sequence, thus preventing the correct hybridization of the DNA sample. For instance, 45,974 OTVs were discovered in a maize population using the 600K Affymetrix® Axiom® SNP array [37]. While these approaches proved to be useful, there is a strong risk of false positive detection of PAVs using OTV patterns, mainly because these arrays were not designed to target PAVs. In order to reduce this risk of false positive detection of PAVs and more largely CNVs, several methods based either on segmentation or Hidden Markov Chain have been developed to use variation of fluorescent intensity signal of contiguous probes along the genome [38–43] These kind of approaches have been used on 600K Affymetrix® Axiom® SNP array to detect several hundreds of CNVs and to explore the contribution of CNV to phenotypic variation [44].

With the emergence of massive parallel sequencing, new methods have been developed to detect structural variations based on the alignment of resequencing reads onto a high quality reference genome sequence. Among these, three have been mainly used [45] : (i) the "read-depth" (RD) method, which can only detect copy number variations; (ii) the "read-pair" (RP) method, which can detect deletions as well as small insertions (up to the size of the library insert); and (iii) the "split-read" (SR) method which can also detect deletions and small insertions (up to the size of a read). Chia et al. (2012) used the RD approach to identify CNVs among 104 maize lines and performed association studies for several traits [10]. However, the RD method does not allow the identification of novel sequences and is error prone, especially regarding the size of the discovered CNVs which greatly depends on the size of the sliding window used. The RP method has been implemented in many computational tools like BreakDancer [46] and has been widely used. Although it has proven to be highly efficient to detect deletions [47–49], this approach suffers from two limitations: it does not allow precise detection of breakpoints, and the size of the insertions which can be detected is directly limited by the library insert size. The SR method, which was first implemented in PInDel [50], has the advantage of defining breakpoints at a single-base resolution, but again the size of the detectable inserted sequence is limited.

The "assembly" (AS) method is able to detect all types of SVs of any size, but is also the most cost and computation-intensive. It is the only method able to detect large insertions with precise breakpoint definition. However, the assembly of large and complex genomes such as maize remains very expensive and computationally intensive, despite recent progress in this area [20, 21, 31]. There has been in the past some attempts to reduce this complexity by reducing the number of sequences to assemble. For instance, Lai et al., (2010) identified 104 deletions and 570 insertions among 6 maize inbred lines by assembling genomic regions from reads that did not map on the B73 reference genome [51]. The sequences assembled by this approach were enriched in erroneous reads or reads coming from external contamination, and they were too short to be anchored to the reference genome B73. Hirsch et al. (2014) identified several putatively expressed genes that were not present within B73 reference genome by assembling and comparing the transcriptome of hundreds of inbred lines [12]. This new approach was limited to the transcribed part of the genome and suffered from a high level of false positives. More recently, Lu et al., (2015) used genotyping by sequencing approaches on 14,129 inbred lines to identify 1.1 million short and unique sequences (GBS tags) that (i) did not align on the B73 reference genome, or were aligned but outside of a 10Mbp windows around their mapped position; or (ii) were mapped at the same location by joint linkage mapping in NAM populations using co-segregation with a SNP and logistic regression between the InDel and the SNP in an association panel [13]. The main drawback of this approach is the high percentage of missing data due to the low depth of sequencing, which requires imputation before being able to perform genetic analysis. Recent whole genome sequence assemblies of PH207 [31], and F2 [20] have allowed the identification of thousands of large InDel and PAV sequences. For instance, 2,500 genes were found either present or absent in PH207 and B73 genomes and 10,735 PAV sequences larger than 1kb were discovered between F2 and B73, including 417 novel genes in F2. These discovery approaches have been limited to a few individuals due to sequencing costs and computational challenges, so they have not been adapted for characterization of SVs on large maize panels. Darracq et al. (2018) developed an interesting approach for the genotyping of PAVs from mapping of low depth (5-20X) resequencing datasets [20]. This method is based on the comparison of reads aligning

to the region found in F2 and in the line of interest. While this method is potentially adapted to genotype
PAVs on any set of line with low resequencing data, it has been so far used for PAV genotyping on a low
(<30) number of maize lines. Moreover, it is restricted to the analysis of PAVs, and is not adapted for
genotyping other types of SVs. To avoid this ascertainment bias due to use of a single reference genome
to genotype SV, other studies proposed to call SV by aligning reads on a pan-genome representing the
combination of several genomes [14, 22, 52]. However, these approaches remained computationally
challenging on a sizable set of individuals, time demanding, and costly for large and complex genomes,
since it requires high-depth sequencing [52]. To our knowledge, no high-throughput genotyping approach
has been developed for genotyping large numbers of InDels, including PAVs, on a large set of individuals.
We have developed, as proof of concept, a new high-throughput and affordable array that is able to
genotype simultaneously large insertions and deletions, with highly variable size and contents that are
previously discovered by different sequencing methods. In this study, we present this approach which is
both (i) comprehensive, as it includes the discovery and localization of deletions as well as insertions
regarding the B73 reference genome at the base pair level and (ii) high-throughput, as it allows genotyping
of thousands of InDels on hundreds of individuals. Our strategy takes advantage of next generation
sequencing (NGS) technologies and recent advances in assembly of complex genomes. It also benefits
from the high efficiency of SNP arrays like the high-throughput Affymetrix® Axiom® technology. In this
paper, we detail how we discovered thousands of small to large InDels, including PAVs, from three maize
inbred lines (F2, PH207 and C103) as compared to the B73 reference genome. We then describe how we
designed and selected 600,000 probes to create a new Maize Affymetrix® Axiom® array to genotype these
InDels. Finally, we describe how we successfully used this array to genotype an association panel of 362
maize inbred lines.

Results

InDel and PAV discovery

To design a comprehensive InDel genotyping array, we first discovered a set of InDels which would be representative of the maize temperate germplasm. We already had access to sequence data for the European flint line F2, and we benefited from a first set of 42,330 F2-specific sequences, larger than 150pb and totaling 16Mbp. This dataset was derived from the *de novo* assembly of an F2 paired-end that failed (at least for one read of the pair) to align onto the B73 AGPv2 sequence, and which were totally devoid of coverage by B73 reads ("Reference guided assembly" in Additional file 2: Figure S1, "no map" approach). We also took advantage of the work done by [20] to add another 10,044 F2-insertions (size >1 kb, total size of 88Mb), with less than 70% of their length covered by B73 reads discovered by a whole genome assembly approach (Additional file 2: Figure S1).

To complement these two datasets of F2/B73 deletions and insertions, we generated and assembled Illumina® paired-end and mate-pair sequences from two other key founders of temperate maize breeding programs: PH207 and C103. We then used this F2, PH207, and C103 sequence data to detect all InDels, including PAVs, at base-pair resolution, between these three lines and B73. As opposed to the "reference guided assembly approach", the "whole genome assembly" methodology allowed us to access both to their sequences and their breakpoints, permitting the genotyping of such InDels in several individuals (more details in Methods). We did not use the "no map" approach for InDel discovery on PH207 and C103, because this approach did not give access to breakpoint resolution, did not allow the discovery of InDels without knowledge of the specific sequence, and was almost redundant with the assembly approach.

We first aligned F2, PH207, and C103 sequences against the B73 reference genome sequence in order to detect deletions. Here, the term "deletion" does not reflect any underlying biological process of DNA excision but refers to a sequence of at least 100bp present in the B73 genome at one locus and absent in another line at the same locus. Deletions were detected for the three lines simultaneously using the "genotyping" option of PInDel [50], generating a set of 26,368 non-redundant deletions with precise identification of their breakpoints (Additional file 2: Figure S2A). The number of deletions found for each line was similar, respectively 12,165, 11,922, and 13,432 for F2, PH207, and C103. 67% of the deletions found were unique to one line, 24% were shared by two lines, and 9% by three lines (Additional file 2: Figure S2A). These results confirm the good complementarity of the lines chosen to discover InDels. The high proportion of unique deletions among 4 lines may also reflect that numerous InDels remain to be discovered in temperate maize germplasm.

Next, we generated a draft genome assembly for each of these lines, which was used as a template for alignment of B73 reads to detect insertions relative to the B73 reference genome (Additional file 1: Table S1). As for deletions, here the term "insertion" does not reflect any underlying biological process of DNA integration, but defines a sequence larger than 100bp that is present in one line at a given locus, and absent from B73 at the same locus. These three draft assemblies cover less than one third of

189   the expected maize genome size but include a large portion of low copy sequences, including genes, as
190   shown by BUSCO results (Table 1).

Table 1: F2, PH207, and C103 de novo assembly metrics.

Maize line	Number of scaffolds	Min size	Max size	Average size	N50	Total (Mb)	% of Ns	Complete BUSCOs (C)	Fragmented BUSCOs (F)	Missing BUSCOs (M)
F2	76,563	892	112,956	16,900	14,042	646.3	9.48%	89.3%	4.9%	5.8%
PH207	81,688	884	2,024,489	29,557	16,860	797.5	8.90%	91.8%	2.7%	5.5%
C103	84,990	886	120,582	19,305	16,146	793	8.21%	90.6%	4.2%	5.2%

Number of scaffold: The number of scaffold sequences assembled, Min Size: the length of the shortest scaffold, Max size: the length of the longest scaffold, Average Size: the average size of scaffolds, N50: N50 of the assembly, Total: the total number of bases included in the assembly, % of Ns: the percentage of Ns present in the assembly; BUSCO statistics included the percentage of complete (C), fragmented (F) and missing (M) BUSCO genes from a total of 1440 BUSCO genes

Detection of insertions was processed separately for each inbred line and generated 28,221 insertions for
F2, 27,904 insertions for C103, and 26,795 insertions for PH207, with their precise breakpoints (Additional
file 2: Figure S2B). The number of insertions is similar between lines, but significantly greater than the
observed deletions. Among these insertions, 26,691 cases could be uniquely anchored at base pair
resolution onto the B73 reference genome sequence (Additional file 2: Figure S2B). Again, a majority of
insertions were unique to one line (72%) confirming the complementarity of the material chosen (Figure
S2B).

Finally, the results from the different approaches were merged into a non-redundant set of
141,325 InDel sequences (see Methods), comprising 52,175 deletions and 89,150 insertions. These
regions were then used for the design of genotyping probes.

Design of the genotyping array

Genotyping strategy

Large InDels can be efficiently genotyped with a SNP array using a combination of two types of
probes: (i) "external" probes, which target breakpoints using the two flanking sequences of a given InDel
(BP probes), and (ii) "internal" probes, which target presence/absence regions (PARs) within the internal
sequence of InDels on polymorphic (OTV probes) or monomorphic sites (MONO probes). We define PARs
as small portions of DNA sequence of at least 35bp that were observed present or absent at the genome
level, when comparing two individuals. They are thus suitable for the design of presence/absence

genotyping probes. Ideally, each InDel should be called by two BP probes on either side and by multiple
internal probes, regularly distributed along the internal sequence of the InDel (Figure 1A). However, in
practice, this combination of different probes is not always possible. For instance, precise breakpoints
were not determined for all PAVs from our "no map" approach and [20], and PARs for internal probes
were not always found in our InDels.

Probe design

BP probes should behave like classical SNP probes where one allele corresponds to the presence
and the other to the absence of the InDel. They are useful to explore the conservation of the localization
of large insertion/deletion events across multiple individuals, even when no internal probe can be
designed due to the absence of PARs. Among the 141,325 selected variants, 86,406 InDels (22,420
deletions and 63,986 insertions as compared to the B73 reference genome sequence) had breakpoints
defined at base-pair resolution and were suitable for BP probe design. Four different breakpoint types
were identified according to the presence of micro-homology and/or shorter non homologous sequence
[53] in place of a complete deleted sequence (Additional file 2: Figure S3): (type I) 3,397 cases with sharp
breakpoints; (type II) 45,987 cases with a micro-homology sequence (8.6 bp on average and no more than
237 bp) which was present in one copy in the reference sequence and duplicated at both extremities of
the novel inserted sequence; (type III) 36,893 cases harboring insertion of a short non-homologous
fragment (42.2 bp on average and up to 892 bp) in place of a large deleted sequence; and (type IV) 156
cases with a combination of type II and type III breakpoints. Following Affymetrix® recommendations,
19,010 InDels with type II breakpoints having a micro-homology sequence longer than 5bp were excluded
from the design process. In the end, 67,396 InDels, representing 48% of all available InDel variants, were
submitted to the Affymetrix® design pipeline. Two probes, one on forward (FW) and one on reverse (REV)
strand, were designed for each breakpoint. These probes were classified as *not possible* (18%), *not*
recommended (33%), *neutral* (15%) and *recommended* (35%) by this automated pipeline (see Methods
for details), leaving 33,430 InDels (51%) that could be targeted by at least one *recommended* probe.

Internal probes, which should behave like "off-target" variants [36] where the hybridization of the
probe indicates presence of the InDel, and the absence of hybridization of the probe indicates absence of
the InDel, are useful to explore the genetic diversity within InDel sequences (Figure 1 D). They will also be
particularly interesting to target InDels for which no breakpoint could be identified (such as PAVs from
the "no map" approach).

For the design of OTV probes, we benefited from the availability of SNPs which had been
previously identified from the alignment of resequencing data from a core collection of 25 temperate
maize inbred lines against the B73-F2 maize pan-genome from [20]. As a consequence, OTV probes have
only been designed for deletions positioned on the B73 reference genome and F2 insertions coming from
[20]. Among these, the context sequences of 436,162 SNPs, corresponding to 21,390 InDels, were
extracted and submitted to the Affymetrix® design pipeline. Two probes, one on forward (FW) and one
on reverse (REV) strand, were designed for each SNP. A total of 872,324 OTV probes could be designed
and scored as *not possible* (0.05%), *not recommended* (71%), *neutral* (14%) and *recommended* (16%),
leaving 17,589 InDels (82%) which could be targeted by at least one *recommended* probe.

For the design of BP and OTV probes we could rely on Affymetrix® design pipeline to identify
probes localized in PARs and thus suitable for the Affymetrix® Axiom® technology. For the design of MONO
probes, we first had to identify such PARs within 141,325 InDels cumulating 133Mbp of sequence. We
used sequence masking methods to exclude repeats based on similarity to known maize repeats or on
occurrence of 17-mers found within the sequencing datasets we had for B73, F2, PH207, and C103 (see
Methods). By doing so, we identified 122,972 PARs, representing a cumulated size of 27Mbp,
corresponding to 20.3% of the initial size and allowing the possibility to design MONO probes for 79,987
InDels (56.5%). These PAR sequences were successfully used for the design of 25,735,797 MONO probes,
among which 59% were scored as *recommended* and allowed to target 62,875 InDels (79%).

With this combined approach, we designed a total of 26,715,361 probes targeting 117,756 InDels,
which represent a cumulated length of 250 Mbp including 27 Mbp of PARs (Table 2).

Table 2: Number of probes and targeted InDels before and after selection for array design and passing the Affymetrix® quality control according to different probes type. Percentages are indicated in brackets

	Before selection		On array		Called by Affymetrix® pipeline	
	Probes	**InDel[+]**	**Probes**	**InDel[+]**	**Probes**	**InDel[+]**
BP Type1	6,648 (0.02%)	3,324 (2.82%)	4,691 (0.71%)	2,751 (2.6%)	2,092 (0.44%)	1,482 (1.66%)
BP Type2	51,770 (0.2%)	25,885 (21.98%)	38,790 (5.85%)	22,662 (21.39%)	20,540 (4.29%)	14,407 (16.12%)
BP Type3	71,820 (0.27%)	35,910 (30.5%)	41,272 (6.23%)	27,897 (26.34%)	23,631 (4.93%)	18,485 (20.68%)
BP Type4	312 (0.001%)	156 (0.13%)	241 (0.04%)	146 (0.14%)	119 (0.02%)	93 (0.1%)
OTV	872,324 (3.26%)	21,390 (18.16%)	163,278 (24.64%)	18,558 (17.52%)	96,867 (20.22%)	15,064 (16.85%)
MONO	25,735,797 (96.25%)	68,573 (58.23%)	414,500 (62.54%)	65,796 (62.11%)	335,778 (70.1%)	63,597 (71.14%)
ALL	26,738,671	117,756	662,772	105,927	479,027	89,393

[+]*Note that a same InDel could be genotyped by several probe types which resulted in the percentage values great than 1.*

259   Among these InDels, 97,748 (83%) can only be targeted with either internal or external probes, but not
260   both (Figure 3A). These results support our overall strategy which includes the discovery of InDels, with
261   precise breakpoints in a preliminary step, and the use of complementary internal/external probes for the
262   genotyping of large InDels.

Array design

We used the Affymetrix® recommendations to select the 700,000 probes to be included in the final array, plus some other criteria depending on the probe type. Nevertheless, because of their added value, we decided to keep all BP probes as long as they had less than 3 hits on the B73 reference genome sequence. This first selection consumed 84,994 probes targeting 53,456 InDels, among which 70% could only be targeted by BP probes. Concerning OTV and MONO probes, we first selected *neutral* and *recommended* probes having no hit at all (for insertions), and only one hit (for deletions), against the B73 reference genome sequence. We then considered their density with the objective to maximize the number of InDels that could be surveyed, as well as to have an even distribution of probes along targeted InDel sequences (see Methods). We then performed a second selection among *not recommended* OTV and MONO probes for 4,541 InDels that were still not targeted. After filtering some duplicated probes, we built a final array design containing 662,772 probes targeting 105,927 InDels that represent a cumulated length of 232 Mbp, including 25.9 Mbp of PARs.

Description of the array content

The final array design allows genotyping InDels with various sizes, ranging from 37 bp to 129.7 kbp, with a median of 501 bp (Figure 2). They are covered by 1 to 482 probes, with a median of 3 probes per InDel (Additional file 2: Figure S4). The number of probes does not always reflect the length of the InDels, as the proportion of PARs within InDels is highly variable (Figure 2A). 8,040 InDels (ranging from 37 bp to 2,409 bp, with a median of 163 bp) were completely covered by PARs and could thus be considered as a proper PAVs, 34,372 InDels (ranging from 101 to 129,700 bp with a median of 320 bp) were not covered by any PAR at all (Figure 2A). The biggest InDels contains more frequently PARs than the little ones (Figure 2B). In fact, the number of internal probes were more strongly correlated to the size of the PARs (r2 = 0.79) rather than to the size of the InDels (r2 = 0.16) (Additional file 2: Figure S5).

As expected, the probe selection process did not impact the overall distribution of probe types among targeted InDels, as 35% of them can exclusively be genotyped by BP probes, and 50% can only be genotyped by internal probes, among which 73% are only targeted by the use of the original MONO probes (Figure 3B). Indeed, a large number of InDels did not contain PARs and cannot be genotyped with 35bp internal probes but only with BP probes. Whereas, others InDels contains PARs but have no BP probes due to the InDel discovery approach ("no map").

Among the 43,117 InDels that could be anchored onto the B73 reference genome sequence and which were included in the array design, 13,737 were located inside a gene, 57 close to a gene (less than 1 kb away), 1,311 inside a pseudo-gene and 2,212 inside a transposable element. InDels and probe density varied across each chromosome (Additional file 2: Figure S6). We observed a higher density in chromosome arms than in peri-centromeric regions (Additional file 2: Figure S6). We also identified clusters of InDels with a large specific sequence at the beginning of chromosome 6 (10-20Mbp) or at the end of chromosome 5 (~190Mbp).

Assessing array quality by genotyping 105,927 InDels on 480 maize DNA samples

InDel calling using dedicated Affymetrix® pipelines

We genotyped 480 maize DNA samples including 440 inbred lines, 24 highly recombinant inbred lines and 16 F1 hybrids. Dedicated Affymetrix® pipelines were implemented for each of the probe types to call genotype of the InDels based on fluorescent intensity and contrast variation of the probes. It included two algorithms already developed by Affymetrix® [36] for BP and OTV probes (Additional file 2: Figure S7A et B) and a third one, which was newly developed for the calling of presence/absence genotypes using MONO probes (Additional file 2: Figure S7C). 35 DNA samples including all F1 hybrids, did not pass Affymetrix® quality control due to their low call rate (<0.9) and were eliminated. Call rate of the 445 remaining samples, which are all inbred lines, varied from 96% to 99% with a median of 98%. The call rate varied according to probe type (median of 90% and 99% for BP and internal probes, respectively). Out of 662,772 probes, 479,027 probes representing 89,393 InDels passed Affymetrix® quality control and were called on 445 DNA samples. Respectively 55%, 59%, and 81% of BP, OTV, and MONO probes were converted into recommended markers after clustering by Affymetrix® pipelines (Additional file 1: Table S2, S3, and S4). 94% of these recommended BP and OTV markers were classified as "PolyHighResolution" (PHR) indicating a high quality of clustering and that these markers were polymorphic (Additional file 2: Figure S8). Note that the criteria defining high quality of clustering for MONO probes called by new Hom2OTV algorithm was not yet implemented in Affymetrix pipeline (Additional file 1: Table S4 and Additional file 2: Figure S7C). As a consequence, classification of MONO probes could not be comparable to BP and OTV probes. Thanks to the 3 probe types and redundancy, 84% of all InDels could be called with an average of 5.4 probes per InDel.

To evaluate the genotyping ability of the 479,027 probes, we first compared the clustering of inbred lines expected for three probe types (BP, OTV, and MONO) with the observed clustering of inbred lines based on fluorescence intensity and contrast of 445 inbred lines genotyped with the array. For BP probes, we expected at least two clusters corresponding to the individuals homozygous either for presence ("AA" or "BB") or absence ("OO"). A third cluster could be observed when individuals were heterozygous individuals for presence/absence ("OA" or "OB" hemizygous) (Figure 1C). For OTV probes, we expected at least 3 different clusters: two cluster corresponding to the individuals homozygous for allele A or B of SNP ("AA", "BB"), and a third "off-target" cluster for the individuals homozygous for absence ("OO"). A fourth cluster could be observed when some individuals were heterozygous at the within-InDel SNPs (AB). For MONO probes, we expected only two clusters corresponding to the individuals for which the sequence was present ("AA" or "BB") or absent ("OO", "AA" or "BB") (Figure 1C). The observed clustering by the three dedicated pipelines was consistent with the expected clustering for 43% of BP, 83% of OTV and 63% of MONO probes (Table 3).

Table 3: Comparison between the clustering expected for BP, MONO, and OTV probe types and the clustering produced by Affymetrix® pipelines based on the fluorescent intensity and contrast of 445 inbred lines for 479,027 probes.

Probe types		Classification based on the clustering produced by Affymetrix® pipelines and genotyping assignment					
BP		**BP**	**OTV**				
	Number (%)	20,370 (43.9%)	26,012 (56.1%)				
	Clustering examples						
	Description	Two homoz. clusters	Two homoz. and one OT clusters				
OTV		**OTV**	**MONO**	**SNP**	**monomorphic**		
	Number (%)	78,799 (81.3%)	502 (0.5%)	17,562 (18.1%)	4 (0.0%)		
	Clustering examples						
	Description	Two homoz. and one OT clusters	One homoz. and one OT clusters	Two homoz. clusters	One cluster		
MONO		**MONO**	**OTV**	**Unexpected MONO 1**	**SNP**	**Unexpected MONO 2**	**monomorphic**
	Number (%)	212,434 (63,3%)	15,690 (4,7%)	68,562 (20,4%)	1,981 (0.6%)	9,525 (2.8%)	27,586 (8.29%)
	Clustering examples						
	Description	One homoz. and one OT clusters	Two homoz. and one OT clusters	One homoz., one OT and one het. clusters	Two homoz. clusters	One homoz. and one het. clusters	One cluster

"Clustering example": typical example of clustering based on the fluorescent intensity (y-axis) and contrast (x-axis). Colors on figure indicate the assignation of the genotype to the individuals based on this clustering; "Number (%)": Number (percentage) of probes displaying the corresponding clustering. "Description": Brief characteristic of each classification based on the clustering of individuals (homoz.= homozygote, het=heterozygous, OT= off-target).

We observed also some unexpected clustering. For 57% of BP probes, we observed an additional off-target cluster (OTV in Table 3). This indicates that some BP probes did not hybridize properly in some inbred lines, which can either be due to the presence of polymorphism within flanking sequences of the targeted InDels or to the existence of more complex rearrangements removing the breakpoints.

Regarding MONO probes, 25% displayed additional cluster(s) when the sequence was present suggesting the presence of single nucleotide polymorphisms at this position. Among these, we were able to distinguish two types of clustering (Table 3). 4.7% of MONO probes exhibited a clustering similar to those observed for OTV probes suggesting that these MONO probes revealed, by chance, a single nucleotide polymorphism. In contrast, 20.4% of MONO probes displayed an unexpected clustering pattern for inbred lines with the presence of a heterozygous cluster but absence of a second homozygous cluster for SNP (Additional file 2: Figure S9B). In the end, 2.8% of MONO probes displayed an additional heterozygous cluster for SNP when the sequence is present but no "off target" cluster corresponding to individuals for which the sequence is absent (Additional file 2: Figure S9D).

For 18% of OTV (Additional file 2: Figure S9A) and 8.3% of MONO probes, clustering displayed no "off target" cluster for absence, suggesting no presence/absence polymorphism at this position (Table 3). Note that some BP were also classified as monomorphic for presence/absence but were filtered out by the BP pipeline ("MonoHighResolution" in Additional file 1: Table S2 and Additional file 2: Figure S8). These monomorphic probes originated from false positive discovery of InDels or PARs within InDels that are not present/absent elsewhere in the genome of four lines (see Discussion). After removing these monomorphic probes for presence/absence, 422,369 probes allowed us to successfully genotype a total of 86,648 InDels (82% of 105,927 InDels targeted by the array) on 445 inbred lines.

Evaluation of genotyping reproducibility and quality

Consistency of genotyping among the four inbred lines used for InDel discovery

We used the 479,027 probes passing Affymetrix® quality controls to evaluate the quality of Presence/Absence genotyping by comparing the genotyping results obtained from our array (GBA: Genotyping By Array) with those from sequencing (GBS: Genotyping by Sequencing) for the 4 lines used for the discovery of InDels (B73, F2, PH207, and C103). Respectively, 97%, 912%, and 88% of the BP, OTV, and MONO probes had a genotyping result consistent with results obtained from BLAST alignments against our three draft genome assemblies and the B73 reference genome. We observed a strong asymmetry for concordance rates for internal probes (OTV and MONO) depending on whether the genotype has been called by sequencing as present or absent (95% vs 80% present and absent, respectively, Table 4). Interestingly, we observed no asymmetry for BP probes that are designed exclusively on B73 genome compared to OTV and MONO probes that are designed from the 4 genome assemblies (Table 4). These low consistencies for internal probes when genotype by sequencing indicated absence could be explained by the use of incompletely assembled genomes of the three lines (PH207, C103, F2) to call the presence/absence genotype from sequencing.

Table 4: Consistency rate between genotyping by sequencing and by array for the 4 individuals used to discover the InDels, for the three probe types and for the two different genotypes observed from sequencing: presence (P) or absence (A).

Probe Types	Genotype by sequencing	B73	F2	C103	PH207	All Lines
BP	A	0,98	0,98	0,98	0,97	0,98
	P	0,97	0,97	0,97	0,96	0,97
	ALL	0,97	0,97	0,97	0,97	0,97
OTV	A	0,85	0,89	0,80	0,78	0,83
	P	0,93	0,97	0,96	0,96	0,96
	ALL	0,90	0,95	0,91	0,90	0,92
MONO	A	0,77	0,81	0,82	0,81	0,80
	P	0,90	0,98	0,94	0,94	0,95
	ALL	0,82	0,94	0,89	0,88	0,88
ALL	A	0,80	0,86	0,84	0,82	0,82
	P	0,92	0,97	0,94	0,95	0,95
	ALL	0,85	0,95	0,90	0,89	0,90

*Note that consistency rate of hemizygous genotypes (heterozygous for presence / absence) were not displayed in the table for BP probes but considered to estimate global consistency rate (ALL). Note that the absence of probe sequence due to absence of hybridization or no alignment on draft sequence of BP probes were considered as missing data. Missing data were not included in the comparison for all probes.

If the genomic region containing the InDels were absent or badly assembled in at least one line, some
probes would not align properly, resulting in false absence calls, instead of presence in GBS. The four
inbred lines showed very similar concordance rates, F2 being the most concordant (95%). This could be
partially explained by the higher proportion of GBS present calls in F2 as compared to the three other lines
since GBS present calls are more consistent with GBA than GBS absent calls. The median consistency rate
of probes within InDels remained relatively high and stable, around 90%, independently of the number of
probes per InDel (Figure S10), suggesting no relationship between the consistency rate of individual
probes and length of PARs within InDels.

Consistency among probes from the same InDel

To estimate the consistency of different probes for typing a given InDel, we analyzed genotyping
results for 50,648 InDels genotyped with at least two probes in a collection of 362 temperate inbred lines.
For each InDel and each inbred line, we calculated the average allelic frequency of presence over all
probes. Frequencies of 1 (presence) and 0 (absence) indicated that all probes displayed consistent
genotyping for the corresponding inbred line (Figure 1D and Figure S11A). Alternatively, frequencies
different from 0 or 1 (FreqDiff01) indicated that at least one probe displayed inconsistent genotyping with
other probes for corresponding inbred lines (Figure S11B). Overall, 75% of the InDel genotyping resulted
in an average allelic frequency for the presence of 1 or 0, meaning that all probes had a consistent
genotyping results for calling the allele at both present or absent states, respectively (Figure 4A).

However, we observed a strong variation of median (average) allelic frequency difference from 0
or 1 (FreqDiff01), according to the number of probe interrogating that InDel (Figure 4B, Additional file 1:
Table S5). Median (average) FreqDiff01 across InDels varied from of 1.2% (9.8%) to 58% (52%) when the

number of probes varied from 0 to 30 (Figure 4B, Additional file 1: Table S5). We compared this variation
to what could be expected for different probe genotyping error rates (1%, 3%, 5%, and 10%). Based on
this comparison, we estimated the probe genotyping error rate is approximately 3% (Figure 4). For InDels
with fewer probes (<10), the average and median FreqDiff01 differed slightly, suggesting that some InDels
with low probe numbers displayed high genotyping inconsistencies among their probes (Figure 4,
Additional file 1: Table S5). In order to evaluate whether probe genotyping error is similar for present or
absent calls, we analyzed the variation of FreqDiff01 with regard to the average frequency of absence of
InDel sequences in 362 lines (Additional file 2: Figure S12A). The median FreqDiff01 was higher for InDels
which have their sequence more frequently absent than present across 362 lines, regardless of the
number of probes (Additional file 2: Figure S12B). It suggested that genotyping was more accurate for
absence than presence. This was logical, considering that polymorphisms within probes would preclude
hybridization of the probes for some lines, and it would result in absent calls with MONO and OTV probes,
while polymorphisms within probes have no impact when the sequences are absent.

Combining genotyping from multiple probes within InDels greatly improved reliability of InDel
genotyping, since it allowed (i) to correct the individual genotyping errors due to polymorphisms within
probe sequences, (ii) to reduce the missing data rate due to bad clustering or probes polymorphisms, and
(iii) to remove probes displaying highly-divergent genotypes compared to other probes for the same InDel,
due, for example, to a bad design of the probes. In order to evaluate the combining of genotypes of several
probes on the accuracy of InDel genotyping, we simulated global genotyping error rates for InDels by
assigning to each inbred line the most frequent allele, based on the average frequency over all probes
from an InDel, with various genotyping error rates (Additional file 1: Table S6). By this approach, the
genotyping error for InDels was greatly reduced. Considering a probe genotyping error of 5%, the
genotyping error of InDels for inbred lines were reduced to 0.2% and 0.1%, when the number of probes
within the InDels were 2 and 5, respectively (Additional file 1: Table S6). Combining genotypes from
several probes also strongly reduced the average missing data rate for InDels; it decreased from 2.3% to
0.2%, when the number of probes increased from 2 to 5 (Additional file 1: Table S5). However, some
contradictory probe genotypes were repeatedly found across the 362 samples (Additional file 2: Figure
S11B), suggesting that some probe inconsistencies could have biological origins (*i.e* more complex
rearrangement), rather than being genotyping errors. Additionally, 35% of InDels called by BP had their
FW and REV probes classified differently (e.g. one as BP and the other as OTV). Altogether, these results
suggest that some calling inconsistencies between probes within InDels could come from polymorphisms
in the flanking sequence while some other could be due to local rearrangements in the genotyped lines
as compared to the lines used for InDels discovery.

Reproducibility and Mendelian inheritance
Genotyping reproducibility was evaluated by comparing genotypes between five DNA replicates
corresponding to unique F1 hybrids derived from a cross between B73 and F72 for all probes type. Median
reproducibility was 95%, 96%, and 97% for BP, OTV and MONO probes respectively. Interestingly, there is
some variation of reproducibility relative to probe clustering (Additional file 1: Table S7). Note that
Affymetrix[©] algorithms were not specified to genotype hemizygote using OTV and MONO probes in this
dataset. We also performed a parent-offspring analysis on 12 F1 hybrids derived from 9 parental lines by

comparing genotypes observed of these F1 hybrids with those predicted from genotypes of their two parental lines for 46,382 BP probes (Additional file 1: Table S8). On average, 95% and 77% of observed genotypes were consistent with those predicted from parental lines for homozygous and hemizygous genotypes, respectively (Additional file 1: Table S8). The consistency rate was slightly higher when genotypes were absent (98%) than present (94.5%). Note that the seed-lot of parental lines used for producing F1 hybrids were different from those genotyped, which could explain lower consistencies rate than for DNA replicate of F1 hybrids. Note also that the genotypes of all F1 hybrids have been initially eliminated by Affymetrix® quality control due to their low call rate and were therefore forced for reproducibility analysis. This low call rates can be attributed to the fact that these samples had different genotype cluster properties (probe intensity profiles) compared to the samples that passed QC. As a consequence, this strongly increased the missing data rate for the F1 hybrids for OTV and MONO probes.

In the end, we evaluated genotyping reproducibility for inbred lines, by comparing the genotyping results of 13 different inbred lines that were replicated in the experiment (Additional file 1: Table S9). Note that these are not perfect biological replicates, as they represent the same variety but come either from different seed lots or from different accessions. These replicates exhibited a genotyping difference varying from 0.6% to 5.2% (Median = 1.7%, Additional file 1: Table S9). This is similar to the amount of inconsistencies obtained on the same material using a 50K SNP array [54], suggesting that InDel genotyping inconsistencies for replicates can be attributed mostly to seed-lot divergences, rather than genotyping errors (Additional file 1: Table S9). However, genotyping reproducibility was higher for these inbred lines than for the DNA replicates of the F1 hybrid, suggesting that errors in F1 hybrids can mostly be attributed to the inability to genotype hemizygous with OTV and MONO probe for this small dataset.

Application: Diversity analysis of 362 maize inbred lines panel

In order to evaluate this new array for genetic analysis, we analyzed genetic diversity using 57,824 polymorphic InDels on a subset of 362 inbred lines, representing genetic variation that has been successfully used to decipher maize genetic structuration and perform genome-wide association studies [55–57]. To represent each InDel in the diversity analysis, we selected one single probe per InDel, based on the probe genotyping quality (see Methods).

We first compared kinship values between 362 inbred lines estimated with 57,824 InDels and with 28,143 prefixed Panzea SNPs from the 50K SNP array. Kinship values between lines obtained with SNPs and InDels were strongly similar and highly correlated (r=0.9), except those for a couple of lines closely related to B73 and F2 (Additional file 2: Figure S13). Then, we performed Principal Coordinate Analysis (PCoA) based on the genetic distance between 362 lines estimated by InDels and SNPs (Figure 5). We included on this PCoA the genetic structuration of these 362 inbred lines, as obtained from the prefixed Panzea SNPs from the 50K SNP array [55]. The global genetic structure developed using two types of polymorphisms are highly similar. The first axis showed good discrimination of European Flint from Corn Belt Dent and Stiff Stalk lines, while the second axis discriminated European Flint and Northern Flint lines.

Overall, the clustering of individuals based on genetic distance estimated with InDels (Figure 5A) was
consistent with those estimated with SNPs (Figure 5B). We observed that B73 and F2, which were used to
discover the majority of InDels, were more contrasted on PCoA when genetic distance was estimated with
InDels, as compared with SNPs from the 50K array, indicating some ascertainment bias. We thus
performed two PCoAs, with InDels and SNPs, excluding B73 and F2 (Additional file 2: Figure S14). The two
PCoAs gave similar patterns, suggesting that this ascertainment bias was largely removed when no close
relative lines from those used for discovering InDels were used in diversity analysis. Due to this
ascertainment bias, result of our array should be therefore interpreted with caution for diversity analysis.

Discussion

1. An original high throughput approach for genotyping InDels

The comparison of whole genome sequence assemblies is in theory the best approach to identify,
precisely and exhaustively, structural variations between two individuals. But even though great progress
has been made recently in this area, high-quality, whole genome assembly is still too costly, time-
consuming, and computationally intensive to be applied to hundreds of individuals, especially when
considering the complexity of the maize genome [20, 58]. Other whole genome sequencing approaches
based on alignment of reads on a single reference, and using either "read-depth", "read-pair", or "split-
read" identification methods [46–50] have mostly been limited to the identification of deletions (i.e.
sequences absent from a reference genome). Liu et al., (2015) partially addressed the lack of insertions
(i.e. novel sequences compared to a reference genome) by the identification 1,973,746 InDels [4].
Although, among these a majority were very small (85% smaller than 11bp), and the use of PCR markers
to genotype them is time-demanding, labor-intensive, and costly at a large-scale level. To avoid this
ascertainment bias due to use of a single reference genome to genotype SVs, other studies proposed to
call SVs by aligning reads from sequencing on a pan-genome representing the combination of several
genomes [14, 20, 22, 52]. However, genotyping InDels with high reliability and call rate by these
approaches required at least 30X-50X coverage of the genome to correctly cover their breakpoint and
their internal sequence, especially to genotype InDels larger than 50bp [52]. Additionally, aligning reads
from a thousand individuals on a pan-genome remained computationally intensive, and therefore
required large informatics facilities [52]. In the end, these approaches required to build a pan-genome of
high-quality, which remains challenging for a complex genome.

In this paper we describe a new approach combining (i) the 'accuracy' of detecting InDels using whole
genome assembly, with the detection of 89,150 insertions and 52,175 deletions from the comparison of
three newly sequenced and assembled maize inbred line (F2, PH207, and C103) genomes and the public
maize B73 AGPv2 reference genome, (ii) and the 'high-throughput' genotyping utility provided by SNP
arrays. This approach allowed us to genotype, for the first time, thousands of insertion/deletion variants,
including PAVs, on a few hundred maize individuals. Genotyping cost per individual using the InDel array
was at least 10-20 fold cheaper than any approach based on sequencing for a species with a genome as

complex as maize, at a similar level of reliability (> 1000€-2000€ for a 30-50X of a 3Gbp genome *vs* 50€-
220€ using Affymetrix® Axiom® array, depending on the number of samples and probes). This genotyping
cost did not include bioinformatics analysis. Calling SVs from a pan-genome of a species with a large and
complex genome, such as maize, was time-consuming and required bioinformatics skills and large
informatics facilities, which are costly and not available in all laboratories. In the contrary, the array could
be analyzed rapidly on a laptop using a pipeline already implemented for analyzing SNPs and the
Hom2OTV R script developed for analyzing MONO probes. Additionally, the array provided a wet-lab
validation of the InDel discovery and allowed the removal of putative genotyping errors from sequencing
(particularly for PAVs), due to incomplete or bad genome assembly, as we observed in our study. In the
end, the probe content of the InDel array can be largely optimized, either to reduce the size of array (and
therefore the cost), or to increase the number of SVs genotyped, without losing reliability (e.g. 200,000
to 300,000 InDels) by filtering out under-performing probes, by strongly reducing the number of probes
per InDel (2-3), and by removing false positive InDels. It would also be easy to design an array combining
probes targeting InDels and more classical SNPs, outside of InDel sequences.

With the use of breakpoint probes for both insertions and deletions, our approach overcomes some
of the limitations of previous CGH or SNP array-based studies, which were only able to call deletions if a
few successive probes had lower fluorescent intensity signals [32–35]. Unterseer et al., (2014) genotyped
specifically 6,759 small deletions, which were discovered by aligning reads of 30 inbred lines against the
B73 genome, but the study did not include any insertions [37]. However, previous CGH and SNP arrays did
not design probes to target breakpoints and detected InDels by analyzing the variation of fluorescent
intensity signals of ordered probes [32–34]. Consequently, these technologies targeted exclusively low
copy regions of the genome, excluding InDels containing repeats, such as transposable elements (TEs) [2,
30, 44]. This is a strong drawback for maize and many other crops since a large part of their sequence is
composed of transposable elements [28, 59] which may be highly variable between individuals [4, 24, 60]
and may impact phenotypes [61–63]. The use of BP probes allows to target Present/Absent Variations,
whose sequence are unique and not present elsewhere in the genome, as well transposable elements,
whose internal sequence can be present/absent at one specific locus but also present elsewhere in the
genome. Another advantage of genotyping breakpoints is that it provides the ability to genotype the same
mutational event across all individuals of the population, as it is highly unlikely that two independent
mutational events could lead to the exact same breakpoint. On the contrary, for InDels detected using
classical CGH or SNP arrays, it is much harder to identify common InDels among a population of
individuals, as we don't know precisely their breakpoints. Genotyping breakpoints is also very cheap since
only one or two probes are needed , which makes the InDel size no longer a limitation for genotyping it
accurately, contrary to previous SNP and CGH arrays that rely on fluorescent intensity variation of probes
covering the entire InDel sequence [45]. The genotyping of breakpoints by sequencing is possible with a
tool like PInDel [50], which has a genotyping mode or BayesTyper [52], but at a much greater cost and
with lower call rate compared to the use of a SNP array. Finally, breakpoint probes are codominant
markers and allow accurate genotyping of hemizygous individuals (Heterozygous for presence/absence),
since their genotyping is based on fluorescent contrast rather than fluorescent intensity variation, which
is known to be noisier as with MONO and OTV probes [45].

Although the use of BP probes is clearly the simplest way to genotype InDels using an SNP array,
breakpoints are not always available ("no map" approach discovery) or "designable" with 35bp probes,
for instance, the cases where sequences of microhomology at breakpoint site were larger than 5bp. In
order to genotype the 52,471 InDels without breakpoints and explore the genetic diversity within InDels,
we also designed 577,778 internal probes both on monomorphic and polymorphic sites in PARs for both
insertions and deletions. To genotype PARs in InDel sequences using SNPs, we took advantage of the
already available Affymetrix® algorithms to call Off-Target Variants (OTVs), which can detect variation of
fluorescent intensity signals for a single probe (Figure 1C) [36]. This approach was used by [37] who was
able to detect 45,974 OTVs on a set of diverse maize inbred lines using a 600K SNP array. Nevertheless,
the array was designed in a classical way to target SNPs, and there was no prior evidence that the probes
called as OTVs would belong to InDels. Additionally, detecting SNPs in insertions required the assembly of
a pan-genome, combining common and specific sequences from different individuals, in order to retrieve
SNPs by aligning reads from sequenced lines [14, 20, 22, 52]. In our case, only using OTV probes would
have resulted in the elimination of many InDels, since 87,372 of them, including 74,648 insertions, did not
have known SNPs within their sequence. In order to avoid this ascertainment bias due to prior knowledge
of the presence of SNPs we designed 414,500 MONO probes on putative monomorphic sites within PARs
of InDel sequences. This permitted the genotyping of 38,134 supplementary InDels that could not be
targeted by OTV or BP probes. This new type of probe required the development of a new algorithm in
order to cluster individuals according to their fluorescent intensity variation only, to be able to assign a
genotype to each individual (Additional file 2: Figure S7C). A limitation of current workflow is that
Affymetrix® algorithms require a larger number of hemizygous individuals to generate high-quality
genotype clusters using the OTV and MONO probes. While it was not an issue for maize inbred lines (or
individuals from autogamous species) that are mostly homozygous, it was an issue for individuals from
allogamous species that are highly heterozygous. By using alternate genotyping techniques or processing
a larger number of hemizygous samples, it should be possible to identify hemizygous clusters according
to fluorescence intensity from OTV and MONO probes. We observed some clusters that seem incorrectly
interpreted as heterozygote for SNPs, although they likely correspond to hemizygous individuals for OTV
and MONO probes (Additional file 2: Figure S9B, see below for a more detailed discussion). Alternatively,
other algorithms/software based on fluorescent intensity variation of either a single probe or several
ordered probes exist and could be used to detect copy number variation for hemizygote individuals [38–
43].

In the end, we observed some ascertainment bias using our array (Figure 5). This was due to the fact
that our four inbred lines do not well represent the whole genomic diversity of maize, notably missing are
tropical lines. As a consequence, it could lead to ascertainment bias by reinforcing the differentiation of
inbred lines genetically close to the four inbred lines used to discover InDels [54, 64, 65] as we observed
in our diversity analysis for lines close to B73 and F2 (Figure 5 and Figure S13). It could be therefore highly
valuable to use more lines for the initial InDel discovery step. Several new individual maize genome
assemblies are now available in the public domain and more and more could become available in the
future. Our approach could easily be applied to these new genome assemblies to discover new InDels on
a larger set of inbred lines representative of maize diversity with the aim to design a new InDel array.

2. Reliability of genotyping / calling results

Our approach provides a reliable and reproducible method for genotyping InDels in inbred lines, since
(i) the genotypes obtained by array and by sequencing were highly consistent for BP probes (97%) and in
a lesser extent with OTV and MONO probes (92% and 88%, respectively), due to the fact that the genome
assembly of sequenced lines were incomplete or incorrect, resulting in high error rates for absent calls
using GBS ; (ii) the average probe genotyping error rate was estimated at 3% (lower for absent calls); (iii)
the InDel genotyping errors could be greatly reduced by combining the genotypes of different probes
within the InDels (0.02% for 5 probes); (iv) the genotyping results were highly reproducible between DNA
replicates of F1 hybrids (95 to 97%, depending on probe type) and between inbred lines (94.8 to 99.4%);
and (v) the call rate for individuals was very high (96 to 99%) and can be increased by combining the
genotypes of the probes within the InDels (97.7 to 99.9% for 2 and 5 probes, respectively).

Our approach is promising as a method to genotype structural variations in maize, as well as other
species with complex genomes. We obtained high metrics, comparable to classical SNP arrays, based on
Affymetrix® Axiom® Technologies, even though InDels are more complex to genotype. First, call rates are
high and quite similar to those obtained for SNP with the 600K SNP Affymetrix® array (98% against 98.1%
in [37]). Nevertheless, we observed a lowest call rate for BP probes (90%). This lowest call rate could be
explained by the usage of more relaxing criterion to filter out probes for building array and by the fact
that polymorphisms in surrounding sequences of InDel breakpoints have not been taken into account
contrary to internal probes. Second, the percentage of BP and OTV probes classified as PHR (94% in both
cases) is similar than for 600K SNP Affymetrix® genotyping array (92%) but higher than for 1.2M screening
Affymetrix® arrays (~65%) that have been used to select best markers for designing the final 600K SNP
Affymetrix® arrays. It is difficult to compare the classification of MONO probes, because the algorithm
used (Hom2OTV) is new and quite different from the one used for BP, OTV, and classical SNPs. Third, the
reproducibility between DNA replicates of F1 hybrids was high (95 to 97%, depending on probe type), but
this is lower than for SNP arrays (~99.5% in [37]). However, the reproducibility was estimated on DNA
replicate of F1 hybrids in our study while it was estimated on inbred lines for 600K SNP Affymetrix® array.
When we compared genotype of 13 inbred lines originated from different seedlots, reproducibility is close
to those of 600K SNP Affymetrix® array (98.3%) and displayed approximatively same reproducibility with
50K SNP Illumina array ([54], Additional file 1: Table S9). This comparison suggested strongly that our
lower reproducibility might not be due to genotyping errors but possibly the divergence between the
samples for inbred lines and the use of F1 hybrids rather than inbred lines for DNA replicate. Fourth, the
Mendelian inheritance between F1 hybrids and their parental lines was lower for our InDel than for SNP
array (88% vs 97.6% in [37]) but quite similar considering only homozygous genotypes (95%). This is likely
due to the presence of a small number of hemizygous samples since the 16 F1 hybrids were eliminated
due to their low call rate (<0.9) and there are only residual hemizygosity for inbred lines. Considering the
F1 hybrids for defining BP cluster could improve the delineation of hemizygous cluster and therefore
Mendelian inheritance. Note that 600K SNP Affymetrix® in maize was designing by selecting the high
confidence probes based on results of a first screening 1.2M SNP Affymetrix® array which could favor
reproducibility for this array. Finally, 72% of probes were converted into markers, which is comparable to
this 1.2 maize Affymetrix® Axiom® SNP screening arrays (74.9% in [37]). Out of these, 88% were

polymorphic for presence/absence. This conversion rate is expected, considering that Affymetrix® Axiom®
array analysis pipelines have been optimized for the detection of bi-allelic SNPs and are more sensitive to
variations in fluorescent contrast (x-axis) compared to variations in fluorescent intensity (y-axis), which is
known to be noisier [36, 45]. Moreover, we did not always follow Affymetrix® recommendations, as we
did not filter out probes with a bad design score.

We identified some inconsistencies between genotyping by array (GBA) and genotyping by
sequencing (GBS) obtained by aligning probes against our genomes (Table 4). These inconsistencies were
higher when GBS called absent for InDels interrogated by OTV and MONO probes (17.1% and 20.2% vs.
4.3% and 5.4%, respectively), although no differences were observed for BP probes (Table 4). These biased
inconsistencies towards absence for internal probes seems very high compared to our analysis on the
consistencies between probes within Indels. Our analysis of consistencies between probes within InDels
showed indeed that genotyping errors produced by the array were close to 3% (Figure 4) and lower for
absent calls (Additional file 2: Figure S12). These results suggested that the higher genotyping
inconsistencies for GBS absent are due to errors in GBS. GBS errors for absence were well explained by
the use of an incomplete genome draft assembly to align probes sequences, and the use of a higher-
quality genome could help to reduce these inconsistencies. The probes targeting sequence regions
present in one line, but not assembled in their draft genome assemblies, were falsely genotyped absent,
but the sample DNA correctly hybridized with the probes, and the InDels were called present with the
array. This could also explain why the number of inconsistencies was higher for B73, as all B73 absence
genotypes were defined in comparison to draft assemblies. Whereas for the other 3 lines, absence
genotypes were defined in comparison with the gold standard B73 genome sequence. The fact that we
obtained a better result on OTV probes interrogating InDels discovered in F2 can be explained because
we used only SNPs discovered on the B73-F2 pan-genome and not in other genomes. And, the fact that
BP probes had similar consistencies for genotyping absent and present calls could be explained by the fact
that the BP probes were designed exclusively on B73 reference genome.

We also found that 20,574 InDels were monomorphic and present across all lines, suggesting they
represented false positives from regions not assembled in our draft genomes. To reduce this false positive
rate, we strongly advise to not only align B73 reads onto each draft genome assembly but to also align
reads from each sequenced genome on each other and against itself. This would have several benefits: (i)
it would allow to discover even more and higher-quality InDels, as each putative deletion discovered in
one sample could potentially benefit from supporting reads from another sample; (ii) this would simplify
the identification of InDels common to more than one genotype; and (iii) it would help to identify and
eliminate false positive deletions by the alignment of each sample on its own draft assembly.

Nevertheless, the use of incomplete draft genomes does not explain all discrepancies between
genotypes obtained by sequencing and by array. First, these discrepancies could also be due to incorrect
clustering and assignment of a genotype call (array errors). This was exemplified by some MONO probes
classified as SNPs, although the clustering pattern looks like a MONO cluster with a large difference of
fluorescence intensity between two clusters (Additional file 2: Figure S9C). A more detailed inspection of
the clustering of MONO probes displayed an unexpected cluster pattern (Table 4, Additional file 2: Figure
S9D), and OTV probes classified as SNPs (Table 4, Additional file 2: Figure S9A) suggests a wrong

assignment of genotypes for the cluster displaying the lowest fluorescent intensity. Similarly, the genome
divergence within probe sequences for some inbred lines could result to group those individuals in an OTV
cluster, and therefore result in an incorrect absent call. However, these genotyping errors due to bad
clustering or genomic divergence between individuals within probes sequences could be strongly reduced
by combining genotypes from several probes. As an InDel called by five different probes has a random
genotyping error of 5%, we showed by simulation that the genotyping error for that InDel would be
reduced to 0.1%, when the most frequent allele among the 5 probes was assigned as genotype of the
InDel (Additional file 1: Table S6).

Surprisingly, 4.7% of MONO probes displayed a classical OTV clustering, suggesting that an unknown
SNP was targeted by these probes by chance. This high level of polymorphism (1 SNP / 21 bp) was slightly
higher than observed by sequencing a small set of diverse lines [66, 67]. It could suggest that PAR genomic
regions might have more divergence than other parts of the genome, because these regions were involved
in local adaptation by maintaining together favorable combinations of alleles as proposed by [68]. These
15,690 new OTVs are very interesting, since they were discovered by chance on a large set of 445 inbred
lines. We could therefore expect that these OTVs have no ascertainment bias, which can be very useful
for analyzing genetic diversity within InDels carrying PAR regions. In addition, 20.4% of MONO probes
displayed unexpected clustering: one off-target cluster, corresponding to absence of the sequence; one
homozygous cluster, corresponding to presence of the sequence; and an unexpected heterozygous cluster
(Unexpected MONO 1 in table 4). Considering these "unexpected MONO 1" as true SNPs would indicate
a density of 1 SNP every 5 bp, which is not compatible with the level of diversity observed in previous
studies of maize [66, 67]. Deeper investigation of these MONO probe clusters identified that for some
probes, the unexpected heterozygous cluster is positioned between the presence and absence clusters
(Additional file 2: Figure S9B). This suggests that these unexpected heterozygous clusters are identifying
inbred samples with only one copy presence (hemizygous genotype). An alternative hypothesis to explain
this unexpected pattern is the presence of divergent duplicated sequences, leading to the existence of an
artificial heterozygous cluster for SNPs corresponding to the presence of two paralogous sequences. This
result suggests therefore that there is probably room to develop genotyping strategies in order to better
identify additional clusters corresponding to the presence of hemizygous individuals for both MONO and
OTV probes and therefore improve the quality of the genotyping of InDels when using a SNP array.

These potential clustering errors, as well as the incorrect design of some probes, can explain some
inconsistent genotypes for presence/absence between probes for the same InDel. Comparison of
genotyping across different probes within InDels could help to identify and remove probes displaying
highly discordant genotypes, due to errors originating from poor clustering or from poor design.
Interestingly, some InDels showed reproducible inconsistent genotypes for presence/absence across their
probes in several inbred lines (Additional file 2: Figure S9B). This suggested that this pattern could have a
biological origin, with possible rearrangements having occurred several times within the same genomic
region in some inbred lines. Following this hypothesis, Gu et al. (2008) observed two different types of
rearrangements which could explain our observations [69]: (i) rearrangements with an unique breakpoint
in population and therefore common size between individuals resulting to two haplotypes in a population
and (ii) rearrangement with non-unique breakpoints, scattered in a genomic region, which resulted in

several haplotypes. This hypothesis is also supported in our experiment by the 56% of BP probes classified
as OTVs, indicating that FW or/and REV flanking sequence did not hybridize in some lines.

The development of a statistical approach to merge either *a posteriori* the calling results of
independent clustering of individual probes or *a priori* the fluorescent intensity signal of successive probes
within a InDel could be interesting in order to improve the robustness of InDel genotyping. This would
have the advantage to limit the effect of genotyping errors due to poor clustering and to reduce the noise
in fluorescent intensity signals. We showed by simulation that assigning the most frequent allele across
probes as the genotype reduced genotyping error to 0.7% and 0.1% when 3 and 5 probes were used,
respectively. Additionally, it increases the InDel call rate (Additional file 1: Table S6). In the end, it would
also help to identify varying haplotypes, representing the complexity of a region in a population. Using
multiple probes for calling InDels is therefore highly valuable for improving reliability of InDel genotyping,
since it allows putatively to reduce random genotyping error, due to genomic divergence or other causes,
removes probes poorly clustered or designed, and identifies more complex rearrangements.

3. Conclusions

Our approach, from the sequencing of a few representative genotypes, their genome assembly, the
insertion/deletion discovery, and to the design and use of the high-throughput genotyping array was
applied to maize as a proof of concept. Our approach allowed us to rapidly create at a reasonable cost a
high-throughput SVs genotyping tool for this species. This approach will remain interesting as long as
calling large InDels from sequencing, for a large set of individuals, remains un-affordable, bioinformatically
challenging, and time-demanding. Nevertheless, our approach could benefit from few improvements
based on the knowledge accumulated from this test on maize. First, it could be highly valuable to use
more lines for the initial InDel discovery step to avoid ascertainment bias [64] as we observed in our
diversity analysis (Figure 5). Using more lines for detecting InDels should also reduce the number of false
positives SVs in array due to poor assembly, genotyping error due to genomic divergence between
individuals, and help to identify complex rearrangement. Second, even though we did not have any
indication that our sequenced data had been contaminated, a contamination cleaning step should be
applied to the sequenced data prior to SVs discovery and genome assembly, in order to avoid potential
false positive SVs in the final array. Third, aligning reads against the internal sequence of InDels, as well as
aligning probes sequences against each genome assemblies, should strongly reduce false positives in the
final array. Fourth, improving the pipeline of MONO and OTV probes to call hemizygous genotype from
variation of fluorescent data would be very valuable, notably for allogamous species. Fifth, capacity of
array could be largely increased to 200,000 or 300,000 InDels without losing reliability by optimizing
number of probes per InDels.

To conclude, we developed a "proof of concept" high-throughput and affordable InDel genotyping
array, based on the InDels discovered by sequencing on four inbred lines. Our "proof of concept" approach
could be easily applied to other species to explore genomic structural variation, notably species with
limited sequence data or few genome assemblies available. This could also be interesting for species with
greater sequencing resources and where genotyping a large set of individuals is required, such as for

breeding purposes, genome wide association studies, genomic selection, or characterizing SVs in large germplasm. Although our array was not designed to genotype duplications and inversions, our approach could be easily extended to genotype breakpoints of inversions, but further development of the pipeline for genotyping duplications using internal probes would be required. This powerful approach opens the way to studying the contribution of InDels and other SVs to trait variation and heterosis in maize [44] and should contribute to decipher the biological impact of InDels and other SVs at a larger scale in different species.

Methods

Sequencing material

Three maize inbred lines, which are key founders of maize breeding programs and originated from three different heterotic groups, had been selected for deep sequencing and InDel discovery: the European Flint line F2 and two American dent lines, PH207 (Iodent) and C103 (Lancaster). For the F2 inbred line, see [20]. For C103 and PH207 inbred lines, DNA was extracted with the NucleoSpin Plant XL, according to the manufacturer's instructions (Macherey Nagel, Düren, Germany). The DNA concentration was estimated by UV measurement and the quality was checked with an agarose gel electrophoresis. Two library types were sequenced: a 180bp overlapping paired-end library and a 3kb mate-pair library. The paired-end libraries and the sequencing were performed by Integragen (Evry, France) on a HiSeq2000 sequencer (Illumina, San Diego, USA). 412 and 377 million 100bp paired-end reads (33x and 30x) were sequenced respectively for C103 and PH207. The mate-pair libraries were prepared and sequenced at BGI (China) also on HiSeq2000 sequencer (Illumina, San Diego, USA). Raw reads were filtered to remove adaptor sequences, contamination, and low-quality reads. 326 and 316 million 100bp mate-pair reads (26x and 25x) were sequenced, respectively for C103 and PH207. A data set of 473 million B73 inbred line 100bp paired-end reads (35x) with an average insert size of 191bp was downloaded from ftp://ftp.sra.ebi.ac.uk/vol1/fastq/SRR404/SRR404240.

InDel and PAV discovery

For the deletion discovery step, F2, PH207, and C103 paired-end reads were aligned against B73 AGPv2 genome sequence using novoalign version 3.01.01 (http://www.novocraft.com) (default parameters). Samtools [70] version 0.1.18 was used to coordinate, sort, and retain reads with a mapping quality of at least Q30. Duplicated reads were eliminated using MarkDuplicate from the picardtools suite (http://broadinstitute.github.io/picard) version 1.48. PInDel [50] version 0.2.5a2 was run in parallel on each chromosome to perform multi-genotype calling of deletions. Raw formatted results were converted to VCF (Variant Calling Format) using the script PInDel2vcf. BreakDancer [46] was used in complement PInDel, but only for F2. Deletions shorter than 100bp were discarded. Deletions spanning a B73 assembly gap or located in regions prone to mis-assemblies, such as telomeric, knob, and centromeric regions, were also excluded from further analysis using IntersectBed BEDTools [71] version 2.16.1.

For whole genome sequence reconstruction of F2, PH207, and C103 inbred lines, paired-end and
mate-pair reads were used together and assembled using ALLPATHs-LG [72] version R41008 (Additional
File 2: Figure S1B). For F2, the script CacheToAllPathsInputs.pl was used to cache the data to use for
assembly: 100% of the non-overlapping 230bp insert paired end data set, 100% of the overlapping 170bp
insert paired end data set, 30% of the non-overlapping 370bp insert paired end data set, and 100% of the
2.4kb insert mate pair data set. Indeed, only overlapping paired end reads are used by ALLPATHs-LG for
building contigs, but the supplementary non-overlapping paired end reads for F2 were used for error
correction. RunAllPathsLG was then run for all three genotypes using optional parameters. Details on the
sequence library usage during the assembly process are given in Additional file 1: Table S1. For each
assembly, the coverage of the gene space was evaluated using BUSCO [73] version 3.0.2 using genome
mode and the maize species (-m geno -sp maize).

B73 paired-end reads were successively aligned to ALLPATHs-LG F2, PH207, and C103 genome
sequence assemblies (Additional File 2: Figure S1B). The same tools and parameters used to call deletions
against the B73 genome were applied to detect B73 deletions against F2, PH207, and C103 genome
sequences. These B73 deletions were reciprocally called insertions of F2, PH207, and C103. Only insertions
smaller than 100bp were discarded, except those spanning real assembly gaps (with approximate size
inferred from paired reads average distance) and not "unsized" gaps like in the B73 genome. When
possible, insertions were anchored onto the B73 AGPv2 genome sequence using a dedicated pipeline
combining Megablast version 2.2.19 [74] and Age version 0.4 [75]. Again, insertions that could be
anchored on the B73 reference and were overlapping regions prone to mis-assemblies such as telomeric,
knob, and centromeric regions, were also excluded from further analysis using IntersectBed.

F2 specific sequences coming either from the no map approach (Additional file 2: Figure S1) or
from the work of [20] were included as such, without any further filtering.

The multiple references and approaches used during the InDel discovery step led to a set of InDels
with various levels of redundancy. Some "intra-tool" redundancy was found (*e.g.* multiple calls found by
one tool within the same genotype at highly polymorphic loci). These "ambiguous" calls were
systematically identified using the Bedtools suite version 2.16.1 [71] and eliminated. Moreover, for F2
deletions, some "inter-approach" redundancy was also expected and eliminated using intersectBed utility
also from the Bedtools suite. When redundancy was found, PInDel calls were preferred to BreakDancer
calls, because they had precise breakpoints and contained the calls for PH207 and C103. The same filter
was applied to all insertions that could be anchored to the B73 genome sequence. Furthermore, for non-
anchored InDels, in order to avoid redundancy in internal genotyping probe design, RepeatMasker
(http://www.repeatmasker.org) was used to mask redundant regions by similarity using an iterative
approach. First, "ALLPATHs-LG assembly" F2 insertions were masked with "ABySS assembly" F2 insertions
(at least 95% of identity) to generate a non-redundant set of F2 insertions. Then C103 insertions were
masked with F2 insertions (at 90% of identity), PH207 insertions were masked with C103 and F2 insertions
(90%), and finally F2 no map specific sequences were masked with PH207, C103, and F2 insertions (90%).

Design of Affymetrix® Axiom® array

Preparation of sequences for probes for design

To identify presence/absence regions (PARs) within InDel sequences suitable for the design of "off-target" probes, we used the genometools Tallymer utility [76] version 1.5.6 to create two indexes for B73, F2, PH207, and C103: one from their genome assemblies (17-mers with a minimal occurrence of 1) and one from a 5x genome equivalent subset of their raw sequenced data (17-mers with a minimal occurrence of 5). Then B73 genome was iteratively annotated with the script tallymer2gff3.plx (options used: -k 17 -min 35 -occ 1|5 depending on the index) to identify regions not covered by F2, PH207, and C103 kmers. Reciprocally, the two F2 draft genomes, PH207 and C103 ALLPATHs-LG draft genomes were run through the same procedure to identify regions not covered by B73 kmers. The gff files generated by this process were then used in combination with gff files of repeats annotated with RepeatMasker to define PARs of a minimum size of 35bp for each type of InDel and each draft genome.

BP preparation

Breakpoints could be targeted by probes (Figure 1A) provided that the nucleotide flanking the breakpoint at the beginning of the deleted sequence was different from the nucleotide right after the end of deleted sequence (and reciprocally on the reverse strand). Type I and type III breakpoints without micro-homology sequence can be submitted for the Affymetrix®' standard design procedure, whereas type II breakpoints have to go through an iterative design process, shifting the sequence by one base on each attempt until reaching a discriminative position. This iterative process stops after 5bp and is also performed by Affymetrix®.

Probes scoring

All potential probes were evaluated in an *in-silico* analysis to predict their microarray performance. A p-convert value, which arises from a random forest model intended to predict the probability that the SNP will convert on the array, was determined for all probes. The model considers factors including probe sequence, binding energies, and the expected degree of non-specific binding and hybridization to multiple genomic regions. This degree of non-specific binding is estimated calculating 16-mer hit counts, which is the number of times all 16 bp sequences in the 30 bp flanking region from either side of the SNP have a matched sequence in the genome. These scores were generated both for forward and reverse probes. A probeset is recommended if p-convert>=0.6 and there are no expected polymorphisms in the flanking region. A probeset is neutral if p-convert>=0.4, the number of expected polymorphisms in the flanking region is less than 3, and the polymorphisms are further than 21 bp of the variant of interest. Probesets not falling into these two categories are scored as *not recommended*. Probesets that cannot be designed are scored as *not possible*.

Probes selection

Concerning OTV and MONO probes, we applied three successive filtering steps. First, we selected only probes classified as recommended and neutral based on their scoring, with no more than one hit on the B73 reference genome for deletion probes, and no hit at all for insertion probes. After this step, 204,213 OTV probes and 18,884,827 MONO probes remained. Secondly, only probes with more than 70%

in PARs were kept. An additional filtering step was implemented specifically for MONO probes to optimize
probe distribution along the targeted PARs. For this step, PARs were split in 75bp windows using
windowmaker (Bedtools) and the MONO probe with the highest p-convert value was selected for each
window. If there were InDels with less than 4 MONO probes selected using 75bp windows, these probes
were eliminated and a second iteration was attempted, using 50bp windows, followed by a last iteration
with 25bp windows. This generated 616,286 probes including BP and OTV probes targeting 108,24 InDels
(90% of InDel selected for design). We completed the design by rescuing 6,219 OTV and 53,441 MONO
probes from InDels or PARs not targeted by any probes, bringing the total number of probes selected to
675,946 to target 109,292 InDel.
At the last step, duplicated probeset were removed based on their sequence by Affymetrix® during the
chip design procedure, leaving 662,772 probeset (105,927 InDels) corresponding to 1,404,570 different
probes to be tiled on the array.

Genotyping of 105k InDels on 480 maize DNA samples

Plant material for genotyping

For genotyping, 480 different DNA samples were extracted from leaves following a NaBisulfite
method modified from [77, 78]. These 480 samples included 440 inbred lines, 24 highly recombinant
inbred lines, and 16 F1 hybrids. Both F1 hybrids (obtained by crossing inbred lines) and their parental
inbred lines were genotyped on the array, but seed lots used to produce F1 hybrids and those used to
extract DNA for genotyping were different. Among these 480 DNAs, 13 inbred lines were genotyped using
two different DNAs from two different seed-lots and were used to evaluate the reproducibility of the
genotyping (Additional file 1: Table S9). DNA samples of one F1 hybrid were also genotyped 6 times.
Mendelian inheritance was estimated between 12 hybrids F1 derived from 9 different parental lines
(Additional file 1: Table S8)

Variant calling using Affymetrix® algorithm

Each type of probe had a dedicated algorithm (Additional file 2: Figure S7) to call genotypes,
according to expected behavior from the probe design. DNA samples from 480 individuals were hybridized
to the array using the Affymetrix® system. The genotyping, sample QC, and marker filtering were
performed according to the Axiom® Best Practice genotyping analysis workflow. Genotype calls and
classifications were generated from the hybridization signals in the form of CEL files using the Affymetrix®
Power Tools (APT) and the SNPolisher package for R, according to the Axiom® Genotyping Solution Data
Analysis Guide, and a custom-made R script, Hom2OTV, implemented the algorithm for calling MONO
probes.

The APT results were then post-processed using SNPolisher, which is an R package specifically
designed by Affymetrix®. Marker metrics were generated using the *Ps_Metrics* function. These marker QC
metrics were used to classify probesets into 14 categories (Additional file 2: Figure S8) using the
Ps_Classification and *Ps_Classification_Supplemental* functions, with all default setting for diploid (e.g.
HetSO.cut=-0.3, HetvMAF.cut=1.9), except for an empirically determined, more stringent heterozygous
variance filter (AB.varY.Z.cut=2.6). Example of clusters from each classification were visualized using the

Ps_Visualization function (Additional file 2: Figure S8). Variants were preferentially selected as
recommended if they were exhibiting stable category assignments with clearly separated clusters. Each
variant was ranked into a category (Additional file 2: Figure S8) at each step of the pipeline.

Algorithms used to convert BP and OTV were similar, as BP and OTV probes behaved like classical
SNPs. For initial genotype calling, a priori (generic) cluster positions were used, since no information about
expected positions was available. A first analysis was performed according to Affymetrix®
recommendations. Secondly, the level of inbreeding was taking into account for a posteriori cluster
definition, because of the high amount of inbred lines in the panel. This parameter took values from 0 for
fully heterozygous to 16 for completely homozygous samples. For OTV and BP algorithms, an inbred
penalty of 4 (lower penalty for inbred species) was applied to try to re-labelled probes that fall into
categories: CallRateBelowThreshold (CRBT), HomHomResolution (HHR), NoMinorHom (NMH), Other and
UnexpectedHeterozygosity, after the first cluster analysis (Additional file 2: Figure S8). Markers that were
classified as OTV may also be considered recommended after the *OTV_caller* function has been used to
re-label the genotype calls. The SNPolisher *OTV_Caller* function performed post-processing analysis to
identify miscalled AB clustering and identify which samples should be in the OTV cluster and which
samples should remain in the AA, AB, or BB clusters. Samples in the OTV cluster were re-labelled as OTV.
Finally, the recommended markers list is created by combining the list of markers that are classified into
the recommended categories (PolyHighResolution (PHR), MonoHighResolution (MHR), and OTV).

BP and OTV probes that exhibited only two clusters (AA or BB and OTV) should fall into the
monomorphic classification and be considered as not recommended. A new MONO algorithm was
developed (Figure 4), because, unlike traditional SNP genotyping, we only expected two clusters for
MONO probes (presence and absence) (Figure 1C). To classify monomorphic sequence genotyping, the
OTV_Caller function was called, and only MHR and NMH were considered as recommended. Other
monomorphic probes are then analyzed with an inbred penalty of 16 (highest level) to re-labelled probes
displaying higher-than-expected levels of heterozygosity. Finally, the new function called *Hom2OTV* was
implemented to classified probes exhibiting two homozygous clusters, with primarily an intensity
difference. This function determined if the intensity difference represents one homozygous cluster (InDel
presence) and one OTV cluster (InDel absence), as we expected. There are no parameters in this function.
The lower intensity homozygous cluster is recalled as OTV.

Evaluation of genotyping quality

We compared the genotyping for 479,027 probes from the InDel array (Genotyping By Array: GBA) with
the genotyping from sequencing (Genotyping By Sequencing: GBS) of 4 inbred lines used to discover the
InDels: B73, F2, PH207, and C103. Genotyping by sequencing was built from the alignment of probe
sequences on the reference genome B73 and the de novo assembly of 3 inbred lines (F2, PH207, and
C103) with Blast software. Sequences were considered present in lines when the probes were aligned
with less than 5% of mismatch or otherwise considered absent.

Genotyping consistency for B73, F2, PH207, and C103 was calculated between GBS and GBA according to
genotype calls "present" or "absent", produced by GBS (Table 4). For this purpose, Affymetrix® genotyping
was converted into these genotypes: present, absent, and hemizygote (1 copy present). Consistency of

Presence/Absence genotypes between sequencing and array genotyping was analyzed for four individuals
(B73, F2, PH207, C103) according to probe types (BP, OTV, MONO): Number of similar genotypes between
GBS and GBA /number of genotype called by GBA and GBS. Note that the seed-lot used for B73 and F2
genotyping is different from the seed-lot used for InDel discovery, but it is the same seed-lot for inbred
lines PH207 and C103.

In order to evaluate the consistency of probe genotyping within InDels (Figure 4), we used 362 inbred
lines from an association panel representing a wide range of genetic diversity (Camus-Kulandaivelu, 2005;
Bouchet et al., 2013). From 479,027 probes, we selected 294,650 polymorphic probes and fully consistent
between GBS and GBA in order to limit the genotyping errors due to sequencing. These probes genotyped
72,555 InDels. We then selected 50,648 polymorphic InDels that are genotyped with at least two probes
(corresponding to 270,581 probes), and calculated the average frequency of the presence allele across all
probes for each InDel and inbred line. For each InDel, we calculated the frequency of inbred lines
displaying fully consistent genotypes between probes, *i.e* the proportion of lines where the average
frequency across all probes is 0 or 1. We also calculated frequency of inbred lines that have a least one
probe with an inconsistent genotype (FreqDiff01), *i.e* the proportion of lines where the average frequency
across all probes is not 0 or 1. To evaluate the effect of the probe numbers on the frequency of lines
inconsistent within InDels, we analyzed the variation of frequency of lines not fully consistent (FreqDiff01)
with relation to the number of probes within the InDels, by estimating median and average FreqDiff01 for
each probe count (Figure 4B, Additional file 1: Table S5). To estimate the probe genotyping error rate, we
compared this variation to what we could expect for different genotyping error rates (1, 3, 5, and 10%) in
362 lines, genotyped by 10,000 Indels, with the number of probes varying from 2 to 50, using a binomial
sampling (Additional file 1: Table S6). For this, we simulated a number of false genotypes among the
probes for each InDel and each line using the rbinom function in R, with the following parameters:
Number of observation = 362 lines x 10,000 Indels; Number of trials for each observation = Number of
probes; Probability of success of each trial = probes genotyping error rate. Using this simulation, we
estimated frequency of inconsistent calls among 362,000 simulated genotypes (FreqDiff01) for each
probes count, varying from 2 to 50, and compared them with the median and average FreqDiff01 (Figure
4). To evaluate the impact of combining multiple probes for a genotype to correct genotype errors, we
used this simulation to estimate the InDel genotyping error rate, if we assign, to an inbred line, the most
frequent allele, based on the average allelic frequency of presence (Additional file 1: Table S6). To
compare accuracy for genotyping absence and presence using this array, we separated the InDels in four
classes, according to their average allelic frequency of absence in 362 inbred lines (0-25%, 25%-50%, 50%-
75%, 75%-100%) and compared their median FreqDiff01 (Additional file 2: Figure S12).

To evaluate the reproducibility of the 479,027 probes on the array, we compared the genotypes between
6 DNA replicates from F1 hybrids that originated from crossing B73 and F72. We also compared the
genotypes of 13 duplicated inbred lines (A554, A632, A654, B73, C103, CO255, D105, EP1, F2, F252, KUI3,
Oh43, and W117) that originated from different seed sources (Additional file 1: Table S9). The genotypes
of these 13 duplicated lines were also compared using 43,982 SNPs from the Illumina 50K SNP array.

To evaluate the quality of genotyping, we also analyzed 12 F1 hybrids derived from 9 parental inbred lines
Additional file 1: Table S8). We first predicted the genotypes of the 12 F1 hybrids, based on the genotyping

of their 2 parental lines, for 46,382 BPs probes, removing OTV calls. These predicted genotypes were then
compared with the observed genotypes of the corresponding hybrids: Number of similar genotypes
(homozygous or hemizygous) between predicted and observed/Number total of genotypes. BP probes
producing missing data or displaying hemizygous genotypes in the parental lies were excluded from the
comparison. Note that the seed-lots of the parental inbred lines genotyped may have been different from
the seed-lots used for producing the F1 hybrids.

Diversity analysis

We performed diversity analysis on 362 inbred lines from an association panel representing a
wide range of diversity [55, 57], obtained using InDels genotyped on our InDels Affymetrix® Axiom® array
and using SNPs genotyped from the Illumina 50K SNP array [54]. The genotypes of InDels were treated as
bi-allelic "present" and "absent".

To perform diversity analysis, we first selected 237,629 probes among the 479,027 probes for
which (i) the clustering observed were consistent with expectation (Table 3) and (ii) for which genotypes
produced by our array for the 4 lines used for discovering the InDels were fully consistent with the
genotyping, based on the alignment of the probes on the genome assemblies using BLAST software. We
filtered out 219,068 probes based on their genotyping quality (missing data rate below 20%, heterozygous
rate below 15% and minor allele frequency above 5%). In the end, we selected a single, best probe for
each InDel, leading to a set of 57,824 probes genotyping 57,824 InDels to analyze diversity in 362 inbred
lines.

We estimated two kinship matrices between 362 lines using "identity by descent" estimators (IBD)
based on 57,824 InDels and on 28,143 prefixed Panzea SNPs from the Illumina 50K (Figure 5). Kinship
matrices were estimated with the "ibd" function in the R package GenABEL [79]. We performed
correlation between IBD values estimated with SNP and InDel polymorphisms. Genetic structuration was
estimated using only the 28,143 Panzea SNPs with admixture software [80]. We selected the admixture
results for five genetic groups (Q=5), since it corresponded to the number of genetics groups defined in
previous studies using the Panzea SNPs from the Illumina 50K [55]. Lines were assigned to one genetic
group, given that the probability of assignment to the groups was greater than 0.6, whereas lines below
this threshold were considered "admixed". In order to compare genetic structuration based on InDels and
SNPs, we performed Principal Coordinate Analysis (PcoA) on genetic distance between lines with (362
lines) and without F2 and B73 (360 lines) based on their dissimilarity (1-IBD) using InDels. Each line was
plotted on the first two planes of PcoA and colored according to the assignment to the 5 genetics groups
(Figure 5).

1008

List of abbreviations

1009

GBA = Genotyping by array

GBS = Genotyping by sequencing

SNP = Single nucleotide polymorphism

InDel = Insertion / Deletion

BP = Breakpoint

MONO = Monomorphic

OTV = Off Target Variant

QC = Quality control

PHR = Poly High Resolyion

VCF = Variant Call Format

PAR = Presence / Absence Region

PAV = Presence / Absence Variant

SV = Structural variant

CNV = Copy Number Variant

TE = Transposable Element

CGH = Comparative Genomic Hybridization

NGS = Next Generation Sequencing

FW = Forward

REV = Reverse

NAM = Nested Association Mapping

DNA = Deoxyribonucleic Acid

PCR = Polymerase Chain Reaction

PcoA = Principal Coordinate Analysis

Mbp = Millions of Base Pairs

bp = base pair

FreqDiff01 = Frequency of lines not fully consistent between probes within InDel

1036

Declarations

Ethics approval and consent to participate

Not applicable

Consent to publish

Not applicable

Availability of data and materials

The array content is available at https://doi.org/10.15454/DWB4UT

Competing interests

Ali Pirani is an employee of Thermo Fisher Scientific (formerly Affymetrix®).

Funding

This work was supported by the project CNV-MAIZE (ANR-10-GENM-003) and the project Investement for the future AMAIZING ANR-10-BTBR-01 (ANR-PIA AMAIZING) and France Agrimer. PhD student C. Mabire is jointly funded by the program CNV4sel in the framework of metaprogram Selgen and by the Plant Biology and Breeding department of the French National Institute for Agricultural Research (INRA).

Authors' contributions

SDN designed and supervised the study and conducted CNVMaize project. CM, JD, and SDN drafted and corrected the manuscript, CV and JJ corrected the manuscript. NR, SDN, JPP, and SP conceived the array, AP, SDN, JJ and JD designed the array. AP developed the Affymetrix® genotype calling software and preformed the InDel genotype calling. JPP, JJ, and CV contributed to the sequencing; JD, AD, HR, and JJ performed the InDel discovery, JD and JJ build genome assemblies, JJ and AD discovered SNP within the InDels. CM evaluated the quality of genotyping and conducted the genetic diversity analysis.DM and VC did DNA extraction and prepared the samples for arrays genotyping. All authors have read and approved the manuscript.

Acknowledgements

We are very grateful to Patrick Schnable and Cheng-Ting "Eddy" Yeh for providing a subset of Presence/Absent Variants coming from their RNAseq and Sequence capture approach and for his helpful

1063 discussion. We are very grateful to Alain Charcosset for their contribution to the choice of inbred lines

1064 genotyped by the array and for his helpful discussion and comments on the manuscript.

References

1. Anderson JE, Kantar MB, Kono TY, Fu F, Stec AO, Song Q, et al. A Roadmap for Functional Structural

Variants in the Soybean Genome. G3amp58 GenesGenomesGenetics. 2014;4:1307–18.

2. Beló A, Beatty MaryK, Hondred D, Fengler KevinA, Li B, Rafalski A. Allelic genome structural variations

in maize detected by array comparative genome hybridization. Theor Appl Genet. 2010;120:355–67.

3. Cao J, Schneeberger K, Ossowski S, Günther T, Bender S, Fitz J, et al. Whole-genome sequencing of

multiple Arabidopsis thaliana populations. Nat Genet. 2011;43:956–63.

4. Liu J, Qu J, Yang C, Tang D, Li J, Lan H, et al. Development of genome-wide insertion and deletion

markers for maize, based on next-generation sequencing data. BMC Genomics. 2015;16:601.

5. Owens GL, Baute GJ, Hubner S, Rieseberg LH. Genomic sequence and copy number evolution during

hybrid crop development in sunflowers. Evol Appl. 2019;12:54–65.

6. Saintenac C, Jiang D, Akhunov ED. Targeted analysis of nucleotide and copy number variation by exon

capture in allotetraploid wheat genome. Genome Biol. 2011;12:R88.

7. Saxena RK, Edwards D, Varshney RK. Structural variations in plant genomes. Brief Funct Genomics.

2014;13:296–307.

8. Springer NM, Ying K, Fu Y, Ji T, Yeh C-T, Jia Y, et al. Maize Inbreds Exhibit High Levels of Copy Number

Variation (CNV) and Presence/Absence Variation (PAV) in Genome Content. PLoS Genet.

2009;5:e1000734.

9. Swanson-Wagner RA, Eichten SR, Kumari S, Tiffin P, Stein JC, Ware D, et al. Pervasive gene content

variation and copy number variation in maize and its undomesticated progenitor. Genome Res.

2010;20:1689–99.

10. Chia J-M, Song C, Bradbury PJ, Costich D, de Leon N, Doebley J, et al. Maize HapMap2 identifies

extant variation from a genome in flux. Nat Genet. 2012;44:803–7.

11. Mace ES, Tai S, Gilding EK, Li Y, Prentis PJ, Bian L, et al. Whole-genome sequencing reveals untapped

genetic potential in Africa's indigenous cereal crop sorghum. Nat Commun. 2013;4:2320.

12. Hirsch CN, Foerster JM, Johnson JM, Sekhon RS, Muttoni G, Vaillancourt B, et al. Insights into the

Maize Pan-Genome and Pan-Transcriptome. Plant Cell. 2014;26:121–35.

13. Lu F, Romay MC, Glaubitz JC, Bradbury PJ, Elshire RJ, Wang T, et al. High-resolution genetic mapping

of maize pan-genome sequence anchors. Nat Commun. 2015;6.

http://dx.doi.org/10.1038/ncomms7914.

14. Montenegro JD, Golicz AA, Bayer PE, Hurgobin B, Lee H, Chan C-KK, et al. The pangenome of
hexaploid bread wheat. Plant J. 2017;90:1007–13.

15. Zhao Q, Feng Q, Lu H, Li Y, Wang A, Tian Q, et al. Pan-genome analysis highlights the extent of
genomic variation in cultivated and wild rice. Nat Genet. 2018;50:278–84.

16. Hardigan MA, Crisovan E, Hamilton JP, Kim J, Laimbeer P, Leisner CP, et al. Genome Reduction
Uncovers a Large Dispensable Genome and Adaptive Role for Copy Number Variation in Asexually
Propagated *Solanum tuberosum*. Plant Cell. 2016;28:388–405.

17. Varshney RK, Saxena RK, Upadhyaya HD, Khan AW, Yu Y, Kim C, et al. Whole-genome resequencing
of 292 pigeonpea accessions identifies genomic regions associated with domestication and agronomic
traits. Nat Genet. 2017;49:1082–8.

18. Belo A, Zheng P, Luck S, Shen B, Meyer DJ, Li B, et al. Whole genome scan detects an allelic variant of
fad2 associated with increased oleic acid levels in maize. Mol Genet Genomics. 2008;279:1–10.

19. Hirsch CN, Hirsch CD, Brohammer AB, Bowman MJ, Soifer I, Barad O, et al. Draft Assembly of Elite
Inbred Line PH207 Provides Insights into Genomic and Transcriptome Diversity in Maize. Plant Cell.
2016;28:2700–14.

20. Darracq A, Vitte C, Nicolas S, Duarte J, Pichon J-P, Mary-Huard T, et al. Sequence analysis of
European maize inbred line F2 provides new insights into molecular and chromosomal characteristics of
presence/absence variants. BMC Genomics. 2018;19:119.

21. Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, et al. Improved maize reference genome with
single-molecule technologies. Nature. 2017. doi:10.1038/nature22971.

22. Pinosio S, Giacomello S, Faivre-Rampant P, Taylor G, Jorge V, Le Paslier MC, et al. Characterization of
the Poplar Pan-Genome by Genome-Wide Identification of Structural Variation. Mol Biol Evol.
2016;33:2706–19.

23. Appels R, Eversole K, Stein N, Feuillet C, Keller B, Rogers J, et al. Shifting the limits in wheat research
and breeding using a fully annotated reference genome. Science. 2018;361:eaar7191.

24. Sun S, Zhou Y, Chen J, Shi J, Zhao H, Zhao H, et al. Extensive intraspecific gene order and gene
structural variations between Mo17 and other maize genomes. Nat Genet. 2018;50:1289–95.

25. Zhou P, Silverstein KAT, Ramaraj T, Guhlin J, Denny R, Liu J, et al. Exploring structural variation and
gene family architecture with De Novo assemblies of 15 Medicago genomes. BMC Genomics.
2017;18:261.

26. Fu H, Dooner HK. Intraspecific violation of genetic colinearity and its implications in maize. Proc Natl
Acad Sci U S A. 2002;99:9573–8.

27. Brunner S, Fengler K, Morgante M, Tingey S, Rafalski A. Evolution of DNA sequence nonhomologies
among maize inbreds. Plant Cell Online. 2005;17:343.

28. Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, et al. The B73 Maize Genome:
Complexity, Diversity, and Dynamics. Science. 2009;326:1112–5.

29. Pinkel D, Segraves R, Sudar D, Clark S, Poole I, Kowbel D, et al. High resolution analysis of DNA copy
number variation using comparative genomic hybridization to microarrays. Nat Genet. 1998;20:207–11.

30. Springer NM, Ying K, Fu Y, Ji T, Yeh C-T, Jia Y, et al. Maize Inbreds Exhibit High Levels of Copy Number
Variation (CNV) and Presence/Absence Variation (PAV) in Genome Content. PLoS Genet.
2009;5:e1000734.

31. Hirsch CN, Hirsch CD, Brohammer AB, Bowman MJ, Soifer I, Barad O, et al. Draft Assembly of Elite
Inbred Line PH207 Provides Insights into Genomic and Transcriptome Diversity in Maize. Plant Cell.
2016;28:2700–14.

32. Cooper GM, Zerr T, Kidd JM, Eichler EE, Nickerson DA. Systematic assessment of copy number
variant detection via genome-wide SNP genotyping. Nat Genet. 2008;40:1199–203.

33. Dellinger AE, Saw S-M, Goh LK, Seielstad M, Young TL, Li Y-J. Comparative analyses of seven
algorithms for copy number variant identification from single nucleotide polymorphism arrays. Nucleic
Acids Res. 2010;38:e105–e105.

34. Wang X, Lebarbier E, Aubert J, Robin S. Variational Inference for Coupled Hidden Markov Models
Applied to the Joint Detection of Copy Number Variations. Int J Biostat. 2019;15. doi:10/gf7323.

35. Wang K, Li M, Hadley D, Liu R, Glessner J, Grant SFA, et al. PennCNV: An integrated hidden Markov
model designed for high-resolution copy number variation detection in whole-genome SNP genotyping
data. Genome Res. 2007;17:1665–74.

36. Didion JP, Yang H, Sheppard K, Fu C-P, McMillan L, de Villena F, et al. Discovery of novel variants in
genotyping arrays improves genotype retention and reduces ascertainment bias. BMC Genomics.
2012;13:34.

37. Unterseer S, Bauer E, Haberer G, Seidel M, Knaak C, Ouzunova M, et al. A powerful tool for genome
analysis in maize: development and evaluation of the high density 600k SNP genotyping array. BMC
Genomics. 2014;15:823.

38. Hupé P, Stransky N, Thiery J, Radvanyi F, Barillot E. Analysis of array CGH data: from signal ratio to
gain and loss of DNA regions. Bioinformatics. 2004.

39. Olshen AB, Venkatraman ES, Lucito R, Wigler M. Circular binary segmentation for the analysis of
array-based DNA copy number data. Biostatistics. 2004;5:557–72.

40. Picard F, Robin S, Lavielle M, Vaisse C, Daudin J. A statistical approach for array CGH data analysis.
BMC Bioinformatics. 2005;6:27.

41. Picard F, Robin S, Lebarbier É, Daudin J. A segmentation/clustering model for the analysis of array
CGH data. Biometrics. 2007;63:758–66.

42. Marioni J, Thorne N, Tavare S. BioHMM: a heterogeneous hidden Markov model for segmenting
array CGH data. Bioinformatics. 2006;22:1144.

43. Stjernqvist S, Ryden T, Skold M, Staaf J. Continuous-index hidden Markov modelling of array CGH
copy number data. Bioinformatics. 2007;23:1006.

44. Lyra DH, Galli G, Alves FC, Granato ÍSC, Vidotti MS, Bandeira e Sousa M, et al. Modeling copy number
variation in the genomic prediction of maize hybrids. Theor Appl Genet. 2018. doi:10.1007/s00122-018-
3215-2.

45. Alkan C, Coe BP, Eichler EE. Genome structural variation discovery and genotyping. Nat Rev Genet.
2011;12:363–376.

46. Chen K, Wallis JW, McLellan MD, Larson DE, Kalicki JM, Pohl CS, et al. BreakDancer: an algorithm for
high-resolution mapping of genomic structural variation. Nat Methods. 2009;6:677–81.

47. Kidd JM, Cooper GM, Donahue WF, Hayden HS, Sampas N, Graves T, et al. Mapping and sequencing
of structural variation from eight human genomes. Nature. 2008;453:56–64.

48. Korbel JO, Urban AE, Affourtit JP, Godwin B, Grubert F, Simons JF, et al. Paired-End Mapping Reveals
Extensive Structural Variation in the Human Genome. Science. 2007;318:420–6.

49. Tuzun E, Sharp AJ, Bailey JA, Kaul R, Morrison VA, Pertz LM, et al. Fine-scale structural variation of
the human genome. Nat Genet. 2005;37:727–32.

50. Ye K, Schulz MH, Long Q, Apweiler R, Ning Z. Pindel: a pattern growth approach to detect break
points of large deletions and medium sized insertions from paired-end short reads. Bioinformatics.
2009;25:2865–71.

51. Lai J, Li R, Xu X, Jin W, Xu M, Zhao H, et al. Genome-wide patterns of genetic variation among elite
maize inbred lines. Nat Genet. 2010;42:1027–30.

52. The Danish Pan-Genome Consortium, Sibbesen JA, Maretty L, Krogh A. Accurate genotyping across
variant classes and lengths using variant graphs. Nat Genet. 2018;50:1054–9.

53. Muñoz-Amatriaín M, Eichten SR, Wicker T, Richmond TA, Mascher M, Steuernagel B, et al.
Distribution, functional impact, and origin mechanisms of copy number variation in the barley genome.
Genome Biol. 2013;14:R58.

54. Ganal MW, Durstewitz G, Polley A, Bérard A, Buckler ES, Charcosset A, et al. A Large Maize (Zea mays
L.) SNP Genotyping Array: Development and Germplasm Genotyping, and Genetic Mapping to Compare
with the B73 Reference Genome. PLoS ONE. 2011;6:e28334.

55. Bouchet S, Servin B, Bertin P, Madur D, Combes V, Dumas F, et al. Adaptation of maize to temperate
climates: mid-density genome-wide association genetics and diversity patterns reveal key genomic
regions, with a major contribution of the Vgt2 (ZCN8) locus. PloS One. 2013;8:e71377.

56. Bouchet S, Bertin P, Presterl T, Jamin P, Coubriche D, Gouesnard B, et al. Association mapping for
phenology and plant architecture in maize shows higher power for developmental traits compared with

growth influenced traits. Heredity. 2016.
https://www.nature.com/hdy/journal/vaop/ncurrent/full/hdy201688a.html. Accessed 21 Jun 2017.

57. Camus-Kulandaivelu L. Maize Adaptation to Temperate Climate: Relationship Between Population
Structure and Polymorphism in the Dwarf8 Gene. Genetics. 2006;172:2449–63.

58. Gabur I, Chawla HS, Snowdon RJ, Parkin IAP. Connecting genome structural variation with complex
traits in crop plants. Theor Appl Genet. 2019;132:733–50.

59. Feschotte C, Jiang N, Wessler SR. PLANT TRANSPOSABLE ELEMENTS: WHERE GENETICS MEETS
GENOMICS. Nat Rev Genet. 2002;3:329–41.

60. Morgante M, De Paoli E, Radovic S. Transposable elements and the plant pan-genomes. Curr Opin
Plant Biol. 2007;10:149–55.

61. Ducrocq S, Madur D, Veyrieras JB, Camus-Kulandaivelu L, Kloiber-Maitz M, Presterl T, et al. Key
impact of Vgt1 on flowering time adaptation in maize: evidence from association mapping and
ecogeographical information. Genetics. 2008;178:2433–7.

62. Salvi S, Sponza G, Morgante M, Tomes D, Niu X, Fengler KA, et al. Conserved noncoding genomic
sequences associated with a flowering-time quantitative trait locus in maize. Proc Natl Acad Sci U A.
2007;104:11376–81.

63. Salvi S, Tuberosa R, Chiapparino E, Maccaferri M, Veillet S, van Beuningen L, et al. Toward positional
cloning of Vgt1, a QTL controlling the transition from the vegetative to the reproductive phase in maize.
Plant Mol Biol. 2002;48:601–613.

64. Clark AG, Hubisz MJ, Bustamante CD, Williamson SH, Nielsen R. Ascertainment bias in studies of
human genome-wide polymorphism. Genome Res. 2005;15:1496–1502.

65. Gouesnard B, Negro S, Laffray A, Glaubitz J, Melchinger A, Revilla P, et al. Genotyping-by-sequencing
highlights original diversity patterns within a European collection of 1191 maize flint lines, as compared
to the maize USDA genebank. Theor Appl Genet. 2017. doi:10.1007/s00122-017-2949-6.

66. Gore MA, Chia JM, Elshire RJ, Sun Q, Ersoz ES, Hurwitz BL, et al. A first-generation haplotype map of
maize. Sci Wash. 2009;326:1115–7.

67. Brandenburg J-T, Mary-Huard T, Rigaill G, Hearne SJ, Corti H, Joets J, et al. Independent introductions
and admixtures have contributed to adaptation of European maize and its American counterparts. PLOS
Genet. 2017;13:e1006666.

68. Yeaman S. Genomic rearrangements and the evolution of clusters of locally adaptive loci. Proc Natl
Acad Sci. 2013;110:E1743–51.

69. Gu W, Zhang F, Lupski JR. Mechanisms for human genomic rearrangements. PathoGenetics.
2008;1:4.

70. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics. 2009;25:2078–9.

71. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26:841–2.

72. Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, et al. High-quality draft
assemblies of mammalian genomes from massively parallel sequence data. Proc Natl Acad Sci.
2011;108:1513–8.

73. Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, et al. BUSCO
Applications from Quality Assessments to Gene Prediction and Phylogenomics. Mol Biol Evol.
2018;35:543–8.

74. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol.
1990;215:403–10.

75. Abyzov A, Gerstein M. AGE: defining breakpoints of genomic structural variants at single-nucleotide
resolution, through optimal alignments with gap excision. Bioinformatics. 2011;27:595–603.

76. Gremme G, Steinbiss S, Kurtz S. GenomeTools: A Comprehensive Software Library for Efficient
Processing of Structured Genome Annotations. IEEE/ACM Trans Comput Biol Bioinform. 2013;10:645–
56.

77. Tai TH, Tanksley SD. A rapid and inexpensive method for isolation of total DNA from dehydrated
plant tissue. Plant Mol Biol Report. 1990;8:297–303.

78. Dellaporta SL, Wood J, Hicks JB. A plant DNA minipreparation: Version II. Plant Mol Biol Report.
1983;1:19–21.

79. Aulchenko Y. GenABEL: an R package for Genome Wide Association Analysis. 2009.

80. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated
individuals. Genome Res. 2009;19:1655–64.

1255

Figures

Figure 1: Genotyping of InDel CNVMAIZE_DEL_12661 using three probe types on 445 individuals. A)
Schematic distribution of the 9 probes along the sequence of InDel CNVMAIZE_DEL_12661 (green line)
and the bordering sequence common between all individuals (blue line) genotyped by the array. Double,
dotted, and full arrows represented the probes designing on the forward and reverse flanking sequences
of the breakpoint sites (BP), at not polymorphic (MONO) and polymorphic sites (OTV) within internal
sequence of InDel. B) Schematic distribution of the 8 probes passing Affymetrix® quality control and called
by the Affymetrix® pipeline C) Clustering produced by the Affymetrix® algorithm for an OTV, MONO, and
BP probe from InDel based on both fluorescence contrast (X axis) and intensity (Y axis) of the 445 inbred
lines. Red, blue and yellow dots indicated the presence of the sequence (genotype "present") either
homozygous for allele A (AA) or allele B (BB) or heterozygous (AB), respectively. Cyan and green indicated
that the sequence was absent in the individual (OO), or only in one copy of the sequence, e.g hemizygous
for presence/absence (OB or OA). Black dots indicated individuals for which no genotype could be
assigned (Missing data) D) Haplotypes displayed by the genotyping using 8 probes (column) on the 445
inbred lines (row). Colors corresponded to the genotype of individuals produced by clustering in C)

A) Probes on array

common region   CNVMAIZE_DEL_12661   common region

B) Probes passing QC

Quality control

C) Clustering

D) Haplotypes

Figure 2: Distribution of 105,927 InDels genotyped by the array according to their size and the cumulated
length of Presence/Absence regions (PARs) in their internal sequence. A) Distribution of the number of
InDels according to the proportion of presence/absence regions (sequence not present elsewhere in the
genome) within their internal sequence. B) Distribution of the number of InDels according to their size
(kbp) and the percentage of internal sequence of InDel covered by PAR(s). Red Color indicates the
proportion of InDels with (red) or without (blue) PARs for the 7 InDel size classes.

1279

1280

1281

Figure 3: Number of InDels interrogated by each probe types or their combination, for which: a probe
could be designed (A) and a probe was finally selected to be included in the final array (B). Vertical bars
indicate number of InDels interrogated by each probe types or their combination. Black points and
connected traits below the vertical bars indicate the corresponding probes types or their combination
that are used for interrogating this subset of InDels. Horizontal bars indicate number of InDels
interrogated by each probe types (OTV, BP, MONO). Number of InDels by probe type, for which: a probe
could be designed (A) and a probe was finally selected to be included in the final array (B). Number of
InDels that could be targeted by each type of probes designed (A) and selected to be included in the final
array (B).

1291

Figure 4: Consistencies among probes within 50,648 InDels with at least two probes genotyped in 362 inbred lines. (A) Distribution of the average allelic frequencies of present calls over all probes. (B) Variation of proportion of genotypes not fully consistent across all probes. The black and gray curves with triangle points represent the variation of the median and average FreqDiff01 across InDels, respectively. Colored curves with circle points represent the expected variation of the proportion for different error rates (1%: red, 3%: green, 5%: light blue, 10%: dark blue). Frequencies of 1 (presence) and 0 (absence) indicate that all probes had consistent genotypes for the corresponding inbred line. Intermediate frequencies (FreqDiff01) indicate that at least one probe was not consistent with the other probes for the same InDel in one inbred line.

1302 **Figure 5:** Principal coordinate analysis on the genetic distance between 362 inbred lines from an association panel estimated by A) 57,824 InDels
1303 and B) 28,143 SNPs. Colors represent the assignment of the inbred lines to the 5 genetic groups defined by admixture using pre-fixed Panzea SNPs
1304 from the 50K Illumina array, when the probability of assignment to a group (membership) was greater than 60%. Inbred lines not assigned to a
1305 group were considered admixed and colored gray. The common names of maize accessions, typical of each genetic group, were used.

1306

1307

1308

Additional Files

Additional file 1:

Table S1: Summary of sequencing data used during the assembly process provided by ALLPATHS-LG; **Table S2:** Classification by the Affymetrix® pipeline of 84,994 BP probes based on cluster number, separation, variance, and call rate. A) Probes recommended for genotyping, B) Probes not recommended for genotyping; **Table S3**: Classification by the Affymetrix® pipeline of 163,278 OTV probes based on cluster number, separation, variance, and call rate. A) Probes recommended for genotyping B) Probes not recommended probes for genotyping; **Table S4:** Classification by the Affymetrix® pipeline of 414,500 MONO probes, based on cluster number, separation, variance, and call rate. A) Probes recommended for genotyping B) Probes not recommended for genotyping; **Table S5:** Effect of probe number within InDels on average percentage of missing data, of genotypes absent and genotypes not fully concordant; **Table S6:** Simulation of genotyping error rates for 362 lines and 10,000 InDels called by various numbers of probes with a probe genotyping error rate ranging from 1% to 10%; **Table S7:** Comparison of reproducibility between 5 DNA replicates of hybrid F1 according to probes type and observed clustering; **Table S8:** Mendelian inheritance of 12 hybrids F1 derived from 9 different parental inbred lines for 46,382 BP probes passing Affymetrix quality control and polymorphic; **Table S9**: Comparison of the reproducibility of InDels and SNP genotyping between 13 maize varieties replicated on 50K Illumina SNP and Affymetrix® Axiom® InDel arrays

Additional file 2:

Figure S1: Description of two approaches used to discover InDels using resequencing data of DNA; **Figure S2:** Number and complementarity of A) deletions and B) insertions regarding B73 reference genome discovered between F2, PH207 and C103 inbred lines and B73; **Figure S3:** Schematic representation of four different breakpoint types identified by PINDEL at InDel breakpoints according to the presence of micro-homology sequence or not in place of the deleted sequence; **Figure S4:** Distribution of probe number per InDel for 105,927 InDels genotyped with the array; **Figure S5:** Relationship between probes number genotyping the InDel and A) the InDel length B) cumulated length of specific sequence (PARs) within InDel; **Figure S6:** Variation of probes and InDels density across the 10 maize chromosomes; **Figure S7:** Three dedicated Affymetrix pipelines used for calling InDel polymorphisms from the fluorescent intensity variation of BP probes (A), OTV probes (B) and MONO probes (C); **Figure S8:** Example of clustering based on probes fluorescence (intensity in y-axis and contrast in x-axis), for 14 different classifications of probes assigned by the Affymetrix® algorithm; **Figure S9:** Example of clustering for 6 randomly probes in different classifications; **Figure S10:** Variation of the distribution of the average consistency rate (%) of InDels between expected and observed genotyping of probes according to number of probes within the InDel; **Figure S11:** Haplotype of two InDels genotyped with multiple probes (in column) for 362 individuals (in rows); **Figure

1335 **S12:** Effect of average frequency of absence across 362 lines on consistencies between probes genotyping within InDels; **Figure S13:** Comparison

1336 of kinship between 362 inbred lines estimated with 57,824 InDels and with 28,143 SNPs from the 50K Illumina genotyping array; **Figure S14:**

1337 Principal coordinate analysis on the genetic distance between 360 inbred lines from an association panel (B73 and F2 were excluded) estimated

1338 by A) 57,824 InDels and B) 28,143 SNPs.