Premeiotic and meiotic failures lead to hybrid male sterility in the *Anopheles*

***gambiae* complex**

Running Head: Cellular basis of hybrid male sterility

Jiangtao Liang[1], Igor V. Sharakhov[1,2*]

[1]Department of Entomology, Virginia Polytechnic Institute and State University, Blacksburg,

Virginia, United States of America

[2]Department of Cytology and Genetics, Tomsk State University, Tomsk, Russian Federation

[*]**Corresponding author**

E-mail: igor@vt.edu

Abstract

Hybrid male sterility contributes to speciation by restricting gene flow between related taxa.

Detailed cytological characterizations of reproductive organs in hybrid males is important for

identifying phenotypes that can help guide searches of speciation genes. To investigate possible

cellular causes of hybrid male sterility, we performed crosses between closely related species of

the *Anopheles gambiae* complex: *An. merus* with *An. gambiae* or *An. coluzzii*. We demonstrate

that hybrid male sterility in African malaria mosquitoes involves two defects in the reciprocal

crosses: a premeiotic arrest of germline stem cells in degenerate testes and a failure of the

reductional meiotic division of primary spermatocytes in normal-like testes. The premeiotic

arrest in degenerate testes of hybrids is accompanied by a strong suppression of meiotic and

postmeiotic genes. Unlike pure species, sex chromosomes in normal-like testes of F1 hybrids are

largely unpaired during meiotic prophase I and all chromosomes show various degrees of

insufficient condensation. Instead of entering reductional division in meiosis I, primary

spermatocytes prematurely undergo an equational mitotic division producing nonmotile diploid

sperm. Thus, our study identified cytogenetic errors in interspecies hybrids that arise during the

early stages of postzygotic isolation.

Keywords: *Anopheles*, chromosome pairing, hybrid male sterility, meiosis, speciation,

spermatogenesis.

32   1. Introduction

When species diverge, hybrid offspring that are produced can suffer from reduced fitness.

Hybrid male fertility is usually one of the first of these postzygotic phenotypes affected [1, 2].

Therefore, the genetic factors, cellular basis, and molecular mechanisms of hybrid male sterility

(HMS) are of considerable interest, as they inform our understanding of both speciation and

normal fertility function. Studies in animals commonly assess testis shape, size, or weight, as

well as sperm morphology, density, or motility to define male sterility phenotypes. However, the

cellular basis of HMS is rarely investigated. Available cytological studies of spermatogenesis in

sterile hybrids indicate that multiple mechanisms of functional sterility are possible. Sterile

hybrids between *Mus musculus domesticus* and *M. m. musculus* have spermatogenic arrest in

early meiosis I with disrupted homoeologous chromosome pairing and meiotic sex chromosome

inactivation [3, 4]. Studies of failed spermatogenesis in different *Drosophila* hybrids observed

arrests at the premeiotic stage [5], reduced chromosome pairing, unequal chromosome

segregation in meiosis [6, 7], characteristic spermiogenic arrests [5], spermatid abnormalities [8],

and problems in sperm bundling and motility [9, 10]. Detailed analyses of cellular phenotypes in

testes of various hybrid organisms could help to establish the order of origin of postzygotic

isolating barriers and to guide the identification of speciation genes.

The *Anopheles gambiae* complex consists of at least nine morphologically nearly

indistinguishable sibling species of African malaria mosquitoes [11-13]. Genome-based

estimations of the age of the *An. gambiae* complex vary from 1.85 [14] to as young as 0.526

million years [15]. Genomic introgression is prevalent in autosomal regions of several species

indicating naturally occurring interspecies hybridization [14, 15]. Experimental crosses of

species from the *An. gambiae* complex often produce sterile F1 hybrid males, conforming to

Haldane's rule of sterility or inviability of the heterogametic sex [16-20]. Early crossing

experiments between members of the complex have found that HMS is associated with various

degrees of testes atrophy and underdevelopment of sperm [18]. A large effect of the X

chromosome on HMS has been demonstrated for crosses between *An. gambiae* and *An.*

arabiensis [16, 21]. However, an introgression of the Y chromosome from *An. gambiae* into the

background of *An. arabiensis* has shown no apparent influence on male fertility, fitness, or gene

expression [17]. Because of the recent evolution and ease of hybridization, sibling species of the

An. gambiae complex offer great opportunities to provide insights into the cellular basis and

molecular mechanisms of HMS.

Here, we investigate possible cellular causes of male sterility in hybrids between sibling species

of the *An. gambiae* complex. We asked three specific questions: (i) What cellular processes are

involved in causing infertility in hybrid mosquito males? (ii) Are the defects leading to HMS

premeiotic, meiotic, or postmeiotic? and (iii) What cytogenetic errors trigger spermatogenic

breakdown? We demonstrate that HMS in malaria mosquitoes involves two cellular defects in

reciprocal crosses: premeiotic arrest in germline stem cells and the failure of the reductional

meiotic division in primary spermatocytes. Our data suggest that meiotic abnormality in hybrid

males stems from the unpairing of the sex chromosomes and insufficient chromatin

condensation. Thus, our study identifies cytogenetic errors in hybrids that arise during the early

stages of postzygotic isolation.

2. Material and Methods

(a) Mosquito strains and crossing experiments

Laboratory colonies of *An. gambiae* ZANU (MRA-594), *An. coluzzii* MOPTI (MRA-763), *An.*

coluzzii MALI (MRA-860), and *An. merus* MAF (MRA-1156) were obtained from the

Biodefense and Emerging Infections Research Resources Repository (BEI). To perform

interspecies crosses, male and female pupae were separated to guarantee virginity of adult

mosquitoes. After the emergence of adults, crossing experiments were performed by combining

30 females and 15 males in one cage. At least two blood meals were fed to females and at least

three repeats of each cross were conducted. Male gonads were photographed using an Olympus

phase contrast microscope BX41 and UC90 digital camera (Olympus, Tokyo, Japan). We

observed the testes of 30 males and took pictures of the testes of five males from each cross. A

movie of sperm motility for five hybrid males from each cross and five males of pure species

was recorded using a UC90 digital camera (Olympus, Tokyo, Japan).

(b) Chromosome preparation and fluorescence *in situ* hybridization (FISH)

Preparations of chromosomes, DNA probe labeling, and FISH were performed as previously

described [22, 23]. Five DNA probes were used for FISH in this study (supplementary material,

table S1). Cytogenetic analyses were performed on at least 10 male individuals of each pure

species and of each hybrid. Lengths of well-spread metaphase I chromosomes of each species

and of each hybrid were measured using the ruler tool in Adobe Photoshop CS6 (Adobe Inc., San

Jose, CA, USA) (supplementary material, table S2). For whole-mount FISH and analysis of

chromosome pairing, testes from one-day-old adults of pure species and hybrids were dissected

in 1×PBS solution and fixed in 3.7% paraformaldehyde in 1×PBS with 0.1% tween-20 (PBST)

for 10 min at room temperature. After adding labeled DNA probes, testes were incubated at 75

℃ for 5 min (denaturation) and then at 37 ℃ overnight (hybridization). Testes from 6

individuals of pure species and of hybrids were scanned with a Zeiss LSM 880 confocal laser

scanning microscope (Carl Zeiss AG, Oberkochen, Germany) to analyze pairing and unpairing of

the sex chromosomes. A total of 418 and 489 nuclei at the early stages of meiotic prophase I

were analyzed in pure species and hybrids, respectively (supplementary material, table S3). The

percentage of cells with pairing and no pairing of sex chromosomes was used to compare

parental species and hybrids. Since the variances for both groups were equal ($s_{max}/s_{min} < 2$), a

statistical two-sample pooled *t*-test was performed using the JMP 13 software (SAS Institute

Inc., Cary, NC, USA). For more details, see the supplementary material and methods.

(c) RNA extraction and reverse transcription polymerase chain reaction (RT-PCR)

We dissected reproductive organs, which include both testes and male accessary glands (MAGs),

from 30 males, only testes or only MAGs from 20 males, and ovaries from 30 females. These

mosquitoes were 0-12-hour-old virgin adults of *An. coluzzii* MOPTI, *An. merus* MAF, and

interspecies hybrids from crosses ♀*An. coluzzii* MOPTI × ♂*An. merus* and ♀*An. merus* × ♂*An.*

coluzzii MOPTI. Total RNA was extracted using a Direct-Zol^{TM} RNA MiniPrep Kit (Zymo

Research, Irvine, California, US). Two-step RT-PCR was performed to analyze gene expression.

cDNA for selected genes (supplementary material, table S1) was generated in a 20-μl reaction

containing 2 μl of 10× RT buffer, 2 μl of 0.1-M dithiothreitol (DTT), 1 μl of 50 μM oligo(dT)$_{20}$

primer, 1 μl of 10 mM dNTP mix solution, 1 μl of 25mM MgCl$_2$, 1 μl of RNaseOUT^{TM}(40 U/μl),

1 μl of SuperScript^{TM} III RT (200 U/μl), 2 μl of RNA, and 6 μl of DEPC-treated water using the

SuperScript^{TM} III First-Strand Synthesis System for RT-PCR (Invitrogen, Carlsbad, CA, US).

Amplification products were visualized in a 2% agarose gel and photographed under the same

parameters for each gene.

3. Results

(a) HMS phenotypes at the cellular level

To obtain interspecies hybrids, we performed reciprocal crosses between *An. merus* MAF and

An. gambiae ZANU or *An. coluzzii* MOPTI and MALI. Backcrossing of F1 males to parental

females resulted in induction of the laying of eggs that did not hatch, which confirmed mating of

sterile F1 hybrid males since seminal fluids but not sperm are required to induce oviposition

[24]. The observed sterility of hybrid males agrees with Haldane's rule [20] for the majority of

interspecies crosses in the *An. gambiae* complex except for crosses between *An. gambiae* and *An.*

coluzzii, which produce fertile hybrids of both sexes [16-19, 25, 26]. To investigate the

developmental phenotypes involved in hybrid sterility, we dissected testes from adult males

obtained from interspecies crosses and from pure species. Normal testes of pure species have a

spindle-like shape (supplementary material, figure S1A). We found obvious differences between

testes morphology/size in one interspecies cross versus its reciprocal. Hybrid males from crosses

between female *An. merus* and male *An. gambiae* or *An. coluzzii* display normal-like testes

(supplementary material, figure S1B). In contrast, F1 males from crosses between female *An.*

gambiae or *An. coluzzii* and male *An. merus* show severely underdeveloped (degenerate) testes

(supplementary material, figure S1C). We then tested whether normal-like and degenerate testes

of interspecies hybrids produce any sperm. Sperm reach maturity within two days after

emergence of *Anopheles* adult males. In squashed testes of 2-5-day-old adults of pure species,

large amounts of mature spermatozoa with long tails can be seen. After we crushed the testes,

spermatozoa with vibrant motility escaped from the ruptures (supplementary material, figure

S1A, movie S1). However, mature sperm or sperm motility could hardly be seen in squashed or

crushed normal-like testes of 2-5-day-old adult hybrids from crosses when *An. merus* was the

mother. Instead, we see fewer spermatids and mostly nonmotile spermatozoa with large heads

and often two short tails growing from opposite ends of the head (supplementary material, figure

S1B, movie S2). Staining with DAPI identified extended premeiotic and spermatogenic stages

that cause delay of spermatogonia amplification, spermatocyte divisions, and spermatid

differentiation in the normal-like testes (supplementary material, figure S2). Only

undifferentiated round cells could be seen in degenerate testes of hybrids from reciprocal crosses

when *An. merus* was the father (supplementary material, figure S1C). Given the small size of the

degenerate testes, these round cells may represent germline stem cells. Thus, neither normal-like

nor underdeveloped testes of the interspecies hybrids produce mature motile spermatozoa.

(b) The progress of meiosis in males of pure species

Here, we provide the first description of chromosome behavior during meiosis of pure *Anopheles*

species. The chromosome complement of *Anopheles* males consists of three chromosome pairs:

two autosomes (2 and 3) and X and Y sex chromosomes. With the help of sex-chromosome-

specific fluorescent probes (supplementary material, table S1), we followed the progress of

meiosis in *An. gambiae*, *An. coluzzii*, and *An. merus*. Figure 1 shows normal activities of meiotic

chromosomes in the testes of *An. gambiae* ZANU. In primary spermatocytes, all homologous

autosomes and sex chromosomes pair and display chiasmata in diplotene/diakinesis of prophase

I, they align with each other at the cell equator in metaphase I, and then move from each other in

anaphase I. In secondary spermatocytes, sister chromatids of each chromosome align with each

other at the cell equator in metaphase II and go to opposite poles of the cell during anaphase II.

Meiotic divisions produce spermatids that contain a haploid set of autosomes and either a Y or X

chromosome. *An. coluzzii* and *An. merus* males show similar behaviors of chromosomes during

meiosis (supplementary material, figures S3, S4).

(c) Meiotic and premeiotic failures in F1 males of interspecies hybrids

To determine possible cytogenetic mechanisms of HMS, we analyzed chromosome behavior in

normal-like testes of hybrids from the ♀*An. merus* × ♂*An. coluzzii/An. gambiae* crosses (Figure

2). In primary spermatocytes, homoeologous autosomes pair and form chiasmata in prophase I as

in pure species. Unlike pure species, X and Y chromosomes do not pair or display chiasmata in

diplotene/diakinesis of prophase I in these hybrids. We found this pattern consistent in all

analyzed hybrid males. Metaphase chromosomes in hybrids are visibly longer than at the same

stage in pure species, indicating insufficient chromatin condensation. Besides, homoeologous

chromosomes in hybrids do not segregate during anaphase. Instead, sister chromatids move to

opposite poles of the dividing cell. Because reductional division does not occur in hybrid males,

both X and Y chromatids move to the same pole during anaphase. As a result, haploid secondary

spermatocytes are not present in these males. Our FISH analysis demonstrated that spermatids in

testes of pure species normally contain either an X or Y chromosome. In contrast, we found both

X and Y chromosomes present in spermatids of the hybrids (supplementary material, figure S5).

Moreover, the abnormal spermatids are larger in size due to insufficient chromatin condensation

and the double chromosome content. Thus, we discovered that chromosomes in normal-like

testes of hybrid males start with meiotic behavior in prophase and then prematurely switch to

mitotic behavior in anaphase, thus, skipping the reductional division. The equational division of

primary spermatocytes results in dysfunctional diploid sperm in hybrids when *An. merus* is the

mother. Degenerate testes of F1s from the reciprocal ♀*An. coluzzii/gambiae* × ♂*An. merus*

crosses have only undifferentiated round germline stem cells. To visualize sex chromosomes, we

performed whole-mount FISH with labeled 18S rDNA (Cy3) and satellite AgY53B (Cy5) to

mark the sex chromosomes of the hybrid (supplementary material, figure S6). Only interphase

sex chromosomes in the nuclei of germline stem cells are detected indicating that meiosis does

not start in the underdeveloped testes of F1 hybrids if *An. merus* is the father. Thus, premeiotic

arrest in degenerate testes is the reason for the lack of spermatids in these hybrids.

(d) Gene expression in pure species and interspecies hybrids

To test if the aforementioned premeiotic or meiotic failures are associated with gene

misexpression, we analyzed the transcript profile of the premeiotic and meiotic *vasa* [27], *SMC2,*

and *SMC4* genes [28, 29], as well as the meiotic and postmeiotic *Spo11*, *Msh4*, *SMC3β* [28, 29],

β2-tubulin [30, 31], *Ams, mts*, and *Dzip1l* [32] genes in F1 hybrid males (supplementary

material, table S1). The RT-PCR results show that all these genes express in the reproductive

tissues of *An. coluzzii*, *An. merus*, and F1 hybrids from the ♀*An. merus* × ♂*An. coluzzii* cross. A

slight but noticeable increase was observed for the *vasa*, *SMC2*, *SMC4*, and *SMC3β* transcripts in

normal-like testes of these hybrids compared with testes of any of the parental species (Figure

3A). This expression profile was consistent between two independent RT-PCR experiments that

involved testes combined with MAGs and testes without MAGs. A separate experiment using

MAGs only shows no *vasa*, *SMC2*, *SMC4*, or *SMC3β* transcripts in this tissue. Although *vasa*,

SMC2, and *SMC4* express in degenerate gonads of ♀*An. coluzzii* MOPTI × ♂*An. merus* F1

hybrids, transcripts of all tested meiotic and postmeiotic genes are virtually absent in these

hybrids (Figure 3A) supporting our observation that meiosis does not occur in degenerate testes

(supplementary material, figure S6). Expression of *vasa* indicated the development of germline

stem cells even in degenerate testes of these interspecies hybrids. To test whether gene

misexpression occurs in fertile female hybrids, we performed RT-PCR for the same genes using

ovaries of pure species and hybrid females (supplementary material, figure S7). Our results show

that, unlike males, all analyzed genes have similar transcript profiles in females of pure species

and hybrids from the reciprocal crosses.

(e) Chromosomal abnormalities in interspecies hybrids

Here, we performed quantitative analyses of chromatin condensation and X-Y chromosome

pairing in pure species and their hybrids. To determine the extent of chromatin condensation in

normal-like testes of interspecies hybrids compared with pure species, we measured the lengths

of metaphase chromosomes (supplementary material, table S2). The results of statistical analysis

with a two-sample pooled *t*-test show that chromosomes in F1 hybrids are typically longer than

chromosomes in *An. coluzzii*, *An. gambiae*, or *An. merus* (supplementary material, figure S8).

For example, chromosomes of the *An. merus* origin in a hybrid background always show

significant (P<0.001) elongation by at least 1.3 times compared to pure *An. merus*. The X

chromosome of *An. merus* suffered the most serious undercondensation in hybrids, exceeding the

length of the X chromosome in the pure species background by 1.6-1.9-fold.

In our cytogenetic study of interspecies hybrids from the crosses ♀*An. merus* × ♂*An. coluzzii/An.*

gambiae, the X and Y chromosomes do not show pairing or chiasmata in diplotene/diakinesis of

prophase I (Figure 2). We hypothesized that sex chromosome pairing is affected in early

prophase I when individual chromosomes cannot be distinguished by direct visualization. To

analyze the X-Y chromosome pairing at the pachytene stage of prophase I, we performed a

whole-mount FISH and examined spatial positions of the X- and Y-specific fluorescent signals

in confocal optical sections of nuclei in testes of pure species and their hybrids (supplementary

material, figure S9). We recorded the number of nuclei with X and Y fluorescent signals

colocalized versus X and Y fluorescent signals located separately (Figure 3B, supplementary

material, table S3). The results of a statistical analysis with a two-sample pooled *t*-test

demonstrate that X and Y chromosomes pair in more than 90% of primary spermatocytes in *An.*

coluzzii, while they pair in less than 30% of primary spermatocytes in F1 hybrids of ♀*An. merus*

× ♂*An. coluzzii* (Figure 3C).

4. Discussion

In this study, we performed the first detailed cytological analysis of spermatogenesis in pure

species and hybrids of mosquitoes. We demonstrate that premeiotic and meiotic defects are

involved in HMS in reciprocal crosses (Figure S10). This observation is at odds with the

commonly accepted view that hybrid males suffer postmeiotic sterility problems more often than

premeiotic or meiotic sterility problems [1, 33]. Although many postmeiotic defects seen in

Drosophila hybrids are indeed related to problems in sperm bundling and motility [5, 9, 33],

some sperm abnormalities may stem from meiotic failures. For example, our data shows that

sperm abnormalities such as nonmotility, two-tailed heads, and chromatin decompaction in

sperm heads can result from impaired meiosis I (Figure 2). Additional studies of interspecies

hybrids of various organisms should determine whether the first meiotic division commonly fails

when fertility is affected.

Our cytogenetic investigation of meiosis in *Anopheles* species demonstrates that the X and Y

chromosomes pair with each other throughout prophase I (Figure 1, 3BC, supplementary

material, figures S3, S4). Unlike pure species, meiotic prophase I in F1 mosquito hybrids shows

cytogenetic anomalies—low percentages of X-Y chromosome pairing and insufficient chromatin

condensation (Figure 3BC, supplementary material, figure S8). The interplay between these two

phenotypes may result in failing a reductional meiotic division and prematurely proceeding to an

equational mitotic division (Figure 4). All chromosomes must achieve synapsis in pachynema,

complete DNA repair, and disassemble their synaptonemal complexes in diplonema as

homologous chromosomes prepare to segregate [34]. The presence of chiasmata and tension

exerted across homologs ensures that cells undergo reductional segregation [35]. Asynapsis of

chromosomes almost invariably triggers pachytene checkpoint and meiotic breakdown [36].

Haldane's rule predicts that meiosis would be aberrant in males but may be normal in females

[20]. This could be due to differences in the mechanisms of X-Y and X-X pairing. For example,

in *D. melanogaster*, X and Y chromosomes pair through specific pairing sites located in

heterochromatin [37], which could be relatively quickly altered during evolution. In contrast, the

X-X and autosome-autosome pairing capacity is widely distributed along the chromosome arm

[38] and is likely more robust to evolutionary changes. A study of introgression of the Y

chromosome from *An. gambiae* into *An. arabiensis* suggests that the Y chromosome does not

cause sterility [17]. This runs counter to our findings that X-Y unpairing underlies sterility in

male hybrids. Identification and analysis of X-Y pairing sites in species of the *An. gambiae*

complex may shed light on this problem.

Reciprocal crosses in malaria mosquitoes and other organisms usually produce hybrids with

sterility phenotypes of different degrees of severity [6, 18, 39]. This observation indicates that

malfunction of multiple stages of spermatogenesis can be involved in postzygotic isolation. At

the early stages of speciation, premeiotic, meiotic, and postmeiotic genes still express in hybrids

and meiotic errors are characterized by decreased pairing between sex chromosomes, insufficient

chromatin condensation, and skipping reductional division. These phenotypes are seen in sterile

hybrids from the ♀*An. merus* × ♂*An. coluzzii/An. gambiae* crosses (Figures 2, 3, 4). Higher

expression levels of the *vasa*, *SMC2,* and *SMC4* genes (Figure 3A) are consistent with our

observation of extended premeiotic and meiotic stages in these hybrids (Figure S2). It is possible

that the observed overexpression of *SMC3β* in the hybrid males may contribute to abandoning

homologous chromosome segregation in favor of sister-chromatid cohesion, recombination, and

premature segregation. This is because the SMC1 and SMC3 proteins function together with

other proteins to control cell cycle transitions by participating in sister-chromatid cohesion,

recombination, and faithful segregation [29, 40, 41]. To support crossover formation between

homologous chromosomes and their subsequent segregation in meiosis, repair from the more

readily available homologous sequences on the sister chromatid must be suppressed [34].

As species continue to diverge, meiotic errors become more prominent, and new hybrid

phenotypes appear as has been seen, for example, in sterile male hybrids from the ♀*D.*

pseudoobscura × ♂*D. persimilis* cross [6, 7]. At this stage of postzygotic isolation, meiosis is

manifested by the unpairing of most of the chromosomes and by malfunction of the abnormally

elongated spindle, resulting in spermatids with unbalanced chromosome content in sterile male

hybrids [6, 7]. In mice hybrids, spermatogenic arrest occurs in early meiosis I when

homoeologous chromosomes fail to pair and meiotic sex chromosome inactivation is disrupted,

resulting in increased apoptosis of spermatocytes [3, 4, 42]. Disruption of synapsis between

heterospecific chromosomes in prophase I has been proposed as a recurrently evolving trigger

for the meiotic arrest of interspecific F1 hybrids [3, 43]. Finally, spermatogenic abnormalities in

hybrids can happen before meiosis starts. A spermatogenic arrest at the premeiotic stage is

characterized by the repression of meiotic and postmeiotic genes and by the lack of

spermatocytes in degenerate testes of hybrid males. These phenotypes have been observed in

sterile hybrids from the ♀*D. mauritiana* × ♂*D. sechellia* cross [5] and in sterile hybrids from the

♀*An. coluzzii/An. gambiae* × ♂*An. merus* crosses (our study).

Charles Darwin, in the chapter on "Hybridism" in his book, the *Origin of Species,* rightly argued

that hybrid sterility ''is not a specially endowed quality, but is incidental on other acquired

differences [44].'' Identification of the germ-line-specific cellular and molecular differences

acquired during the early stages of postzygotic isolation between species is crucial to explaining

both speciation and normal fertility function. Our study suggests that meiotic abnormalities in

hybrid males stem from the unpairing of the sex chromosomes and insufficient chromatin

condensation. Cytological analyses of spermatogenesis in interspecies hybrids from diverse

groups of organisms may highlight general patterns and mechanisms in the origin and evolution

of postzygotic isolation. Recently evolved members of the *An. gambiae* complex represent an

excellent, new system for studying the genetic basis and molecular mechanisms of species

incompatibilities at the early stages of postzygotic reproductive isolation.

Acknowledgments

The following mosquito strains were obtained through BEI Resources, NIAID, NIH: *An.*

coluzzii, Strain Mali-NIH, Eggs, MRA-860 contributed by Nora J. Besansky; *An. coluzzii*, Strain

MOPTI, Eggs, MRA-763 contributed by Gregory C. Lanzaro; *An. gambiae*, Strain ZANU,

MRA-594 contributed by Hilary Ranson and Frank H. Collins; *An. merus*, Strain MAF, MRA-

1156 contributed by Maureen Coetzee. We thank Kristin Rose and Janet Webster for editing the

text.

References

1.   Powell JR: Progress and Prospects in Evolutionary Biology. New York - Oxford: Oxford

University Press; 1997.

2.   Turissini DA, McGirr JA, Patel SS, David JR, Matute DR: The Rate of Evolution of

Postmating-Prezygotic Reproductive Isolation in Drosophila. *Mol Biol Evol* 2018,

35(2):312-334.

3.   Bhattacharyya T, Gregorova S, Mihola O, Anger M, Sebestova J, Denny P, Simecek P,

Forejt J: Mechanistic basis of infertility of mouse intersubspecific hybrids. *Proc Natl*

Acad Sci U S A 2013, 110(6):E468-477.

4.   Forejt J, Ivanyi P: Genetic studies on male sterility of hybrids between laboratory and

wild mice (Mus musculus L.). *Genet Res* 1974, 24(2):189-206.

5.   Kulathinal R, Singh RS: Cytological Characterization of Premeiotic Versus Postmeiotic

Defects Producing Hybrid Male Sterility among Sibling Species of the Drosophila

Melanogaster Complex. *Evolution* 1998, 52(4):1067-1079.

6.   Dobzhansky T: Studies on hybrid sterility. I. Spermatogenesis in pure and hybrid

Drosophila pseudoobscura. *Zeitschr Zelljorsch U Mikrosk Anat* 1934, 21(2):169-223.

7.   Dobzhansky T: On the sterility of the interracial hybrids in Drosophila pseudoobscura. *P*

Natl Acad Sci USA 1933, 19:397-403.

8.   Hardy RW, Lougheed A, Markow TA: Reproductive tract and spermatid abnormalities of

hybrid males from reciprocal crosses between Drosophila mojavensis and D. arizonae.

Fly (Austin) 2011, 5(2):76-80.

9.   Coyne JA: Genetic-Basis of Male-Sterility in Hybrids between 2 Closely Related Species

of Drosophila. *P Natl Acad Sci-Biol* 1984, 81(14):4444-4447.

10.   Masly JP, Jones CD, Noor MA, Locke J, Orr HA: Gene transposition as a cause of hybrid

sterility in Drosophila. *Science* 2006, 313(5792):1448-1450.

11.   Coluzzi M, Sabatini A, della Torre A, Di Deco MA, Petrarca V: A polytene chromosome

364   analysis of the Anopheles gambiae species complex. *Science* 2002, 298(5597):1415-1418.

365   12.   Coetzee M, Hunt RH, Wilkerson R, Della Torre A, Coulibaly MB, Besansky NJ:

366   Anopheles coluzzii and Anopheles amharicus, new members of the Anopheles gambiae

367   complex. *Zootaxa* 2013, 3619(3):246-274.

368   13.   Barron MG, Paupy C, Rahola N, Akone-Ella O, Ngangue MF, Wilson-Bahun TA, Pombi

369   M, Kengne P, Costantini C, Simard F *et al*: A new species in the Anopheles gambiae

370   complex reveals new evolutionary relationships between vector and non-vector species.

371   *bioRxiv* 2018, https://doi.org/10.1101/460667.

372   14.   Fontaine MC, Pease JB, Steele A, Waterhouse RM, Neafsey DE, Sharakhov IV, Jiang XF,

373   Hall AB, Catteruccia F, Kakani E *et al*: Extensive introgression in a malaria vector

374   species complex revealed by phylogenomics. *Science* 2015, 347(6217):42-+.

375   15.   Thawornwattana Y, Dalquen D, Yang Z: Coalescent Analysis of Phylogenomic Data

376   Confidently Resolves the Species Relationships in the Anopheles gambiae Species

377   Complex. *Mol Biol Evol* 2018, 35(10):2512-2527.

378   16.   Slotman M, Della Torre A, Powell JR: The genetics of inviability and male sterility in

379   hybrids between Anopheles gambiae and An. arabiensis. *Genetics* 2004, 167(1):275-287.

380   17.   Bernardini F, Galizi R, Wunderlich M, Taxiarchi C, Kranjc N, Kyrou K, Hammond A,

381   Nolan T, Lawniczak MNK, Papathanos PA *et al*: Cross-Species Y Chromosome Function

382   Between Malaria Vectors of the Anopheles gambiae Species Complex. *Genetics* 2017,

383   207(2):729-740.

18. Davidson G, Paterson HE, Coluzzi M, Mason GF, Micks DW: The Anopheles gambiae

Complex. In: *Genetics of Insect Vectors of Diesease.* Edited by Wright JW, Pal R.

Amsterdam-London-New York: Elsevier Publishing Company; 1967: 211-249.

19. Presgraves DC, Orr HA: Haldane's rule in taxa lacking a hemizygous X. *Science* 1998,

282(5390):952-954.

20. Haldane JBS: Sex ratio and unisexual sterility in hybrid animals. *J Genet* 1922,

12(2):101-109.

21. Curtis CF: The mechanism of hybrid male sterility from crosses in the Anopheles

gambiae and Glossina morsitans complexes. In: *Recent Developments in the Genetics of*

Disease Vectors. Edited by Steiner WM, Tabachnick WJ, Rai KS, Narang S. Champaign,

IL: Stipes Publishing Company; 1982: 290-312.

22. Timoshevskiy VA, Sharma A, Sharakhov IV, Sharakhova MV: Fluorescent in situ

Hybridization on Mitotic Chromosomes of Mosquitoes. *J Vis Exp* 2012(67).

23. Sharakhova MV, George P, Timoshevskiy V, Sharma A, Peery A, Sharakhov IV:

Mosquitoes (Diptera). In: *Protocols for Cytogenetic Mapping of Arthropod Genomes.*

Edited by Sharakhov IV. Boca Raton, FL: CRC Press, Taylor & Francis Group; 2015:

93-170.

24. Thailayil J, Magnusson K, Godfray HC, Crisanti A, Catteruccia F: Spermless males elicit

large-scale female responses to mating in the malaria mosquito Anopheles gambiae. *Proc*

Natl Acad Sci U S A 2011, 108(33):13677-13681.

25. Diabate A, Dabire RK, Millogo N, Lehmann T: Evaluating the effect of postmating

isolation between molecular forms of Anopheles gambiae (Diptera: Culicidae). *J Med*

Entomol 2007, 44(1):60-64.

26.   Aboagye-Antwi F, Alhafez N, Weedall GD, Brothwood J, Kandola S, Paton D, Fofana A,

408   Olohan L, Betancourth MP, Ekechukwu NE *et al*: Experimental swap of Anopheles

409   gambiae's assortative mating preferences demonstrates key role of X-chromosome

410   divergence island in incipient sympatric speciation. *PLoS Genet* 2015, 11(4):e1005141.

411   27.   Papathanos PA, Windbichler N, Menichelli M, Burt A, Crisanti A: The vasa regulatory

412   region mediates germline expression and maternal transmission of proteins in the malaria

413   mosquito Anopheles gambiae: a versatile tool for genetic control strategies. *BMC Mol*

414   *Biol* 2009, 10:65.

415   28.   Malik SB, Pightling AW, Stefaniak LM, Schurko AM, Logsdon JM, Jr.: An expanded

416   inventory of conserved meiotic genes provides evidence for sex in Trichomonas vaginalis.

417   *PLoS One* 2007, 3(8):e2879.

418   29.   Losada A, Hirano T: Dynamic molecular linkers of the genome: the first decade of SMC

419   proteins. *Genes Dev* 2005, 19(11):1269-1287.

420   30.   Catteruccia F, Crisanti A, Wimmer EA: Transgenic technologies to induce sterility.

421   *Malar J* 2009, 8 Suppl 2:S7.

422   31.   Kemphues KJ, Kaufman TC, Raff RA, Raff EC: The testis-specific beta-tubulin subunit

423   in Drosophila melanogaster has multiple functions in spermatogenesis. *Cell* 1982, 31(3 Pt

424   2):655-670.

425   32.   Krzywinska E, Krzywinski J: Analysis of expression in the Anopheles gambiae

426   developing testes reveals rapidly evolving lineage-specific genes in mosquitoes. *BMC*

427   *Genomics* 2009, 10:300.

428   33.   Coyne JA, Orr HA: Speciation. Sunderland, Mass.: Sinauer Associates; 2004.

429   34.   Subramanian VV, Hochwagen A: The meiotic checkpoint network: step-by-step through

430   meiotic prophase. *Cold Spring Harb Perspect Biol* 2014, 6(10):a016675.

431   35.   Miyazaki S, Kim J, Sakuno T, Watanabe Y: Hierarchical Regulation of Centromeric

432   Cohesion Protection by Meikin and Shugoshin during Meiosis I. *Cold Spring Harb Symp*

433   *Quant Biol* 2017, 82:259-266.

434   36.   Bolcun-Filas E, Schimenti JC: Genetics of meiosis and recombination in mice. *Int Rev*

435   *Cell Mol Biol* 2012, 298:179-227.

436   37.   McKee BD, Yan R, Tsai JH: Meiosis in male Drosophila. *Spermatogenesis* 2012,

437   2(3):167-184.

438   38.   McKee BD: Homolog pairing and segregation in Drosophila meiosis. *Genome Dyn* 2009,

439   5:56-68.

440   39.   Good JM, Handel MA, Nachman MW: Asymmetry and polymorphism of hybrid male

441   sterility during the early stages of speciation in house mice. *Evolution* 2008, 62(1):50-65.

442   40.   Revenkova E, Eijpe M, Heyting C, Hodges CA, Hunt PA, Liebe B, Scherthan H,

443   Jessberger R: Cohesin SMC1 beta is required for meiotic chromosome dynamics, sister

444   chromatid cohesion and DNA recombination. *Nat Cell Biol* 2004, 6(6):555-562.

445   41.   Biswas U, Stevense M, Jessberger R: SMC1alpha Substitutes for Many Meiotic

446   Functions of SMC1beta but Cannot Protect Telomeres from Damage. *Curr Biol* 2018,

447   28(2):249-261 e244.

448   42.   Schwahn DJ, Wang RJ, White MA, Payseur BA: Genetic Dissection of Hybrid Male

449   Sterility Across Stages of Spermatogenesis. *Genetics* 2018.

450   43.   Bhattacharyya T, Reifova R, Gregorova S, Simecek P, Gergelits V, Mistrik M,

451   Martincova I, Pialek J, Forejt J: X chromosome control of meiotic chromosome synapsis

452   in mouse inter-subspecific hybrids. *PLoS Genet* 2014, 10(2):e1004088.

453   44.   Darwin C: On the origin of species by means of natural selection or, The Preservation of

454   favoured races in the struggle for life. In. London: Electric Book Co.; 2001: 1 online

455   resource (422 p.).

Figure Legends

Figure 1. Chromosome behavior during meiosis in testes of *An. gambiae.* X chromosomes are

labeled with 18S rDNA (red); Y chromosomes are labeled with retrotransposon *zanzibar* (green).

Chromosomes are counterstained with DAPI (blue). Scale bar – 5 μm.

Figure 2. Chromosome behavior during meiosis in testes of interspecies hybrids. (A) ♀*An. merus*

× ♂*An. coluzzii* MALI. (B) ♀*An. merus* × ♂*An. coluzzii* MOPTI. (C) ♀*An. merus* × ♂*An.*

gambiae ZANU. X chromosomes are labeled with 18S rDNA (red) and Y chromosomes are

labeled with retrotransposon *zanzibar* (green). Chromosomes are counterstained with DAPI

(blue). Scale bar – 5 μm.

Figure 3. Molecular and chromosomal abnormalities in interspecies hybrids. (A) Gene

expression in male reproductive organs of *An. coluzzii* MOPTI, *An. merus*, F1 hybrids of ♀*An.*

merus × ♂*An. coluzzii* MOPTI (F1 M×C), and F1 hybrids of ♀*An. coluzzii* MOPTI × ♂*An.*

merus (F1C×M) analyzed by RT-PCR. *AgS7* – an endogenous control gene. (B) Confocal

images of primary spermatocytes with FISH showing pairing and unpairing of X (red) and Y

(green) chromosomes in *An. coluzzii* MOPTI males and F1 hybrid males from the ♀*An. merus* ×

♂*An. coluzzii* MOPTI cross. (C) Percentage of the X and Y chromosome pairing in primary

spermatocytes of six *An. coluzzii* MOPTI males and six F1 hybrid males from the ♀*An. merus* ×

♂*An. coluzzii* MOPTI cross. Statistical significance was assessed with a two-sample pooled *t*-

test.

Figure 4. Scheme of male meiosis in pure species and interspecies hybrids of the *An. gambiae*

complex. Unlike pure species, X and Y chromosomes in meiotic prophase I of hybrids are

largely unpaired. The absence of chiasmata between the sex chromosomes and insufficient

chromatin condensation reduce tension exerted across the homoeologous chromosomes,

triggering premature sister chromatid segregation. As a result, instead of a reductional division,

primary spermatocytes in interspecies hybrids undergo an equational mitotic division leading to

diploid nonmotile sperm.

18S rDNA (red)		
zanzibar (green)		
Prophase I	Metaphase I	Anaphase I
Metaphase II	Anaphase II	Spermatid
Metaphase II	Anaphase II	Spermatid

Normal male meiosis in pure species

Prophase I   Anaphase I   Normal sperm

X   Y

Failed male meiosis in F1 hybrids

Prophase I   Anaphase   Diploid sperm

X   Y