Premeiotic and meiotic failures lead to hybrid male sterility in the *Anopheles*

***gambiae* complex**

Jiangtao Liang[1], Igor V. Sharakhov[1,2,*]

[1]Department of Entomology, Virginia Polytechnic Institute and State University, Blacksburg,

Virginia, United States of America

[2]Department of Cytology and Genetics, Tomsk State University, Tomsk, Russian Federation

[*]**Corresponding author**

E-mail: igor@vt.edu

Abstract

Background: Hybrid male sterility contributes to speciation by restricting gene flow between

related taxa at the beginning stages of postzygotic isolation. The genetic factors, cellular basis,

and molecular mechanisms of hybrid male sterility are of considerable interest as they inform our

understanding of both speciation and normal fertility function. However, we have limited

knowledge about precise cellular phenotypes that come into play when fertility is first affected.

Hybrids between closely related species of the *Anopheles gambiae* complex offer opportunities

to identify spermatogenic errors that arise early during speciation.

Results: To investigate possible cellular causes of hybrid male sterility, we performed crosses

between sibling species of the *An. gambiae* complex. Our results demonstrate that testes are

severely underdeveloped in hybrids between male *An. merus* and female *An. gambiae* or *An.*

coluzzii. No meiotic chromosomes are identified and suppressed expression of germline-specific

genes *β2-tubulin, Ams, mts*, and *Dzip1l* is observed in these hybrid males. However, testes have

nearly normal morphologies and sizes but produce mostly nonmotile spermatozoa in hybrids

from the reciprocal crosses. Using chromosome X- and Y-specific fluorescent probes, we

followed the process of meiosis in each species and their F1 hybrids between female *An. merus*

and male *An. gambiae* or *An. coluzzii*. Unlike for pure species, sex chromosomes in meiotic

prophase I of F1 hybrids are largely unpaired and all chromosomes show various degrees of

insufficient condensation. Instead of entering the reductional division in meiosis I, primary

spermatocytes undergo an equational mitotic division producing abnormal diploid sperm.

Meiotic chromosomes of some F1 hybrid individuals are involved in *de novo* genome

rearrangements. Yet, the germline-specific genes *β2-tubulin, Ams, mts*, and *Dzip1l* express

normally in these hybrid males.

Conclusions: Our study demonstrates that hybrid male sterility in the malaria mosquitoes

involves two defects in reciprocal crosses: premeiotic arrest in germline stem cells and the failure

of the reductional meiotic division in primary spermatocytes. We identified cytogenetic errors in

hybrids that arise during the early stages of postzygotic isolation. This knowledge will inform the

development of innovative mosquito control strategies based on population suppression by

manipulating reproduction via genetic technologies.

Keywords: *Anopheles*, chiasmata, chromosome pairing, gene expression, hybrid male sterility,

meiosis, microscopy, mitosis, mosquito, speciation, spermatogenesis.

Background

Hybrid male fertility is among the first phenotypes affected as the postzygotic isolation between species is being established [1]. Therefore, genetic factors, cellular basis, and molecular mechanisms of hybrid male sterility are of considerable interest, as they inform our understanding of both speciation and normal fertility function. Genetic loci responsible for hybrid male sterility in plants and animals are often being identified using quantitative trait loci (QTL) mapping [1-8]. In a variety of organisms, hybrid sterility can be caused by the genetic factors on the X chromosome or by epistasis between X-linked and autosomal alleles [4, 8-15]. Studies in animals commonly assess testis shape, size or weight, as well as sperm morphology, density or motility to define male sterility phenotypes. However, a cellular basis of hybrid male sterility is rarely investigated. Cytological studies of spermatogenesis in sterile hybrids indicate that multiple mechanisms of functional sterility are possible. Sterile hybrids between *Mus musculus domesticus* and *M. m. musculus* have spermatogenic arrest in early meiosis I with disrupted both homoeologous chromosome pairing and meiotic sex chromosome inactivation [16, 17], as well as reductions in spermatocyte and spermatid number, increased apoptosis of primary spermatocytes, and more multinucleated syncytia [18]. Studies of failed spermatogenesis in different *Drosophila* hybrids observed arrests at the premeiotic stage [19], reduced chromosome pairing and unequal chromosome segregation in meiosis [20, 21], characteristic spermiogenic arrests [19], spermatid abnormalities [22], and problems in sperm bundling and motility [23, 24]. Detailed analyses of cellular and developmental abnormalities in testes of various hybrid organisms could help better define the causes of infertility. Cytological studies of spermatogenesis in hybrids may link different sterility phenotypes to the early or advance stages

of postzygotic isolation. Moreover, identification of first-evolved cytogenetic errors in hybrids

between closely related species could improve our understanding of speciation.

The *Anopheles gambiae* complex consists of at least eight morphologically nearly

indistinguishable sibling species of African malaria mosquitoes [25, 26]. Genome-based

estimations of the age of the *An. gambiae* complex vary from 1.85 [27] to as young as 0.526

million years [28]. Genomic introgression has been found prevalent in autosomal regions of

several species indicating naturally occuring interspecies hybridization [27, 28]. Experimental

crosses of species from the *An. gambiae* complex often produce sterile F1 hybrid males

confirming Haldane's rule of sterility or inviability of the heterogametic sex [13, 29-31]. Early

crossing experiments between members of the complex have found that hybrid male sterility is

associated with various degrees of atrophies of testes and underdevelopment of sperms [30].

Examination of hybrid sterility in backcrosses of *An. gambiae* and *An. arabiensis* using an X-

linked white-eye marker demonstrated a large effect of the X chromosome on male hybrid

sterility [32]. The large X effect on hybrid sterility has been supported with a QTL mapping

approach [13]. However, an introgression of the Y chromosome from *An. gambiae* into a

background of *An. arabiensis* has shown no apparent influence on male fertility, fitness, or gene

expression [29]. A recent study of hybridization in nature has shown that postmating isolation is

positively associated with ecological divergence of *An. coluzzii, An. gambiae*, and *An. arabiensis*

[33]. However, the cellular basis and molecular mechanisms of hybrid male sterility in

mosquitoes are unknown. Because of the recent evolution and ease of hybridization, sibling

species of the *An. gambiae* complex offer great opportunities to provide insights into

mechanisms of speciation.

A recent World Health Organization report has shown that after a successful period of global

malaria control, progress has stalled since 2015. There are still 219 million cases and 435,000

related deaths from malaria without a tendency to decrease [34]. In the 60s-70s, the sterile insect

technique (SIT) was one of the first methods used to control the malaria-transmitting *Anopheles*

mosquitoes as it aimed to interfere with their reproduction by introducing sterile males into

natural populations. However, because of the low competitiveness of X-ray-irradiated males and

the lack of institutional commitments, the traditional SIT largely failed in controlling malaria

mosquitoes [35]. Novel genetic approaches based on CRISPR-Cas9 gene drives [36-40] show

potential toward the generation of self-sustainable and species-specific mosquito control

strategies. A better understanding of the normal male fertility function and mechanisms of

naturally occurring hybrid male sterility will inform the development of novel SIT tools and

implementation of pre-existing technologies for the control of malaria-transmitting mosquitoes.

Here, we investigate possible cellular causes of male sterility in hybrids between sibling species

of the *An. gambiae* complex. We asked three specific questions: (i) Which cellular processes are

involved in causing infertility in hybrid mosquito males? (ii) Are the defects leading to hybrid

male sterility premeiotic, meiotic, or postmeiotic? and (iii) What cytogenetic errors trigger the

spermatogenic breakdown? We demonstrate that hybrid male sterility in malaria mosquitoes

involves two cellular defects in reciprocal crosses: premeiotic arrest in germline stem cells and

the failure of the reductional meiotic division in primary spermatocytes. Our data suggest that the

meiotic abnormalities in hybrid males stem from the unpairing of the sex chromosomes and

chromatin decondensation. Thus, our study identified first cytogenetic errors in hybrids that arise

during the early stages of postzygotic isolation.

Results

Hybrid male sterility phenotypes at the cellular level

To obtain sterile males, we performed reciprocal crosses between *An. merus* MAF and *An.*

gambiae ZANU or *An. coluzzii* MOPTI and MALI. Backcrossing of F1 males to parental

females resulted in no progeny demonstrating sterility of F1 hybrid males. This result confirms

Haldane's rule for the majority of interspecies crosses in the *An. gambiae* complex, except for

crosses between *An. gambiae* and *An. coluzzii*: F1 females are fertile while F1 males are sterile

[13, 29-31, 41, 42]. To investigate the developmental phenotypes involved in hybrid sterility, we

dissected testes from adult males obtained from interspecies crosses and from pure species.

Normal testes of pure species have a spindle-like shape (Fig. 1A). We found obvious asymmetry

in morphology and sizes of testes when reciprocal crosses were compared. Hybrid males from

crosses between female *An. merus* and male *An. gambiae* or *An. coluzzii* display normal-like

reproductive organs (Fig. 1B). In contrast, F1 males from crosses between female *An. gambiae*

or *An. coluzzii* and male *An. merus* show severely underdeveloped testes (Fig. 1C). We then

tested if normal-like and underdeveloped testes of interspecies hybrids produce any sperm. In

squashed testes of 2-day-old adults of pure species, large amounts of mature spermatozoa with

long tails can be seen. After we crushed the testes, spermatozoa with vibrant motility escaped

from the ruptures (Fig. 1A, Additional file 1: Movie S1). However, mature sperm or sperm

motility hardly could be seen in squashed or crushed normal-like testes of 2-day-old adult

hybrids from crosses when *An. merus* was the mother. Instead, we see delayed spermatid

differentiation, fewer spermatids, and mostly nonmotile spermatozoa with large heads and often

two short tails growing from opposite ends of the head (Fig. 1B, Additional file 2: Movie S2).

Only undifferentiated round cells could be seen in underdeveloped testes of hybrids from

reciprocal crosses when *An. merus* was the father (Fig. 1C). Given the small size of the

degenerate testes, these round cells may represent germline stem cells. Thus, neither normal-like

nor underdeveloped testes of the interspecies hybrids produce mature motile spermatozoa.

The normal progress of meiosis in males of pure species

Before we investigate meiosis of interspecies hybrids, we provide the first description of

chromosome behavior in meiosis of pure *Anopheles* species. The chromosome complement of

Anopheles males consists of three chromosome pairs: two autosomes (2 and 3) and X and Y sex

chromosomes. With the help of sex-chromosome-specific fluorescent probes, we followed the

progress of meiosis in *An. gambiae*, *An. coluzzii*, and *An. merus.* The following DNA sequences

were used as the probes for fluorescence *in situ* hybridization (FISH). Retroelement *zanzibar* is

specific to the Y chromosome of *An. gambiae* and *An. coluzzii* but is absent in *An. merus*; 18S

ribosomal DNA (rDNA) labels only the X chromosome in *An. gambiae* or *An. coluzzii* but labels

both X and Y chromosomes in *An. merus*; and satellite AgY53B [43] labels both X and Y

chromosomes in all three species [44] (Table 1). Fig. 2 shows normal activities of meiotic

chromosomes in testes of *An. gambiae* ZANU. In primary spermatocytes, all homologous

autosomes and sex chromosomes pair and display chiasmata in diplotene/diakinesis of prophase

I, they align with each other at the cell equator in metaphase I, and then move from each other in

anaphase I. In secondary spermatocytes, sister chromatids of each chromosome align with each

other at the cell equator in metaphase II and go to opposite poles of the cell during anaphase II.

Meiotic divisions produce spermatids that contain a haploid set of autosomes and either a Y or X

chromosome. *An. coluzzii* males show similar morphology and behavior of chromosomes during

meiosis (Additional file 3: Figure S1). Because retroelement *zanzibar* is absent in *An. merus*, we

used satellite AgY53B and 18S rDNA to label sex chromosomes in this species. Each of these

probes hybridized with both X and Y chromosomes in *An. merus*, making discrimination

between X and Y more difficult in this species. However, we could differentiate metaphase sex

chromosomes by the euchromatic arm of the X chromosome and by the slightly larger distal

heterochromatic block of the Y chromosome (Additional file 4: Figure S2). Heterochromatic

parts of the X and Y chromosomes are relatively large and structurally similar in *An. merus* in

comparison with *An. gambiae* or *An. coluzzii*. In the latter two species, the heterochromatic parts

of the X and Y chromosomes are relatively small and substantially different from each other both

in size and genetic content [44]. Despite these differences between *An. merus* and either *An.*

gambiae or *An. coluzzii*, the activities of meiotic sex chromosomes in testes are similar in all

three species.

Table 1. Labeled oligos and primer sequences used for FISH and RT-PCR.

Sequence name	GenBank accession number	VectorBase gene ID	Labeled oligos and primer sequences	Reference to oligo design
18S rDNA	AM157179	AGAP028978	F: AACTGTGGAAAAGCCAGAGC	This study
			R: TCCACTTGATCCTTGCAAAA	
zanzibar	KP878482		F: TTCTTCGATGTTGTGCTGGA	[44]
			R: ATGGAGAAACAGGGCAACAA	
			F: ATGCATGCTTGGATTCCTTC	

			R: GGTTTCTATGATCGCCTGGA	
			F: TTGGCATTCATCTGTCCAAA	
			R: GCACCCTTGATCTCATGTCA	
AgY53B	AY754156		F: CCTTTAAACACATGCTCAAATT	[43]
			R: GTTTCTTCATCCTTAAAGCCTAG	
AgY477-AgY53B junction region	AY754151		TTCTAAGTTTCTAGGCTTTAAGGATG AAGAAACCGACTATTC[Cyanine5]	This study
Contig_240	SRS667972, SRR1509742, SRR1508169		CAATAAATTTCCTTTTTAATGATGCA AAATCTACGTCTCTAGC[Cyanine3]	This study
Tandem-42345	ERX008245		GGCTTCATTTGGATGAAACATGGCCG TAACAACGATAATAGAT[Cyanine3]	This study
vasa	XM_314684	AGAP008578	F: TTCTGCTGAGGTGCTTAGCG	This study
			R: CGTCTCCGCTCATGTTTCCT	
β2-tubulin	XM_314718	AGAP008622	F: GTACGTGCCGGATCATTTCG	This study
			R: GGCCAGTTTGCAAATGCACTA	
Ams	FJ869235	AGAP029148	F: CATACGGGAGGTGAGGAAAT	[45]
			R: CCCCTTCATGCTTCATCTT	
mts	FJ869236	AARA006451	F: TGGGATCCAAATTATTTCGTG	
			R: CTGTTCGGTTCAACAATGGA	
Dzip1l	FJ869237	AGAP001165	F: GGCCAAAGTGATACAAATTGTTT	
			R: CGTTTCCAATAGGGACTTCG	
AgS7	L20837	AGAP010592	F: AGAACCAGCAGACCACCATC	[46]
			R: GCTGCAAACTTCGGCTATTC	

Meiotic failures in F1 males of interspecies hybrids

To determine possible cytogenetic mechanisms of hybrid male sterility, we analyzed

chromosome behavior in testes of hybrids from interspecies crosses. Meiotic chromosomes were

present in normal-like testes of hybrids from the ♀*An. merus* × ♂*An. coluzzii/An. gambiae*

crosses and we identified important abnormalities of meiosis in these males (Fig. 3). In primary

spermatocytes, homoeologous autosomes pair and form chiasmata in prophase I as in pure

species, but X and Y chromosomes do not display chiasmata in diplotene/diakinesis of prophase

I. We found this pattern consistent in all analyzed hybrid males. In hybrids, metaphase

chromosomes are visibly longer than at the same stage in pure species, indicating insufficient

chromatin condensation. Besides, homoeologous chromosomes in hybrids do not segregate

during anaphase. Instead, sister chromatids move to opposite poles of the dividing cell. Because

reductional division does not occur in hybrid males, both X and Y chromatids move to the same

pole during anaphase. As a result, haploid secondary spermatocytes do not form in these males.

Our FISH analysis demonstrated that each spermatid in testes of pure species normally contains

either an X or Y chromosome. In contrast, we found that both X and Y chromosomes present in

each spermatid of the hybrids (Additional file 5: Figure S3). Moreover, the abnormal spermatids

are larger in size due to insufficient chromatin condensation and the double chromosome content.

Thus, we discovered that chromosomes in normal-like testes of hybrid males start with a meiotic

behavior in prophase and then switch to a mitotic behavior in anaphase. The equational division

of primary spermatocytes results in dysfunctional diploid sperm in hybrids if *An. merus* is the

mother. In contrast, degenerate testes of F1s from the reciprocal ♀*An. coluzzii/gambiae* × ♂*An.*

merus crosses have only undifferentiated round germline stem cells. To visualize sex

chromosomes, we performed whole-mount FISH with labeled 18S rDNA (Cy3) and satellite

AgY53B (Cy5) to mark the sex chromosomes of the hybrid (Additional file 6: Figure S4). Only

interphase sex chromosomes in nuclei of germline stem cells are detected indicating that meiosis

does not start in the underdeveloped testes of F1 hybrids if *An. merus* is the father. Thus, the

premeiotic arrest in the degenerate testes is the reason for the lack of spermatids in these hybrids.

Chromosomal and molecular abnormalities in interspecies hybrids

Here, we performed quantitative analyses of chromatin condensation and the X-Y chromosome

pairing in pure species and their hybrids. We also tested if germline-specific genes express in

sterile male hybrids from the reciprocal crosses. To determine the extent of chromatin

condensation in normal-like testes of interspecies hybrids in comparison with pure species, we

measured the lengths of metaphase chromosomes (Additional file 7: Table S1). The results of a

statistical analysis with a two-sample pooled *t*-test show that chromosomes in F1 hybrids are

typically longer than chromosomes in *An. coluzzii*, *An. gambiae*, or *An. merus* (Fig. 4A,

Additional file 8: Figure S5). For example, chromosomes of the *An. merus* origin in a hybrid

background always show significant (P<0.001) elongation by at least 1.3 times in comparison

with pure *An. merus*. The X chromosome of *An. merus* suffered the most serious

undercondensation in hybrids exceeding the length of the X chromosome in the pure species

background by 1.6-1.9-fold.

In our cytogenetic study of interspecies hybrids from the crosses ♀*An. merus* × ♂*An. coluzzii/An.*

gambiae, the X and Y chromosomes do not show pairing or chiasmata in diplotene/diakinesis of

prophase I (Fig. 3). We hypothesized that sex chromosome pairing is affected in the early

prophase I when individual chromosomes cannot be distinguished by direct visualization. To

analyze the X-Y chromosome pairing at the pachytene stage of prophase I, we performed a

whole-mount FISH and examined spatial positions of the X- and Y-specific fluorescent signals

in confocal optical sections of nuclei in testes of pure species and their hybrids (Additional file 9:

Figure S6). The *An. merus* X chromosomes were labeled with 18S rDNA (Cy3) and a satellite

from Contig_240 (Cy3), while the *An. coluzzii* MOPTI Y chromosomes were labeled with

satellite AgY53B (Cy5) and satellite AgY477-AgY53B junction region (Cy5). We recorded the

number of nuclei with X and Y fluorescent signals colocalized versus X and Y fluorescent

signals located separately (Fig. 4B). The results of a statistical analysis with a two-sample pooled

t-test demonstrate that X and Y chromosomes pair in more than 90% of primary spermatocytes

in the *An. coluzzii*, while they pair in less than 30% of primary spermatocytes in F1 hybrids of

♀*An. merus* × ♂*An. coluzzii* (Fig. 4C).

To test if premeiotic or meiotic failures are associated with the misexpression of germline-

specific genes, we analyzed the presence of the postmitotic germline transcripts *Ams, mts*, *Dzip1l*

[45], and *β2-tubulin* [47, 48] in F1 hybrid males (Table 1). The reverse transcription polymerase

chain reaction (RT-PCR) results show that these genes express at similar levels in the

reproductive tissues of *An. coluzzii* and F1 hybrids from the ♀*An. merus* × ♂*An. coluzzii* MOPTI

cross (Fig. 4D). However, *Ams, mts*, *Dzip1l,* and *β2-tubulin* are strongly down regulated in

gonads of the ♀*An. coluzzii* MOPTI × ♂*An. merus* F1 hybrids, supporting our observation that

meiosis does not occur in these hybrids. In contrast, a pre-meiotic gene, *vasa* [49] (Table 1),

expresses at similar levels in reproductive tissues of all hybrids and pure species, indicating that

germline stem cells present even in degenerate testes of interspecies hybrids.

Besides the abnormalities that affect all progeny of the ♀*An. merus* × ♂*An. coluzzii/An. gambiae*

crosses, chromosomes of some hybrid individuals display structural abnormalities. In several

cases, chromosomes were involved in *de novo* genome rearrangements (Fig. 5). For example, a

large fragment of the *An. merus* X chromosome, including the rDNA locus, was translocated to

the 2L arm of *An. coluzzii* in the male hybrid from the ♀*An. merus* × ♂*An. coluzzii* MALI cross

(Fig. 5A). In addition, we detected a duplication of a chromosomal segment involving the rDNA

locus within the X chromosome of *An. merus* in the ♀*An. merus* × ♂*An. gambiae* ZANU cross

(Fig. 5B). These rearrangements were not observed in pure species, and their occurrence in F1

hybrid males suggests an increased genome instability as a result of interspecies hybridization. In

addition, the *An. merus* X chromosomes in the hybrid karyotype can be involved in formation of

chromatin bridges during the abnormal anaphase (Additional file 10: Figure S7). Finally, we

noticed that heterochromatic regions of hybrid males have elevated fragility compared with

heterochromatic regions of pure species males. When squashed chromosome preparations are

made, pericentromeric heterochromatin of meiotic chromosomes in the ♀*An. merus* × ♂*An.*

coluzzii MOPTI hybrids, unlike pure species, can be stretched in prophase and uncoiled in

metaphase due to normal mechanical pressure (Additional file 11: Figure S8), suggesting

malfunctioning of heterochromatin proteins.

Discussion

Cellular and molecular defects involved in hybrid male sterility

In this study, we performed the first detailed cytological analysis of spermatogenesis in pure

species and hybrids of mosquitoes. Hybrid male sterility phenotypes in the *An. gambiae* complex

are clearly asymmetric between the reciprocal crosses (Fig. 6), as has been demonstrated earlier

for mosquitoes and other animals [8, 20, 30, 50]. The observed cellular and molecular

abnormalities suggest the following developmental scenarios in testes of the interspecies hybrids.

In hybrids from crosses with *An. merus* as the father, mitotic divisions of germline cells are

impaired, no meiosis occurred, and expression of postmitotic germline-specific genes is

repressed (Fig. 4D). As a result, these hybrids have degenerate testes with arrested germline stem

cells. Degenerate testes have also been observed in F1 hybrids between sibling species of the *An.*

albitarsis [51] and the *An. barbirostris* [52] complexes. However, in hybrids from the reciprocal

crosses with *An. merus* as the mother, mitotic divisions of germ cells occur as normally as in

pure species. Moreover, germ cells in these hybrids go through meiotic prophase I as evidenced

from our cytogenetic analyses (Fig. 3) and expression of the postmitotic germline genes (Fig.

4D). However, the downstream developmental processes in these hybrids are impaired. The

abnormalities begin in meiosis I where homoeologous chromosomes fail to segregate. Testes of

these hybrids show delayed spermatid differentiation, a smaller number of these cells, and

eventual formation of immotile abnormal (including two-tailed) sperm with insufficient

chromatin condensation (Fig. 6). Thus, we demonstrate that premeiotic and meiotic defects are

involved in hybrid male sterility in malaria mosquitoes. This observation is at odds with the

commonly accepted view that more often hybrid males suffer postmeiotic sterility problems [1,

2]. Although many postmeiotic defects seen in *Drosophila* hybrids are indeed related to

problems in sperm bundling and motility [2, 19, 23], some sperm abnormalities may stem from

meiotic failures. For example, our results demonstrate that such sperm abnormalities as

nonmotility, two-tailed heads, and chromatin decompaction in sperm heads result from the

impaired meiosis I. However, very few studies have been devoted to the detailed cytological

investigation of meiosis in interspecies hybrids, especially in dipteran insects. For the first time,

Theodosius Dobzhansky described pronounced defects in meiosis I in sterile male hybrids from

the ♀*D. pseudoobscura* × ♂*D. persimilis* cross [20, 21]. Also, a histological investigation of

spermatogenesis in hybrid mice identified defects in meiosis I as a primary barrier to

reproduction [18]. Additional studies of interspecies hybrids of various organisms should

determine if the first meiotic division commonly fails when fertility is affected.

Cytogenetic mechanisms of hybrid male sterility

Our cytogenetic investigation of meiosis in pure *Anopheles* species revealed that the X and Y

chromosomes normally pair tightly with each other throughout prophase I (Fig.s 2, 4BC, and

Additional files 3, 4). Sex chromosomes in malaria mosquito males genetically recombine [44]

highlighting an important difference from male meiosis in fruit flies [53-55]. As a result of the

recombination, some mosquito species, such as *An. merus*, have homomorphic heterochromatic

parts in their X and Y chromosomes (Additional file 4: Figure S2). Future sequencing and

assembly of the heterochromatin in malaria mosquitoes using long reads may yield important

insights into structural and functional organization of the sex chromosomes, as it has been

demonstrated for *Drosophila* [56].

Unlike pure species, the meiotic prophase I in F1 mosquito hybrids shows cytogenetic

anomalies—low percentage of the X-Y chromosome pairing and insufficient chromatin

condensation (Fig. 7). Disruption of chromosome pairing and synapsis is frequently observed in

interspecies hybrids of various organisms. Similar to the mosquitoes, pairing of X and Y

chromosomes is more adversely affected than that of autosomes during the prophase I in male

hybrids between Campbell's dwarf hamster and the Djungarian hamster [57]. A recent work has

clarified that the autosomes of male hybrids between these hamster species undergo paring and

recombination as normally as their parental forms do, but the heterochromatic arms of the X and

Y chromosomes show a high frequency of asynapsis and recombination failure [58]. Another

study has demonstrated a high rate of synaptic aberrations in multiple chromosomes of male

hybrids between two chromosome races of the common shrew [59]. It has been proposed that

asynapsis of heterospecific chromosomes in prophase I may provide a recurrently evolving

trigger for the meiotic arrest of interspecific F1 hybrids of mice [8, 16].

The normal mechanics of meiotic segregation require that homologous chromosome pairs with

each other because crossovers ensure their proper orientation on the meiotic spindle [60]. All

chromosomes must achieve synapsis in pachynema, complete DNA repair, and disassemble their

synaptonemal complexes in diplonema as homologous chromosomes prepare to segregate [61].

The presence of chiasmata and tension exerted across homologs ensures that cells undergo

reductional segregation [62]. Whenever it occurs, an asynapsis of chromosomes almost

invariably triggers pachytene checkpoint and meiotic breakdown [63]. The genetic determinants

of chromosome pairing and synapsis in both plants and animals are of great interest. In bread

wheat, the *Ph1* locus has the largest effect on preventing homeologous pairing in meiosis [64]. In

mice, a null mutation in the *PRDM9* gene causes chromosome asynapsis, arrest of

spermatogenesis at pachynema, impairment of double-strand break repair, and disrupted sex-

body formation [65]. The *PRDM9*-null phenotype resembles observed univalents and frequent

X-Y dissociation in interspecies hybrids of mice [17]. In fact, infertility in male hybrids has been

rescued with bacterial artificial chromosomes (BACs) carrying the *PRDM9* gene from a strain

with the "fertility" allele [66]. In mice hybrids, *PRDM9*-mediated meiotic arrest causes sterility

by a mechanism which relies on the DNA binding properties of PRDM9 [67]. A recent study has

demonstrated a genetic link between meiotic recombination and hybrid male sterility [12]. It has

been suggested that within species there may be selection to maintain sequence homozygosity

for meiosis genes because their divergence can lead to reproductive isolation through failures of

double-strand break formation and synapsis in hybrids [64].

Our data from male hybrids of the crosses ♀ *An. merus* × ♂ *An. coluzzii/An. gambiae* suggest

that meiosis is unaffected until diplonema when autosomes pair and form chiasmata. However,

the downstream processes of meiosis in F1 hybrids are abnormal probably due to the failure of X

and Y chromosomes to synapse. It is possible that the absence of chiasmata between the sex

chromosomes causes failure of reductional segregation in the mosquito hybrids. Unlike hybrids

of mice, primary spermatocytes in the mosquito hybrids do not undergo arrest in meiosis I.

Instead, they continue dividing by mitosis, in which sister chromatids move to the opposite poles

of the dividing cell (Fig. 7). This meiosis-mitosis switch is likely caused by the bi-orientation

instead of the normal mono-orientation of sister kinetochores, in which tension across the

centromere regions and proteins shugoshins plays a key role [53, 68, 69]. It is known that

kinetochore-microtubules attachment errors are more commonly found in meiosis I than in

mitosis [70, 71]. A mathematical model explains why kinetochore-microtubules attachment

errors occur more frequently in the first meiotic division than in mitosis. The model suggests that

the gradual increase of microtubules may help turn off the spindle assembly checkpoint in

meiosis I leading to chromosome mis-segregation errors [72]. In mosquito hybrids, change in the

orientation of sister kinetochores may be caused by reduced tension across the centromeres of

homologous chromosomes due to chromatin decompaction. Changes in chromatin condensation

have been documented in interspecies hybrids of diverse groups of organisms [73, 74]. In a study

of F1 hybrids between *Arabidopsis thaliana* and *A. lyrata*, the *A. thaliana* chromatin became

more compact, whereas the *A. lyrata* chromatin became moderately less compact [74]. An

intriguing spatial arrangement model suggests that that Y chromosome-linked variation may alter

spatial position and packaging of other chromosomes in the nucleus [75]. Chromosome

condensation and spatial position could also be affected in mosquito hybrids by improper

function of condensins. A study of *Drosophila* male meiosis demonstrated that condensin II

subunits, Cap-H2 and Cap-D3, are required to promote chromosome territory formation in

primary spermatocyte nuclei. Moreover, anaphase I is abnormal in Cap-H2 mutants as chromatin

bridges are formed between segregating heterologous and homologous chromosomes [76]. Thus,

the interplay between the two phenotypes in mosquito hybrids—sex chromosome unpairing and

chromatin decompaction—may result in failing a reductional meiotic division and proceeding to

an equational mitotic division.

Evolution of spermatogenesis errors in interspecies hybrids

Reciprocal crosses in malaria mosquitoes and other organisms usually produce hybrids with

sterility phenotypes of different degrees of severity [20, 30, 50]. This observation suggests that

multiple stages of postzygotic isolation exist based on malfunction of the spermatogenesis. At

the early stages of evolution, germline-specific genes still express normally and meiosis proceeds

until metaphase I and then switches to an equational anaphase producing diploid sperm cells in

hybrids. In this case, meiotic errors are characterized by the dramatically decreased pairing

between sex chromosomes and insufficient chromatin condensation, which are seen in sterile

hybrids from the ♀*An. merus* × ♂*An. coluzzii/An. gambiae* crosses (Fig.s 3, 7). The observed

incidental chromosome rearrangements in mosquito hybrids (Fig. 5) support the notion that

genome instability is a common characteristic trait of hybrid incompatibility that may be

associated with increased transposable element activity, ectopic recombination, and double-

strand DNA breaks [77]. A recent work demonstrated that biogenesis of Piwi-interacting RNA

(piRNA) is enhanced in testes of hybrids between *Drosophila buzzatii* and *D. koepferae*. The

study argues that interspecies hybridization causes a genomic stress that can activate the piRNA

response pathway to counteract transposable element deregulation [78]. Because piRNAs

predominantly target long terminal repeats (LTR) retrotransposons in both *Anopheles* and

Drosophila [79], future studies of the piRNA expression in mosquito hybrids may identify

common and specific responses to the genomic stress in dipteran insect species. As species

continue to diverge, meiotic errors become more prominent, and new hybrid phenotypes appear

as has been seen, for example, in sterile male hybrids from the ♀*D. pseudoobscura* × ♂*D.*

persimilis cross [20, 21]. At this stage of postzygotic isolation, an abnormal anaphase is

characterized by unequal segregation of chromosomes resulting in spermatids with unbalanced

chromosome content. The more advanced interspecies divergence of meiosis is manifested by

the unpairing of the most chromosomes and by the malfunction of the abnormally elongated

spindle in sterile male hybrids [20, 21]. The next stage of evolution is spermatogenic arrest in

early meiosis I with disrupted homoeologous chromosome pairing, meiotic sex chromosome

inactivation, and increased apoptosis of spermatocytes as seen in mice hybrids [16-18].

Interestingly, chromosome unpairing and asynapsis in male hybrids stand out as common

spermatogenic phenotypes associated with sterility. It has been hypothesized that nongenic

repetitive sequences, as the fastest diverging component of the genome, may facilitate asynapsis

in hybrids, thus, representing suitable candidates for a Dobzhansky-Muller incompatibility [16].

Findings in *Drosophila* hybrids demonstrate that rapid evolution of heterochromatin may indeed

result in hybrid incompatibilities [73, 77, 80, 81]. Also, the centromere drive model has been

proposed to explain how paired chromosomes at meiosis I can be subject to nondisjunction

leading to infertility in *Drosophila* males [82, 83]. Accordingly, incompatibilities between

rapidly evolving centromeric components of emerging species, such as co-evolving the CENP-A

histone variant and its chaperone CAL1 [84], may account for species incompatibility between

centromeric histones and for postzygotic reproductive isolation. Finally, spermatogenic

abnormalities in hybrids can happen even before meiosis starts. A spermatogenic arrest at the

premeiotic stage is characterized by the repression of germline-specific genes and by the lack of

spermatocytes or spermatids in degenerate testes of hybrid males. These phenotypes have been

observed in sterile hybrids from the ♀*D. mauritiana* × ♂*D. sechellia* cross [19] and in sterile

hybrids from the ♀*An. coluzzii/An. gambiae* × ♂*An. merus* crosses (our study). Thus, cytological

analyses of spermatogenesis in interspecies hybrids can associate sterility phenotypes with early

or advance stages of speciation. Such studies of interspecies hybrids from diverse groups of

organisms may highlight general patterns and mechanisms in the origin and evolution of

postzygotic isolation.

Conclusions

Hybrid male sterility contributes to speciation by restricting gene flow between related taxa at the

beginning stages of postzygotic isolation. Charles Darwin in the chapter "Hybridism" of his

Origin of Species rightly argued that hybrid sterility ''is not a specially endowed quality, but is

incidental on other acquired differences [85].'' Identification of the cellular and molecular

differences acquired during the early stages of postzygotic isolation between species is crucial to

explaining both speciation and normal fertility function. Studies of sterility in male hybrids

between recently evolved species offer opportunities to identify developmental errors that arise

early in speciation. The cross between a female *An. merus* and a male *An. gambiae* or *An.*

coluzzii produces sterile hybrid males with spermatogenic abnormalities in the first meiotic

division. Our study uncovered differences in chromosome behaviors between pure *Anopheles*

species and their hybrids. The obtained data suggest that the meiotic abnormalities in hybrid

males stem from the unpairing of the sex chromosomes and chromatin decondensation. The

meiotic abnormalities also include unpairing of sex chromosomes, chromatin decompaction, and,

in some cases, *de novo* genome rearrangements. Therefore, these malaria mosquitoes represent a

great new system for studying a genetic basis and molecular mechanisms of species

incompatibilities at early stages of postzygotic reproductive isolation.

Knowledge of the mechanisms of reproductive isolation in *Anopheles* has important implications

not only for evolutionary biology but also for malaria control. The success of malaria

transmission highly depends on the rate of mosquito reproduction. The development of novel

approaches to control the reproductive output of mosquitoes must include the understanding of

how fertility is regulated. Determining the mechanisms of sterility in male hybrids between

closely related species of mosquitoes can also empower the sterile insect technique. In this

respect, our results shed new light on the cellular processes and possible mechanisms in

spermatogenesis that first appear when fertility is affected.

Methods

Mosquito maintenance and crossing experiments

The laboratory colonies of *An. gambiae* ZANU (MRA-594), *An. coluzzii* MOPTI (MRA-763),

An. coluzzii MALI (MRA-860), and *An. merus* MAF (MRA-1156) were obtained from the

Biodefense and Emerging Infections Research Resources Repository (BEI). Authentication of

the species was performed by a cytogenetic analysis and by PCR diagnostics [86, 87].

Mosquitoes were reared at $27\pm1°C$, with 12-hour photoperiod and $70\pm5\%$ relative humidity.

Larvae were fed fish food, and adult mosquitoes were fed 1% sugar water. To induce

oviposition, females were fed defibrinated sheep blood (Colorado Serum Co., Denver, Colorado,

USA) using artificial blood feeders. To perform interspecies crosses, male and female pupae

were separated to guarantee virginity of adult mosquitoes. We differentiated males and females

at the pupal stage using sex-specific differences in the shape of their terminalia [88]. After the

emergence of adults, crossing experiments were performed by combining 30 females and 15

males in one cage. Five days after random mating, the females were fed sheep blood. Two days

later, an egg dish, covered with moist filter paper to keep the eggs from drying out, was put into

the cage. Backcrosses of F1 males and parental females were done using a similar method. At

least two blood meals were fed to females with three repeats of each crosses.

Male gonad and sperm observation

Male gonads were dissected and photographed using an Olympus SZ 61 stereoscopy microscope

(Olympus, Tokyo, Japan) with an Olympus Q-Color 5 digital camera (Olympus, Tokyo, Japan).

We observed the testes of 30 males and took pictures of the testes of five males from each cross.

For sperm observation, testes were mounted in 20 μl sperm assay buffer containing 4 mM KCl,

1.3 mM $CaCl_2$, 145 mM NaCl, 5 mM D-glucose, 1 mM $MgCl_2$, and 10 mM 4-(2-hydroxyethyl)-

1-piperazineethanesulfonic acid (Hepes) [89]. After gently covering testes with a coverslip, they

were crushed, and sperm motility was observed under a phase contrast microscope BX 41

(Olympus, Tokyo, Japan). A movie of sperm motility for at least five males of hybrids from each

cross and five males of pure species was recorded using a digital camera UC90 (Olympus,

Tokyo, Japan).

Chromosome preparation

Testes with male accessory glands were dissected from male pupae and 0-12 hours-old adults in

0.075% potassium chloride (KCl) hypotonic solution on a frosted glass slide (Thermo Fisher

Scientific, Waltham, MA, USA) under an Olympus SZ 61 dissecting microscope (Olympus,

Tokyo, Japan). To observe meiotic chromosomes, male accessory glands and other tissues were

removed, and only testes were left on the slide. Immediately after dissection, a drop of 50%

propionic acid was added to the testes, and they were covered with a 22×22-mm coverslip

(Thermo Fisher Scientific, Waltham, MA, USA). Preparations were tapped using the flat rubber

end of a pencil and observed under an Olympus CX41 phase-contrast microscope (Olympus,

Tokyo, Japan). The slides were frozen in liquid nitrogen, and a sharp razor was used to take off

the coverslips. The preparations were placed in 50% ethanol chilled at -20℃ for a minimum of 2

hours. Later, serial dehydrations were performed in 70%, 90%, and 100% ethanol for 5 min each

at room temperature (RT). Subsequently, the best preparations from at least 10 slides/individuals

with desired meiotic stages were chosen for further studies.

DNA probe labeling

Six DNA probes were used for FISH in this study (Table 1): retroelement *zanzibar* specific to the

Y chromosome of *An. gambiae* and *An. coluzzii*, 18S rDNA specific to the X chromosome of *An.*

gambiae and *An. coluzzii* but hybridized to both X and Y chromosomes of *An. merus,* satellite

AgY53B derived from the AgY53B/AgY477 satellite array and hybridized to X and Y

chromosomes of all three species [44]*,* satellite AgY477-AgY53B junction region specific to the

Y chromosome of *An. gambiae* and *An. coluzzii*, a satellite from Contig_240 identified from *An.*

gambiae sequencing data (SRS667972, SRR1509742, SRR1508169) [44] using the Redkmer

pipeline [90] and specific to the X chromosomes of *An. coluzzii* and *An. merus*, and a satellite

tandem-42345 [91] specific to autosomal pericentric heterochromatin in *An. coluzzii* and to

pericentric heterochromatin of all chromosomes in *An. merus*. The 18S rDNA, AgY53B and

zanzibar probes were labeled by Cyanine3 (Cy3) or Cyanine5 (Cy5) fluorochromes using PCR

with genomic DNA as a template. Genomic DNA was extracted using DNeasy Blood & Tissue

Kits (Qiagen, Hilden, Germany) from virgin males of *An. gambiae* ZANU or *An. coluzzii*

MOPTI for labeling *zanzibar*, virgin males of *An. merus* MAF for labeling satellite AgY53B,

and virgin females of *An. gambiae* or *An. coluzzii* for labeling 18S rDNA. Each 25 µl of a PCR

mix consisted of 1-µl genomic DNA, 12.5-µl ImmoMix^{TM} 2x reaction mix (Bioline USA Inc.,

Taunton, MA, USA), 1 µl of 10-µM forward and reverse primers, and water. PCR labeling was

performed in the Mastercycler® pro PCR thermocycler (Eppendorf, Hamburg, Germany)

starting with a 95°C incubation for 10 min followed by 35 cycles of 95°C for 30 sec, 52°C for 30

sec, 72°C for 45 sec; 72°C for 5 min, and a final hold at 4°C. Oligonucleotides with 3'-end-

labeling by Cy5 or Cy3 fluorochromes (Sigma-Aldrich, St. Louis, MO, USA) were designed

based on sequences of the satellite AgY477-AgY53B junction region, satellite from Contig_240,

and satellite tandem-42345.

FISH

FISH was performed as previously described [92, 93]. Briefly, slides with good preparations

were treated with 0.1-mg/ml RNase at 37℃ for 30 min. After washing with 2×SSC for 5 min

twice, slides were digested with 0.01% pepsin and 0.037% HCl solution for 5 min at 37°C. After

washing slides in 1×PBS for 5 min at RT two times, preparations were fixed in 3.7%

formaldehyde for 10 min at RT. Slides were then washed in 1×PBS and dehydrated in a series of

70%, 80%, and 100% ethanol for 5 min at RT. Then 10 μl of probes were mixed, added to the

preparations, and incubated at 37℃ overnight. After washing slides in 1×SSC at 60℃ for 5 min,

4×SSC/NP40 solution at 37℃ for 10 min, and 1×PBS for 5 min at RT, preparations were

counterstained with a DAPI-antifade solution (Life Technologies, Carlsbad, CA, USA) and kept

in the dark for at least 2 hours before visualization with a fluorescent microscope.

Cytogenetic analyses

Cytogenetic analyses of meiotic and mitotic stages and of behaviors of FISH-labeled

chromosomes were performed on at least 10 male individuals of each pure species and of each

hybrid. To visualize and photograph meiotic chromosomes after FISH, we used an Olympus

BX61 microscope (Olympus, Tokyo, Japan) with a connected camera Olympus U-CMAD3

(Olympus, Tokyo, Japan). Lengths of well-spread 10 metaphase I chromosomes of each species

and of each hybrid were measured using the ruler tool in Adobe Photoshop CS6 (Adobe Inc., San

Jose, CA, USA). Since the variances for both groups were equal ($s_{max}/s_{min} < 2$), a statistical two-

sample pooled t-test was done with the JMP 13 software (SAS Institute Inc., Cary, NC, USA).

Whole-mount FISH and analysis of chromosome pairing

Testes from one-day-old adults of pure species and hybrids were dissected in 1×PBS solution

and fixed in 3.7% paraformaldehyde in 1×PBS with 0.1% tween-20 (PBST) for 10 min at room

temperature. After incubation with 0.1-mg/ml RNase for 30 min at 37 ℃, testes were penetrated

with 1% triton/0.1M HCl in PBST for 10 min at room temperature. After adding labeled DNA

probes, testes were incubated at 75 ℃ for 5 min (denaturation) and 37 ℃ overnight

(hybridization). Later, testes were washed with 2×SSC and mounted with a DAPI-antifade

solution (Life Technologies, Carlsbad, CA, USA). Testes from 6 individuals of pure species and

of hybrids were scanned and analyzed. Visualization and z-tack 3D scanning were performed on

the whole testes with an interval of 1.25 μm between two optical sections under a 63× oil lens of

Zeiss LSM 880 confocal laser scanning microscope (Carl Zeiss AG, Oberkochen, Germany).

Based on the size of the testis cells and on the limitations of microscope imaging, we chose from

the first to sixteenth optical layers with strong and clearly detected fluorescent signals to analyze

sex-chromosome-pairing events. Specifically, cell nuclei at the early stages of meiotic prophase I

from $4^{th}\pm1$, $8^{th}\pm1$, and $12^{th}\pm1$ optical layers were used to count paring events. A total of 418 and

489 nuclei at the early stages of meiotic prophase I were analyzed for paring and unpairing of the

sex chromosomes in pure species and hybrids, respectively (Additional file 12: Table S2). The

percentage of the numbers of cells with pairing and no paring of sex chromosomes was used to

compare parental species and hybrids. Since the variances for both groups were equal ($s_{max}/s_{min} <$

2), a statistical two-sample pooled *t*-test was done using the JMP 13 software (SAS Institute Inc.,

Cary, NC, USA).

RNA extraction and RT-PCR

RNA was extracted from 40 abdomens (only distal segments that include testes) and from 20

carcasses of 1- to 2-day-old virgin adult males of *An. coluzzii* MOPTI, as well as of interspecies

hybrids from crosses ♀*An. coluzzii* MOPTI × ♂*An. merus* and ♀*An. merus* × ♂*An. coluzzii*

MOPTI using a Direct-Zol^{TM} RNA MiniPrep Kit (Zymo Research, Irvine, California, US).

cDNA for selected genes (Table 1) was generated in a 40-μl reaction that contained 8 μl of 5×

first-strand buffer, 4 μl of 0.1-M dithiothreitol (DTT), 2 μl of 50-μM random hexamer primer

(Thermo Fisher Scientific, Waltham, MA, USA), 1 μl of 25-mM dNTP mix solution (Thermo

Fisher Scientific, Waltham, MA, USA), 1 μl of 40-U/μl RNAsin (Promega, Madison, WI, USA),

2 μl of 200-U/μl M-MLV Reverse Transcriptase (Thermo Fisher Scientific, Waltham, MA,

USA), 1ng to 2μg of RNA and water. After incubation at 42°C for 1h, the reaction was stopped

by incubation at 95°C for 5 min. Two-step RT-PCR was performed to analyze the gene

expression. Each 25-μl PCR mix consisted of 1-μl cDNA, 12.5-μl ImmoMix^{TM} 2× reaction-mix

(Bioline USA Inc., Taunton, MA, USA), 1 μl of 10-μM forward and reverse primers, and water.

PCR was performed in the Mastercycler® pro PCR thermocycler (Eppendorf, Hamburg,

Germany) starting with a 95°C incubation for 10 min followed by 35 cycles of 95°C for 30 sec,

57°C for 30 sec, 72°C for 30 sec; 72°C for 10 min; and a final hold at 4°C. Amplification

products were visualized in a 1-1.5% agarose gel.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and its

additional files.

Authors' contributions

589   IVS conceived the study. JL conducted mosquito maintenance and crosses. JL and IVS

590   performed mosquito dissection and sperm motility assays. JL conducted FISH, microscopy, and

591   RT-PCR. JL and IVS did data analysis. IVS led the writing of the manuscript with critical

592   contributions from JL. All authors read and approved the final manuscript.

Competing interests

595   The authors declare that they have no competing interests.

Acknowledgments

598   We thank the Malaria Research and Reference Reagent Resource (MR4) at the BEI for providing

599   us the laboratory colonies of malaria mosquitoes and Melissa Wade for proofreading the text.

References

1.   Powell JR: **Progress and Prospects in Evolutionary Biology**. New York - Oxford:

Oxford University Press; 1997.

2.   Coyne JA, Orr HA: **Speciation**. Sunderland, Mass.: Sinauer Associates; 2004.

3.   Larson EL, VaLnderpool D, Sarver BAJ, Callahan C, Keeble S, Provencio LL, Kessler

MD, Stewart V, Nordquist E, Dean MD *et al*: **The Evolution of Polymorphic Hybrid**

Incompatibilities in House Mice. *Genetics* 2018, **209**(3):845-859.

4.   Good JM, Dean MD, Nachman MW: **A complex genetic basis to X-linked hybrid male**

609   **sterility between two species of house mice**. *Genetics* 2008, **179**(4):2213-2228.

610   5.   White MA, Stubbings M, Dumont BL, Payseur BA: **Genetics and evolution of hybrid**

611   **male sterility in house mice**. *Genetics* 2012, **191**(3):917-934.

612   6.   Fishman L, Stathos A, Beardsley PM, Williams CF, Hill JP: **Chromosomal**

613   **rearrangements and the genetics of reproductive barriers in mimulus (monkey**

614   **flowers)**. *Evolution* 2013, **67**(9):2547-2560.

615   7.   Leppala J, Savolainen O: **Nuclear-cytoplasmic interactions reduce male fertility in**

616   **hybrids of Arabidopsis lyrata subspecies**. *Evolution* 2011, **65**(10):2959-2972.

617   8.   Bhattacharyya T, Reifova R, Gregorova S, Simecek P, Gergelits V, Mistrik M,

618   Martincova I, Pialek J, Forejt J: **X chromosome control of meiotic chromosome**

619   **synapsis in mouse inter-subspecific hybrids**. *PLoS Genet* 2014, **10**(2):e1004088.

620   9.   Dobzhansky T: **Studies on Hybrid Sterility. II. Localization of Sterility Factors in**

621   **Drosophila Pseudoobscura Hybrids**. *Genetics* 1936, **21**(2):113-135.

622   10.   Turissini DA, Liu G, David JR, Matute DR: **The evolution of reproductive isolation in**

623   **the Drosophila yakuba complex of species**. *J Evol Biol* 2015, **28**(3):557-575.

624   11.   Hu XS, Filatov DA: **The large-X effect in plants: increased species divergence and**

625   **reduced gene flow on the Silene X-chromosome**. *Mol Ecol* 2016, **25**(11):2609-2619.

626   12.   Balcova M, Faltusova B, Gergelits V, Bhattacharyya T, Mihola O, Trachtulec Z, Knopf

627   C, Fotopulosova V, Chvatalova I, Gregorova S *et al*: **Hybrid Sterility Locus on**

628   **Chromosome X Controls Meiotic Recombination Rate in Mouse**. *PLoS Genet* 2016,

629   **12**(4):e1005906.

13.   Slotman M, Della Torre A, Powell JR: **The genetics of inviability and male sterility in**

hybrids between Anopheles gambiae and An. arabiensis. *Genetics* 2004, **167**(1):275-

287.

14.   Phadnis N, Orr HA: **A Single Gene Causes Both Male Sterility and Segregation**

Distortion in Drosophila Hybrids. *Science* 2009, **323**(5912):376-379.

15.   Ting CT, Tsaur SC, Wu ML, Wu CI: **A rapidly evolving homeobox at the site of a**

hybrid sterility gene. *Science* 1998, **282**(5393):1501-1504.

16.   Bhattacharyya T, Gregorova S, Mihola O, Anger M, Sebestova J, Denny P, Simecek P,

Forejt J: **Mechanistic basis of infertility of mouse intersubspecific hybrids**. *Proc Natl*

Acad Sci U S A 2013, **110**(6):E468-477.

17.   Forejt J, Ivanyi P: **Genetic studies on male sterility of hybrids between laboratory**

and wild mice (Mus musculus L.). *Genet Res* 1974, **24**(2):189-206.

18.   Schwahn DJ, Wang RJ, White MA, Payseur BA: **Genetic Dissection of Hybrid Male**

Sterility Across Stages of Spermatogenesis. *Genetics* 2018.

19.   Kulathinal R, Singh RS: **Cytological Characterization of Premeiotic Versus**

Postmeiotic Defects Producing Hybrid Male Sterility among Sibling Species of the

Drosophila Melanogaster Complex. *Evolution* 1998, **52**(4):1067-1079.

20.   Dobzhansky T: **Studies on hybrid sterility. I. Spermatogenesis in pure and hybrid**

Drosophila pseudoobscura. *ZEITSCHR ZELLJORSCH U MIKROSK ANAT* 1934,

21(2):169-223.

21.   Dobzhansky T: **On the sterility of the interracial hybrids in Drosophila**

pseudoobscura. *P Natl Acad Sci USA* 1933, **19**:397-403.

22.   Hardy RW, Lougheed A, Markow TA: **Reproductive tract and spermatid**

abnormalities of hybrid males from reciprocal crosses between Drosophila

mojavensis and D. arizonae. *Fly (Austin)* 2011, **5**(2):76-80.

23.   Coyne JA: **Genetic-Basis of Male-Sterility in Hybrids between 2 Closely Related**

Species of Drosophila. *P Natl Acad Sci-Biol* 1984, **81**(14):4444-4447.

24.   Masly JP, Jones CD, Noor MA, Locke J, Orr HA: **Gene transposition as a cause of**

hybrid sterility in Drosophila. *Science* 2006, **313**(5792):1448-1450.

25.   Coluzzi M, Sabatini A, della Torre A, Di Deco MA, Petrarca V: **A polytene**

chromosome analysis of the Anopheles gambiae species complex. *Science* 2002,

298(5597):1415-1418.

26.   Coetzee M, Hunt RH, Wilkerson R, Della Torre A, Coulibaly MB, Besansky NJ:

Anopheles coluzzii and Anopheles amharicus, new members of the Anopheles

gambiae complex. *Zootaxa* 2013, **3619**(3):246-274.

27.   Fontaine MC, Pease JB, Steele A, Waterhouse RM, Neafsey DE, Sharakhov IV, Jiang

XF, Hall AB, Catteruccia F, Kakani E *et al*: **Extensive introgression in a malaria**

vector species complex revealed by phylogenomics. *Science* 2015, **347**(6217):42-+.

28.   Thawornwattana Y, Dalquen D, Yang Z: **Coalescent Analysis of Phylogenomic Data**

Confidently Resolves the Species Relationships in the Anopheles gambiae Species

Complex. *Mol Biol Evol* 2018, **35**(10):2512-2527.

29.   Bernardini F, Galizi R, Wunderlich M, Taxiarchi C, Kranjc N, Kyrou K, Hammond A,

Nolan T, Lawniczak MNK, Papathanos PA *et al*: **Cross-Species Y Chromosome**

Function Between Malaria Vectors of the Anopheles gambiae Species Complex.

Genetics 2017, **207**(2):729-740.

30.   Davidson G, Paterson HE, Coluzzi M, Mason GF, Micks DW: **The Anopheles gambiae**

Complex. In: *Genetics of Insect Vectors of Diesease.* Edited by Wright JW, Pal R.

Amsterdam-London-New York: Elsevier Publishing Company; 1967: 211-249.

31.   Presgraves DC, Orr HA: **Haldane's rule in taxa lacking a hemizygous X**. *Science* 1998,

282(5390):952-954.

32.   Curtis CF: **The mechanism of hybrid male sterility from crosses in the Anopheles**

gambiae and Glossina morsitans complexes. In: *Recent Developments in the Genetics*

of Disease Vectors. Edited by Steiner WM, Tabachnick WJ, Rai KS, Narang S.

Champaign, IL: Stipes Publishing Company; 1982: 290-312.

33.   Pombi M, Kengne P, Gimonneau G, Tene-Fossog B, Ayala D, Kamdem C, Santolamazza

F, Guelbeogo WM, Sagnon N, Petrarca V *et al*: **Dissecting functional components of**

reproductive isolation among closely related sympatric species of the Anopheles

gambiae complex. *Evol Appl* 2017, **10**(10):1102-1120.

34.   **World malaria report**. In. Geneva, Switzerland: World Health Organization; 2018: 165.

35.   Dame DA, Curtis CF, Benedict MQ, Robinson AS, Knols BG: **Historical applications**

of induced sterilisation in field populations of mosquitoes. *Malar J* 2009, **8 Suppl**

2:S2.

36.   Hammond A, Galizi R, Kyrou K, Simoni A, Siniscalchi C, Katsanos D, Gribble M, Baker

D, Marois E, Russell S *et al*: **A CRISPR-Cas9 gene drive system targeting female**

reproduction in the malaria mosquito vector Anopheles gambiae. *Nat Biotechnol*

2016, **34**(1):78-83.

37.   Macias VM, Ohm JR, Rasgon JL: **Gene Drive for Mosquito Control: Where Did It**

Come from and Where Are We Headed? *Int J Env Res Pub He* 2017, **14**(9).

38.   Kyrou K, Hammond AM, Galizi R, Kranjc N, Burt A, Beaghton AK, Nolan T, Crisanti

699   A: **A CRISPR-Cas9 gene drive targeting doublesex causes complete population**

700   **suppression in caged Anopheles gambiae mosquitoes**. *Nat Biotechnol* 2018,

701   **36**(11):1062-1066.

702   39.   Gantz VM, Jasinskiene N, Tatarenkova O, Fazekas A, Macias VM, Bier E, James AA:

703   **Highly efficient Cas9-mediated gene drive for population modification of the**

704   **malaria vector mosquito Anopheles stephensi**. *Proc Natl Acad Sci U S A* 2015,

705   **112**(49):E6736-6743.

706   40.   Hammond AM, Kyrou K, Bruttini M, North A, Galizi R, Karlsson X, Kranjc N, Carpi

707   FM, D'Aurizio R, Crisanti A *et al*: **The creation and selection of mutations resistant to**

708   **a gene drive over multiple generations in the malaria mosquito**. *PLoS Genet* 2017,

709   **13**(10):e1007039.

710   41.   Diabate A, Dabire RK, Millogo N, Lehmann T: **Evaluating the effect of postmating**

711   **isolation between molecular forms of Anopheles gambiae (Diptera: Culicidae)**. *J*

712   *Med Entomol* 2007, **44**(1):60-64.

713   42.   Aboagye-Antwi F, Alhafez N, Weedall GD, Brothwood J, Kandola S, Paton D, Fofana A,

714   Olohan L, Betancourth MP, Ekechukwu NE *et al*: **Experimental swap of Anopheles**

715   **gambiae's assortative mating preferences demonstrates key role of X-chromosome**

716   **divergence island in incipient sympatric speciation**. *PLoS Genet* 2015,

717   **11**(4):e1005141.

718   43.   Krzywinski J, Sangare D, Besansky NJ: **Satellite DNA from the Y chromosome of the**

719   **malaria vector Anopheles gambiae**. *Genetics* 2005, **169**(1):185-196.

44.   Hall AB, Papathanos PA, Sharma A, Cheng C, Akbari OS, Assour L, Bergman NH,

Cagnetti A, Crisanti A, Dottorini T *et al*: **Radical remodeling of the Y chromosome in**

a recent radiation of malaria mosquitoes. *Proc Natl Acad Sci U S A* 2016,

113(15):E2114-2123.

45.   Krzywinska E, Krzywinski J: **Analysis of expression in the Anopheles gambiae**

developing testes reveals rapidly evolving lineage-specific genes in mosquitoes. *BMC*

Genomics 2009, **10**:300.

46.   Balabanidou V, Kampouraki A, MacLean M, Blomquist GJ, Tittiger C, Juarez MP,

Mijailovsky SJ, Chalepakis G, Anthousi A, Lynd A *et al*: **Cytochrome P450 associated**

with insecticide resistance catalyzes cuticular hydrocarbon production in Anopheles

gambiae. *Proc Natl Acad Sci U S A* 2016, **113**(33):9268-9273.

47.   Catteruccia F, Crisanti A, Wimmer EA: **Transgenic technologies to induce sterility**.

Malar J 2009, **8 Suppl 2**:S7.

48.   Kemphues KJ, Kaufman TC, Raff RA, Raff EC: **The testis-specific beta-tubulin**

subunit in Drosophila melanogaster has multiple functions in spermatogenesis. *Cell*

1982, **31**(3 Pt 2):655-670.

49.   Papathanos PA, Windbichler N, Menichelli M, Burt A, Crisanti A: **The vasa regulatory**

region mediates germline expression and maternal transmission of proteins in the

malaria mosquito Anopheles gambiae: a versatile tool for genetic control strategies.

BMC Mol Biol 2009, **10**:65.

50.   Good JM, Handel MA, Nachman MW: **Asymmetry and polymorphism of hybrid male**

sterility during the early stages of speciation in house mice. *Evolution* 2008, **62**(1):50-

65.

51.   Fontoura NG, Araki AS, Azevedo RV, Galardo AKR, Peixoto AA, Lima JBP: **Hybrid**

sterility in crosses between two Brazilian sibling species of the Anopheles albitarsis

complex. *Parasite Vector* 2014, **7**.

52.   Suwannamit S, Baimai V, Otsuka Y, Saeung A, Thongsahuan S, Tuetun B,

Apiwathnasorn C, Jariyapan N, Somboon P, Takaoka H *et al*: **Cytogenetic and**

molecular evidence for an additional new species within the taxon Anopheles

barbirostris (Diptera: Culicidae) in Thailand. *Parasitol Res* 2009, **104**(4):905-918.

53.   McKee BD, Yan R, Tsai JH: **Meiosis in male Drosophila**. *Spermatogenesis* 2012,

2(3):167-184.

54.   Bonaccorsi S, Gatti M: **Drosophila Male Meiosis**. *Methods Mol Biol* 2017, **1471**:277-

288.

55.   Hawley RS: **Meiosis: How male flies do meiosis**. *Current Biology* 2002, **12**(19):R660-

R662.

56.   Chang CH, Larracuente AM: **Heterochromatin-Enriched Assemblies Reveal the**

Sequence and Organization of the Drosophila melanogaster Y Chromosome.

Genetics 2018.

57.   Ishishita S, Tsuboi K, Ohishi N, Tsuchiya K, Matsuda Y: **Abnormal pairing of X and Y**

sex chromosomes during meiosis I in interspecific hybrids of Phodopus campbelli

and P. sungorus. *Sci Rep* 2015, **5**:9435.

58.   Bikchurina TI, Tishakova KV, Kizilova EA, Romanenko SA, Serdyukova NA,

Torgasheva AA, Borodin PM: **Chromosome Synapsis and Recombination in Male-**

Sterile and Female-Fertile Interspecies Hybrids of the Dwarf Hamsters (Phodopus,

Cricetidae). *Genes (Basel)* 2018, **9**(5).

59. Belonogova NM, Polyakov AV, Karamysheva TV, Torgasheva AA, Searle JB, Borodin

PM: **Chromosome Synapsis and Recombination in Male Hybrids between Two**

Chromosome Races of the Common Shrew (Sorex araneus L., Soricidae,

Eulipotyphla). *Genes (Basel)* 2017, **8**(10).

60. Miller MP, Amon A, Unal E: **Meiosis I: when chromosomes undergo extreme**

makeover. *Curr Opin Cell Biol* 2013, **25**(6):687-696.

61. Subramanian VV, Hochwagen A: **The meiotic checkpoint network: step-by-step**

through meiotic prophase. *Cold Spring Harb Perspect Biol* 2014, **6**(10):a016675.

62. Miyazaki S, Kim J, Sakuno T, Watanabe Y: **Hierarchical Regulation of Centromeric**

Cohesion Protection by Meikin and Shugoshin during Meiosis I. *Cold Spring Harb*

Symp Quant Biol 2017, **82**:259-266.

63. Bolcun-Filas E, Schimenti JC: **Genetics of meiosis and recombination in mice**. *Int Rev*

Cell Mol Biol 2012, **298**:179-227.

64. Bomblies K, Higgins JD, Yant L: **Meiosis evolves: adaptation to external and internal**

environments. *New Phytol* 2015, **208**(2):306-323.

65. Hayashi K, Yoshida K, Matsui Y: **A histone H3 methyltransferase controls epigenetic**

events required for meiotic prophase. *Nature* 2005, **438**(7066):374-378.

66. Mihola O, Trachtulec Z, Vlcek C, Schimenti JC, Forejt J: **A mouse speciation gene**

encodes a meiotic histone H3 methyltransferase. *Science* 2009, **323**(5912):373-375.

67. Davies B, Hatton E, Altemose N, Hussin JG, Pratto F, Zhang G, Hinch AG, Moralli D,

Biggs D, Diaz R *et al*: **Re-engineering the zinc fingers of PRDM9 reverses hybrid**

sterility in mice. *Nature* 2016, **530**(7589):171-176.

68.   Gutierrez-Caballero C, Cebollero LR, Pendas AM: **Shugoshins: from protectors of**

cohesion to versatile adaptors at the centromere. *Trends Genet* 2012, **28**(7):351-360.

69.   Pinto BS, Orr-Weaver TL: **Drosophila protein phosphatases 2A B' Wdb and Wrd**

regulate meiotic centromere localization and function of the MEI-S332 Shugoshin.

Proc Natl Acad Sci U S A 2017, **114**(49):12988-12993.

70.   Kitajima TS, Ohsugi M, Ellenberg J: **Complete kinetochore tracking reveals error-**

prone homologous chromosome biorientation in mammalian oocytes. *Cell* 2011,

146(4):568-581.

71.   Jones KT, Lane SI: **Molecular causes of aneuploidy in mammalian eggs**. *Development*

2013, **140**(18):3719-3730.

72.   Saka Y, Giuraniuc CV, Ohkura H: **Accurate chromosome segregation by probabilistic**

self-organisation. *Bmc Biol* 2015, **13**:65.

73.   Blum JA, Bonaccorsi S, Marzullo M, Palumbo V, Yamashita YM, Barbash DA, Gatti M:

The Hybrid Incompatibility Genes Lhr and Hmr Are Required for Sister

Chromatid Detachment During Anaphase but Not for Centromere Function.

Genetics 2017, **207**(4):1457-1472.

74.   Zhu W, Hu B, Becker C, Dogan ES, Berendzen KW, Weigel D, Liu C: **Altered**

chromatin compaction and histone methylation drive non-additive gene expression

in an interspecific Arabidopsis hybrid. *Genome Biol* 2017, **18**(1):157.

75.   Francisco FO, Lemos B: **How do y-chromosomes modulate genome-wide epigenetic**

States: genome folding, chromatin sinks, and gene expression. *J Genomics* 2014,

2:94-103.

76. Hartl TA, Sweeney SJ, Knepler PJ, Bosco G: **Condensin II resolves chromosomal**

associations to enable anaphase I segregation in Drosophila male meiosis. *PLoS*

Genet 2008, **4**(10):e1000228.

77. Dion-Cote AM, Barbash DA: **Beyond speciation genes: an overview of genome**

stability in evolution and speciation. *Curr Opin Genet Dev* 2017, **47**:17-23.

78. Romero-Soriano V, Modolo L, Lopez-Maestre H, Mugat B, Pessia E, Chambeyron S,

Vieira C, Garcia Guerreiro MP: **Transposable Element Misregulation Is Linked to the**

Divergence between Parental piRNA Pathways in Drosophila Hybrids. *Genome Biol*

Evol 2017, **9**(6):1450-1470.

79. George P, Jensen S, Pogorelcnik R, Lee J, Xing Y, Brasset E, Vaury C, Sharakhov IV:

Increased production of piRNAs from euchromatic clusters and genes in Anopheles

gambiae compared with Drosophila melanogaster. *Epigenetics Chromatin* 2015, **8**:50.

80. Ferree PM, Barbash DA: **Species-specific heterochromatin prevents mitotic**

chromosome segregation to cause hybrid lethality in Drosophila. *PLoS Biol* 2009,

7(10):e1000234.

81. Bayes JJ, Malik HS: **Altered heterochromatin binding by a hybrid sterility protein in**

Drosophila sibling species. *Science* 2009, **326**(5959):1538-1541.

82. Henikoff S, Ahmad K, Malik HS: **The centromere paradox: Stable inheritance with**

rapidly evolving DNA. *Science* 2001, **293**(5532):1098-1102.

83. Malik HS, Henikoff S: **Adaptive evolution of cid, a centromere-specific histone in**

drosophila. *Genetics* 2001, **157**(3):1293-1298.

84.   Rosin L, Mellone BG: **Co-evolving CENP-A and CAL1 Domains Mediate**

Centromeric CENP-A Deposition across Drosophila Species. *Developmental Cell*

2016, **37**(2):136-147.

85.   Darwin C: **On the origin of species by means of natural selection or, The**

Preservation of favoured races in the struggle for life. In. London: Electric Book Co.,;

2001: 1 online resource (422 p.).

86.   Scott JA, Brogdon WG, Collins FH: **Identification of single specimens of the**

Anopheles gambiae complex by the polymerase chain reaction. *Am J Trop Med Hyg*

1993, **49**(4):520-529.

87.   Fanello C, Santolamazza F, della Torre A: **Simultaneous identification of species and**

molecular forms of the Anopheles gambiae complex by PCR-RFLP. *Med Vet*

Entomol 2002, **16**(4):461-464.

88.   Benedict M, Dotson EM: **Methods in Anopheles Research**. In.:

https://www.beiresources.org/Portals/2/VectorResources/2016%20Methods%20in%20An

opheles%20Research%20full%20manual.pdf; 2015: 408.

89.   Pitts RJ, Liu C, Zhou X, Malpartida JC, Zwiebel LJ: **Odorant receptor-mediated sperm**

activation in disease vector mosquitoes. *Proc Natl Acad Sci U S A* 2014, **111**(7):2566-

2571.

90.   Papathanos PA, Windbichler N: **Redkmer: An Assembly-Free Pipeline for the**

Identification of Abundant and Specific X-Chromosome Target Sequences for X-

Shredding by CRISPR Endonucleases. *The CRISPR Journal* 2018, **1**(2):88-98.

91.   Melters DP, Bradnam KR, Young HA, Telis N, May MR, Ruby JG, Sebra R, Peluso P,

Eid J, Rank D *et al*: **Comparative analysis of tandem repeats from hundreds of**

species reveals unique insights into centromere evolution. *Genome Biol* 2013,

14(1):R10.

92.   Timoshevskiy VA, Sharma A, Sharakhov IV, Sharakhova MV: **Fluorescent in situ**

857   **Hybridization on Mitotic Chromosomes of Mosquitoes**. *J Vis Exp* 2012(67).

858   93.   Sharakhova MV, George P, Timoshevskiy V, Sharma A, Peery A, Sharakhov IV:

859   **Mosquitoes (Diptera)**. In: *Protocols for Cytogenetic Mapping of Arthropod Genomes.*

860   Edited by Sharakhov IV. Boca Raton, FL: CRC Press, Taylor & Francis Group; 2015:

861   93-170.

Figure Legends

Fig. 1. Morphology and cell content of testes from 2-day-old adult males. (A) Whole testis,

squashed testis, and mature spermatozoa in crushed testis from *An. coluzzii* MOPTI. (B) Normal-

like whole testis, squashed testis, and immature spermatozoa and spermatids in crushed testis

from an F1 hybrid of the cross ♀*An. merus* × ♂*An. coluzzii* MOPTI. (C) Underdeveloped whole

testis, squashed testis, and undifferentiated cells in crushed testis from an F1 hybrid of the cross

♀*An. coluzzii* MOPTI × ♂*An. merus.* Vertical scale bars – 50 and 20 μm.

Fig. 2. Chromosome behavior during meiosis in testes of *An. gambiae.* X chromosomes are

labeled with 18S rDNA (red), and Y chromosomes are labeled with retrotransposon *zanzibar*

(green). Chromosomes are counterstained with DAPI (blue). Scale bar – 5 μm.

Fig. 3. Chromosome behavior during meiosis in testes of interspecies hybrids. (A) ♀*An.*

merus × ♂*An. coluzzii* MALI. (B) ♀*An. merus* × ♂*An. coluzzii* MOPTI. (C) ♀*An. merus* × ♂*An.*

gambiae ZANU. X chromosomes are labeled with 18S rDNA (red), and Y chromosomes are

labeled with retrotransposon *zanzibar* (green). Chromosomes are counterstained with DAPI

(blue). Scale bar – 5 μm.

Fig. 4. Chromosomal and molecular abnormalities in interspecies hybrids. (A) Left panel:

Lengths of metaphase chromosomes in *An. gambiae* ZANU males (filled boxplots) and in F1

hybrid males from the cross ♀*An. merus* × ♂*An. gambiae* ZANU (open boxplots). Right panel:

Ratios of the lengths of chromosomes between F1 hybrids and *An. gambiae* ZANU. X and Y –

sex chromosomes. 2 and 3 – autosomes. Statistical significance was assessed with a two-sample

pooled *t*-test. (B) Confocal images of primary spermatocytes with FISH to demonstrate pairing

and unpairing of X (red) and Y (green) chromosomes in *An. coluzzii* MOPTI males and F1

hybrid males from the ♀*An. merus* × ♂*An. coluzzii* MOPTI cross. (C) Percentage of the X and Y

chromosome pairing in primary spermatocytes of six *An. coluzzii* MOPTI males and six F1

hybrid males from the ♀*An. merus* × ♂*An. coluzzii* MOPTI cross. Statistical significance was

assessed with a two-sample pooled *t*-test. (D) Expression of meiotic (*Ams, mts, Dzip1l, β2-*

tubulin) and pre-meiotic (*vasa*) germline-specific genes in testes of *An. coluzzii*, F1 hybrids of

♀*An. merus* × ♂*An. coluzzii* MOPTI (F1M×C), and F1 hybrids of ♀*An. coluzzii* MOPTI × ♂*An.*

merus (F1C×M) analyzed by RT-PCR. *AgS7* – an endogenous control gene. R – reproductive

tissues, B – rest of mosquito body.

Fig. 5. Genomic rearrangements of meiotic chromosomes in testes of interspecies hybrids.

(A) A translocation of the fragment of the *An. merus* X chromosome to the *An. coluzzii* autosome

2 in a hybrid from the ♀*An. merus* × ♂*An. coluzzii* MALI cross. The red arrow points to the

translocated X-chromosome ribosomal locus labelled by 18S rDNA (red). (B) A duplication of

the segment involving the rDNA locus within the *An. merus* X chromosome in a hybrid from the

cross ♀*An. merus* × ♂*An. gambiae* ZANU. The red arrow indicates the new positions of the

segment with the ribosomal locus labelled by 18S rDNA (red). Chromosomes are counterstained

with DAPI.

Fig. 6. Schematic representation of testis development in pure species and reciprocal F1

hybrids of the *An. gambiae* complex.

Fig. 7. Scheme of male meiosis in pure species and interspecies hybrids of the *An. gambiae*

complex. Unlike pure species, X and Y chromosomes in meiotic prophase I of hybrids are

largely unpaired. Instead of a reductional division, primary spermatocytes in interspecies hybrids

undergo an equational mitotic division leading to diploid nonmotile sperms.

Supporting Information

Additional file 1: Movie S1. Sperm motility in a crushed testis from a 2-day-old adult of *An.*

merus.

Additional file 2: Movie S2. Sperm motility in a crushed normal-like testis from a 2-day-old

adult hybrid of the cross ♀*An. merus* × ♂*An. coluzzii* MOPTI.

Additional file 3: Figure S1. Chromosome behavior during meiosis in testes of *An. coluzzii*

MOPTI. X chromosomes are labeled with 18S rDNA (red), and Y chromosomes are labeled

with retrotransposon *zanzibar* (green). Chromosomes are counterstained with DAPI (blue). Scale

bar – 5 μm.

Additional file 4: Figure S2. Chromosome behavior during meiosis in testes of *An. merus*

MAF. Sex chromosomes are labeled with 18S rDNA (red) and satellite AgY53B (green).

Chromosomes are counterstained with DAPI (blue). Scale bar – 5 μm.

Additional file 5: Figure S3. Sizes and sex chromosome content of spermatids in pure

species and interspecies hybrids. (A) Spermatids from a 2-day-old adult of *An. coluzzii*

MOPTI. (B) Spermatids from a 5-day-old adult F1 hybrid from the cross ♀*An. merus* × ♂*An.*

coluzzii MOPTI. The X and Y chromosomes are visualized with 18S rDNA (red) and

retrotransposon *zanzibar* (green), respectively. Chromatin is counterstained with DAPI (blue).

The insets show magnified images of spermatids.

Additional file 6: Figure S4. Visualization of sex chromosomes in a degenerate testis of a

hybrid adult from the cross ♀*An. coluzzii* MOPTI × ♂*An. merus*. (A) DAPI-stained male

accessory glands and an underdeveloped testis of a one-day-old hybrid adult. (B) Whole-mount

FISH of the degenerate testis with 18S rDNA (red) and satellite AgY53B (green) detects the

interphase sex chromosomes. Chromatin is counterstained with DAPI (blue).

Additional file 7: Table S1. Sizes of metaphase chromosomes in pure species and

interspecies F1 hybrids of the *An. gambiae* complex.

Additional file 8: Figure S5. Increased lengths of chromosomes in testes of interspecies F1

hybrids. Left panel: Lengths of metaphase chromosomes in *An. coluzzii* MOPTI (filled boxplots)

and in F1 hybrid males from the cross *An. merus* × ♂*An. coluzzii* MOPTI (open boxplots). Right

panel: Ratios of the lengths of chromosomes between hybrids and pure species. X and Y – sex

chromosomes. 2 and 3 – autosomes. Statistical significance was assessed with a two-sample

pooled *t*-test.

Additional file 9: Figure S6. Visualization of pairing between sex chromosomes in testes of

a pure species and interspecies F1 hybrid. (A, B) Testis of a one-day-old adult of *An. coluzzii*

MOPTI. (C, D) Testis of a one-day-old adult F1 hybrid of the cross ♀*An. merus* × ♂*An. coluzzii*

MOPTI. Whole-mount FISH with 18S rDNA (Cy3), a satellite from Contig_240 (Cy3), satellite

AgY53B (Cy5), and satellite AgY477-AgY53B junction region (Cy5) is performed to detect the

X (red) and Y (green) chromosomes. Chromatin is counterstained with DAPI (blue).

Additional file 10: Figure S7. Chromatin bridges formed by the *An. merus* X chromosomes

during anaphase in primary spermatocytes of F1 hybrid males from the cross *An. merus* ×

♂*An. gambiae* ZANU. The arrows show chromatin bridges between the X chromosomes.

Additional file 11: Figure S8. Differences in condensation of heterochromatin between *An.*

coluzzii MOPTI and F1 hybrid of the cross ♀*An. merus* × ♂*An. coluzzii* MOPTI. (A) A

squashed preparation of meiotic chromosomes from *An. coluzzii* MOPTI male. (B) A squashed

preparation of meiotic chromosomes from the ♀*An. merus* × ♂*An. coluzzii* MOPTI hybrid male.

The *An. coluzzii* MOPTI Y chromosome is labeled by the satellite AgY477-AgY53B junction

region (Cy5, red). The *An. merus* X chromosome and autosomes of both species are labeled by

the satellite tandem-42345 (Cy3, green). The arrows point to the fibers of uncoiled

heterochromatin on the grayscale inverted image. Chromatin is counterstained with DAPI (blue).

Additional file 12: Table S2. Numbers and percentages of primary spermatocytes with

paired and unpaired sex chromosomes.

	Whole testis	Squashed testis	Crushed testis

A *An.coluzzii* MOPTI

50 μm | 50 μm | 20 μm

B ♀*An. merus* x ♂*An. coluzzii* MOPTI

50 μm | 50 μm | 20 μm

C ♀*An. coluzzii* MOPTI x ♂*An. merus*

50 μm | 20 μm | 20 μm

18S rDNA
zanzibar

Prophase I
Metaphase I
Anaphase I
Metaphase II
Anaphase II
Spermatid
Metaphase II
Anaphase II
Spermatid

Testis of *An. coluzzii*
- Stem cell divisions
- Gonial cell amplificaiton
- Growth and gene expression
- Meiotic divisions
- Spermatid differentiation
- Mature haploid spermatozoa

Testis of F1♀ *An. merus* ×♂ *An. coluzzii*
- Stem cell divisions
- Gonial cell amplificaiton
- Growth and gene expression
- Meiotic prophase I is followed by mitotic anaphase
- Delayed spermatid differentiation and partially developed diploid spermatozoa

Testis of F1♀ *An. coluzzii* ×♂ *An. merus*
- Arrested spermatogenesis with undifferentiated cells

Hub

Gonial cells

Premeiotic cells

Spermatids

Cyst cells

Meiotic prophase I with X-Y chromosome pairing

Meiotic prophase I with no X-Y chromosome pairing

Anaphase I with homologous chromosome segregations

Abnormal anaphase with both X and Y sister chromatid segregations in hybrids

Anaphase II with sister chromatid segregations

Diploid spermatozoa

Haploid spermatozoa

Normal male meiosis in pure species

Prophase I   Anaphase I   Normal sperm

Failed male meiosis in F1 hybrids

Prophase I   Anaphase   Diploid sperm