A new Approximate Bayesian Computation framework to distinguish among complex evolutionary models using whole-genome data

Silvia Ghirotto[*§], Maria Teresa Vizzari[*], Francesca Tassi, Guido Barbujani and Andrea Benazzo[§]

Department of Life Sciences and Biotechnology, University of Ferrara, 44121 Ferrara, Italy

[*] these authors contributed equally to this work

[§] correspondence should be addressed to andrea.benazzo@unife.it and silvia.ghirotto@unife.it

Abstract

Inferring past demographic histories is crucial in population genetics, and the amount of complete genomes now available for many species should in principle facilitate this inference. In practice, however, the available inferential methods suffer from severe limitations. Although hundreds complete genomes can be simultaneously analyzed, complex demographic processes can easily exceed computational constraints, and there are no standard procedures to make sure that the estimates obtained are reliable. Here we present ABC-SeS, a new Approximate Bayesian Computation (ABC) framework, based on the Random Forest algorithm, to infer complex past population processes using complete genomes. All possible pairs of populations are compared, and the data are summarized by the full genomic distribution of the four mutually exclusive categories of segregating sites, a set of statistics fast to compute even from unphased genome data. We constructed an efficient ABC pipeline and tested how accurately it allows one to recognize the true model among models of increasing complexity, using simulated data and taking into account different sampling strategies in terms of number of individuals analyzed, number and size of the genetic loci considered. Once assessed the power of ABC-SeS in the comparison of even complex models, we applied it to the analysis of real data, testing models on the dispersal of anatomically modern humans out of Africa and exploring the evolutionary relationships of the three species of

26 Orangutan inhabiting Borneo and Sumatra. The flexibility of our ABC framework, combined with

27 the power provided by the set of statistics proposed pave the way for reliable inference of past

28 population processes for any species for which high coverage genomes are available.

Authors' summary

We propose a new method for inferring demographic history from whole-genome data, ABC-SeS.

The idea is to analyze the distributions of segregating sites within an Approximate Bayesian

Computation framework, using a Random Forest approach. Extensive simulation shows that ABC-

SeS can handle very large datasets and identify the model that generated the data, even when

population relationships are complicated by factors such as admixture. We also provide guidelines

concerning the sampling strategies leading to the most accurate inference. As examples of possible

applications, we show: (1) that complex alternative models on human dispersal out of Africa can

now be formally tested and compared by ABC-SeS; and (2) that the evolutionary relationships

among three orangutan species can now be better assessed by ABC-SeS than by alternative

methods.

Introduction

A robust reconstruction of the demographic dynamics of a species is important both to

improve our knowledge about the past and to disentangle the effects of demography from those of

natural selection [1–3]. In recent years, thousands of modern and ancient complete genome

sequences have become available, potentially containing vast amounts of information about the

evolutionary history of populations [4–10]. However, these genomes do not speak by themselves; to

extract the evolutionary information they contain, appropriate inferential statistical methods are

required. Some methods based on the Sequential Markovian Coalescent (SMC) model [11], became

popular among population geneticists due to their ability to infer population size changes through

time (PSMC [12]) and divergence times (MSMC [13]), and to scale well on whole genome

sequences. Under these approaches, the local density of heterozygote sites along chromosomes is

used to estimate the times of the most recent common ancestor (TMRCA) of genomic regions

separated by recombination, thus providing insight into ancestral population sizes and the timing of

past divergence processes. These estimates are often used to indirectly support hypotheses

regarding the evolution of the studied organisms. Albeit sophisticated, these methods present some

limitations; the temporal resolution of the inferred demographic events seems to be strongly

dependent on the number of individuals included, with poor performance in the recent past

especially when analyzing single individuals. Moreover, these methods assume no gene flow among

the investigated populations, which in many case is plainly implausible. The consequences on the

inferential process of violation of this assumption are still under investigation [14].

Other methods infer demographic parameters via the diffusion approximation [15], or

coalescent simulations [16], from the site frequency spectrum (SFS) computed on large genomic

datasets. The SFS records the observed number of polymorphisms segregating at different

frequencies in a sample of *n* individuals and it is generally computed over a certain number of

genomic regions where no influence of natural selection is assumed. The expectation of the SFS

under different evolutionary scenarios could be approximated by the diffusion theory (as

implemented e.g. in *dadi*) or directly via coalescent simulations (as in *fastsimcoal*); alternative

demographic histories can be explicitly compared via e.g. AIC [17]. Still, there are limits to the

complexity of models that can be analyzed, and AIC-like approaches can only be used to

understand which modifications significantly improve the model. Therefore, through these

approaches, it is not easy to evaluate whether and to what extent the compared models can actually

be distinguished from each other, and hence quantify the strength of the support associated to the

best model. Indeed, the only available procedure to assess the model identifiability requires the

analysis of many datasets simulated under known demographic conditions, which can be

computationally prohibitive, in particular for complex evolutionary scenarios [16].

Recently, an inferential method that couples the ability of the SMC to deal with whole

genome sequences and the population signal gathered from the SFS has been developed (SMC++

[18]). Under this inferential framework, both the genomic and the SFS variation are jointly used to

estimate population size trajectories through time, as well as the divergence time between pairs of

populations. Although this approach seems to scale well on thousands of unphased genomes, it is

based on the same assumption of classical SMC methods (with populations evolving

independently), which severely limits its use whenever gene flow cannot be ruled out.

One powerful and flexible way to quantitatively compare alternative models and estimating

model's parameters relies on the Approximate Bayesian Computation (ABC) methods. Under these

methods, the likelihood functions do not need be specified, because posterior distributions can be

approximated by simulation, even under complex (and hence realistic) population models,

incorporating prior information. The genetic data, both observed and simulated, are summarized by

the same set of "sufficient" summary statistics, selected to be informative about the genealogic

processes under investigation. The ability of the framework to distinguish among the alternative

demographic models tested and the quality of the results can be evaluated with rather limited

additional effort (for a review see e.g. [19]). Although ABC has the potential to deal with complex

and realistic evolutionary scenarios, its application to the analysis of large genomic datasets, such as

complete genomes, is still problematic. In its original formulation, indeed, the ABC procedure

requires the simulation of millions data sets of the same size of those observed, which becomes

computationally very expensive as the dataset increases in size, or when many models need be

compared. In addition, there is a problem with the choice of the summary statistics describing both

observed and simulated data, as recognized since the first formal introduction of ABC [20,21].

Increasing the number of summary statistics, indeed, makes it easier to choose the best model, but

inevitably reduces the accuracy of the demographic inference (this problem is referred to as the

"curse of dimensionality" [22]). Ideally, the good practice would be to select a set of summary

statistics that is both low-dimensional and highly informative on the demographic parameters

defining the model. In practice, however, this problem has never really been solved, despite several

serious attempts [23].

Recently, a new ABC framework has been developed based on a machine-learning tool

called Random Forest (ABC-RF [24]). Under ABC-RF, the Bayesian model selection is rephrased

as a classification problem. At first, the classifier is constructed from simulations from the prior

distribution via a machine learning RF algorithm. Once the classifier is constructed and applied to

the observed data, the posterior probability of the resulting model can be approximated through

another RF that regresses the selection error over the statistics used to summarize the data. The RF

classification algorithm has been shown to be insensitive both to the correlation between the

predictors (in case of ABC, the summary statistics) and to the presence of relatively large numbers

of noisy variables. This means that even choosing a large collection of summary statistics, the

correlation between some of them and others (which may be uninformative about the models

tested), have no consequences on the RF performance, and hence on the accuracy of the inference.

Moreover, compared to the standard ABC methods, the RF algorithm performs well with a radically

lower number of simulations (from millions to tens of thousands per model). These properties make

the new ABC-RF algorithm of particular interest for the statistical analysis of massive genetic

datasets.

In this paper we present ABC-SeS (where SeS stands for Segregating Sites), a new

framework for demographic inference via ABC-RF, suitable for the analysis of complete genomes.

ABC-SeS compares all possible pairs of populations, summarizing both the observed and the

simulated genetic variation by the distribution over the genome of the four mutually exclusive

categories of segregating sites (i.e. private polymorphisms in either population, shared

polymorphisms and fixed differences) known to be informative about processes of divergence and

admixture [25], and not requiring phased data. These statistics have already been successfully used

in a standard ABC context [26], but only in the form of the first four moments of the distribution

across loci. Here, for the first time, and thanks to the ABC-RF procedure, we analyze the full

genomic distribution of each statistic. We performed a power analysis, to evaluate how accurately

this ABC pipeline can recognize the true model among models of increasing complexity, using

simulated data. We also explored the inferential power of the presented procedure with respect to

the experimental conditions, evaluating the consequences of sampling strategies involving different

numbers of genomes, different numbers of loci, and different locus lengths. Finally, we applied our

method to two case studies. First, we analyzed the demographic history of anatomically modern

humans and the dynamics of migration out of the African continent, explicitly comparing the

models proposed by [27,28]. Secondly, we reconstructed the past demographic history and the

interaction dynamics among the three orangutan species inhabiting Borneo and Sumatra, revising

the models presented by [29].

Results

Summary Statistics

We summarize the genetic data by means of the genomic distributions of the four mutually

exclusive categories of segregating sites in two populations, namely 1- segregating sites private of

the first population, 2- segregating sites private of the second populations, 3- segregating sites that

are polymorphic in both populations and 4- segregating sites fixed for different alleles in both

populations [25]. To calculate the complete genomic distribution of these statistics we considered

the genome as subdivided in a certain number of independent fragments of a certain length, and for

each fragment we counted the number of sites belonging to each of the four above-mentioned

categories. The final vector of summary statistics is thus composed by the truncated frequency

distribution of fragments having from 0 to n segregating sites in each category, for each pair of

populations considered. We refer to this set of statistics as FDSS (frequency distribution of

segregating sites) hereafter. The maximum number of segregating sites in a locus of a certain length

is fixed to n, and hence the last category contains all the observations higher than n. In the one

population models, the four distributions described above collapse in a single distribution

describing the frequency of loci showing specific levels of within-population segregating sites.

Power Analysis

To determine the power of the ABC-SeS procedure in distinguishing among alternative evolutionary trajectories, we performed an extensive power analysis, as described below.

We simulated genetic data considering different experimental conditions, thus testing all the possible combinations of locus length (bp) {200; 500; 1,000; 2,000; 5,000}, number of loci {1,000; 5,000; 10,000} and number of chromosomes {2, 4, 10, 20}, for a total of 60 combinations of sampling conditions tested. The data were generated using the *msms* simulator according to each of these combinations for three sets of non-nested models of increasing complexity, namely one-population models (four alternative models, Fig 1A), two-population models (three alternative models, Fig 2A) and multi-populations models (two alternative models, Fig 3A); for a more comprehensive description of the models see the Materials and Methods section.

For each combination of experimental conditions, we compared alternative models within the three sets tested treating 1,000 simulated datasets for each model as pseudo-observed data (pods). The models were compared through ABC-RF, and we counted the number of times the model selection procedure correctly assigned a pod to the true model, i.e. to the model that actually generated that pseudo-observed dataset. The proportion of true positives (TP) thus calculated is a measure of the power of the whole procedure, considering all its features (model selection approach, alternative models compared, statistics summarizing the data, genomic parameters simulated). The TP results are summarized in Figs 1-3B.

One-population models

We designed four one-population models, respectively considering a population of constant size through time, a bottleneck, an exponential growth and a structured population. In each of the 50,000 simulations (per model and per combination of experimental parameters) demographic and evolutionary parameter values are drawn from prior distributions, detailed in S1 Table. The four

plots of Fig 1B report the results of the power analyses. In each plot, we see the proportion of times

each model was correctly recognized as the most likely one. In general, the percentage of true

positives is quite high, ranging from almost 80% to 100% depending on the model generating the

pod and on the combination of experimental conditions tested. The bottleneck model has the highest

rate of identification, with most combinations of experimental conditions yielding nearly 100% true

positives. By contrast, the least identifiable model seems the one considering a structured

population, with 0.76 to 0.86 true positives, depending on the combination of experimental

conditions. However, we observed that the decrease in the power is actually linked to the extent of

gene flow among demes, and to the number of demes sampled; as increasing gene flow and

decreasing the number of demes sampled, the structured and the panmictic models converge, hence

becoming harder to distinguish (S2 Fig). As expected, we observed a general increase in the power

with the increase of both the locus length and the number of loci considered. The number of

sampled chromosomes does not appear to be directly linked to the increase of the proportion of true

positives. For some sampling conditions, we observed instead a decrease in the TP rate going from

2 to 20 chromosomes (see Fig 1B). However, we showed that this unexpected behavior can be

explained with the overlap of the FDSS generated by the constant and the structured models, which

increases as increasing the number of chromosomes sampled (S3 Fig). We repeated the whole

analysis considering intra-locus recombination (S4 Fig), and the general pattern did not change. We

observed a reduced power for the structured model (TP from 75 to 90%) with respect to the other

models tested (TP from 80 to 100%), and a stronger effect of the locus length and of the number of

loci, than of the number of chromosomes simulated.

Two-populations models

The two-population scenarios include a divergence model with isolation after the divergence

(no gene flow); a divergence model with continuous migration from the separation until present

times; and a divergence model with a single event of bidirectional migration (i.e. pulse of

admixture), for a total of three models compared for each combination of experimental parameters

(Fig 2A, prior distributions reported in S2 Table). The plots in Fig 2B clearly show that the most

recognizable model is the one including divergence and migration, with proportions of true

positives close to 0.95, regardless of the combination of experimental conditions tested. For the

other two models, the power was clearly lower than that estimated for the one-population models,

with values ranging from 40% to 65% (for the model without migration) and from 60% to 80% (for

the model with a pulse of admixture). This low power might be due to the fact that, when the pulse

of admixture of the third model occurs right after the divergence, the two models become

essentially identical. To verify this prediction, we calculated the proportion of TP for the pulse of

admixture model, subdividing the pods in seven categories, depending on the time interval between

the divergence and the admixture event (S5 Fig). The proportion of the pods correctly assigned to

the pulse of admixture model increases as the intervals between divergence and admixture increase,

reaching values of 85% for some combination of parameters. In this case, we did not observe

remarkable differences with respect to specific experimental parameters. The performance of the

method improved when we included intra-locus recombination in the simulations (S6 Fig). In this

case, we observed an increase of the power with increasing locus length, both for the no migration

and for the pulse of admixture model, respectively reaching values of 95% and 100%.

We also ran many parallel simulations, using either the whole FDSS or its first four

moments only. In all cases (a selection of the results is in S3 Table) the FDSS's ability to

distinguish among different evolutionary scenarios was greater, and sometimes substantially so,

under the former conditions.

Multi-populations models

In most realistic cases, interactions among several populations need be considered. Among

the many possible scenarios, we chose to initially focus on the alternative hypotheses proposed to

explain the expansion of anatomically modern humans out of the African continent. We shall refer

to the two models (shown in Fig 3A and detailed in Materials and Methods) as Single Dispersal

(proposed e.g. in [27]) and the Multiple Dispersal (proposed e.g. in [28,30,31]). The main

demographic events are shared between the two models, and here we shall refer to the description

given by [27]: three archaic branches (unknown, Denisova and Neandertal), three modern branches

(Africans, Eurasians and Papuans) and a specific pattern of gene flow. The difference between the

two alternative models lies in the details of the expansion from Africa. Under the Single Dispersal

model (SDM), Eurasians and Papuans derived from the same expansion from Africa, through the

Near East. Under the Multiple Dispersal model (MDM), the Papuans derived from an earlier

dispersal from Africa, independent from that giving rise to the Eurasian populations.

The prior distributions associated with these models are reported in S4 and S5 Table. Fig 3B

summarizes the power analysis. The proportion of true positives ranges between 0.65 and 0.70 for

the SDM, regardless of the combination of experimental parameters tested, and between 0.6 and 0.8

for the MDM, in this case with a slight increase of the power with the size of the fragments

simulated. Being the SDM and the MDM quite similar, in particular when the time interval between

the first and second exit in the MDM is short, we also evaluated the ABC-SeS ability to identify the

correct model as a function of the time span between the divergence time of the African ghost

populations and the second exit in the Multiple Dispersal model. To do this, we considered 10,000

pods from the MDM. We then subdivided these 10,000 pods in 6 bins of increasing interval

between these two events (up to 60,000 years), and, within each bin, we separately calculated the

proportion of times in which the MDM is correctly recognized by the ABC-RF procedure. As might

be expected, the proportion of true positives increases with increasing intervals between the

divergence of the two African ghost populations and the time of the second exit (S7 Fig), reaching

values of 90% for some combinations of experimental parameters.

For the multi-populations models we tested the effect of intra-locus recombination only for

4 key sampling schemes that combine few (1,000) and many (10,000) genetic loci with either short

(500 bp) or long (2,000 bp) DNA fragments. We selected these combinations because they might

represent extremely favorable (10,000 loci, 2,000 bp length) or unfavorable (1,000 loci 500bp

length) conditions that can be encountered when analyzing real data. Compared with the results

obtained for the same experimental parameters without considering intra-locus recombination, we

observed a slight increase in the TP rate when analyzing high number of fragments, or when

considering longer locus lengths (S6 Table).

Real Case: out of Africa dynamics

Simulations in the previous section basically show that alternative models can be

distinguished by ABC-SeS, except when they differ for parameters approaching 0, thus creating

overlaps between their expected consequences. We then moved to applying the ABC-SeS method to

estimate posterior probabilities of alternative models about early human expansion from Africa.

Whether human demographic history is better understood assuming one [5,27] or two [28,30,31]

major episodes of African dispersal is still an open question. While concluding that indigenous

Australians and Papuans seem to derive their ancestry from the same African wave of dispersal as

most Eurasians, Mallick et al. [5] admitted that these inferences change depending on the

computational method used for phasing haplotypes. Therefore, it made sense to compare the SDM

and the MDM by ABC-SeS. For this purpose, we analyzed the high coverage Neandertal genome

[7], the high coverage Denisova genome [6], and a large set of modern individuals published in [28]

(S7 Table). To make sure that the choice of the genomes does not affect the results, we repeated the

model selection analysis replacing the Papuan individuals with those published by [27] (S7 Table).

We extracted from all these genomes 10,000 shared independent fragments of 500 base pairs length

(see Material and Methods for details). Since we knew that the number of chromosomes has a

minor impact on the analysis, we considered two chromosomes per population, so that the sample

size would be the same in ancient and modern samples. The proportion of true positives for the

combination of experimental parameters here considered (i.e. 10,000 loci of 500 bp length and 2

chromosomes per population) was 0.66 for the SDM, and between 0.69 and 0.85 for the MDM (Fig

3A and S7 Fig). Considering recombination, we observed an increase of the proportion of TP for

the MDM, which (averaged over the whole range of possible divergence times between the African

ghost populations and the second exit) reached a value of 0.75.

As the SDM and the MDM mainly differ for the origin of the Papuan population (whether or

not descending from the same ancestor of Eurasians), to test for the robustness of the results we

used the genomic information of all the six Papuan individuals sequenced by [28]. To this purpose,

we repeated the ABC-RF model selection procedure six times, each time considering a different

Papuan individual. Fig 4 and S8-S11 Tables presents the results of the ABC-RF model selection,

showing that, in all the comparisons, the results supported the MDM, with posterior probabilities

ranging from 0.67 to 0.69. We repeated the whole analysis considering the 25 Papuan individuals in

[27], this time repeating the ABC-RF model selection procedure 25 times, and always found results

favoring the MDM over the SDM. The posterior probabilities estimated for the MDM were

comparable to those estimated with the Pagani dataset [28] (Fig 4). Considering intra-locus

recombination, MDM received additional support, with posterior probabilities ranging from 0.70 to

0.76 for both datasets (Fig 4).

Real Case: Orangutan evolutionary history

As a second application of our framework to real data, we investigated the past demographic

and evolutionary dynamics of the three known orangutan species. In addition to the orangutan

species previously recognized in Borneo (*Pongo pygmeus*) and in Sumatra, North of Lake Toba

(*Pongo abelii*), [29] described a new species of Sumatran orangutan, *Pongo tapanuliensis*, South of

Lake Toba. To reduce the otherwise excessive computational effort in their ABC analysis, [29] had

to compare simplified versions of the alternative models. Factors such as bottlenecks and population

structure were considered only when estimating the parameters of the best-performing model,

namely the one with an earlier split between the two Sumatran species, and an origin of the Bornean

species from *Pongo tapanuliensis* ancestors. The ABC-SeS method does not suffer from this

shortcoming. We compared four complex scenarios, designed following the models presented in

[29] (Fig 5A). While considering the same substructure within the three orangutan species, the four

models differ in the genealogical relationships assumed among species (Fig 5A and Materials and

Methods for details). From the whole genomic dataset presented in [29], we selected seven

individuals, one from each of the populations emerged from the study (S12 Table, see Materials and

Methods for details). Following the procedure described in Materials and Methods, we extracted

9,000 independent sites 1kb length shared among the 7 selected individuals, on which we calculated

the FDSS. We generated 50,000 simulations per model considering the above mentioned genomic

structure (9,000 loci, 1kb length and 2 chromosomes per population), with prior distributions

detailed in S13-S16 Tables. We first assessed the ability to correctly recognize the four alternative

models tested through a power analysis, considering 1,000 pods per model. The results are

presented in Fig 5A. The most identifiable model (TP=0.81) appeared to be the model 2b, under

which there is a first separation of South Toba from Borneo Orangutan, followed by the divergence

of North Toba from South Toba. The model assuming an early separation of South Toba form

North Toba, followed by the separation of Borneo from South Toba, actually showed the lowest

proportion of true positives (0.45). The application to real data favored the model 1a, (also

associated with the highest posterior probability in [29]), with a posterior probability of 0.49. Under

the most supported model both the North Toba (first) and Borneo (later) separated from *Pongo*

tapanuliensis Fig 5B.

Discussion

The cost of typing large genomic datasets has dramatically decreased lately, making

population-scale genomic data available for a large set of organisms [4,9,10,32,33]. The main

challenge now is how to extract as much information as possible from these data, developing

flexible and robust statistical methods of analysis [12,13,16]. Approximate Bayesian Computation,

explicitly comparing alternative demographic models and estimating the models' probabilities,

represents a powerful inferential tool for about past population dynamics [34]. Another advantage

of such a simulation-based approach is the possibility to easily check whether the models being

compared are actually distinguishable, hence quantifying the reliability of the estimates produced

[35]. Nevertheless, only recently, with the development of the Random Forest procedure for ABC

model selection [24], it has become possible to apply ABC to large genomic datasets. With this

work, we took advantage of this newly proposed algorithm to develop ABC-SeS, an ABC-based

flexible framework to compare demographic models. The novelty of this framework mainly lies in

the summary statistics used to summarize genomic data, namely the complete genomic distribution

of the four mutually exclusive categories of segregating sites for pairs of populations [25].

Power Analysis

Initially, we analyzed sets of models with increasing levels of complexity, simulating

genetic data under a broad spectrum of experimental conditions. This extensive power analysis

showed that ABC-SeS can often identify the model under which the data were generated, with some

uncertainties only when two models are just marginally different. This was the case for both simple

(one or two-population scenarios, Figs 1 and 2) and complex (multi-populations scenarios, Fig 3)

demographies. When we compared one-population scenarios, to summarize the data we necessarily

considered only one of the four distributions of polymorphic sites. Nonetheless the model

identifiability, calculated as the proportion on TPs over 1,000 pods, reached values between 80%

and 100%, with slightly lower values only for the structured model. This reduction in the power was

always due to the levels of gene flow among demes; when it is high, the structured model becomes

hardly distinguishable from the panmictic model (S2 Fig), as has already been known since

Wright's times [36]. We also showed that the power depends on the number of demes; indeed, the

proportion of TPs increases in parallel with the number of demes considered in the structured model

(S2 Fig). Among the two-populations demographies, the model with bi-directional migration at a

constant rate proved easiest to identify, with almost 100% TPs, regardless of the combination of

experimental parameters tested. Predictably (see e.g. [37]), the pulse of admixture model produced

the most accurate results when the two populations were highly genetically divergent and the time

interval between admixture and divergence was large (S5 Fig). Our results showed that models are

difficult to discriminate only when the divergence is recent, and/or close the admixture event. Pods

generated under these conditions, indeed, are often erroneously assigned to the other models.

Even when more complicated scenarios were compared (e.g., the multi-populations models),

the ABC-SeS framework recognizes the true model, with about 70% TPs. It is worth noting that in

the MDM the first and the second exit from Africa could sometimes occur at very close times. In

these cases, the SDM and the MDM models become extremely similar. Indeed, we observed an

increase in the power of the test at increasing intervals between the African divergence and the

second exit (S7 Fig), reaching values close to 90%.

In the first round of comparisons, we assumed complete linkage among all sites within the

same fragment. Even if acceptable for some experimental conditions (i.e. for short locus lengths),

this is generally simplistic. We thus repeated the power tests, this time generating data that could

undergo intra-locus recombination. In all the comparisons, we found that, with recombination, the

results are generally better, regardless of the experimental parameters considered, and particularly

when the loci are long. In particular, we observed a substantial increase of the proportion of TPs for

the MDM, but not for the SDM (S6 Table). One possible explanation for this finding is, in a tree

generated under SDM, all loci of the African, European, Asian and Papuan populations, belong to a

branch that has a unique topology (T1 in S8 Fig) and recombination will not alter this condition. By

contrast, under MDM, the topologies are three (T1, T2 and T3 in S8 Fig) a recombination event will

generate a transition from one of these baseline topologies to the recombinant topology (denoted as

R in S8 Fig). Thus, recombination under MDM reduces the fraction of loci showing the T1

topology, i.e. the topology shared with the alternative SDM, thus increasing the possibility that the

two models be discriminated. As could be expected, under most conditions tested, the model

identifiability increases increasing the locus length, and the number of loci considered. Indeed, by

analyzing more loci and longer loci, the probability of observing recombination increases, resulting

in a higher fraction of the R topologies. Thus, when one is analyzing many loci, or long stretches of

DNA, recombination should be considered, although this is not necessarily leading to an increase in

TPs, because the specific features of the demographic models compared are also a factor.

As it is intuitively reasonable, and how it has been already pointed out e.g. by [38], the

accuracy of the model selection seems to be more dependent on the number of loci considered and

on the locus length rather than on the number of individuals sampled per population. In absence of

recombination, the locus length appeared to be crucial to correctly identify simple demographies,

such as those generated under one-population models, whereas its effect was minor when complex

scenarios were compared (Figs 1 and 3). When recombination was included in the modeling, the

locus length seemed to become the main factor leading to a good discrimination between models. In

the two-populations scenarios we observed a significant increase of the proportion of TPs at

increasing lengths of the fragments simulated, reaching values close to 100% (S6 Fig). It is anyway

worth noting that the power to correctly identify the true model was quite good even when we

simulated short fragments, as well as in the comparison of complex demographies (Fig 3). This

finding means that the ABC-SeS procedure is particularly suitable for the analysis of ancient data

[6] and of RAD sequencing data [39], where short DNA fragments are more the rule than the

exception.

Applications to real datasets

We compared two demographic models about the anatomically modern humans expansion

out of Africa, combining ancient (Neandertal and Denisova) and modern genomes data. Because we

were including ancient data, we restricted the analysis to short fragments (500 bp) to maximize the

number of independent loci retrievable. This combination of experimental parameters (10,000 loci

500 bp length, 2 chromosomes per population) showed anyway a good ability to distinguish

between models (Fig 3). For the first comparison, we used as modern genomes those published by

[28], and repeated the model selection procedure six time, each time considering one of the six

Papuan genomes in the dataset. All such comparisons supported the MDM (Fig 4), i.e. a first

expansion of the ancestors of the current Austro-Melanesians, followed by a second expansion

leading to the peopling of Eurasia. We obtained the same conclusion by using Papuan genomes

coming from another dataset [27] (Fig 4). Considering different modern individuals from African,

European and Asian populations did not change the support for the MDM. These results raise

several questions; indeed, it was the SDM that showed the best fit in [27], whereas the MDM

appeared to account for the data only when the analysis was restricted to modern populations.

However, our findings are in agreement with those by [28], who estimated that at least 2% of the

Papuan genome derive from an earlier, and distinct, dispersal out of Africa. Other genomic studies

[31], but not all [5], and phenotypic analyses [30] appear support the MDM, which calls for further

research in this area. Be that as it may, in no other study besides the present one (i) the alternative

hypotheses are explicitly compared analyzing complete genomes; (ii) posterior probabilities are

estimated for each model, and (iii) the accuracy of the estimates is assessed by power analysis.

We then moved to investigating the evolutionary history of the three extant Orangutan

subspecies, the last of which (located south of the Lake Toba) has been recently described by [29].

We basically repeated the ABC analysis performed by [29] applying our ABC-SeS framework.

Nater and colleagues [29] started comparing simple evolutionary scenarios, and considered

population substructure and gene flow only when estimating parameters, but not in the phase of

model choice. ABC-SeS allowed us to analyze complex demographies from the very beginning, i.e.

from the model selection phase, thus comparing more realistic evolutionary scenarios. We basically

obtained the same results presented by [29]; in both cases the most supported model was the one

accounting for a first separation of the North Toba and a later separation of the Borneo species from

the newly identified South Toba species (Fig 5B). The main difference was about the strength of the

support associated to this model. While Nater and colleagues [29] estimated high posterior

probabilities for the best-fitting model (73% when comparing the 4 models and 98% when

comparing the two best scenarios), our procedure associated to the same model a posterior probability of 49% (Fig 5A). Moreover, the power analysis that we conducted (absent in [29]), revealed that the ability to correctly distinguish among the four tested models is between 45% and 81%, with the selected model that can be erroneously recognized as the most probable one in the 38% of cases. These results emphasize (i) the importance of including complex demographic histories in the model selection step, so as to evaluate the real posterior probability associated to the best model, on which the parameter estimation will be performed and (ii) the importance of performing a power analysis of the models tested, so as to be aware of the level of uncertainty about the conclusions of the study.

Perspectives

In this paper we presented a new framework based on ABC-RF to explicitly compare complex demographic models exploiting whole-genome data. We proposed to summarize the data with a set of statistics that were already known to be informative about past population dynamics [25], but for the first time, and thanks to the Random Forest algorithm, we used the entire genomic distribution of the four categories of segregating sites between pairs of populations in an ABC context. It is clear that considering the whole FDSS, rather than its first four moments, significantly improves our ability to distinguish among different evolutionary scenarios. How large this improvement, depends on the amount of genetic loci available, and, of course, on how large are the differences between the alternative models. At present, the ABC-SeS is optimized for high-coverage genotypes [5,10,33]. A natural extension of this work will thus be to implement the use of low coverage data, developing an approach able to retrieve the FDSS taking into account the genotype uncertainty and sequencing errors, for instance through the use of the genotype likelihood (as, e.g., in ANGSD [40]).

The flexibility of the ABC-RF model selection approach, combined with the inferential power proven by the summary statistics that we proposed to calculate on genomic data, pave the

way for a detailed and comprehensive study of complex demographic dynamics for any species for

which few high coverage genomes are available.

Materials and Methods

ABC Random Forest

Approximate Bayesian computation is a flexible framework, based on coalescent

simulations, to compare different evolutionary hypotheses and identify the one that with the highest

probability generated the observed data [20,34]. Despite its potential, the application of the classical

ABC approach to the analysis of large genomic datasets and complex evolutionary models, has

been quite limited. Possible reasons are (i) the levels of arbitrarity in the choice of the sufficient set

of summary statistics and (ii) the high number of simulations required to obtain good quality

posterior estimates, resulting in an unacceptable increase of computational time when multiple

complex models are simultaneously analyzed. Both these issues appear to have been solved with the

development of a new ABC approach, based on a machine learning procedure (Random Forest

[24]). Random Forest uses the simulated datasets for each model in a reference table to predict the

best fitting model at each possible value of a set of covariates. After selecting it, another Random

Forest obtained from regressing the probability of error of the same covariates estimates the

posterior probability. This procedure allows one to overcome the difficulties traditionally associated

with the choice of summary statistics, while gaining a larger discrimination power among the

competing models. Moreover, a satisfying level of accuracy in the estimates is achieved with about

30-50 thousand simulations per model [24], significantly reducing the computational cost of the

analysis of complex demographic histories. All the ABC-RF estimates have been obtained using the

function *abcrf* from the package abcrf and employing a forest of 500 trees, a number suggested to

provide the best trade-off between computational efficiency and statistical precision [24]. We

compared all models and obtained the posterior probabilities using the function *predict* from the

same package.

Power Analysis

For the power analysis, we generated data under different combinations of experimental parameters, varying the number of loci (calculating FDSS on 1,000, 5,000 and 10,000 loci), the locus length (200; 500; 1,000; 2,000; 5,000 bp) and the number of chromosomes sampled per population (2, 4, 10, 20), for a total of 60 combinations per model. We evaluated the power considering three sets of models of increasing complexity, detailed below.

One-population models.

We started by considering four demographic models (Fig 1). The first model represents a constantly evolving population with an effective population size $N1$, drawn from a uniform prior distribution (S1 Table). Under the second model, the population experienced a bottleneck of intensity i, T generations ago. The intensity and the time of the bottleneck, and the ancient effective population size Na are drawn from uniform prior distributions, showed in S1 Table. The third model represents an expanding population. The expansion (of intensity i) is exponential and starts T generations ago, with the effective population size increasing from $N1/i$ to $N1$ (prior distributions in S1 Table). Under the last model, the population is structured in different demes, exchanging migrants at a certain rate. The actual number of demes d, the migration rate m and the effective population size $N1$ are drawn from prior distributions (S1 Table).

Two-populations models.

We then moved to considering three demographic models with two populations (Fig 2). The first one is a simple split model without gene flow after the divergence. Under this model, an ancestral population of size $Nanc$ splits $Tsep$ generation ago into two populations. These two derived populations evolve with a constant population size ($N1$ and $N2$) until the present time (priors for these free parameters are shown in S2 Table). The second model also includes a continuous and bidirectional migration, all the way from the divergence moment to the present. The

per generation migration rates *m12* and *m21* are drawn from priors defined in S2 Table. The third

and last model assumes a single pulse of bidirectional admixture at time *Tadm* after divergence.

Admixture rates *adm12 adm21*, and the time of admixture are drawn from priors (S2 Table).

Multi-populations models.

Finally, we compared two demographic models, representing the two alternative hypotheses

that have been proposed to explain the dispersal of anatomically modern humans out of the African

continent (i.e single and multiple dispersals, Fig 3). To design the models we followed the

parametrization proposed by [27], with some minor modifications. Both models share the main

demographic structure: on the left the archaic groups (i.e. Neandertal, Denisova and an unknown

archaic source), and on the right the anatomically modern humans (with a first separation between

Africans and non-Africans and subsequent separations among population that left Africa). Given

the evidence for admixture of Neandertals and Denisovans with non-African modern human

populations [6,7], we allowed for genetic exchanges from archaic to modern species, indicated in

Fig 3 by the colored arrows. It has to be noted that the archaic populations actually sending

migrants to modern humans are two ghost populations that diverged from the Denisovan and the

Neandertal Altai samples 393 kya and 110 kya, respectively [27]. This way, we took into account

that the archaic contributions to the modern gene pool did not necessarily come from the archaic

populations that have been genotyped so far. We modeled bidirectional migration between modern

populations along a stepping-stone, thus allowing for gene flow only between geographically

neighboring populations. Under the Single Dispersal model (SDM) a single wave of migration

outside Africa gave rise to both Eurasian and Austromelanesian populations, whereas under the

Multiple Dispersal model (MDM) there are two waves of migration out of Africa, the first giving

rise to Austromelanesians and the second to Eurasians. We took into account the presence of

genetic structure within Africa modeling the expansion from a single unsampled "ghost" population

under the SD model, and from two separated unsampled "ghost" populations for the MD model.

The prior distributions for all the parameters considered in these models are in S4 and S5 Tables.

We simulated both demographic models under all possible combinations of experimental parameters. We ran 50,000 simulations per model and combination of experimental parameters, using the *ms/msms* software [41,42]. To evaluate the quality of the estimates of the model selection procedure we calculated the proportion of True Positives (TP) as follows. We randomly sampled 1,000 datasets from each simulated combination of demographic model and experimental parameters. We treated each of these sampled datasets as pseudo-observed data (pod) and performed a models comparison within the correspondent set of models tested. For each model and combination of experimental parameters, we calculated the proportion of TPs as the proportion of times the true model was the most supported by the model selection procedure.

Real Case: out of Africa dynamics

We considered the ancient high-coverage genomes of Denisova and Neandertal [6,7], together with modern human samples from [28]. A detailed description of the samples is in S7 Table. All the individuals were mapped against the human reference genome hg19 build 37. To calculate the observed FDSS we only considered autosomal regions found outside known and predicted genes +/- 10,000 bp and outside CpG islands and repeated regions (as defined on the UCSC platform, [43]). We also took into consideration only the fragments found in genomic regions that passed a set of minimal quality filters used for the analysis of the ancient genomes (map35_50%, [6,7]). From the resulting regions we extracted 10,000 independent fragments of 500 bp length, separated by at least 10,000 bps. All comparisons involved a single individual (i.e. two chromosomes) per population, and so each run of the analysis took into account the Denisova, the Neandertal, one African, one European one Asian and one Papuan individual (detailed in S7 Table). We calculated the observed FDSS six times, each time considering a different Papuan individual present in the Pagani dataset [28]. We repeated the whole procedure substituting the Papuan individuals with those published by [27]. To do this, we downloaded the corresponding alignments in CRAM format from https://www.ebi.ac.uk/ega/datasets/EGAD00001001634. The *mpileup* and

call commands from *samtools-1.6* [44], were used to call all variants within the 10,000 neutral genomic fragments, using the *--consensus-caller* flag, without considering indels. We then filtered the initial call set according to the filters reported in [27] using *vcflib* and *bcftools* [44]. We calculated the FDSS for each of these Papuan individuals, hence obtaining 25 observed FDSS, that have been separately analyzed through the ABC-RF model selection procedure.

Real Case: Orangutan evolutionary history

We selected seven Orangutan individuals, one from each of the populations defined by [29], choosing the genomes presenting the highest coverage (S12 Table). We downloaded the FASTQ files from https://www.ncbi.nlm.nih.gov/sra/PRJEB19688, and mapped the reads to the ponAbe2 reference genome (http://genome.wustl.edu/genomes/detail/pongo-abelii/) using the BWA-MEM v0.7.15) [45]. We used picard-tools-1.98 (http://picard.sourceforge.net/) to add read groups and to filtered out duplicated reads from the BAM aligments. We performed local realignment around indels by the Genome Analysis Toolkit (GATK) v2.7-2 [46]. To obtain genomic fragments suitable to calculate the FDSS, we generated a mappability mask (identified with the *GEM-mappability* module from the *GEM* library build [47]) so as to consider only genomic positions within a uniquely mappable 100-mer (up to 4 mismatches allowed). We then excluded from this mask all the exonic regions +/- 10,000 bp, repeated regions (as defined in the *Pongo abelii* Ensembl gene annotation release 78), as well as loci on the X chromosome and in the mitochondrial genome. We then generated the final mask calculating the number of fragments separated by at least 10 kb, thus obtaining 9,000 fragments of 1,000 bp length. We called the SNPs within these fragments using the *UnifiedGenotyper* algorithm from *GATK*; the filtering step has been performed as reported in [29] through *vcflib*. We finally calculated the observed FDSS from the quality filtered VCF file.

To investigate past population dynamics of the three Orangutan species, we designed competitive scenarios following the demographic models reported in [29]. Nater [29] first compared models assuming simple demographies, and then estimated the parameters from a more complicated

model. By contrast, we directly compared complex demographies, designing the within-species substructure as described by [29], (Fig 5A). The four competing models indeed share the same within species features (four populations for the Bornean group, two Sumatran populations north of Lake Toba, and a single population south of Lake Toba), while differing for the topology, i.e. for the evolutionary relationships among the three species, as reported in Fig 5A. We modeled bidirectional migration both among populations within a species and between neighboring species. A detailed description of the models' parameters and of the priors are in S13-S16 Tables. We ran 50,000 simulations per model using the *ms* software [41], generating two chromosomes per population (4 Bornean, 1 south of Lake Toba and 2 north of Lake Toba), and 9,000 independent fragments of 1kb length per chromosome. We first assessed the power to distinguish among the four models calculating the proportion of TPs as described above, and then explicitly compared the simulated variation with the FDSS calculated on the observed data (Fig 5B).

Software and data availability

All the scripts of the ABC-SeS framework can be found at https://github.com/anbena/ABC-SeS.

Acknowledgments

We would like to thank the DFG Center for Advanced Studies, ''Words, Bones, Genes, Tools,'' at University of Tübingen, that hosted AB and SG during the first phase of the project. FT was supported by ERC Advanced Grant 295733, 'LanGeLin'.

Author Contributions

AB conceived the study; AB and SG designed the experiments; MTV, AB, SG and FT analyzed the data; SG, MTV, FT, GB and AB discussed the results; SG, GB and AB wrote the paper with inputs from all coauthors.

References

625   1.   Akey JM, Eberle MA, Rieder MJ, Carlson CS, Shriver MD, Nickerson DA, et al. Population

626   history and natural selection shape patterns of genetic variation in 132 genes. PLoS Biol.

627   2004;

628   2.   Meyer D, Single RM, Mack SJ, Erlich HA, Thomson G. Signatures of demographic history

629   and natural selection in the human major histocompatibility complex loci. Genetics. 2006;

630   3.   Lohmueller KE. The Impact of Population Demography and Selection on the Genetic

631   Architecture of Complex Traits. PLoS Genet. 2014;

632   4.   1000 Genomes Project Consortium. An integrated map of genetic variation. Nature. 2012;

633   5.   Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F, et al. The Simons Genome

634   Diversity Project: 300 genomes from 142 diverse populations. Nature. 2016;

635   6.   Meyer M, Kircher M, Gansauge MT, Li H, Racimo F, Mallick S, et al. A high-coverage

636   genome sequence from an archaic Denisovan individual. Science (80-). 2012;

637   7.   Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, et al. The complete

638   genome sequence of a Neanderthal from the Altai Mountains. Nature. 2014;

639   8.   Moreno-Mayar JV, Vinner L, de Barros Damgaard P, de la Fuente C, Chan J, Spence JP, et

640   al. Early human dispersals within the Americas. Science (80-) [Internet]. 2018 Dec

641   7;362(6419):eaav2621.   Available   from:

642   http://science.sciencemag.org/content/362/6419/eaav2621.abstract

643   9.   Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ, et al.

644   Butterfly genome reveals promiscuous exchange of mimicry adaptations among species.

645   Nature. 2012;

646   10.   De Manuel M, Kuhlwilm M, Frandsen P, Sousa VC, Desai T, Prado-Martinez J, et al.

647   Chimpanzee genomic diversity reveals ancient admixture with bonobos. Science (80-).

648   2016;

649   11.   McVean GAT, Cardin NJ. Approximating the coalescent with recombination. Philos Trans R

650   Soc B Biol Sci. 2005;

651   12.   Li H, Durbin R. Inference of human population history from individual whole-genome

652   sequences. Nature. 2011;

653   13.   Schiffels S, Durbin R. Inferring human population size and separation history from multiple

654   genome sequences. Nat Genet. 2014;

655   14.   Heller R, Chikhi L, Siegismund HR. The Confounding Effect of Population Structure on

656   Bayesian Skyline Plot Inferences of Demographic History. PLoS One. 2013;

657   15.   Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Diffusion Approximations

658   for Demographic Inference : DaDi. Nat Preced. 2010;

659   16.   Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. Robust Demographic

660   Inference from Genomic and SNP Data. PLoS Genet. 2013;

661   17.   Akaike H. A New Look at the Statistical Model Identification. IEEE Trans Automat Contr.

662   1974;

663   18.   Terhorst J, Kamm JA, Song YS. Robust and scalable inference of population history from

664   hundreds of unphased whole genomes. Nat Genet. 2017;

665   19.   Bertorelle G, Benazzo A, Mona S. ABC as a flexible framework to estimate demography

666   over space and time: Some cons, many pros. Molecular Ecology. 2010.

667   20.   Beaumont MA, Zhang W, Balding DJ. Approximate Bayesian computation in population

668   genetics. Genetics. 2002;

669   21.   Marjoram P, Molitor J, Plagnol V, Tavare S. Markov chain Monte Carlo without likelihoods.

670   Proc Natl Acad Sci. 2003;

671   22.   Blum MGB, François O. Non-linear regression models for Approximate Bayesian

672   Computation. Stat Comput. 2010;

673   23.   Blum MGB, Nunes MA, Prangle D, Sisson SA. A Comparative Review of Dimension

674   Reduction Methods in Approximate Bayesian Computation. Stat Sci. 2013;

675   24.   Pudlo P, Marin JM, Estoup A, Cornuet JM, Gautier M, Robert CP. Reliable ABC model
676   choice via random forests. Bioinformatics. 2015;

677   25.   Wakeley J, Hey J. Estimating ancestral population parameters. Genetics. 1997;

678   26.   Robinson JD, Bunnefeld L, Hearn J, Stone GN, Hickerson MJ. ABC inference of multi-
679   population divergence with admixture from unphased population genomic data. Mol Ecol.
680   2014;

681   27.   Malaspinas AS, Westaway MC, Muller C, Sousa VC, Lao O, Alves I, et al. A genomic
682   history of Aboriginal Australia. Nature. 2016.

683   28.   Pagani L, Lawson DJ, Jagoda E, Mörseburg A, Eriksson A, Mitt M, et al. Genomic analyses
684   inform on migration events during the peopling of Eurasia. Nature. 2016;

685   29.   Nater A, Mattle-Greminger MP, Nurcahyo A, Nowak MG, de Manuel M, Desai T, et al.
686   Morphometric, Behavioral, and Genomic Evidence for a New Orangutan Species. Curr Biol.
687   2017;

688   30.   Reyes-Centeno H, Ghirotto S, Detroit F, Grimaud-Herve D, Barbujani G, Harvati K.
689   Genomic and cranial phenotype data support multiple modern human dispersals from Africa
690   and a southern route into Asia. Proc Natl Acad Sci. 2014;

691   31.   Tassi F, Ghirotto S, Mezzavilla M, Vilaça ST, De Santi L, Barbujani G. Early modern human
692   dispersal from Africa: Genomic evidence for multiple waves of migration. Investig Genet.
693   2015;

694   32.   Benazzo A, Trucchi E, Cahill JA, Maisano Delser P, Mona S, Fumagalli M, et al. Survival
695   and divergence in a small group: The extraordinary genomic history of the endangered
696   Apennine brown bear stragglers. Proc Natl Acad Sci. 2017;

697   33.   Miller W, Schuster SC, Welch AJ, Ratan A, Bedoya-Reina OC, Zhao F, et al. Polar and
698   brown bear genomes reveal ancient admixture and demographic footprints of past climate
699   change. Proc Natl Acad Sci. 2012;

34.   Beaumont MA. Approximate Bayesian Computation in Evolution and Ecology. Annu Rev

Ecol Evol Syst. 2010;

35.   Csilléry K, Blum MGB, Gaggiotti OE, François O. Approximate Bayesian Computation

(ABC) in practice. Trends in Ecology and Evolution. 2010.

36.   Wright S. Evolution in Mendelian Populations. Genetics [Internet]. 1931 Mar;16(2):97–159.

Available from: https://www.ncbi.nlm.nih.gov/pubmed/17246615

37.   Medina P, Thornlow B, Nielsen R, Corbett-Detig R. Estimating the timing of multiple

admixture pulses during local ancestry inference. Genetics. 2018;

38.   Felsenstein J. Accuracy of coalescent likelihood estimates: Do we need more sites, more

sequences, or more loci? Mol Biol Evol. 2006;

39.   Rowe HC, Renaut S, Guggisberg A. RAD in the realm of next-generation sequencing

technologies. Molecular Ecology. 2011.

40.   Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation

Sequencing Data. BMC Bioinformatics. 2014;

41.   Hudson RR. ms: a Program for Generating Samples Under Neutral Models. Bioinformatics.

2002;

42.   Ewing G, Hermisson J. MSMS: A coalescent simulation program including recombination,

demographic structure and selection at a single locus. Bioinformatics. 2010;

43.   Hinrichs AS, Raney BJ, Speir ML, Rhead B, Casper J, Karolchik D, et al. UCSC Data

Integrator and Variant Annotation Integrator. Bioinformatics. 2016;

44.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;

45.   Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform.

Bioinformatics. 2010;

46.   Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et

al. From fastQ data to high-confidence variant calls: The genome analysis toolkit best

practices pipeline. Curr Protoc Bioinforma. 2013;

47.   Derrien T, Estellé J, Sola SM, Knowles DG, Raineri E, Guigó R, et al. Fast computation and

applications of genome mappability. PLoS One. 2012;

Figures Legend:

Fig 1. One-population models and proportion of True Positives. A) Demographic models compared: Constant, Bottleneck, Expansion, Structured population. N_1 is the effective population size, i the intensity of the bottleneck or of the expansion, T the time of the bottleneck or of the start of the expansion, m is the migration rate. B) True Positives rates. The plot below each of the four models represents the proportion of TPs obtained analyzing pods coming from the above model under 60 combinations of experimental parameters. Different locus lengths are in the x-axes, number of loci is represented by different colors and the number of chromosomes is represented by different symbols.

Fig 2. Two-populations models and proportion of True Positives. A) Demographic models compared: Divergence with isolation, Divergence with migration, Divergence with a single pulse of admixture. N_{anc} is the effective population size of the ancestral population, N_1 and N_2 are the effective population sizes of the diverged populations, T_{sep} is the time of the split, m_{12} and m_{21} the migration rates, T_{adm} is the time of the single pulse of admixture, adm_{12} and adm_{21} the proportions of admixture. B) True Positives rates. The plots have the same features of Fig 1.

Fig 3. Multi-populations models and proportion of True Positives. A) Demographic models compared: Single Dispersal and Multiple Dispersals. The populations sampled are indicated in bold. B) True Positives rates. The plots have the same features of Fig 1.

Fig 4. Posterior Probabilities for the MDM. Left panel: posterior probabilities obtained analyzing 6 Papuan individuals from [28], simulating data without (PWR) or considering (PR) intra-locus

recombination. Right panel: posterior probabilities obtained analyzing 25 Papuan individuals from

[27], simulating data without (MWR) or considering (MR) intra-locus recombination.

Fig 5. Demographic models tested to study the evolutionary history of Orangutan species. A-

Four demographic models compared. The numbers in the black boxes indicate the proportion of TP

calculated analyzing 1,000 pods coming from that demographic model. NT, Sumatran populations

north of Lake Toba; ST, the Sumatran population south of Lake Toba; BO, Bornean populations. B-

Number of votes associated to each model by ABC-RF and posterior probability of the most

supported model (model 1a).

Supporting Informations

Supplementary Figures:

S1 Fig. Example of a frequency distribution of segregating sites. On the x-axes: the number of

segregating sites (here from 0 to 20). On the y-axes: the number of genomic loci showing a certain

number of segregating sites. We calculated four FDSS for each pair of populations compared

(representing the genomic distribution of segregating sites private of the first population, private of

the second population, shared, and fixed for different alleles). In the one-population models we used

a single FDSS.

S2 Fig. Proportion of True positives for the one-population structured model as a function of

the migration rate (A) and the number of demes considered (B). (A) Each plot represents the

proportion of pods from the structured model assigned to each of the four one-population models

with the migration rates among demes in the structured model constrained at ranges of increasing

values (from $1*10^{-5}$ to $1*10^{-1}$). All the plots consider two chromosomes and a specific combination

of locus length and number of loci; the number of demes in the structured model is fixed to four. In

general, the TP rate (in dark blue) decreases as increasing the migration rate among demes, with the

constant model erroneously recognize as the true model for higher migration rates. (B) Proportion

of pods from the structured model assigned to each of the four one-population models as a function

of the number of demes (from 2 to 10). The TP rate increase with the number of demes, regardless

of the level of migration among demes.

S3 Fig. FDSS generated under the one-population models for each number of chromosomes

tested. Each plot represents the FDSS simulated under each of the four one-population models

considering 1,000 fragments of 1,000 base pair length, for a specific number of chromosomes

sampled. Going from two to twenty chromosomes, we observe an increase of the overlapping

between the FDSS generated under the Constant and the Structured model, thus possibly explaining

the decrease in the models' identifiability as increasing the number of chromosomes considered.

S4 Fig. Proportion of True Positives for the one-population models with recombination. The

plots have the same features of Fig 1, we considered a per nucleotide per generation recombination

rate of $1*10^{-8.}$

S5 Fig. Proportion of True Positives for the Pulse of Admixture model as a function of the

time span between the divergence and the admixture. We expressed the x-axes as (T_{sep}-

T_{adm})/T_{sep}, so as to normalize the results respect to the age of the divergence. Each plot represents

the results for a different locus length.

S6 Fig. Proportion of True Positives for the two-population models with recombination. The

plots have the same features of Fig 1, we considered a per nucleotide per generation recombination

rate of $1*10^{-8}$

S7 Fig. Proportion of True Positives for the MDM a function of the time span between the

divergence time of the African ghost populations and the second exit (Delta tdYG-OOA2). The

time difference between the divergence time of the African ghost populations and the second exit

from Africa is on the x-axes and it is expressed in years (considering a generation time of 29 years).

Each plot reports the results for a different locus length.

S8 Fig. Expected topologies under the SDM and MDM. T1, T2 and T3 are baseline topologies,

of which T1 is shared between SDM and MDM, whereas T2 and T3 are exclusively originated by

MDM. R is the recombinant topology expected under MDM.

Supplementary Tables:

S1 Table. Demographic parameters and prior distributions of one-population models.

Mutation and Recombination rates are expressed per nucleotide per generation.

S2 Table. Demographic parameters and prior distributions of two-population models.

Mutation and Recombination rates are expressed per nucleotide per generation. Time is in

generations.

S3 Table. RF Prior Error Rates using the complete FDSS or its first four moments. We

calculated the Prior Error Rates under the complete SDFF and under a summary of the distribution

(i.e. the first four moments). We analyzed ten combinations of experimental parameters (indicated

in the "ID" column), comparing the three two-population divergence models. ll= locus length, nl=

number of loci, r= recombination rate, nc= number of chromosomes. The Prior Error Rate is a

measure of the global quality of the RF classifier (the lower the better), which depends on the

summary statistics used. The complete FDSS always generate lower values of Prior Error Rates.

S4 Table. Demographic parameters and prior distributions of multi-population models:

Single Dispersal model. Migration and admixture rates are expressed per generation, times in

years. We considered a generation time of 29 years as in [27]. Per nucleotide per generation

mutation and recombination rates are fixed as in [27].

S5 Table. Demographic parameters and prior distributions of multi-population models:

Multiple Dispersals model. Migration and admixture rates are expressed per generation, times in

years. We considered a generation time of 29 years as in [27]. Per nucleotide per generation

mutation and recombination rates are fixed as in [27].

S6 Table. Power Analysis of the Multi-Populations models. Comparison of the results with

and without intra-locus recombination for the four combinations of experimental parameters

tested. The first four rows present the results without considering intra-locus recombination,

whereas the last four rows refer to the same combinations of experimental parameters simulated

considering a per nucleotide per generation recombination rate of $1.12*10^{-8}$. TP% columns

represent the percentage of True Positives calculated analyzing 1,000 pods coming from SDM and

MDM. SDM mpp and MDM mpp represent the mean posterior probability value obtained for the

datasets that have been correctly assigned to the true model.

S7 Table. Genomes used for the comparison of SDM and MDM using real data.

S8 Table. Model selection results using Papuan individuals from [28] without intra-locus

recombination. The first column represents the id of the Papuan sample used in that comparison as

reported in the dataset. The second column shows the model selected by the ABC-SeS procedure.

The third and the fourth columns represent the proportion of votes assigned to SDM and MDM by

the RF algorithm. The last column is the posterior probability of the most supported model.

S9 Table. Model selection results using Papuan individuals from [28] considering intra-locus

recombination. The first column represents the id of the Papuan sample used in that comparison as

reported in the dataset. The second column shows the model selected by the ABC-SeS procedure.

The third and the fourth columns represent the proportion of votes assigned to SDM and MDM by

the RF algorithm. The last column is the posterior probability of the most supported model.

S10 Table. Model selection results using Papuan individuals from [27] without intra-locus

recombination. The first column represents the id of the Papuan sample used in that comparison as

reported in the dataset. The second column shows the model selected by the ABC-SeS procedure.

The third and the fourth columns represent the proportion of votes assigned to SDM and MDM by

the RF algorithm. The last column is the posterior probability of the most supported model.

S11 Table. Model selection results using Papuan individuals from [27] considering intra-locus

recombination. The first column represents the id of the Papuan sample used in that comparison as

reported in the dataset. The second column shows the model selected by the ABC-SeS procedure.

The third and the fourth columns represent the proportion of votes assigned to SDM and MDM by

the RF algorithm. The last column is the posterior probability of the most supported model.

S12 Table. Genomes used for the comparison of the four Orangutan evolutionary scenarios.

S13 Table. Demographic parameters and prior distributions for Model 1a. Migration rates are

expressed per generation, times in years. We used a generation time of 25 years as in [29]. The per

nucleotide per generation mutation rate is fixed as in [29].

S14 Table. Demographic parameters and prior distributions for Model 2a. Migration rates are

expressed per generation, times in years. We used a generation time of 25 years as in [29]. The per

nucleotide per generation mutation rate is fixed as in [29].

S15 Table. Demographic parameters and prior distributions for Model 1b. Migration rates are

expressed per generation, times in years. We used a generation time of 25 years as in [29]. The per

nucleotide per generation mutation rate is fixed as in [29].

S16 Table. Demographic parameters and prior distributions for Model 2b. Migration rates are

expressed per generation, times in years. We used a generation time of 25 years as in [29]. The per

nucleotide per generation mutation rate is fixed as in [29].

A

B

Selected Model	Votes model 1A	Votes model 2A	Votes model 1B	Votes model 2B	PP
1A	0.398	0.190	0.292	0.120	0.489

Figure 5

Figure 1

Figure 2

Figure 3

Figure 4