Distinct molecular etiologies of male and female hepatocellular carcinoma

Heini M. Natri,[1]* Melissa A. Wilson,[1] Kenneth Buetow[1]

[1] Center for Evolution and Medicine, School of Life Sciences, Arizona State University, Tempe,
Arizona, USA

Corresponding author:

E-mail: hnatri@asu.edu (H.M.N)

The authors declare no potential conflicts of interest.

Abstract

Sex-differences in cancer occurrence and mortality are evident across tumor types; men exhibit higher rates of incidence and often poorer responses to treatment. Targeted approaches to the treatment of tumors that account for these sex-differences require the characterization of the biological mechanisms that differentiate them. Hepatocellular carcinoma (HCC) is the second leading cause of cancer death worldwide, with the incidence rapidly rising. HCC exhibits a male-bias in occurrence and mortality, but previous studies have failed to explore the sex-specific dysregulation of gene expression in HCC. Here, we characterize the sex-shared and sex-specific regulatory changes in HCC. By sex-specific analysis of tumor and tumor-adjacent samples, we uncovered etiologically relevant genes, pathways, and networks differentiating male and female HCC. While both sexes exhibited activation of pathways related to apoptosis and p53 signaling, pathways involved in innate and adaptive immunity showed differential activation between the sexes. Using eQTL analyses, we discovered germline regulatory variants with differential effects on tumor gene expression between the sexes. 24.3% of the discovered eQTLs exhibit differential effects between the sexes, illustrating the role of sex in modifying the effects of regulatory variants in HCC. The genes that showed sex-specific dysregulation and harbored a sex-specific eQTL converge in canonical pathways, suggesting that the molecular etiologies of male and female HCC are partially driven by differential genetic effects on gene expression. Overall, our results provide insights into the role of genetic regulation of transcription in modulating sex-differences in HCC etiology and provide a framework for future studies on sex-biased cancers.

Introduction

Differences in cancer occurrence and mortality between sexes are evident across tumor types; males exhibit higher rates of cancer incidence and often poorer response to treatment, including some forms of chemotherapy and immunotherapy (1,2). While differences in risk factors may explain some portion of the sex-bias, the bias remains after appropriate adjustment for these factors (3,4). A recent study examining the mutational profiles of tumors from males and females across The Cancer Genome Atlas (TCGA) found sex-differences in cancer driver genes and biomarkers, calling for the consideration of sex as a biological variable in studies on cancer

occurrence, etiology, and treatment (5). Despite these underlying molecular differences, sex is

rarely considered in the development of cancer therapies.

Across tumor types analyzed, the largest sex-differences in autosomal mutational profiles were

seen in liver hepatocellular carcinoma (HCC), indicating that male and female HCC are

etiologically distinct (5). Furthermore, HCC exhibits sex-bias in occurrence, with a male-to-

female incidence ratio between 1.3:1 and 5.5:1 across populations (6,7). The sexes also differ in

the clinical manifestation of HCC, males exhibiting an earlier onset and more/larger nodules (8).

HCC is the second leading cause of cancer mortality worldwide, accounting for 8.2% of all

cancer deaths (2), and the incidence in the US has doubled in the last 3 decades, attributable to

increased rates of obesity (7), calling for the development of new interventions and targeted

therapies.

Sex-specific gene regulation may partially underlie differences between the sexes in disease

prevalence and severity (9,10). Previous work observed extensive sex-biased signatures in gene

expression in HCC and other sex-biased cancers (11). However, this study focused solely on

comparing male and female tumor samples, while ignoring sex-differences in healthy and tumor-

adjacent tissues. To understand cancer-specific processes, it is necessary to contrast the sex-

differences in gene expression identified in HCC with those in healthy and tumor tissues. For the

targeted treatment of tumors, it is necessary to understand whether sex differences in cancer

reflect unique cancer-specific changes, or are reflective of healthy sex-differences that may

underlie observed sex-bias in cancer occurrence and disease etiology.

In addition to sex differences in overall gene expression due to the wide effects of sex as a

biological variable, genetic variants may alter gene expression in a sex-specific manner. A pan-

cancer analysis of the TCGA dataset identified 128 germline variants altering gene expression

levels (eQTLs) in HCC (12). However, this study purposefully controlled for and removed the

effect of sex and, to date, a sex-specific eQTL analysis in HCC has not been performed. Sex-

stratified analyses can reveal sex-biased genetic effects on gene expression that may be obscured

in a joint analysis of both sexes - e.g. cases where the regulatory variant has a zero or very small

effect in one sex, or the eQTL exhibits an opposite effect direction in the two sexes (13). eQTLs

that are discovered in one sex but not in the whole sample analysis are likely to affect gene

expression in a sex-dependent manner, and while a combined analysis of both sexes achieves a

greater statistical power to detect sex-shared effects, it dilutes the signal of sex-dependent effects

(14).

Targeted approaches to the treatment of male and female HCC require the characterization and

understanding of the fundamental biological mechanisms that differentiate them. Here, we

analyzed data from TCGA and The Genotype-Tissue Expression project (GTEx) to examine the

sex-specific patterns of gene expression and regulation in HCC. Here, we have contrasted the

sex-biased patterns of gene expression in HCC tumors with healthy and tumor-adjacent liver

tissues, allowing us to detect sex-differences in gene expression shared between and specific to

the different tissues. We show that male and female HCC exhibit differences in the dysregulation

of genes and the germline genetic regulation of tumor gene expression. Importantly, these

orthogonal approaches identify genes that converge in shared pathways, indicating sex-specific

regulatory etiology in HCC. The results presented here have implications for the development of

targeted therapies for male and female HCC.

Materials and Methods

Data

GTEx (release V6p) whole transcriptome (RNAseq) data (dbGaP accession #8834) were

downloaded from dbGaP. TCGA LIHC Affymetrix Human Omni 6 array genotype data, whole

exome sequencing (WES) and RNAseq data (dbGaP accession #11368) were downloaded from

NCI Genomic Data Commons (15). In total, RNAseq data from 91 male and 45 female GTEx

donors, germline genotypes and tumor RNAseq data from 248 male and 119 female TCGA

LIHC donors, as well as paired tumor and tumor-adjacent samples from 28 male and 22 female

TCGA LIHC donors were utilized in this study. FASTQ read files were extracted from the

TCGA LIHC WES BAM files using the *strip_reads*() function of *XYAlign* (16). We used

FastQC (17) to assess the WES and RNAseq FASTQ quality. Reads were trimmed using

TRIMMOMATIC IlluminaClip (18), with the following parameters: seed mismatches 2,

palindrome clip threshold 30, simple clip threshold 10, leading quality value 3, trailing quality

value 3, sliding window size 4, minimum window quality 30 and minimum read length of 50.

Read mapping and read count quantification

Sequence homology between the X and Y chromosomes may cause the mismapping of short
sequencing reads derived from the sex chromosomes and affect downstream analyses (16). To
overcome this, reads were mapped to custom sex-specific reference genomes using *HISAT2* (19).
Female samples were mapped to the human reference genome GRCh38 with the Y-chromosome
hard-masked. Male samples were mapped to the human reference genome with Y-chromosomal
pseudoautosomal regions hard-masked. Gene-level counts from RNAseq were quantified using
Subread featureCounts (20). Reads overlapping multiple features (genes or RNA families with
conserved secondary structures) were counted for each feature.

Germline variant calling and filtering

BAM files were processed according to Broad Institute *GATK* (*Genome Analysis Toolkit*) best
practices (21–23): Read groups were added with *Picard Toolkit*'s *AddOrReplaceReadGroups*
and optical duplicates marked with *Picard Toolkit*'s *MarkDuplicates* (v.2.18.1,
http://broadinstitute.github.io/picard/). Base quality scores were recalibrated with *GATK*
(v.4.0.3.0) *BaseRecalibrator*. Germline genotypes were called from whole blood Whole Exome
Sequence samples from 248 male and 119 female HCC cases using the scatter-gather method
with *GATK HaplotypeCaller* and *GenotypeGVCFs* (21). Affymetrix 6.0 array genotypes were
lifted to GRCh38 using the UCSC *LiftOver* tool (24) and converted to VCF. Filters were applied
to retain variants with a minimum quality score >30, minor allele frequency >10%, minor allele
count >10, and no call rate <10% across all samples.

Clinical characteristics and cellular content of tumor samples

Confounding effects, e.g. differences in clinical and pathological characteristics or cell type
composition of the sequenced samples, may contribute to the observed effect modification when
utilizing stratified analyses. We examined the differences in the clinical characteristics between
males and females in the TCGA LIHC cohort. We used a *t*-test to test for the equality of means
in patient age, and Fisher's exact test to test to detect differences in risk factors and pathological
classifications (Supplementary Table S1). To examine the cellular composition of tumor

124   samples, we performed cell type enrichment analysis for 64 immune and stromal cell types using

125   *xCell* (25).

126   Filtering of gene expression data

FPKM (Fragments Per Kilobase of transcript per Million mapped reads) expression values for

each gene were obtained using *EdgeR* (26). Each expression dataset was filtered to retain genes

with mean FPKM>0.5 and read count of >6 in at least 10 samples across all samples under

investigation. In the comparative analysis of differentially expressed genes (DEGs) between the

tumor vs. tumor-adjacent samples in males, females, and both sexes, genes that reached the

previously described expression thresholds in at least one tissue in at least one sex were retained.

This assures that the DEGs detected in the sex-specific and combined analyses are not due to

filtering.

135   Differential expression analysis

For differential expression (DE) analysis, filtered, untransformed read count data were quantile

normalized and logCPM transformed with *voom* (27). From the TCGA LIHC dataset, paired

tumor and tumor-adjacent samples were available for 22 females and 28 males. From the GTEx

liver dataset, 91 male and 45 female samples were used in the DE analysis. A multi-factor design

with sex and tissue type as predictor variables were used to fit the linear model.

duplicateCorrelation function was used to calculate the correlation between measurements made

on the same subject, and this inter-subject correlation was accounted for in the linear modeling.

DEGs between comparisons were identified using the *limma*/*voom* pipeline (27) by computing

empirical Bayes statistics with *eBayes*. An FDR-adjusted *p*-value threshold of 0.01 and an

absolute $\log_2$ fold-change (FC) threshold of 2 were used to select significant DEGs.

To reliably detect genes that are expressed in a sex-biased way in HCC but not in healthy liver or

in tumor-adjacent tissue, we examined genes that were DE in the male vs. female tumor

comparison using the previously described significance thresholds, but not in the male vs. female

comparisons of normal or tumor-adjacent samples with a relaxed significance threshold of FDR-

adjusted *p*-value <0.1 and absolute $\log_2(FC) > 0$.

To detect genes that are dysregulated in tumors compared to matched tumor-adjacent samples in

each sex, we identified DEGs in the tumor vs. tumor-adjacent comparison of males, females, and

in the whole sample. DEGs that were identified in one sex but not in the other or in the combined

analysis of both sexes were considered sex-specific. DEGs identified in the combined analysis

were considered sex-shared. This approach allows the identification of high-confidence sex-

specific events that are a result of the underlying biological differences as opposed to sampling

or statistical power. ANOVA and Kruskal-Wallis tests were used to test for equality of fold

changes of sex-shared and sex-specific DEGs across male, female, and all samples.

Overrepresentation of biological functions and canonical pathways

We further analyzed the sex-shared and sex-specific tumor vs. tumor-adjacent DEGs as well as

the sex-specific eQTL target genes (eGenes) to identify sex-shared and sex-specific pathways

driving HCC etiology. We used the Gene Ontology Consortium's web tool (28) which utilizes

the PANTHER classification system (29) and a binomial test (30) to determine whether a list of

genes is overrepresented in pathways in regards to molecular functions, biological processes, and

cellular functions. Fisher's exact test was used to calculate overrepresentation scores. We used

the *NetworkAnalyst* webtool (31), which utilizes a hypergeometric test to compute *p*-values for

the overrepresentation of genes in regards to KEGG and Reactome pathways. An FDR-adjusted

p-value threshold of 0.01 was used to select significantly overrepresented GO terms and

canonical pathways.

eQTL analysis

We used eQTL analyses to detect germline genetic effects on tumor gene expression. Similar to

the DE analysis, we utilized combined and sex-stratified analyses to detect sex-shared and sex-

specific effects. Germline genotypes and tumor gene expression data from 248 male and 119

female donors in the TCGA LIHC cohort were used in the eQTL analysis. Filtered count data

was normalized by fitting the FPKM values of each gene and sample to the quantiles of the

normal distribution. To account for technical confounders and population structure, 10 *de novo*

PEER factors and three genotype principal components were used as covariates (Supplementary

Methods). *Cis*-acting (proximal) eQTLs were detected by linear regression as implemented in

QTLtools v.1.1 (32). Variants within 1Mb of the gene under investigation were considered for

testing. We used the permutation pass with 10,000 permutations to get adjusted *p*-values for

associations between the phenotypes and the top-variants in *cis*. FDR-adjusted *p*-values were

calculated to correct for multiple phenotypes tested, and an adjusted *p*-value threshold of 0.01

was used to select significant associations. To allow the comparison of effect sizes of sex-

specific and sex-shared eQTLs across the sexes, effects of each variant located within the 1Mb

cis-window were obtained using the *QTLtools* nominal pass.

Similarly to the tumor vs. tumor-adjacent DEGs, eQTLs that were detected in one sex but not in

the other or in the combined analysis were considered sex-specific, while eQTLs detected in the

combined analysis were considered sex-shared. ANOVA and Kruskal-Wallis tests were used to

test for equality of effect sizes of sex-shared and sex-specific eQTLs across male, female, and all

samples.

191   Results

192   Sex-specific patterns of gene expression in HCC

We identified sex-differences in gene expression in healthy liver (GTEx; 21 sex-biased genes),

tumor-adjacent tissue (TCGA LIHC; 72), and HCC (TCGA LIHC; 24) to characterize the shared

and unique sex-differences that may drive the observed sex-biases in HCC occurrence and

etiology (Fig. 1, Supplementary Tables S2-4). X-linked *XIST* and Y-linked genes were expressed

in a sex-biased way across all tissues. While sex-biased gene expression in healthy and tumor-

adjacent tissues may contribute to the sex-differences in cancer occurrence, sex-biased

expression in tumors is suggestive of distinct molecular etiologies of male and female HCC. We

identified 13 genes, including 7 protein-coding genes (*CD24*, *CYP4F22*, *DTX1*, *HRCT1*,

HSD11B1, *PIGR*, *ZCCHC16)* and 6 non-coding RNAs or pseudogenes (*ADORA2BP1*, *CTD-*

2325A15.3, *RP11-431J24.2*, *RP11-495P10.5*, *RP11-495P10.7*, *RP11-495P10.9*) that show sex-

differences in expression in HCC, but not in tumor-adjacent tissue or healthy liver, even with a

relaxed significance threshold (Fig. 1A). Notably, Notch-regulating *DTX1* (Fig. 1B) and signal

transducer *CD24* were downregulated in male HCC.

To further examine the sex-shared and sex-specific mechanisms driving HCC etiology, we

detected DEGs between tumor and tumor-adjacent samples in males and females, as well as in

the combined analysis of both sexes. Dimensionality reduction of gene expression data shows

that variation among the tumor and tumor-adjacent samples is driven by tissue type and sex (Fig.

1C). When inspecting the tumor samples only, the first dimension is driven by sex

(Supplementary Fig. S1). In the combined analysis of male and female samples, we detected 581

tumor vs. tumor-adjacent DEGs (Supplementary Table S5). In male- and female-specific

analyses, we detected 606 and 416 tumor vs. tumor-adjacent DEGs, respectively (Supplementary

Tables S6-7). Out of the total of 756 unique DEGs, 77% were shared between the sexes. We

identified 83 female-specific and 92 male-specific tumor vs. tumor-adjacent DEGs. Notably,

substantially more DEGs were detected in sex-specific analyses than in the unstratified analysis

(Fig. 1D). Specifically, DEGs that showed different magnitudes in fold change between the sexes

(based on ANOVA/Kruskal-Wallis tests) were detected in the sex-specific analyses, while DEGs

with similar fold changes across all comparisons were detected in the combined analysis as well

as the sex-specific analyses (Supplementary Fig. S2). Sex-shared DEGs that were only detected

in the combined analysis, and not in the sex-specific analyses, showed a large variance in

expression and, due to limited power, were not detected as statistically significant DEGs in sex-

specific analyses (Supplementary Fig. S2). Tumor-infiltrating immune cells may produce

spurious signals in DE analyses, which is evident from the detection of various immunoglobulin

genes in tumor vs. tumor-adjacent comparisons (Supplementary Tables S5-7). However, cell

content analyses based on gene expression data did not exhibit sex-differences in the prevalence

of tumor-infiltrating immune cells in this data (Supplementary Fig. S3), and thus such spurious

signals are unlikely to affect male-female comparisons. Furthermore, male and female cases did

not differ in terms of tumor pathology (Supplementary Table S1). The observed differences in

gene expression are thus likely to reflect actual sex-differences rather than confounding

differences in sample characteristics or composition.

To put these results in a broader context, we analyzed the male- and female-specific DEGs

(tumor vs. tumor-adjacent) for the overrepresentation of functional pathways and canonical

networks. We found that the sex-shared and sex-specific DEGs were enriched in pathways

relevant to oncogenesis and cancer progression (Supplementary Tables 8-10). Most of the sex-

specific top pathways were non-overlapping, indicating that male and female HCC are driven by

different mechanisms and processes (Fig. 1E-F).

Differential cis-eQTL effects in male and female HCC

To further investigate the mechanisms of sex-difference in HCC etiology, we used eQTL analyses to detect germline genetic effects on tumor gene expression in both the joint and sex-stratified analyses (Fig. 2A). We detected 1,204, 761, and 245 eQTLs in the combined, male-specific, and female-specific analyses, respectively (Supplementary Tables S11-13). As expected, genomic annotations show that most eQTLs are located on non-coding regions (Supplementary Methods, Fig. 2B). Consistent with previous reports, most *cis*-eQTLs were located near transcription start sites (TSSs), with 63% of all eQTLs across the combined and sex-specific analyses being located within 20kb of TSSs. On average, 384 variants were tested per gene. 31% of the unique shared and sex-specific cis-eQTLs in HCC were also identified as eQTLs in the liver data in the GTEx project, indicating shared tissue origin. Out of the total of 1,595 unique associations, 75.7% were shared between the sexes. We detected 295 male-specific and 92 female-specific eQTLs. Since these associations were not detected in the unstratified analysis, they are likely not a result of differential power to detect associations due to different sample sizes, but exhibit effect modification by sex. Sex-specific associations exhibited differences in effect size between the sexes (based on ANOVA/Kruskal-Wallis tests), and the sex-specific effect is diluted in the combined analysis (Supplementary Fig. S6). Sex-shared large effect eQTLs were detected in sex-specific and combined analyses, and, due to the larger sample size, sex-shared low-effect eQTLs are detected in the combined analysis only (Supplementary Fig. S6).

We detected 27 shared eGenes that were regulated by independent regulatory variants in males and females. This could be due to actual biological differences in gene regulation, or due to technical constraints, in particular, missing genotypes in one sex affecting the permutation scheme to select the top-variant for each target gene. To overcome this and to detect high confidence instances of differential gene regulation between the sexes, we further examined the sex-shared and sex-specific eGenes: we found 24 genes that are under germline regulatory control in only male HCC (Fig. 2C), including *POGLUT1*, which is an essential regulator of Notch signaling (Fig. 2D). No genes were found to be under regulatory control in females only, likely due to reduced power to detect associations in females (Supplementary Fig. S5). Male-specific eGenes were overrepresented in pathways related to cell cycle, apoptosis, and cancer

(Supplementary Table S17). Concordant with previous studies (14,33), none of the male-specific

eGenes were differentially expressed between male and female HCC, indicating that the male-

specific eQTLs are not a result of differences in overall gene expression levels between males

and females, but are likely to arise from factors such as differential chromatin accessibility or

transcription factor activity. The observation that the sex-biased autosomal genes in tumors did

not harbor significant *cis*-eQTLs (Supplementary Table S18) also suggests that the sex-biased

gene expression is a result of trans-effects, e.g. sex-chromosomal effects on autosomal gene

expression, or, more widely, a result of sex as a biological variable, e.g. hormonal effects.

Discussion

Distinct molecular etiologies of male and female HCC

It is well established that patterns of gene expression vary between the sexes across different

tissues. Previous studies have confounded these differences with those which may be driving

etiological differences between male and female tumors. For example, Yuan et al. previously

reported extensive sex-biased signatures in gene expression in HCC and other strongly sex-

biased cancers (11). While they identified immunity and cancer-associated enriched pathways

based on sex-biased genes detected in HCC tumors, their approach was limited as it did not

include the examination of tumor-adjacent tissues. From the results presented here, we are able

to distinguish the differences detected in comparisons of male and female HCC from those

reflecting sex-differences in the healthy liver or in tumor-adjacent tissue, as well as to detect

genes that are dysregulated in HCC in a sex-shared or sex-specific manner.

We characterized differences in gene expression between male and female HCC cases. Notably,

sex-differences in gene expression were the largest in the tumor-adjacent tissue, with 72 genes

(including 20 autosomal genes) being expressed in a sex-biased way. These sex-differences point

to distinct mechanisms underlying HCC oncogenesis between the sexes, and may partially

underlie the observed sex-biases in HCC occurrence and onset. We detected 13 genes that were

expressed in a sex-biased way in HCC tumors, but not in healthy or tumor-adjacent liver tissues.

Some of these genes are of particular interest in the context of HCC: Notch-regulating *DTX1*,

found here to be underexpressed in males compared to females, has been identified as a putative

tumor suppressor gene in head and neck squamous cell carcinoma (34). Another female-biased

gene detected here, *CD24*, has a crucial role in T cell homeostasis and autoimmunity (35). The

opposing roles of *CD24* expression in cancer and autoimmune diseases raise interesting

questions on the role of sex-differences in immunity underlying sex-differences in cancer. Future

studies will focus on better understanding the differential regulation of immune functions

between the sexes, and how these differences contribute to the observed biases in disease

occurrence and etiology.

By sex-specific analyses of matched tumor and tumor-adjacent samples, we detected genes that

are uniquely dysregulated in male and female HCC. Further examination of these genes revealed

sex-differences in the pathway and network activation, indicating that the molecular etiologies of

male and female HCC are partly driven by distinct functional pathways. Males and females

differed in oncogenic processes with females showing RAS pathway enrichment while males

showed PI3K/AKT pathway enrichment (Fig. 1E, Supplementary Tables 9-10). While both sexes

exhibited an enrichment of genes involved in cell cycle, apoptosis and p53 signaling, only males

showed activation of pathways regulating to natural killer cell cytotoxicity, and females showed

an activation of pathways involved in NF-kB (nuclear factor kappa-light-chain-enhancer of

activated B cell) activation. Furthermore, female-specific DEGs were enriched in genes involved

in innate immunity Toll-like receptor (TLR) signaling, while male-specific DEGs exhibited an

enrichment of genes involved in adaptive immunity and Notch signaling. TLR signaling has a

crucial role in the regulation of the immune system by evoking an inflammatory response (36).

As TLR and Notch signaling pathways are notable targets for anti-cancer and anti-metastasis

therapies (37,38), the results presented here have implications to the targeted treatment of male

and female HCC.

Sex-specific germline genetic effects on tumor gene expression may drive the molecular

etiologies of male and female HCC

Sex-specific regulatory functions may underlie sex-differences in cancer etiology, progression,

and outcome. We detected sex-differences in the germline genetic regulation of tumor gene

expression in HCC, including 24 genes that were under germline regulatory control only in male

HCC (Fig. 2). Functional annotations of these male-specific eGenes provide insight into possible

regulatory mechanisms contributing to the observed male-bias in HCC and sex-differences in

HCC etiology. Protein O-glucosyltransferase 1 (*POGLUT1*) was found to be under germline

regulation in male HCC, but not in female HCC or in the joint analysis of both sexes (Fig. 3D).

The eQTL associated with *POGLUT1* is located on a promoter region of its target

(Supplementary Table S15). *POGLUT1* is an enzyme that is responsible for O-linked

glycosylation of proteins. Altered glycosylation of proteins has been observed in many cancers

(39,40), including liver cancer (41,42). *POGLUT1* is an essential regulator of Notch signaling

and is likely involved in cell fate and tissue formation during development. Genes involved in

Notch signaling were also found to be expressed in a sex-biased way in HCC tumors and highly

overrepresented among the male-specific DEGs detected in the tumor vs. tumor-adjacent

comparison, showing that sex-specific eQTLs and sex-specific dysregulated genes converge in

canonical pathways. While the Notch signaling pathway was not detected as statistically

significantly overrepresented (FDR-adj. p-value < 0.01) among the 24 male-specific eGenes,

these genes exhibited an enrichment of genes belonging to the KEGG Notch signaling pathway

(4 out of 47 genes detected, 0.542 expected, FDR-adj. $p=0.06$) and Reactome term "NOTCH1

Intracellular Domain Regulates Transcription" (4 out of 50 genes detected, 0.412 expected,

FDR-adj. $p=0.05$). These results point to a possible role of Notch in the development of HCC in

males. Notch signaling is of particular interest in the context of HCC, as it is involved in liver

development (43,44), the development of sexually dimorphic traits (45), and tumorigenesis (46).

The role of Notch signaling in HCC may differ between early and late-stage tumors and among

molecular subtypes, and further studies are necessary to understand the possible oncogenic

properties of Notch among HCC subtypes and between the sexes.

In summary, we discovered differential genetic and regulatory functions in HCC tumors between

the sexes. This work provides a framework for future studies on sex-biased cancers. We are

calling for the generation of suitable large datasets and further studies to identify and validate

sex-specific genetic effects on tumor gene expression and its consequences in HCC and other

sex-biased cancers.

Acknowledgments

353 This study was supported by ASU Center for Evolution and Medicine postdoctoral fellowship

354 and the Marcia and Frank Carlucci Charitable Foundation postdoctoral award from the Prevent

Cancer Foundation for HMN, ASU School of Life Sciences startup funds for MAW, and ASU Center for Evolution and Medicine Venture funds. We thank Dr. Nicholas Banovich and three anonymous reviewers for their feedback on previous versions of this manuscript.

Citations

1. Clocchiatti A, Cora E, Zhang Y, Dotto GP. Sexual dimorphism in cancer. Nat Rev Cancer. 2016;16:330.

2. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin [Internet]. 2018; Available from: http://dx.doi.org/10.3322/caac.21492

3. Wisnivesky JP, Halm EA. Sex differences in lung cancer survival: do tumors behave differently in elderly women? J Clin Oncol. 2007;25:1705–12.

4. OuYang P-Y, Zhang L-N, Lan X-W, Xie C, Zhang W-W, Wang Q-X, et al. The significant survival advantage of female sex in nasopharyngeal carcinoma: a propensity-matched analysis. Br J Cancer. 2015;112:1554–61.

5. Li CH, Haider S, Shiah Y-J, Thai K, Boutros PC. Sex Differences in Cancer Driver Genes and Biomarkers. Cancer Res. 2018;78:5527–37.

6. Wands J. Hepatocellular carcinoma and sex. N Engl J Med. 2007;357:1974–6.

7. Petrick JL, Braunlin M, Laversanne M, Valery PC, Bray F, McGlynn KA. International trends in liver cancer incidence, overall and by histologic subtype, 1978-2007. Int J Cancer. 2016;139:1534–45.

8. Ladenheim MR, Kim NG, Nguyen P, Le A, Stefanick ML, Garcia G, et al. Sex differences in disease presentation, treatment and clinical outcomes of patients with hepatocellular carcinoma: a single-centre cohort study. BMJ Open Gastroenterol. 2016;3:e000107.

9. Gilks WP, Abbott JK, Morrow EH. Sex differences in disease genetics: evidence, evolution, and detection. Trends Genet. 2014;30:453–63.

10. Morrow EH. The evolution of sex differences in disease. Biol Sex Differ. 2015;6:5.

11. Yuan Y, Liu L, Chen H, Wang Y, Xu Y, Mao H, et al. Comprehensive Characterization of Molecular Differences in Cancer between Male and Female Patients. Cancer Cell. 2016;29:711–22.

12. Gong J, Mei S, Liu C, Xiang Y, Ye Y, Zhang Z, et al. PancanQTL: systematic identification
of cis-eQTLs and trans-eQTLs in 33 cancer types. Nucleic Acids Res. 2018;46:D971–6.

13. Behrens G, Winkler TW, Gorski M, Leitzmann MF, Heid IM. To stratify or not to stratify:
power considerations for population-based genome-wide association studies of quantitative
traits. Genet Epidemiol. 2011;35:867–79.

14. Dimas AS, Nica AC, Montgomery SB, Stranger BE, Raj T, Buil A, et al. Sex-biased genetic
effects on gene regulation in humans. Genome Res. 2012;22:2368–75.

15. Grossman RL, Heath AP, Ferretti V, Varmus HE, Lowy DR, Kibbe WA, et al. Toward a
Shared Vision for Cancer Genomic Data. N Engl J Med. 2016;375:1109–12.

16. Webster TH, Couse M, Grande BM, Karlins E, Phung TN, Richmond PA, et al. Identifying,
understanding, and correcting technical biases on the sex chromosomes in next-generation
sequencing data [Internet]. bioRxiv. 2018 [cited 2018 Dec 16]. page 346940. Available from:
https://www.biorxiv.org/content/early/2018/07/18/346940

17. Andrews S. FastQC A Quality Control tool for High Throughput Sequence Data [Internet].
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. 2010. Available from:
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

18. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30:2114–20.

19. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. Nat Methods. 2015;12:357–60.

20. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. Bioinformatics. 2014;30:923–30.

21. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20:1297–303.

22. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet.
2011;43:491–8.

23. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, et
al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best
practices pipeline. Curr Protoc Bioinformatics. 2013;43:11.10.1–33.

24. Hinrichs AS, Karolchik D, Baertsch R, Barber GP, Bejerano G, Clawson H, et al. The UCSC
Genome Browser Database: update 2006. Nucleic Acids Res. 2006;34:D590–8.

25. Aran D, Hu Z, Butte AJ. xCell: digitally portraying the tissue cellular heterogeneity
landscape. Genome Biol. 2017;18:220.

26. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26:139–40.

27. Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model analysis
tools for RNA-seq read counts. Genome Biol. 2014;15:R29.

28. Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, et al. PANTHER version 11:
expanded annotation data from Gene Ontology and Reactome pathways, and data analysis
tool enhancements. Nucleic Acids Res. 2017;45:D183–9.

29. Mi H, Muruganujan A, Casagrande JT, Thomas PD. Large-scale gene function analysis with
the PANTHER classification system. Nat Protoc. 2013;8:1551–66.

30. Cho RJ, Campbell MJ. Transcription, genomes, function. Trends Genet. 2000;16:409–15.

31. Xia J, Gill EE, Hancock REW. NetworkAnalyst for statistical, visual and network-based
meta-analysis of gene expression data. Nat Protoc. 2015;10:823–44.

32. Delaneau O, Ongen H, Brown AA, Fort A, Panousis N, Dermitzakis E. A complete tool set
for molecular QTL discovery and analysis. bioRxiv [Internet]. 2016; Available from:
http://biorxiv.org/content/early/2016/08/10/068635.abstract

33. Kukurba KR, Parsana P, Balliu B, Smith KS, Zappala Z, Knowles DA, et al. Impact of the X
Chromosome and sex on regulatory variation. Genome Res. 2016;26:768–77.

34. Gaykalova DA, Zizkova V, Guo T, Tiscareno I, Wei Y, Vatapalli R, et al. Integrative
computational analysis of transcriptional and epigenetic alterations implicates DTX1 as a
putative tumor suppressor gene in HNSCC. Oncotarget. 2017;8:15349–63.

35. Liu Y, Zheng P. CD24: a genetic checkpoint in T cell homeostasis and autoimmune diseases.
Trends Immunol. 2007;28:315–20.

36. Gu J, Liu Y, Xie B, Ye P, Huang J, Lu Z. Roles of toll-like receptors: From inflammation to
lung cancer progression. Biomed Rep. 2018;8:126–32.

37. Li L, Tang P, Li S, Qin X, Yang H, Wu C, et al. Notch signaling pathway networks in cancer
metastasis: a new target for cancer therapy. Med Oncol. 2017;34:180.

38. Braunstein MJ, Kucharczyk J, Adams S. Targeting Toll-Like Receptors for Cancer Therapy.
Target Oncol. 2018;13:583–98.

39. Pinho SS, Reis CA. Glycosylation in cancer: mechanisms and clinical implications. Nat Rev
Cancer. 2015;15:540–55.

40. Stowell SR, Ju T, Cummings RD. Protein glycosylation in cancer. Annu Rev Pathol.
2015;10:473–510.

41. Mehta A, Herrera H, Block T. Chapter Seven - Glycosylation and Liver Cancer. In: Drake
RR, Ball LE, editors. Advances in Cancer Research. Academic Press; 2015. page 257–79.

42. Liang K-H, Yeh C-T. O-glycosylation in liver cancer: Clinical associations and potential
mechanisms. Liver Research. 2017;1:193–6.

43. McDaniell R, Warthen DM, Sanchez-Lara PA, Pai A, Krantz ID, Piccoli DA, et al. NOTCH2
mutations cause Alagille syndrome, a heterogeneous disorder of the notch signaling pathway.
Am J Hum Genet. 2006;79:169–73.

44. Oda T, Elkahloun AG, Pike BL, Okajima K, Krantz ID, Genin A, et al. Mutations in the
human Jagged1 gene are responsible for Alagille syndrome. Nat Genet. 1997;16:235–42.

45. Deng H, Jasper H. Sexual Dimorphism: How Female Cells Win the Race. Curr Biol.
2016;26:R212–5.

46. Ntziachristos P, Lim JS, Sage J, Aifantis I. From fly wings to targeted cancer therapies: a
centennial for notch signaling. Cancer Cell. 2014;25:318–34.

Supporting Information

Supporting Information contains Supplementary Methods, six figures, and 18 tables.

Fig S1. A multi-dimensional scaling plot of the TCGA LIHC tumor samples of each sex.
Euclidean distances between samples were calculated based on 100 genes with the largest
standard deviations between samples.

Fig S2. Absolute $\log_2$-fold changes of DEGs detected from tumor vs. tumor-adjacent
comparisons in the sex-specific analyses and combined analysis of both sexes. Absolute $\log_2$-fold
changes are given for female samples, male samples, and across all samples. Global p-values for
ANOVA are shown for each DEG type. Adjusted p-values based on Kruskal-Wallis tests are
shown for each pairwise comparison.

Fig S3. Cellular content of TCGA LIHC tumor samples based on cell-type enrichment analysis from bulk-RNAseq data. Heatmap is based on scaled and mean centered estimates of cell type proportions.

Fig S4. PEER factors associated with the continuous and categorical sample and patient covariates. **A-B:** R^2 and FDR-adjusted p-values between PEER factors and continuous technical and biological covariates. **C-D:** ANOVA test statistic and FDR-adjusted p-values between PEER factors and categorical technical and biological covariates.

Fig S5. Estimation of statistical power in the combined (grey), male-specific (blue), and female-specific (red) eQTL analyses with a p-value level 0.01 and 384 variants. Increased power in the combined analysis allows the detection of sex-shared low-effect eQTLs.

Fig S6. Absolute effect sizes of sex-shared and sex-specific eQTLs in males, females, and the whole study sample. Due to the larger sample size, sex-shared low-effect eQTLs are only detected as significant in the combined analysis. "Sex-shared" facet shows eQTLs that were identified as significant in both sex-specific analyses and in the combined analysis of both sexes. Sex-specific eQTLs exhibit a larger effect in one sex than the other, and the effect is diluted in the combined analysis. Sex-shared large effect eQTLs can be detected in sex-specific and combined analyses. Global p-values for ANOVA are shown for each DEG type. Adjusted p-values based on Kruskal-Wallis tests are shown for each pairwise comparison.

Table S1. Clinical characteristics of male and female patients in the TCGA LIHC cohort. Bonferroni adjusted p-value threshold for Fisher's exact test = 0.0006.

Table S2. Sex-biased gene expression in the GTEx liver tissue samples with an FDR-adjusted p-value < 0.1 and absolute $\log_2$ fold-change > 0. N males = 91, N females = 45.

Table S3. Sex-biased gene expression in the TCGA LIHC tumor-adjacent samples with an FDR-adjusted p-value < 0.1 and an absolute $\log_2$ fold-change > 0. N males = 28, N females = 22.

Table S4. Sex-biased gene expression in TCGA LIHC tumor samples with an FDR-adjusted p-value < 0.01 and an absolute $\log_2$ fold-change > 2. N males = 28, N females = 22. Empirical p-value based on sex-label shuffling with 1000 permutations.

Table S5. Differentially expressed genes between matched tumor and tumor-adjacent samples in the combined analysis of male and female samples with an FDR-adjusted p-value < 0.01 and an absolute $\log_2$ fold-change > 2. Each tissue N=50, N female = 22, N male = 28.

Table S6. Differentially expressed genes between matched male tumor and tumor-adjacent samples with an FDR-adjusted p-value < 0.01 and an absolute $\log_2$ fold-change > 2. N of sample pairs = 28.

Table S7. Differentially expressed genes between matched female tumor and tumor-adjacent samples with an FDR-adjusted p-value < 0.01 and an absolute $\log_2$ fold-change > 2. N of sample pairs = 22.

Table S8. Overrepresented GO terms and canonical pathways in the sex-shared tumor vs. tumor-adjacent DEGs. Significant terms and pathways were selected based on an FDR-adjusted p-value threshold 0.01.

Table S9. Overrepresented GO terms and canonical pathways in the male-specific tumor vs. tumor-adjacent DEGs. Significant terms and pathways were selected based on an FDR-adjusted p-value threshold 0.01.

Table S10. Overrepresented GO terms and canonical pathways in the female-specific tumor vs. tumor-adjacent DEGs. Significant terms and pathways were selected based on an FDR-adjusted p-value threshold 0.01.

Table S11. *cis*-eQTLs detected in the combined analysis of both sexes. N=367 (N males = 248, N females = 119). Significant eQTLs were selected based on an FDR-adjusted p-value threshold 0.01.

Table S12. *cis*-eQTLs detected in the male-specific analysis. N=248. Significant eQTLs were selected based on an FDR-adjusted p-value threshold 0.01.

Table S13. *cis*-eQTLs detected in the female-specific analysis. N=119. Significant eQTLs were selected based on an FDR-adjusted p-value threshold 0.01.

Table S14. Genomic annotations of eQTLs detected in the combined analysis of both sexes.

Table S15. Genomic annotations of eQTLs detected in the male-specific analysis.

Table S16. Genomic annotations of eQTLs detected in the female-specific analysis.

Table S17. Overrepresented canonical pathways in the male-specific eQTL target genes.

Significant terms and pathways were selected based on an FDR-adjusted *p*-value threshold 0.01.

Table S18. Top-variants associated with autosomal genes that were expressed in a sex-biased

way in HCC tumor samples.

Figure Captions

Fig. 1. Patterns of gene expression and molecular etiologies of male and female HCC. A:
Sex-biased gene expression in HCC. A volcano plot of DEGs between male (N=28) and female
(N=22) HCC tumor samples. X-linked genes are indicated in orange, Y-linked in teal, and

autosomal in black. Significant genes were selected based on an FDR-adjusted *p*-value threshold

of 0.01 and absolute $\log_2$(FC) threshold of 2. Multiple transcripts of the long non-coding RNA

XIST are independently expressed. Genes that were not expressed in a sex-biased way in healthy

liver (GTEx) or in the tumor-adjacent tissues are indicated in italics and underlined. **B:** An

example of a gene exhibiting a sex-bias in HCC but not in healthy liver or tumor-adjacent

tissues. *DTX1* expression in log(CPM) is shown for male and female samples in each tissue. **C:**

A multi-dimensional scaling plot of the paired TCGA LIHC tumor and tumor-adjacent samples

of each sex. Euclidean distances between samples were calculated based on 100 genes with the

largest standard deviations between samples. Tissue type (dimension 1) and sex (dimension 2)

drive the overall patterns of gene expression in HCC. **D:** Venn-diagram of the overlap of DEGs

in the sex-specific and combined analysis of matched tumor and tumor-adjacent samples.

Substantially more DEGs were identified in the sex-specific analyses. **E:** Sex-specific and sex-

shared DEGs were analyzed for the overrepresentation of pathways and canonical networks.

Sex-specific patterns of pathway enrichment point to differential processes driving the etiology

of male and female HCC. **F:** Examples of sex-specific and sex-shared pathways.

Fig. 2. Sex-specific genetic effects on tumor gene expression in HCC. A: QQ-plot of eQTL associations in the combined analysis of both sexes (grey), male-specific analysis (blue), and female-specific analysis (red). **B:** Genomic annotations of eQTLs in the combined analysis of both sexes, male-specific analysis, and female-specific analysis. **C:** Overlap of eGenes detected in combined and sex-specific analyses. **D:** An example of a male-specific eQTL. *POGLUT1* expression in tumors is modulated by a germline variant in *cis* in male HCC, but not in female HCC nor in the combined analysis of both sexes, indicating effect modification by sex.