Title: Revisiting the dimensionality of biological diversity

Abstract

Biodiversity can be represented by a variety of different dimensions (e.g. functional

diversity, phylogenetic diversity, genetic diversity and taxonomic diversity). While

the many different representations of biodiversity can lead to a more complete

description of communities, they also lead to a degree of 'fragmentation' of the

biodiversity concept. Developing a unified measure that integrates all the dimensions

of biodiversity is a theoretical solution for this problem, however, it remains an

operationally utopian task. Alternatively, quantifying the dimensionality of

biodiversity is a way to integrate the different dimensions of biological diversity. This

concept is explored in the present study. We define the dimensionality of diversity as

the number of different diversity metrics needed to effectively describe biodiversity.

We provide a solution that joins the two components of dimensionality (correlation

and the variation of each diversity metric) in the same framework, allowing us to

quantify dimensionality. We show, through a simulation study, that considering the

correlation among dimensions with the variation of each dimension offers important

information that is neglected when only correlation information (the component used

in the current methodology to measure dimensionality) is considered. We demonstrate

how to apply the proposed framework by investigating the dimensionality of South

American small mammal communities (cricetids and marsupials). Our example

demonstrates that the taxonomic dimension of diversity captures most of the spatial

variation among these communities. We conclude by highlighting the greater

understanding of dimensionality that is obtained through this method, as well as the

insights it provides into the importance of each diversity metric to the whole

biodiversity space, insights which can not be obtained through the current common

method.

Keywords: biodiversity metrics, communities, biodiversity measurement, correlation,

variation.

Introduction

Introduction

Biodiversity consists of the fundamental source of information that underlies all

ecological studies, which justifies the efforts of ecological research to develop

measures that properly operationalize its concept (Wilson 1989). These efforts are

reflected in the immensurable number of diversity metrics that have appeared in an

attempt to encompass all the variation in biodiversity. As the number of diversity

metrics grows and promotes the description of the different dimensions of variation,

the same proliferation promotes confusion around the definition of biodiversity,

which was expressed well by Hurlbert (1971) in his seminal work on the "non-

concept of diversity".

A possible way to resolve the confusion created from the many operational

solutions (diversity metrics) of such a unique concept (biodiversity), exists in the

search for a unifying diversity metric that encompass all the dimensions of variation

that define biodiversity. However, consolidating the many dimensions represented by

diversity metrics in a reliable way appears to be a utopian task. This consolidation can

be achieved by investigating the relationships among the existing metrics, which

essentially means assessing the dimensionality of biological diversity.

Dimensionality is an inherent characteristic of biological diversity that can be

defined as the minimum set of diversity metrics needed to sufficiently describe the

whole variation that is present at a given level of biological organization (e.g.

populations and communities) (Lyashevska and Farnsworth 2010; Stevens and

Gavilanez 2015). Which metrics of diversity must be used to sufficiently express the

variation in a given community? What is the minimum set of metrics needed to

characterize a community (Lyashevska and Farnsworth, 2010)? Do functional,

phylogenetic and taxonomic diversity consist of independent dimensions of variation

when defining the diversity of communities? These questions that can be answered by

quantifying the dimensionality of biological diversity. The search for answers to these

questions leads to the development of statistical tools that are able to capture the level

of redundancy (or correlation) in a set of diversity metrics calculated for communities.

In this multidimensional context, ordination techniques offer the basic machinery to

solve problems that involve multiple dimensions however, when considered on their

own, they do not provide a global measure that indicates the dimensionality of a data

set.

A sequence of recent publications by Stevens and Tello (2014), Stevens and

Gavilanez (2015) and Stevens and Tello (2018) follows the investigation of the

dimensionality of bat communities using an index called the Evenness of Eigenvalues

(hereafter EE). This metric is obtained by applying a Principal Ordination Analysis

(PCA) over a matrix of diversity metrics calculated for a set of communities (the

calculation details are explained later in the methods section). The logic behind EE is

that, if the diversity metrics used to characterize a given set of communities are very

redundant, this indicates low dimensionality and produces low EE values. On the

other hand if diversity metrics are completely orthogonal with each other, this

indicates a high dimensionality and the EE value of the community will be high (close

to 1).

The EE metric certainly accomplishes, in a simple and effective way, the

operationalization of the correlation component presented in dimensionality. However

it ignores the amount of variation, or importance, of each metric present in the

multidimensional biodiversity space. Let´s suppose a situation in which diversity

metrics are correlated, but each one accounts for similar amounts of variation in the

total multidimensional biodiversity space. In this situation, despite the high

correlation among dimensions and the low dimensionality, according to the EE

calculation, we will need to rely on a great number of diversity metrics in order to

effectively represent the variation in biodiversity present in this set of communities.

Wilsey et al (2005) was the first to quantify the amount of information that each

metric accounted for in a multidimensional space, using a metric called Importance

Values (hereafter IV, with detailed calculations presented in the methods section).

Through this metric the authors accessed the variation component of dimensionality

in communities.

The common way to quantify dimensionality within the ecological literature

(Stevens and Tello, 2015) provides an incomplete picture of the inherent

characteristics of biodiversity, since it only relies on correlation and does not consider

the variation component of the dimensionality concept. In Figure 1 we illustrate the

problem associated with only considering the correlation component of

dimensionality. Circles and triangles represent two metacommunities for which two

diversity metrics (which could be imagined as functional diversity and phylogenetic

diversity) were calculated and plotted against each other (panel A). In the first

scenario diversity measures are highly correlated (HiC in Figure 1), while in the other

scenario the same biodiversity measures are minimally correlated with each other

(LoC in Figure 1). Thus, communities in scenarios HiC and LoC present low and high

dimensionality, respectively, when calculated according to the EE (panel B).

However, in scenario HiC we note that, for triangles, diversity metric 2 has a much

lower variance than diversity metric 1 (high correlation but a different amount of

variation among metrics – HiC/DifV). In contrast, both diversity metric 1 and

diversity metric 2 present similar variation for the circles (high correlation and similar

variation among metrics – HiC/EqV). Despite these different patterns, the EE cannot

distinguish the differential contribution of the two diversity metrics, since it only

relies on the correlation component of dimensionality. A similar situation is expressed

by the LoC scenario, in which metric 1 has a lower variation than metric 2 for triangle

communities (low correlation and differential variation [Lo/DifV]), and variation does

not differ between the two metrics for the circle communities (low correlation and

equal variation [Lo/EqV]).

To overcome the problem illustrated in Figure 1, which is that the EE do not

capture all the components that make up the concept of dimensionality, a framework

that combines a measure that captures the correlation among metrics with the

variation of each diversity metric must be used. Operationally, this implies combining

the EE and IV measures into one framework which adequately characterizes the

dimensionality of biodiversity. Despite the existence of these two metrics that are

capable of measuring the two components of the dimensionality concept, they have so

far only been used separately, providing a limited perspective of the dimensionality of

biodiversity.

In this paper we propose an integrative framework for the dimensionality of

biodiversity and provide the most well-rounded characterization of dimensionality to

date. Within this framework we integrate the EE and IV metrics in order to conciliate

the correlation and variation components of the dimensionality of diversity. We start

by showing, through a simulation study, how using the EE and IV together can

overcome the problem presented in Figure 1. Following this we present an empirical

example of the investigation of dimensionality by applying the proposed framework

of dimensionality to communities of small mammals (crycetids and marsupials).

Methods

Investigating the dimensionality of biodiversity: obtaining EE and IV

Our framework to investigate the dimensionality of biodiversity is comprised

of three steps. The first step is to calculate the matrix of diversity metrics, hereafter

called matrix **M** (*sensu* Ricotta 2005). Here matrix **M** will contain three metrics of

diversity: a measure of functional diversity (FD [Petchey & Gaston, 2006)]),

phylogenetic diversity (PD [Faith, 1992]) and richness. Despite our choices of these

three metrics we highlight that the matrix **M** can be assembled with any other

diversity metrics. We chose these three metrics since they are some of the most

commonly used diversity metrics, representing the most frequently applied

dimensions of biological diversity in the ecological literature (Magurran 2011). The

second step involves performing a Principal Component Analysis (PCA) of matrix **M**.

As will be shown next, the standardization method applied to matrix **M** prior to the

PCA must differ between the calculation of EE and IV. The third step is the

calculation of the dimensionality metrics EE and IV. We calculate EE using the

Camargo´s evenness index:

$$EE = 1 - \left(\sum_{P}^{A(A-1)} |e_{ip} - e_{jp}|\right)\Big/_{A} \quad \text{Equation 1}$$

Camargo's evenness index (Camargo 1983) was calculated Using the axes (A) and

their respective eigenvalues (e_{ih} and e_{jh}), which were produced from a PCA (p)

performed on a standardized matrix **M** in which metrics were scaled to have zero

mean and equal variances. This is the procedure used by Stevens and Tello (2015).

EE values range from 0 (low dimensionality) to 1 (high dimensionality). The greater

the value of EE, the less correlated communities are in relation to the diversity metrics

that characterize these communities. On the contrary, low EE values indicate that the

diversity metrics that characterize the communities are highly correlated. IV was

calculated by modifying the method proposed by Wilsey et al., (2005), which uses a

matrix (**M**) that has been standardized by the maximum values of each diversity

metric. This standardization removes the effect that the different units of each

diversity metric can have, without modifying their original variation. To obtain the IV

of each diversity metrics in matrix **M** we apply Equation 2, in which IV_i represents

the IV of the diversity metric, i, r^2_{ij} is the correlation of diversity metric i with PCj,

and R^2_j is the amount of variation that PC$_j$ accounts for in the ordination space.

161            $$IV_i = \sum r_{ij}^2 \times R_j^2 \qquad \text{Equation 2}$$

162   PC varies from 1 to j and corresponds to the number of significant

eigenvectors in the PCA, evaluated by Kaiser-Gutmann criterion. The greater the IV

value the more variation the diversity metric accounts for in the multidimensional

biodiversity space. IV values approach 1 when the diversity metric accounts for

almost all variation and approach zero when the metric presents low values of

variation.

Testing the assessment of the dimensionality of diversity using EE and IV

169   We calculated both EE and IV values for metacommunities simulated

according to the patterns of diversity metrics illustrated in the scenarios presented in

Figure 1 (HiC/EqV, HiC/DifV, LoC/EqV and LoC/DifV). They were calculated by

using the matrix **M**, which was composed of three metrics: richness, PD and FD. An

illustration of the general framework used to investigate the dimensionality of

diversity is presented in Figure 2. Here, the matrix **M** is shown to have three

dimensions, however it can be assembled by more than three dimensions and any

other dimension of diversity can be used inside the framework presented in this work.

177   We were interested in determining whether EE values could correctly detect

the levels of information redundancy present in matrix **M** (high or low), and if IV

values could identify differences in the contribution of each diversity metric to the

multidimensional biodiversity space. To prove that EE and IV values can do these

things, the following conditions must be met. EE values must not differ in scenarios

with equal and differential variation (when comparing between scenarios HiC/EqV

and HiC/DifV or LoC/EqV and LoC/DifV), but they can differ between highly

correlated and poorly correlated scenarios (when comparing between scenarios

HiC/EqV and LoC/EqV or HiC/DifV and LoC/DifV). The differences between the

values of IV must be lower in communities in which all diversity metrics possess

similar variation (Scenario HiC/EqV and LoC/EqV, represented by the circle

communities in Figure 1) and they must be greater in scenarios in which only two

diversity metric accounts for most of the variation in matrix **M** (Scenarios HiC/DifV

and LoC/DifV, represented by the triangle communities in Figure 1).

Simulation methodology

Figure 1 represents the theoretical scenarios in which the performance of the

EE and IV values were tested. We simulated metacommunities based on a pattern-

oriented procedure, producing diversity metrics with patterns of correlation and

variation that resembled the scenarios presented in Figure 1. In the HiC/EqV scenario

the diversity metrics are highly correlated but have a similar variation in the

dimensional space (illustrated in Fig.1A, HiC with circles). The HiC/DifV scenario

presents diversity metrics that are highly correlated but vary in relation to the

variation that each one account in the multidimensional space, which is mainly

determined by two diversity metric. In these two scenarios we expect to obtain low

values of EE (since the diversity metrics are very correlated), but we expect similar

values of IV in the HiC/EqV scenario and different values of IV in the HiC/DiFV

scenario. In the HiC/DiFV scenario we expect two diversity metrics to have higher

values than the other one of the diversity metrics.

The biological reasoning of the HiC/EqV and HiC/DifV scenarios followed a

pattern that can be predicted by a species sorting model (Leibold et al. 2004). Within

this model, species possess traits that allow them to persist in a community according

to the availability of resources (represented along a gradient here). The simulation

starts with a phylogeny that is simulated by a birth-death processes (function

sim.bdtree from the package geiger [Harmon, Weir, Brock, Glor, & Challenger,

2008]) where a species, chosen randomly, colonizes the community and starts the

assembling procedure. The addition of a species to the community depends on the

species that are already present in that community. At one extreme of the gradient the

communities will only contain species that are phylogenetically closely related to

each other (the 10% of species that are the most phylogenetically related to each

other), with the phylogenetic filter becoming less restrictive until at the other extreme

the communities contain species that have the same probability of being selected to

coexist (less selective conditions). This creates a phylogenetic diversity gradient in

the metacommunity (a pattern expected by a process of phylogenetic habitat filtering,

sensu Duarte [2011]).

In scenario HiC/EqV we simulated a trait that was continuously conserved

over the phylogenetic tree (a trait that evolved according to a Brownian motion model

of evolution, determined by using the function *rTraitCont* (Paradis, 2012) with the ρ

[*rho*] parameter set to the value of 3), with the number of species in each community

increasing gradually (less phylogenetic filter, more species). This created a gradient in

phylogenetic, functional and taxonomic diversity in a metacommunity that contained

high levels of correlation and similar values of variation in diversity metrics. In

scenario HiC/ DifV we simulated a trait that evolves according to a stabilizing

selection regime (Ornstein-Uhlebeck model with the strength of selection set by

giving the α parameter a value of 0.8) that restricts trait variation to within an optimal

range (represented by a θ [theta] value of 0). This allowed us to generate a

metacommunity in which the diversity metrics are highly correlated, but the variation

of FD is much lower than that of the richness and PD metrics since the traits that were

used in the calculation of FD were restricted by the selection process.

In scenario LoC/DifV we start with a phylogeny and then generate traits

following the same procedures described above for HiC/EqV scenario. However, the

trait generated here to had a low phylogenetic signal and the phylogenetic tree used to

calculate PD for each community was modified to simulate a process of evolution in

which most of the speciation occurs near the root (a star-like phylogeny). This

procedure resulted in poorly correlated diversity metrics since the relationship

between phylogeny and traits was disrupted. Additionally, a low variability for PD

and richness metrics was obtained since we set the simulations to produce

communities with the same number of species. Consequently most of the variation in

this scenario is due to the FD metric. Finally, scenario LoC/EqV was simulated in a

manner by which all species present in the phylogenetic tree had an equal probability

of being present in any community within the metacommunity (no phylogenetic

habitat filtering acting on the assembly of the metacommunity). This procedure

generated metacommunities with low correlation and similar amounts of variation for

the diversity metrics.

We generated 999 metacommunities for each scenario described above

(HiC/EqV; HiC/DifV; LoC/EqV and; LoC/DifV). In all scenarios the

metacommunities were composed of 50 communities, with each community

possessing a minimum of 20 and a maximum of 200 species. The phylogenetic filter

was set to act gradually through the communities, increasing by the order of 10% (*i.e*,

start by selecting the top 10% of species that are the most phylogenetically related,

followed by the top 20% and so on until 90% of the species have been selected from

the pool). The details and an illustration of the simulation procedures and scenarios

are presented in the Supplementary Material S1. Here we also describe the interactive

module we produced to illustrate the simulation procedure used in this work.

Finally, we tested whether the EE and IV values met our expectations in the

simulated communities. To analyze the EE values we checked if they were different

in scenarios with low correlation and high correlation (scenarios HiC/DifV and

HiC/EqV differing from LoC/EqV and LoC/DifV). To effectively capture

dimensionality the EE metric must be higher in scenarios with low correlation than in

scenarios that presented high correlation among diversity metrics. To test for

differences among the IV values of each metric in the scenarios presented above we

compared the IV values using a graphical tool called profile of importance (*sensu*

Wilsey et al. 2005) and quantified the differences by calculating F values from a

linear model (Equation 3). These values compare the IV of the three dimensions (PD,

FD and richness) and test if the IV in the scenarios with differential variation in

diversity metrics (scenarios HiC/DifV and LoC/DifV) presented greater differences

than IV calculated for the scenarios that present similar variation in diversity metrics

(scenarios HiC/EqV and LoC/EqV).

Assessing the dimensionality of biodiversity in small mammal communities

To illustrate the application of the dimensionality framework we used a

database of small mammal communities (marsupial and cricetid mammals) distributed

throughout the South American continent. We considered the same three diversity

metrics used to assemble the matrix **M** in the simulation analysis (richness, PD and

FD,)**.** The EE and IV metrics were calculated as previously described in the section

'Investigating the dimensionality of biodiversity: obtaining EE and IV'*,* although we

only used the first axis of variation provided by the PCA of matrix **M** to calculate the

IV metric, since this was the only axis that was significant according to the Kaiser-

Gutmann stop criterion.

We also compared the observed values of EE with a null distribution

composed of 999 EE values, generated by two null models. The first null model

randomizes a species incidence matrix while preserving the differences in richness

among sites but mixing species frequency (performed with the *sim3* function from the

EcoSimR package [Gotelli, 2000]). The second null model randomizes an incidence

matrix while preserving both differences in species richness among sites and species

frequency (*sim9* function in *EcoSimR* package). We tested the null hypothesis that the

observed EE values do not differ from EE expected by variation in richness values.

We implemented a function called *dimensionality* to calculate the EE values from the

matrix **M**. The function allows the user to choose the evenness method that will be

used in the calculation. It can be accessed at

https://github.com/GabrielNakamura/dimensionality_function.

To calculate IV for the metacommunities of small mammals we applied the R

code for the IV function, *ImportanceVal*, (the function can be accessed at

https://github.com/GabrielNakamura/IV_function/blob/master/ImportanceVal.R). We

used the Kaiser-Gutmann stop criterion and a bootstrap procedure to generate

confidence intervals for the IV value of each diversity metric. We performed all the

calculations with a standardized matrix **M** (scaled to zero mean and unit variance for

the calculation of EE values and standardized by the maximum values of each metric

for the calculation of IV values. Explanations were provided in the previous section).

The bootstrapped IV values were submitted to an Ordinary Least Square (OLS) linear

model:

$IV_i = \alpha + \beta * dimension_i + \varepsilon_i$,   Equation 3

where IV_i represents the predicted value of IV for the diversity metric i (the sum of

the intercept α and the effect of the dimension i (β_i)), and ε_i represents the error term

associated with the residuals, which follow a gaussian distribution. This tests the null

hypothesis that IV values do not differ among the dimensions of diversity.

Results

Simulated data

Our simulation shows that EE and IV behave as we predicted, according the scenarios

tested. EE correctly captured the correlation component of dimensionality, evidenced

by the higher values presented in the LoC/DifV and LoC/EqV than in the HiC/DifV

and HiC/EqV scenarios. The IV metric also behaved as we expected; showing greater

differences among IV values in scenarios where the variation in the multidimensional

biodiversity space was only affected by two or one metric (PD and richness in

HiC/DifV and FD in LoC/DifV). This result was evidenced in Figure 3 C and D, with

much greater F-values in the HiC/DifV and LoC/DifV scenario (indicating greater

differences in contribution) than in the HiC/EqV and LoC/EqV scenario.

Small mammal communities

Calculation of the EE value for the small mammal communities produced an observed

value of 0.46. Figure 4 reveals that the correlation component of dimensionality, at

least for the three dimensions analyzed (functional, phylogenetic and taxonomic),

may be a consequence of the spatial gradients of species richness.

IV values for the three diversity metrics were illustrated in Figure 5 through the

Importance Values profile (*sensu* Willig and Hollander, 1995). Analysis of the IV

values showed that richness represents most of the variation in the biodiversity space

(composed of the three diversity metrics), with an importance value of 0.32. The

phylogenetic dimension (PD) represents the second largest quantity of variation, with

a value of 0.29, and the functional dimension (FD) represents the smallest quantity

with a value of 0.22. These values represent, respectively, 38%, 34% and 26% of the

total variation along the significant axis selected for the IV analysis of matrix **M** (only

the first axis was selected according to the Kaiser-Guttman criterion). Furthermore,

the ANOVA model showed that IV differed significantly (F-value= 3.428; p<0.05),

reinforcing the greater quantity of variation that is represented by richness in the

dimensionality of the communities of cricetids and marsupials.

Discussion

Our simulation results demonstrate that it is necessary to use EE and IV values in a

complementary manner, since a similar value of EE can be found for communities

presenting diversity metrics with different levels of importance. Given this finding,

our framework represents a step beyond the method proposed by Stevens and Tello

(2014) to quantify the dimensionality of diversity. Additionally, our work promotes

an integration of different tools already used to quantify separate aspects of

dimensionality in a single and comprehensive framework. Finally, we provide the first

results (to our knowledge) showing the effectiveness of using EE and IV values to

capture patterns of the dimensionality of biodiversity.

Despite only using three dimensions in the present work, the methods used

here are highly flexible in the sense that we can use a matrix (**M**) that contains many

more dimensions (Ricotta 2005). Therefore, we could represent diversity in a much

more complete manner, with metrics that capture other quantifiable dimensions such

as the genomic dimension (*e.g.* Nei 1978) and many others (*e.g.* Gotelli et al. 2012).

Including other dimensions would allow the contributions of these other metrics to

biodiversity to be assessed, and the complementary or redundant nature of these

metrics, in relation to the phylogenetic, functional and taxonomic dimensions, could

be discovered.

Our application of the dimensionality framework to small mammal

communities showed that cricetids and marsupials possess low dimensionality and

that the observed value of EE can be explained by the variation in the taxonomic

dimension (represented by richness). These findings differ from those of Stevens and

Gavilanez (2015), who found the EE of bat communities to be significantly higher

than expected by null models (the same null models used in the present work),

although their observed EE value is lower than the observed value of the present

study. The difference between the present study and the study of Stevens and

Gavilanez (2015) is evidence that the correlation component of the dimensionality of

diversity can change depending on the taxonomic group being investigated.

IV values for small mammal communities contrasted the findings of Wilsey et

al (2005) and Lyashevska and Farnsworth (2002), who concluded that richness was

the least important diversity metric representing the variation in the structure of

communities (grassland and marine benthic communities). Therefore we have

evidence that the variation component of dimensionality, in addition to the correlation

component, can present specific patterns depending on the taxonomic group being

investigated. This highlights the need to understand the contingencies in the

correlation and variation components of the dimensionality of different communities.

Conclusion and future directions

This work provides an update of the definition of the concept of

dimensionality presented by previous works. We demonstrate how to include the

variation components of dimensionality, through the use of EE and IV values, in the

same framework. Since the evidence presented in the literature is limited, and based

only on specific groups of organisms, some questions still need to be addressed to

provide a more complete understanding and generalization of the role that some

factors play in the dimensionality of ecological communities. For instance, we might

wonder if some dimensions of diversity are consistently more informative than others

when describing diversity patterns among different taxa, or if distinct factors

(historical, evolutionary and/or ecological) generate predictable higher or lower levels

of diversity dimensionality across communities.

References

Duarte, L. D. S. 2011. Phylogenetic habitat filtering influences forest nucleation in
grasslands. - Oikos 120: 208–215.

Faith, D. P. 1992. Conservation evaluation and phylogenetic diversity. - Biol.
Conserv. 61: 1–10.

Gaston, K. J. 1996. What is biodiversity?. - In: Gaston K.J. (ed.), Biodiversity: a

biology of numbers and difference. Blackwell Science, pp. 1–9.

Gotelli, N. J. et al. 2013. Environmental proteomics, biodiversity statistics, and food-

web structure. – Trends Ecol. Evol. 27: 436-442.

Lyashevska, O. and Farnsworth, K.D. 2012. How many dimensions of biodiversity do

we need?. - Ecol. Indic. 18: 485–492.

Magurran, A. and McGill, B.J. 2010. Biological Diversity: frontiers in measurement

and assessment. - Oxford Univ. Press.

Magurran, A.E. 1988. Ecological Diversity and its Measurement. - Princeton Univ.

Press.

Naeem, S. 2002. Ecosystem consequences of biodiversity loss: the evolution of a

paradigm. – Ecology 83: 1537-1552.

Nanney, D. 2004. No Trivial Pursuit. - Bioscience 54: 720–721.

Petchey, O.L. and Gaston, K.J. 2002. Functional diversity (FD), species richness and

community composition. – Ecol. Lett. 5: 402-411.

Purvis, A. and Hector, A. 2000. Getting the measure of biodiversity. - Nature 405:

212–219.

Ricotta, C. 2005. Through the jungle of biological diversity. - Acta Biotheor. 53: 29-

38.

Stevens, R.D. and Gavilanez, M.M. 2015. Dimensionality of community structure:

phylogenetic, morphological and functional perspectives along biodiversity and

environmental gradients. – Ecography. 38: 861–875.

Stevens, R.D. and Tello, J.S. 2014. On the measurement of dimensionality of

biodiversity. - Glob. Ecol. Biogeogr. 23: 1115-1125.

Tucker, C.M. and Cadotte, M.W. 2013. Unifying measures of biodiversity:

understanding when richness and phylogenetic diversity should be congruent. -

Divers. Distrib. 19:845–854.

Whittaker, R.H. 1972. Evolution and Measurement of Species Diversity. - Taxon 21:

213–251.

Willig, M.R. and Hollander, R.R. 1995. Secondary sexual dimorphism and

phylogenetic constraints in bats: a multivariate approach. - J. Mammal. 76: 981–992.

Wilsey, B.J. et al. 2005. Relationships among indices suggest that richness is an

incomplete surrogate for grassland biodiversity. - Ecology 86: 1178–1184.

Figure captions

Figure 1: A) Relationships between two diversity metrics from two metacommunities

– circles represent the Equal Variation (EqV) and triangles represent the Different

Variation (DifV) in diversity metrics. These metacommunities can have a high

correlation among metrics (HiC) or a low correlation among metrics (LoC). B)

Evenness of eigenvalues (EE). C) Importance Values (IV).

Figure 2: General framework used to investigate the dimensionality of diversity

through the calculation of the metrics EE and IV. Starting with a set of communities,

metrics of diversity are calculated to assemble matrix **M** (step 1). A PCA of matrix **M**

is then performed (step 2) and the dimensionality measures of diversity are calculated

(step 3). Details about the calculation of IV and EE are provided in the text of the

Methods section 'Investigating the dimensionality of biodiversity: obtaining EE and

IV'.

Figure 3: Profile of importance of the IV and EE metrics (bar plots A and B),

calculated for simulated metacommunities according to the simulated scenarios. For

each of these scenarios we presented situations in which the metrics either contributed

similarly or contributed unequally to dimensionality (respectively HiC/EqV,

LoC/EqV and HiC/DifV, LoC/DifV). Boxplots C and D represent the F-value

distributions from linear models for scenarios HiC/DifV, HiC/EqV, LoC/DifV and

LoC/EqV, evidencing the difference in contribution of one or two diversity metrics in

scenarios HiC/DifV and LoC/DifV.

Figure 4: Histogram of null EEs. The dotted lines represent the 25% and 95% values

of the null distribution. The solid lines represent the observed EE, indicating that the

observed EE does not differ from the EE values presented by the null models.

Figure 5: IV profile of marsupial and cricetid mammal communities from South

America. Bar height corresponds to the observed IV for each diversity metric while

the lines represent the 95% confidence intervals, generated by a bootstrap analysis.

Dimensions of biodiversity

Taxonomic   Phylogenetic   Functional

metric 1   metric 2   metric 3

Comm 1
Comm 2
Comm 3
.
Comm n

Matrix **M**

Communities

Step 1

PCA of
matrix **M**

PC1

PC2

calculation of
dimensionality
metrics

Step 2

Evenness of
Eigenvalues (EE)

Importance
Values (IVs)

Step 3