31° South: Dietary niche of an arid-zone endemic passerine

Ângela M. Ribeiro[1][§], Ben Smit[2,4], and M. Thomas P. Gilbert[1,3]

[1] Natural History Museum of Denmark, University of Copenhagen, Copenhagen, Denmark

[2] Department of Zoology, Nelson Mandel Metropolitan University, Port Elizabeth, South Africa

[3] Norwegian University of Science and Technology, University Museum, Trondheim, Norway

[4] *Current address:* Department of Zoology and Entomology, Rhodes University, Grahamstown,

South Africa

[§] E-mail: ribeiro.angela@gmail.com

ABSTRACT

Balancing energy budgets is thought to be challenging for birds living in arid ecosystems because food supplies are low and unpredictable, and climatic conditions extreme. Thus, to ensure they obtain sufficient energy to fuel daily energetic budgets, birds may need to adjust their diets and become less selective (generalist) as conditions become harsher. To test this hypothesis, we used DNA metabarcoding to characterize both the prey availability and the dietary content of several conspecific populations of a semi- and arid-endemic insectivorous bird, the Karoo-scrub-robin (*Cercotrichas coryphaeus*) across a climatic gradient. Our results showed that Coleoptera, Hymenoptera, Orthoptera, and Lepidoptera were the main prey. When accounting for their presence as available prey, Coleoptera and Hymenoptera were preferred in all regions, whereas robins avoided Orthoptera and Lepidoptera in all but the most arid region. Although the different populations live in regions that vary with regards to productivity and thermoregulatory demands, we found that the dietary niche breadth (Bs) of the three populations was intermediate to low, and did not differ significantly. As a whole, our findings show that regardless of environmental harshness these insectivores have similar dietary niches, suggesting that large dietary plasticity is fundamental for their survival in energy-depauperated ecosystems.

Keywords: birds, arid ecosystems, dietary niche, arthropods, DNA-barcoding, high-throughput sequencing

INTRODUCTION

Food acquisition is a basic need of all animals: it provides the energy and nutrients required to sustain life in any given environment. Because of the intricate relationship of animals with the environment (where food is obtained from), there has long been an interest in describing patterns of dietary variation in order to predict the underlying causes. The classical Optimal Foraging Theory model (MacArthur and Pianka 1966) (Levins and MacArthur 1969) (Schoener 1971) predicts that animals seek to maximize energy intake, so as to meet their daily energy costs. Consequently, both prey availability in the environment and animal physiological status can affect diet variation (e.g.: Burgar et al. 2014).

Endotherms living in landscapes characterized by dramatic change in rainfall and temperature face two main problems: firstly, frequent changes in food webs; and secondly, thermal challenges that require them to adjust their physiology (energetically expensive tasks) to maintain body temperatures within tolerable limits. As a consequence of both, some populations may undergo dietary niche shifts in order to maximize fitness (Roches et al. 2016). In response to limited resources, selection can either lead to generalist strategies such as dietary niche flexibility to use the panoply of resources available (Grant et al. 2008), or drive the evolution of specialization so as to exploit underused resources (Martin and Pfennig 2009). Moreover, if resources vary greatly through time or space, being a generalist may allow fitness to be maintained (Stephens and Krebs 1986).

Arid-zone endemic birds are good models for testing the role of environmental temperature and energetic physiology in dietary strategies, because they face continual challenges from their energy-depauperated ecosystems, while seeking to fulfill the dietary requirements needed to fuel the physiological processes associated with thermoregulation. In this study we molecularly characterized both the diet and available arthropod prey community (Pompanon et al. 2012) (Bohmann et al. 2011) of conspecific populations of the Karoo-scrub robin (*Cercotrichas*

coryphaeus; hereafter: robin), a ground-feeding insectivore. This southern African passerine

provides a good system with which to explore the avian dietary strategy to ensure adequate

energy intake, because its range spans an environmental gradient of primary productivity

variation (Figure 1A). Primary productivity ultimately determines arthropod biomass in the

regions, which in turn affects the food available for insectivores (Maclean 1969) (Lloyd 1999).

Furthermore, these robins also face dramatic temperature changes, which lead thermoregulatory

challenges that the birds must adjust their rates of energy expenditure to (Figure 1B).

Our objectives were therefore two-fold. Firstly, to determine whether environmental

heterogeneity resulted in differential prey availably. And secondly, to assess whether this prey

availably and physiological status lead to differences in the birds' niche breadths, a proxy for

their dietary flexibility and prey selectivity. In particular, we hypothesized that populations

resident in the area with the lowest productivity and highest metabolic expansibility, should

exhibit the most generalist feeding strategy, as their means for coping with the harshness of the

arid-conditions.

2. MATERIAL AND METHODS

2.1 Study sites, arthropods and fecal sample collection

The study was conducted along a gradient of (i) annual temperature amplitude (T_{amp}) and (ii)

primary productivity (measured as normalized difference vegetation index – NDV) (Figure 1).

We selected these two metrics with the rationale that: wide ambient temperature range implies

increased energy expenditure so as to deal with extreme climatic conditions (cold and hot), and

consequent incremental intake of energy sources (Bicudo et al. 2010); and that NDVI is a good

predictor of plant biomass in the arid-zone of southern Africa (Borer et al. 2012).

We used the GPS coordinates of our pitfall grid to obtain T_{amp} and NDVI. T_{amp} was estimated

using data from WorldClim2 at 2.5min resolution (Fick and Hijmans 2017) as: maximum

temperature in December – minimum temperature in July. NDVI data, the surrogate for primary

productivity, was extracted from eMODIS dataset hosted by NASA-LPDAAC (USGS, USA).

Beside environmental variables, we gathered physiological phenotypic data (Ribeiro et al. *in*

prep) for populations inhabiting the same localities. We used these metrics to define three

operational regions: *Coastal, Central,* and *Inland.*

We sampled available prey arthropods using 27 pitfall traps left open for seven consecutive days,

at each of three locations (Figure 1) in the summer of 2015 and winter of 2016. The pitfall traps

were placed within an area we previously checked to be used by robins, and then set them in

three grids. Each grid comprised of nine traps forming a square (3m x 3m), with a distance of one

meter between individual traps, and a distance of 100 m between grids. The traps were made

from two nested transparent plastic cups of 8 cm diameter, and 250 ml of volume; we partially

filled them with a solution of water and soap to reduce superficial tension (Cheli and Corley

2010). We collected the arthropods in the traps on the fourth and seventh (last) day of sampling.

All arthropods caught per site were pooled to represent the arthropod community of the region in

the given season. Post sampling, arthropods were washed with water and ethanol to remove

debris, and stored in 96% ethanol until further use.

In the same above-mentioned periods we collected 49 bird fecal pellets ($N_{Coastal}$= 16, $N_{Central}$= 16,

N_{Inland}= 17) using single use filter-paper bags lining cloth bags in which captured birds were

placed. Upon collection, the fecal pellets were placed in tubes containing beads and lysis buffer

as provided in the Soil/Fecal DNA MiniPrep kit (ZymoResearch, USA), homogenized by

vortexing at 2700rpm (45HZ) for 1 min, and stored at 4˚C until further processing in the lab. The

sampling periods were selected to avoid the breeding season; in fact, we found no sign of

breeding activity while in the field.

2.2 DNA extraction

Prior to DNA extraction, arthropods were sorted by size into three categories: small (<1cm),

medium (1-3cm) and large (>3cm). The rational was two fold: first, large-sized arthropods could

be easily identified morphologically (supplementary Table S1); and second, we wanted to extract

DNA from several arthropods at a time (mixed sample), hence it is crucial that the animals in the

same pool have similar sizes so not to biases the proportion of DNA in the final volume.

Arthropod samples were drained and let dried at 56°C to remove all ethanol. In order to maintain

the arthropod exoskeletons intact, we extracted DNA from arthropod samples following (Gilbert

et al. 2007) and (Nielsen 2015). Briefly, this consisted of immersing the arthropods in a digestion

buffer with incubation overnight at 56°C with gentle agitation, after which the digest was purified

using the QiaQuick Purification kit (Qiagen, USA). Extracted DNA was stored at -20°C until

further use.

We homogenized fecal samples prior to DNA extraction using a tissue lyser (Qiagen Tissue

Lyser II, USA) for 30s at 30Hz, followed by 30s at 20Hz. DNA extractions used the Soil/Fecal

DNA MiniPrep kit (ZymoResearch, USA), with one minor change to the manufacturer's

guidelines: after homogenization, we added Proteinase K and incubated the samples at 60 °C for

30 min. The DNA extracted was stored at -20°C until further use. We specifically chose this kit

because it contains beads in a bead beating tube, a feature that may help crush both the

exoskeleton fragments of arthropods and seeds in fecal pellets, hence increase the yield of DNA.

Negative controls were included in each extraction to check for potential contamination. These

negative controls proceeded in the workflow (from PCR to sequencing) as with all other extracts.

2.3 Metabarcoding and sequencing

To maximize detection and identification of arthropods we used two different primers targeting

different portions of the COI gene in the mitochondrial genome of arthropods: i) ZBJ-

ArtF1c/ZBJ-ArtR2c (Zeale et al. 2011) and, ii) FormiF/FormiR. The latter set was included

because the ZBJ-Art primers have been reported to poorly detected ants and termites (Hamad et

al. 2014) (Brandon-Mong et al. 2015). Thus we specifically designed this set to ensure detection

of ants (supplementary Table S2)., by first downloading all arthropod COI sequences from South

Africa archived on GenBank in January 2016, then designing them by hand in Geneious 7.0.6

(Biomatters, New Zealand) to meet the following criteria: a) 20-22bp length; b) produce a

fragment of 150-210bp (including primers); and c) C/G as the first two 3' end bases (alignment

provided as Supplementary Material).

In addition, because robins have been seen eating berries (P. Lloyd, pers. com.), we targeted the

chloroplast rbcL gene using primers rbcL-h1aF/rbcL-h2aR (Poinar et al. 1998). The ZBJ-Art

primers amplify a 211bp region in the 5' section of COI gene, while the Formi primers target a

168bp region in the 3' of same gene. All primers were modified to include a unique 6 to 8

nucleotide sequence (nucleotide multiplex identifiers; MIDs) on the 5' end to allow individual

identification following (Binladen et al. 2007). MIDs were designed with a custom script using R

(R version 3.2.1): we generated random sequences of 6-8 nucleotides, where the probability of

each base per site was 0.25, and selected those with pairwise differences >50% as our MIDs.

Prior to the metabarcoding PCR amplifications, we compared the efficiency of ZBJ-Art and

Formi primers, and generated pilot data to help decide the clustering thresholds needed to obtain

the most accurate taxonomic richness, by performing PCRs on an artificial positive control. This

sample contained a mixture of DNA derived from four species covering a broad taxonomic

breadth (1 Coleoptera: Carabidae, 1 Diptera: Sarcophagida, 2 Hymenoptera: *Tetramorium sp* and

Camponotus fulvopilosus).

PCR amplifications (25 μl) contained 1-3 μl of template DNA, 25 nM $MgCl_2$ (stock 25nM), PCR

buffer (stock 1x), 2 nM each dNTP, 0.6 nM each primer, 0.2U AmpliTaqGold DNA polymerase;

thermal cycling conditions for arthropods amplifications (primers Formi and ZBJ) were: 95 °C

for 5 min, followed by 33 cycles of 95 °C for 30 s, 48 °C for 30 s, and 72 °C for 30 s. Thermal

cycling conditions for plant amplifications were 95 °C for 5 min, followed by 38 cycles of 95 °C

for 30 s, 50.0 °C for 30 s and 72 °C for 30 s. A final extension step of 72 °C for 5 min was

included in all reactions. The optimal number of PCR cycles was established using a real-time

PCR assay in a subset of samples as the inflection point in the amplification curve and it

represents a compromise to obtain sufficient prey DNA while reducing clonality and consequent

bias in prey diversity. Real time PCR was performed in 20ul reaction containing 1-2 µl of

template, 1x buffer, 2.5 nM MgCl2, 2 nM each dNTP, 0.5U Amplitaq Gold, 1 nM of each

primer, and 1 µl SYBR Green/Rox mix (Invitrogen, USA). PCR amplifications were detected by

gel electrophoresis on 2% agarose gel stained with Gelred. All PCR preparations used aerosol

resistant filter tips, and were prepared in UV-sterilized laminar flow hoods. We note that

arthropods and fecal samples were processed independently, and the same applied to summer and

winter samples. This differential processing of samples was implemented as to avoid cross-

contamination.

All PCR products were visualized on a gel to confirm amplification success, and then pooled at

an approximately equimolar ratio into 11 pools (5 for summer and 6 for winter). We included the

extraction blanks in the pools, using 5 µl, despite no actual product visualized in agarose.

We removed unspecific fragments and primer dimer using Ampure XP SPRI beads, following a

double-sided protocol. Shortly, by adjusting the ratio of beads to DNA (0.7x) we first captured

the fragments larger than 300bp, keep the supernatant, which was in turn used to sequester

fragments larger than 120bp (beads:DNA ratio = 1.8x). The concentration of the purified pools

was determined in Qubit and efficiency of purification evaluated in and 2% agarose gel stained

with Gelred.

The PCR pools were converted into sequencing libraries using the NEBNext 6070 blunt end

library build kit following the manufacture's protocol. Index PCRs were carried out in 25 µl

reaction volumes, and each library was amplified in three replicates to maximize amplicon

diversity. Each indexing PCR consisted of 2.5 μl of library, 1x Gold buffer, 2.5 U AmpliTaqGold

DNA polymerase, 2.5 nM MgCl2 and 2 nM each dNTP, 0.2 μM forward indexed primer

(InPE1.0), and 0.2 μM reverse index primer such that library contained a different reverse index.

PCR was performed at 95˚C for 5min, followed by 8 cycles at 95˚C for 30 sec, 60˚C for 30 sec,

72˚C for 30 sec, and final extension at 72˚C for 5 minutes. We combined the three replicates of

each library and purified them using the QiaQuick columns (Qiagen, USA). To determine the

quality and quantity (concentration) of our 11 libraries, we used the TapeStation System (Agilent

Technologies, USA). We equalized the amplified library concentrations, then sequenced them on

an Illumina MiSeq platform using PE250 chemistry and spiked with 30% of PhiX.

Metabarcoding amplifications were performed in triplicate in order to enable us to: i) discard

potential errors (i.e. remove sequences only present in one PCR), and ii) optimize diversity (i.e.

keep sequences only if they were present in at least two PCRs). All PCRs included one negative

control to check for contamination and one positive control (DNA from an ant or plant species

not present in southern Africa) to ascertain amplification. PCR products were incorporated into

sequencing libraries (details in supplementary material) for generation of metabarcoding data

using an Illumina MiSeq platform with PE250 chemistry.

2.4 Sequence processing and taxonomic assignment

Post sequencing, the raw reads were processed with TRIMMOMATIC-0.36 (Bolger et al. 2014)

to first remove adapters (ILLUMINACLIP module), then bases with low quality. The quality of

the surviving reads (mean of surviving reads across pools: 93%) was checked with FASTQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and then the overlapping paired-end

reads were merged into contigs using PEAR (Hebert et al. 2003). These contigs were then

demultiplexed and filtered with DAMe (Zepeda-Mendoza et al. 2016) to only retain fragments

that matched the following criteria: i) present in at least two PCR replicates, ii) conform to the

expected size after primer trimming, and iii) present in at least 25 copies (abundance cut-off),

thus removing low abundance contigs that could be potential artefacts.

Clustering of sequences (within sample) into molecular operational taxonomic units (MOTUs)

was carried out with USEARCH (Edgar 2010) and followed the 3% sequence divergence criteria.

This clustering criteria was selected taking into account our empirical distribution of pairwise

percentage of identity between arthropod sequences (supplementary Figure S1) as well as the

reported variability in insect populations (Hebert et al. 2003). The MOTUs were aligned to the

National Center for Biotechnology Information nucleotide sequence database (NCBI; website:

http://www.ncbi.nlm.nih.gov/; accessed July 2017) the software Basic Local Alignment Search

Tool (BLAST 2.2.6+; (Camacho et al. 2009)). The high-scoring hits obtained with BLASTn (e-

value < 1e-30) were then parsed with MetaGenome Analyzer software (MEGAN v6.0; (Huson et

al. 2007)) to assigning the MOTUS to their lowest common ancestor (LCA) in the NCBI

taxonomic tree, as to meet the stringency criteria: min score = 130 for COI and 100 for rbcL, max

expected = 0.01, min percent identity = 0, top percent = 10, min support percent = 0 (off), min

support = 1, min complexity filter = 0.01. The MOTUs for which the LCA taxonomic assignment

fell below the thresholds or for which a match was not found in the NCBI database was

categorized as "unknown". Not all MOTUs could be assigned to species level due to the

incompleteness of the reference database for our study area. Therefore we present results

identified to family-level. We contend this still provides a good grasp of the dietary components

as well as the arthropod communities.

2.5 Statistical Analyses

All analyses were implemented in R v3.3.2 (R Development Core Team). We estimated

Shannon-Wiener's (H) and Simpon's (1-D) indices to quantify both consumed and available prey

diversity using *vegan* package (Oksanen 2017). Diversity indices were compared among sites

using the non-parametric Kruskal-Wallis test, as the data was not normally distributed (Shapiro-

Wilk test, $p < 0.05$). Changes in composition of prey were inspected using a non-metric

multidimensional scaling (NMDS) with Bray-Curtis dissimilarity. To test whether region

(*Coastal, Central, Inland;* proxy for primary productivity and energetic demands) is shaping

diversity in available and consumed prey we used a Mantel test (Mantel 1967) with 999 random

permutations as implemented in *vegan* (Oksanen 2017). Briefly, we compared the observed

dissimilarity matrix (Bray-Curtis dissimilarity) against a conceptual matrix for regional effects,

coded as zero for same region comparisons and one for different region comparisons.

Trophic niche breadth was estimated using a standardized Levin's index (Bs; Levins 1968).

Standardized Levin's index (Bs) was estimated as: $Bs=(1/\sum p_i^2)-1/n-1$, where p_i is the proportion

of individuals consuming arthropod family i and n is the total number of families available. This

index measures the uniformity of distribution of individuals (birds) among resource states

(arthropod families in each region) and it is closer to zero when diets are dominated by few items

(specialist) and closer to one for a generalist diet. We created Bs 95% confidence intervals for

each region by bootstrapping with 10,000 replicates. In addition, to test whether the observed Bs

values were significantly different from a scenario of "no regional difference" we used a

permutational approach. All occurrences (p_i) were pooled regardless of their regional origin, then

we randomly sampled n arthropod families (from the observed range: 37 – 45) and finally

estimated Bs_random. This procedure was repeated 9999 times to generate the null distribution.

The estimated probability of obtaining a result that exceeded the observed value under the null

hypotheses of "no region difference" was estimated as p = [(number Bs_random >

Bs_observed)/total number randomizations].

For the most common arthropod MOTUs in the diet of robins, we tested whether selectivity of

prey changed among regions, by calculating Ivlev's electivity index (Ivlev 1961) for each region

as implemented in *selectapref* package. This index varies from "-1" (avoidance of prey) to "+1"

(total reliance on prey), and "0" indicates prey selection is proportional to availability.

3. RESULTS

3.1 Arthropod availability

Our survey of potential Arthropoda prey availability across the gradient revealed 519 MOTUs,

with the majority (83%) identified to class Insecta (Table S3). Coleoptera, Diptera, Hymenoptera

and Lepidopera were the most abundant orders (Figure2). The most diverse community was

found in the *Central* region, the area with largest annual primary productivity (192 MOTUS, 45

Families, Shannon-Wiener's H = 3.44, Simpson's 1-D = 0.95; Table 1), whereas the in region

with lowest NDVI and highest temperature amplitude, *Inland,* the arthropod community was least

diverse (157 MOTUs, 42 Families, Shannon-Wiener's H = 3.30, Simpson's 1-D = 0.94; Table 1).

However, when these observations were tested statistically, we were unable to reject the null

hypothesis that there is no regional effect on arthropod community (Mantel$_{region}$, p = 0.430).

3.2 Arthropods and plants consumed

After data filtering criteria, 82% of fecal samples provided arthropod MOTUs (170). Insecta was

the richest Arthropoda class (127 MOTUs) with the orders Coleoptera, Hymenoptera,

Lepidoptera and Orthoptera being the most abundant (Figure 2, Table S3). Although the Karoo

scrub-robin is viewed as insectivorous, we detected plant DNA in 94% of the fecal samples.

Overall, 86 MOTUs were assigned to Spermatophyta (seed Plants) with Solanales and

Aspargales being the most abundant (Table S4).

Our data showed no evidence that dietary composition was associated with region whether

considering only arthropod prey (Shannon-Wiener's H: Kruskal-Wallis X^2 = 1.855, df = 2, p =

0.396; Simpson´s 1-D: Kruskal-Wallis X^2 = 1.885, df = 2, p = 0.390; Mantel$_{region}$, p = 0.072;

Figure 3) or combing arthropod prey and plant items (Shannon-Wiener's H: Kruskal-Wallis X^2

= 0.745, df = 2, p = 0.689; Simpson's 1-D: Kruskal-Wallis X^2 = 0.422, df = 2, p = 0.810;

Mantel$_{region}$ p = 0.433; Figure 3).

3.3. Trophic niche and prey selection

When accounting for available arthropod prey, we found that birds living in the *Inland* region -

that which was least productive and exerted high energetic demands on the robins – appeared to

have the narrowest dietary niche breadth (Bs = 0.372, 95% CI = 0.241 – 0.677, Figure 3A). This

contrasts with the *Coastal* population, whose individuals presented the widest niche (Bs = 0.479,

95% CI = 0.328 – 0.667; Figure 3A). However, again when statistically tested, we found that the

observed Bs values for each of the three study regions was not significantly different from the

values expected under the null hypothesis of no regional difference (mean Bs_random = 0.445,

95% CI = 0.247 - 0.592, p$_{Coastal}$= 0.658, p$_{Central}$ = 0.568, p$_{Inland}$ = 0.738; Figure S2).

An electivity analysis restricted to the five most abundant potential prey items (Coleoptera,

Diptera, Hymenoptera, Lepidoptera and Orthoptera), showed that robins from all regions

positively selected for Hymenoptera:Formicidae (Ivlev´s electivity > 0; Figure 4). Electivity was,

however, largest in the *Inland* population, with 55% of sampled birds consuming ants. Although

in *Central* region relatively more Coleoptera families were potentially available as prey (Figure

S3), robins did not positively selected on this prey type (Figure 4). We also found that Dipterans,

another widely available potential prey source, were negatively selected on in all regions (Ivlev´s

electivity < 0; Figure 4)

4. DISCUSSION

Species living in spatially and temporally heterogeneous environments must adjust their behavior and physiology in order to persist. Dietary plasticity may thus be an advantageous trait in energy-depauperated ecosystems as switching from alternative resources allows them to deal with environmental change (e.g.: (Varner and Dearing 2014)). Here, we investigated dietary patterns among conspecific Karoo scrub-robin populations living in regions with different productivity and thermoregulatory demands. Overall our findings revealed that: i) although Karoo scrub-robins are generalist insectivores, some plant items seem to be included in its diet; ii) dietary composition was not associated with region; iii) when accounting for available arthropod prey the dietary niche breadth of each population was not different from a scenario of no regional effect (i.e., similar productivity and thermoregulatory demands); iv) Formicidae (ants) were positively selected in the three populations, yet *Inland* robins showed the largest electivity.

Dietary niche breadth

The concomitant assessment of available and consumed prey in conspecific populations revealed that the foraging behavior of robins is not simply based on encounter rate: Coleopterans, Hymenopterans and Orthopeterans were preferred prey. While coleopterans represent less than 16% of available prey (7%, 16% and 11% of MOUTUs in *Coastal, Central* and *Inland*, respectively) they were clearly preferentially preyed upon (representing 33% of total dietary MOTUs in the *Coastal* region, 23% in the *Central* region and 20% in the *Inland* region). A similar pattern of preference was found for Hymenopterans in the *Central* and *Inland* regions: Hymenopetans were consumed at a higher rate than their relative availability (comprising 11% and 19% of the dietary MOTUs ,*vs* 7% and 11% of available MOTUs). The exception was found in the area with intermediate primary productivity and temperature amplitude – the *Coastal*

region – where Hymenopetans were only 4% of available prey and 5% of consumed MOTUs. In

contrast, Diptera and Lepidopetra were consumed at a rate inversely proportional to their

presence, i.e. they were avoided by the birds. These results are in accordance to prey descriptions

based on Karoo scrub-robins stomach contents (invasive method) sampled throughout the year in

a region East of our *Inland* site (Free State; Oatley 1970). Specifically, Oatley revealed that ants

(Hymenoptera) and beetles (Coleoptera) were present in 55% and 14% of the stomachs,

respectively. We also noted that while terrestrial arthropods are the bulk of robin's prey, they

appear to supplement their diets with plant matter (94% of samples yielded plant DNA).

Although, this finding could in theory simply indicate that robins are plant secondary consumer

through herbivore arthropods, we contend the latter is unlikely given both the anticipated

degradation to the plant DNA that would be incurred through two rounds of ingestion and

degradation, and perhaps more importantly, two additional lines of evidence. Firstly, there are

unpublished observations (P. Lloyd, pers. comm.) of birds foraging on berries and plant shoots,

and secondly, seeds and berries were detected in the stomachs contents or robins (Oatley 1970).

Thus together, it suggests that that plant matter is not simply a serendipitous food item, but may

in fact be selected for.

Our study sites are embedded within the Southern African semi- and arid-regions, where seasonal

pulses of rainfall are known to determine arthropod densities (Dean and Milton 1999) and

subsequently food supplies for insectivores (Lloyd 1999). Surprisingly, despite the differences in

primary productivity among our study sites, the dietary niche breadth (Bs) among populations

was not different from a scenario of homogeneous environment (i.e., no difference in

productivity or thermal demands). This plasticity in diet may be favorable in environments where

prey is patchily distributed in space and time, such as the Southern African semi- and arid-

regions (Dean and Milton 1999). Moreover, this finding corroborates the expectations of Optimal

Foraging Theory that animals can only afford to be specialist (small niche breadth) where

resources are readily available (Stephens and Krebs 1986).

Life in arid environments

Life in arid environments is harsh for endotherms in general, and in particular for ground-feeding

insectivore birds as they need to fuel their typically high avian metabolic rates in energy-

depauperated ecosystems. We predicted the robins living in the most demanding regions (*Inland*)

would exhibit the most generalist feeding strategy, under the rationale that as this was the most

energy-depauperated ecosystem and energetically demanding environment, their diet should

become more generalist as ignoring potential sources of energy is not an option (Levins and

MacArthur 1969; Schoener 1971). Specifically, we expected robins living in the *Inland* region to

prey on the greatest diversity of ground arthropods, so as to obtain sufficient energy to fuel the

thermogenic demands of winter (burn energy to produce heat as a means to warm-up

(thermogenesis; (Hothola 2004)) as well as actively cool down in summer through evaporative

cooling (Williams and Tieleman 2005). In contrast however, we found that the *Inland* robins prey

on an array of arthropods that is similar to robins living in the less demanding region (*Coastal*).

We also noted they exhibit a preference for ants (Formicidae), a finding that lends support to the

notion that dependable food sources such as ants are critical for arid endemic birds (Dean and

Milton 2017), yielding the necessary energy, nutrients and water (robins do not drink) to

overcome the challenges that the environment exerts.

We also hypothesize that supplementation of their diet with plant items such as berries (the most

common plant in diet, Solanaceae, produces berries) may provide the carbohydrates and hence

calories required to cover their general energetic budget, when sufficient animal protein and fat is

not available. Furthermore beside sugars, fruits would also provide extra water, a critical

component when dealing with hot temperatures: birds loose water to cool down and avoid

hyperthermia in the hot days. Although we did not test for seasonal differences, it is worth noting

that plant DNA was mostly found in samples collected in winter. We hypothesize this may be

related to high energy demands to deal with cold as well as short time-window to actively search

for arthropod prey on the ground (shorter periods of day light).

eDNA metabarcoding in trophic ecology

Investigating foraging strategies by applying molecular methods to environmental and non-

invasive samples is increasingly facilitating the study of highly-mobile and evasive animals (e.g.:

Jedlicka et al. 2013; Mallott et al. 2014; Jedlicka et al. 2017). Nevertheless, there are several

caveats underlying inferences of the dietary components and arthropod communities based on

DNA recovered from fecal and environmental samples: i) data is typically qualitative allowing

only presence/absence inferences, ii) primer specificity may bias detection of some taxa, iii) the

differential digestibility of hard- *vs* soft-bodied prey may bias results, iv) incompleteness of

reference databases can hinder species-level taxonomic assignment, v) samples, either fecal or

environmental, represent a snapshot of the dietary diversity available and preyed on, and iv)

inferences of herbivory may be confounded if animals are actively consuming herbivorous

arthropods. Nevertheless, we contend the dietary patterns obtained in this study are robust and

provide a first grasp on how arid endemic birds may be ensuring sufficient energy yields in a

rather challenging environment. Thus we hope our findings provide a stepping-stone for further

research into the role of plasticity in foraging strategies. Moreover, studies like this one are both

relevant for evolutionary biologists because describing the factors that lead to dietary variation in

natural populations is fundamental for understanding local adaptation and speciation (e.g.: Grant

et al. 2008; Svanbäck and Schluter 2012), as well as for conservation biologists because the on-

going global change is expected to increase aridity and consequently alter food-webs, which in

turn affect avian communities (e.g.: Iknayan and Beissinger 2018).

REFERENCES

Bicudo, J. E. P. W., W. A. Buttemer, M. A. Chappell, J. T. Pearson, and C. Bech. 2010. Ecological and environmental physiology of birds. Oxford University Press, Oxford.

Binladen, J., M. T. P. Gilbert, J. P. Bollback, F. Panitz, C. Bendixen, R. Nielsen, and E. Willerslev. 2007. The use of coded PCR primers enables high-throughput sequencing of multiple homolog amplification products by 454 parallel sequencing. PLoS ONE 2:e197.

Bohmann, K., A. Monadjem, C. L. Noer, M. Rasmussen, M. R. K. Zeale, E. Clare, G. Jones, E. Willerslev, and M. T. P. Gilbert. 2011. Molecular Diet Analysis of Two African Free-Tailed Bats (Molossidae) Using High Throughput Sequencing. PLoS ONE 6:e21441.

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120.

Borer, E. T., E. W. Seabloom, D. Tilman, and V. Novotny. 2012. Plant diversity controls arthropod biomass and temporal stability. Ecology Letters 15:1457–1464.

Brandon-Mong, G. J., H. M. Gan, K. W. Sing, P. S. Lee, P. E. Lim, and J. J. Wilson. 2015. DNA metabarcoding of insects and allies: an evaluation of primers and pipelines. Bull. Entomol. Res. 105:717–727.

Burgar, J. M., D. C. Murray, M. D. Craig, J. Haile, J. Houston, V. Stokes, and M. Bunce. 2014. Who's for dinner? High-throughput sequencing reveals bat dietary differentiation in a biodiversity hotspot where prey taxonomy is largely undescribed. Mol. Ecol. 23:3605–3617.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, and T. L. Madden. 2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421. BioMed Central.

Cheli, G. H., and J. C. Corley. 2010. Efficient sampling of ground-dwelling arthropods using pitfall traps in arid steppes. Neotropical entomology.

Dean, W. R. J., and S. J. Milton. 2017. Ants (Formicidae) as food for birds in southern Africa: opportunism or survival? Ostrich 89:1–4.

Dean, W. R. J., and S. Milton. 1999. The Karoo. Cambridge University Press.

Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics
26:2460–2461.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces
for global land areas. Int. J. Climatol. 37:4302–4315.

Gilbert, M. T. P., W. Moore, L. Melchior, and M. Worobey. 2007. DNA Extraction from Dry
Museum Beetles without Conferring External Morphological Damage. PLoS ONE 2:e272.

Grant, P. R. and B. R. Grant. 2008. How and why species multiply: the radiation of Darwin's
finches. Princeton, NJ [etc.]: Princeton University Press.

Hamad, I., E. Delaporte, D. Raoult, and F. Bittar. 2014. Detection of termites and other insects
consumed by African great apes using molecular fecal analysis. Sci Rep 4:4478.

Hebert, P. D. N., A. Cywinska, S. L. Ball, and J. R. deWaard. 2003. Biological identifications
through DNA barcodes. Proceedings of the Royal Society B: Biological Sciences 270:313–321.

Hothola, E. 2004. Shivering Thermogenesis in Birds and Mammals. Pp. 241–252 *in* B. M. Barnes
and H. V. Carey, eds. *Life in the cold: evolution, mechanisms, adaptation, and application.* 12th
International Hibernation Symposium.

Huson, D. H., A. F. Auch, J. Qi, and S. C. Schuster. 2007. MEGAN analysis of metagenomic
data. Genome Research 17:377–386.

Iknayan, K. J., and S. R. Beissinger. 2018. Collapse of a desert bird community over the past
century driven by climate change. Proceedings of the National Academy of Sciences 115:8597–
8602.

Ivlev, V. S. 1961. Experimental Ecology of the Feeding of Fishes. Yale University Press.

Jedlicka, J. A., A. M. Sharma, and R. P. P. Almeida. 2013. Molecular tools reveal diets of
insectivorous birds from predator fecal matter. Conservation Genet Resour 5:879–885.

Jedlicka, J. A., A.-T. E. Vo, and R. P. P. Almeida. 2017. Molecular scatology and high-
throughput sequencing reveal predominately herbivorous insects in the diets of adult and nestling

Western Bluebirds (Sialia mexicana) in California vineyards. The Auk 134:116–127.

Levins, R. 1968. Evolution in Changing Environments. Princeton University Press.

Levins, R., and R. MacArthur. 1969. An Hypothesis to Explain the Incidence of Monophagy.
Ecology 50:910–911. Ecological Society of America.

Lloyd, P. 1999. Rainfall as a breeding stimulus and clutch size determinant in South African arid-
zone birds. Ibis 141:637–643. Wiley Online Library.

MacArthur, R. H., and E. R. Pianka. 1966. On the optimal use of a patchy environment. The
American Naturalist 100:603–609.

Maclean, G. L. 1969. The breeding seasons of birds in the south-western Kalahari. Ostrich.

Mallott, E. K., R. S. Malhi, and P. A. Garber. 2014. High-throughput sequencing of fecal DNA to
identify insects consumed by wild Weddell's saddleback tamarins (Saguinus weddelli, Cebidae,
Primates) in Bolivia. Am. J. Phys. Anthropol. 156:474–481.

Mantel, N. 1967. The detection of disease clustering and a generalized regression approach.
Cancer Res. 27:209–220.

Martin, R. A., and D. W. Pfennig. 2009. Disruptive selection in natural populations: the roles of
ecological specialization and resource competition. The American Naturalist 174:268–281.

Nielsen, M. 2015. Insect soup, eDNA and Robots. University of Copenhagen, Copenhagen.

Oatley, T. B. 1970. Observations on the food and feeding habits of some African robins
(Aves:Turdinae). Ann. Natal Mus. 20:293–327.

Oksanen, J. 2017. Package "vegan." 1–292.

Poinar, H. N., M. Hofreiter, W. G. Spaulding, P. S. Martin, B. A. Stankiewicz, H. Bland, R. P.
Evershed, G. Possnert, and S. Pääbo. 1998. Molecular coproscopy: dung and diet of the extinct
ground sloth Nothrotheriops shastensis. Science 281:402–406.

Pompanon, F., B. E. Deagle, W. O. C. Symondson, D. S. Brown, S. N. Jarman, and P. Taberlet.
2012. Who is eating what: diet assessment using next generation sequencing. Mol. Ecol.

21:1931–1950.

Roches, Des, S., L. J. Harmon, and E. B. Rosenblum. 2016. Colonization of a novel depauperate
habitat leads to trophic niche shifts in three desert lizard species. OIKOS 125:343–353.

Schoener, T. W. 1971. Theory of Feeding Strategies. Annu. Rev. Ecol. Evol. Syst. 2:369–404.

Stephens, D. W., and J. R. Krebs. 1986. Foraging Theory. Princeton University Press.

Svanbäck, R., and D. Schluter. 2012. Niche specialization influences adaptive phenotypic
plasticity in the threespine stickleback. Am. Nat. 180:50–59.

Varner, J., and M. D. Dearing. 2014. Dietary plasticity in pikas as a strategy for atypical resource
landscapes. J Mammal 95:72–81. Oxford University Press.

Williams, J. B., and B. I. Tieleman. 2005. Physiological adaptation in desert birds. BioScience
55:416–425. BioOne.

Zeale, M. R. K., R. K. Butlin, F. K. Barker, D. C. Lees, and G. Jones. 2011. Taxon-specific PCR
for DNA barcoding arthropod prey in bat faeces. Mol Ecol Resour 11:236–244.

Zepeda-Mendoza, M. L., K. Bohmann, A. Carmona Baez, and M. T. P. Gilbert. 2016. DAMe: a
toolkit for the initial processing of datasets with PCR replicates of double-tagged amplicons for
DNA metabarcoding analyses. BMC Res Notes 9:255.

Acknowledgments

We thank the Southern African authorities for providing permits, the Danish National High-

Throughput DNA Sequencing Centre for generating sequence data, Nick Pattinson for his help in

the field, Lara Puetz and Sarah Mak for useful lab hints, Shyam Gopalakrishnan for assistance

with bioinformatics, and Rute Fonseca for valuable discussions.

Ethics and Permits

Capture permits were issued by Cape Nature (0056-AAA008-00057) and the Department of

Environment and Nature Conservation (1611/2015), South Africa. Ethical approval was obtained

from the Animal Research Ethics Committee at Nelson Mandela University (A15-SCI-ZOO-

005).

Author contributions

AMR and MTPG conceived the study. AMR and BS collected samples. AMR performed

molecular work and conducted analyses. AMR, BS and MTPG wrote the paper.

Funding

This study was funded by a Marie Sklodowska-Curie Individual fellowship to AMR (Grant

655150 - BARREN).

Data accessibility

Data reported are available as supplementary material.

1. Fasta file with Formi primers aligned against Formicidae.

2. Fasta files with available and consumed MOTUs obtained with different primer sets.

    TABLE

Table1. Available arthropod diversity and dietary diversity (Arthropod and Spermatophyta: seed plants) among populations of Karoo scrub-robin. T_{amp}: annual temperature range. In parenthesis we denote relative arthropod consumption, i.e. % of prey consumed in relation to prey available.

	Coastal	Central	Inland
	NDVI: 0.43	NDVI: 0.50	NDVI: 0.30
	T_{amp}: 21.5 °C	T_{amp}: 27.7 °C	T_{amp}: 32.1 °C
Available Arthropod MOTUs	170	192	157
Available Families	37	45	44
Shannon-Wiener's H	3.109	3.422	3.300
Simpson's 1-D	0.930	0.953	0.943
Consumed Arthropod MOTUs	36 (21%)	57 (30%)	64 (40%)
Consumed Families	20 (54%)	25 (56%)	28 (64%)
Shannon-Wiener's H	2.880	2.706	2.947
Simpson's 1-D	0.936	0.878	0.906
Consumed Spermatophyta MOTUs	21	22	43
Consumed Families	12	14	21
Shannon-Wiener's H	3.404	3.684	3.727

Figures

Figure 1. Energetic landscape of the three study sites and populations of Karoo scrub-robin (*Cercotrichas coryphaeus*). (A) Map of annual primary productivity (NDVI; the darker the pixel the larger the productivity) with depiction of the three study sites. Karoo scrub-robin range limit is delimited by a dashed line. (B) Primary productivity (NDVI), Temperature amplitude (ºC) and energetic physiology (metabolic expansibility) for the study populations (from Ribeiro et al unpublished data) three study sites. Karoo scrub-robin range limit is delimited by a dashed line.

Figure 2: Diversity of Arthropod MOTUs per Family in pitfalls and faecal samples at three study locations: *Coastal*, *Central* and *Inland*. Bubble size is proportional to the number of MOTUs found in each family. Arthropod Orders are annotated with initials as a prefix to Family names (e.g.: Ar_Agelenidae): Ar: Araneae, As: Astigmata, Di: Diptera, En: Entomobryomorpha, He: Hemiptera, Hy: Hymenoptera, Is: Isopetra, Od: Odonata, Or: Orthtoptera, Pl: Pleurostigmophora, Po: Polydesmida , Ps: Psocoptera, Sc: Scutigemorpha, Sp: Spiristrepida, Th: Thysanoptera,

Figure 3. Diversity in arthropods potentially available to, and consumed by, the Karoo-scrub robin. (A) Simpson's 1-D diversity index and Levin's Bs niche breadth ($Bs_{Coastal}$ 95% CI = 0.328 − 0.667, $Bs_{centrall}$ 95% CI = 0.285 − 0.659, Bs_{Inland} 95% CI = 0.241 − 0.677). (B) NMDS for arthropod prey and (C) NMDS for plant items consumed; $STRESS_{arthropods}$ = 0.17, $STRESS_{plants}$ = 0.14.

Figure 4. Ivlev's electivity index for the main arthropod orders consumed: Coleoptera (Co) Diptera (Di), Hymenoptera (Hy), Lepidoptera (Le) and Orthoptera (Or). Positive values indicate prey selection whereas negative values indicate prey avoidance.

Figure 1

Figure 2

Figure 3

(A)

consumed   available

Simpson's 1-D

0.94 0.93   0.95   0.91 0.94
      0.88

Coastal   Central   Inland

Levin's Bs

0.48   0.43   0.37

(B)

NMDS2

0.2

0.0

−0.2

Coastal
Central
Inland

−0.25   0.00   0.25

NMDS1

(C)

NMDS2

−1

Coastal
Central
Inland

−4   −2   0   2

NMDS1

Figure 4