Title:

Proteasome inhibitor-induced modulation reveals the spliceosome as a

specific therapeutic vulnerability in multiple myeloma

Authors:

Hector H. Huang[1], Ian D. Ferguson[1], Alexis M. Thornton[2], Christine Lam[1], Yu-

Hsiu T. Lin[1], Prabhakar Bastola[1], Priya Choudhry[1], Margarette C. Mariano[1],

Makeba Marcoulis[1], Julia Malato[3], Paul Phojanakong[3], Thomas G. Martin[4],

Jeffrey L. Wolf[4], Sandy W. Wong[4], Nina Shah[4], Byron C. Hann[3], Angela N.

Brooks[2], Arun P. Wiita[1,3,*]

Affiliations:

[1]Dept. of Laboratory Medicine, University of California, San Francisco, CA [2]Dept.

of Biomolecular Engineering, University of California, Santa Cruz, CA, [3]Helen

Diller Family Comprehensive Cancer Center, University of California, San

Francisco, CA, [4]Dept. of Medicine, University of California, San Francisco, CA

Running Title:

PI response reveals spliceosome as therapeutic target in MM

Keywords:

myeloma, proteasome inhibitor, phosphoproteomics, splicing, mass spectrometry

Financial support:

This work was supported by NIH/NCI P30CA083103 supporting UCSF Preclinical

Therapeutic Core facility managed by B.C.H.), NIH/NHGRI T32HG008345 (to

A.M.T.), the Damon Runyon Cancer Research Foundation Dale Frey

Breakthrough Award (DFS 14-15), NIH/NCI K08CA184116, NIH/NIGMS

DP2OD022552, and the UCSF Stephen and Nancy Grand Multiple Myeloma

Translational Initiative (to A.P.W.) and NIH/NCI R01CA226851 (to A.N.B. and

A.P.W.)

Declaration of Interests

The authors declare no relevant conflicts of interest.

*Corresponding Author Contact information:

Arun P. Wiita, MD, PhD

tel: (415) 514-6238

fax: (415) 353-4877

Arun.wiita@ucsf.edu

UCSF Dept. of Laboratory Medicine

185 Berry St., Ste. 290

San Francisco, CA 94107

Abstract:

Enhancing the efficacy of proteasome inhibitors is a central goal in myeloma therapy. We proposed that signaling-level responses after PI would reveal new mechanisms of action that could be therapeutically exploited. Unbiased phosphoproteomics after the PI carfilzomib surprisingly demonstrated the most prominent phosphorylation changes on splicing related proteins. Spliceosome modulation was invisible to RNA or protein abundance alone. Transcriptome analysis after PI demonstrated broad-scale intron retention, suggestive of spliceosome interference, as well as specific alternative splicing of protein homeostasis machinery components. These findings led us to evaluate direct spliceosome inhibition in myeloma, which synergized with carfilzomib and showed potent anti-tumor activity. Functional genomics and exome sequencing further supported the spliceosome as a specific vulnerability in myeloma. Our results propose splicing interference as an unrecognized modality of PI mechanism, reveal additional modes of spliceosome modulation, and suggest spliceosome targeting as a promising therapeutic strategy in myeloma.

Significance:

New ways to enhance PI efficacy are of major interest. We combine systems-level analyses to discover that PIs specifically interfere with splicing and that myeloma is selectively vulnerable to spliceosome inhibition. We reveal a new approach to advance myeloma therapy and uncover broader roles of splicing modulation in cancer.

Keywords:

Mass Spectrometry, Proteomics, Splicing, Alternative Splicing, Phosphoproteomics, Myeloma, Spliceosome, SF3B1, SRSF1

INTRODUCTION

Multiple myeloma is a clonal malignancy of plasma cells with no known cure. Like normal plasma cells, myeloma cells produce and secrete incredible amounts of immunoglobulin. This unique function may be exploited by therapeutically inhibiting the proteasome using the FDA-approved proteasome inhibitors (PIs) bortezomib, carfilzomib, and ixazomib. Proteotoxic stress caused by these first-line therapeutic agents has been proposed to induce the apoptotic function of the unfolded protein response (UPR) (1), leading to plasma cell death while largely sparing normal tissues (2, 3). However, despite the appealing simplicity of this mechanism, the canonical UPR is not always strongly induced in myeloma cells by PIs (4) and is unlikely to be the sole mode of PI cytotoxicity in MM. Indeed, many additional mechanisms of action of PIs have also been proposed, ranging from NF-kB inhibition to immune microenvironment effects to aberrant recycling of cytosolic amino acids (5, 6).

Identifying the full range of PI mechanisms of action remains relevant given that acquired PI resistance is clinically widespread but its origins remain unclear (7, 8) and finding new methods to specifically target PI-resistant disease, or molecules to synergize with PIs to avoid resistance by driving deeper remissions, remains a long-standing goal. As one approach to achieving this goal, we and others have studied the response of malignant plasma cells to PIs using both gene expression and proteomic methods (9-11). Notably, one of the most prominent features of the cellular response to PIs is the activation of the heat shock response (12). This mechanism leads to significant induction of cytosolic protein-folding chaperones, possibly to assist in protein refolding and decrease in unfolded protein stress. We and others (9, 12, 13) have therefore proposed targeting mediators of the heat shock response as potential combination therapies with PIs.

However, one unresolved question is whether proteasome inhibition may carry additional effects on plasma cells that are not revealed by mRNA or protein abundance analysis alone. We hypothesized that additional modalities of response, and thereby new myeloma-relevant therapeutic targets, may be

revealed by studying the signaling-level response to PIs with unbiased mass spectrometry-based phosphoproteomics. The large majority of therapy-relevant investigations using this technique have focused on elucidating the effects of kinase inhibitors (14). However, we reasoned that a significant cellular perturbation such as proteasome inhibition would likely also indirectly perturb kinase and phosphatase signaling in a broad fashion.

Here, we used unbiased phosphoproteomics to quantify >5000 phosphopeptides in myeloma cells exposed to the irreversible PI, carfilzomib (Cfz). Surprisingly, we found the greatest increases in phosphorylation occurred in proteins associated with the spliceosome machinery. A link between these processes was invisible at the gene expression level. We further evaluated this link from a mechanistic and therapeutic perspective, finding that PIs lead to specific disruption of normal splicing. We suggest interference of splicing as an additional mechanism of action of PIs not previously explored. Inhibition of splicing has recently become a promising therapeutic strategy in other hematologic malignancies (15). Our results reveal an intersection of cellular stress and the splicing machinery, which may have broad relevance in biology. Furthermore, we propose the spliceosome as a new and potentially selective therapeutic target in myeloma.

RESULTS

Proteasome inhibition results in sustained phosphorylation of splicing factors in myeloma plasma cells

We first used unbiased phosphoproteomics to examine the signaling-level response of MM.1S multiple myeloma cells to Cfz. We chose time points across 24 hours for analysis based on our prior results demonstrating that the transcriptional and proteomic response to proteasome inhibition evolves over many hours (9). This is in contrast with the majority of prior perturbation phosphoproteomic studies, which have typically examined direct effects on kinase activation or inhibition on a timescale of minutes (14). Here, we instead

consider the indirect effects on phosphorylation induced by PI exposure. Using

label-free quantification of immobilized metal affinity chromatography (IMAC)-

isolated phosphopeptides, we indeed found that altered phosphorylation

signatures were most prominent 24 hours after treatment (**Fig. 1 and Fig. S1**). In

total, we quantified 5791 phosphosites in at least one technical replicate of the

time course, with >99% of phosphosites representing Ser or Thr phosphorylation

events, as expected using this enrichment technique. Notably, with 30 nM Cfz,

cell viability was approximately 30% of baseline, indicating significant drug-

induced cytotoxicity by this final time point.

At each time point we simultaneously performed single-end RNA-seq to

determine gene expression to compare with our phosphoproteomic results. **Fig.**

1B shows 58 upregulated (red) and 75 downregulated (blue) phosphopeptides

from proteins with largely unchanged RNA transcript abundance as detected by

unsupervised hierarchical clustering. Upon this initial analysis, we were

encouraged to find decreased phosphorylation of the translation factor EIF4E-

BP1 as well as the ribosomal subunit RPS6 (**Fig. 1B**). These phosphorylation-

level responses related to suppressed translation are expected upon PI-induced

cellular stress (9). While other downregulated phosphopeptides did not suggest a

specific highly-enriched biological function, upon manual inspection of

upregulated phosphosites we were surprised to find that 14 of 58 were present

on proteins related to pre-mRNA splicing. These primarily included

phosphopeptides deriving from the heterogeneous ribonucleoprotein (HNRNP)

family of proteins as well as phosphopeptides belonging to the SRSF family of

splicing factors (**Fig. 1B**). In particular, the arginine- and serine-rich "RS" domain

of the SRSF proteins are known to have their splicing activity modulated by

phosphorylation (16). Notably, these prominent signaling-level effects on splicing

factors were invisible to prior gene expression studies of PI response and have

not been investigated previously. We therefore chose to further explore the

interaction between PIs and the splicing machinery.

To validate this initial result from label free quantitative proteomics, we

prepared independent samples using a stable isotope labeling (SILAC)

phosphoproteomics approach. Based on our results above, we examined only the 24 hr time point in MM.1S cells. We evaluated both a low dose (10 nM, $n = 2$ biological replicates) and a moderate dose (18 nM, $n = 2$) of Cfz (**Fig. 2A-B**). With this lot of Cfz, 10 nM drug elicited ~20% cell death after 24 hr, while 18 nM killed ~85% of cells (**Fig. S3A**). Of the 520 phosphosites significantly ($p < 0.05$; ≥ 2-fold-change) upregulated in MM.1S treated with 18 nM Cfz in **Fig. 2A**, 127 (24.4%) are associated with splicing-related proteins, with 23 of these as part of the SRSF protein family of splicing factors. Background-corrected Gene Ontology (GO) analysis confirms that all of the top enriched biological processes involve RNA splicing regulation and mRNA processing (**Fig. S1B, 2E**). At 10 nM Cfz, though, this signaling response is much weaker with only 25 upregulated phosphosites; none of these are splicing-related. These results suggest that there is a strong dose-response effect of phosphorylation changes after proteasome inhibition, both across splicing factors and the broader proteome.

  To compare these changes at the signaling level to changes at the protein level, unenriched peptides were also analyzed by LC-MS/MS (**Fig. S2A-B**). Confirming expected responses to proteasome inhibition, the most upregulated proteins included heat shock-induced chaperones as well as SQSTM1/p62 associated with autophagy (9). In contrast, splicing factors with increased phosphorylation sites do not significantly change in abundance, confirming that phosphosite increases are due to changes at the signaling level and not protein copy number.

Melphalan induces a similar but not identical phosphorylation response

  We next investigated whether this broad splicing factor phosphorylation phenotype was unique to proteasome inhibition or was also seen under a different drug mechanism of action. We chose to compare the response of MM.1S cells to melphalan, a DNA alkylating agent and clinically-used myeloma therapeutic. In parallel, we also treated another MM cell line, AMO-1, with Cfz to determine if the phosphorylation response to proteasome inhibitor is consistent across cell line models.

For these experiments we again used a single-timepoint SILAC approach. Here, both 10 µM melphalan and 15 nM Cfz led to ~20% cell death in MM.1S and AMO-1, respectively, at 24 hr (**Fig. S3A**). Western blot confirmed induction of DNA damage by melphalan and proteotoxic stress response for Cfz (**Fig. S3C-D**). Compared to 18 nM Cfz, we saw largely decreased phosphorylation-level responses to both of these agents (**Fig. 2C-D**). Of 113 phosphosites significantly upregulated in AMO-1, 7 belong to splicing related proteins (SRSF2, SRSF6, SRRM1, HNRNPH1, TRA2A, DDX1). This result is consistent with the MM.1S results in **Fig. 2A-B**, where greater PI response correlates with more prominent phosphorylation changes.

Under 10 µM melphalan, 93 phosphosites were significantly upregulated, with 8 sites on splicing related proteins (HNRNPK, TRA2A, SRRM2, and WDR77), although none are SRSF family members (**Fig. 2D**). Furthermore, as expected, both unenriched shotgun proteomics and RNA-seq for gene expression confirm that proteasome inhibition and DNA damage elicit different responses (**Fig. S2A-E**). Again, no splicing factors with altered phosphosites under either drug treatment were changed at the protein abundance level.

We further performed Kinase Set Enrichment Analysis (KSEA) (17) on our MM.1S datasets to identify kinases whose activity may regulate differential phosphorylation found by phosphoproteomics. While this tool is limited by its reliance on well-characterized kinase-substrate relationships, and despite the different number of phosphosites upregulated under each condition, within this framework this tool identified similar kinases active under both 18 nM Cfz and 10 µM melphalan treatment (**Fig. 2F**). Notably, both drugs are predicted to induce activity of cdc2-like kinase 1 (CLK1), a kinase known to phosphorylate SRSF family splicing factors among other proteins (18). However, in line with the specific biology of PIs, Cfz also strongly induced inhibitory kappa B kinase (IKBKB) activity, a kinase leading to NF-kB inhibition after PI treatment (19). Taken together, these results indicate that drug-induced stress may broadly lead to phosphorylation of splicing factors, though precise patterns of phosphorylation may differ in a drug-specific manner.

SRSF splicing factors appear highly phosphorylated at baseline in MM cells

To investigate our phosphoproteomic results via an orthogonal method, we performed Western blots to evaluate for phosphorylation-induced gel mobility shift after Cfz treatment of SRSF1, SRSF3, and SRSF6. After Cfz treatment and isolation of the cytoplasm, we initially saw no discernable shift of these proteins. However, treatment of lysate with calf alkaline phosphatase resulted in a substantial shift of SRSF proteins but not actin (**Fig. S3F**). Therefore, these SRSF factors exist in a highly phosphorylated state even at baseline in MM plasma cells. Upregulated phosphorylation post-Cfz identified by mass spectrometry may therefore represent additional phosphorylation at only selected phosphosites. While these changes in phosphorylation may still result in biological effects, Cfz-induced modulation does not appear to reflect a dramatic shift in the overall phosphorylation status of these SRSF proteins in this system.

To further investigate baseline phosphorylation status of SRSF proteins, we treated MM.1S cells with 50 µM KH-CB19 (20), a reported highly selective inhibitor of the SRSF kinases CLK1 and CLK4 (K_D = 20 nM vs. CLK1). We did not observe any viability effects in MM.1S even at this high concentration (**Fig. S3A**). Unbiased phosphoproteomics after 24 hr of KH-CB19 treatment surprisingly showed no significant change in phosphorylation status of any quantified SRSF phosphosites, except one upregulated (**Fig. S2F**). These results suggest that other kinases also play a role in maintaining SRSF phosphorylation in this system, either at baseline or via feedback mechanisms after sustained CLK1 inhibition.

Proteasome inhibition induces intron retention in MM cells

Given our results demonstrating splicing factor phosphorylation, we next investigated whether pre-mRNA splicing itself was altered after drug treatment. We obtained paired-end sequencing data from polyA-enriched RNA on the same samples used for phosophoproteomics, plus one additional biological replicate (*n* = 3 total) of each of the following: MM.1S treated with 18 nM Cfz, with 10 µM

melphalan, and with DMSO as control; and AMO-1 treated with 15 nM Cfz and with DMSO as control. We used JuncBASE (21) to process the aligned sequencing data by identifying and quantifying both annotated and novel splice junctions. Data for each alternative splicing event was evaluated using the standard measure of "percent spliced in" or PSI (ψ) (**Fig. 3A**).

Comparative analysis of differential PSI (ΔPSI) between 18 nM Cfz- and DMSO-treated MM.1S were considered according to categories including alternative exon acceptor (3' splice site selection), alt. donor (5' splice site selection), alt. last exon, alt. first exon, alt. exon cassette, and intron retention (IR) (**Fig. 3B, Supplementary Table S3**). The ΔPSI distribution for IR demonstrated the greatest positive shift after Cfz treatment (n = 25,807 total IR events measured; median = 2.54). ΔPSI medians for alternative splice site selection also demonstrated a significant shift (alt. donor median = 1.64, n = 9,188 and alt. acceptor median = 0.81, n = 9,352). All other categories were closer to a median of zero (p < 2.2E-16 for median of IR distribution, alt. donor, and alt. acceptor vs. median of cassette by Mann-Whitney test, **Supplementary Table S3**). Intriguingly, PIs are well known to induce a strong heat shock response (12) and prior work in non-cancer cells demonstrated that heat shock alone could impair splicing and induce IR without broadly affecting other alternative splicing events (22, 23). In general, intron-retained transcripts may be subject to nonsense-mediated decay or retained in the nucleus where they remain untranslated. Our results suggest that a similar splicing impairment may be present in MM cells exposed to PI.

We then considered the possibility that the IR phenotype results from a global dysfunction of the splicing machinery during drug-induced apoptosis, which is likely occurring with ~85% cell death at our high-dose Cfz treatment. Our prior proteomic data indicated that SF3B1 and U2AF2, core components of the splicing machinery, are some of the earliest substrates cleaved by caspases during PI-induced apoptosis (9). Indeed, by Western blotting we validated that SF3B1 and U2AF2 are proteolytically cleaved after Cfz treatment and this cleavage can be blocked by the pan-caspase inhibitor zVAD-fmk (**Fig. S3E**).

These caspase cleavage events, then, could be responsible for the IR

phenomenon.

However, we found a similar shift in IR distribution in AMO-1 cells treated

with 15 nM Cfz (n = 27,386; median = 2.2) (**Fig. 3C**) despite much less cytoxicity

(~20%) than the 18 nM Cfz treatment in MM.1S. As caspase cleavage correlates

with degree of cell death, it therefore appears unlikely that cytotoxicity alone is

responsible for IR. Notably, an even smaller shift was observed in IR for MM.1S

with 10 µM melphalan treatment, also at ~20% cytotoxicity (n = 24,247; median =

0.44; p < 2.2E-16 for IR distribution MM.1S 18 nM Cfz vs 10 µM melphalan).

Instead, after melphalan the greatest ΔPSI shift occurred with alt. exon cassettes

(single cassette median = 1.06, n = 12,267, coordinated cassette median = 1.75,

n = 1,417, **Fig. 3D**). Even if we only consider statistically significant IR events (p

< 0.05), the ΔPSI distributions for the drug responses remain distinct (**Fig. S4A-**

B). **Fig. 3E** compares the GO enrichment of all significant ASEs induced by Cfz

and by melphalan and shows divergent classes of genes alternatively spliced.

Therefore, while melphalan also affects alternative splicing it appears to do so via

a different mechanism than Cfz (24).

Intriguingly, in the case of AMO-1 treated with 15 nM Cfz, we noticed the

ΔPSI shift for alternative splice sites (alt. acceptor median = 0.40, n = 9,620; alt.

donor median = 0.61, n = 8,116, **Fig. 3C**, **Supplementary Table S3**) were

decreased compared to MM.1S treated with 18 nM Cfz. Notably, this finding also

correlates with the lesser degree of splicing factor phosphorylation (**Fig. 2**).

These findings illustrate a more pronounced IR phenotype after PI than DNA

damage, irrespective of the amount of cell death, while alternative exonic splice

site determination may be a dose-response behavior.

Exogenous expression of SRSF1 wildtype and RS-domain mutants do not

significantly alter splicing patterns

Having shown that proteasome inhibition can lead to both robust splicing

factor phosphorylation as well as widespread IR of pre-mRNA, we next

considered whether these processes are causally linked or whether they instead

occur via parallel mechanisms. To initially investigate this question, we considered SRSF1 (also known as SF2 or ASF), a well-characterized member of the SR family of splicing factors and a putative proto-oncogene (16, 25). All members of this family contain RNA recognition motifs (RRM) and arginine- and serine-rich domains (RS) (16). In general, SR proteins recognize *cis*-acting splice enhancers on pre-mRNA and work to promote splicing by initially recruiting the spliceosome to these intron-exon junctions (16). We found that SRSF1 demonstrates upregulated phosphorylation at sites in both the RS1 and RS2 domain when MM cells are treated with Cfz (**Fig. 2A, Supplementary Table S4**). The current model of SRSF1 function suggests that 1) SR-protein kinases (SRPK)-mediated phosphorylation of RS domain leads to translocation into the nucleus, 2) further hyperphosphorylation by CLK1 causes association with the U1 spliceosome, and 3) partial dephosphorylation is required for splicing catalysis (16, 26, 27).

To study the effects of SRSF1 phosphorylation in MM, we exogenously expressed a wildtype (SRSF1-WT), phosphomimetic (SRSF1-SD), or phosphodead (SRSF1-SA) variant in AMO-1 plasma cells. We assumed an all-or-none model of SR protein phosphorylation, where exogenous SRSF1 mutants have all 20 serines in the RS1 and RS2 domains replaced with either an aspartate (SD) or an alanine (SA). Exogenously expressed SRSF1 proteins are tagged with a C-terminal mCherry, nuclear localization signal (NLS) and 3x FLAG-peptide (**Fig. 4A**). It is known that phosphorylation of the RS1 domain is necessary for nuclear localization (28, 29); the attempted forced nuclear localization of the SA mutant was chosen to probe potential splicing-level effects of phospho-dead SRSF1 interacting with the spliceosome. Immunoblot confirms expression of exogenous SRSF1 constructs, which migrate higher than endogenous SRSF1 (**Fig. S4E**), and demonstrates lower expression than the high-abundance endogenous protein. Epi-fluorescent images in **Fig. 4B** show the distribution of exogenous SRSF1-WT, SD, and SA mutants. Notably, most of WT and SD signal is localized to the nucleus, suggestive of functional protein product and consistent with expected biology. However, a much larger fraction of SA

mutant is trapped in the cytosol despite NLS tagging. Consistent with prior work (30, 31), this finding suggests that phosphorylation of RS domains is a major requirement for entry into the nucleus.

Upon JuncBASE analysis of poly-A RNA-seq data from DMSO-treated WT, SD, and SA construct (n = 3 for each), we saw remarkably few global differences in PSI as a function of modeled SRSF1 phosphorylation status (**Fig. 4D**). Notably, our results in **Fig. 1B** suggested that phosphorylation of multiple splicing factors, including other SRSF proteins, occurs simultaneously under Cfz-induced stress; we find that altered phosphorylation of SRSF1 alone may not carry any significant effects.

SRSF1 RS-domain phosphomimetic mutant demonstrates weakened interaction with the spliceosome

Though we cannot draw a direct link between SRSF1 phosphorylation status and specific alternative splicing events, we further investigated the diverse biological roles of SRSF1. In addition to modulating pre-mRNA splicing, these include regulating nuclear export of spliced mRNAs and translational regulation in the cytosol via interaction with the ribosome (32-34). Using the 3x-FLAG tag on constructs we performed affinity purification mass spectrometry (AP-MS) with label-free quantitative proteomics vs. an mCherry-NLS-[FLAG]$_3$ control. We specifically evaluated differential binding partners of SRSF1 as a function of phosphorylation status across both the nuclear and cytoplasmic compartments.

While clear differences were observed between the nuclear and cytosolic interactome for each construct, overall biological signatures based on GO analysis were surprisingly similar across WT, SD, and SA within each compartment (**Fig. S5B, D, F**). Notably, in the cytosol we found consistent interactions between both SRSF1-WT and SRSF1-SD with several RNA-binding proteins as well as components of the translational machinery. We do note one stark difference between WT and the phosphomimetic mutant in the nuclear fraction: the WT construct showed direct evidence of interaction with several small nuclear ribonucleoprotein (snRNP) polypeptides, core components of the

U1-U2 spliceosome (**Fig. 4E**). Unexpectedly, these nuclear interactions were not

enriched in the SD construct, which instead interacted with other splicing-related

factors such as TRA2A, TRA2B, and PABPN (**Fig. 4F**). This interactome

mapping may help refine the current model of SRSF1 biology, which suggests

that hyperphosphorylation of RS domains leads to preferential integration with

the U1 spliceosome (35, 36) and would explain the lack of change seen in global

alternative splicing in the SD expressing cells.

Proteasome inhibition of MM cells results in both stochastic intron

retention and specific alternative exon usage

We next explored the splicing-level effects of 15 nM Cfz treatment on

AMO-1 cells expressing the WT, SD, and SA constructs. Notably, in this setting

cytotoxicity at 24 hr was <10% in 8 of 9 total replicates (**Fig. S4D**). Compared to

DMSO-treated samples (**Fig. 4C**), Cfz again elicited a response consistent with

that found in **Fig. 3C**: despite minimal cell death, we observed a clear shift in the

median ΔPSI toward increased global IR (n = 12,139; median = 2.45, p < 2.2E-16

for one-sample Wilcoxon summed rank test). These findings in the absence of

apoptosis underscore that caspase cleavage of splicing factors is unlikely to be a

primary mechanism of IR after PI.

The combined RNA-seq dataset of all Cfz-treated samples vs. DMSO for

these additional SRSF1 constructs were analyzed together (**Fig. 5A**) with

JuncBASE (n = 24 replicates total across all AMO-1, including data in **Fig. 3C,**

4C). With this increased statistical power, we were able to identify *CNNM3*, which

encodes a divalent metal cation transporter, as showing among the strongest

signatures of IR across all events (FDR-corrected p = 0.032) (**Fig. S4G**).

However, despite detecting n = 22,559 IR events by JuncBASE (**Fig. 5B, left**),

very few individual transcripts (n = 43, including *CNNM3*) showed statistically

significant (FDR-corrected p < 0.05) IR across replicates (**Fig. 5B, right**). This

finding suggests that Cfz-induced IR may be a stochastic process, perhaps

resulting from general interference with the splicing machinery without a coherent

selection for specific transcripts.

In contrast, alternative exon splice site usage (alternative exon donor (n = 1134) and alternative exon acceptor (n = 810)) emerged as the dominant type of alternative splicing when considering only statistically significant events (**Fig. 5B**). We investigated whether these consistently observed alternative splicing events may carry some biological relevance. Interestingly, GO enrichment analysis of all the genes undergoing significant alternative splicing after Cfz (n = 2,575 events total across all categories in **Fig. 5B, right**) revealed 'proteasome-mediated ubiquitin-dependent protein catabolic process' (p = 2.08E-16) and 'protein polyubiquitination' (p = 1.39E-13) as highly enriched (**Fig. 5C**). Notably, multiple proteasome subunits (*PSMA3/5/7, PSMB4/5, PSMC1/4/5, PSMD1-4, PSME2*), the protein homeostasis node p97 (*VCP*), and ubiquitin (*UBB, UBC*) all undergo some degree of alternative splicing with Cfz (example in **Fig. 5D**). These findings raise the possibility that alternative splicing may modulate the protein homeostasis machinery in response to therapeutic proteasome inhibition.

       Taken together, our results offer a model for the effects of proteasome inhibition on the splicing machinery in myeloma (**Fig. 5E**). Upon therapeutic insult, the stress response induces phosphorylation of multiple splicing factors. Though the effect of this phosphorylation on specific splicing events remains unclear, these events may relate to specific alterations in exon usage based on known SRSF biochemistry. Our analysis of specific exon usage suggests that modification of the proteasome itself via alternative splicing may play a role in adaptation or resistance to proteasome inhibitor. In parallel, we observe a broad increase in the number of stochastically distributed IR events. These IR events, expected to reduce the number of functional protein products, may work to generally reduce proteotoxic stress and conserve cellular resources normally devoted to protein synthesis, thereby playing a role in adaptation to proteasome inhibition. Alternatively, the intron retention phenotype may indicate malfunction of the spliceosome, an essential process whose loss reduces tumor cell fitness. Interference with splicing may therefore be a previously unappreciated part of the PI mechanism of action.

The spliceosome inhibitor E7107 is broadly potent versus MM cells and synergistic with proteasome inhibitor

Extending from this potential new mechanism of action of PIs as interfering with splicing, we further investigated the therapeutic potential of more dramatic spliceosome disruption in myeloma. For our preclinical studies we employed the tool compound E7107, a pladienolide B analog and direct inhibitor of the core U2 catalytic spliceosome component SF3B1 (15). This molecule has recently been described to induce extreme IR and strong cytotoxic effects versus models of myeloid malignancy, particularly those carrying mutations within splicing factors (37).

Using both qPCR validation of canonical IR events after SF3B1 inhibition (38) (**Fig. S6B**) as well as JuncBASE analysis of RNA-seq data (**Fig. 6A**), as expected we identified very significant IR after 6h of 10 nM E7107 treatment in MM.1S cells (ΔPSI median = 13.79, n = 30,666). There was no noted cytotoxicity at this early time point (**Fig. S6A**). This finding supports the previous conclusion that splicing impairment, not apoptosis, induces IR. However, unlike the PI response, we also observed massive global loss of cassette exon splicing under E7107 (ΔPSI median = -16.6, n = 24,053, **Fig. S6D**). Furthermore, the number of significant ($p < 0.05$) IR events remained very high with E7107 (n = 7,171), unlike the apparently stochastic IR events seen with PI (**Fig. S6C**). Altogether, this suggests that PI-induced impairment of splicing is a partial interference of normal splicing operations, unlike the total abrogation of splicing seen with E7107. Underscoring the potential of splicing inhibition as a therapeutic strategy in MM, E7107 was extremely potent versus a panel of seven MM cell lines treated for 48 hr, with LC_{50}'s ranging from <1 nM to 30 nM (**Fig. 6B**). In addition, a PI-resistant AMO-1 cell line (39) showed very similar sensitivity to E7107 as the parental line (**Fig. 6C**). This finding suggests the potential for clinical utility of splicing inhibition even in PI-refractory disease.

We noted that our MM cell line sensitivities appeared essentially bimodal, with one group of more sensitive lines with LC_{50}'s of <1 nM and another slightly less sensitive group of cell lines with LC_{50} of 20-50 nM. In an attempt to identify

potential determinants of this differential drug sensitivity, we examined publicly

available RNA-seq data of baseline gene expression in MM cell lines

(www.keatslab.org). We were intrigued to find that the more sensitive lines

demonstrated significantly higher RNA expression of *SF3B1* (**Fig. 6D**). This

outcome hinted that more sensitive cell lines may somehow be more "addicted"

to SF3B1, leaving them more vulnerable to splicing inhibition, as well as

revealing a potential biomarker that could be used for patient stratification.

Unfortunately, this result was not confirmed at the protein level (**Fig. 6E**),

suggesting that SF3B1 may undergo post-transcriptional regulation. We found no

other candidates for markers of sensitivity or resistance to E7107 based on

available DNA or RNA sequencing data from this limited cohort of cell lines.

We further explored the hypothesis that interfering with splicing via two

different mechanisms may lead to synergistic MM cell death. Indeed,

combination studies with Cfz and E7107 showed strong synergy across the

dosing landscape based on ZIP synergy scoring (40) (**Fig. 6F-G**). In contrast,

melphalan, which induced much less IR than PI (**Fig. 3B-C**), showed much

weaker synergy in combination with E7107 (**Fig. 6H-I**). These findings support

the approach of using splicing inhibitors in combination with PIs in MM treatment.

Also, this result strengthens the hypothesis that splicing interference is a part of

the PI mechanism of action.

E7107 is a highly potent versus myeloma both *in vivo* and *ex vivo*

Based on this encouraging *in vitro* data, we then moved into a standard *in*

vivo MM model of luciferase-labeled MM.1S cells implanted intravenously into

NOD *scid* gamma (NSG) immunocompromised mice. These cells home to the

murine bone marrow, partially recapitulating the tumor microenvironment in

human disease (41). We found that E7107 was generally well-tolerated with no

appreciable weight loss (**Fig. S7A**). At 3 mg/kg E7107 I.V., a relatively low dose

compared to prior studies in other malignancies (37), we still found pronounced

anti-MM effect after a brief 2 week treatment (**Fig. 7A-C**). This suppression of

tumor translated into a significant survival benefit (p = 0.01, log-ranked test; n = 6

per arm).

We next turned to *ex vivo* evaluation versus primary patient samples.

Fresh bone marrow mononuclear cells from seven PI-refractory MM patients

were treated for 48 hr with varying doses of E7107. Based on flow cytometry

analysis of CD138+ plasma cells (**Fig. S7B**), we found similar high sensitivity of

patient tumor cells to E7107 as found in cell lines, with estimated LC_{50}'s in the

low-nM range (**Fig. 7D**). Notably, non-plasma cell bone marrow mononuclear

cells (CD138- fraction) showed remarkably little cytotoxicity at these same doses,

supporting a potential therapeutic index for splicing inhibitors in MM.

Functional genomics and whole exome sequencing suggests clinical

applications of splicing inhibition in MM

Analysis of CRISPR essentiality screen data in the Cancer Dependency

Map (www.depmap.org; Avana library public 18Q4 (42)), across over 400 cancer

cell lines, demonstrated that myeloma has among the strongest genetic

dependencies on the target of E7107, *SF3B1* (**Fig. S7C**). This genetic ablation

data further supports the ability to pharmacologically eliminate MM tumor cells

via splicing inhibition while sparing normal cells. We further extended this

analysis to other core components of the U1-U2 spliceosome found to be

"common essential" genes per DepMap (43). By aggregating DepMap rankings,

we found that MM cell lines are the most sensitive tumor cell type to genetic

ablation of these central snRNP protein components, necessary for association

with pre-mRNA and splicing catalysis (**Fig. 7E**). Compared to the essential

subunits of the 20S proteasome (including the direct PI target *PSMB5*) (**Fig.**

S7D), we surprisingly found more favorable genetic evidence for targeting the

spliceosome in MM than the proteasome.

Furthermore, a recent study validated the sulfonamide indisulam as an

inhibitor of splicing via targeted degradation of RBM39, another component of the

spliceosome with high homology to U2AF2 (44). In this work, hematopoeitic

malignancy cell lines were broadly more sensitive to indisulam than solid tumor

cell lines. We confirmed cytotoxicity of indisulam versus a panel of MM cell lines

(**Fig. S7E**), though LC_{50}'s (0.3 - >20 µM) were much higher than those for the

SF3B1 inhibitor E7107. In DepMap data, MM was again among the more

sensitive tumor type to RBM39 ablation (**Fig. S7F**). Indisulam may therefore

represent another approach to targeting the spliceosome in this disease, though

given lower potency the potential for clinical translation is less clear.

We next took advantage of genomic and transcriptomic data from isolated

malignant plasma cells from newly-diagnosed MM patients in the Multiple

Myeloma Research Foundation CoMMpass study (research.themmrf.org; version

IA11). First evaluating gene expression data, we intriguingly found significantly

decreased progression-free survival among patients in the top quartile of *SRSF1*

expression versus those in the bottom quartile (*p* = 0.0081 by log-ranked test)

and a trend toward similarly decreased overall survival for patients in the top vs.

bottom quartile of *SF3B1* expression (*p* = 0.087) (**Fig. S7G**). These results raise

the possibility of poorer outcomes in patients whose disease is more dependent

on the spliceosome.

However, we note that both E7107 (37) and the recently described

splicing inhibitor H3B-8800 (38) have both been shown to have the greatest

potency versus hematopoietic malignancies carrying mutations in splicing factors

such as *SF3B1*, *SRSF2*, *U2AF2*, and *ZRSR2* (45). These mutations are seen

frequently in myelodysplastic syndromes (MDS), acute myeloid leukemia, and

chronic lymphocytic leukemia, appearing in up to 50% of MDS patients (45). We

therefore examined exome sequencing data available in CoMMpass and found

that 28.0% of MM patients (268 of 956) were found to carry missense mutations

within at least one of 119 splicing-associated factors recently proposed to be

most relevant to tumorigenesis across a survey of the The Cancer Genome Atlas

(**Fig. 7F, Supplementary Table S6**) (46). While only a small minority of these

identified mutations has been functionally validated to affect splicing, the most

common single mutation was at the known "hotspot" *SF3B1* K666T, found in

three patients. Variant allele frequencies for these expected heterozygous

mutations were 42%, 35%, and 22%, suggestive of a prominent subclonal

fraction of the tumor cell population. Among well-characterized genes, mutations were found in *SF3B1* (*n* = 10 patients, including K666T mutations), *SRSF2* (*n* = 2), *U2AF1* (*n* = 4), and *ZRSR2* (*n* = 1). Unfortunately we were unable to obtain rare primary patient samples containing mutations in these genes, and no myeloma cell lines are known to carry hotspot mutations in these well-characterized splicing factors (www.keatslab.org). While our data suggest that spliceosome inhibition should be considered a therapeutic option for MM patients of any genotype, recent work in other malignancies (37, 38) supports the potential for particular benefit in the subset of patients carrying pathogenic splicing factor mutations.

DISCUSSION:

Our results demonstrate that PI therapy in myeloma leads to both specific alterations in splice site usage and broad-scale interference with spliceosome function. This observation, initially generated through unbiased phosphoproteomics, led us to explore the spliceosome itself as a MM vulnerability. Our preclinical evaluation and analysis of functional genomics and exome sequencing data further reinforced the spliceosome as a therapeutic target in MM.

These results raise a number of intriguing questions. From a mechanistic perspective, prior work examining SR phosphorylation after cellular perturbation using Western blotting did not consistently show a broad hyperphosphorylation signature (47-49). Our results therefore illustrate the utility of unbiased phosphoproteomics to elucidate cancer drug response. Recent work suggests that additional kinases beyond the well-characterized SRPKs and CLKs may be involved in SR phosphorylation (50, 51). However, in the context of drug-induced stress in cancer, the mechanism that leads to coordinated, upregulated phosphorylation across multiple splicing factors, whether via kinase activation or phosphatase inhibition, will be an important topic for future investigation.

We also found a correlation between PI-induced stress and both SR factor phosphorylation and the degree of alternative exon selection. This finding

appears consistent with prior studies (16) suggesting that SR factors play a role

in exon selection and their activity can be modulated by phosphorylation. Why

these effects specifically lead to alternative splicing of protein homeostasis genes

remains to be investigated. In contrast, we found IR to be largely independent of

the degree of PI-induced stress. This result suggests that the IR phenotype is

mediated by a different mechanism and uncoupled from SR factor

phosphorylation.

In attempting to model the relationship between SR phosphorylation and

splicing, we recognize our phosphomimetic construct likely does not fully

recapitulate the complex phosphorylation biology of SRSF1 within cells (27), nor

does it fully match the level of expression of endogenous SRSF1. In general,

causal links have been noted between SRSF1 phosphorylation and splicing in

single transcript, *in vitro* systems (34, 52, 53), but isolating global effects of SR

phosphorylation on splicing within cells have remained elusive. Furthermore,

using genetic approaches we cannot readily model phosphorylation changes on

multiple SR proteins simultaneously, which may be necessary to elicit broader

phenotypic effects. Despite these limitations, however, our WT expression

studies provide a landscape of the SRSF1 cytosolic and nuclear interactome,

which may inform future studies of SR protein biology in myeloma and other

systems.

This PI-induced interference with normal splicing even at minimal

cytotoxicity, much greater than that found with melphalan, may relate to the

activation of the heat shock response. We found prominent heat shock

chaperone induction even under a non-cytotoxic dose of the PI bortezomib in

MM.1S cells (10). As previously shown in non-cancer cells, heat shock alone,

without cell death, can lead to significant intron retention (23). One hypothesis is

that this broad-scale inhibition of splicing acts in a similar fashion to translational

inhibition after drug-induced stress: a way to conserve cellular resources and

focus on only producing genes required for survival and the stress response.

However, as described in our model of **Fig. 5E**, and evidenced by our mRNA-seq

data after E7107 treatment (**Fig. 6A**), another possible result of widespread

intron retention and downstream loss of normal protein production is significant

decrease in cellular fitness and ultimately, cell death. There may be a

quantitative threshold effect between these two outcomes that remains to be

elucidated.

Here, we propose that the loss-of-fitness modality of drug-induced IR

constitutes a previously unexplored mechanism of action of PIs. We further

performed a preclinical evaluation of splicing inhibition in myeloma using E7107,

finding potent anti-myeloma effects *in vitro*, *in vivo*, and *ex vivo* versus primary

patient samples. From a therapeutic perspective, one of the major questions is

the potential toxicity of targeting such an essential process as the catalytic

spliceosome. However, our analysis of genetic dependency data and our *ex vivo*

data with E7107 clearly demonstrates the potential to target core spliceosome

subunits in MM while largely sparing normal cells. In fact, based on this analysis

the spliceosome appears to be an even more promising target than the clinically-

validated approach of targeting essential subunits of the proteasome.

Furthermore, presumed efficacious doses (based on measured blood

concentrations in the nM range) of E7107 were largely well tolerated in a Phase I

clinical trial (54). While this molecule is no longer in clinical development, it is

thought that E7107 visual toxicity was molecule-specific and is not a function of

targeting the spliceosome in general (15). Our genomic analysis suggests that

mutations in splicing factors are found in a substantial fraction of MM patients.

Newer generations of splicing inhibitors are currently in clinical trials for other

hematologic malignancies (38) (NCT02841540) and may be of particular benefit

for these patients. Our results support clinical investigation of these compounds

in MM either alone or to enhance PI efficacy as combination therapy.

METHODS

Cell culture

All cell lines were grown in suspension at 37°C, 5% CO_2 in complete media:

RPMI 1640 medium (Gibco, 22400105, UCSF CCFAE002), supplemented with

10% FBS (Atlanta Biologicals, S11150) for proteomics experiments and

Benchmark FBS (Gemini Bio-products, 100-106) for drug viability experiments

and 1% penicillin-streptomycin (UCSF, CCFGK003). INA6 cell media was

supplemented with 90 ng/mL recombinant human IL-6 (ProSpec Bio, CYT-213).

Drug cytotoxicity assay

For dose-response cell toxicity assays, 1E+3 myeloma cells were seeded per

well in 384 well plates (Corning) using the Multidrop Combi (Thermo Fisher) and

incubated for 24 hr. In monotherapy cytotoxicity assays, cells were treated with

drug or DMSO and incubated for 48 hr, while cells were further incubated with

E7107 (H3) for an additional 24 hr in E7107 dual therapy combination assays.

Carfilzomib (SelleckChem, S2853-50mg), melphalan (Sigma, S2853-50mg), and

E7107 (H3 Biomedicine, CAS:630100-90-2), and KH-CB19 (sc-362756) were

solubilized in DMSO at 10 mM.

All cell viability was determined with Cell-Titer Glo reagent (Promega, G7573)

using a Glomax Explorer (Promega) luminescence plate reader. For the drug

titration cytotoxicity assays, measurements were performed in quadruplicate,

while measurements were performed in triplicate in all other assays, and

viabilities are reported as mean (+/- S.D.) ratio normalized to DMSO-treated

controls or measurements at 0 hr. For ZIP synergy calculations, normalized

viability data was submitted to SynergyFinder web application (40).

Drug dosing for proteomics and RNA-seq experiments

Proteomic/phosphoproteomic/RNA-seq experiments were performed at a cell

density of 1E6 cells/mL. For timecourse studies, ~20E6 cells were grown in

complete media for each timepoint (0, 8, 16, and 24 hr), whereas for single-

timepoint experiments, 15-20E+6 cells in light SILAC media were treated with

drug compound and cells in heavy SILAC media (L-Lysine-$^{13}C_6,^{15}N_2$, L-Arginine-

$^{13}C_6,^{15}N_4$ (Cambridge Isotope, CNLM-291-H-1, CNLM-539-H-1) were treated with

DMSO for 24 hr. 1-3E+6 cells were set aside for RNA-seq. Cells were washed in

PBS and cell pellets were frozen in liquid nitrogen (LN2) and stored in -80°C. 1

biological replicate for the timecourse experiment, 2 biological replicates for each

single-timepoint condition (with a third only for RNA-seq), and 3 biological

replicates for all AP-MS were gathered and analyzed.

Cloning and lentiviral transduction

SRSF1 and mCherry genes, along with 3X FLAG sequences and nuclear

localization signal (NLS) were cloned into pLV-416G second generation lentiviral

plasmid (UCSF HMTB) by Gibson Assembly. SRSF1 constructs were transfected

into Lenti-X 293T(Takara Bioscience, 632180) packaging cells with Gag-Pol

expressing pCMV-dR8.91 (Addgene, Plasmid#2221) and VSV-G envelope

expressing pMD2.G (Addgene, Plasmid#12259) plasmids. Viral particles were

harvested, concentrated with Lenti-X concentrator (Takara Bioscience, 631231)

and viral titers were incubated with AMO-1 cells. Positively transduced cells were

selected with selection drug, G418 (VWR, 970-3-058), for several passages, then

by mCherry expression with Fluorescence Activated Cell Sorting (FACS, Sony

SH800). Protocol details are found in Supplementary Information.

Phosphoproteomic peptide preparation

Frozen pellets of ~15-20E6 cells were lysed in 8 M urea, 0.1 M Tris pH 8.0, 150

mM NaCl and 1X HALT phosphatase/protease inhibitor cocktail (Pierce, 78442)

for timecourse experiments or 8 M Guanadine-Cl (Gdn, Chem Impex Intl., 00152-

1KG), 0.1 M Tris pH 8.5, 10 mM tris(2-carboxyethyl)phosphine (TCEP, Pierce,

20491), 40 mM 2-chloroacetamide (2-CAA, Sigma, 22790-250G-F), 1X HALT for

SILAC samples and lysed with probe sonicator (BRANSONIC). In the case of

single-timepoint SILAC samples, equal part light and heavy labeled lysate

samples were combined (~ 2.5–3 mg total). Lysate is diluted with 0.1 M Tris pH

8.0 to a final concentration of 1.3 M Gdn or urea. Proteome is digested with

1:100 dilution of trypsin overnight for 22-24 hr at room temperature. Peptides are

extracted with SEP-PAK C18 cartridges (WATERS). For single-timepoint SILAC

samples, ~100 µg of eluted peptides were dried and analyzed separately by LC-

MS/MS as unenriched "global proteomics." Remainder of eluate was diluted 3-4

fold with water, lyophilized, then resuspended in 80% ACN, 0.1% TFA and

enriched on $FeCl_3$ charged NTA-agarose beads sitting atop a C18 matrix in a

stage-tip platform (Nest). Eluted phosphopeptides are dried and stored at -80°C.

Affinity Purification

For each replicate, frozen cell pellets were gently lysed on ice with 200 µl

hypotonic lysis buffer (20 mM Tris (pH 7.4@4°C), 10 mM KCl, 0.1 mM EDTA

(Fisher, BP120-500), 0.5% NP-40 alternative (EMD, 492016-100ML), 1 mM DTT

(Gold Biotech, DTT50), 1 mM PMSF (RPI, P20270-1.0), 1x HALT

protease/phosphatase inhibitor cocktail (Pierce, 78442), 300mM Sucrose, 0.03

U/mL aprotinin (RPI, A20550-0.001)), underwent 3 X freeze-thaw cycles, and

clarified with 5 passes through an 18-gauge syringe needle. Lysate was

centrifuged at 5,000 rcf, 4°C for 10 min and supernatant was reserved as

cytoplasmic fraction, while nuclear fraction was washed and resuspended in 60

µl of 20 mM HEPES (pH 7.9), 420 mM NaCl, 25% glycerol, 1 mM EDTA, 1 mM

DTT, 1 mM PMSF, 0.03 U/mL aprotinin, 1x protease/phosphatase inhibitor

cocktail (HALT), 25 U Benzonase/mL and clarified with 10 passes through 18-

gauge syringe needle. Both fractions were adjusted to 50 mM Tris pH 7.4, 150

mM NaCl, 1mM EDTA (binding buffer) and combined with M2 anti-FLAG

magnetic beads (Sigma, M8823). Bound lysate was washed with binding buffer +

0.05% NP-40, then binding buffer, then twice with 20 mM Tris pH 8.0, 2 mM

$CaCl_2$. Proteins are denatured and cystines are reduced and alkylated with 6 M

Gdn, 40 mM 2-CAA, 5 mM TCEP, 100 mM Tris pH 8.0, then trypsinized on-bead

with ~0.75 µg trypsin/ sample, ~ 20h at 37°C, and peptides were desalted with

homemade C18 stagetips and dried and stored at -80°C.

LC-MS/MS

~1 µg peptides were analyzed for each sample by "shotgun-" LC-MS/MS on a

Dionex Ultimate 3000 RSLCnano with 15 cm Acclaim PEPMAP C18 (Thermo,

164534) reverse phase column and Thermo Q-Exactive plus mass spectometer.

Samples were analyzed with either a 3h 15 min non-linear gradient or a 1h 23

min linear gradient from 2.4% acetonitrile (ACN, Sigma, 34998-4L), 0.1% FA to 32% ACN. Experiment specific LC-MS/MS settings are listed in Supplementary Information.

Proteomic data analysis and quantification

Initial timecourse unlabeled phosphoproteomics data were processed together on Maxquant v1.5.1.2 (55) and searched against the human proteome (Uniprot downloaded 2014/12/3, with 89,706 entries). All AP-MS samples were processed together with similar settings. All SILAC samples (phospho- and unenriched peptides) were processed together with similar settings. SILAC quantification for global proteomics at the protein level requires 1 minimum razor or unique peptide. A one-sample T-test was applied to the log-2 transform of the normalized SILAC-labeled peptide ratios (heavy:light) for single-timepoint analysis, while for AP-MS data, two-sample T-test was applied to the log-2 transform of the median-normalized MaxQuant label-free quantification (LFQ) values of protein groups. The number of total entries (phosphosites, protein groups, significance is $p < 0.05$, |t-test difference| ≥ 1), along with correlation statistics between replicates, are summarized in **Supplementary Table S4** and shown in **Fig. S3B**. See Supplementary Information for specific search and analysis settings.

RNA-seq library preparation

RNA was extracted from frozen cell pellets with RNeasy Mini-prep kit (Qiagen, 74104). For timecourse experiments, cDNA library of expression transcripts was carried out with RNA Hyper Prep kit with RiboErase (Kapa, KK8560) to enhance transcript reads above ribosomal reads, while single-timepoint experiments assessing splicing required mRNA enrichment with magnetic mRNA Isolation kit poly-dT beads (NEB), then RNA Hyper Prep kit (Kapa, KK8540) for cDNA construction of 200-400 bp library with Illumina platform TruSeq indexed adaptors (**Supplementary Table S1**). RNA and DNA quantified at all steps by Nanodrop (Thermo Scientific) and cDNA library size and quality were evaluated

on a Bioanalyzer 2100 (Agilent) with High Sensitivity DNA Kit (Agilent, 5067-

4626), before being submitted for next generation sequencing on a HiSeq4000

(Illumina) at the UCSF Center for Advanced Technologies core facility.

JuncBASE alternative splicing analysis

Alternative splicing events were identified and quantified with JuncBASE v1.2-

beta using default parameters (21). Intron-exon junction database was created

from hg19 annotations. T-test was used to compare number of inclusion and

exclusion reads and p-values were adjusted with Benjamini-Hochberg correction.

For ΔPSI histograms in Fig. **3B-C, 4C-D, and 5A**, JuncBASE output included a

subset of alternative splice events with median PSI = 0.00 in both conditions or

median PSI = 100.00 in both conditions, resulting in ΔPSI = 0.00. These events

were manually removed for downstream analyses. Histograms and splicing

statistics were determined with statistical computing program R (v3.5.1) and a

summary is listed in **Supplementary Table S3**.

Gene Ontology enrichment analysis

Gene Ontology (GO) enrichment analysis of upregulated phosphosites and

enriched SRSF1 interactors was performed in STRING (v10.5, https://string-

db.org/) (56), searching against a background of all quantified protein entries.

Enrichment analysis of all significantly alternative spliced genes (raw $p < 0.05$)

was performed using web-based enrichment analysis tool, Enrichr

(http://amp.pharm.mssm.edu/Enrichr/) (57). Reported combined score is

calculated by multiplying the natural log transform of the p-value with the Fisher's

exact test of expected rank deviation (Z-score). Functional GO analysis is limited

to biological processes and compiled in **Supplementary Table S5**.

Xenograft mouse model and *in vivo* luminescence imaging

1E6 MM.1S-luc cells, stably expressing luciferase, were transplanted via tail vein

injection into 12 NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Wjl}*/SzJ (NSG) mice from The Jackson

Laboratory (cat# 005557). All the mice were female, 6-8 wks old at start of

studies, and typically weigh 20-25 g. NSG mice were handled with aseptic

techniques and housed in pathogen free environments at the UCSF Laboratory

Animal Research Center. All mouse studies were performed according to UCSF

Institutional Animal Care and Use Committee-approved protocols. Tumor burden

was assessed through weekly bioluminescent imaging in the UCSF preclinical

therapeutic core on a Xenogen In Vivo Imaging System (IVIS), beginning 13 days

after implantation, which is the same day as treatment initiation. Tumor implanted

humanized mice were randomized and sorted into control arm and treatment

arm, 6 mice/arm. Mice were treated for two weeks (five days on, two days off)

with vehicle or 3 mg/kg E7107, formulated in vehicle (10% Ethanol, 5% Tween-

80, QS with Saline) and administered by continuous subcutaneous infusion. Mice

were kept and observed until survival endpoint; final timepoint was 54 days after

MM.1S transplant. Acquired luciferase intensities were quantified with Living

Image Software (PerkinElmer) in units of radiance (photons/s/cm^2/sr). Kaplan-

meier survival curves along with log-ranked test to determine significance were

calculated in GraphPad Prism 6 software.

Patient bone marrow aspirate, CD138 labeling and flow cytometry analysis

Fresh de-identified primary multiple myeloma patient bone marrow (BM) samples

were obtained from the UCSF Hematologic Malignancies Tissue Bank in

accordance with the UCSF Committee on Human Research-approved protocols

and the Declaration of Helsinki. Bone marrow mononuclear cells were isolated by

density gradient centrifugation with Histopaque-1077 (Sigma Aldrich), and

washed with 10 mL D-PBS 3 times. Mononuclear cells were resuspended in a

small volume (~1.5 mL) of media (RPMI1640, 10% FBS, 1%

penicillin/streptomycin, 2 mM glutamine) and incubated at 37°C, 5% CO_2 for

15min. Isolated mononuclear cells from multiple myeloma patient bone marrow

were adjusted to 2E5 cells/well in a 96 well plate. Cells were stimulated with 50

ng/ml recombinant human IL-6 (ProsPec) for 17 hr before treatment with E7107

or DMSO for 24 hr. Cells were then stained with 10 µL Alexa-Fluor 647 mouse

anti-human CD138 antibody (BD Pharmingen, cat# 562097;

RRID:AB_10895974) or Alexa-Fluor 647 IgG κ isotype (BD Pharmingen, cat#

557714; RRID:AB_396823) control and 2 μL SyTOX Green (Thermo, S34860)

per 1 mL FACS buffer (D-PBS, 5% FBS). Resuspended cells are characterized

with a CytoFLEX fluorescence cytometer (BD).

Statistical Analyses

All data are presented as mean +/- standard deviation, unless otherwise stated.

Statistical significance in proteomics comparisons was determined by Student's t-

test: One-sample t-test with null hypothesis that $\log_2$-transform of the normalized

SILAC ratio = 0, or a two-sample T-test with null hypothesis that the difference in

$\log_2$-transform of the intensities is equal to. A $p < 0.05$ is considered statistically

significant. For all Kaplan-Meier survival analysis, log-ranked test was used to

determine statistical significance.

Data Availability

The mass spectrometry proteomics data and MaxQuant analysis results have

been deposited to the ProteomeXchange Consortium via the PRIDE (58) partner

repository with the dataset identifier PXD012172. Datasets are private during

review; reviewers may access datasets with following credentials:

Username: reviewer92428@ebi.ac.uk

Password: Xxe5zQtD

Raw RNA-seq data, processed analysis files, and JuncBASE results may be

downloaded from the Gene Expression Omnibus, GEO

(https://www.ncbi.nlm.nih.gov/geo/) with the accession number: GSE124510.

Reviewers may access this data during review with the token: ifwxqgsgptcbtgz

Acknowledgements

We thank Dr. Silvia Buonamici at H3 Biomedicine for providing E7107 and

insightful discussions and Jacob Runyan for assisting in quality control of

sequencing data. We thank Drs. Renate Burger and Christoph Driessen for

providing INA-6 and AMO-1 parental and bortezomib resistant cell lines,
respectively. We also thank the lab of James Wells for use of Zeiss Z1 Observer
microscope, Center for Advanced Technology at UCSF for HiSeq sequencing,
Dr. Jane Gordon and the Laboratory for Cell Analysis at Helen Diller Family
Comprehensive Cancer Center for use and assistance of Sony SH800, and Dr.
Danielle Swaney for discussion of AP-MS. This work was supported by NIH/NCI
P30CA083103 (Cancer Center Support Grant, supporting UCSF Preclinical
Therapeutic Core facility managed by B.C.H.), NIH/NHGRI T32HG008345 (to
A.M.T.), the Damon Runyon Cancer Research Foundation Dale Frey
Breakthrough Award (DFS 14-15), NIH/NCI K08CA184116, NIH/NIGMS
DP2OD022552, and the UCSF Stephen and Nancy Grand Multiple Myeloma
Translational Initiative (to A.P.W.) and NIH/NCI R01CA226851 (to A.N.B. and
A.P.W.).

Author Contributions

H.H.H., I.F., C.L., P.B., M.C.M., M.M., and A.P.W. performed experiments and
analyzed experimental data. A.M.T., Y-H.L., and A.B. analyzed transcriptomic
and genomic data. J.M., P.P., and B.C.H. performed *in vivo* studies. T.G.M.,
J.L.W., S.W.W., and N.S. consented patients and obtained primary specimens.
H.H.H. and A.P.W. wrote the manuscript with input from all authors.

Declaration of Interests

The authors declare no relevant conflicts of interest.

REFERENCES

1. Walter P and Ron D. The unfolded protein response: from stress pathway
to homeostatic regulation. *Science* 2011;334(6059):1081-6 doi
10.1126/science.1209038.

2. Obeng EA, Carlson LM, Gutman DM, Harrington WJ, Jr., Lee KP and
Boise LH. Proteasome inhibitors induce a terminal unfolded protein response in

multiple myeloma cells. *Blood* 2006;107(12):4907-16 doi 10.1182/blood-2005-08-3531.

3.   Lee AH, Iwakoshi NN, Anderson KC and Glimcher LH. Proteasome inhibitors disrupt the unfolded protein response in myeloma cells. *Proc Natl Acad Sci USA* 2003;100(17):9946-51 doi 10.1073/pnas.1334037100.

4.   Le Moigne R, Aftab BT, Djakovic S, Dhimolea E, Valle E, Murnane M*, et al.* The p97 inhibitor CB-5083 is a unique disrupter of protein homeostasis in models of Multiple Myeloma. *Mol Cancer Therap* 2017 doi 10.1158/1535-7163.MCT-17-0233.

5.   Gandolfi S, Laubach JP, Hideshima T, Chauhan D, Anderson KC and Richardson PG. The proteasome and proteasome inhibitors in multiple myeloma. *Cancer Metas Rev* 2017;36(4):561-84 doi 10.1007/s10555-017-9707-8.

6.   Suraweera A, Munch C, Hanssum A and Bertolotti A. Failure of amino acid homeostasis causes cell death following proteasome inhibition. *Mol Cell* 2012;48(2):242-53 doi 10.1016/j.molcel.2012.08.003.

7.   Wallington-Beddoe CT, Sobieraj-Teague M, Kuss BJ and Pitson SM. Resistance to proteasome inhibitors and other targeted therapies in myeloma. *Br J Haematol* 2018;182(1):11-28 doi 10.1111/bjh.15210.

8.   Mitra AK, Harding T, Mukherjee UK, Jang JS, Li Y, HongZheng R*, et al.* A gene expression signature distinguishes innate response and resistance to proteasome inhibitors in multiple myeloma. *Blood Cancer J* 2017;7(6):e581 doi 10.1038/bcj.2017.56.

9.   Wiita AP, Ziv E, Wiita PJ, Urisman A, Julien O, Burlingame AL*, et al.* Global cellular response to chemotherapy-induced apoptosis. *eLife* 2013;2:e01236 doi 10.7554/eLife.01236.

10.   Liu TY, Huang HH, Wheeler D, Xu Y, Wells JA, Song YS*, et al.* Time-Resolved Proteomics Extends Ribosome Profiling-Based Measurements of Protein Synthesis Dynamics. *Cell Syst* 2017;4(6):636-44 e9 doi 10.1016/j.cels.2017.05.001.

11. Mitsiades N, Mitsiades CS, Poulaki V, Chauhan D, Fanourakis G, Gu X, *et al.* Molecular sequelae of proteasome inhibition in human multiple myeloma cells. *Proc Natl Acad Sci USA* 2002;99(22):14374-9 doi 10.1073/pnas.202445099.

12. Shah SP, Lonial S and Boise LH. When Cancer Fights Back: Multiple Myeloma, Proteasome Inhibition, and the Heat-Shock Response. *Mol Cancer Res* 2015;13(8):1163-73 doi 10.1158/1541-7786.MCR-15-0135.

13. Li X, Colvin T, Rauch JN, Acosta-Alvear D, Kampmann M, Dunyak B*, et al.* Validation of the Hsp70-Bag3 protein-protein interaction as a potential therapeutic target in cancer. *Mol Cancer Therap* 2015;14(3):642-8 doi 10.1158/1535-7163.MCT-14-0650.

14. Casado P, Hijazi M, Britton D and Cutillas PR. Impact of phosphoproteomics in the translation of kinase-targeted therapies. *Proteomics* 2017;17(6) doi 10.1002/pmic.201600235.

15. Lee SC and Abdel-Wahab O. Therapeutic targeting of splicing in cancer. *Nat Med* 2016;22(9):976-86 doi 10.1038/nm.4165.

16. Long JC and Caceres JF. The SR protein family of splicing factors: master regulators of gene expression. *Biochem J* 2009;417(1):15-27 doi 10.1042/BJ20081501.

17. Casado P, Rodriguez-Prados JC, Cosulich SC, Guichard S, Vanhaesebroeck B, Joel S*, et al.* Kinase-substrate enrichment analysis provides insights into the heterogeneity of signaling pathway activation in leukemia cells. *Sci Signaling* 2013;6(268):rs6 doi 10.1126/scisignal.2003573.

18. Colwill K, Pawson T, Andrews B, Prasad J, Manley JL, Bell JC*, et al.* The Clk/Sty protein kinase phosphorylates SR splicing factors and regulates their intranuclear distribution. *EMBO J* 1996;15(2):265-75.

19. Traenckner EB, Wilk S and Baeuerle PA. A proteasome inhibitor prevents activation of NF-kappa B and stabilizes a newly phosphorylated form of I kappa B-alpha that is still bound to NF-kappa B. *EMBO J* 1994;13(22):5433-41.

20. Fedorov O, Huber K, Eisenreich A, Filippakopoulos P, King O, Bullock AN*, et al.* Specific CLK inhibitors from a novel chemotype for regulation of alternative splicing. *Chem Biol* 2011;18(1):67-76 doi 10.1016/j.chembiol.2010.11.009.

21.   Brooks AN, Yang L, Duff MO, Hansen KD, Park JW, Dudoit S, *et al.* Conservation of an RNA regulatory map between Drosophila and mammals. *Genome Res* 2011;21(2):193-202 doi 10.1101/gr.108662.110.

22.   Yost HJ and Lindquist S. RNA splicing is interrupted by heat shock and is rescued by heat shock protein synthesis. *Cell* 1986;45(2):185-93.

23.   Shalgi R, Hurt JA, Lindquist S and Burge CB. Widespread inhibition of posttranscriptional splicing shapes the cellular transcriptome following heat shock. *Cell Rep* 2014;7(5):1362-70 doi 10.1016/j.celrep.2014.04.044.

24.   Marchesini M, Ogoti Y, Fiorini E, Aktas Samur A, Nezi L, D'Anca M, *et al.* ILF2 Is a Regulator of RNA Splicing and DNA Damage Response in 1q21-Amplified Multiple Myeloma. *Cancer Cell* 2017;32(1):88-100 e6 doi 10.1016/j.ccell.2017.05.011.

25.   Karni R, de Stanchina E, Lowe SW, Sinha R, Mu D and Krainer AR. The gene encoding the splicing factor SF2/ASF is a proto-oncogene. *Nat Struct Mol Biol* 2007;14(3):185-93 doi 10.1038/nsmb1209.

26.   Misteli T, Caceres JF, Clement JQ, Krainer AR, Wilkinson MF and Spector DL. Serine phosphorylation of SR proteins is required for their recruitment to sites of transcription in vivo. *J Cell Biol* 1998;143(2):297-307.

27.   Ghosh G and Adams JA. Phosphorylation mechanism and structure of serine-arginine protein kinases. *FEBS J* 2011;278(4):587-97 doi 10.1111/j.1742-4658.2010.07992.x.

28.   Caceres JF, Screaton GR and Krainer AR. A specific subset of SR proteins shuttles continuously between the nucleus and the cytoplasm. *Genes Dev* 1998;12(1):55-66.

29.   Misteli T, Caceres JF and Spector DL. The dynamics of a pre-mRNA splicing factor in living cells. *Nature* 1997;387(6632):523-7 doi 10.1038/387523a0.

30.   Caceres JF, Misteli T, Screaton GR, Spector DL and Krainer AR. Role of the modular domains of SR proteins in subnuclear localization and alternative splicing specificity. *J Cell Biol* 1997;138(2):225-38.

31.   Aubol BE, Serrano P, Fattet L, Wuthrich K and Adams JA. Molecular

interactions connecting the function of the serine-arginine-rich protein SRSF1 to

protein phosphatase 1. *J Biol Chem* 2018;293(43):16751-60 doi

10.1074/jbc.RA118.004587.

32.   Das R, Yu J, Zhang Z, Gygi MP, Krainer AR, Gygi SP*, et al.* SR proteins

function in coupling RNAP II transcription to pre-mRNA splicing. *Mol Cell*

2007;26(6):867-81 doi 10.1016/j.molcel.2007.05.036.

33.   Huang Y, Gattoni R, Stevenin J and Steitz JA. SR splicing factors serve as

adapter proteins for TAP-dependent mRNA export. *Mol Cell* 2003;11(3):837-43.

34.   Xiao SH and Manley JL. Phosphorylation of the ASF/SF2 RS domain

affects both protein-protein and protein-RNA interactions and is necessary for

splicing. *Genes Dev* 1997;11(3):334-44.

35.   Cho S, Hoang A, Sinha R, Zhong XY, Fu XD, Krainer AR*, et al.* Interaction

between the RNA binding domains of Ser-Arg splicing factor 1 and U1-70K

snRNP protein determines early spliceosome assembly. *Proc Natl Acad Sci USA*

2011;108(20):8233-8 doi 10.1073/pnas.1017700108.

36.   Ngo JC, Chakrabarti S, Ding JH, Velazquez-Dones A, Nolen B, Aubol BE*,*

et al. Interplay between SRPK and Clk/Sty kinases in phosphorylation of the

splicing factor ASF/SF2 is regulated by a docking motif in ASF/SF2. *Molecular*

Cell 2005;20(1):77-89 doi 10.1016/j.molcel.2005.08.025.

37.   Lee SC, Dvinge H, Kim E, Cho H, Micol JB, Chung YR*, et al.* Modulation

of splicing catalysis for therapeutic targeting of leukemia with mutations in genes

encoding spliceosomal proteins. *Nat Med* 2016;22(6):672-8 doi

10.1038/nm.4097.

38.   Seiler M, Yoshimi A, Darman R, Chan B, Keaney G, Thomas M*, et al.*

H3B-8800, an orally available small-molecule splicing modulator, induces

lethality in spliceosome-mutant cancers. *Nat Med* 2018;24(4):497-504 doi

10.1038/nm.4493.

39.   Soriano GP, Besse L, Li N, Kraus M, Besse A, Meeuwenoord N*, et al.*

Proteasome inhibitor-adapted myeloma cells are largely independent from

proteasome activity and show complex proteomic changes, in particular in redox and energy metabolism. *Leukemia* 2016 doi 10.1038/leu.2016.102.

40.   Yadav B, Wennerberg K, Aittokallio T and Tang J. Searching for Drug Synergy in Complex Dose-Response Landscapes Using an Interaction Potency Model. *Comp Struct Biotechnol J* 2015;13:504-13 doi 10.1016/j.csbj.2015.09.001.

41.   McMillin DW, Negri JM and Mitsiades CS. The role of tumour-stromal interactions in modifying drug response: challenges and opportunities. *Nat Rev Drug Discov* 2013;12(3):217-28 doi 10.1038/nrd3870.

42.   Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS*, et al.* Defining a Cancer Dependency Map. *Cell* 2017;170(3):564-76 e16 doi 10.1016/j.cell.2017.06.010.

43.   Wahl MC, Will CL and Luhrmann R. The spliceosome: design principles of a dynamic RNP machine. *Cell* 2009;136(4):701-18 doi 10.1016/j.cell.2009.02.009.

44.   Han T, Goralski M, Gaskill N, Capota E, Kim J, Ting TC*, et al.* Anticancer sulfonamides target splicing by inducing RBM39 degradation via recruitment to DCAF15. *Science* 2017;356(6336) doi 10.1126/science.aal3755.

45.   Saez B, Walter MJ and Graubert TA. Splicing factor gene mutations in hematologic malignancies. *Blood* 2017;129(10):1260-9 doi 10.1182/blood-2016-10-692400.

46.   Seiler M, Peng S, Agrawal AA, Palacino J, Teng T, Zhu P*, et al.* Somatic Mutational Landscape of Splicing Factor Genes and Their Functional Consequences across 33 Cancer Types. *Cell Rep* 2018;23(1):282-96 e4 doi 10.1016/j.celrep.2018.01.088.

47.   Shin C, Feng Y and Manley JL. Dephosphorylated SRp38 acts as a splicing repressor in response to heat shock. *Nature* 2004;427(6974):553-8 doi 10.1038/nature02288.

48.   Jakubauskiene E, Vilys L, Makino Y, Poellinger L and Kanopka A. Increased Serine-Arginine (SR) Protein Phosphorylation Changes Pre-mRNA

Splicing in Hypoxia. *J Biol Chem* 2015;290(29):18079-89 doi

10.1074/jbc.M115.639690.

49.   Ninomiya K, Kataoka N and Hagiwara M. Stress-responsive maturation of

Clk1/4 pre-mRNAs promotes phosphorylation of SR splicing factor. *The Journal*

of cell biology 2011;195(1):27-40 doi 10.1083/jcb.201107093.

50.   Long Y, Sou WH, Yung KWY, Liu H, Wan SWC, Li Q*, et al.* Distinct

mechanisms govern the phosphorylation of different SR protein splicing factors. *J*

Biol Chem 2018 doi 10.1074/jbc.RA118.003392.

51.   Shinde MY, Sidoli S, Kulej K, Mallory MJ, Radens CM, Reicherter AL*, et*

al. Phosphoproteomics reveals that glycogen synthase kinase-3 phosphorylates

multiple splicing factors and is associated with alternative splicing. *J Biol Chem*

2017;292(44):18240-55 doi 10.1074/jbc.M117.813527.

52.   Prasad J, Colwill K, Pawson T and Manley JL. The protein kinase Clk/Sty

directly modulates SR protein activity: both hyper- and hypophosphorylation

inhibit splicing. *Mol Cell Biol* 1999;19(10):6991-7000.

53.   Cao W, Jamison SF and Garcia-Blanco MA. Both phosphorylation and

dephosphorylation of ASF/SF2 are required for pre-mRNA splicing in vitro. *RNA*

1997;3(12):1456-67.

54.   Hong DS, Kurzrock R, Naing A, Wheler JJ, Falchook GS, Schiffman JS*, et*

al. A phase I, open-label, single-arm, dose-escalation study of E7107, a

precursor messenger ribonucleic acid (pre-mRNA) splicesome inhibitor

administered intravenously on days 1 and 8 every 21 days to patients with solid

tumors. *Investig New Drugs* 2014;32(3):436-44 doi 10.1007/s10637-013-0046-5.

55.   Tyanova S, Temu T, and Cox J. The MaxQuant computational platform for

mass spectrometry-based shotgun proteomics, *Nat Protocols* 2016;11:2301-19.

56.   Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, *et al*.

The STRING database in 2017: quality-controlled protein-protein association

networks, made broadly accessible. *Nucleic Acids Res.* 2017;45:D362-D368.

57.   Chen E Y, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, *et al*. Enrichr:

interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC*

Bioinformatics 2013;128:1-14.

1   58.   Vizcaíno JA, Csordas A, del-Toro N, Dianes JA, Griss J, Lavidas I, *et al*.

2   2016 update of the PRIDE database and related tools. *Nucleic Acids Res.*

3   2016;44:D447-56.

Main Figure Titles and Legends

Figure 1. Unbiased phosphoproteomic timecourse analysis of MM.1S cells treated

with the PI carfilzomib (Cfz). A. Workflow of timecourse treatment of MM.1S cells with

Cfz. Cells were allotted for both RNA-seq analysis and LC-MS/MS. **B.** Downregulated

(left) and upregulated (right) $\log_2$ transformed phospho-peptide MS1 intensities for two

technical replicates of proteins with unchanged transcript levels (RNA-seq). Labels

highlight dephosphorylation of RPS6 and EIF4EBP1 on the left and phosphorylation of

splicing-related proteins on the right.

Figure 2. Cfz induces phosphorylation of splicing factors in a dose-responsive

manner. A-D. Volcano plots of log₂ transformed ratios of phosphosite abundances

between **A.** MM.1S treated with18 nM Cfz, **B.** 10 nM Cfz, or **D.** or 10 μM melphalan

compared to DMSO and **C.** AMO-1 with 15 nM Cfz compared to DMSO. Significant

upregulated sites are in red, while downregulated are in blue (>2-fold change, $p < 0.05$).

SRSF related sites are in orange. Circle size corresponds to summed SILAC intensities.

E. Top-ranked GO terms for genes with significantly upregulated phosphosites in MM.1S

cells treated with 18 nM Cfz. **F**. Top-ranked KSEA activated kinases (with at least 5

substrates) for MM.1S treated with 18 nM Cfz (top) and 10 μM melphalan (bottom).

Figure 3. Cfz treatment leads to prominent intron retention. A. Cartoon description

of alternative splicing event (ASE) types and description of ΔPSI. **B.** Histograms of ΔPSI

for JuncBASE identified ASEs in MM.1S treated with 18 nM Cfz stratified according to

type of splicing event (IR = yellow, alt. exon cassette = teal, alt. exon donor (5' splice

site) = pink, alt. exon acceptor (3' splice site) = blue, alt. first exon = green, alt. last exon

= purple). Bin = 2, red line indicates ΔPSI = 0. **C.** Histograms of ΔPSI for all IR events

(top) and Alt. Exon Donor events (bottom) in AMO-1 treated with 15 nM Cfz **D.**

Histograms of ΔPSI for all IR events (top) and Alt. cassette events (bottom) in MM.1S

treated with 10 µM melphalan (right). **E.** Top ranked GO enrichment terms for genes with

significant (*p* < 0.05) ASEs for MM.1S cells treated with 18 nM Cfz (left) or with 10 µM

melphalan (right).

Figure 4. Modeling SRSF1 phosphorylation in MM drives interactome dynamics

but not global splicing changes. A. Cartoon of protein architecture for exogenous

SRSF1-NLS-mCherry-[FLAG]$_3$. **B.** Epi-fluorescent imaging of DAPI stained AMO-1

expressing mCherry labeled SRSF1-WT (top), SRSF1-SD (middle), and SRSF1-SA

(bottom). Scalebar represents 10 µm. **C and D.** Histograms of ΔPSI for IR, alt. exon

donor, alt. exon acceptor, alt. first exon, alt. last exon ASEs when comparing differential

splicing of AMO-1 expressing **C.** SRSF1-WT treated with 15 nM Cfz to DMSO and **D.** SD

to WT. **E.** Volcano plots indicating differential interactors of SD compared to WT in both

the nucleus and cytoplasm. **F.** Volcano plots of WT or SD when compared to control

1   (NLS-mCherry-[FLAG]$_3$) in AMO-1 nucleus reveal SD exclusion from spliceosome.

2   Significant enriched proteins in pink, unenriched proteins in cyan ($p < 0.05$, ≥2-fold

3   change). Circle size corresponds to summed LFQ intensities. mCherry ratio is red and

4   SRSF1 ratio is yellow.

Figure 5. Combined SRSF1 constructs validate the splicing phenotype after Cfz. A.

Histograms of ΔPSI in pooled analysis of parental AMO-1, AMO-1 SRSF1-WT, SD, and

SA expressing cells treated with 15 nM Cfz compared to DMSO. **B.** Graph shows total

number of events (n = 62,474) for each ASE type (left) and only the significant (FDR-

corrected p < 0.05) events (n = 2,575) for each type (right). **C.** Top ranked GO

enrichment terms of all genes involved in significant ASEs, regardless of type. **D.**

Snapshot of UCSC Genome Browser Bar graph compares RNA-seq counts for the

proteasomal subunit *PSMC5* between AMO-1 treated with 15 nM Cfz (bottom) and with

1 DMSO (top). Inset displays sequencing counts showing alternative first exon. **E.** Model

2 of new PI mechanism of action found in MM.

Figure 6. The catalytic spliceosome inhibitor E7107 induces IR and has potent

anti-MM activity *in vitro*. A. ΔPSI histogram of IR events for MM.1S treated with 10 nM

E7107 for 6 hr with respect to DMSO. Bin = 2 and red line at ΔPSI = 0. **B.** Cell viability

curves compare a panel of 7 MM cell lines treated with E7107 for 48 hr (n = 4; mean +/-

S.D.). **C.** Cell viability comparison of WT AMO-1 and PI-resistant AMO-1 cell line (BtzR)

when treated with E7107 (n = 4; mean +/- S.D.) **D.** Evaluation of SF3B1 expression by

RNA-seq (www.keatslab.org, mean +/- S.D.) and **E.** Western blot across more E7107-

sensitive (AMO-1, INA6, L363, and RPMI8266) and less E7107-sensitive (MM.1S,

KMS34, JJN3) cell lines. **F.** Cell viability curves of MM.1S combination therapy with

E7107 and Cfz (n = 4; mean +/- S.D.) **G.** 2-D heatmap of ZIP synergy-scored landscape

from Cfz and E7107 combination study. Red = synergy; green = antagonism; white =

1   additive. Overall ZIP score of 6.295 suggests strong synergy. **H and I.** Same as **F.** and

2   **G.** but for melphalan + E7107 combination. Overall ZIP synergy score of 2.663 denotes

3   weaker synergy than with Cfz.

Figure 7. Inhibition of the spliceosome is a promising therapeutic strategy in

myeloma. A-B. Bioluminescence imaging of luc-labeled MM.1S cells implanted in mice

treated with either vehicle (left, n = 6) or 3 mg/kg E7107 (right, n = 6). Green arrows

indicate days when drug was administered (14-18, 21-25). **C.** E7107 leads to significant

improvement in murine survival (*p* = 0.01 by log-ranked test) **D.** Treatment of primary

bone marrow aspirate samples from PI-refractory myeloma patients at various doses of

E7107 for 24 hr shows significant cytotoxicity to CD138+ MM plasma cells at low-nM

concentrations but minimal effects on other (CD138-) hematopoietic cells (n = 2

technical replicates; mean +/- S.D.). **E.** Heatmap of CRISPR-Cas9 essentiality screen

data analysis in the Cancer Dependency Map (www.depmap.org; **Avana 18Q4** release)

of core spliceosomal subunits among all tested tumor cell types. **F.** Analysis of MMRF

CoMMpass data (research.themmrf.org; release **IA11**) summarizing mutations with

possible functional effects in numerous splicing-related factors, as defined by Seiler *et*

al. (46), within MM patient plasma cells.