Title: Removal of a subset of non-essential genes fully attenuates a highly virulent *Mycoplasma*

strain

Authors: Joerg Jores*[1,2]; Li Ma[3#]; Paul Ssajjakambwe[2,4#]; Elise Schieck[2]; Anne Liljander[2];

Suchismita Chandran[3]; Michael Stoffel[5]; Valentina Cippa[1]; Yonathan Arfi[6,7]; Nacyra Asad-

Garcia[3]; Laurent Falquet[8]; Pascal Sirand-Pugnet[6,7]; Alain Blanchard[6,7]; Carole Lartigue[6,7];

Horst Posthaus[9]; Fabien Labroussaa[1]; and Sanjay Vashee[3]

To whom correspondence should be addressed: Joerg Jores, Tel: +41 31 631 2414,

joerg.jores@vetsuisse.unibe.ch

# equal contribution

[1] Institute of Veterinary Bacteriology, University of Bern, PO Box, CH-3001 Bern, Switzerland

[2] International Livestock Research Institute, Old Naivasha Road, PO Box 30709, 00100

Nairobi, Kenya

[3] J. Craig Venter Institute, 9704 Medical Center Drive, MD 20850 Rockville, USA

[4] Makerere University Kampala, P.O Box 7062, Kampala Uganda.

[5] Division of Veterinary Anatomy, University of Bern, PO Box, CH-3001 Bern, Switzerland

[6] INRA, UMR 1332 de Biologie du Fruit et Pathologie, F-33140 Villenave d'Ornon, France

[7] Univ. Bordeaux, UMR 1332 de Biologie du Fruit et Pathologie, F-33140 Villenave d'Ornon,

France

[8] Biochemistry Unit, University of Fribourg and Swiss Institute of Bioinformatics, Fribourg,

Switzerland

[9] Institute of Veterinary Pathology, University of Bern, PO Box, CH-3001 Bern, Switzerland

Keywords: Mycoplasmas ; Attenuation ; Genome Engineering; *in vivo* challenge

ABSTRACT:

Mycoplasmas are the smallest free-living organisms and cause a number of economically important diseases affecting humans, animals, insects and plants. Here, we demonstrate that highly virulent *Mycoplasma mycoides* subspecies *capri* (*Mmc*) can be fully attenuated *via* targeted deletion of non-essential genes encoding, among others, potential virulence traits. Five genomic regions, representing approximately ten percent of the original *Mmc* genome, were successively deleted using *Saccharomyces cerevisiae* as an engineering platform. Specifically, a total of 68 genes out of the 432 genes verified to be individually nonessential in the JCVI-Syn3.0 minimal cell, were excised from the genome. *In vitro* characterization showed that this mutant was similar to its parental strain in terms of its doubling time, even though ten percent of the genome content were removed. A novel *in vivo* challenge model in goats revealed that the wild-type parental strain caused marked necrotizing inflammation at the site of inoculation, septicemia and all animals reaching endpoint criteria within seven days after experimental infection. This is in contrast to the mutant strain, which caused no clinical signs nor pathomorphological lesions. These results highlight, for the first time, the rational design, construction and complete attenuation of a *Mycoplasma* strain via synthetic genomics tools. Trait addition using the yeast-based genome engineering platform and subsequent *in vitro* or *in vivo* trials employing the *Mycoplasma* chassis will allow us to dissect the role of individual candidate *Mycoplasma* virulence factors and lead the way for the development of an attenuated designer vaccine.

IMPORTANCE

Members of the *Mycoplasma mycoides* cluster cause important animal plaques in Africa and Asia, which impact animal welfare, provision of food and the life of thousands of small-scale farmers. We applied synthetic biology tools to *Mycoplasma mycoides* in order to design and create a fully attenuated *Mycoplasma* strain that was subsequently confirmed *in vivo* using a novel caprine infection model. This is the first time that a *Mycoplasma* mutant developed by applying synthetic biology tools has been tested *in vivo* to pin point candidate virulence traits. The mutant strain is similar to "apathogenic *E. coli* K12" strains that boosted the research on host-pathogen interactions for the genus *Escherichia* and other bacterial genera.

INTRODUCTION

Bacteria belonging to the genus *Mycoplasma* are wall-less bacteria that cause massive economic losses in the livestock sector (chickens, ruminants and pigs) and are responsible for human pneumonia and sexually transmitted diseases (STDs). Currently, there is an absence of commercial vaccines against infections with the human pathogens *Mycoplasma pneumoniae* and *Mycoplasma genitalium* (1). In contrast, many livestock vaccines are commercialized, which rely either on adjuvanted killed bacteria or on attenuated strains obtained after successive rounds of sub-culturing or chemical mutagenesis (2). Due to these empirical approaches, the exact mechanism triggering the attenuation is unknown for many of the previously developed live attenuated *Mycoplasma* vaccines. Strikingly, these vaccines are far from being optimal since they often display short durations of immunity and limited efficacy (3-5). A better understanding of pathogenicity and the identification of virulence traits would foster next generation vaccines.

For many years, the lack of genetic tools has limited our basic understanding of *Mycoplasma* pathogenicity. Due to their regressive evolution by gene loss , mycoplasmas appear to lack many of the common bacterial effectors and toxins used to interact with their hosts or to escape the hosts' immune systems (6, 7). Lipoproteins have been proposed to be involved in both aspects by using their cytoadherent properties and allowing antigenic variability through phase or sequence variation (8). Other candidate virulence traits, such as the *Mycoplasma* Ig binding protein-*Mycoplasma* Ig protease (MIB-MIP) system (9) and the hydrogen peroxide production system (10) have been suggested, but not confirmed *in vivo*.

The availability of a genome engineering platform that allows directed and precise mutagenesis for *Mycoplasma mycoides* is undoubtedly a new starting point towards better understanding of host-pathogen interactions. In this work, we engineered a *Mmc* strain by deleting approximately one tenth of the genome, including candidate virulence traits. The resulting mutant retains

77   almost wild-type like growth characteristics and was attenuated both *in vitro* and *in vivo*. The

78   construction of this fully attenuated and safe laboratory *Mycoplasma* strain paves the way for

79   research into host-pathogen interactions and is a good starting point to revisit the actual role of

80   suggested virulence determinants in *Mycoplasma*.

RESULTS

Generation of the mutant strain GM12::YCpMmyc1.1-*Δ68*

To demonstrate attenuation of *Mmc* by rational design, five genomic regions were targeted in this study. The precise localizations in the *Mmc* GM12 genome are shown in Figure 1. The first two target deletion regions contained genes encoding the glycerol-dependent hydrogen peroxide metabolic pathway and its suggested ABC transporter encoded by the *gtsABCD* operon (11). This pathway has been suggested to be a main virulence mechanism for *M. mycoides* (11), but *in vivo* confirmation is still missing and in *M. gallisepticum* the pathway does not seem to be linked to virulence (12). Thus, the genes *glpF*, *glpK* and *glpO* (MMCAP2_0217-0219; 2,984-bp region; D1) and the *gts* gene region that includes the gene *lppB* (MMCAP2_0456-0459; 4,950-bp region; D2) were deleted in the *Mmc* genome by the yeast-based engineering method (13). As previously mentioned, lipoproteins were another target of interest since they likely trigger not only host-pathogen interactions but also, overwhelming immune reactions that result in inflammation (14). The lipoproteins encoded in the D3 (MMCAP2_0014-0016; 4,677-bp region; D3) as well as the D5 region (*lppQ*, MMCAP2_0889-0904; 24,906-bp region, D5) were also excised employing again the yeast-based engineering method. Finally, we deleted a large genomic region that contained the *Mycoplasma*-specific F1-likeX0 ATPase (15), the MIB-MIP system (9) that has been shown *in vitro* to modulate the action of immunoglobulins through specific degradation, an integrative and conjugative element (ICE) (16) and several genes encoding other lipoproteins. The ICE was targeted in an effort to reduce mobile elements from the *Mmc* genome. In this case, about 70 kbp (MMCAP2_0550-0591; 69,220-bp region; D4) were targeted and deleted from the *Mmc* genome using the yeast-based engineering method in one stretch.

Figure 1: Design of deletion mutant *Mmc* strain. (A) Schematic displaying the genomes of the isogenic parental strain GM12::YCpMmyc1.1 and its derivative, the deletion mutant GM12::YCpMmyc1.1-*Δ68*.

After each cycle of deletions, the modified *Mmc* genome was isolated from yeast cells and

transplanted back into *M. capricolum* subsp. *capricolum* (*Mcc*) recipient cells to confirm the

viability of each mutant *Mmc* strain. Overall, the final mutant strain, named *Mmc*

GM12::YCpMmyc1.1-*Δ68,* was generated in five sequential deletion cycles (Figure 1). The

gene knock-outs were verified by amplifying across each deleted region (Figure S1). Genomic

DNA from the GM12::YCpMmyc1.1-*Δ68* was isolated and analyzed by sequencing to confirm

the deletions (Figure1B). The genome sequence of GM12::YCpMmyc1.1-Δ68 was deposited

at the ENA database under the accession number LS483503.

The mutant strain GM12::YCpMmyc1.1-Δ68 is viable and unaffected in its morphology or growth in axenic medium

The colonies of *Mmc* GM12::YCpMmyc1.1-Δ68 were of similar size to those of

GM12::YCpMmyc1.1 and GM12. Cell morphology of the GM12, the isogenic parental strain

GM12::YCpMmyc1.1 and GM12::YCpMmyc1.1-Δ68 strains was evaluated using scanning

electron microscopy (Figure 2A). All strains tested were globular in shape and lacked any

special morphological features. The diameter of the microorganisms was in the range of 500

nm, as expected for a *Mycoplasma* cell. The mutant GM12::YCpMmyc1.1-Δ68 grew with a

doubling time somewhat similar to that of the parental strains GM12 and GM12::YCpMmyc1.1

(Figure 2B). Together, these results strongly suggest that the deletion of approximately 100 kbp

of genomic content from the *Mmc* genome did not adversely affect structural integrity or *in*

vitro growth of the mutant GM12::YCpMmyc1.1-Δ68.

Inability of the mutant strain GM12::YCpMmyc1.1-Δ68 to produce hydrogen peroxide in the presence of glycerol *in vitro*

This pathway was completely deleted in the construction of the mutant strain

GM12::YCpMmyc1.1-Δ68. Therefore to phenotypically confirm the deletion, we measured and

compared hydrogen peroxide production levels between the control GM12,

GM12::YCpMmyc1.1 and GM12::YCpMmyc1.1-Δ68 *in vitro*. In the presence of the glycerol

substrate, the mutant GM12::YCpMmyc1.1-Δ68 shows a significant decrease in hydrogen

peroxide production when compared to its parental strains (Figure 2C). Indeed, while GM12

and GM12::YCpMmyc1.1 produced >0.3μM of H_2O_2, the mutant strain produced very low

amounts of H_2O_2 (0.01μM), at least 30-fold lower under these conditions.

Fig. 2

A

Figure 2: *In vitro* characteristics of the parental strains GM12, GM12::YCpMmyc1.1 and its deletion mutant GM12::YCpMmyc1.1-*Δ68* with respect to their morphology revealed by scanning electron microscopy (A), their doubling time (B), their production of hydrogen peroxide in the presence of glycerol (C), the bars display the standard deviation in B and C

Inability of the mutant strain GM12::YCpMmyc1.1-*Δ68* to degrade immunoglobulin *in

*vitro***

Another potential virulence trait encoded by mycoplasmas is the MIB-MIP system, which may

play a role in immune evasion by cleavage of immunoglobulins (Figure 3A) (9). Incubation of

caprine IgG with GM12, GM12::YCpMmyc1.1 and GM12::YCpMmyc1.1-*Δ68* showed a clear

difference in the strains' abilities to degrade IgG (Figure 3B). The two bands at 25 kDa and

50kDa corresponds to the IgG light and heavy chains. The mutant strain GM12::YCpMmyc1.1-

Δ68 exhibited no degradation of IgG, as noted by the lack of the 44 kDa band (Figure 3B, black

asterisk). This band, clearly visible in the other two strains, is indicative of proteolytic cleavage

of the IgG heavy chain. Another pattern of degradation, with a band at a size of about 30 kDa,

is visible in the three strains. It was previously reported that this IgG cleavage is not specific or

directly linked to the MIB-MIP system (9).

Fig. 3

Figure 3: *In vitro* and *in vivo* ability of the *Mmc* MIB-MIP system to degrade caprine immunoglobulin: (A) cartoon displaying the suggested action of the MIB-MIP system, (B) *in vitro* ability of the parental strains GM12, GM12::YCpMmyc1.1 and their deletion mutant GM12::YCpMmyc1.1-Δ*68* to degrade caprine immunoglobulin, 1-GM12 + 5 µg caprine IgG, 2-GM12::YCpMmyc1.1 + 5 µg caprine IgG, 3-GM12::YCpMmyc1.1-Δ*68* + 5 µg caprine IgG; the black asterisk marks the 44 kDa cleaved fragment (C) *In vivo* detection of degradation of the *Mmc* MIB-MIP system using serum samples from the animal CK51 that succumbed from disease and had septicemia, 4-5 µg caprine IgG, 5-serum from goat CK51: -1 dpi (GM12 group), 6-serum from goat CK51: 4 dpi (GM12 group), 7-serum from goat CK45: -1 dpi (GM12::YCpMmyc1.1-Δ*68* group), 8-serum from goat CK45: 4 dpi (GM12::YCpMmyc1.1-Δ*68* group)

The mutant strain GM12::YCpMmyc1.1-Δ68 is fully attenuated *in vivo*.

We next tested whether GM12::YCpMmyc1.1-Δ68 was able to cause disease in its native host.

Sixteen male outbred goats (*Capra aegagrus hircus*) were used in this animal infection trial.

The animals were separated into two groups of equal numbers. After the infection, no

immediate clinical signs of disease were observed. Two animals in the GM12::YCpMmyc1.1-

Δ68 group had to be removed from the experiment, because of acquired wounds unrelated to

the infectious agent. Animal euthanasia was planned 28 days post infection (dpi). However, all

eight animals inoculated with the GM12 strain developed severe clinical signs, with pyrexia

starting 2-3 dpi (Figure 4B). Their body temperature continued to increase, up to 41-41.5°C,

during the following days. The animals stopped feeding, were apathic and showed signs of pain.

According to the endpoint criteria stated in the animal experiment protocol, they had to be

euthanized between 5-6 dpi (Figure 4A, red line). In sharp contrast, animals inoculated with the

GM12::YCpMmyc1.1-Δ68 mutant strain did not develop any clinical signs of disease and were

all monitored until the end of the trial (Figure 4A). Their body temperatures fluctuated between

38°C and 39°C, with a few isolated cases where animals showed temperatures above 39°C but,

never for more than 2 consecutive days (Figure 4B). The animals appeared to remain healthy

and gained weight during the experiment (Figure 4C). Their heartbeat and respiratory rates,

between 80-110 beats/min and 20-30 breaths/min, respectively, remained constant over the

course of experimentation.

Fig. 4

Figure 4: Comparison of the clinical parameters monitored during the in vivo challenge trial between the animals that received the GM12 and its derivative GM12::YCpMmyc1.1-Δ*68*. (A) Kaplan Meier survival curve based on animals reaching endpoint criteria, (B) Average body temperatures during experimental infection. Values were generated using daily rectal temperatures from the two groups. (C) Average weight gain during experimental infection. Values were generated using interval measures from the two groups. The standard deviations are displayed as bars in B and C.

In all animals infected with the GM12 strain, the main pathological lesions were similar, with

a severe and extensive inflammation of the soft tissues of the neck around the site of inoculation.

Additional macroscopic findings were severe pulmonary edema and congestion. Histologically,

there was extensive coagulation necrosis of the connective tissue (Figure 5B, thick arrows) and

musculature surrounding the trachea, in the vicinity of the inoculation site (Figure 5B and D,

diamonds). A marked infiltration of mainly degenerate neutrophilic granulocytes (Figure 5B

and D, asterisks) was always found associated with the necrosis. The necrotizing process

extended to the trachea, the subcutis and skin. In addition, all animals showed multifocal acute

necrosis with infiltration of neutrophilic granulocytes in liver and kidney and for three animals,

in the lung. These lesions were indicative of an acute septicemia. Among all animals infected

with the mutant strain GM12::YCpMmyc1.1-*Δ68*, and upon euthanasia 28-29 dpi, no

pathological lesions were found around the inoculation site. The soft tissue around the

inoculation site was within normal limits. Additionally, neither inflammation nor necrosis

associated with the epithelium, the submucosa or any cartilage tissues was histologically

observed (Figure 5, panels A and C).

Mmc GM12 was re-isolated from the blood of all animals experimentally infected with the

wild-type parental strain. Bacteremia was characterized by 10^6 up to 10^9 CCU.ml^{-1} of blood, as

measured by serial dilutions (Table 1). Blood samples collected from goats infected with

GM12::YCpMmyc1.1-Δ*68* did not reveal any bacteremia.

Fig. 5

Figure 5: Composite figure displaying representative histological results from tracheal tissue at the inoculation site. Tissues were stained with hematoxylin and eosin. Low (40x) and high (200x) magnification of trachea of a goat inoculated with *Mmc* GM12::YCpMmyc1.1-Δ*68* (panels A and C) depicting unaffected epithelium (circle), submucosa (diamond) and cartilage (asterics). or of a goat inoculated with *Mmc* GM12 (panels B and D), depicting ulceration of the epithelium (thick arrow, panel B), massive extension of the submucosa due to extensive areas of necrosis (diamond, panels B and D) and infiltration with large numbers of degenerate neutrophilic granulocytes (asterisk, panels B and D). Size standards are displayed in each picture.

The MIB-MIP system in *Mmc* GM12 is functional *in vivo*

The MIB-MIP system was shown to be active *in vitro*, and to be present in large amounts at the

cell surface (17) during infection (18). Two animal sera were selected from the *in vivo* infection

trial: CK51 from the GM12 group and CK45 from the GM12::YCpMmyc1.1-Δ*68* group and

tested for cleavage of IgG. The pre-infection sera from both animals demonstrated no

proteolytic cleavage of IgG when compared to the IgG control (Figure 3C). Conversely, the

post-infection serum of CK51, which had succumbed to disease and had a high titer of bacteria

in the venous blood, clearly exhibited the typical band at 44 kDa, consistent with the size of a

cleaved IgG heavy chain by the MIB-MIP system. As expected, no band at 44 kDa was seen in

the post-infection serum of CK45. This clearly demonstrates, for the first time, that the MIB-

MIP system is functional within the caprine host and that its proteolytic activity is triggered

during an *Mmc* infection.

DISCUSSION

The first aim of this work was to fully attenuate a highly pathogenic strain of

Mycoplasma mycoides following a rational deletion design. The second aim was to verify this

attenuation *in vivo* using the native host, since no rodent animal models for highly virulent *M.*

mycoides exist (3). Many candidate virulence factors of *Mycoplasma mycoides* have been

suggested but, all except capsular polysaccharide (19) have never been confirmed according to

Falkow's postulates *in vivo* (20).

First, we relied on previous knowledge to select five genomic regions that encode candidate

virulence traits. These regions, distributed around the *Mmc* genome, consisted of 68 genes. The

first two regions (D1 and D2) encode enzymes and putative glycerol transporters involved in

the production of hydrogen peroxide using the glycerol-dependent metabolism of mycoplasmas

(10). The region D3 encodes major antigens (LppA/P72) in *Mycoplasma mycoides* (21) that

induced T cell responses early in infection (22). The region D4 includes an integrative

conjugative element (ICE), the MIB-MIP system (9) and the F1-likeX0 ATPase (15). The last

region (D5) encodes six lipoproteins including the lipoprotein Q (LppQ), which has been

suspected to be involved in exacerbating immune responses and other virulence determinants

(23). Interestingly, among these 68 genes deleted in our mutant, 67 were defined as non-

essential in the JCVI-Syn3.0 minimal cell, a study in which 432 genes were classified as non-

essential to sustain the life of a minimal *Mycoplasma* cell (24). Only one gene encoding the

glycerol phosphate kinase (GlpK) (MMCAP2_0218, region D1), which was retained in the

minimal JCVI-Syn3.0 cell but, classified as a "quasi-essential" gene. This means that the

function encoded by the glycerol kinase, i.e. the transfer of a phosphate group on the glycerol

molecule to produce glycerol-3-phosphate, could be compensated for by another gene which

encodes a similar function in the full-length genome. The compensatory gene for *glpK* is

probably   absent   in   the   minimal   JCVI-Syn3.0   cell,   but   present   in   our   mutant

GM12::YCpMmyc1.1-*Δ68*, allowing the deletion of *glpK* without any defect in growth.

Therefore, our results are consistent with the quasi-essentiality of *glpK*. Altogether we retained

69 out of the 87 lipoproteins (D1-0, D2-1, D3-3, D4-8, D5-6) present in GM12, while the

minimal cell retained only 15 lipoproteins (24).

It was paramount for us that the mutant strain GM12::YCpMmyc1.1-*Δ68* maintains a doubling

time similar to its parental strain, since we wanted to create a 'K12-like' strain that can be

further used as a cellular platform to introduce antigens or stretches of DNA. It was interesting

for us to observe that, despite complete removal of the glycerol pathway, which maybe

important for cell metabolism, there was no significant impact on *in vitro* growth. It is known

that there is a trade-off between genome size and growth rate. The drastic deletions in the

genome of JCVI-syn3.0 strain led to a substantial increase in the generation time, from ~60 min

to ~180 min (24). Recently, we also have shown that the deletion of a gene encoding an enzyme

important for synthesis of carbohydrates can subsequently lead to an increase in the generation

time (25). However, in this work, we significantly reduced the genome of GM12 by more than

100 kbp (i.e. 106,737 bp) without seeing any significant difference in the growth rate of

GM12::YCpMmyc1.1-Δ*68* in comparison to the wild-type strain. Still, this reduction represents

~10% of the initial genome size confirming that, in addition to the size of the deletions, the

nature of the genes deleted is also very likely to influence the generation time of mycoplasmas.

The main goal of this work was to construct a fully attenuated *Mmc* strain, that is safe to handle

in the laboratory. The first confirmation of the attenuation of the GM12::YCpMmyc1.1-Δ*68*

strain was obtained *in vitro*. The production of hydrogen peroxide was almost completely

abolished in the mutant strain, confirming the participation of the *glpFKO* and/or *gtsABCD*

pathways in the metabolism of glycerol. In addition, the loss of the IgG specific cleavage band

at 44 kDa in GM12::YCpMmyc1.1-Δ*68* confirmed the role of the MIB-MIP system in the

degradation of the host immunoglobulins.

To confirm the mutant strain's attenuation *in vivo*, we developed an animal challenge model

using Kenyan goats, outbred animals derived from different herds. The use of such animals

increased variability to get a better idea of reproducibility and significance of the results (26).

Animals infected with the GM12 strain developed specific symptoms (fever, heavy breathing,

septicemia, etc.) and were all euthanized by 6 dpi. Strikingly, none of the goats infected with

GM12::YCpMmyc1.1-Δ*68* developed such symptoms and were healthy for the entire course of

the experimentation. These results exceeded our expectations and confirmed the complete

abolishment of pathogenicty of the mutant strain GM12::YCpMmyc1.1-Δ*68*. In addition, the

massive septicemia associated with very high titers of *Mycoplasma* observed in animals

infected with the GM12 strain prompted us to investigate whether the MIB-MIP system would

leave signatures of its action on immunoglobulins (Ig) in the serum of an animal (CK51) that

had a titer of 10^9 CFU/ml. Specific IgG cleavage, characteristic of the MIB-MIP system (9),

was observed. No such cleavage was observed in the serum of animals infected with the mutant

strain. This work shows, for the first time, that the MIB-MIP system of *Mycoplasma mycoides*

is functional *in vivo*. *Mycoplasmas* have been viewed as stealth pathogenic organisms because

they lack most of the immune activators or PAMPs found in other bacteria (27). Indeed, the

lack of a cell wall or the capacity to produce either LPS or flagellins likely contribute to the

chronicity of infection. The only PAMP that has been described for several *Mycoplasma* species

is the surface lipoproteins, abundant components of their membrane (8). In the present study,

we suggest another mechanism that could contribute to activating the immune system. Indeed,

Ig cleaved by several bacteria, including those generated by *Mycoplasma hyorhinis*, have been

described as ligands of the innate immune receptor LILRA2 (28). Once bound to this receptor,

it triggers the activation of the innate immune system. It is also possible that this cleavage is in

line with the 'nutritional virulence' of the parasite (29). The exact significance of the Ig cleavage

regarding *Mycoplasma* infection of mucosal surfaces remains to be studied.

Interestingly, we observed severe inflammation around the site of injection in the animals that

received GM12 whereas animals that were injected with the strain GM12::YCpMmyc1.1-*Δ68*

developed no such pathomorphological lesions. Overwhelming immune reactions at the site of

vaccination have been reported from immunizations against contagious bovine

pleuropneumonia using live *M. mycoides* subsp. *mycoides* based vaccines such as T1/44, which

is the closest relative of *Mmc* (30). Therefore, it is likely that any one or several of the deleted

genes encode proteins that drive this overwhelming immune reaction in the GM12.

To conclude, we confirmed, *in vitro* and *in vivo*, our ability to design a fully attenuated strain

via the precise reduction of ~10 % of the *Mmc* genome. However, we cannot currently pinpoint

the weight of each deletion on the observed attenuation. The total clearance of the pathogen

and the absence of a compelling humoral immune response, even at the inoculation point, is

surprising and supports the total abolishment of pathogenicity. Now it is necessary to test more

defined mutants such as a *glpOKF* mutant strain to get clarity about its real role in

pathogenicity.

In addition, the design of next generation vaccines for, but not restricted to, *Mycoplasma*

diseases will benefit from this study since a chassis that is fully attenuated and able to

accommodate antigens for vaccine delivery that can be constructed based on our deletion

mutant is now within reach. To induce a proper immune response via such a chassis, we have

the option to add genes that appropriately stimulate an inflammatory immune response or

alternatively, we can construct different chassis that direct responses towards Th1 or Th2 using

TLR agonists. We consider a genetically modified *Mycoplasma* less problematic than other

potential chassis since the survival time of *Mycoplasma* in general in the environment is very

short. In addition, the unconventional codon usage (where UGA encodes tryptophan) and high

AT content of *Mycoplasma* minimizes the risk of spread of genes to other bacteria. Additional

experiments are necessary to decipher the role of individual virulence traits to understand these

minimal bacterial pathogens better and to develop next generation rationale vaccines.

Regardless, this study provides an attractive blueprint towards these goals, especially for those

that are needed in low and middle-income countries.

MATERIALS AND METHODS

Mycoplasma strains

The *Mycoplasma mycoides* subsp. *capri* outbreak strain GM12 was used as positive control in the *in vivo* experiment (31). A modified *Mycoplasma capricolum* subsp. *capricolum* strain CK was used as recipient strain in genome transplantation protocols (13).

Yeast strain and media

The yeast *Saccharomyces cerevisiae*, strain VL6-48N (MATαhis3-Δ200 trp1-Δ1 KlURA3-Δ1 lys2 ade2-101 met14) containing the 1.08 Mb genome of *Mycoplasma mycoides* subsp. *capri* (*Mmc*) strain GM12 with an integrated yeast centromeric plasmid (YCp) (13) was used for construction of the mutants. Yeast cells were grown and maintained in either synthetic minimal medium containing dextrose (SD, Takara Bio) (13), or in standard rich medium containing glucose (YPD, Takara Bio) or galactose (YPG, Takara Bio) (32). SD medium was supplemented with 5-fluoroorotic acid (5-FOA), for KlURA3 counter-selection (13, 33).

Preparation of mutagenesis cassettes

Sixty eight genes that encode candidate virulence traits were seamlessly deleted from the genome of *Mmc* GM12::YCpMmyc1.1 in five consecutive cycles (D1, D3, D4 and D5) using Tandem Repeat coupled with Endonuclease Cleavage [TREC] as described (32, 34) or a variation of TREC involving the Cre-lox system for the D2 deletion, see below. Primer sequences to target and confirm the insertion of the mutagenesis cassette into each target site and to verify seamless deletion of the targeted genes are shown in Table S 1.

The *gts* gene cluster (D2) was targeted and deleted in the *Mmc* genome in the back-ground of the *glpFKO* deletion strain by employing a derivative of the *Mmc* synthetic cell JCVI-syn1.0 (35). Primers RC0905 and RC0906 (Table S1) were used to amplify the mutagenesis cassette from the synthetic cell derivative and was targeted to the *gts* region. Specific primers were used to confirm correct insertion at the target site by amplifying the junctions between the

341 GM12::YCpMmyc1.1 genome and the inserted cassette. Galactose induction resulted in the

342 Cre-mediated deletion of the *gts* region, leaving 13 bp of the 5' end of the *gtsA* region, the 34

343 bp *loxP* site, and 27 bp of the 3' end of the *lppB* gene. Specific primers were used to verify the

344 knock-out.

Transformation and PCR analysis

Transformation of the CORE3 cassette was performed by the lithium acetate method as

described previously (36). Transformed yeast were plated on appropriate selection media [SD

medium minus His (Teknova, CA) or SD medium (minus His and minus Ura)] and incubated

at 30°C for 48 hours. Yeast colonies were patched on appropriate selective media and total

DNA was isolated for PCR screening (37). The correct insertion of the mutagenesis cassette

was verified by PCR amplification using upstream and downstream specific primers (Integrated

DNA Technologies, Coralville, IA, USA) (Table S1).

Transplantation

354 The modified GM12::YCpMmyc1.1 genomes (D1 to D5) were transplanted into *Mycoplasma*

355 *capricolum* subsp. *capricolum* (*Mcc*) recipient cells with polyethylene glycol and selected for

356 tetracycline resistance as described previously (13, 37). The resulting mutant strains were

357 subjected to multiplex PCRs and pulsed-field gel electrophoresis as described elsewhere (38)

358 to confirm integrity of the genome.

Confirmation of the mutants using next generation sequencing and mapping assembly

Total DNA of the strains GM12, GM12::YCpMmyc1.1 and GM12::YCpMmyc1.1-Δ*68* was

isolated as described before (39). DNA was sheared using sonication and subjected to Illumina

sequencing using a MiSeq machine by University of California Santa Cruz, CA (USA). Reads

were mapped to the designed genome sequences based on the parental strains GM12 and

GM12::YCpMmyc1.1 and GM12::YCpMmyc1.1-Δ*68*. The raw reads (300bp PE) were QC

with FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). The corrected

reads were mapped onto the reference genome WT-YCP.fa with bwa mem (40) and converted

to sorted bam with samtools (41). The bam files were analyzed for deletions using Delly2 (42)

and Sprites (43), and the predictions validated visually using IGV (44). The list of strains and

their deleted regions is summarized in Table 2.

Scanning electron microscopy of *Mycoplasma*

Unless stated otherwise, chemicals were obtained from Merck (Schaffhausen, Switzerland).

Mycoplasmas were washed with distilled water (dH$_2$O) and fixed with 4% para-formaldehyde

(Life Technologies, Thermo Fisher, Zug, Switzerland; Cat. No. 28906) in dH$_2$O for five days

at 4°C. Thereafter, samples of 40 µl of cell suspension were centrifuged onto gold-sputtered

poly-L-lysine coated coverslips (high molecular poly-L-Lysine hydrobromide) at 125 *rcf* for 4

min. Coverslips were washed 1x with PBS and 2x with 0.1% bovine serum albumin in PBS

(BSA/PBS). Free aldehydes were blocked with 0.05M glycine in 0.1% BSA-c/PBS (Aurion,

ANAWA Trading, Wangen, Switzerland) for 15 min. at room temperature. After 3x washes

with 0.1% BSA/PBS, cells were fixed with 2.5% glutaraldehyde (Merck 104239) in 0.1M

cacodylate buffer (dimethylarsinic acid sodium salt trihydrate), washed 3x with dH$_2$O and

postfixed with 1% OsO$_4$ (Polysciences, Warrington, PA, USA) in 0.1M cacodylate buffer for

15 min. at room temperature. Another 5x washes with dH$_2$O were followed by dehydration in

an ascending ethanol series. Samples were then transferred to hexamethyldisilazane (Merck

814051) for 10 min, air-dried, mounted onto aluminum stubs with carbon conductive adhesive

tabs (Ted Pella Inc., Redding, CA, USA) and coated with approximately 25 nm of gold in an

SCD004 (Leica Microsystems, Heerbrugg, Switzerland). Secondary electron micrographs and

corresponding backscattered images were obtained with a fully digital field emission scanning

electron microscope DSM 982 Gemini (Zeiss, Oberkochen, Germany) at an accelerating

voltage of 5 kV, a working distance of 6–8 mm and primary magnifications ranging from

30,000 to 50,000×.

Growth assay

Overnight cultures of *Mycoplasma* strains were grown at 37°C in SP4 medium containing streptomycin (GM12) or tetracycline (GM12::YCpMmyc1.1, GM12::YCpMmyc1.1-*Δ68*) for about 16 hours. Doubling times of the Mycoplasma strains were then determined as described elsewhere (24), except that time interval samples were collected and processed at 0, 1, 2, 3, 4, 5, 6, 7, 9, 12, 15, and 24 h.

***In vitro* hydrogen peroxide assay**

Overnight cultures of *Mycoplasma* strains were grown as described above. When the pH of overnight cultures reached 6.0 - 6.5, they were inoculated into fresh SP4 medium at 1:200 dilution and incubated at 37 °C for different time intervals of 0, 5, 7, and 24 h. At each time interval, an aliquot of culture was taken for DNA extraction (24) and another aliquot was taken to determine hydrogen peroxide levels.

To determine hydrogen peroxide levels, the aliquots were spun at 14,000 rpm for 10 minutes at 4°C. The pellets were washed with 1 ml of cold PBS, pH 7.5 to remove traces of media, then resuspended in 400 µl of cold PBS and stored at 4°C. Hydrogen peroxide levels were determined using the Amplex Red Hydrogen Peroxide Assay Kit (Life Technologies, NY) according to the manufacturer's instructions. Briefly, 50 µl of diluted samples (1:5 in PBS) was aliquoted onto 96-well plates and warmed to 37°C for 1 h prior to starting the assay. 100 µM final concentration of glycerol (Sigma Aldrich, MO) or GPC (Sigma Aldrich, MO) was then added to the diluted sample and incubated at 37°C for 1 h. 50 µl of the Amplex Red reagent was added to the samples, incubated at room temperature in the dark for 30 minutes and fluorescence was measured using a spectrophotometer (SpectraMax M5, Molecular Devices, CA). Three technical replicates were performed for each sample and normalized to their respective DNA concentrations.

Detection of immunoglobulin degradation by *Mycoplasma in vitro* and *in vivo*

In vitro functionality of the MIP-MIP system was tested using the strains GM12,

GM12::YCpMmyc1.1 and GM12::YCpMmyc1.1-Δ*68*. Each strain was grown overnight at

37°C in 3 mL modified SP5 medium (containing 5% FBS). 250 μL of each culture was

harvested and centrifuged for 10 minutes at 4,000 *g*. The cells were then resuspended in 15 μL

modified SP5 medium (containing 5% FBS) and incubated with 5 μg of purified caprine IgG

(Sigma) at 37°C for 3 hours. Bacterial CFUs were estimated for each strain by serial dilutions

and were $3.2*10^9$ CFUs *Mmc* GM12, $8.5*10^8$ CFUs *Mmc* GM12::YCpMmyc1.1 and $3.5*10^8$

CFUs *Mmc* GM12::YCpMmyc1.1-Δ*68*. A sample consisting of 5 μg caprine IgG in dH$_2$O was

included as a control. The incubated samples were mixed with 2x Laemmli Sample Buffer (Bio-

Rad,) at a 1:1 ratio, boiled for 10 minutes at 98°C and separated onto a 12% SDS-PAGE gel.

They were subsequently transferred onto a 0.2 μm nitrocellulose membrane (Bio-Rad) using a

Bio-Rad Trans-Blot® Turbo™ Transfer System (25 volts, 1.0 A, 30 minutes). Next, a Western

Blot was performed using PBS supplemented with 0.1% Tween-20 and 2% BSA as a blocking

buffer, and mouse anti-goat IgG (H+L) (Jackson Immuno Research, 205-005-108) and goat

anti-mouse (Fc) labelled with horseradish peroxidase (Sigma, A0168) as primary and secondary

antibodies. The antibodies were diluted in blocking buffer at 1:2000 and 1:70,000, respectively,

and incubated with the membrane for 1 hour each. In between the antibody incubations, the

membrane was washed once with PBS – 0.1% Tween-20 + 3.2% NaCl and twice with PBS –

0.1% Tween-20, for 10 minutes each time. The results were visualized using the Fujifilm LAS-

3000 Luminescent Image Analyzer.

Serum samples derived from the *in vivo* challenge were diluted in water to achieve a total load

of 10-20 μg of protein per sample. Western Blots were performed and analyzed using the same

protocol as described above.

Animal experiment setup

All protocols of this study were designed and performed in strict accordance with the Kenyan and US American legislation for animal experimentation and were approved by the institutional animal care and use committee at both institutions (JCVI and ILRI, IACUC reference number 2014.08).

Sixteen male outbred goats (*Capra aegagrus hircus*), 1-2 years of age and randomly selected in Naivasha, were transferred to the ILRI campus in Nairobi and kept under quarantine for 6 months. After arrival at the campus, all animals were dewormed twice using levamisole and treated prophylactically against babesiosis and anaplasmosis using imidocarb. Upon entry to ILRI, the goats were vaccinated against anthrax & blackleg (Blanthax®, Cooper), Foot and Mouth Disease (FOTIVAX®) and Peste des Petits Ruminants (Live attenuated strain Nig. 75/1). All animals were tested negative for presence of antibodies against CCPP, using a competitive ELISA (IDEXX). Two weeks before experimental infection, all animals were transferred to the animal biosafety level two (ABSL2) unit. *Mycoplasma* cells were cultivated in PPLO medium supplemented with horse serum (45) to early logarithmic phase, aliquoted and stored at -80°C. Afterwards, we determined the CFU using two aliquots. Just before infection we thawed the vials and adjusted the concentration of *Mycoplasma* to 10^9CFU per mL^{-1} using broth. All 16 goats were infected transtracheally by needle puncture 5-10 cm distal to the larynx. Each animal received 1 mL of *Mmc* GM12 or GM12::YCpMmyc1.1-*Δ68* liquid culture (10^9 colony forming units per animal), followed by 5 mL of phosphate buffered saline (PBS). The animals were allowed to move freely within the ABSL2 unit and had *ad libitum* access to water. They were fed *ad libitum* with hay and received pellets each morning. Three veterinarians monitored the health status of the animals throughout the experiment. Rectal temperature, oxygen blood saturation, heart rate & breathing frequency were measured daily in the morning hours using the GLA M750 thermometer (GLA agricultural electronics, USA), VE-H100B oximeter (Edan, USA), and a stethoscope classic II (Littmann, USA) with a water-

resistant wrist watch Seamaster (Omega, Switzerland), respectively. Blood samples for

subsequent analysis were taken twice a week by jugular vein puncture. Nasal swabs (Flocked

swabs, Copan, Italy) were taken twice a week. Swabs were transferred into cryotubes filled

with media and stored at -80°C until further processing. Goats were euthanized when they

developed severe disease associated with unwarranted moderate to severe pain. Therefore, they

received an intravenous injection of Lethabarb Euthanasia Injection (Virbac, USA) of 200

mg.kg^{-1} body weight. Severe disease and pain were determined by a fever of $\geq$41ºC for >3

consecutive days, an oxygen saturation of $\leq$92% and a lateral recumbency of $\geq$1 day without

the ability to feed or intake water. Goats that were not put down because of ethical reasons were

euthanized on 28 dpi.

Pathomorphological and histology analysis

A complete necropsy was performed on all animals. Tissue samples of the neck region around

the inoculation site and all internal organs were fixed in 10 % buffered formalin for 72 hours

and subsequently routinely processed for paraffin embedding. Tissue sections were cut at 3 μm

and stained routinely with hematoxylin and eosin (H&E) and evaluated by a board-certified

pathologist.

Microbiology

Venous blood samples, lung samples, carpal joint fluid, and pleural fluid specimens taken at

necropsy were used for isolation of *Mmc* as described elsewhere (46) using *Mycoplasma* liquid

medium (Mycoplasma Experience, Ltd., United Kingdom). Lung samples and pleural fluid

were used for screening of *Pasteurella* and *Mannheimia* spp. using standard methods (47).

Statistical analysis

Exact and normal approximation binomial tests were used to compare the two groups using

GenStat 12th Edition (48). P values for differences in parameters were estimated using a 2-

sided 2-sample t-test comparing average levels between both groups at 5% level of significance.

ACKNOWLEDGMENTS

The National Science Foundation [grant number IOS-1110151 (to S.V., C.L., and J.J.)] provided the funds for this work. Additional support was received from the University of Bern and the CGIAR research program on Livestock and Fish. Elise Schieck was supported by the German Federal Ministry for Economic Cooperation and Development (Project No. 09.7860.1-001.00 Contract no. 81136800, http://www.bmz.de). Anne Liljander was supported by the Centrum for International Migration and Development in Germany. We thank Ray-Yuan Chuang, Vladimir Noskov, Steven Weber, Pamela Nicholson, Helga Mogel, Elisabeth Cook and the ILRI animal caretakers for their excellent advice and help.

AUTHOR CONTRIBUTIONS

J.J. and S.V. designed research; L.M., N.G-A., S.C., J.J., A.L., P.S., M.S., E.S., V.C.,Y.A., H.P. and F.L. performed research; J.J., F.L., S.C., Y.A., L.F., P.S-P., C.L., A.B. and SV., analyzed data; J.J. and S.V. wrote the paper

REFERENCES

506   1.   Linchevski I, Klement E, Nir-Paz R. 2009. *Mycoplasma pneumoniae* vaccine protective

507   efficacy and adverse reactions--Systematic review and meta-analysis. Vaccine 27:2437-

508   46.

509   2.   Browning GF, Whithear KG, Geary SJ. 2005. Vaccines to control Mycoplasmosis, p

510   569-597. *In* Blanchard A, Browning G (ed), Mycoplasmas : molecular biology,

511   pathogenicity and strategies for control. Horizon Bioscience, Wymondham, Norfolk,

512   U.S.A.

513   3.   Jores J, Mariner JC, Naessens J. 2013. Development of an improved vaccine for

514   contagious bovine pleuropneumonia: an African perspective on challenges and

515   proposed actions. Vet Res 44:122.

516   4.   Nicholas R, Churchward C. 2012. Contagious caprine pleuropneumonia: new aspects

517   of an old disease. Transbound Emerg Dis 59:189-96.

518   5.   Maes D, Segales J, Meyns T, Sibila M, Pieters M, Haesebrouck F. 2008. Control of

519   *Mycoplasma hyopneumoniae* infections in pigs. Veterinary microbiology 126:297-309.

520   6.   Chopra-Dewasthaly R, Spergser J, Zimmermann M, Citti C, Jechlinger W, Rosengarten

521   R. 2017. Vpma phase variation is important for survival and persistence of Mycoplasma

522   agalactiae in the immunocompetent host. PLoS Pathog 13:e1006656.

523   7.   Citti C, Nouvel LX, Baranowski E. 2010. Phase and antigenic variation in

524   mycoplasmas. Future Microbiol 5:1073-85.

525   8.   Chambaud I, Wroblewski H, Blanchard A. 1999. Interactions between mycoplasma

526   lipoproteins and the host immune system. Trends Microbiol 7:493-499.

527   9.   Arfi Y, Minder L, Di Primo C, Le Roy A, Ebel C, Coquet L, Claverol S, Vashee S, Jores

528   J, Blanchard A, Sirand-Pugnet P. 2016. MIB-MIP is a mycoplasma system that captures

529   and cleaves immunoglobulin G. Proc Natl Acad Sci U S A 113:5406-5411.

10.   Blotz C, Stulke J. 2017. Glycerol metabolism and its implication in virulence in

Mycoplasma. FEMS Microbiol Rev 41:640-652.

11.   Pilo P, Frey J, Vilei EM. 2007. Molecular mechanisms of pathogenicity of *Mycoplasma*

mycoides subsp. *mycoides* SC. Vet J 174:513-21.

12.   Szczepanek SM, Boccaccio M, Pflaum K, Liao X, Geary SJ. 2014. Hydrogen peroxide

production from glycerol metabolism is dispensable for virulence of *Mycoplasma*

gallisepticum in the tracheas of chickens. Infect Immun 82:4915-20.

13.   Lartigue C, Vashee S, Algire MA, Chuang RY, Benders GA, Ma L, Noskov VN,

Denisova EA, Gibson DG, Assad-Garcia N, Alperovich N, Thomas DW, Merryman C,

Hutchison CA, 3rd, Smith HO, Venter JC, Glass JI. 2009. Creating bacterial strains

from genomes that have been cloned and engineered in yeast. Science 325:1693-6.

14.   Browning GF, Marenda MS, Noormohammadi AH, Markham PF. 2011. The central

role of lipoproteins in the pathogenesis of mycoplasmoses. Veterinary microbiology

153:44-50.

15.   Beven L, Charenton C, Dautant A, Bouyssou G, Labroussaa F, Skollermo A, Persson

A, Blanchard A, Sirand-Pugnet P. 2012. Specific evolution of F1-like ATPases in

mycoplasmas. PLoS One 7:e38793.

16.   Tardy F, Mick V, Dordet-Frisoni E, Marenda MS, Sirand-Pugnet P, Blanchard A, Citti

C. 2015. Integrative conjugative elements are widespread in field isolates of

Mycoplasma species pathogenic for ruminants. Appl Environ Microbiol 81:1634-43.

17.   Krasteva I, Liljander A, Fischer A, Smith DG, Inglis NF, Scacchia M, Pini A, Jores J,

Sacchini F. 2014. Characterization of the *in vitro* core surface proteome of *Mycoplasma*

mycoides subsp. *mycoides*, the causative agent of contagious bovine pleuropneumonia.

Vet Microbiol 168:116-23.

18. Weldearegay YB, Pich A, Schieck E, Liljander A, Gicheru N, Wesonga H, Thiaucourt

F, Kiirika LM, Valentin-Weigand P, Jores J, Meens J. 2016. Proteomic characterization

of pleural effusion, a specific host niche of *Mycoplasma mycoides* subsp. *mycoides* from

cattle with contagious bovine pleuropneumonia (CBPP). J Proteomics 131:93-103.

19. Jores J, Schieck E, Liljander A, Sacchini F, Posthaus H, Lartigue C, Blanchard A,

Labroussaa F, Vashee S. 2018. Capsular polysaccharide in *Mycoplasma mycoides* is a

virulence factor. J Infect Dis doi:10.1093/infdis/jiy713.

20. Falkow S. 1988. Molecular Koch's postulates applied to microbial pathogenicity. Rev

Infect Dis 10 Suppl 2:S274-6.

21. Monnerat MP, Thiaucourt F, Poveda JB, Nicolet J, Frey J. 1999. Genetic and serological

analysis of lipoprotein LppA in *Mycoplasma mycoides* subsp. *mycoides* LC and

Mycoplasma mycoides subsp. *capri*. Clin Diagn Lab Immunol 6:224-30.

22. Dedieu L, Totte P, Rodrigues V, Vilei EM, Frey J. 2010. Comparative analysis of four

lipoproteins from *Mycoplasma mycoides* subsp. *mycoides* Small Colony identifies

LppA as a major T-cell antigen. Comp Immunol Microbiol Infect Dis 33:279-90.

23. Mulongo M, Frey J, Smith K, Schnier C, Wesonga H, Naessens J, McKeever D. 2015.

Vaccination of cattle with the N terminus of LppQ of *Mycoplasma mycoides* subsp.

mycoides results in type III immune complex disease upon experimental infection.

Infection and immunity 83:1992-2000.

24. Hutchison CA, 3rd, Chuang RY, Noskov VN, Assad-Garcia N, Deerinck TJ, Ellisman

MH, Gill J, Kannan K, Karas BJ, Ma L, Pelletier JF, Qi ZQ, Richter RA, Strychalski

EA, Sun L, Suzuki Y, Tsvetanova B, Wise KS, Smith HO, Glass JI, Merryman C,

Gibson DG, Venter JC. 2016. Design and synthesis of a minimal bacterial genome.

Science 351:aad6253.

25.   Schieck E, Lartigue C, Frey J, Vozza N, Hegermann J, Miller RA, Valguarnera E,

Muriuki C, Meens J, Nene V, Naessens J, Weber J, Lowary TL, Vashee S, Feldman

MF, Jores J. 2016. Galactofuranose in *Mycoplasma mycoides* is important for

membrane integrity and conceals adhesins but does not contribute to serum resistance.

Mol Microbiol 99:55-70.

26.   Richter SH, Garner JP, Auer C, Kunert J, Wurbel H. 2010. Systematic variation

improves reproducibility of animal experiments. Nat Methods 7:167-8.

27.   Mogensen TH. 2009. Pathogen recognition and inflammatory signaling in innate

immune defenses. Clin Microbiol Rev 22:240-73.

28.   Hirayasu K, Saito F, Suenaga T, Shida K, Arase N, Oikawa K, Yamaoka T, Murota H,

Chibana H, Nakagawa I, Kubori T, Nagai H, Nakamaru Y, Katayama I, Colonna M,

Arase H. 2016. Microbially cleaved immunoglobulins are sensed by the innate immune

receptor LILRA2. Nat Microbiol 1:16054.

29.   Abu Kwaik Y, Bumann D. 2013. Microbial quest for food *in vivo*: 'nutritional virulence'

as an emerging paradigm. Cell Microbiol 15:882-90.

30.   Fischer A, Shapiro B, Muriuki C, Heller M, Schnee C, Bongcam-Rudloff E, Vilei EM,

Frey J, Jores J. 2012. The Origin of the '*Mycoplasma mycoides* Cluster' Coincides with

Domestication of Ruminants. PLoS One 7:e36150.

31.   DaMassa AJ, Brooks DL, Adler HE. 1983. Caprine mycoplasmosis: widespread

infection in goats with *Mycoplasma mycoides* subsp *mycoides* (large-colony type).

American journal of veterinary research 44:322-5.

32.   Noskov VN, Segall-Shapiro TH, Chuang RY. 2010. Tandem repeat coupled with

endonuclease cleavage (TREC): a seamless modification tool for genome engineering

in yeast. Nucleic acids research 38:2570-6.

33. Boeke JD, LaCroute F, Fink GR. 1984. A positive selection for mutants lacking

orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance.

Mol Gen Genet 197:345-6.

34. Chandran S, Noskov VN, Segall-Shapiro TH, Ma L, Whiteis C, Lartigue C, Jores J,

Vashee S, Chuang R. 2014. TREC-IN: gene knock-in genetic tool for genomes cloned

in yeast. BMC Genomics 15:1180.

35. Gibson DG, Glass JI, Lartigue C, Noskov VN, Chuang RY, Algire MA, Benders GA,

Montague MG, Ma L, Moodie MM, Merryman C, Vashee S, Krishnakumar R, Assad-

Garcia N, Andrews-Pfannkoch C, Denisova EA, Young L, Qi ZQ, Segall-Shapiro TH,

Calvey CH, Parmar PP, Hutchison CA, 3rd, Smith HO, Venter JC. 2010. Creation of a

Bacterial Cell Controlled by a Chemically Synthesized Genome. Science 329:52-56.

36. Gietz D, St Jean A, Woods RA, Schiestl RH. 1992. Improved method for high efficiency

transformation of intact yeast cells. Nucleic Acids Res 20:1425.

37. Noskov V, Kouprina N, Leem SH, Koriabine M, Barrett JC, Larionov V. 2002. A

genetic system for direct selection of gene-positive clones during recombinational

cloning in yeast. Nucleic Acids Res 30:E8.

38. Labroussaa F, Lebaudy A, Baby V, Gourgues G, Matteau D, Vashee S, Sirand-Pugnet

P, Rodrigue S, Lartigue C. 2016. Impact of donor-recipient phylogenetic distance on

bacterial genome transplantation. Nucleic Acids Res 44:8501-11.

39. Fischer A, Santana-Cruz I, Hegerman J, Gourle H, Schieck E, Lambert M, Nadendla S,

Wesonga H, Miller RA, Vashee S, Weber J, Meens J, Frey J, Jores J. 2015. High quality

draft genomes of the *Mycoplasma mycoides* subsp. *mycoides* challenge strains Afadé

and B237. Stand Genomic Sci 10:89.

40. Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler

transform. Bioinformatics 26:589-95.

41.   Li H. 2011. Improving SNP discovery by base alignment quality. Bioinformatics

27:1157-8.

42.   Rausch T, Zichner T, Schlattl A, Stutz AM, Benes V, Korbel JO. 2012. DELLY:

structural variant discovery by integrated paired-end and split-read analysis.

Bioinformatics 28:i333-i339.

43.   Zhang Z, Wang J, Luo J, Ding X, Zhong J, Wang J, Wu FX, Pan Y. 2016. Sprites:

detection of deletions from sequencing data by re-aligning split reads. Bioinformatics

32:1788-96.

44.   Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer

(IGV): high-performance genomics data visualization and exploration. Brief Bioinform

14:178-92.

45.   Sacchini F, Naessens J, Awino E, Heller M, Hlinak A, Haider W, Sterner-Kock A, Jores

J. 2011. A minor role of CD4+ T lymphocytes in the control of a primary infection of

cattle with *Mycoplasma mycoides* subsp. *mycoides*. Vet Res 42:77.

46.   Liljander A, Yu M, O'Brien E, Heller M, Nepper JF, Weibel DB, Gluecks I, Younan M,

Frey J, Falquet L, Jores J. 2015. A field-applicable recombinase polymerase

amplification assay for rapid detection of *Mycoplasma capricolum* subsp.

capripneumoniae. J Clin Microbiol 53:2810-5.

47.   Carter GR, Cole JR (ed). 1990. Diagnostic procedures in veterinary bacteriology and

mycology. Academic Press, San Diego, Calif.

48.   Payne RW, Murray DA, Harding SA, Baird DB, Soutar DM. 2012. *GenStat for*

Windows (15th Edition) Introduction., VSN International, Hemel Hempstead, UK.

Table 1: Summary of post mortem records of goats

Animal No.	Date of euthanasia	Bacteremia ccu/ml	Inflammation of the neck around the injection site	Pulmonary congestion	Pulmonary oedema	Congested kidneys	Mucoid enteritis & congestion	Pleural fluid in thoracic cavity	Adherent lung to rib cage	Liver abscess
CK032	6 dpi	10^7	X	X	X	X	X	X		
CK034	5 dpi	10^7	X	X	X	X	X			
CK035	29 dpi									
CK040	5 dpi	10^8	X	X	X	X	X			X
CK043	5 dpi	10^8	X	X	X	X	X			
CK045	28 dpi									
CK046	5 dpi	10^8	X	X	X	X	X			
CK047	29 dpi									
CK048	5 dpi	10^9	X	X	X	X	X			
CK049	28 dpi									
CK051	5 dpi	10^9	X	X	X	X	X			
CL001	28 dpi									
CL002	5 dpi	10^8	X	X	X	X	X		X	
CL003	28 dpi									

Animals displayed in bold were infected with GM12::YCpMmyc1.1-*Δ68*, the other animals were infected with wtGM12; dpi - days post infection,

ccu – color changing unit

Table 2: Table showing the results of the Illumina sequencing-based mapping assemble of the strains used in this study

Region (size in bp)	YCP (9,192)	D1 (2,984 bp)	D2 (4,950 bp)	D3 (4,677 bp)	D4 (69,220 bp)	D5 (24,906 bp)
GM12	Not present	Present (I)	Present (I)	Present (I)	Present (I)	Present (I)
GM12::YCpMmyc1.1 (position)	Present (25,426-34,618)	Present (283,950-286,934)	Present (569,098-574,048)	Present (18,716-23,393)	Present (665,418-734,638)	Present (1,055,169-1,080,075)
GM12::YCpMmyc1.1-*Δ68*	Present (I)	Deleted (DIS)	Deleted (IS)	Deleted (DIS)	Deleted (DIS)	Deleted (DIS)

Legend: D = Detected by Delley; I = Validated visually by IGV; S = Detected by Sprites

Table S1: Oligonucleotide primers used to construct the mutagenesis cassettes and to confirm the deletions

Primer Name	Primer sequence (5'→ 3')	PCR product and notes
D1: Deletion using TREC method (modified CORE3 cassette):		
FKO-F1	**GTT GAT TCA GTC TTC ATA ACG TTT TTT ACC TAA AAC ATT TAA TTT TGC ATT** *TAG GGA TAA CAG GGT AA TAC GGA TTA G*	To amplify the 2.4 kb CORE3 mutagenesis cassette with 50 bp homology overhangs at the target site.
FKO-R	**AAT GCA AAA TTA AAT GTT TTA GGT AAA AAA CGT TAT GAA GAC TGA ATC AA***T CGG TAC ATA AAT ATA TGT GAT TCT*	
FKO-F2	CTG AAG ATT TAG CAG CAG TTA ATC CTC AAA TAC CAA CTA ATA GAA CTA CAG **TTG ATT CAG TCT TCA TAA CGT TT**	
FKO-DG-F	CCA GTT GTT CCA CCT AAT G	To amplify 235 bp with RC0421 to confirm the left junction of the glpFKO-CORE3 insertion. To amplify 250 bp with FKO-DG-R to confirm pop-out and seamless deletion of the glpFKO genes.
RC0421	CTT CGG AGG GCT GTC ACC	
FKO-DG-R	TCG CAT TAG AAA CAA GTA GTC	To amplify 283 bp with RC0430 to confirm the right junction of the glpFKO-CORE3 insertion.
RC0430	AAG TGT CAC CAT GAA CGA CA	
D2: Deletion using TREC method (Cre/ URA3 cassette):		
RC0905	CTA ATT AAT ATA AAT GAA AGA ATT AAA	To amplify the 2.4 kb Cre-Ura3 mutagenesis cassette with 50 bp homology overhangs at the target site.
RC0906	TTA AAG AAT TAT TAC TTG GTA TAT TA	
RC0350	CGG CTA AAG CTA AAA TCA AAA GAA C	To amplify 615 bp to confirm the left junction of the gts-CRE/URA3 insertion. To amplify 650 bp with RC0350 and RC0332 to confirm pop-out and seamless deletion of the gts-5kb region.
RC0373	ATA TTT GAG AAG ATG CGG CCA GC	
RC0332	AAG AGC ATA TTA GTA AAA TAT TGT CA	
D3: Deletion using TREC method (CORE3 cassette):		
lppA123-F1	**TAA AAC TGG AAA AAA CTA TGA AAA AAG CAA TTA AAT TAT TAC TAT CTA TT***T AGG GAT AAC AGG GTA ATA CGG ATT A*	To amplify the 2.5 kb CORE3 mutagenesis cassette with 50 bp homology overhangs at the target site.

lppA123-R	**AAT AGA TAG TAA TAA TTT AAT TGC TTT TTT CAT AGT TTT TTC CAG TTT TA**T CGG TAC ATA AAT ATA TGT GAT TCT	
LppA123-F2	TAG CTA TGT TGT TAT AAT ATT TTC GAA AAA TTT TTT GTA GGT ATA TTA TT**T AAA ACT GGA AAA AAC TAT GAA AAA AG**	
lppA123-DG-F	CTA GAT GTA AAT TAC CAC TAG G	To amplify 360 bp with RC0421 to confirm the left junction of the lppA-CORE3 insertion. To amplify 436 bp with lppA-DG-R to confirm pop-out and seamless deletion of the lppA region.
lppA123-DG-R	GGT TTT GAT GGT GTT TTT GGA G	To amplify 237 bp with RC0430 to confirm the left junction of the lppA-CORE3 insertion.
D4: Deletion using TREC method (CORE3 cassette):		
50CORE-F1	**TTTTAATCCTCCAACCTATTAATATTTTAAATAAGAT AAAACATTGTTGG**TAGGGATAACAGGGTAATACGGATTA G	To amplify the 2.5 kb CORE3 mutagenesis cassette with 50 bp homology overhangs at the target site.
50CORE-R	CCAACAATGTTTTATCTTATTTAAAATATTAATAGGTTG GAGGATTAAAATCGGTACATAAATATATGTGATTCTG	
50CORE-F2	AGTATAAAACTTTTATCCTAACCGATTTTAAGTTTTATA CAGGAGGAATT**TTTTAATCCTCCAACCTATTAATATTT**	
RC0863	TCAACATATTCTGGTATGTCT	To amplify 425 bp with RC0421 to confirm the left junction of the ICE-CORE3 insertion. To amplify 480 bp with RC0878 to confirm pop-out and seamless deletion of the ICE region.
RC0878	TAATCTAACACAACCGGTAG	To amplify 325 bp with RC0430 to confirm the left junction of the ICE-CORE3 insertion.
D5: Deletion using TREC method (CORE3 cassette):		
TREC37-F1	**TAATTCATTAGTCTCTAATTCTTATTAAAAGATTTAG AGATTTTTTATTT**TAGGGATAACAGGGTAATACGGATTAG	To amplify the 2.5 kb CORE3 mutagenesis cassette with 50 bp homology overhangs at the target site.
TREC37-R	**AAATAAAAAATCTCTAAATCTTTTAATAAGAATTAGA GACTAATGAATTA**TCGGTACATAAATATATGTGATTCT	
TREC37-F2	CTAGAAATATATTAATATATCATTAAAAATAAAAATAG CATTTAAGATT**TAATTCATTAGTCTCTAATTCTTATTA A**	

TREC37 -DG-F	TGGTGTTGCTACTGAAATATG	To amplify 250 bp to confirm pop-out and seamless deletion of the D5 region.
TREC37 -DG-R	ACTTAAAAAAGAACGACACCG	

Figure S1: Specific PCRs confirming the deletions in GM12::YCpMmyc1.1-*Δ68*. Genomic DNA from GM12::YCpMmyc1.1-*Δ68* (clone 1711) and its parental strain GM12 was used as template. Specific primers were used to amplify across the deleted regions. DNA from GM12::YCpMmyc1.1-*Δ68* (clone 1711) yielded expected amplicons that are a product covering the flanking regions of the deleted subgenomic fragment. Diagnostic primers FKO-DG-F and FKO-DG-R result in a 250 bp amplicon across the D1 region in clone 1711, primers RC0350 and RC0332 result in a 650 bp amplicon across the D2 region, primers LppA123-DG-F and LppA123-DG-R result in a 436 bp amplicon across the D3 region, primers RC0863 and RC0878 result in a 480 bp amplicon across the D4 region, and primers TREC37-DG-F and TREC37-DG-R result in a 250 bp region across the D5 region. L denotes the DNA ladder.