ENVIRONMENTAL PERTURBATIONS AND TRANSITIONS BETWEEN

ECOLOGICAL AND EVOLUTIONARY EQUILIBRIA: AN ECO-EVOLUTIONARY

FEEDBACK FRAMEWORK

Tim Coulson

Department of Zoology, University of Oxford, The Zoology Research and Administration

Building, 11a Mansfield Road, Oxford, OX1 3SZ, UK

Email: tim.coulson@zoo.ox.ac.uk

Abstract

I provide a general framework for linking ecology and evolution. I start from the fact that

individuals require energy, trace molecules, water, and mates to survive and reproduce, and

that phenotypic resource accrual traits determine an individual's ability to detect and acquire

these resources. Optimum resource accrual traits, and their values, are determined by the

dynamics of resources, aspects of the environment that hinder resource detection and

acquisition by imposing risks of mortality and reproductive failure, and the energetic costs of

developing and maintaining the traits – part of an individual's energy budget. These budgets

also describe how individuals utilize energy by partitioning it into maintenance, development

and/or reproduction at each age and size, age and size at sexual maturity, and the size and

number of offspring produced at each reproductive event. The optimum energy budget is

consequently determined by the optimum life history strategy that describes how resources are

utilized to maximize fitness by trading off investments in maintenance, development, and

reproductive output at each age and size. The optimum life history in turn determines body

size. An eco-evolutionary feedback loop occurs when resource accrual traits evolve to impact

the quality and quantity of resources that individuals accrue, resulting in a new optimum life

history strategy and energy budget required to deliver it, a change in body size, and altered

population dynamics that, in turn, impact the resource base. These feedback loops can be

complex, but can be studied by examining the eco-evolutionary journey of communities from

one equilibrium state to another following a perturbation to the environment.

KEYWORDS: body size evolution, dynamic energy budgets, global environmental change,

integral projection models, life history theory, paradox of stasis, quantitative genetics,

stochastic demography, Trinidadian guppy.

Introduction

In the lowland streams of Northern Trinidad, guppies (*Poecilia reticulata*) live in complex,

stable, communities (Reznick and Endler 1982). They predominantly feed on abundant stream

invertebrates of high nutritional quality that are also food for a number of other fish species.

Guppies themselves are prey for predatory fishes, such as wolf fish (*Hoplias malabaricus*) and

pike cichlids (*Crenicichla frenata*). Predation is the primary cause of death for guppies living

in these communities, such that their populations are limited by predation. In response to this

limitation, guppies have evolved streamlined bauplans, live in shoals, have high maximum

swimming speeds, and dull coloration to aid with the acquisition of resources in a predator-

rich environment. They have also evolved fast metabolic rates, small sizes and low ages at

sexual maturity, large litters of small offspring, and fast life histories to rapidly utilize the

energy accrued (Reznick and Endler 1982, Reznick and Bryga 1987, Reznick and Yang 1993,

Reznick et al. 2001, Travis et al. 2014). Evolution has consequently enabled these guppies to

optimally detect, acquire, and utilize food in an environment where the risk of death in the jaws

of a predator is near constant and high.

Above waterfalls, on the same streams where high-predation guppies are found, predatory fish

are largely absent. Guppies in these streams live at high density in simpler, stable, communities,

with only one major competitor, Hart's killifish (*Rivulus hartii*), which is also an intra-guild

predator. The guppy populations are food-limited, the limiting resource is therefore food

availability, and the primary cause of death and failure to breed is starvation. A consequence

of this is that high-quality invertebrates are rapidly consumed and are largely absent, so that

guppy diets contain a high percentage of low-quality food such as algae and bacteria, that are

largely absent from the diets of high-predation guppies. Guppies adapted to these low predation

environments have evolved brighter male coloration, a less-streamlined bauplan, reduced

shoaling behaviour, altered mouth morphology, a lower metabolic rate, greater age and larger

size at sexual maturity, smaller litter sizes, larger offspring and a slower life history (Reznick

and Endler 1982, Reznick and Bryga 1987, Reznick and Yang 1993, Reznick et al. 2001, Travis

et al. 2014). These traits have evolved to enable guppies to detect, acquire, and utilize food in

a relatively safe environment compared to the high predation one, but where the major

challenge is securing enough resources to thrive.

When high predation guppies are introduced to low predation environments, their population

increases in size, the age, sex and phenotypic trait structure of the population changes, and they

rapidly evolve, with the community moving from the high predation equilibrium to the low

predation one. Because these changes are repeated across streams, they are predictable in that

we know what will happen when guppies are released from predation (Reznick and Endler

1982, Reznick and Bryga 1987, Reznick and Yang 1993, Reznick et al. 2001, Travis et al.

2014).

What characterises the Trinidadian freshwater ecosystems when they are at ecological and

evolutionary (quasi)-equilibrium in either the high- or low-predation state? The size and

genetic-, phenotypic trait-, size-, age-, and sex- structure of all populations in the community

show no persistent temporal trends with time. I consequently use the term 'equilibrium' to

describe a stochastic but stationary state of populations and the community. This means there

may be temporal fluctuations in the sizes and structures of populations within the community,

but none will exhibit a long-term temporal trend (Lundberg, Ranta, and Kaitala 2000). I use

the prefix 'quasi' to acknowledge that some slow temporal patterns in genotype frequencies

may occur in cases of slow co-evolution between species, but, that in general, natural systems

frequently achieve states where statistics describing each population within the community

remain within bounds for protracted periods of time (Figure 1; (Lewontin 1969, Bronstein et

al. 2004). A consequence of this is there will also be no persistent temporal trends in

community size and structure, or energy and nutrient flows.

The existence of a (quasi)-equilibrium state raises a paradox. When systems are in (quasi)-

equilibrium states, all heritable phenotypic traits in all species are expected to be at, or very

close to, their optimal values, as otherwise phenotypic evolution would be expected to occur

and the system would not be at an evolutionary equilibrium. When I refer to phenotypic traits,

I define them as any attribute that can be characterised at the individual level, ranging from

molecules within cells to lifetime reproductive success. In the wild, many traits including

measures of body size, lifetime reproductive success, and those that describe the timing of

events such as birth, are under directional selection, are heritable, but do not evolve as predicted

(Merilä et al. 2001a, Estes and Arnold 2007, Haller and Hendry 2014, Rollinson and Rowe

2015, Bonamour et al. 2017). Instead they exhibit stasis that is consistent with the system being

at equilibrium. The paradox is why do we predict phenotypic evolution but not see it? I address

this paradox later in the paper.

An environmental perturbation, such as the loss of a predator species, leads to a period of

transience within a community as population sizes and structures change (Hastings 2004,

Georgelin et al. 2015). In time, the system will achieve a new (quasi)-equilibrium state (Figure

1) (Brown and Vincent 1992, Lundberg et al. 2000, Johansson and Dieckmann 2009) The

observation of almost immediate evolution following a biotic or abiotic shift raises a second

paradox. Evolution requires additive genetic variation within a population, but if phenotypic

traits are at their optimum values, they are subject to stabilising selection, which is expected to

erode additive genetic variation (Crow and Kimura 1970). The evolutionary equilibrium

attained by a system should consequently exhibit low levels of additive genetic variance. Yet

populations in ecosystems in the wild in (quasi)-equilibrium states express substantial amounts

of additive genetic variance for many traits (Crow and Kimura 1970, Kruuk et al. 2000). The

maintenance of genetic variance is a second paradox that requires a solution if we are to

understand why perturbed systems move from one evolutionary equilibrium to another. I also

consider this paradox later in the paper.

The genetic diversity paradox has similarities with another problem that has long interested

biologists – why do so many species coexist when they appear to share niches and rely upon

the same resources (Chesson 1986, Wright 2002)? The coexistence problem is also relevant to

the problems I address in this paper, and in particular why the ecological and evolutionary

(quasi)-equilibria contain so many coexisting species. I return to this topic later in the paper.

Before reviewing properties of ecological and evolutionary equilibria, and considering ways to

predict the ecological and evolutionary transients between contrasting stationary states

following a biotic or abiotic change, I do two things. First, I consider how the different levels

of biological organisation – alleles, genotypes, phenotypes – are linked by a number of

fundamental processes (Figure 2) (Coulson et al. 2017).

Parental alleles combine, through patterns of mate choice, to generate the distribution of

offspring genotypes, which in turn develop into a distribution of offspring phenotypic traits.

These traits determine individual abilities to acquire resources from the ecological environment

in which they find themselves, which in turn determines continued rates of development and

chances of survival, reproduction and dispersal at each age and size. Both the population

dynamics, and the strength of selection that can alter the distribution of phenotypic traits,

genotypes and alleles, are inevitable emergent properties of these rates. Recombination and

mutation determine the distribution of alleles among individuals selected to be parents (Crow

and Kimura 1970). The fundamental biological processes of mate choice; development;

survival, reproduction, and dispersal; and mutation, can all be influenced by aspects of the

biotic and abiotic environment. The biotic environment experienced by an individual consists

of the numbers and attributes of all individuals it interacts with whether they are con- or hetero-

specifics. These interacting individuals include its parents, who can use their attributes to

influence offspring phenotypes via controlling the developmental environment.

Second, I introduce the concept of resource accrual traits. These are the set of traits associated

with the detection and acquisition of resources required for life (Abrams and Chen 2002):

energy, trace molecules and water required for survival, and mates necessary for reproduction

(at least in sexually reproducing diploid species) (Brown et al. 2004). Within the set of resource

accrual traits are resource detection traits such as the size and structure of sense organs and the

brain, and resource acquisition traits such as foraging behaviours and aspects of morphology.

I will argue that these traits are likely to be subject to stabilising selection, yet that individual

heterogeneity in the types of resources individuals specialise on (Bolnick et al. 2007), act to

maintain additive genetic variation underpinning these traits. Desirable resource accrual traits,

and their optimum values, are specific to the environment in which individuals find themselves.

For example, the traits expressed by high-predation guppies in high-predation environments

(Reznick and Endler 1982) are well-adapted to an environment where the limiting factor is

predation and the primary cause of death is predation. In contrast, in the low-predation

environments, guppies express traits that make them well-adapted to an environment where the

primary cause of death is starvation and the population is food-limited.

In addition to optimising resource accrual traits, evolution also optimizes energy utilisation to

maximise fitness (Kooijman and Kooijman 2010). It does this by optimising the life history via

evolving an energy budget to optimally allocate energy to maintenance, development, and

reproduction, at each age and size (Hin and de Roos 2019). The energy budget determines age

and size at sexual maturity, the number and size of offspring produced at each reproductive

attempt, and life expectancy (Stearns 1976).

I primarily focus on the guppies as they provide a well-characterised example of a general

phenomenon: if the primary causes of death or failures to breed in a population are altered by

a biotic or abiotic perturbation, so too is the limiting factor, the population dynamics (Schindler

1974, Sibly and Hone 2002, Rohr et al. 2003), selection, and the course of evolution (Walters

and Juanes 1993, Fishman and Willis 2008). Such a perturbation results in rapid change in

population size and selective regimes in a focal species and the species with which it interacts.

The new regime selects for phenotypic traits and trait values that allow organisms to optimally

detect and acquire energy, and a life history that allows these resources to be optimally utilised

given the primary causes of death and failures to breed in the new environment in which they

find themselves. These processes determine the transient dynamics between ecological and

evolutionary (quasi)-equilibria.

In the remainder of this paper I provide 1) a description of the conditions required for a system

to be at ecological and evolutionary equilibria, 2) discussion of how these conditions can be

achieved in nature, and the observation of two paradoxes that need to be addressed: the paradox

of stasis for size-related phenotypic traits, and the paradox of standing additive genetic

variation, 3) solutions to the paradoxes, 4) a more detailed treatment of the evolution of life

history and body size, which I consider to be central to solving the paradox of stasis, 5) an

examination of models that can describe systems in ecological and evolutionary equilibrium,

6) a focus on how systems transition between different equilibria states as a consequence of

abrupt environmental change, and 7) a brief concluding section.

Characterising Populations at Ecological and Evolutionary Equilibria

I start by considering a single population. I define a population as being stationary and at a

stochastic ecological and evolutionary equilibrium when both its size and genetic, phenotypic,

sex, and age structure fluctuate within bounds but show no persistent temporal trends (Caswell

et al. 2004). In formal demography, this equilibrium is a special case, as populations can

achieve structural stationarity, yet can also change in size with time as the population grows at

a constant stochastic rate (Tuljapurkar 1989). Because I require a stationary distribution of

population sizes N_t, the special case I consider requires the long-run stochastic growth rate a

to equal zero, where

$$a = \overline{\log(\lambda_t)} = \frac{1}{T}\sum_{t=1}^{T} \log(\lambda_t) \qquad (1)$$

with t being time (Tuljapurkar and Orzack 1980). The equation defines the long-run stochastic

growth rate of the population as the arithmetic mean of the logged per-time step population

growth rates λ_t calculated across a time series of length T where $\lambda_t = \frac{N_{t+1}}{N_t}$. But how does this

formulation relate to the ecological and evolutionary equilibria I describe above, where there

is also no persistent temporal trend in allele and genotype frequencies, and phenotypic trait

means and variances? Examining the determinants of λ_t helps answer this question.

I consider a population as consisting of individuals, each with a very large number of

phenotypic trait and genotypic attributes. From these individuals, I can construct a multivariate

distribution describing the number of individuals in each individual attribute (or structural)

class at time t (Ellner et al. 2016). For example, in a very simple case where a population is

structured by the individual attributes sex, age, weight, and genotype at a locus X, there may

be 23 individuals within the population at a particular point in time that are male, aged 35 years,

weigh 82kg and express genotype AA at locus X, but only seven individuals that are female,

aged 74, weigh 52kg with genotype BB at locus X. All other sex, age, weight and genotype

classes will also contain a particular number of individuals (that could be zero). Over time, the

numbers of individuals in each structural class can fluctuate (Ellner and Rees 2006), but as

long as there are no persistent temporal trends in these time series of numbers, the population

will be at a stationary equilibrium. It is possible to estimate the long-run stochastic growth rate

for each structural class using equation (1) above; at ecological and evolutionary equilibrium,

these rates will all be zero (Tuljapurkar and Orzack 1980).

The per-time step growth rate of the entire population $\lambda_t = \frac{N_{t+1}}{N_t}$ can be expressed as $\lambda_t = S_t +$

R_t where S_t is the annual per capita survival rate, and R_t is the annual per capita recruitment

rate (defined as the per capita production of offspring that are produced between times t and

$t + 1$ that survive to enter the population at time $t + 1$ weighted by the relatedness of offspring

to parents (Coulson et al. 2006)). If we ignore dispersal, change in population size over a time

step is consequently caused by any difference in the numbers of individuals that enter the

population via birth, and the numbers that leave it via death. But what determines S_t and R_t at

a particular time t? Both rates vary across structural classes (e.g. individual attributes), and

with the environment (Leslie 1945, Caswell 2001). The values of S_t and R_t at any particular

time t consequently depend upon the population's genetic, phenotypic trait, age and sex

structure.

Survival, development, reproduction, mutation, recombination, mate choice, and genetic and

non-genetic inheritance are all processes that can alter the population's structure between time

t to $t + 1$ (Coulson et al. 2017). With the exception of genetic inheritance, all these processes

can be influenced by aspects of the biotic or abiotic environment, and it is environmental

variation that continually generates variation in population structure via impacting these

processes (Figure 2). Structured demographic models based on Figure 2 can be constructed to

capture the effects of the environment on each of these processes, and used to iterate

distributions of population structure and size from one-time step to the next (Coulson et al.

2017).

Structured demographic models have been constructed for small numbers of individual

attributes such as one or two genetic loci (Charlesworth 1994), one or two phenotypic traits

(Easterling et al. 2000), age (Leslie 1945), or sex (Schindler et al. 2015). Despite a focus on

only a small number of attributes, these models have provided significant ecological and

evolutionary insight. Models consist of functions that capture the processes (Figure 2) that can

alter distributions of individual attributes by moving individuals into and out of each structural

class at a point in time (Ellner et al. 2016). For example, survival functions determine the

proportion of individuals within a class that remain within a population, development functions

determine the rates at which surviving individuals move between structural classes,

reproductive functions determine the expected number of offspring produced by individuals

within each structural class, while mutation, mate choice, and genetic and non-genetic

inheritance functions, determine the attributes of offspring and the structural classes in which

they enter the population. I cumulatively refer to survival and reproductive functions as

demographic functions, and mutation, mate choice, and genetic and non-genetic inheritance

functions as inheritance functions. Any function, with the exception of those that describe

genetic inheritance, can contain terms describing the influence of biotic and abiotic

environmental drivers (see section **Population models of ecological and evolutionary stasis**

for additional details).

Structured demographic models do not need to include all levels of biological organization

found within a population (Figure 2). For example, models have been constructed that structure

populations by only age, genotype, or phenotypic trait, or by just two of these three levels of

organization. When these simple models are constructed, they make assumptions about the

processes that link levels of biological organization that are not explicitly incorporated into

models (Traill et al. 2014). For example, models that include genotypes, but not phenotypes,

assume that each genotype always develops with perfect fidelity into the same phenotype trait

value, such that fitness can be directly predicted from the genotype without having to model

phenotypic development (Charlesworth 1994). In this paper I primarily focus on cases where

this assumption is not made, and populations are structured by alleles, genotypes and

phenotypic traits (Coulson et al. 2011, Coulson et al. 2017).

As previously mentioned, each genotype (and consequently allele) will have a long run

stochastic growth rate of zero at ecological and evolutionary equilibrium. There is another

equivalence that individuals in particular structural classes will share: when genotypes are

additive, and in the absence of mutation, genotypes within offspring cohorts are expected to

have identical reproductive values (Sæther and Engen 2015). The expected reproductive value

is the expected number of descendants of an individual in a particular structural class at time

t, at some distant time point in the future (Fisher 1930, Caswell 2001). There are two ways

expected reproductive value can be calculated for genotypes from a model. First, by keeping

track of the number of descendent copies of each allele within the population with time (Barton

and Etheridge 2011). Second, by recording the number and relatedness of descendants alive in

future time steps of individuals that carried a particular allele at time t (Fisher 1930).

Descendants are included in the calculation regardless of whether they carry alleles found in

the original genotype themselves; genetic inheritance in sexually reproducing species means

that descendants do not necessarily carry ancestral alleles. The two quantities are expected to

be equivalent, although stochasticity in survival, reproduction and inheritance can result in an

individual with a particular allele having very many descendants with none, or very few,

carrying copies of it. Reproductive value can be calculated for any structural class – genotype,

phenotype, age, sex. However, at evolutionary equilibria, reproductive values of offspring

genotype or phenotypic trait classes are only equivalent when genotypes and phenotypes are

additive functions of underlying alleles. This requires a little more explanation.

In structured models of sexually reproducing species, alleles do not form structural classes.

Instead, these classes consist of genotypes or phenotypic traits (Charlesworth 1994, Childs et

al. 2016, Coulson et al. 2017). The dynamics of alleles, and hence microevolution, can be

determined from the dynamics of the number of individuals within genotypic classes. A

genotype is considered additive, if its contribution to a phenotypic trait is determined solely by

the sum of the contribution of each of its alleles to a phenotypic trait (Falconer 1960). If a

phenotypic trait value is determined by an interaction between alleles at a locus (dominance)

or by interactions between genotypes at different loci (epistasis), the assumption of additivity

is violated. A consequence of this is that microevolution cannot be inferred from the dynamics

of phenotypic trait values.

If all genotypes at all loci that contribute to a phenotypic trait are additive, then the dynamics

of the phenotypic trait, the genotypes that contribute to the trait, and the alleles that constitute

the genotypes, are asymptotically predicted by their reproductive values (Fisher 1930). Under

these circumstances, when different genotypes and phenotypic trait values expressed in

offspring cohorts have different reproductive values, then microevolution will occur and allele

frequencies will change. When they all have identical reproductive values, the population is at

evolutionary equilibrium.

Expected reproductive value is a key concept in biology as it is the expected asymptotic fitness

of an individual with particular attributes (Fisher 1930). The reproductive value of a structural

class is determined by the life history of individuals with those attributes. A life history

describes key events within life – age and size at sexual maturity, the timing of reproductive

events, the size and number of offspring produced at each successful reproductive attempt, and

the age and size at death (Stearns 1976). Life history strategies can consequently be

characterized with size- and age-specific survivorship, development, fertility and inheritance

schedules (Figure 4) (Stearns 1976) – all of which can be calculated from the functions that

constitute structured demographic models (Childs et al. 2004).

A life history strategy plays out on a per generation time scale, and can be described with per

generation summary statistics such as generation length, mean lifetime reproductive success

R_0, and life expectancy (Caswell 2001, Tuljapurkar et al. 2009a, Steiner et al. 2014). R_0

describes the rate of population growth per generation and is a per generation version of λ_t. In

a deterministic population that exhibits no temporal trend in numbers $\log(R_0) = 0$.

R_0 can be calculated for individuals within each genotypic or phenotypic structural class

(Tuljapurkar et al. 2009b). When additivity is assumed, individuals in all structural classes will

have equal expected values of R_0 when the population is at ecological and evolutionary

equilibria. When the assumption of additivity is violated, this will not necessarily be the case.

Biologists are often interested in the evolution of life histories, without focusing on evolution

of the resource accrual traits that influence energy acquisition and the optimal life history

(Stearns 1976). To do this, life history strategies are assumed to be clonally and near perfectly

inherited, rather than as arising as a consequence of underlying resource accrual traits (Childs

et al. 2004). Under this assumption, the reproductive value of a life history will be associated

with its long-run stochastic growth rate when measured in the presence of competing, clonally

inherited, life histories. R_0 can usually (but not always) also predict which clonally inherited

life history will grow to dominate all others. Such approaches rely on the assumption of clonal

inheritance. The approach is useful as invasion approaches can be used to identify the

evolutionarily stable life history strategy, and in particular the optimal value of life history

trade-offs such as those between offspring size and offspring number. However, genetic

architecture significantly influences ecological and evolutionary dynamics (Schreiber et al.

2018), except in very specific cases (Geritz and Éva 2000, Metz and de Kovel 2013), and I

consequently focus on cases where inheritance is not assumed to be clonal.

The conditions for a stochastic ecological and evolutionary equilibrium within a population –

a long-run stochastic growth rate of zero for each structural class – are relatively

straightforward, but they impose constraints on the values of reproductive value and the life

history when additivity is assumed. Structured demographic models of populations at

stochastic ecological and evolutionary equilibrium can be constructed from demographic,

development, and inheritance functions that describe the rates of movement of individuals

between structural classes from time t and $t + 1$. From these models, it is then possible to

calculate long-run stochastic growth rates, reproductive values, temporal dynamics of the

number of individuals, and descriptors of the life history for each class (Coulson et al. 2011,

Coulson et al. 2017).

Structured demographic models provide a remarkably powerful tool to study the dynamics of

populations, allele and genotype frequencies, and phenotypic trait distributions. The properties

of models at ecological and evolutionary equilibria are also well understood. However, for

these equilibria to be achieved, particular conditions are necessary.

Conditions Required for Ecological and Evolutionary Stationarity

A long-run stochastic growth rate of $a = 0$ in a single species population requires density-

dependence: a pattern that describes a decline in the per time step population growth rate λ_t as

population size increases (Engen and Sæther 1998). Density-dependence occurs when a direct

species interaction, such as predation, herbivory, or rarity of a resource species, limits a

population's growth rate (Hassell 1975). Populations are consequently limited by aspects of

the environment that prevent organisms detecting, acquiring and utilizing sufficient resources

to survive and reproduce (Murray 1982). Limiting factors determine key causes of mortality or

failures to breed within a population (Stearns 1976, Gaillard et al. 2000). For example, prey

species that are predator-limited are limited by their ability to accrue resources while avoiding

predation.

Ecologists have long been interested in how species that rely upon the same resources or

limiting factor can coexist (Gause 1932). The general solution to this problem is that

coexistence of competing species occurs when individuals within a species compete more

strongly for a resource than individuals between species. In contrast, if interspecific

competition is stronger than intraspecific competition, then the better competitor species will

drive the other extinct (MacArthur and Levins 1967).

Four different mechanisms can contribute to the relative strengths of intra- (density

dependence) and inter-specific competition for a resource or limiting factor. They act by either

increasing the strength of intraspecific competition or by reducing the strength of interspecific

competition (Chesson 2000b). The first mechanism, termed density-fitness covariance, does

the former. It describes any mechanism that acts to increase the strength of intraspecific

competition relative to interspecific competition, such that each population increases in density

when its numbers drop (Chesson 2000a). An example of this is when competing species utilise

the same resource, but are segregated in space, such that they tend to compete with conspecifics

rather than with hetero-specifics. The second set of mechanisms is termed "stabilising" and

works by decreasing the strength of interspecific competition by ensuring each species has a

refuge that protects it from extinction (MacArthur and Levins 1967). An example of a

stabilising mechanism is two species that do not have completely overlapping niches, but

instead partition some resources or limiting factors between themselves. The third coexistence

mechanism is termed the storage effect, which occurs when different species respond to

environmental variation in space or time in contrasting ways, with different species being more

competitive in some types of environment than others (Warner and Chesson 1985). For

example, one species may be a better competitor in winter, while the other may gain an

advantage in summer. The final mechanism is termed relative non-linearity where different

species benefit in different ways from fluctuations in the abundance of the limiting resource

via non-linear effects of environmental variation on their population growth rates (Chesson

1994). These last two mechanisms can contribute to coexistence when they act to increase the

strength of intraspecific competition to a greater extent than they increase the strength of

interspecific competition.

To incorporate these ecological coexistence mechanisms into demographic models it is

necessary to construct a model for each interacting species, and to couple them (Adler et al.

2010, Bassar et al. 2017). This can be achieved by making the demographic, development, or

inheritance functions of each species a function of the size or structure of both the focal species

and of the competitor species. Such models capture the indirect effects of two species

competing for a resource or other limiting factor; the resource or shared predator are not

explicitly included in models. Direct effects can be incorporated into models by formulating

demographic, development, and inheritance functions of focal species to be functions of the

size and structure of interacting resource or consumer species on neighbouring trophic levels

(Lachish et al. in press).

Ecological coexistence of multiple species requires direct interactions to limit each population

such that its long-run stochastic growth rate is zero. This might sound like a big ask, but most

species only interact strongly with a relatively small number of others, and when this happens,

interacting species often coexist by limiting one another (Allesina and Tang 2012). Consider,

for example, the case of the lynx and snowshoe hare, where both species cycle, with the

predator dynamics lagging a quarter of a cycle behind that of the prey (Keith and Windberg

1978), a pattern often termed Lotka-Volterra predator-prey dynamics. The predatory lynx is

limited by availability of the snowshoe hare prey, particularly in periods following high lynx

abundances. In contrast, the hare is limited by lynx predation, particularly when lynx are

abundant. The strength of these limiting factors fluctuate (reasonably) predictably with time,

and the two populations cycle. Their long-run stochastic growth rates are both zero, but the

numbers of both species cycle, in an endless entwined pattern, where each species limits the

other, at least some of the time (Krebs et al. 1986). Because many limiting factors are biotic, it

appears that ecosystem stability may be the norm, particularly when limitation of each

population operates primarily via a small number of strongly interacting species.

The case of the lynx and snowshoe hare is a special one, and few systems are this simple.

However, different species that share a resource or limiting factor are able to coexist via one

or more of the mechanisms described above because they have adaptations that allow them to

do so. In particular, different species may have adaptations that increase their fitness by making

them specialists in particular environments (e.g. the storage effect) or by allowing them to

utilise a unique resource that is unavailable to individuals of competitor species (e.g. a

stabilising mechanism) (MacArthur and Levins 1967, Warner and Chesson 1985). Successful

species are often those that have been able to innovate in a way that allows them to utilise a

previously unobtainable or under-utilised resource.

Rotifers feeding on algae provide a particularly interesting example of how phenotypic

variation that allows adaptation to specific environments permits coexistence. Algae-rotifer

dynamics deviate from classic Lotka-Volterra predator-prey expectations, with the predatory

rotifer population lagging a half cycle, rather than the expected quarter cycle, behind that of

their prey (Yoshida et al. 2003). The reason for this is that the algae come in two different

phenotypic forms – clumpers and soloists (Becks et al. 2010). The two forms are clonal, and

rarely reproduce sexually. They consequently act like two different species, even though they

are in fact two genetically distinct clones of the same species. Individuals of the soloist clones

have high fitness when the predators are rare. They can grow quickly, efficiently utilising

abundant resources, and they outcompete the clumper clone. Survival is high, as rotifers are

rare, and the population of soloists increases rapidly. The fitness of the soloist individuals starts

to decline as predator numbers increase, and their population becomes predator limited. Rotifer

numbers increase because the soloists make easy prey. However, the decline of soloists is

simultaneous with an increase in the fortunes of the clumpers. Individuals of this clone band

together into groups that are too large to be easily predated. They are not as efficient at turning

resources into new clones as the soloists are, but when soloists are largely absence, such

inefficiency is not an issue. As the clumpers dominate, the predatory rotifer population starts

to decline as accessible prey abundance declines, and the time of the soloist dominance comes

around once again. The two clones are able to coexist because the clumper has an adaptation

that favours them in the high predation environment, while the soloists dominate in a low

predation environment (Hiltunen et al. 2014). Population limitation of the two clones via

resource limitation and predation allow coexistence of different phenotypes: the two clones

coexist via the storage effect which is facilitated by the adaptive ability to clump or not.

Mechanisms such as this must be abundant in nature give so many species coexist.

The algae-rotifer system provides a link between coexistence of two species that share a

predator but cannot reproduce, and the maintenance of genetic variation in sexually

reproducing diploid species. The observation is particularly useful, as it reveals that the

mechanisms that allow ecological coexistence of species in communities at equilibrium, may

also contribute to evolutionary equilibria – another problem that needs solving if we are to

understand stationary ecosystems.

Trivial evolutionary equilibria occur when all genetic variation is eroded such that all

individuals are genetically identical (Wright 1969, Crow and Kimura 1970). However, such

equilibria are not informative when considering the types of evolutionary transients observed,

for example, in Trinidadian freshwater streams when predators are removed and the

community moves between stationary states quite rapidly (see introduction). More interesting

equilibria maintain additive genetic variance. For this to happen, either stabilising selection

and mutation-selection balance, a particular form of phenotypic plasticity, or frequency-

dependent selection, must be in operation (Gillespie 1994). But how are these related to the

coexistence mechanisms that ecologists focus on when considering species coexistence?

The strength of selection is determined by the association between phenotypic trait values and

fitness (e.g. survival and reproduction) (Fisher 1930, Price 1970). It is a within generation

process, most easily measured over a time step using viability and fertility differentials (Lynch

and Walsh 1998). How selection translates to the underlying genotypes and alleles is

determined by the genotype-phenotype map that determines how phenotypic traits develop.

Directional selection requires increasing or decreasing functions between phenotypic trait

values and fitness (e.g. survival and reproduction), while stabilising selection requires humped-

shaped associations, where the highest survival or reproductive rates are observed at

intermediate phenotypic trait values (Barton and Partridge 2000). Many evolutionary models

assume simple additive genotype-phenotype maps described in the previous section that allow

microevolution to be inferred directly from the dynamics of the phenotypic trait. This

assumption is known as the phenotypic gambit (Lynch and Walsh 1998).

If the phenotypic gambit is assumed (and frequency-dependence is not operating), evolutionary

equilibria require stabilising selection. Yet stabilising selection (as well as directional

selection) erodes additive genetic variance unless some other process replaces it.   When

selection is weak (i.e. the hump of the association between the phenotype and fitness is quite

flat), mutations can generate novel additive genetic variance at a rate that counters losses due

to selection in a process termed selection-mutation balance (Bürger and Lande 1994).

Phenotypic plasticity occurs when the same genotype can generate different phenotypes in

different environments (Pigliucci et al. 2006). If multiple genotypes can generate the optimal

phenotypic trait value, each genotype will have maximal fitness and will be able to coexist.

The third process that can maintain additive genetic variance is frequency-dependent selection

(Ayala and Campbell 1974). This occurs when a phenotypic trait value, and the underlying

genotypes and alleles, experience higher survival and reproductive rates the rarer they become,

such that they increase in frequency when rare. There are a number of processes that can

generate frequency-dependent selection including heterozygote advantage, disassortative mate

choice, spatial variation in the optimum phenotypic trait value, and heritable individual

specialisation in resource use or limiting factor (Hartl et al. 1997). This last process occurs

when the strength of competition increases with phenotypic similarity between competing

individuals, with coexistence of individual variation being facilitated by the four coexistence

mechanisms that operate at the species level (Bassar et al. 2016). For example, fitness-density

covariation can occur when individuals with different phenotypic trait values feed on the same

resources but in different locations – i.e. there is small-scale genetic variation underpinning

phenotypic variation. Stabilising mechanisms occur when populations consist of individual

specialists that exhibit different foraging phenotypes and feed on different resources (Chesson

1994). In the Cocos Island finch, some individuals exclusively forage on seeds and

invertebrates found on the ground, while others forage solely on arboreal insects, rarely

venturing onto the ground (Werner and Sherry 1987). When survival, reproductive, or

development rates of individuals with different phenotypic trait values are impacted by

environmental variation in different ways, then the storage effect can occur, and depending

upon the way that this environmental variation impacts demographic, development or

inheritance functions, relative non-linearity can also arise (Warner and Chesson 1985, Chesson

1994, 2000a).

The same processes can consequently maintain genetic diversity within a population, and

species diversity within a community. When these processes are operating, ecological and

evolutionary equilibria can be achieved, whereby each species, and each structural class in each

population, has a long-run stochastic growth rate of zero. At these equilibria, a number of other

properties of stationarity may be observed, including equivalence of reproductive values of

genotypes in the structural classes of offspring, and equivalent mean lifetime reproductive

successes of individuals regardless of their phenotypic trait values. But such patterns are not

always observed, and this conveniently leads me to the paradox of stasis.

A solution to the paradox of stasis

When the phenotypic gambit is assumed, phenotypic variation between individuals can be

partitioned into a contribution attributable to additive genetic variation (Falconer 1960, Lynch

and Walsh 1998). This can be estimated by statistically modelling phenotypic variation as a

function of relatedness between individuals (Kruuk 2004). The ratio of the additive genetic

variance of a phenotypic trait, to the variance of the trait itself, is the narrow sense heritability,

h^2. The closer h^2 is to one, the larger the proportion of phenotypic variance that is attributable

to additive genetic differences between individuals. When $h^2 = 0$, then none of the phenotypic

variation is attributable to additive genetic differences between individuals. The phenotypic

variance that is not attributable to the additive genetic variance can be partitioned into various

other contributions, such as dominance and epistatic variances, but the one I focus on is the

environmental variance. I consequently assume that an individual's phenotypic trait Z_i is equal

to the sum of an additive genetic contribution A_i (termed its breeding value) and an

environmental component E_i such that $Z_i = A_i + E_i$ (Lynch and Walsh 1998).

In populations at evolutionary stasis, distributions of phenotypic traits are stationary and show

no persistent temporal trends in their means or variances (Lande 2009). If additivity is assumed,

and the traits are heritable (e.g. $h^2 > 0$), this means they should not be subject to persistent

directional selection (Merilä et al. 2001b, Estes and Arnold 2007, Haller and Hendry 2014,

Rollinson and Rowe 2015, Bonamour et al. 2017). Paradoxically, many size-related traits, such

as body mass in fish (O'Sullivan et al. 2019), and horn length and antler size in ungulates

(Kruuk et al. 2002, Traill et al. 2014, Coulson et al. 2018), are heritable, under directional

selection, but do not evolve as predicted (Merilä et al. 2001b). Groups of individuals with larger

trait values have higher reproductive values, and greater values of R_0, but they have long-run

stochastic growth rates of $a = 0$. Structured demographic models that do not explicitly

incorporate evolution, capture these dynamics well (Traill et al. 2014). These models predict

no change in mean body size with time, even though the trait has an estimated non-zero

heritability from both observation and from model predictions (Coulson et al. 2010). Put

another way, evolution should occur, but the size-related traits are in fact at evolutionary

equilibria. Why does this happen? In short, the assumptions of additivity and the phenotypic

gambit may be violated, the demographic functions that determine phenotypic selection may

be incorrectly characterised, or the non-zero heritabilities may be an illusion. I argue here that,

for size-related traits exhibiting stasis, the assumption of the phenotypic gambit is violated.

However, to understand why, it is useful to address a question that is yet to be satisfactorily

answered: why are organisms the size that they are (Smith and Lyons 2013, Bhat et al. 2020)?

Body size is a fundamental component of life history strategies. Within a species, individuals

that are able to grow to large sizes are typically able to invest proportionally more energy in

reproduction than those that grow to smaller sizes. Such within-species hyper-allometric

scaling of reproductive investment is widespread, and may be universal (Barneche et al. 2018).

Achieving large size is consequently desirable because it results in a fitness return via an

increase in reproductive output. But if this is the case, why are not all species large? There is a

risk to individuals of growing to large sizes before starting reproduction: development takes

time, and delaying reproduction to older ages increases the risk of death before sexual maturity

is reached (Blanckenhorn 2000). There is consequently a time to maturity-reproductive output

trade-off that is determined by the associations of body size with survival, reproductive output,

and development rate throughout the life course (Childs et al. 2004, Bhat et al. 2020). At some

point on this trade-off, fitness will be maximised, and that determines size and age at sexual

maturity. The optimal age and size at sexual maturity, and adult body size, is consequently

determined by the shape of age- and size-specific fitness and transition functions (Kentie et al.

2020). Given this, the evolution of body size can be predicted by integrating across the entire

life cycle assuming trade-offs are appropriately identified (Childs et al. 2004). Reproductive

output can be divided into two other aspects of the life history strategy: offspring number and

offspring size (Fleming and Gross 1990). These two quantities also trade-off against one

another, with there being optimal values for both litter size and mean offspring size.

Rates of development are largely determined by the ability of individuals to acquire resources

from the environment (Reznick et al. 2000). Individuals utilise the energy they obtain by

allocating it to maintenance (staying alive and acquiring more resources), development, and

reproduction (De Roos et al. 2003, Kooijman and Kooijman 2010, Smallegange et al. 2017).

The way that resources are partitioned varies with age and size. The optimum body size and

life history strategy are achieved by adopting a specific energy budget that partitions energy

into each component of the life history at each age and size (Bhat et al. 2020). There is

consequently a very close association between energy budgets and life history; this has led

some authors to redefine fitness in terms of energy (Brown et al. 1993). Although at one level

such an approach is appealing, it is inconsistent with standard evolutionary theory where

selection, and fitness, are strictly defined in terms of numbers of descendants produced (Fisher

1930). Energy accrual needs to be linked to survival, development, reproduction, and

inheritance to understand body size evolution (Bhat et al. 2020).

Within in a population there is usually one dominant cause of death that kills the largest

proportion of individuals. However, individuals of different ages, or sizes, may experience

different causes of death. Nonetheless, at the level of the population there is usually a primary

cause of death associated with the limiting factor. Individuals within a population compete for

energy in the presence of a limiting factor that determined the primary cause of death and

failure to breed (Berryman 2004). For example, if prey (food) availability limits a predator

population, individual predators compete over prey. In order to be competitive at predation,

individuals need to express phenotypic traits that aid successful prey capture and consumption.

Such traits could include camouflaged pelage, fast running speeds, weaponry associated with

killing prey, and an ability to consume prey quickly. Selection is consequently expected to

operate most strongly on phenotypic traits associated with prey capture and consumption as

prey availability limits the population. In contrast, prey living in a predator-limited population

are selected to acquire food resources without being predated. They experience selection for

the accrual of enemy-free energy. In general, selection is expected to operate most strongly on

traits associated with the accrual of energy given a population's limiting factors and the primary

causes of death and failures to breed.

Those individuals with phenotypic trait values that are well-matched to the environment in

which they find themselves are better able to accrue resources than those with less optimal

phenotypic trait values (Charlesworth 1989). They consequently grow faster, to larger age-

specific sizes, have lower age-specific mortality rates, and higher rates of age-specific

reproductive output. Those individuals that accrue most resources consequently perform well

at all aspects of life compared to those that accrue fewer resources, and this heterogeneity in

resource accrual between individuals can mask life-history trade-offs (Van Noordwijk and de

Jong 1986, Metcalf 2016). A focus on resource accrual traits is consequently likely key to

understanding how individuals obtain utilise energy in as an optimal way as possible to

optimise life history and energy budgets, and maximise fitness (Bhat et al. 2020).

We can break down the accrual of resources into two separate processes: detection and

acquisition. The senses, and the ability to process the information they reveal, are associated

with the detection of resources, while many behaviors and some aspects of morphology are

associated with their acquisition. The optimal set of phenotypic traits, and their values,

associated with the detection and acquisition of resources will vary as a function of the

population's limiting factor and the primary causes of death and failures to breed.

Once resources are accrued, they need to be utilised. To do this, at each age and size,

individuals need to allocate the resources they have accrued to maintenance, development, and

reproduction (Kooijman and Kooijman 2010), in a manner that is expected to maximise their

reproductive value via optimising the life history (Goodman 1982, Bhat et al. 2020). This is a

hard-to-solve state-dependent decision-making problem that is constrained (McNamara and

Houston 1996). Life history constraints mean that the total amount of energy an individual can

allocate to a particular activity remains within bounds. For example, a long-lived iteroparous

species with low reproductive effort, cannot suddenly switch strategies and invest all its energy

into a single reproduction event, sacrificing its own survival. Energy budgets are plastic, but

within limits that are set by past fluctuations in energy availability that populations have

adapted to (DeWitt et al. 1998).

Phenotypic traits associated with resource detection, acquisition and utilisation are likely

subject to stabilising selection. This is because a trait value below a particular threshold will

likely carry survival or reproductive costs as it makes it challenging to accrue resources, while

an over-engineered trait carries a cost. For example, consider eye sight in Mexican cavefish.

Vision can be critical for detecting predators, mates, and energy. Invest too little in developing

vision, and the risk is an early death or a failure to find a mate. However, vision is expensive,

commanding up to 15% of an individual's energy budget (Moran et al. 2015). Visual acuity

beyond being able to observe a mate, a predator, or food, is over-engineered and carries an

energetic cost, and this is why the loss of cave fish vision in predator-free environments is

adaptive. There is consequently a sweet spot, where vision is good enough to thrive, but not

too good to be too expensive. Individuals with trait values at the sweet spot will acquire most

resources and get the most out of them given the limiting environment, are expected to develop

at a faster rate, and will have highest survival and reproductive rates, while those with smaller

or larger values will achieve smaller sizes and have lower expected fitness (Figure 3).

If the resource accrual traits are heritable (and random mating is assumed), individuals with

non-optimal resource accrual trait values and small body sizes will be expected to produce

offspring that also have non-optimal trait values and small body sizes. In contrast, those

individuals with resource accrual traits at the optimal value will be more likely to produce

offspring that resemble them in these traits. Body size will consequently appear heritable and

subject to directional selection (Figure 3), yet will not evolve to larger values because the mean

of the resource accrual trait will be at the optimal value once the population has achieved

evolutionary equilibrium. In addition, life history trade-offs will be hard to detect, because

individuals with more optimal resource accrual traits and energy budgets acquire more energy

and utilise it more effectively in development and reproduction than those with less optimal

trait values (Van Noordwijk and de Jong 1986).

For the arguments above to hold, genetic variation needs to be maintained in the resource

accrual trait. These traits may be determined by many loci of small effect, which means additive

genetic variation is lost slowly (Lande 1982). Selection-mutation balance could theoretically

contribute to the maintenance of genetic variation for resource accrual traits in such a scenario.

Frequency-dependent selection, operating via individual heterogeneity in resource use

(Svanbäck and Bolnick 2006), could also maintain genetic and phenotypic variation via fitness-

density covariance, stabilising mechanisms, the storage effect, or relative non-linearity

(Bolnick et al. 2011). For example, if a new mutation generates a phenotypic innovation that

allows an individual to utilise an under-utilised, or new, resource, that individual will accrue

more resources, grow quickly, have high survival, and produce many offspring. If some of this

individual's descendants are also equipped to utilise the resource, the frequency of the new

innovation will increase, but the novel resource being utilized will decline in abundance.

Eventually the frequency of the new innovation will achieve an equilibrium point where it, and

the resource, are in equilibrium. Random fluctuations in the biotic and abiotic environment can

generate fluctuations in the numbers of genotypes and phenotypic traits, and this could generate

frequency-dependent selection in resource accrual traits, contributing to maintain additive

genetic variation.

It is possible that frequency-dependent and stabilising selection can operate simultaneously on

resource accrual traits. If stabilising mechanisms contribute to the maintenance of genetic and

phenotypic diversity, different phenotypic optima may exist among individuals that specialise

on different resources. This would manifest as a flat-humped fitness function for the resource

accrual trait when averaged across multiple time steps.

In this section I have argued that body size is determined by an interaction between breeding

values A_i and the environmental component of the phenotype E_i such that body size is not

determined by $Z_i \neq A_i + E_i$ but rather by $Z_i = A_i + E_i|A_i$ (Coulson et al. 2017). I propose that

this is plausible, because A_i is in fact a breeding value for a resource accrual trait subject to

stabilising selection that impacts body size by determining the quantities of resources accrued

by each individual. This conclusion is general: when a phenotypic trait appears heritable, is

subject to directional selection, but does not evolve, then the assumption of additivity must be

violated. Size-related traits appear to frequently violate this assumption. But in some cases, the

amount of resources accrued may be determined by body size. How might this affect evolution?

Body size as a resource accrual trait

In some instances, individuals with large body size are able to access more resources than those

that are smaller. For example, large lions can prevent smaller members of the pack feeding on

a kill, while large individuals of many species may be able to dominate smaller individuals in

competition for a territory. Alternatively, it may be aggression that is the resource accrual trait

determining the quantity of resources accrued, such that body size is a consequence of how

aggressive an individual is, rather than body size per se being the resource accrual trait.

Body size also has the potential to play a key role as a resource accrual trait in seasonal

environments where resource availability varies within a year as can be seen by returning to

the long-run stochastic growth rate. The long run stochastic growth rate can be approximated

as

$$a \sim \log(\overline{\lambda}_t) - V_{E,t} \tag{2}$$

where the first term is the log of the mean of the per time-step population growth rates, and the

second term, $V_{E,t}$, is a function of the logged variance in the per time-step population growth

rates (Lande et al. 2003). When $a = 0$, then obviously $\log(\overline{\lambda}_t) = V_{E,t}$. Mean fitness can

consequently be increased when evolution operates on traits that either increase $\log(\overline{\lambda}_t)$ or

decrease $V_{E,t}$ (Lande et al. 2017). Body size, hibernation, and migration are all traits that can

evolve to decrease $V_{E,t}$ in seasonal environments. For example, individuals of large size of

warm-blooded species have a higher body mass to volume ratio than their smaller counterparts

and consequently use relatively less energy to keep warm in cold environments (Coulson et al.

2001). A consequence of this is that they may be able to survive winters more easily at higher

latitudes than smaller individuals. Bigger individuals can also store larger fat reserves than

smaller individuals to utilise during winter when energy is scarce. Large body size may

consequently allow non-hibernating species to exist in areas where smaller competitor species

cannot survive, thus accessing resources that are only available to them. However, the flip side

to this, is that large size also requires more energy for maintenance (Brown et al. 1993, West

et al. 1997), so resources must be sufficiently abundant during the summer months for

individuals to accrue sufficient resources to lay down enough fat to survive the winter. It is

possible that productivity in the growing season, and the harshness of the winter season, may

both select for larger size within species as distance to the poles decreases, a phenomenon

termed Bergmann's rule (Bergmann 1848). It is also possible that climate change resulting in

shorter and warmer winters size may allow smaller individuals to survive, leading to a

reduction in body size within populations (Ozgul et al. 2009).

In some cases when body size is a resource accrual trait, it can conform to the additivity

assumptions of quantitative genetics, and can evolve as predicted by the Breeder's equation.

Evidence for this comes from the observation that when body size is selected for in artificial

settings it can evolve as predicted using methods from quantitative genetics. This insight might

help explain why quantitative genetics so often works well in artificial environments but fails

to do so in natural ones (Merilä et al. 2001b).

In artificial environments, energy is easy to detect and to acquire *ad libitum*. Selection is

consequently largely relaxed on resource accrual traits, but is instead targeted at phenotypic

traits we consider desirable. Only those individuals with desirable trait values are permitted to

breed. This targeting selects for individuals that budget the largest proportions of the energy

they accrue to the traits we desire. Artificial selection consequently selects for particular energy

budgets that result in the traits we wish to evolve.

In the above sections I argue that populations and communities achieve ecological and

evolutionary quasi-equilibria for protracted periods of time over which species composition

remains constant. I then explain the set of conditions required for these equilibria to be attained.

I now consider how models of the phenomena described above can be constructed.

Population models of ecological and evolutionary stasis

In the section "**Characterising Populations at Ecological and Evolutionary Equilibrium**" I

briefly explain that models can be structured to track the dynamics of distributions of individual

attributes based on functions that describe associations of these attributes with survival,

reproduction, development, and inheritance. I now expand on these earlier comments.

Structured demographic models can be constructed in discrete or continuous time. Discrete-

and continuous-time modelling has proceeded largely independently, but has arrived at

remarkably similar conclusions. Both approaches combine demographic, development, and

inheritance functions to track the dynamics of distributions of genotypes and phenotypic traits

(Figure 4). Models can be structured or individual-based (e.g. (Dunlap et al. 2006, Romero-

Mujalli et al. 2019).

Discrete time models called integral projection models (IPMs) of phenotypic trait distributions

were initially developed by (Easterling et al. 2000). Since then, models have been extended to

incorporate explicit genotype-phenotype maps (Coulson et al. 2011, Childs et al. 2016, Coulson

et al. 2017, Rees and Ellner 2019), frequency-dependence (Schindler et al. 2015), density-

dependence (Rees et al. 2014), abiotic variation (Ellner and Rees 2006), and species

interactions (Adler et al. 2010, Bassar et al. 2017, Lachish et al. in press). These models have

been used to study the dynamics of populations and distributions of phenotypic traits, breeding

values and additive genotypes, non-additive genotypes, and life history strategies. The work

has identified circumstances when eco-evolutionary feedbacks occur (Coulson et al. 2011). In

these models, fitness, population dynamics and evolution are all emergent properties of the

demographic, development and inheritance functions used to construct the models.

Continuous time models that track distributions of individual attributes have started to receive

significant attention more recently. For example, (DeLong and Gibert 2016) introduced

Gillespie eco-evolutionary models that are a form of continuous time individual-based model,

and (Doebeli et al. 2017) argue that evolutionary models developed from birth and death rate

functions to track distributions of heritable phenotypic traits are useful because fitness is an

emergent property that does not need to be defined. Along similar lines, (Lowe et al. 2017)

argue that demography and population genetics need to be explicitly combined to understand

eco-evolutionary dynamics.

Because the two separate literatures have arrived at the same conclusions, it suggests that

models based on demographic, development and inheritance functions offer great promise in

providing novel ecological, evolutionary, and eco-evolutionary insight. I now focus on discrete

time models that track distributions of alleles, genotypes, breeding values, and phenotypes for

three reasons. First, the theory is more developed than for continuous time models, second, the

approach has gained more traction amongst empiricists, and third, I am more familiar with this

approach than with continuous time equivalents. However, the conclusions will extend to

continuous time models.

IPMs are of the general canonical form:

753
$$N_{t+1}(\mathbf{Z'}) = \int [D_t(\mathbf{Z}, \mathbf{\Theta}) S_t(\mathbf{Z}, \mathbf{\Theta}) + H_t(\mathbf{Z}, \mathbf{\Theta}) R_t(\mathbf{Z}, \mathbf{\Theta})] N_t(\mathbf{Z}) d\mathbf{Z} \tag{3}$$

where $\mathbf{Z}$ is the set of individual attributes being iterated forwards, $N_t(\mathbf{Z})$ is the distribution of

these attributes, the functions $S_t(\mathbf{Z}, \mathbf{\Theta})$ and $R_t(\mathbf{Z}, \mathbf{\Theta})$ are demographic functions describing how

the individual attributes and the environment $\mathbf{\Theta}$ at time t influence the rates of survival and

reproduction respectively, and the functions $D_t(\mathbf{Z}, \mathbf{\Theta})$ and $H_t(\mathbf{Z}, \mathbf{\Theta})$ are transition functions

describing development and inheritance of the individual attributes. These functions are

probability density functions such that they can move individuals between individual attribute

classes but they do not alter the size (or volume) of the distribution. The transition functions

can become quite complicated when different attributes included in Z have different

development and inheritance rules (Schindler et al. 2013, Childs et al. 2016, Coulson et al.

2017). Feedbacks occur when properties of $N_t(\mathbf{Z})$ contribute to the environment $\mathbf{\Theta}$ either

directly or indirectly via interactions with other species.

How can models be constructed that achieve ecological and evolutionary equilibrium? First,

these models need to iterate forward the dynamics of genotypes and phenotypic trait

distributions. Demographic functions describe the growth rate of each part of the distribution

being iterated but do not move individuals between categories; transition functions − e.g.

development and inheritance functions − describe these movements (Coulson 2012). Second,

models need to include ecological feedbacks such as species interactions, and stabilizing or

frequency-dependent selection operating on genetic variation.

Demographic and transition functions are consequently central to constructing structured

models of ecological and evolutionary dynamics and feedbacks. A key challenge in

constructing models for multivariate distributions is that different types of individual attributes,

or structural classes, have different development and inheritance properties, and consequently

require different functional forms for their transition functions (Childs et al. 2016, Coulson et

al. 2017) (Figure 4). Genotypes and breeding values, for example, remain fixed throughout life

such that development functions simply keep individuals in the same class throughout

life(Charlesworth 1994). For models containing these attributes to achieve a non-trivial

evolutionary equilibrium, demographic functions need to capture a process such as frequency-

dependent selection that maintains additive genetic variance, while simultaneously preventing

directional evolution occurring indefinitely. Transition functions are also required to capture

genetic inheritance, and the formation of offspring genotypes and breeding values as a function

of parental values and patterns of mate choice (Figures 4(A-D)).

If I assume that resource accrual traits are additive and heritable and retain a fixed shape

throughout life, but scale with body size, demographic and genetic inheritance functions that

are similar to those for genotypes and breeding values are required, but so too is a development

function that specifies how the size of the trait changes across the lifespan via changes to the

environmental component of the phenotype (Figures 4(E-H)). These development functions

keep the breeding values or genotypes associated with the resource accrual traits constant

throughout life, while allowing the trait to develop with age, via change in the value of the

environmental component of the phenotype.

Traits that exhibit the paradox of stasis such as size-related traits at equilibrium, need to

develop with age, may show directional selection, but are not specified with an additive

genotype-phenotype map (Coulson et al. 2010). Large parents may produce large offspring,

and traits may statistically appear to be heritable but do not evolve (Figures 4(L)). Traits such

as this require demographic, development, and inheritance functions, but the inheritance

functions do not need to explicitly include rules for genetic inheritance for the trait itself

because these traits are emergent consequences of an individual's resource accrual trait value

and the environment. Similarly, development functions can be specified to be determined by

the value of resource accrual traits and the quantity of resources accrued (Lachish et al. in

press).

The final trait type is life history traits such as size and age at sexual maturity. These traits are

often referred to as latent traits, and they are only expressed once during an individual's life

(Childs et al. 2016). If these traits are under genetic control, the underlying genotypes need to

be tracked throughout life, with the trait only expressed at one point in time, determining when

reproduction occurs for the first time. The genetic architecture underpinning the trait can be

subject to selection when the trait is not expressed if the trait genetically covaries with

phenotypic traits that are expressed and are subject to selection.

IPMs have been developed to track the dynamics of each trait type, but all trait types are yet to

be combined into a single model. One way to do this would be to construct a model of i) a

resource accrual trait that is determined by a breeding value and an environmental component

of the phenotype, ii) body size, where the development function is determined by the value of

the resource accrual trait and the amount of resources accrued, and iii) age and size at sexual

maturity is determined by a breeding value and an environmental component of the phenotype.

An initial model might focus on a resource species and a consumer species. The resource

accrual trait in the resource species would define the ability to escape consumption, while the

resource accrual trait in the consumer species would define success at consuming resources.

Lachish *et al.* (in press) have developed a model of a vegetation resource that renews

seasonally, and stochastically. The resource is consumed by elk (*Cervus canadensis*), with

individuals in each attribute class partitioning the energy they accrue into maintenance,

development, and, if individuals have reached sexual maturity, reproduction. A proportion of

the maternal energy reserve of breeding individuals is invested into offspring. In this

bioenergetic model based on dynamic energy budget theory (Kooijman and Kooijman 2010),

the life history is determined by the energy budget at each age and size. This work builds on a

previous model of Smallegange *et al.* (2017) that was the first to incorporate energy budgets

into IPMs.

In such a model where the initial conditions are away from the evolutionary and ecological

equilibrium, the breeding values for the resource accrual traits will change at each time step in

a way determined by the covariance between breeding value and annual survival and

reproductive rates at each age (Coulson et al. 2017). The dynamics of these traits will be

accurately predicted by their per time-step selection differentials. Body sizes will change with

time as the distribution of the resource accrual traits and the numbers of resources and

consumers change. Body size dynamics will not necessarily be well-predicted by their per-time

selection differentials. Neither will be the dynamics of age- and size at sexual maturity (Childs

et al. 2011). In order to understand the dynamics of the resource accrual trait it is necessary to

understand how evolution optimizes the trade-off between the risk of mortality prior to sexual

maturity against any benefits of delaying reproduction to a later stage of development (Childs

et al. 2004). The optimization of the trade-off maximizes reproductive value. A feedback now

occurs: as reproductive value is maximized, age-specific survival and reproductive rates

change, which means that the distribution of annual fitness changes, which alters the strength

of selection on the resource accrual traits. However, eventually an ecological and evolutionary

equilibrium will be achieved.

Transitions between equilibria

Sudden shifts in the environment Θ be they biotic, such as the addition or removal of a species,

or abiotic, such as a volcanic eruption, can alter the causes of death or failures to breed in some

species. These changes will alter predicted values of the demographic and transition functions.

For example, the loss of a predator species may result in a shift from top-down to bottom-up

regulation of a prey species (Figure 5(A)-(B)) reflected by a reduction in the influence of a

predator population on survival, reproduction, development or inheritance of a focal species,

coupled with an increase in the role of a resource species on these functions. Such a change in

a limiting factor can result in rapid change in population size of affected species (Figure 1(A)),

and alter selection pressures on phenotypic traits associated with resource accrual (Figure 5(a)-

(B)) and utilization (Figure 5(C)). Novel selection on these traits can result in a shift in

phenotypic trait means via plasticity, (Figure 1(B)), and a slower rate of evolution, altering the

frequency of alleles associated with resource accrual traits. Evolution occurs at a slower rate

to the initial change in population size, such that the long-run stochastic growth rate of the

population rapidly returns to being close to zero following the initial environmental

perturbation. The novel selective regime on resource accrual traits can alter the optimal form

of associations between body size and survival, reproduction, development, and non-genetic

inheritance, leading to evolution of the life history. Change in the life history may result in

additional novel selection pressures on resource accrual traits – a form of eco-evolutionary

feedback.

If a change in a limiting factor, population size, and selection pressures on resource accrual

traits impacts the size of "keystone species" within the system, trophic cascades can occur

where populations of other species within the ecosystem are impacted (Paine 1966, Bronstein

et al. 2004, Georgelin et al. 2015). In this case, the distances between the population structures

of the species within the community at the initial and new equilibrium will be large (Johansson

and Dieckmann 2009). In contrast, if the changes in one species do not have substantial effects

on keystone species, then a major shift in ecological and evolutionary equilibrium may not be

observed (Brown and Vincent 1992).

The most pronounced trophic cascades occur when the composition of the dominant producer

species changes. For example, in Yellowstone, it is argued that the decline in elk numbers due

to increased wolf predation has led to forests replacing grasslands (Fortin et al. 2005). The

effect is currently localized to small areas, as the two decades since wolves were reintroduced

is not sufficient time for a forest to displace a grassland. However, if the effects observed on

small spatial scales are eventually repeated on a larger scale, the reintroduction of wolves will

be responsible for a pronounced trophic cascade as forests replace grasslands on the Northern

range.

Less pronounced trophic cascades can occur when only a small part of a food web is affected,

that does not impact primary producers. For example, a decline in the density of a predatory

bird may influence the dynamics of its passerine prey and the insects on which they feed, but

effects on the species structure of the primary producers is likely to be minimal (Borer et al.

2005). Those species directly impacted may experience altered causes of death or failures to

reproduce, and novel selection pressures, but community structure and ecosystem function is

unlikely to be significantly impacted to a sufficient extent to generate a pronounced trophic

cascade.

Transitions between equilibria, whether they produced pronounced trophic cascades or not, can

be captured with structured demographic models of interacting species that have achieved

ecological and evolutionary stasis by setting the population of one species in the community to

zero, by introducing a new species, or by changing the distribution of an abiotic driver. Any

function that includes population size of the removed or added species, or the abiotic driver,

will be directly impacted. Indirect effects will then cascade to other species whose demography

might not be directly impacted by environmental perturbation, but might impacted by species

that are directly impacted.

Conclusions

The motivation for this paper was understanding the observations that i) communities often

appear to be in equilibrium for protracted periods of time, yet ii) maintain additive genetic

variance for many traits, allowing them iii) to evolve following an environmental perturbation,

yet iv) many species exhibit traits, such as size, that do not evolve as predicted but instead

exhibit stasis. When systems are perturbed from their equilibrium by the addition, or removal,

of a species, they tend to a new ecological and evolutionary equilibrium (Bronstein et al. 2004,

Georgelin et al. 2015). Individuals that are well-matched to their environment can exhibit

different life history strategy strategies and resource accrual traits, or trait values, in each of

the equilibrium states, with selection on these traits determined by the limiting environment. I

have focused on traits associated with the accrual of food, but similar arguments hold for traits

associated with the acquisition of mates. In order to understand observed dynamics, I have

focused on the properties of ecological and evolutionary equilibria, the conditions required for

these properties to be observed, how systems are expected to move from one state to another,

and how one could go about modelling these dynamics using existing, well-developed,

approaches (see also Johansson & Dieckmann 2009).

In this paper I have linked population limitation to community dynamics to evolution via biotic

interactions and the accrual of resources. The ideas I present are consistent with many

observations from a range of systems, but some are speculative. This manuscript should be

seen as a perspective, rather than a comprehensive review. I hope that rather than being the last

word, the paper generates discussion, tests of some of the hypotheses proposed, and acts to

stimulate new models of the interplay between ecology and evolution.

ACKNOWLEDGEMENTS

The ideas in this paper have arisen from numerous discussions and arguments with friends,

colleagues, and adversaries over the years. In the friend and colleague camp I would like to

thank Ron Bassar, Tim Benton, Ottar Bjornstad, Mark Boyce, Sonya Clegg, Tim Clutton-

Brock, Sasha Dall, Andy Dobson, Jean-Michel Gaillard, Alan Grafen, Bryan Grenfell, Pete

Hudson, Bruce Kendall, Loeske Kruuk, Virpi Lummaa, Dan MacNulty, Dan Nussey,

Josephine Pemberton, David Reznick, Andy Russell, Bernt-Erik Saether, Isabel Smallegange,

Doug Smith, Dan Stahler, Joe Travis, Shripad Tuljapurkar, Alastair Wilson and all past and

present members of my research group, for stimulating science discussions. I also owe a debt

of gratitude to Marco Festa-Bianchet, Michael Morrissey, Erik Postma, and Norman Owen-

Smith who I have not always agreed with, but whose critiques influenced my thinking; thank

you, and apologies if our arguments caused any stress. I would also like to thank NERC,

BBSRC, the Wellcome Trust, and the ERC for funding. Having been funded to work on

systems as diverse as the guppies of Trinidad (thanks Ron, Joe and David), the wolves of

Yellowstone (thanks Doug, Dan and Dan), the Soay sheep and red deer of Scotland (thanks

Tim CB, Josephine, Dan N, Alastair, and Loeske), and the mites of Holland (thanks Isabel) is

a real privilege, and each system has contributed to the ideas described here.

Figure 1. The (A) population, (B) phenotypic trait, and (C) micro-evolutionary dynamics of populations as they move from one stationary state to another. At ecological and evolutionary equilibrium, there are no temporal trends in population size, phenotypic trait distributions, or allele frequencies (black lines). Following an environmental perturbation (dotted vertical line) ecological and evolutionary change occurs (red and green lines) until a new stationary state is achieved (blue lines). In this example, population size changes quickly following the perturbation, while phenotypic and evolutionary changes occur more slowly. There will be a period when phenotypic plasticity is likely stronger than microevolution, which is reflected in quite rapid phenotypic change (B). Evolution occurs more slowly, resulting in gradual change in population size, the mean of phenotypic traits, and allele frequencies. I have given an example of ecological and evolutionary change in a single population, but similar patterns will be observed in other species within the community.

Figure 2. Complex eco-evolutionary feedbacks can arise from the dynamics of distributions of

alleles, genotypes and phenotypes (levels of biological organization). These levels are

mechanistically linked by key biological processes. Mate choice determines how alleles form

genotypes in offspring, with development determining how phenotypes develop from

genotypes. Phenotypes impact which resources individuals acquire and can influence survival,

reproduction, dispersal and life history (demography). Acquired resources, such as a territory,

can also influence demography directly, while also feeding back to impact further phenotypic

development (including state-dependent behaviors). Selection and population dynamics are

emergent consequences of which phenotypes survive and reproduce. The size and structure of

the focal species and those species with which it interacts, constitute the biotic environment.

The biotic environment, along with the abiotic environment, can directly influence the

distribution of resources, but also biological processes in both the focal species and other

species within the community. Recombination and mutation generate genetic variation and can

be influenced by the environment.

965   **Figure 3.** (A) A positive association between body size and fitness arises as (B) individuals

966   that accrue most resources are larger and have higher fitness, which (C) is due to an n-shaped

967   association between resource accrual and the resource accrual trait values. The dotted lines

968   help visualize the map across functions.

970
Figure 4. The form of survival (A, E, I), development (B, F, J), reproduction (C, G, K) and

inheritance (D, H, L) functions for different individual attribute types. (A-D) form of

functions for a breeding value or genotype that is subject to stabilizing viability (A) and

fertility (C) selection, that does not develop throughout life (B), and exhibits genetic

inheritance (D). The two narrower grey distributions in (D) represent breeding values or

genotypes in parents, while the flatter black distribution represents the distribution in

offspring following genetic inheritance. Similar dynamics are observed for a resource

accrual trait that maintains its shape throughout life but increases in size with age as the

environmental component of the phenotype develops (F). The trait is genetically inherited

(H). Body size (I-L) is subject to directional viability (I) and fertility (K) selection, develops

with age, and appears heritable (L) with parents and offspring have more similar values than

non-related individuals. Different colored lines represent different ages, showing how

functions can vary with age. In (B), (D) and (H) the same functions are observed at each age.

Figure 5. An environmental perturbation, such as the loss of a predator, can result in a

period of transience between ecological and evolutionary equilibria. The removal of a

predator can result in a change in limiting factor and a switch from top down (A) to bottom

up (B) regulation. The change in the primary cause of death and failure to reproduce can

relax selection on phenotypic traits associated with the original limiting factor and impose

selection on traits associated with the new one (3D surfaces in A and B). These changes can

alter demographic (C – survival as an example) and transition (D – development as an

example) functions for size-related traits. In this example, a switch in limiting factor an

increase in prime adult survival occurs with a simultaneous decrease in juvenile survival (C,

gray arrows), an increase in the mean and variance in senescent survival (C black arrows),

and a slowing in development rates (D, black arrow). In the original and novel equilibrium

these functions take values that result a long stochastic growth rate of zero, although the

mean and variance in population size may have changed (Figure 1(A)).

1000

Conflict of Interest Disclosure

The author of this article declares that he has no financial conflict of interest with the content

of this article. Tim Coulson is one of the founding members of PCIEcology and acts as a

recommender for it.

References

Abrams, P. A., and X. Chen. 2002. The evolution of traits affecting resource acquisition and predator vulnerability: character displacement under real and apparent competition. The American Naturalist **160**:692-704.

Adler, P. B., S. P. Ellner, and J. M. Levine. 2010. Coexistence of perennial plants: an embarrassment of niches. Ecology Letters **13**:1019-1029.

Allesina, S., and S. Tang. 2012. Stability criteria for complex ecosystems. Nature **483**:205.

Ayala, F. J., and C. A. Campbell. 1974. Frequency-dependent selection. Annual Review of Ecology and Systematics **5**:115-138.

Barneche, D. R., D. R. Robertson, C. R. White, and D. J. Marshall. 2018. Fish reproductive-energy output increases disproportionately with body size. Science **360**:642-645.

Barton, N., and L. Partridge. 2000. Limits to natural selection. BioEssays **22**:1075-1084.

Barton, N. H., and A. M. Etheridge. 2011. The relation between reproductive value and genetic contribution. Genetics **188**:953-973.

Bassar, R. D., D. Z. Childs, M. Rees, S. Tuljapurkar, D. N. Reznick, and T. Coulson. 2016. The effects of asymmetric competition on the life history of Trinidadian guppies. Ecology Letters **19**:268-278.

Bassar, R. D., J. Travis, and T. Coulson. 2017. Predicting coexistence in species with continuous ontogenetic niche shifts and competitive asymmetry. Ecology **98**:2823-2836.

Becks, L., S. P. Ellner, L. E. Jones, and N. G. Hairston Jr. 2010. Reduction of adaptive genetic diversity radically alters eco-evolutionary community dynamics. Ecology Letters **13**:989-997.

Bergmann, C. 1848. Über die verhältnisse der wärmeökonomie der thiere zu ihrer grösse. Göttinger Studien **3**:595-708.

Berryman, A. A. 2004. Limiting factors and population regulation. Oikos **105**:667-670.

Bhat, U., C. P. Kempes, and J. D. Yeakel. 2020. Scaling the risk landscape drives optimal life-history strategies and the evolution of grazing. Proceedings of the National Academy of Sciences **117**:1580-1586.

Blanckenhorn, W. U. 2000. The evolution of body size: what keeps organisms small? The Quarterly Review of Biology **75**:385-407.

Bolnick, D. I., P. Amarasekare, M. S. Araújo, R. Bürger, J. M. Levine, M. Novak, V. H. Rudolf, S. J. Schreiber, M. C. Urban, and D. A. Vasseur. 2011. Why intraspecific trait variation matters in community ecology. Trends in Ecology & Evolution **26**:183-192.

Bolnick, D. I., R. Svanbäck, M. S. Araújo, and L. Persson. 2007. Comparative support for the niche variation hypothesis that more generalized populations also are more heterogeneous. Proceedings of the National Academy of Sciences **104**:10075-10079.

Bonamour, S., C. Teplitsky, A. Charmantier, P. A. Crochet, and L. M. Chevin. 2017. Selection on skewed characters and the paradox of stasis. Evolution **71**:2703-2713.

Borer, E. T., E. W. Seabloom, J. B. Shurin, K. E. Anderson, C. A. Blanchette, B. Broitman, S. D. Cooper, and B. S. Halpern. 2005. What determines the strength of a trophic cascade? Ecology **86**:528-537.

Bronstein, J. L., U. Dieckmann, and R. Ferrière. 2004. Coevolutionary dynamics and the conservation of mutualisms. in: Evolutionary Conservation Biology, eds. Ferrière R, Dieckmann U & Couvet D, pp. 305–326. Cambridge University Press.

Brown, J. H., J. F. Gillooly, A. P. Allen, V. M. Savage, and G. B. West. 2004. Toward a metabolic theory of ecology. Ecology **85**:1771-1789.

Brown, J. H., P. A. Marquet, and M. L. Taper. 1993. Evolution of body size: consequences of an energetic definition of fitness. The American Naturalist **142**:573-584.

Brown, J. S., and T. L. Vincent. 1992. Organization of predator-prey communities as an evolutionary game. Evolution **46**:1269-1283.

Bürger, R., and R. Lande. 1994. On the distribution of the mean and variance of a quantitative trait under mutation-selection-drift balance. Genetics **138**:901-912.

Caswell, H. 2001. Matrix Population Models: Construction, Analysis, and Interpretation. Sinauer Associates, Sunderland, MA.

Caswell, H., T. Takada, and C. M. Hunter. 2004. Sensitivity analysis of equilibrium in density-dependent matrix population models. Ecology Letters **7**:380-387.

Charlesworth, B. 1994. Evolution in age-structured populations. Cambridge University Press Cambridge.

Charlesworth, D. 1989. Evolution of low female fertility in plants: pollen limitation, resource allocation and genetic load. Trends in Ecology & Evolution **4**:289-292.

Chesson, P. 1994. Multispecies competition in variable environments. Theoretical Population Biology **45**:227-276.

Chesson, P. 2000a. General theory of competitive coexistence in spatially-varying environments. Theoretical Population Biology **58**:211-237.

Chesson, P. 2000b. Mechanisms of maintenance of species diversity. Annual Review of Ecology and Systematics **31**:343-366.

Chesson, P. L. 1986. Environmental variation and the coexistence of species. Community Ecology **240**:54.

Childs, D., T. Coulson, J. Pemberton, T. Clutton-Brock, and M. Rees. 2011. Predicting trait values and measuring selection in complex life histories: reproductive allocation decisions in Soay sheep. Ecology Letters **14**:985-992.

Childs, D. Z., M. Rees, K. E. Rose, P. J. Grubb, and S. P. Ellner. 2004. Evolution of size–dependent flowering in a variable environment: construction and analysis of a stochastic integral projection model. Proceedings of the Royal Society of London B: Biological Sciences **271**:425-434.

Childs, D. Z., B. C. Sheldon, and M. Rees. 2016. The evolution of labile traits in sex-and age-structured populations. Journal of Animal Ecology **85**:329-342.

Coulson, T. 2012. Integral projections models, their construction and use in posing hypotheses in ecology. Oikos **121**:1337-1350.

Coulson, T., T. G. Benton, P. Lundberg, S. R. X. Dall, B. E. Kendall, and J.-M. Gaillard. 2006. Estimating individual contributions to population growth: evolutionary fitness in ecological time. Proceedings of the Royal Society B: Biological Sciences **273**:547-555.

Coulson, T., E. A. Catchpole, S. D. Albon, B. J. T. Morgan, J. M. Pemberton, T. H. Clutton-Brock, M. J. Crawley, and B. T. Grenfell. 2001. Age, sex, density, winter weather, and population crashes in Soay sheep. Science **292**:1528-1531.

Coulson, T., B. E. Kendall, J. Barthold, F. Plard, S. Schindler, A. Ozgul, and J.-M. Gaillard. 2017. Modeling adaptive and nonadaptive responses of populations to environmental change. The American Naturalist **190**:313-336.

Coulson, T., D. R. MacNulty, D. R. Stahler, R. K. Wayne, and D. W. Smith. 2011. Modeling effects of environmental change on wolf population dynamics, trait evolution, and life history. Science **334**:1275-1278.

Coulson, T., S. Schindler, L. Traill, and B. E. Kendall. 2018. Predicting the evolutionary consequences of trophy hunting on a quantitative trait. The Journal of Wildlife Management **82**:46-56.

Coulson, T., S. Tuljapurkar, and D. Z. Childs. 2010. Using evolutionary demography to link life history theory, quantitative genetics and population ecology. Journal of Animal Ecology **79**:1226-1240.

Crow, J. F., and M. Kimura. 1970. An introduction to Population Genetics Theory. The Blackburn Press, NJ.

De Roos, A. M., L. Persson, and E. McCauley. 2003. The influence of size-dependent life-history traits on the structure and dynamics of populations and communities. Ecology Letters **6**:473-487.

DeLong, J. P., and J. P. Gibert. 2016. Gillespie eco-evolutionary models (GEM s) reveal the role of heritable trait variation in eco-evolutionary dynamics. Ecology and Evolution **6**:935-945.

DeWitt, T. J., A. Sih, and D. S. Wilson. 1998. Costs and limits of phenotypic plasticity. Trends in Ecology & Evolution **13**:77-81.

Doebeli, M., Y. Ispolatov, and B. Simon. 2017. Point of view: Towards a mechanistic foundation of evolutionary theory. Elife **6**:e23804.

Dunlap, K. A., M. Palmarini, M. Varela, R. C. Burghardt, K. Hayashi, J. L. Farmer, and T. E. Spencer. 2006. Endogenous retroviruses regulate periimplantation placental growth and differentiation. Proceedings of the National Academy of Sciences **103**:14390-14395.

Easterling, M. R., S. P. Ellner, and P. M. Dixon. 2000. Size-specific sensitivity: applying a new structured population model. Ecology **81**:694-708.

Ellner, S. P., D. Z. Childs, and M. Rees. 2016. Data-driven modelling of structured populations: A Practical Guide to the Integral Projection Model. Springer.

Ellner, S. P., and M. Rees. 2006. Integral projection models for species with complex demography. The American Naturalist **167**:410-428.

Engen, S., and B.-E. Sæther. 1998. Stochastic population models: some concepts, definitions and results. Oikos:345-352.

Estes, S., and S. J. Arnold. 2007. Resolving the paradox of stasis: models with stabilizing selection explain evolutionary divergence on all timescales. The American Naturalist **169**:227-244.

Falconer, D. S. 1960. Introduction to Quantitative Genetics. Oliver And Boyd, Edinburgh / London.

Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Clarendon, Oxford.

Fishman, L., and J. H. Willis. 2008. Pollen limitation and natural selection on floral characters in the yellow monkeyflower, *Mimulus guttatus*. New Phytologist **177**:802-810.

Fleming, I. A., and M. R. Gross. 1990. Latitudinal clines: a trade-off between egg number and size in Pacific salmon. Ecology **71**:1-11.

Fortin, D., H. L. Beyer, M. S. Boyce, D. W. Smith, T. Duchesne, and J. S. Mao. 2005. Wolves influence elk movements: behavior shapes a trophic cascade in Yellowstone National Park. Ecology **86**:1320-1330.

Gaillard, J.-M., M. Festa-Bianchet, N. G. Yoccoz, A. Loison, and C. Toigo. 2000. Temporal variation in fitness components and population dynamics of large herbivores. Annual Review of Ecology and Systematics **31**:367-393.

Gause, G. F. 1932. Experimental studies on the struggle for existence: I. Mixed population of two species of yeast. Journal of Experimental Biology **9**:389-402.

Georgelin, E., G. Kylafis, and N. Loeuille. 2015. Eco-Evolutionary Dynamics of Plant–Insect Communities Facing Disturbances: Implications for Community Maintenance and Agricultural Management. Pages 91-114 Advances in Ecological Research. Elsevier.

Geritz, S. A., and K. Éva. 2000. Adaptive dynamics in diploid, sexual populations and the evolution of reproductive isolation. Proceedings of the Royal Society of London. Series B: Biological Sciences **267**:1671-1678.

Gillespie, J. H. 1994. The causes of molecular evolution. Oxford University Press.

Goodman, D. 1982. Optimal life histories, optimal notation, and the value of reproductive value. The American Naturalist **119**:803-823.

Haller, B. C., and A. P. Hendry. 2014. Solving the paradox of stasis: squashed stabilizing selection and the limits of detection. Evolution **68**:483-500.

Hartl, D. L., A. G. Clark, and A. G. Clark. 1997. Principles of population genetics. Sinauer associates Sunderland, MA.

Hassell, M. 1975. Density-dependence in single-species populations. Journal of Animal Ecology:283-295.

Hastings, A. 2004. Transients: the key to long-term ecological understanding? Trends in Ecology & Evolution **19**:39-45.

Hiltunen, T., N. G. Hairston Jr, G. Hooker, L. E. Jones, and S. P. Ellner. 2014. A newly discovered role of evolution in previously published consumer–resource dynamics. Ecology Letters **17**:915-923.

Hin, V., and A. M. de Roos. 2019. Evolution of size-dependent intraspecific competition predicts body size scaling of metabolic rate. Functional Ecology **33**:479-490.

Johansson, J., and U. Dieckmann. 2009. Evolutionary responses of communities to extinctions.

Keith, L. B., and L. A. Windberg. 1978. A demographic analysis of the snowshoe hare cycle. Wildlife Monographs:3-70.

Kentie, R., S. M. Clegg, S. Tuljapurkar, J.-M. Gaillard, and T. Coulson. 2020. Life history strategy varies with the strength of competition in a food-limited ungulate population. Ecology Letters **23**:811-820.

Kooijman, B., and S. A. L. M. Kooijman. 2010. Dynamic energy budget theory for metabolic organisation. Cambridge University Press, Cambridge.

Krebs, C. J., B. Gilbert, S. Boutin, A. R. Sinclair, and J. N. Smith. 1986. Population biology of snowshoe hares. I. Demography of food-supplemented populations in the southern Yukon, 1976-84. Journal of Animal Ecology:963-982.

Kruuk, L. E., T. H. Clutton-Brock, J. Slate, J. M. Pemberton, S. Brotherstone, and F. E. Guinness. 2000. Heritability of fitness in a wild mammal population. Proceedings of the National Academy of Sciences **97**:698-703.

Kruuk, L. E., J. Slate, J. M. Pemberton, S. Brotherstone, F. Guinness, and T. Clutton-Brock. 2002. Antler size in red deer: heritability and selection but no evolution. Evolution **56**:1683-1695.

Kruuk, L. E. B. 2004. Estimating genetic parameters in natural populations using the ‘animal model’. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences **359**:873-890.

Lachish, S., E. Brandell, M. Craft, A. Dobson, P. Hudson, D. MacNulty, and C. T. in press. Investigating the dynamics of elk population size and body mass in a seasonal environment using a mechanistic integral projection model. The American Naturalist.

Lande, R. 1982. A quantitative genetic theory of life history evolution. Ecology **63**:607-615.

Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. Journal of Evolutionary Biology **22**:1435-1446.

Lande, R., S. Engen, and B.-E. Saether. 2003. Stochastic Population Dynamics in Ecology and
Conservation. Oxford University Press, Oxford.
Lande, R., S. Engen, and B.-E. Sæther. 2017. Evolution of stochastic demography with life
history tradeoffs in density-dependent age-structured populations. Proceedings of the
National Academy of Sciences:201710679.
Leslie, P. H. 1945. On the use of matrices in certain population mathematics. Biometrika:183-
212.
Lewontin, R. C. 1969. The meaning of stability. Pages 13-24 *in* Brookhaven Symposia in
Biology.
Lowe, W. H., R. P. Kovach, and F. W. Allendorf. 2017. Population genetics and demography
unite ecology and evolution. Trends in Ecology & Evolution **32**:141-152.
Lundberg, P., E. Ranta, and V. Kaitala. 2000. Species loss leads to community closure. Ecology
Letters **3**:465-468.
Lynch, M., and B. Walsh. 1998. Genetics and Analysis of Quantitative Traits. Sinauer,
Sunderland ,MA.
MacArthur, R., and R. Levins. 1967. The limiting similarity, convergence, and divergence of
coexisting species. The American Naturalist **101**:377-385.
McNamara, J. M., and A. I. Houston. 1996. State-dependent life histories. Nature **380**:215-221.
Merilä, J., B. Sheldon, and L. Kruuk. 2001a. Explaining stasis: microevolutionary studies in
natural populations. Genetica **112**:199-222.
Merilä, J., B. C. Sheldon, and L. E. B. Kruuk. 2001b. Explaining stasis: microevolutionary
studies in natural populations. Genetica **112**:199-222.
Metcalf, C. J. E. 2016. Invisible trade-offs: Van Noordwijk and de Jong and life-history
evolution. The American Naturalist **187**:iii-v.
Metz, J. A., and C. G. de Kovel. 2013. The canonical equation of adaptive dynamics for
Mendelian diploids and haplo-diploids. Interface Focus **3**:20130025.
Moran, D., R. Softley, and E. J. Warrant. 2015. The energetic cost of vision and the evolution
of eyeless Mexican cavefish. Science Advances **1**:e1500363.
Murray, B. G. 1982. On the meaning of density dependence. Oecologia **53**:370-373.
O'Sullivan, R. J., T. Aykanat, S. E. Johnston, A. Kane, R. Poole, G. Rogan, P. A. Prodöhl, C.
R. Primmer, P. McGinnity, and T. E. Reed. 2019. Evolutionary stasis of a heritable
morphological trait in a wild fish population despite apparent directional selection.
Ecology and Evolution.
Ozgul, A., S. Tuljapurkar, T. G. Benton, J. M. Pemberton, T. H. Clutton-Brock, and T.
Coulson. 2009. The dynamics of phenotypic change and the shrinking sheep of St.
Kilda. Science **325**:464-467.
Paine, R. T. 1966. Food web complexity and species diversity. The American Naturalist
100:65-75.
Pigliucci, M., C. J. Murren, and C. D. Schlichting. 2006. Phenotypic plasticity and evolution
by genetic assimilation. Journal of Experimental Biology **209**:2362-2367.
Price, G. R. 1970. Selection and covariance. Nature **227**:520-521.
Rees, M., D. Z. Childs, and S. P. Ellner. 2014. Building integral projection models: a user's
guide. Journal of Animal Ecology **83**:528-545.
Rees, M., and S. P. Ellner. 2019. Why so variable: can genetic variance in flowering thresholds
be maintained by fluctuating selection? The American Naturalist **194**:E13-E29.
Reznick, D., M. J. Butler IV, and H. Rodd. 2001. Life-history evolution in guppies. VII. The
comparative ecology of high-and low-predation environments. The American
Naturalist **157**:126-140.
Reznick, D., and J. A. Endler. 1982. The impact of predation on life history evolution in
Trinidadian guppies (*Poecilia reticulata*). Evolution **36**:160-177.

Reznick, D., L. Nunney, and A. Tessier. 2000. Big houses, big cars, superfleas and the costs of
reproduction. Trends in Ecology & Evolution **15**:421-425.
Reznick, D., and A. P. Yang. 1993. The influence of fluctuating resources on life history:
patterns of allocation and plasticity in female guppies. Ecology **74**:2011-2019.
Reznick, D. N., and H. Bryga. 1987. Life-history evolution in guppies (*Poecilia reticulata*): 1.
Phenotypic and genetic changes in an introduction experiment. Evolution **41**:1370-
1385.
Rohr, J. R., A. A. Elskus, B. S. Shepherd, P. H. Crowley, T. M. McCarthy, J. H. Niedzwiecki,
T. Sager, A. Sih, and B. D. Palmer. 2003. Lethal and sublethal effects of atrazine,
carbaryl, endosulfan, and octylphenol on the streamside salamander (*Ambystoma*
barbouri). Environmental Toxicology and Chemistry: **22**:2385-2392.
Rollinson, N., and L. Rowe. 2015. Persistent directional selection on body size and a resolution
to the paradox of stasis. Evolution **69**:2441-2451.
Romero-Mujalli, D., F. Jeltsch, and R. Tiedemann. 2019. Individual-based modeling of eco-
evolutionary dynamics: state of the art and future directions. Regional Environmental
Change **19**:1-12.
Sæther, B.-E., and S. Engen. 2015. The concept of fitness in fluctuating environments. Trends
in Ecology & Evolution **30**:273-281.
Schindler, D. W. 1974. Eutrophication and recovery in experimental lakes: implications for
lake management. Science **184**:897-899.
Schindler, S., J. M. Gaillard, A. Grüning, P. Neuhaus, L. W. Traill, S. Tuljapurkar, and T.
Coulson. 2015. Sex-specific demography and generalization of the Trivers–Willard
theory. Nature **526**:249.
Schindler, S., P. Neuhaus, J.-M. Gaillard, and T. Coulson. 2013. The influence of nonrandom
mating on population growth. The American Naturalist **182**:28-41.
Schreiber, S. J., S. Patel, and C. terHorst. 2018. Evolution as a coexistence mechanism: Does
genetic architecture matter? The American Naturalist **191**:407-420.
Sibly, R. M., and J. Hone. 2002. Population growth rate and its determinants: an overview.
Philosophical Transactions of the Royal Society of London B: Biological Sciences
357:1153-1170.
Smallegange, I. M., H. Caswell, M. E. M. Toorians, and A. M. Roos. 2017. Mechanistic
description of population dynamics using dynamic energy budget theory incorporated
into integral projection models. Methods in Ecology and Evolution **8**:146-154.
Smith, F. A., and S. K. Lyons. 2013. Animal body size: linking pattern and process across
space, time, and taxonomic group. University of Chicago Press.
Stearns, S. C. 1976. Life-history tactics: a review of the ideas. The Quarterly review of biology
51:3-47.
Steiner, U. K., S. Tuljapurkar, and T. Coulson. 2014. Generation time, net reproductive rate,
and growth in stage-age-structured populations. The American Naturalist **183**:771-783.
Svanbäck, R., and D. I. Bolnick. 2006. Intraspecific competition drives increased resource use
diversity within a natural population. Proceedings of the Royal Society B: Biological
Sciences **274**:839-844.
Traill, L. W., S. Schindler, and T. Coulson. 2014. Demography, not inheritance, drives
phenotypic change in hunted bighorn sheep. Proceedings of the National Academy of
Sciences **111**:13223-13228.
Travis, J., D. Reznick, R. D. Bassar, A. López-Sepulcre, R. Ferriere, and T. Coulson. 2014. Do
eco-evo feedbacks help us understand nature? Answers from studies of the Trinidadian
guppy. Advances in Ecological Research **50**:1-40.
Tuljapurkar, S. 1989. An uncertain life: demography in random environments. Theoretical
Population Biology **35**:227-294.

Tuljapurkar, S., J.-M. Gaillard, and T. Coulson. 2009a. From stochastic environments to life
histories and back. Philosophical Transactions of the Royal Society of London B:
Biological Sciences **364**:1499-1509.
Tuljapurkar, S., and S. H. Orzack. 1980. Population dynamics in variable environments I.
Long-run growth rates and extinction. Theoretical Population Biology **18**:314-342.
Tuljapurkar, S., U. K. Steiner, and S. H. Orzack. 2009b. Dynamic heterogeneity in life
histories. Ecology Letters **12**:93-106.
Van Noordwijk, A. J., and G. de Jong. 1986. Acquisition and allocation of resources: their
influence on variation in life history tactics. The American Naturalist **128**:137-142.
Walters, C. J., and F. Juanes. 1993. Recruitment limitation as a consequence of natural
selection for use of restricted feeding habitats and predation risk taking by juvenile
fishes. Canadian Journal of Fisheries and Aquatic Sciences **50**:2058-2070.
Warner, R. R., and P. L. Chesson. 1985. Coexistence mediated by recruitment fluctuations: a
field guide to the storage effect. The American Naturalist **125**:769-787.
Werner, T. K., and T. W. Sherry. 1987. Behavioral feeding specialization in Pinaroloxias
inornata, the "Darwin's finch" of Cocos Island, Costa Rica. Proceedings of the National
Academy of Sciences **84**:5506-5510.
West, G. B., J. H. Brown, and B. J. Enquist. 1997. A general model for the origin of allometric
scaling laws in biology. Science **276**:122-126.
Wright, J. S. 2002. Plant diversity in tropical forests: a review of mechanisms of species
coexistence. Oecologia **130**:1-14.
Wright, S. 1969. Evolution and the Genetics of Populations: Vol. 2. The Theory of Gene
Frequencies. University of Chicago Press, Chicago.
Yoshida, T., L. E. Jones, S. P. Ellner, G. F. Fussmann, and N. G. Hairston Jr. 2003. Rapid
evolution drives ecological dynamics in a predator–prey system. Nature **424**:303.
1318