Title: Intermittent hormonal therapy shows similar outcome than SOC in ER+ breast cancer preclinical model.

Authors: Pedro M. Enriquez-Navas[1], Libia Garcia[1], Mahmoud Abdalah[2], Olya Stringfield[2], Kim Luddy[1], Sabrina Hassan[1], Robert J. Gillies[1], and Robert A. Gatenby[1].

Affiliations: (1) Department of Cancer Physiology and (2) Image Response Assessment Team (IRAT) Core, H. Lee Moffitt Cancer Center & Research Institute. 12902 Magnolia Dr. Tampa (FL) 33612.

Corresponding Author

Pedro M. Enriquez-Navas. Pedro.enriqueznavas@mofitt.org. 813-745-1417.

Abstract:

Clinical breast cancers in which at least 10% of cells express the estrogen receptor are labeled as "ER positive." First line therapy for these patients is typically continuous administration of anti-estrogen drugs at maximum tolerated dose (MTD) until progression. In the vast majority of patients, resistance to hormone therapy evolves in the breast cancer cells within 2 years leading to treatment failure and tumor progression. In prior studies, we have demonstrated continuous application of MTD chemotherapy results in evolutionary dynamics (termed "competitive release") that accelerates proliferation of treatment-resistance populations. In contrast, evolution-informed application of treatment reduces drug administration to maintain substantial populations of therapy-sensitive cells to reduce proliferation of resistant phenotypes. Prior pre-clinical and clinical studies have shown this strategy can delay or prevent proliferation of resistant cells and prolong time to progression (TTP). We hypothesize that similar dynamics may be observed in hormonal therapy of ER+ breast cancers. Here we address two important dynamics. First, we consider a clinical scenario in which symptoms are sufficiently severe or life-threatening to require rapid and substantial tumor reduction. Can this be achieved while retaining evolutionary dynamics to subsequently delay proliferation of resistance? A second, related question is defining the cost of resistance to anti-estrogen therapy. Here, we investigated the evolutionary dynamics of resistance to anti-estrogen therapy using ER+ MCF-7 orthotropic xenografts treated with both continuous Tamoxifen as well as cycles in which estrogen stimulation is combined with estrogen suppression. As expected, continuous administration of anti-estrogen drugs successfully suppressed tumor growth. However

we found that brief interruptions in drug administration permitted equal tumor control

while administering up to 50% less drug and maintaining cell phenotypes that retained

high levels of ER expression and lower levels of MDR1 expression. In follow-on

experiments combining hormonal and chemo- therapies; we obtained similar tumor

control to hormonal therapy alone but with more necrosis and significantly lower ER

expression in the surviving population.

Introduction:

In 2017 about 300,000 women in the United States were diagnosed with breast

cancer and, among these, almost 90% were characterized as estrogen receptor positive

(ER+) [1, 2]. The criteria for ER+ requires that at least 10% of the tumor cells express

ER on immunohistochemical staining so that, in many breast cancers, a significant

fraction of the cancer cells do not express ER and may be resistant to anti-estrogen

therapy [3].

Initial treatment for ER+ breast cancer includes blockade of the effect of estrogen

by selective estrogen receptor modulator (SERM) drugs.[4] Tamoxifen, which blocks

the interaction between estrogen and its receptor impeding cell replication, is among the

most widely used SERM drugs and is typically administered at maximum tolerated dose

daily until tumor progression [5, 6]. Alternative strategies, such as aromatase inhibitors,

block the synthesis of estrogen by normal cells. While nearly all ER+ tumors initially

respond to anti-estrogen therapy, evolution of resistance with treatment failure and

tumor progression typically is observed within a few months to a few years[7].

There are three major drawbacks in anti-estrogen treatment: (1) cost and side

effects reduce compliance in up to 25% of the patients [8]; (2) about 10% will develop

one or more side-effects that require dose adjustments or treatment cessation [1, 7, 9];

(3) prolonged continuous treatment may significantly increase risk for endometrial

cancer. Nevertheless, the vast majority of patients initially respond to anti-estrogen

therapy but development of resistance leading to treatment failure and progression is

virtually inevitable [10-12] and, among the multiple mechanisms of resistance, evolution

of estrogen-independent growth is most common[11].

A common evolution-based strategy to delay tumor progression focuses on the

phenotypic cost of resistance. This is readily apparent in resistance to chemotherapy

based on increased expression of MDR1 (Multi-Drug Resistance system-1); a

membrane glycoprotein (PgP) that is an active ATPase pump extruding lipophilic

cationic xenobiotics. In some studies, up to 40% of a cell's energy budget must be used

to synthesize, maintain, and operate MDR membrane pumps – an obvious cost of

resistance [13, 14]. Thus, a strategy termed "adaptive therapy" explicitly limits cancer

treatment to maintain a significant population of treatment-sensitive cells. Therapy is

then withdrawn. However, in the subsequent tumor regrowth, the sensitive cells, in the

absence of the cost of resistance, outcompete the resistant cells. Thus, through multiple

cycles, the tumor population remains sensitive to the primary treatment. In an ongoing

clinical trial in metastatic castrate-resistant prostate cancer, we have found that

treatment that only reduces the serum PSA to half of its pre-treatment value can both

substantially increase the time to progression while decreasing the cumulative drug

does [15].

A number of questions regarding optimal evolution-based treatments remain.

Among these, perhaps the most urgent is the apparently conflicting demands for

treatment in which a patient presents with highly symptomatic or potentially life-

threatening conditions. Here, rapid and significant reduction of the tumor burden is

clinically necessary but could also result in competitive release of resistant clones that

result in rapid proliferation leading to tumor failure and tumor progression with recurrent

symptoms. Here we examine potential treatment strategies that can both rapidly

diminish tumor burden to very low levels while maintaining evolutionary dynamics that

can prolong tumor control and reduce the cumulative drug dose to reduce toxicity and

cost.

A second question in this study is the cost of resistance to hormonal therapy.

Although drug efflux by the MDR proteins is a mechanism of resistance in ER+ cells,

clinical studies have found that, in general, durable resistance to estrogen therapy is

most commonly obtained in breast cancer cells through expression of alternative

pathways that permit estrogen-independent survival and proliferation. However, unlike

the dynamics of PgP, the evolutionary cost of estrogen-independence is not obvious.

Nevertheless, that such a cost exists can be inferred using a concept termed

"evolutionary triage." Briefly, "evolutionary triage"[15] simply states that, among

competing populations, the fittest phenotype will be the most proliferative and, in

general, be the largest population. Therefore, in general, the relative fitness of each

cancer subpopulation can be estimated by their relative abundance within the tumor.

This, however, yields puzzling results for ER+ breast cancers in which greater than 50%

of the cells do not express the ER on immunohistochemical stains. Despite ER+ cells

being in the minority, these tumors still typically respond, at least initially, to anti-

estrogen therapy suggesting some component of the treatment dynamics is not being

captured in the IHC results. Thus, it is not clear if evolution-based treatment strategies

could be successfully applied in clinical treatment of ER+ metastatic breast cancer.

Here, we address these questions in pre-clinical studies. Our results showed that

(i) intermittent therapies can control tumor growth with less Tamoxifen (using as little as

50% of the standard dose), (ii) the ER expression is maintained at or above the levels of

SOC, (iii) expression of MDR1 was reduced in tumors treated with intermittent

tamoxifen therapy, and (iv) the combination of hormonal- and chemo- therapies in the

presence of high levels of estrogen kept the tumor at the same volume than the

standard therapy, although showing more tumor necrosis.

We conclude that evolution-based administration of anti-estrogen drugs in

patients is likely to benefit patients with metastatic ER+ breast cancer compared to

current strategies of continuous MTD dosing until progression. Our results, however,

also suggest the evolutionary dynamics that govern estrogen-related fitness in breast

cancer cells and the clinical efficacy of anti-estrogen therapy are not fully understood

and require further investigation.

Materials and Methods

In-vivo Experiments

Different cohorts (n= 10, 14, 4, and 39) of nude (nu/nu) mice were injected in the

mammary fat pad with $5*10^6$ MCF7 cells (injected in a mixture 1:1 with phenol-red free

Matrigel) tagged with GFP (cells were obtained from ATCC, and were grown following

its guidelines, cell culture media and supplies were obtained from Thermo Fisher

Scientific). Prior to the cell injections, an estrogen pellet (Innovative Research of

America) of 0.72 mg of β-estrogen with 90 days slow release was subcutaneously

implanted in each mouse, giving a continuous dose of 400 pg of estrogen per milliliter of

blood. If the experiments lasted longer than 90 days, a similar pellet was implanted to

the mice.

When tumors reached 300 mm^3 approximately, mice were randomly distributed

in groups and treated following one of the subsequent treatments: *Controls* (Ctrl), no

treatment; *Tamoxifen standard* (TamST), 0.5 mg of tamoxifen per mouse daily;

Tamoxifen-vacation (TamVac), 2 weeks of TamST followed by one week of no

treatment (vacation); *Tamoxifen 2 weeks* (Tam2weeks), two weeks of TamST followed

by 2 weeks of vacation; *Tamoxifen 3 weeks* (Tam3weeks), three weeks of TamST

followed by three weeks of vacation; *Tamoxifen and Paclitaxel* (TamPac), 2 weeks of

TamST followed by one week in which Paclitaxel is applied in two non-consecutive days

at a concentration of 20mg/kg (ip) (PacST); *Vacation and Paclitaxel* (VacPac), 2 weeks

of no treatment followed by one week in which 20 mg/kg of Paclitaxel in applied in two

non-consecutive days; *Paclitaxel and Tamoxifen* (PacTam), one week of PacST

followed by two weeks of TamST.

Tamoxifen (Caiman Chemical Company) was suspended in peanut oil (Sigma

Aldrich) and given to animals by either i.p or gavage routes; initially Tamoxifen was

administered in 200 µl of peanut oil through i.p. injection. However, we changed the

route of administration to gavage because mice were not able to metabolize the peanut

oil and all of them died in a short period of time; maybe due to peritonitis (when tumors

were collected, we noticed that peanut oil was accumulated in mice abdomen).

Paclitaxel (LC laboratories) was dissolved in a mixture of Koliphor oil and ethanol (1:1,

both solvents were obtained from Sigma Aldrich) and it was given via i.p. injection;

before injection, the corresponding dose was diluted twice with PBS.

Mice were visually monitored, and their weights monitored once a week, to address any

treatment toxicity issues. Tumor growth was measured once a week by either caliper,

using the formula: $vol = \frac{\pi}{6} * \frac{(short\ distance)^2}{long\ distance}$, or by MRI; magnetic resonance data were

acquired with a 7 T horizontal magnet Agilent ASR 310 (Agilent Technologies Inc.)

equipped with nested 205/120/HDS gradient insert and a bore size of 310 mm. Before

imaging, the animals were placed in an induction chamber and anesthetized with 2 -3%

isoflurane delivered in 1.5 liter/min oxygen ventilation. After complete induction, animals

were restrained in a custom-designed holder and inserted into the magnet while

constantly receiving isoflurane (1 to 3%) within the same oxygen ventilation. Body

temperature (37° ± 1°C) and respiratory functions were monitored continuously (SAII

System) during the experimental time. A 35 mm Litzcage coil (Doty Scientific) was used

to carry out axial T2-weighted fast spin-echo multislice experiments (acquired with

TE/TR [echo time/repetition time] = 72 ms/1000 ms, field of view (FOV) = 35 × 35 mm^2,

matrix = 128 × 128, yielding a spatial in-plane resolution of 273 μm, slice thickness of

1.5 mm).

At the completion of the experiments (either tumor under control or when tumor

volume reached ~2000 mm^3) tumors were collected for histological analysis. After

collection, tumors were processed for histological studies by soaking in formalin, during

24 h at least, followed by embedding in paraffin blocks. Consecutive histological slices

(5 μm thickness) were cut from each tumor to study the necrosis percentage (H&E),

vascularity density and functionality (CD31 and SMA, respectively), estrogen receptor

(ER) expression, and resistance mechanisms (by means of MDR1 expression). Once

stained, the slices were imaged at the Moffitt Cancer Center microscope core facilities,

using Aperio ScanScope XT microscope and Aperio Spectrum version 10.2.5.2352

image analytic software (Leica Biosystems Inc.). To optimize the image analysis, we

trained the analytic algorithm by using ROIs that were selected manually to represent

ROIs that are positive or negative for each stain. After initial algorithm training, the

software developed a final algorithm, which was used to automatically analyze the

slides with a pixel-size resolution (5 μm x 5 μm).

Imaging analysis

To analyze the homogeneity in the ER expression, a radiomics analysis was

performed on IHC estrogen receptor images. Features extraction was done on IHC

digital images with the same magnification (x20). Color images were segmented using a

thresholding method to identify the pixels that were positively stained. Pixels outside the

segmented range (background and/or unstained cells) were discarded and not used for

analyses. Masks from positively stained cells were used to generate neighborhood

maps. For each pixel in the mask, the number of immediate neighbors that were also in

the mask (positively stained) was counted (up to 8) and that number is the pixel's

neighborhood coefficient. Thus, for all samples, the masks of positively stained pixels

were replaced by corresponding neighborhood maps consisting of neighborhood

coefficients. The neighborhood maps were computed to capture the distribution and the

density of positively (ER+) stained pixels.

Following the generation of neighborhood maps, 202 2D image features were

calculated from them, including statistical, shape, and texture variables. These features

were then reduced by including only one feature from subsets of inter-correlated

features (Pearson correlation coefficient > |0.8|). Among the final features list, only

those features correlated with heterogeneity in the tissue texture were used to perform

the analysis; table 1 shows a brief description of these features.

All the animal work during this project was done following the IACUC regulations of

University of South Florida (Tampa) at the Moffitt Cancer Center facilities.

Statistical calculations were performed using the Excel software. Student's t-tests

were performed considering two-tailed distribution and two samples with unequal

variance.

Results

This project was designed to understand the evolutionary dynamics of resistance

in ER+ breast tumors and, consequently, to improve first-line treatment of estrogen

receptor positive (ER+) breast cancers with SERMs, such as Tamoxifen.

Treatment algorithms used in this study were suggested by preliminary *in-vitro* data in

which MCF7 cells were grown under different microenvironment conditions to study the

expression of the estrogen receptor following addition of Tamoxifen and/or Paclitaxel

media.

In the first cohort of mice (n = 10) bearing orthotopic MCF7 tumors, Tamoxifen

treatment was administered by i.p. injections. As it is shown in **Figure 1A**, treatment

algorithm with a one-week vacation (TamVac) maintained tumors at similar volumes as

standard treatment (TamST). No increase in tumor volume was observed during the

vacation period. Furthermore, at completion of the study, the remaining tumor cells

demonstrated greater ER expression compared to the control and equal or greater

expression the continuous dose cohort **(Figure 2A).**

TamST treatment achieved the same level of control as continuous high dose

tamoxifen. This is in contrast to prior reports in which this combination was

unsuccessful. [16-18]. It is possible this difference is due to treatment schedule. We

treated the animals 7 days per week (matching the daily dose used clinically) and it

appears that in the prior studies treatment was not administered on weekends. While

the results were encouraging, we noted that all of the mice developed increased

peritoneal fluid that appeared to be caused by the peanut oil used in the Tamoxifen

injections.

In a second cohort (n=14) we used higher concentration of Tamoxifen (50 mg/ml)

with 20 µl of the suspension injected i.p. daily. This reduced the peritoneal fluid

collection and showed the same outcome as the prior cohort (**Figure 1B**).

Histological analysis of ER expression showed no significant difference between the

intermittent and standard tamoxifen therapies (**Figure 2B**). However, the vascularity

(density and functionality) was decreased in TamVac therapy (**Figure 3**, columns A and

B). MDR1 was expressed in the Tamoxifen therapy groups suggesting membrane

extrusion play a significant role in evolution of resistance in this setting (**Figure 4B**).

Finally, because the ip injection of Tamoxifen was associated with peritoneal lipid

collections, we examined an additional cohort of mice (n=41) in which Tamoxifen was

administered by gavage (no significant difference was found between ip and gavage

treatments, **Sup. Figure 1**). This cohort included prior treatment (continuous Tamoxifen

dosing and one-week vacation period) but also examined longer "vacation" periods

during which treatment was suspended for 2 or 3 weeks (Tam2weeks and Tam3weeks).

We also examined alternative sequences with Tamoxifen first followed by Paclitaxel or

vice-versa (TamPac and PacTam, respectively).

After 130 days of treatment, no significant differences in tumor control were

noted in the groups (**Figure 1C**). However, TamVac, Tam2weeks, and Tam3weeks

groups had a cumulative dose reduction of 33, 50, and 50%, respectively. No

significant tumor growth during these vacation periods was noted (**Figure 1** (grey zones

represents vacations periods)). At necropsy, the tumors treated with vacation periods

had slightly higher expression of estrogen receptor compared to continuous Tamoxifen

(**Figure 2C and Sup. Figure 2**). Interestingly, vessel density and functionality were

increased in tumors in which hormonal therapy was combined with a cytotoxic drug

(**Figure 3**) but these tumors also demonstrated relatively larger fractions of necrosis

(**Figure 3** and **Sup. Figure 3**) when compared to the other cohorts.

We studied the expression of MDR1 systems for cohorts B and C using

immunohistochemistry (IHC) (**Figure 4**). In general, cohorts with vacation periods

showed lower expression of MDR1 than the continuous Tamoxifen group (**Figure 4**, B

and C) reflecting the diminished selection pressure for resistance during treatment

vacation.

We analyzed the homogeneity in the ER expression in the treatment cohorts

using a "neighborhood" imaging analysis, which examined variations in ER expression

in physically adjacent cell groups. To do so, we created a mask for the tumor slices

following IHC staining for ER expression. An algorithm calculates probability that each

ER+ pixels will have similar adjacent pixels (up to 8). This analysis found increased

numbers of ER+ similar "neighbors" in treated tumors with either Tamoxifen SOC or

intermittent therapies compared to tumors also treated with chemotherapy (**Figure 5**).

Discussion

Cancer cells, like all living systems, evolve to adapt to local environmental

selection forces. When clinical therapy is applied, evolution of resistance is commonly

observed leading to treatment failure and tumor progression. Here we examine the

evolutionary dynamics of ER+ breast cancer treated with anti-estrogen therapy, the

typical first line clinical treatment for ER+ breast cancers. In prior pre-clinical and

theoretical studies we have found that continuous application of therapy at MTD

maximally selects for resistance – a well-known phenomenon in pest management

termed "competitive release."   By periodically withdrawing therapy, we reduced the

environmental selection forces for resistance by permitting survival of some treatment-

sensitive cells. In the absence of treatment, the fitness advantage of the sensitive cells

tended to suppress proliferation of the resistant phenotype thus prolonging tumor

response. Here we addressed two potential barriers for applying this strategy to ER+

clinical breast cancers. First, in patients who are symptomatic, optimal therapy must

reduce the tumor burden below some symptomatic threshold before it is withdrawn.

Second, we were concerned with the possibility that the tumor might rapidly progress

after treatment withdrawal leading to rapid loss of control and thus decreased TTP.

In these pre-clinical experiments with ER+ breast cancers, we demonstrated

intermittent application anti-estrogen drugs could achieve complete tumor control

identical to that obtained with continuous MTD treatment. No tumor growth was

observed even during a 3-week interval during which therapy was not applied.

Advantages of this therapy included a significant (up to 50%) cumulative dose reduction

and decreased evidence tumor cell resistance (based on ER and MDR1 expression).

Our study does not demonstrate that Tamoxifen intermittent therapy can prolong

response as it we have found in chemotherapy for breast cancer in pre-clinical studies

and hormone therapy in prostate cancer. Here we were primarily focused on

experiments that address a clinical scenario in which the patient presents with highly

symptomatic or life-threatening disease requiring rapid and significant reductions of the

tumor burden. We demonstrate that such treatment can be administered to

substantially reduce the tumor burden while also using interruptions of therapy to

reduce evolutionary selection for resistant populations while still maintaining tumor

control. Furthermore, these outcomes can be achieved while substantially reducing (by

up to 50%) the total dose of Tamoxifen thus reducing toxicity and cost.

Finally, by imaging analysis (radiomics), we have demonstrated that

evolutionary-based Tamoxifen therapies develop tumors with the same ER

homogeneity than SOC. These results suggest that imaging biomarkers that correlate

with intratumoral evolution during treatment may ultimately prove to be useful guides for

evolution-based treatments.

Acknowledgements: This work has been supported in part by the SAIL Core, Tissue Core

Facility, the Analytic Microscopy Core Facility, and by the IRAT Core Facility at the H. Lee

Moffitt Cancer Center & Research Institute, an NCI designated Comprehensive Cancer Center

(P30-CA076292).

References:

1. http://www.breastcancer.org/. 2016.
2. ACS, *www.cancer.org.* 2016.
3. Viale, G., *The current state of breast cancer classification.* Ann Oncol, 2012. **23 Suppl 10**: p. x207-10.
4. Lumachi, F., et al., *Treatment of estrogen receptor-positive breast cancer.* Curr Med Chem, 2013. **20**(5): p. 596-604.
5. Burstein, H.J., et al., *Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: american society of clinical oncology clinical practice guideline focused update.* J Clin Oncol, 2014. **32**(21): p. 2255-69.
6. Lumachi, F., D.A. Santeufemia, and S.M. Basso, *Current medical treatment of estrogen receptor-positive breast cancer.* World J Biol Chem, 2015. **6**(3): p. 231-9.
7. Shapiro, C.L. and A. Recht, *Side effects of adjuvant treatment of breast cancer.* N Engl J Med, 2001. **344**(26): p. 1997-2008.
8. Partridge, A.H., et al., *Nonadherence to adjuvant tamoxifen therapy in women with primary breast cancer.* J Clin Oncol, 2003. **21**(4): p. 602-6.
9. www.cancerresearchuk.org. *Tamoxifen*. 2016.
10. Colleoni, M., et al., *Annual Hazard Rates of Recurrence for Breast Cancer During 24 Years of Follow-Up: Results From the International Breast Cancer Study Group Trials I to V.* J Clin Oncol, 2016. **34**(9): p. 927-35.
11. Chang, M., *Tamoxifen Resistance in Breast Cancer.* Biomolecules & Therapeutics, 2012. **20**(3): p. 256-267.
12. Hayes, E.L. and J.S. Lewis-Wambi, *Mechanisms of endocrine resistance in breast cancer: an overview of the proposed roles of noncoding RNA.* Breast Cancer Res, 2015. **17**: p. 40.
13. Kam, Y., et al., *Sweat but no gain: Inhibiting proliferation of multidrug resistant cancer cells with "ersatzdroges".* Int J Cancer, 2014.
14. Silva, A.S., et al., *Evolutionary approaches to prolong progression-free survival in breast cancer.* Cancer Res, 2012. **72**(24): p. 6362-70.
15. Zhang, J., et al., *Integrating evolutionary dynamics into treatment of metastatic castrate-resistant prostate cancer.* Nature Communications, 2017. **8**(1): p. 1816.
16. Osborne, C.K., K. Hobbs, and G.M. Clark, *Effect of estrogens and antiestrogens on growth of human breast cancer cells in athymic nude mice.* Cancer Res, 1985. **45**(2): p. 584-90.
17. Osborne, C.K., et al., *The importance of tamoxifen metabolism in tamoxifen-stimulated breast tumor growth.* Cancer Chemother Pharmacol, 1994. **34**(2): p. 89-95.
18. Whitesell, L., et al., *HSP90 empowers evolution of resistance to hormonal therapy in human breast cancer models.* Proc Natl Acad Sci U S A, 2014. **111**(51): p. 18297-302.

Figures:

Figure 1: Tumor volumetric data of mice under different treatments. Grey zones

correspond to either vacation periods or paclitaxel application (more information about

each treatment can be found in Material and Methods section). Tumor volumes were

measured by MRI. In parenthesis is the number of mice in each group. Data are shown

by mean and error bars represent the standard deviation.

Figure 2: IHC analysis of the ER expression under different treatments. (Significance calculations were done by using a Student t-test with two-tailed distribution and considering two samples with unequal variance).

Figure 3: IHC analysis of viable and necrotic tissues (H&E), vessel density (CD31) and

functionality (SMA) in tumors under different treatments. Data are shown by mean

values with the standard deviation (error bars). p value calculated using Student t- test

with two-tailed and unequal variance.

Figure 4: IHC analysis of the expression of MDR1 systems for cohorts B and C. Data is

shown by mean with the standard deviation (error bars).

385 **Figure 5:** Radiomics analysis of ER expression. The neighbor map (ER+ pixels
386 surrounded by similar ones) showing the mean value. Mean±SD is represented in bar
387 graph. p-values are shown to compare the treated groups with the control (untreated)
388 one.

Sup. Figure 1: Fold of change in tumor volume in mice treated with tamoxifen by either

391   i.p. injections or gavage.

394 **Sup. Figure 3:** Cohort *C* IHC analysis of the ER expression under different treatments.

395 (Significance calculations were done by using a Student t-test with two-tailed distribution

396 and considering two samples with unequal variance).

Sup. Figure 4: Cohort *C* IHC analysis of viable and necrotic tissues (H&E), vessel

density (CD31) and functionality (SMA) in tumors under different treatments.