1 **Regeneration in the absence of a blastema requires cell**
2 **division but is not tied to wound healing in the ctenophore**
3 *Mnemiopsis leidyi*
4
5

6 Authors:
7 Julia Ramon-Mateu[1], Tori Ellison[2], Thomas E. Angelini[2,3], Mark Q. Martindale[1*]
8
9

Affiliation:
[1]The Whitney Laboratory for Marine Bioscience, 9505 N, Ocean Shore Blvd, St.
Augustine, FL 32080-8610, USA
[2]Department of Materials Science and Engineering, Herbert Wertheim College of
Engineering, University of Florida, Gainesville, Florida, 32611, USA
[3]Department of Mechanical and Aerospace Engineering, Herbert Wertheim College of
Engineering, University of Florida, Gainesville, Florida, 32611, USA
17
18
19 *Corresponding author
20 Email: mqmartin@whitney.ufl.edu
21

ABSTRACT

Background:

The ability to regenerate is a widely distributed but highly variable trait among metazoans. A variety of modes of regeneration has been described for different organisms, however, many questions regarding the origin and evolution of these strategies remain unanswered. Most species of ctenophore (or "comb jellies"), a clade of marine animals that branch off at the base of the animal tree of life, possess an outstanding capacity to regenerate. However, the cellular and molecular mechanisms underlying this ability are unknown. We have used the ctenophore *Mnemiopsis leidyi* as a system to study wound healing and adult regeneration and provide some first-time insights of the cellular mechanisms involved in the regeneration of one of the most ancient extant group of multicellular animals.

Results:

We show that cell proliferation is activated at the wound site and is indispensable for whole-body regeneration. Wound healing occurs normally in the absence of cell proliferation forming a scar-less wound epithelium. No blastema-like structure is generated at the cut site, rather undifferentiated cells assume the correct location of missing structures and differentiate in place. Pulse-chase experiments and surgical intervention show that cells originating in the main regions of cell proliferation (the tentacle bulbs) do not seem to contribute to the formation of new structures after surgical challenge, suggesting a local source of cells during regeneration. While exposure to cell-proliferation blocking treatment inhibits regeneration, the ability to regenerate is recovered when the treatment ends (days after the original cut), suggesting that ctenophore regenerative capabilities are constantly ready to be triggered and they are somehow separable of the wound healing process.

Conclusions:

Ctenophore regeneration takes place through a process of cell proliferation-dependent non blastemal-like regeneration and is temporally separable of the wound healing process. The remarkable ability to replace missing tissue, the many favorable experimental features (e.g. optical clarity, high fecundity, rapid regenerative performance, stereotyped cell lineage, sequenced genome), and the early branching phylogenetic position in the animal tree, all point to the emergence of ctenophores as a new model system to study the evolution of animal regeneration.

Keywords: Ctenophore, regeneration, wound healing, wound epidermis, blastema, stem cells, dedifferentiation, *Mnemiopsis leidyi.*

BACKGROUND

Regeneration, the ability to re-form a body part that has been lost, is a widely shared property of metazoans (1). However, the contribution of cell proliferation, the source of regenerating tissue, and the mechanisms which pattern the replaced tissues vary greatly among animals with regenerative ability, resulting in a collection of different "modes" of regeneration (2,3). Based on the involvement of cell proliferation, there have been described cases of regeneration in which the restoration of the missing body part is accomplished in the absence of cell proliferation, by the remodeling of pre-existing cell tissues; on the other hand, and also more common, is the strategy of regeneration which requires active cell proliferation. Cell proliferation-based regeneration can involve the production of a regeneration-specific structure, the blastema. Several different definitions of a "blastema" have been proposed depending on the model system and its regenerative strategy. For example, in amphibians, the blastema is described as an unpigmented outgrowth consisting of a mass of undifferentiated progenitor cells that forms at the wound site from where cells proliferate and differentiate to form the missing structures (4,5); while in planarians, the blastema is composed of post-mitotic progeny of proliferating cells that differentiate to reform the lost tissue (6). Given this lack of consensus around the word "blastema" and based on the biology, morphological features and regenerative response of our organism of study, we define the regeneration blastema as a "field" of undifferentiated cells that accumulate at the wound site and are later patterned to give rise to the appropriate set of missing structures but remains agnostic about their origin or their proliferative status.

The classical example of cell proliferation-independent regeneration is provided by the freshwater cnidarian polyp *Hydra*, which is able to regenerate the head after decapitation through remodeling of the pre-existing tissue without a significant contribution from cell proliferation (7-11). While documented cases of strict cell proliferation-independent regeneration are very few, most of the organisms with regenerative potential rely on cell proliferation – or a combination of both cell proliferation and tissue remodeling – to re-form lost structures. Regenerative abilities also appear to be diverse even within individual evolutionary clades. For example, regeneration of oral structures in another member of the phylum Cnidaria – *Nematostella vectensis* – is characterized by high levels of cell proliferation thus, differing from the cell proliferation-independent regeneration potential in *Hydra* (12). In planarians, whole-body regeneration is accomplished by the proliferation of pluripotent stem cells (neoblasts), the only cells in the adult with proliferative potential, which form a mass of undifferentiated cells known as the regenerating blastema (13–15). Annelid

regeneration provides examples of both blastema-based regeneration and tissue-
remodeling based regeneration (16,17), showing diversity within the Lophotrochozoa.
Moreover, evidence of cell migration has been documented during regeneration of
several annelid species such as the freshwater annelid *Pristina leidyi* (18) and the
marine annelid worm *Capitella teleta,* in which local (proliferating cells close to the
wound site) and distant (stem cell migration) sources of cells contribute to the
formation of the regenerating blastema (19). Evidence of cell migration during
regeneration is also provided by the hydrozoan *Hydractinia echinata* in which stem
cells (i-cells) from a remote area migrate to the wound site and contribute in the
formation of the blastema (20). In vertebrates, regenerative potential is limited primarily
to the structural or cellular level. Urodele amphibians are known for being the only
vertebrate tetrapods that can regenerate amputated limbs as adults. Similar to the
previous examples of cell proliferation-based regeneration, they require cell
proliferation and the formation of a blastema. However, the urodele blastema is not
generated from or composed of cells of a single type, but consists of a heterogeneous
collection of lineage-restricted progenitors (21).

Among the animals with impressive whole-body regenerative capabilities are
lobate ctenophores (comb jellies), fragile holopelagic marine carnivores that represent
one of the oldest extant metazoan lineages. Ctenophora is latin for "comb bearer",
referring to eight longitudinally oriented rows of locomotory ctene (or comb) plates
which they coordinately beat to propel through the water column. Ctenphores possess
a highly unique body plan characterized by a biradial symmetry (with no planes of
mirror symmetry) and two epithelial layers: the ectoderm and the endoderm, separated
by a thick mesoglea mostly composed of extracellular matrix, but also containing
several types of individual muscle and mesenchymal cells. The oral-aboral axis is their
major body axis and it is characterized by the mouth at one (oral) pole and the apical
sensory organ containing a statocyst at the opposite (aboral) pole. Most ctenophores
possess a pair of muscular tentacles that bear specialized adhesive cells called
colloblasts, used to capture prey (22) (**Figure 1C**). One of the best studied species of
ctenophores is the lobate ctenophore *Mnemiopsis leidyi,* which is emerging as a new
model system in evolutionary-developmental biology (23-28). *M. leidyi*'s life cycle is
characterized by a rapid development including a highly stereotyped cleavage program
and two adult stages: the juvenile tentaculate cydippid, distinguishable for having a pair
of long branching tentacles (**Figure 1A, 1B**), and the larger lobate adult form which
possess two oral feeding lobes. A particular feature of ctenophore embryogenesis is
that they undergo mosaic development, meaning that embryos cannot compensate for
cells/structures derived from cells killed or isolated during early development. If

blastomeres are separated at the two-cell stage, each will generate a "half-animal,"
possessing exactly half of the normal set of adult features (29,30). This lack of ability to
replace missing parts during embryogenesis contrasts with the outstanding capacity to
regenerate as adults. Both the tentaculate larval and lobate adult life stages of *M. leidyi*
readily regenerate and are capable of whole-body regeneration from only a single body
quadrant (30).

It has been known for well over 80 years that ctenophores have the capacity to
replace missing body parts (30-33) but the cellular and molecular mechanisms
underlying this ability are poorly understood. Is cell proliferation required for ctenophore
regeneration? Is any kind of blastema-like structure formed during regeneration? What
is the source and nature of cells that contribute to the regenerated structures? What is
the role of the wound epidermis in regulating the future regenerative outcome? We
have studied wound healing and adult regeneration in the ctenophore *Mnemiopsis*
leidyi and show that cell proliferation is activated at the wound site several hours after
wound healing is complete and is indispensable for the regeneration of all the
structures of the cydippid's body. Wound healing occurs normally in the absence of cell
proliferation forming a scar-less wound epithelium only a few hours after amputation. In
both animals cut in half along the oral-aboral axis and those in which the apical organ
is removed, anlage of all missing structures occurs within 24-48 hours and complete
replacement of all cell types by 72 hours after the injury. No blastema-like structure is
generated, rather undifferentiated cells assume the correct location of missing
structures and differentiate in place. EdU (5-ethynyl-2'-deoxyuridine) labeling shows
that in uncut animals the majority of cell divisions occur in the tentacle bulbs where the
tentacles are continuously growing. In surgically challenged animals, cell division is
stimulated at the wound site between 6-12 hours after injury and continues until 72
hours after injury. EdU pulse and chase experiments after surgery together with the
removal of the two main regions of active cell proliferation suggest a local source of
cells in the formation of missing structures. Although the appearance of new structures
is completely dependent on cell division, surprisingly, the ability to regenerate is
recovered when exposure to cell-proliferation blocking treatment ends, suggesting that
the onset of regeneration is constantly ready to be triggered and it is somehow
temporally separable from the wound healing process. This study provides some first-
time insights of the cellular mechanisms involved in ctenophore regeneration and
paves the way for future molecular studies that will contribute to the understanding of
the evolution of the regenerative ability throughout the animal kingdom.
168
169   **RESULTS**

Whole-body regeneration in *Mnemiopsis leidyi* cydippids

Although the regenerative response has been studied previously in *M. leidyi* (23, 30-33) we first characterized the sequence of morphogenic events during cydippid wound healing and regeneration to provide a baseline for further experimental investigations. For this, two types of surgeries – representing the replacement of all the structures and cell types of the cydippid's body (e.g. apical organ, comb rows, tentacle bulbs and tentacles) – were performed: 1) Bisection through the oral aboral axis keeping the piece with an intact apical organ which required the remaining piece to regenerate 4 comb rows and a tentacle apparatus (bisected cydippids with a complete intact apical organ regenerate into whole animals in a higher percentage of the cases compared to bisected animals with a half apical organ (30), 2) Apical organ amputation consisting in the removal of the apical organ, requiring the remaining piece to regenerate dome cilia, balancing cilia, lithocytes, polar fields, and neural cells of the apical organ floor (**Figure 1D**). The timing and order of formation of missing structures was assessed by *in vivo* imaging of the regenerating animals at different time points along the regeneration process.

Wound healing

To assess the mechanism of wound healing, juvenile cydippids were punctured generating a small epithelial gap (**Figure 2A**). (Imaging of larger wound healing events provided to be too difficult to document visually). Within minutes after puncture, the edges of the cells lining the gap increased their thickness indicating the start of the wound closure. The next phase of wound closure was characterized by the migration of a small number of cells coming from deep levels of the mesoglea (underneath the epithelial layer) to the edges of the wound (**Figure 2B, Additional file 1**). Interestingly, while the migration of cells from the mesoglea to the wound site was quite evident, the migration of epithelial cells across the wounded area was not observed. Once the migrating deep cells adhered to the edges of the gap, they started to extend spike-shaped cytoskeleton extensions (filopodia) laterally towards the adjacent cells. The diameter of the gap was progressively reduced as the connections between filopodia of marginal cells pulled the edges of the wound together (**Figure 2C**). When the diameter of the gap was significantly reduced, the cells at the margind of the gap started to extend filopodia not only to adjacent cells, but also to cells from the opposite edge of the wound. At this stage, multiple filopodia were detected emerging from a single cell (**Figure 2D**). Filopodia from all the edges of the wound eventually met forming a network of filaments that sealed the gap (**Figure 2E**) resulting in a scar-free epithelium within approximately 1 -1.5 hours after the procedure.

Events during half-body regeneration of Mnemiopsis leidyi cydippids following bisection

through the oral-aboral axis

Cydippid stage animals were bisected through the oral-aboral axis with one half

retaining the intact whole apical organ. Bisected cydippids containing half of the set of

structures present in intact cydippids (four comb rows and one tentacle) and a

complete apical organ were left to regenerate in 1x filtered sea water (1x FSW) (n>100)

at 22ºC. Wound closure was initiated rapidly after bisection with the edges of the

wound forming a round circumference that continued to reduce in diameter until

meeting and was completed within 2 hours after bisection (hab). No scar or trace of the

original wound was evident after this time. About 16 hab, four ciliated furrows – which

connect the apical organ with the comb rows – appeared on a surface epithelium at the

aboral end of the cut site (**Figure 3B**). A large mass of cells (blastema-like structure)

never appeared at the cut site. Rather, accumulations of cells were detected forming

the primordia of all four of the future comb rows in a deeper plane at the end of each

ciliated furrow. By 24 hab, the first comb plates appeared, first in the two most external

(closer to existing comb rows) comb rows and later in the two internal rows (**Figure

3C**). Comb plate formation did not follow a consistent pattern initially. The correct

orientation of comb plates and coordination of their beating was accomplished after a

number of comb plates were formed (**Figure 3D** surface, deep), as has been described

previously (33). Within 40 hab, coordinated comb plates were beating in all four

regenerating comb rows and the primordia of tentacle bulb had emerged in the middle

of the four comb rows. By 48 hab, regeneration of the missing structures of the

cydippid body was essentially completed including the formation of the tentacle

growing from the tentacle bulb (**Figure 3E**). At 96 hours after bisection, the

regenerated tentacle was long enough to actively catch prey. The cut side continued to

grow and within a day or two it was indistinguishable from the uncut side (**Figure 3F**).

Events during regeneration of Mnemiopsis leidyi cydippids following apical organ

amputation

Cydippids in which the apical organ was amputated were left to regenerate in

1x FSW (n>100) at 22ºC. The cut edges of the wound met and sealed within 30-60

minutes of the operation and the lesion was completely healed around 2 hours post

amputation (hpa). Between 6 and 12 hpa, cells congregated under the wounded

epithelium forming the primordia of the future apical organ (**Figure 4E-F'**). Extension of

the ciliated furrows from each comb row towards the wound site could be spotted

around 12 hpa. Within 24 hpa, cells at the wound site started to differentiate into the

floor of the apical organ and its supporting cilia (**Figure 4G-H'**). At 48 hpa all the components of the statolith, including the supporting cilia, the balancing cilia and lithocytes, were formed (**Figure 4I-J'**). At approximately 60 hpa the complete set of structures forming the apical organ were regenerated with the exception of the polar fields (**Figure 4K-L'**). Within 3 days after surgery, the polar fields had formed, and animals were indistinguishable from control animals of the same size.

Cell proliferation in intact cydippids

To identify areas of cell proliferation in juvenile *M. leidyi*, intact cydippids between 1.5 – 3.0 mm in diameter were labelled with the thymidine analog 5-ethynyl-20-deoxyuridine (EdU), which is incorporated into genomic DNA during the S-phase of the cell cycle (27,34-36) (**Figure 5A**). Cydippids incubated with EdU during a 15-minute pulse showed a pattern of cell division characterized by two main regions of active cell proliferation corresponding to the two tentacle bulbs (**Figure 5B-B'**). Higher magnifications of these structures showed EdU staining specifically concentrated at the lateral and median ridges of the tentacle bulb. Two symmetrical populations of densely packed cells were observed at the aboral extremity of the lateral ridges, previously characterized by Alié et al. (2011) as the aboral/external cell masses (a.e.c.) (**Figure 5C'**). EdU labeling was also found in some cells of the apical organ and few isolated cells along the pharynx and under the comb rows (n=20, **Figure 5B-D'**). To detect dividing cells in M phase of the cell cycle we performed anti-phospho-histone 3 (anti-PH3) immunolabelings in intact cydippids. The spatial pattern and distribution of PH3 labeling closely matched the one described for EdU incorporation, although PH3+ cells were always about 10% less numerous than the EdU labeled cells, suggesting that the duration of the M phase is much shorter than the S phase (n=10, **Figure 5B'',5C'',5D''**).

In order to track the populations of proliferating cells over time in intact animals we performed EdU pulse-chase experiments consisting in a 15-minute EdU incubation (pulse) and a chase of different times followed by visualization (**Figure 5A**). After a 24h chase, the pools of proliferating cells had migrated from the tentacle bulb through the proximal region of the tentacles, although some EdU+ cells were still detected at the tentacle sheath. Increased labeling of nuclei in the apical organ, pharynx and comb rows was also observed (n=10, **Figure 5E-F'**). Following a 48h chase, the population of proliferating cells that was originally in the tentacle bulbs at the time of labeling had migrated to the most distal end of the tentacles, but only a few cells associated with the tentacle bulb showed long-term EdU retention, suggesting that there is a resident population of slowly dividing stem cells in the tentacle bulb as previously reported by

Alié et al. (2011). The number of EdU+ nuclei along the pharynx, the apical sensory
organ (specifically in the apical organ floor) and comb rows was considerably
increased compared to the 24h chase condition (n=10, **Figure 5G-G'**), suggesting that
there are either small populations of EdU labeled cells restricted to those areas that
had proliferated during the chase period, or that cells migrated in to those regions from
regions of high mitotic density, or a combination of both events. Attempts to quantify
and compare brightness levels of EdU positive cells to infer additional rounds of cell
division during the chase period proved inconclusive (data not shown).

Cell proliferation is activated during ctenophore regeneration

In order to determine the role of cell proliferation in ctenophore regeneration we
performed a series of EdU experiments in regenerating cydippids. A 15-minute
exposure to EdU at different times after surgical cutting was evaluated after two types
of surgeries that required different regenerative responses: a bisection through the
oral-aboral axis and an apical organ ablation (**Figures 6A** and **7A**). The dynamics of
cell proliferation at the wound site were quantified by calculating the ratio of EdU+
nuclei to total nuclei (DAPI stained) at different time-points following surgery (**Figures
6B** and **7B**).

Following oral-aboral bisection, EdU+ nuclei were first detected at the wound
site between 6 and 12 hours after bisection (hab). There was some variability in the
presence of EdU+ nuclei at 6 hab – with some specimens having fewer EdU+ nuclei at
the wound site than others – however the presence of EdU+ cells was consistent in all
the analyzed individuals by 12 hab. The few EdU+ cells at the early stages were
scattered all along the cut site, but no aggregation of cells was observed (n=7, **Figure
6C-C''**). The number of EdU+ nuclei at the wound site slightly increased between 12
and 24 hab reaching a maximum at 24hab (**Figure 6B**), when EdU+ cells appeared
concentrated in discrete areas corresponding to the forming primordia of the
regenerating tissues (the tentacle bulb and comb rows) (n=27, **Figure 6D-D''**). By 48
hab, the % of EdU+ nuclei had decreased as the cells started to differentiate into the
final structures. EdU+ nuclei appeared confined into the regenerating comb rows and
tentacle bulb, already distinguishable by nuclear staining (n=12, **Figure 6E-E''**). At 72
hab, the number of EdU+ nuclei at the comb rows was considerably reduced and these
were concentrated at the oral end of the regenerating structures, where oral portions of
structures are generated later than aboral regions. For example, proliferative cells were
no longer detected at the aboral end of the comb rows where cells had already
differentiated into comb plates. In contrast, EdU+ cells at the regenerating tentacle bulb
were abundant but appeared organized at the aboral extremity forming the two

symmetrical populations of cells characteristic of the structure of the tentacle bulb
(n=15, **Figure 6F-F'**). By 96 hab, when major repatterning events of regeneration were
completed, EdU+ cells were only detected at the regenerated tentacle sheath forming
the pattern of cell proliferation previously described in the tentacle bulbs of intact
cydippids (**Figure 5**) (n=5, **Additional file 3A-A''**). In combination with EdU
incorporation experiments, anti-PH3 immunostaining was performed at selected time-
points following bisection. PH3+ cells were detected in the regenerating comb rows
and tentacle bulb at 24 hab and 48 hab (**Additional file 4A-B''**) consistent with the
EdU incorporation pattern, although the number of PH3+ cells was always less
numerous than the EdU+ cells.

EdU labeling was also detected at the wound site of regenerating cydippids
after apical organ amputation. Consistent with the oral-aboral bisection surgeries,
EdU+ cells were first detected at 12 hpa suggesting that the start of the cell
proliferation response occurred between the 6 and 12 hpa time points. A peak of cell
proliferation was also observed at 24 hpa (**Figure 7B**), with EdU+ cells localized at the
primordia of the apical organ, specifically in the apical organ floor and in the
surrounding tissue including the regenerating comb rows adjacent to the cut site (n=15,
Figure 7E-F''). The number of proliferating cells slightly decreased at 48 hpa when
EdU+ cells were concentrated in the regenerating apical organ and were no longer
found in the tissues near the wound site (n=20, **Figure 7G-H''**). By 72 hpa, the EdU+
nuclei were scarce and localized mostly along the polar fields in some specimens,
while EdU+ nuclei were completely absent in other individuals at the same time-point
(n=6, **Additional file 3B-C''**). Anti-PH3 immunostaining showed presence of M-phase
cells at the regenerating area at both 24 hpa and 48 hpa. Similar to half body
regeneration, the pattern of anti-PH3 was consistent with the EdU labeling with PH3+
cells being more numerous at 24 hpa than 48 hpa. However, it should also be noted
that as in all of our other observations, the number of PH3+ cells at any given time-
point was always lower compared to the EdU+ cells (**Additional file 4C-D'**).

Interestingly, for both types of surgeries, proliferating cells were not organized
in a compacted mass of "blastema-like" cells from were new tissue formed. In contrast,
proliferating cells were very few and scattered throughout the wound site at early time-
points after surgery – when a blastema is normally formed in animals with cell-
proliferation based regeneration – and appeared more abundant and directly confined
at the correct location of missing structures at later stages of regeneration, where they
differentiated in place.

Cells participating in the regenerative response appear to arise locally

To investigate the source of cells that contribute to the formation of new tissue during ctenophore regeneration we performed a series of EdU pulse and chase experiments in regenerating cydippids. This technique has been successfully used in different model systems as a strategy to indirectly track populations of proliferating cells and determine their contribution to the formation of new structures (19,37). With the aim of determining whether cells proliferating before amputation contribute to the formation of new tissues, uncut cydippids were incubated in EdU, which was incorporated into cells undergoing the S-phase of cell cycle. After a 15-minute pulse, animals were removed from EdU, and EdU incorporation was blocked ("chased") with several washes of thymidine and 1x FSW. Following the washes, apical organ amputations and oral-aboral bisections were performed and animals were left to regenerate in 1x FSW. The location of EdU+ cells was subsequently visualized at 24 and 48 hours after injury. In combination with EdU detection, an immunostaining against PH3 was performed in order to detect cells that were actively dividing in the animal immediately before fixation (**Figure 8A**).

No EdU+ cells were detected at the wound site at 24h (n=30) nor 48h (n=10) after bisection (**Figure 8B-C''**). EdU labeling at the tentacle bulb resembling the pattern of cells migrating from the tentacle bulb distally along the tentacle previously described (**Figure 5F-F'**) confirmed that the chase worked properly (**Figure 8B**). Moreover, presence of PH3+ cells were observed at the regenerating area indicating active cell division at the moment of fixation (**Figure 8B'' and 8C''**). Following apical organ amputation, few EdU+ nuclei were detected at the area of apical organ regeneration although the EdU signal was very weak, suggesting that these cells were the result of multiple rounds of division (n=13, **Figure 8D-D''**). After a 48h chase, few bright EdU+ nuclei were detected at the apical organ suggesting that S-phase cells from the uncut tissue might contribute to the formation of the apical sensory organ at later stages of regeneration (n=12, **Figure 8E-E''**). Presence of PH3+ cells at the regenerating apical organ confirmed active cell division at the apical organ area (**Figure 8E'-E''**). Taken together, these results show a minor contribution of proliferative cells originating in distant pre-existing proliferative tissue such as the tentacle bulbs to the formation of new structures.

Expression patterns determined through *in-situ* hybridization have reveled spatially restricted expression of the stem cell gene markers *Piwi, Vasa, Nanos and Sox* within areas of cell proliferation including the tentacle bulbs, in both juvenile cydippid and adult stages (27,28,36). On the other hand, the ctenophore group of Beroids do not possess tentacles at any stage of their life cycle and they are the only group of ctenophores that have lost the ability to regenerate (24). Based on these

observations, it was hypothesized that the tentacle bulbs served as putative "stem cell
niche" source of new cells involved in the regeneration process (23). To test this
hypothesis, we surgically removed both tentacle bulbs from juvenile cydippids and
assessed their ability to regenerate. Two days after amputation all animals had
regenerated all the cell types of the tentacle bulb (n>100, **Figure 9A-C'**). EdU labeling
at different time-points after amputation showed activation of cell proliferation during
tentacle bulb regeneration, consistent with the other two types of surgeries analyzed.
EdU+ nuclei were first detected at the distal end of the endodermal canals that connect
the tentacles to the gastrovascular system at 18 hpa (n=10, **Figure 9E-E''**). At 24 hpa
the number of EdU+ cells had increased, and they were mainly organized forming the
primordia of tentacle bulbs although some EdU+ cells were still detected at the tip of
the endodermal canal connecting to the tentacle bulbs in formation (n=20, **Figure 9F-**
F''). By 48 hpa, EdU+ nuclei appeared organized in the characteristic pattern of intact
tentacle bulbs (**Figure 5B'** and **5C'**), and they were not detected at the endodermal
canals any more (n=20, **Figure 9G-G''**). In addition, animals in which both tentacle
bulbs and apical organ were removed, were able to regenerate all the missing
structures (data not shown). These data argue strongly that the tentacle bulbs are not
the source of multipotent stem cells required for the successful regenerative response
in tentaculate ctenophores and point to a local source of cells in the formation of new
structures.
412

There is recruitment of slowly-dividing cells at the regenerating structures
After having discarded the tentacle bulbs as the "stem cell niche" source of new cells
for the regeneration of all missing structures, we questioned whether there is a
population of slowly-dividing cells scattered among the ctenophore body which get
stimulated after injury and contribute to the regenerative process. In order to target for
slowly-cycling cells, we performed a long pulse-chase experiment consisting of a 2-
hour EdU incubation pulse (in contrast to the 15-minute EdU pulse performed in
previous pulse-chase experiments) followed by a 5-day chase (**Figure 10A**).
EdU+ cells just after the 2-hour EdU pulse were found very densely compacted
in two main areas corresponding to the tentacle bulbs (**Figures 10D-D'** and **10G-G'**),
and sparser although also abundant in the apical sensory organ (specifically the apical
organ floor) (**Figures 10C-C'**, **10G-G'**and **Additional file 8**), under the comb rows
(**Figure 10I-I'**) and the pharynx (**Figure 10F'**) (n=22, **Figure 10B-I**). This EdU labeling
is consistent with the EdU pattern observed in intact cydippids after 15-minute EdU
pulse (**Figure 5B-B'**), however, in contrast to the 15-minute pulse EdU pattern, the
amount of EdU+ cells detected at the apical organ, comb rows and pharynx was

considerably higher after the 2-hour EdU incubation. Moreover, EdU+ cells were also
detected through the epidermal surface (**Additional file 6 and 7**) and endodermal
canals (**Additional file 9**), locations where EdU+ cells were not detected after the 15-
minute EdU pulse. After the 5-day chase, EdU+ cells were located around the apical
organ, pharynx, comb rows and epidermis but they were no longer detected at the
tentacle bulbs nor the tentacles (n=21, **Figure 10J-O'** and **Additional file 10** and **11**).
This result is consistent with the idea that a population of protected slowly-dividing cells
does not exist confined in a concrete location (tentacle bulbs) but, rather, slowly-cycling
cells are found scattered among several structures of the cydippid body.
To determine whether this population of slowly-dividing cells contribute to the
process of regeneration, we amputated the apical organ structure from cydippids
exposed to the 2-hour EdU pulse and 5-day chase. The location of EdU+ cells was
subsequently visualized at 24 hpa (**Figure 10A**). EdU+ cells were detected at the
regenerating apical organ 24 hours post amputation (n=10, **Figure 10P-S'**) indicating a
contribution of slowly-dividing cells originated at the pre-existing tissue to the
regenerating structure.
445

Cell proliferation is strictly required for ctenophore regeneration
Having demonstrated that cell proliferation is activated during ctenophore
regeneration, our next aim was to address the requirement of cell proliferation in the
process of regeneration. Juvenile cydippids were exposed to hydroxyurea (HU)
treatments, a drug that inhibits cell proliferation by blocking the ribonucleotide
reductase enzyme and thereby preventing the S-phase of cell cycle (38). We first
performed a dose-response test experiment in order to set the working concentration of
HU in which animals could be continuously incubated during the complete period of
regeneration with no significant disruption of their fitness. Concentrations of 20, 10 and
5mM HU were tested over a 72-hour time course. Incubations in 20 and 10mM HU
were toxic and caused the degeneration and eventually death of most of the animals
during the first 24 hours of incubation (data not shown). Incubations in 5mM HU were
much less harmful; cydippids maintained a good condition, swimming normally with no
cell death over the 72-hour time course. We therefore decided to set 5mM HU as the
working concentration for the cell proliferation inhibitor experiments. We then assessed
the efficacy of that drug concentration in blocking cell proliferation in intact cydippids.
Intact cydippids were incubated in 5mM HU for 24 and 72 hours and then incubated for
15 minutes with EdU as previously described (**Additional file 12A**). At 24 hours of HU
incubation, there was no detectable incorporation of EdU as compared with control
cydippids, which showed the characteristic pattern of cell proliferation described in

Figure 5 (**Additional file 12B-C'**). Inhibition of cell proliferation was maintained 72 hours after continuous HU incubation, as shown by the total absence of EdU+ cells in treated cydippids (**Additional file 12D-E'**). Finally, we evaluated the effect of the drug during regeneration in dissected cydippids. Cydippids bisected through the aboral-oral axis and cydippids in which apical organ was amputated were exposed to a continuous incubation of 5mM HU from 0 to 72 hours after surgery. None of the bisected cydippids had regenerated at 72 hours following HU treatment (n=75, **Figure 11D-E'**). Wound closure and healing occurred normally as shown by the continuous epidermal layer covering the wound (**Figure 11E**), but no sign of formation of the missing structures (tentacle bulb and comb rows) was observed. Moreover, bisected cydippids in which HU was added 4 hab (n=25) – when wound healing is already completed – and 12 hab (n=25) – when cells at the wound site have already begun to cycle – fail to regenerate the missing structures (data not shown). Likewise, none of the apical organ amputated cydippids had regenerated any of the structures/cell types of the missing apical organ at 72 hours following HU treatment, although the wound had correctly healed (n=55, **Figure 11H-J'**). Although HU treated animals failed to regenerate any of their missing structures, an aggregation of cells could be observed at the wound site (**Figures 11E** and **11J'**). These accumulations of quite large round-shaped cells could correspond to undifferentiated cells ready to re-form the missing structures but not able to proceed due to the blocking of cell proliferation. Importantly, the absence of EdU incorporation in dissected cydippids treated with HU confirmed that cell proliferation was completely suppressed (**Additional file 12I-J'**). From these observations we conclude that regeneration was impaired due to the absence of cell proliferation, therefore, cell proliferation is indispensable for normal ctenophore regeneration.

Regenerative ability is recovered after HU treatment ends

Hydroxyurea has been shown to be reversible in cell culture following removal of the inhibitor (33) (**Figure 12A**). HU treatments on cut cydippids showed that wound healing occurs normally without cell division. In order to determine whether regeneration could be initiated in HU treated animals we took surgically cut cydippids that had been exposed to HU over 48 hours, washed them in 1x FSW to remove the inhibitor, and then followed their development for 48 hours to check for any ability to regenerate missing cell types (**Figures 12B** and **12E**). Surprisingly, 36 out of 94 bisected cydippids (38%) had regenerated all the missing structures (comb rows, tentacle bulb and tentacle) 48 hours after HU had been removed (**Figure 12D-D''**). 58 out of 94 bisected cydippids (62%) showed some signs of regeneration but ultimately remained as "half animals", suggesting that these animals were not healthy enough to

complete the regeneration process (40). (Note that these animals were not fed during

the treatment (2 days) or recovery period (2 additional days)). On the other hand,

100% of the cydippids in which the apical organ was surgically removed and had been

treated with HU for 48 hours, regenerated all the normal cell types of the apical organ

(n=51, **Figure 12H-I'**). Altogether, these results show that ctenophore regeneration can

be initiated over 48 hours after wound healing is complete, hence, wound healing and

regeneration appear to be two relatively independent events which can be temporally

decoupled.

DISCUSSION

In this study, we provide a detailed morphological and cellular characterization

of wound healing and regeneration in the ctenophore *Mnemiopsis leidyi*. Wound

closure is initiated immediately after injury, with the edges of the wound forming a

round circumference that moves over the underlying mesoglea as it continues to

reduce in diameter until they meet and forming a scar-less wound epithelium by 2

hours following injury. Two main mechanisms seem to be pivotal for ctenophore wound

closure: active cell migration of cells from the mesoglea underneath the epithelium

upwards to the edges of the wound; and dynamic extension of filopodia by the leading-

edge epithelial cells in order to zipper the wound edges together. Cell migration and

formation of actin-based cellular protrusions have been described during wound

closure in multiple systems (41), however, slight differences in those mechanisms have

been observed in ctenophore wound healing. First, cell migration takes place in a

"deep to surface" direction instead of a lateral direction, suggesting that only specific

cell-types from the mesoglea, such as mesenchymal cells, have the ability to migrate

and contribute to gap closure. Second, wound-edge cells in ctenophores organize their

cytoskeleton in spike-shaped filopodia rather than in plate-like extensions

(lamellipodia), which happen to be the most common type of cellular protrusions

among different model systems of wound healing, including the cnidarian Clytia (42).

Despite these minor differences, the fact that common mechanisms of wound closure

are shared between early branching phyla like ctenophores and cnidarians and

bilaterians (including vertebrates) proves the ancient origin of wound healing

mechanisms as a strategy to maintain epithelium integrity. Wound healing in *M. leidyi*

takes place through changes in cell behavior and occurs normally in the absence of

cell proliferation. This observation is consistent with the majority of animal models of

regeneration found in cnidarians (12,20,42-44) as well as with the more

phylogenetically distantly-related marine annelid worm *Platynereis dumerilii* (37).

Following wound healing and prior to activation of cell proliferation in *M. leidyi*, there is

remodeling of the tissue surrounding the wound and small numbers of round-shaped
cells sparsely congregate at the wound site suggesting a reorganization of the tissue in
order to prepare it for regeneration. Ctenophore regeneration, however, is strictly
dependent on cell proliferation since none of the missing structures can be reformed in
the absence of cell proliferation as proved by cell proliferation blocking treatments.
Indeed, a combination of both tissue remodeling and cell proliferation-based strategies
has been previously described in the regeneration of other animals including annelids
(17,45), although in those cases tissue remodeling takes place simultaneously with cell
proliferation – or even subsequent to activation of cell proliferation – and is involved in
the regeneration of a specific structures such as parapodia (46) or the gut (47).

Cell proliferation in *M. leidyi* is first detected at the wound site between 6-12
hours after surgery. The percentage of proliferating cells increases progressively
during the first 12 hours following injury and reaches a maximum around 24 hours
when the primordia of the missing structures are clearly delineated. Following this peak
of cell proliferation, the percentage of cells undergoing cell division (S-phase)
decreases while cells start to differentiate into their final structures. Comparing the
kinetics of cell proliferation during regeneration of *M. leidyi* with the anthozoan
cnidarian *Nematostella vectensis* (12), the percentage of dividing cells at the wound
site is lower and the peak of maximum cell proliferation occurs earlier in ctenophore
regeneration. In intact cydippids, cell proliferation is concentrated in two main areas of
the cydippid's body corresponding to the tentacle bulbs. Some actively cycling cells are
also found in the apical organ as well as few isolated dividing cells along the pharynx
and under the comb rows. These results are consistent with previous EdU analysis
performed in *M. leidyi* cydippids (27,28) and adult ctenophores of the species
Pleurobrachia pileus (36) where EdU labeling has been detected in the same spatially
restricted populations identified as stem cell pools, specialized in the production of
particular cell types. Pulse-chase experiments show concrete areas of active cell
proliferation in the tentacle bulb and progressive migration of these proliferating cells
from the tentacle bulb to the distal tips of the tentacle. These observations fit with
histological and cellular descriptions of the tentacle apparatus (36,48) which identified
different populations of undifferentiated progenitors source of all cell types found in the
tentacle tissue. Surprisingly, long-term EdU retention is not detected in any of the cells
of the tentacle apparatus suggesting that a population of "protected" slowly-dividing
stem cells might not exist in the tentacle bulb. This opens the question of the nature
and origin of the progenitor cells responsible for the maintenance of the homeostasis of
the tentacle structure. Perhaps tentacle bulb stem cells are continuously recruited from

adjacent somatic cells rather than being derived from a uniquely committed set of slow
dividing "set aside" stem cells.

Interestingly, proliferating cells during regeneration do not organize to form a
single large blastema-like structure from which a field of cells are reorganized to form
the missing structures. Rather, small numbers of undifferentiated cells assume the
correct location of all missing structures simultaneously and differentiate in place.
Considering the early branching phylogenetic position of ctenophores in the tree of life
(49,50), the absence of blastema during ctenophore regeneration questions whether
the formation of a blastema – which so far appears to have been reported in
representatives of all phyla of regenerating animals (51) – is a conserved trait
throughout the evolution of animal regeneration.

The strict requirement of cell proliferation and the absence of blastema
formation could make ctenophore regeneration a case of non-blastemal cell
proliferation-dependent regeneration. Although far less common than the blastemal
based regeneration, isolated cases of non-blastemal regeneration have been reported
such as lens regeneration by transdifferentiation in newts (52) or liver regeneration by
compensatory proliferation in humans (53). EdU pulse-chase experiments after
amputation show little to no contribution of cells originating in the main regions of active
cell proliferation, including the tentacle bulbs, to the formation of missing structures.
Moreover, the removal of these structures (tentacle bulbs), which have been reported
to be localized areas of expression of genes involved in stem cell maintenance and
regulation of cell fate (27,28,36) – and thus proposed to act as stem cell niches for
regeneration – do not prevent regeneration. These observations argue against the
contribution of discrete stem cell pools that migrate to and give rise to the re-formation
of lost structures, suggesting that new structures are generated from a local source of
cells that become activated to give rise to missing structures/cell types. Longer pulse-
chase experiments in which the animals where incubated in EdU for an extended
period of time (2 hours vs 15 min) and then followed by a much longer chase allowed
the identification of a population of slowly-cycling (potentially stem) cells which could
have escaped the initial short 15-minute pulse. Three main observations can be done
from this experiment: 1) While the pattern of EdU+ proliferating cells after the long 2-
hour pulse coincides with the one observed after the short 15-minute pulse, EdU+ cells
after the chase are no longer found in any of the cells forming the tentacle apparatus.
This observation is consistent with the results obtained in the pulse-chase experiments
after amputation and thus strongly discards the existence of a stem cell niche in the
tentacle bulbs source of cells for the regenerative process; 2) Long retaining EdU+
cells (referred as slow-dividing cells), are found uniformly distributed around the

cydippid's body instead of being organized in discrete pools. This distribution pattern of
cell proliferation could be comparable to the neoblast distribution pattern characteristic
of planarians (6,54,55), with the difference that neoblasts are known to reside in the
planarian parenchyma – a mesenchymal tissue surrounding organ systems – (6), while
these potential stem cells in ctenophores are found in both ectodermal and endodermal
structures but not in the mesenchymal cells of the mesoglea. 3) EdU+ long retaining
cells are detected at regenerating structures after amputation indicating that these
slow-cycling cells contribute to the re-formation of new structures.

It is however important to note that our experiments do not give a definitive
answer to the question of the origin of cells that give rise to new structures. There is
the possibility that wound healing activates the dedifferentiation of cells at the wound
site that are reprogrammed to give rise to whatever the appropriate set of cell types are
needed to reconstitute the missing structures. The accumulation of large, round,
apparently undifferentiated cells at the wound site during HU treatment is at least
consistent with this scenario. On the other hand, wound healing could activate a
dormant population of slowly-dividing pluripotent stem cells located uniformly around
the body that could migrate to the wound site and drive the regeneration process,
which could have escaped the short pulse of EdU incorporation and re-entered the cell
cycle as a consequence of injury. Although we saw some evidence of cells migrating
from the underlying mesoglea to repair the wound site, we did not see any evidence of
long range cell migration to the site of new cell type formation. Nonetheless, an early
study leveraging the combination of cell-lineage and specific cell-deletion experiments
in *M. leidyi* showed that comb plate regeneration cannot occur when the entire
complement of cell lineage comb plate progenitors are killed during embryogenesis,
suggesting that, at least for comb plate regeneration, a semi-committed somatic stem
cell population is set-aside during embryogenesis for comb plate replacement (32,56).
These data are premature and need to be extended to other cell types and later stages
of the regenerative process, however the stereotyped cell lineage seen in ctenophores
provides exciting opportunities to pursue the origins of stem cells in the regenerative
process in living animals.

Overall, our data, together with evidences from previous studies in ctenophores,
is consistent with a model of ctenophore regeneration based on a combined
contribution of active cell proliferation at the wound site together with the recruitment of
slowly-dividing cells to the regenerating structure (**Figure 13**). The fact that no discrete
pools of neoblast-type stem cells are identified in the ctenophore body, and the
absolute requirement for cell proliferation supports the hypothesis that proliferating
cells at the wound site are the result of dedifferentiation events. On the other hand,

whether these potential slowly dividing stem cells migrate to the wound site from close
(local) or distant locations remains an open question. Gene expression data during the
process of *M. leidyi* regeneration combined with cell tracing experiments will contribute
to address these questions and thus refine our working model of the origin of cells
during ctenophore regeneration. Molecular data during regeneration will also be very
valuable for performing comparisons of gene expression profiles between *M. leidyi*
development (57) and regeneration and thus determine whether the molecular basis of
ctenophore regeneration is similar to that deployed during development.

It is quite accepted that cells that re-epithelialize the wound (i.e the "wound
epithelium") provide the signals necessary to initiate regeneration (58,59). In
vertebrates, local thrombin activation is a signal for regeneration as shown by the study
in which cultured newt myotubes returned to the cell cycle by the activity of a thrombin-
generated ligand (60). On the other hand, cellular interactions also seem to be
important for the initiation of the regenerative response. One such case is the
dorsoventral interaction between the wounded tissues during wound healing in
planarians which has been shown to play a key role in the formation of the blastema
and, hence, initiation of regeneration (61). These observations suggest that wound
healing and regeneration are two closely related processes which need to take place
sequentially in time. Our results, however, show that ctenophore regeneration can be
initiated over 48 hours after wound healing is completed, suggesting that regeneration
can be initiated without direct signaling induced by the wounded epithelium.
Regeneration of the missing structures is not initiated until the cell proliferation blocking
treatment is removed. Hence, another case scenario is that the wound epithelium
produces persistent signaling necessary for triggering regeneration at the time of
wound healing, but the process cannot be initiated due to the blocking of cell
proliferation. This is consistent with the proposed hypothesis for *Nematostella* that the
key transition from wound healing to a state of regeneration is the activation of cell
proliferation (62). Studying and comparing the molecular signaling involved in both
ctenophore wound healing and regeneration will be very useful to get further insight
into the relationship between these two processes.
680

CONCLUSIONS

In conclusion, this study provides a thorough description of the morphological
and cellular events during ctenophore wound healing and regeneration and compares
them with the regenerative strategies followed by other metazoans. The early
branching phylogenetic position of ctenophores together with their rapid, highly

stereotyped development and remarkable ability to regenerate make them a key
system to gain a better understanding of the evolution of animal regeneration.
688
METHODS
Animal care
Regeneration experiments were performed on juvenile *Mnemiopsis leidyi*
cydippid stages due to their small size and ease of visualization and because their
power of regeneration is the same as adults (30). *M. leidyi* cydippids were obtained
from spawning adults collected from either the floating docks located around Flagler
Beach area, FL. USA, or from the floating docks at the east end of the Bridge of Lions
on Anastasia Island, St. Augustine, FL. USA. For spawning, freshly collected adults
were kept in constant light for at least two consecutive nights and then individual
animals transferred into 6" diameter glass culture dishes filled with 1x FSW and placed
in total darkness. After approximately 3-4 hours in the dark at 22-24ºC, these self-fertile
hermaphroditic animals had spawned and embryos were collected by pipetting them
into a new dish of UV treated 1.0 micron filtered full strength seawater (1x FSW) using
a transfer pipette. Embryos were raised at 22-24ºC for approximately 5-7 days and fed
once a day with rotifers (*Brachionus plicatilis*, 160µm) (Reed Mariculture, Campbell,
CA. USA).
705
Animal surgeries
Operations were done in 35 mm plastic petri dishes with 2 mm thick silicon-
coated bottoms (SYLGARD–184, Dow Corning, Inc.) on cydippids 1.5-3.0 mm in
diameter. Cydippids were transferred in to the operation dishes in 0.2 µm-filtered
seawater and cut using hand pulled glass needles from Pyrex capillaries (30). Three
types of operations were performed: 1) Oral-aboral bisections, in which animals were
cut longitudinally through the esophageal plane generating two "half-animals". The
operations were performed such that one half retained an intact apical organ while the
remaining half lacked the apical organ. Only the halves retaining the apical organ were
studied here as these halves regenerate to normal animals in a high percentage of the
cases (30). 2) Apical organ amputations, involving the removal of the apical organ by
cutting perpendicular to the oral-aboral axis above the level of the tentacle bulbs. 3)
Tentacle bulb amputations, consisting in the removal of both tentacle bulbs (**Figure
1D**). Following surgery, halves containing the apical organ, amputated cydippids
without apical organ and amputated cydippids without tentacle bulbs were returned to
35 mm plastic Petri dishes filled with 0.2 µm filtered 1x FSW for the desired length of
time without feeding. All the regenerating experiments were performed at 22-24ºC.

To study the wound healing process, juvenile cydippids were punctured
generating a round-shaped wound of approximately 200-400 µm of diameter. Animals
were placed in a small drop of water on a siliconized (Rain-X, Inc.) treated microscope
slide and punctures were performed by pinching the epithelium layer using a pair of
sharp needles (World Precision Instruments, Sarasota, FL. USA, Cat#500341). After
puncture, animals were checked for the presence of an epithelial gap with the edges of
the wound forming a small circumference exposing the mesoglea, and then they were
immediately mounted for live imaging (see below).

Tissue labeling and cell counts

Detection of cell proliferation by incorporation of EdU

To label proliferating cells, cydippids were fixed and processed for fluorescent
detection of incorporated EdU using the Click-iT EdU labeling kit (Invitrogen Thermo
Fisher Scientific, Waltham, MA. USA, Cat #C10424), which incorporates EdU in cells
that are undergoing the S phase of the cell cycle. Specifically, intact cydippids between
1.5-3.0 mm in diameter or bisected/amputated cydippids were incubated in EdU
labeling solution (100 µM of EdU in 1x FSW) for 15 minutes. For pulse-chase
experiments cydippids were incubated with 100 µM EdU in 1x FSW for 15 minutes,
washed 3 times with 100 µM thymidine in 1x FSW, and maintained in increasing
volumes of 1x FSW until fixation. Control or operated cydippids were embedded in
1.2% low melt agarose (25ºC melting temperature, USB, Inc Cat #32830) in a 35 mm
plastic petri dish (Fisher, Inc. Cat #08757100A) and fixed in ice-cold 100mM HEPES
pH 6.9; 0.05M EGTA; 5mM MgSO4; 200mM NaCl; 1x PBS; 3.7% Formaldehyde; 0.2%
Glutaraldehyde; 0.2% Triton X-100; and 1x FSW (0.2 µm filtered) for 1 hour at room
temperature with gentle rocking (63). Animals were then washed several times in PBS-
0.02% Triton X-100, then one time in PBS-0.2% Triton X-100 for 20 min, and again
several times in PBS-0.02% Triton X-100. The EdU detection reaction was performed
according to manufacturer instructions using the Alexa-567 reaction kit. Following
detection, cydippids were washed three times in PBS-0.02% Triton X-100, and
subsequently all nuclei were counterstained with DAPI (Invitrogen, Carlsbad, CA. USA,
Cat. #D1306) at 1.43 µM in 1x PBS for 2 hours. Cydippids were mounted in TDE
mounting media (97% TDE: 970µl 2,2'-thiodiethanol (Sigma-Aldrich, St. Louis, MO.
USA); 30µl PBS) for visualization. To quantify the percentage of EdU labeled cells at
the wound site, Zeiss 710 confocal z-stack projections of operated cydippids were
generated using Fiji software (Image J) and individual cells were digitally counted using
Imaris, Inc. software (Bitplane, Switzerland). Only the area and z-stacks surrounding
the wound site were used for the analysis. EdU+ cells and nuclei were counted

separately in 5 to 10 specimens for each time-point. The number of EdU-positive nuclei

were divided by the total number of nuclei stained with DAPI generating a ratio

corresponding to the % of EdU+ cells.

Immunofluorescence

Proliferating cells in M phase were detected using an antibody against phospho

Histone 3 (PH3 – phospho S10). Control or operated cydippids were fixed as

mentioned above. Fixed cydippids were washed several times in PBS-0.02% Triton X-

100 (PBT 0.02%), then one time in PBS-0.2% Triton X-100 (PBT 0.2%) for 10 min, and

again several times in PBT 0.02%. They were then blocked in 5% normal goat serum

(NGS; diluted in PBT 0.2%) for 1 hour at room temperature with gentle rocking. After

blocking, specimens were incubated in anti-phospho histone H3 antibody (ARG51679,

Arigo Biolaboratories, Taiwan) diluted 1:150 in 5% NGS overnight at 4ºC. The day

after, specimens were washed at least five times with PBS-0.2% Triton X-100.

Secondary antibody (Alexa Fluor 488 goat anti-rabbit IgG (A-11008, Invitrogen,

Carlsbad, CA. USA) was diluted 1:250 in 5% NGS and incubated over night at 4ºC with

gentle rocking. After incubation, specimens were washed three times with PBT 0.02%

and incubated with DAPI (0.1µg/µl in 1x PBS; Invitrogen, Carlsbad, CA. USA, Cat.

#D1306) for 2 hours to allow nuclear visualization. Samples were then rinsed in 1x

PBS and mounted in TDE mounting media (97%TDE: 970µl 2,2'-thiodiethanol (Sigma-

Aldrich, St. Louis, MO. USA); 30µl PBS) for visualization.

Cell proliferation inhibitor treatment with Hydroxyurea (HU)

Cell proliferation was blocked using the ribonucleotide reductase inhibitor hydroxyurea

(HU) (Sigma-Aldrich, St. Louis, MO. USA). Incubations with hydroxyurea were

performed at a concentration of 5 mM in 1x FSW. Operated cydippids were exposed to

continuous incubations of 5mM HU for 48-72 hours. HU solution was exchanged with

freshly diluted inhibitor every 12 hours. For washing experiments, the effect of HU was

reversed by removal and replacement of the drug with 1x FSW.

Imaging

Visualization of living animals

In order to immobilize live animals for scoring and time lapse observations we utilized a

custom made optically transparent jammed microgel. A solution of

azobisisobutyronitrile (Sigma Aldrich, Inc.) and N'-Methylene Bisacrylamide (Sigma

Aldrich, Inc) in ethanol is prepared at a 99:1 AAM:MBA molar ratio (64,65). The

solution is sparged with nitrogen for 30 minutes, then placed into a preheated 60°C oil

bath. After approximately 30 minutes, the solution becomes hazy and a white
precipitate begins to form. The reaction mixture is heated for an additional 4 hours, the
precipitate collected by vacuum filtration and rinsed with ethanol on the filter. The
microparticles are triturated with 500 mL of ethanol overnight. The solids are again
collected by vacuum filtration and dried on the filter for ~10 minutes. The particles are
dried completely in a 50°C vacuum oven to yield a loose white powder. The purified
microgel powder is dispersed in 0.2 µm-filtered seawater at a concentration of 7.5%
(w/w) and mixed at 3500 rpm in a centrifugal speed mixer (64,66) in five-minute
intervals until no aggregates are apparent. The microgel is then left to swell overnight,
yielding mounting medium made from packed hydrogel microparticles. The 7.5%
microgel was placed around the operated/punctured cydippids mounted in a
hydrophobic-treated slide (Rain-Ex, Inc). For short live imaging, a cover slip with clay
corners was placed over the specimens; for long time-lapse live imaging, the corners of
the cover slip were sealed using Vaseline in order to maintain the humidity of the
preparation.

Microscopes and image analysis
In vivo differential interference contrast (DIC) images were captured using a Zeiss Axio
Imager M2 coupled with an AxioCam (HRc) digital camera. Fluorescent confocal
imaging was performed using a Zeiss LSM 710 confocal microscope (Zeiss, Gottingen,
Germany) using either a 10x/0.3 NA dry objective, a 20x/0.8 NA dry objective or a
40x/1.3 NA water immersion objective. For time-lapse imaging, DIC images were
captured using a Zeiss Axio Imager M2 coupled with a Rolera EM-C2 camera (Surrey,
BC. Canada). Stacks were taken every minute. Generation of Z-stack projections, time-
lapse movies and image processing was performed using Fiji software (67).

ABBREVIATIONS
1x FSW: 1x filtered sea water; hab: hours after bisection; hpa: hours post amputation;
EdU: 5-ethynyl-20-deoxyuridine; PH3: phospho-histone 3; HU: hydroxyurea; DIC:
differential interference contrast.

DECLARATIONS
Acknowledgements
We thank the owners and staff of Marker 8 Hotel and Marina in St. Augustine for
allowing us access to their floating docks for animal collection, and all the members of
our lab for assistance and discussions.

Funding

This work was supported by the National Science Foundation (NSF) grant (IOS-

1755364). The funders had no role in study design, data collection and interpretation,

or the decision to submit the work for publication.

Availability of data and materials

All data generated or analysed during this study are included in this published article

[and its supplementary information files].

Authors' contributions

Julia Ramon Mateu, Conceptualization, Investigation, Writing—original draft, Writing—

review and editing; Tori Ellison and Thomas E. Angelini, design and synthesis of

microgels; Mark Q. Martindale, Conceptualization, Supervision, Writing—original draft,

Writing—review and editing. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

Not applicable.

REFERENCES

1.   Bely AE, Nyberg KG. Evolution of animal regeneration: re-emergence of a field.

Trends Ecol Evol. 2010;25(3):161–70.

http://dx.doi.org/10.1016/j.tree.2009.08.005

2.   Alvarado AS, Tsonis PA. Bridging the regeneration gap: Genetic insights from

diverse animal models. Nat Rev Genet. 2006;7(11):873–84.

http://dx.doi.org/10.1038/nrg1923

3.   Tanaka EM, Reddien PW. The Cellular Basis for Animal Regeneration. Dev Cell.

2011;21(1):172–85. http://dx.doi.org/10.1016/j.devcel.2011.06.016

4.   Bryant S V, Endo T, Gardiner DM. Vertebrate limb regeneration and the origin of

limb stem cells. Int J Dev Biol. 2002;46:887–96.

5.   McCusker C, Bryant S V., Gardiner DM. The axolotl limb blastema: cellular and

molecular mechanisms driving blastema formation and limb regeneration in

tetrapods. Regeneration. 2015;2(2):54–71. http://dx.doi.org/10.1002/reg2.32

6.   Reddien PW, Alvarado AS. Fundamentals of Planarian Regeneration. Annu Rev

Cell Dev Biol. 2004;20(1):725–57.

http://www.annualreviews.org/doi/10.1146/annurev.cellbio.20.010403.095114

7.   Park HD, Ortmeyer AB, Blankenbaker DP. Cell Division during Regeneration in
Hydra. Nature. 1970;228(October 17th):227–31.

8.   Cummings SG, Bode HR. Head regeneration and polarity reversal in Hydra
attenuata can occur in the absence of DNA synthesis. Wilhelm Roux's Arch Dev
Biol. 1984;194(2):79–86.

9.   Dübel S, Schaller HC. Terminal Differentiation of Ectodermal Epithelial Stem
Cells of Hydra Can Occur in G2 without Requiring Mitosis or S Phase. J Cell
Biol. 1990;110(April):939–45.

10.   Holstein TW, Hobmayer E, David CN. Pattern of epithelial cell cycling in hydra.
Dev Biol. 1991;148(2):602–11.

11.   Chera S, Ghila L, Dobretz K, Wenger Y, Bauer C, Buzgariu W, et al. Apoptotic
Cells Provide an Unexpected Source of Wnt3 Signaling to Drive Hydra Head
Regeneration. Dev Cell. 2009;17(2):279–89.
http://dx.doi.org/10.1016/j.devcel.2009.07.014

12.   Passamaneck YJ, Martindale MQ. Cell proliferation is necessary for the
regeneration of oral structures in the anthozoan cnidarian Nematostella
vectensis. BMC Dev Biol. 2012;12. http://dx.doi.org/10.1186/1471-213X-12-34

13.   Baguna J, Salo E, Auladell C. Regeneration and pattern formation in planarians.
III. that neoblasts are Regeneration and pattern formation in planarians. III. that
neoblasts are totipotent stem cells and the cells totipotent stem cells and the
cells. Development. 1989;107(September):77–86.

14.   Newmark PA, Sánchez Alvarado A. Bromodeoxyuridine specifically labels the
regenerative stem cells of planarians. Dev Biol. 2000;220(2):142–53
http://dx.doi.org/53.10.1006/dbio.2000.9645

15.   Wagner DE, Wang IE, Reddien PW. Clonogenic neoblasts are pluripotent adult
stem cells that underlie planarian regeneration. Science (80-).
2011;332(6031):811–16. http://dx.doi.org/10.1126/science.1203983.

16.   Bely AE. Early events in annelid regeneration: a cellular perspective. Integr
Comp Biol. 2014;54(4):688–99. http://dx.doi.org/10.1093/icb/icu109

17.   Özpolat BD, Bely AE. Developmental and molecular biology of annelid
regeneration: a comparative review of recent studies. Curr Opin Genet Dev.
2016;40:144–53. http://dx.doi.org/10.1016/j.gde.2016.07.010

18.   Zattara EE, Turlington KW, Bely AE. Long-term time-lapse live imaging reveals
extensive cell migration during annelid regeneration. BMC Dev Biol.
2016;16(1):1–21. http://dx.doi.org/10.1186/s12861-016-0104-2

19.   de Jong DM, Seaver EC. Investigation into the cellular origins of posterior

regeneration in the annelid *Capitella teleta*. Regeneration. 2017;(October):1–17.
http://doi.wiley.com/10.1002/reg2.94

20. Bradshaw B, Thompson K, Frank U. Distinct mechanisms underlie oral vs aboral
regeneration in the cnidarian hydractinia echinata. Elife. 2015;2015(4):1–19
http://dx.doi.org/10.7554/eLife.05506

21. Kragl M, Knapp D, Nacu E, Khattak S, Maden M, Epperlein HH, et al. Cells keep
a memory of their tissue origin during axolotl limb regeneration. Nature.
2009;460(7251):60–5. http://dx.doi.org/10.1038/nature08152

22. Pang K, Martindale MQ. Comb jellies (Ctenophora): A model for basal metazoan
evolution and development. Cold Spring Harb Protoc. 2008;3(11):1–11
http://dx.doi.org/10.1101/pdb.emo106

23. Henry JQ, Martindale MQ. Regulation and regeneration in the ctenophore
Mnemiopsis leidyi. Dev Biol. 2000;227(2):720–33.
http://dx.doi.org/10.1006/dbio.2000.9903

24. Martindale MQ. The onset of regenerative properties in ctenophores. Vol. 40,
Current Opinion in Genetics and Development. 2016. p. 113-19
http://dx.doi.org/10.1016/j.gde.2016.06.017

25. Jager M, Manuel M. Ctenophores: an evolutionary-developmental perspective.
Curr Opin Genet Dev. 2016;39:85-92
http://dx.doi.org/10.1016/j.gde.2016.05.020

26. Fischer AHL, Pang K, Henry JQ, Martindale MQ. A cleavage clock regulates
features of lineage-specific differentiation in the development of a basal
branching metazoan, the ctenophore Mnemiopsis leidyi. Evodevo. 2014;5(1).
http://dx.doi.org/10.1186/2041-9139-5-4

27. Schnitzler CE, Simmons DK, Pang K, Martindale MQ, Baxevanis AD.
Expression of multiple Sox genes through embryonic development in the
ctenophore Mnemiopsis leidyi is spatially restricted to zones of cell proliferation.
Evodevo. 2014;5(1):1–17. http://dx.doi.org/10.1186/2041-9139-5-15

28. Reitzel AM, Pang K, Martindale MQ. Developmental expression of "germline"-
and "sex determination"-related genes in the ctenophore Mnemiopsis leidyi.
Evodevo. 2016;7(1):1–16. http://dx.doi.org/10.1186/s13227-016-0051-9

29. Freeman G. Studies on regeneration in the creeping ctenophore, Vallicula
multiformis. Vol. 123, Journal of Morphology. 1967. p. 71–83
http://dx.doi.org/10.1002/jmor.1051230107

30. Martindale MQ. The ontogeny and maintenance of adult symmetry properties in
the ctenophore, Mnemiopsis mccradyi. Vol. 118, Developmental Biology. 1986.
p. 556–76. http://dx.doi.org/10.1016/0012-1606(86)90026-6

31. Coonfield BR. Regeneration in Mnemiopsis Leidyl Agassiz. Biol Bull.
1936;71:421–8.

32. Martindale MQ, Henry JQ. Development and Regeneration of Comb Plates in
the Ctenophore Mnemiopsis leidyi. Biol Bull. 1996 Oct 1;191(2):290–2.
https://doi.org/10.1086/BBLv191n2p290

33. Tamm SL. Regeneration of Ciliary Comb Plates in the Ctenophore Mnemiopsis
leidyi. I. Morphology. J Morphol. 2012;(273):109–20.
https://doi.org/10.1002/jmor.11016

34. Salic A, Mitchison TJ. A chemical method for fast and sensitive detection of DNA
synthesis in vivo. PNAS. 2008;105(7). https://doi.org/10.1073/pnas.0712168105

35. Chehrehasa F, Meedeniya ACB, Dwyer P, Abrahamsen G, Mackay-Sim A. EdU,
a new thymidine analogue for labelling proliferating cells in the nervous system.
J Neurosci Methods. 2009;177(1):122–30.
https://doi.org/10.1016/j.jneumeth.2008.10.006

36. Alié A, Leclère L, Jager M, Dayraud C, Chang P, Le Guyader H, et al. Somatic
stem cells express Piwi and Vasa genes in an adult ctenophore: Ancient
association of "germline genes" with stemness. Dev Biol. 2011;350(1):183–97.
http://dx.doi.org/10.1016/j.ydbio.2010.10.019

37. Planques A, Malem J, Parapar J, Vervoort M, Gazave E. Morphological, cellular
and molecular characterization of posterior regeneration in the marine annelid
Platynereis dumerilii. Dev Biol. 2019;445(2):189–210.
https://doi.org/10.1016/j.ydbio.2018.11.004

38. Young CW, Hodaas S, 1964. Hydroxyurea: inhibitory effect on DNA metabolism.
Science. 146:1172–1174.

39. Adams RLP, Lindsay J. Hydroxyurea. Reversal of inhibition and use as a cell-
synchronizing agent. J Biol Chem. 1967;242(6):1314–17.

40. Bading KT, Kaehlert S, Chi X, Jaspers C, Martindale MQ, Javidpour J. Food
availability drives plastic self-repair response in a basal metazoan-case study on
the ctenophore Mnemiopsis leidyi A. Agassiz. Sci Rep. 2017;7(1):1–9.
http://dx.doi.org/10.1038/s41598-017-16346-w

41. Begnaud S, Chen T, Delacour D, Mège RM, Ladoux B. Mechanics of epithelial
tissues during gap closure. Curr Opin Cell Biol. 2016;42:52–62.
http://dx.doi.org/10.1016/j.ceb.2016.04.006

42. Kamran Z, Zellner K, Kyriazes H, Kraus CM, Reynier JB, Malamy JE. In vivo
imaging of epithelial wound healing in the cnidarian Clytia hemisphaerica
demonstrates early evolution of purse string and cell crawling closure
mechanisms. BMC Dev Biol. 2017;17(1):1–14.

http://dx.doi.org/10.1186/s12861-017-0160-2

43. Singer II. Tentacular and oral-disc regeneration in the sea anemone, Aiptasia
diaphana. J Embryol Exp Morphol. 1971;26(2):253–70.
http://dx.doi.org/10.1007/BF00597293

44. Amiel AR, Johnston HT, Nedoncelle K, Warner JF, Ferreira S, Röttinger E.
Characterization of morphological and cellular events underlying oral
regeneration in the sea anemone, Nematostella vectensis. Int J Mol Sci.
2015;16(12):28449–71. http://dx.doi.org/10.3390/ijms161226100

45. De Jong DM, Seaver EC. A Stable Thoracic Hox Code and Epimorphosis
Characterize Posterior Regeneration in Capitella teleta. PLoS One. 2016;11(2).
http://dx.doi.org/10.1371/journal.pone.0149724

46. Berrill NJ, 1931. Regeneration in Sabella pavonina (Sav.) and other sabellid
worms. J Exp Zool. 58:495-523.

47. Zattara EE, Bely AE. Evolution of a novel developmental trajectory: Fission is
distinct from regeneration in the annelid Pristina leidyi. Evol Dev. 2011;13(1):80–
95. http://dx.doi.org/10.1111/j.1525-142X.2010.00458.x

48. Borisenko I, Ereskovsky A V. Tentacular apparatus ultrastructure in the larva of
Bolinopsis infundibulum (Lobata: Ctenophora). Acta Zool. 2013;94(2):193-202
http://dx.doi.org/10.1111/j.1463-6395.2011.00542.x

49. Dunn CW, Hejnol A, Matus DQ, Pang K, Browne WE, Smith SA, et al. Broad
phylogenomic sampling improves resolution of the animal tree of life. Nature.
2008;452(7188):745–49. http://dx.doi.org/10.1038/nature06614

50. Ryan JF, Pang K, Schnitzler CE, Nguyen AD, Moreland RT, Simmons DK, et al.
The genome of the ctenophore Mnemiopsis leidyi and its implications for cell
type evolution. Science. 2008;452(7188):745-49.
http://dx.doi.org/10.1038/nature06614

51. Sánchez Alvarado A. Regeneration in the metazoans: why does it happen?
BioEssays. 2000;22(6):578–90.

52. Tsonis PA, Del Rio-Tsonis K. Lens and retina regeneration: Transdifferentiation,
stem cells and clinical applications. Exp Eye Res. 2004;78(2):161–72.
http://dx.doi.org/10.1016/j.exer.2003.10.022

53. Michalopoulos GK, DeFrances M. Liver regeneration. Science.
1997;276(5309):60-66 http://dx.doi.org/10.1007/b99968

54. Forsthoefel DJ, Newmark PA. Emerging patterns in planarian regeneration. Curr
Opin Genet Dev. 2009;19(4):412–20. 10.1016/j.gde.2009.05.003.Emerging

55. Baguñà J. The planarian neoblast: The rambling history of its origin and some
current black boxes. Int J Dev Biol. 2012;56(1–3):19–37. 10.1387/ijdb.113463jb

56.   Martindale MQ, Henry JQ. Intracellular Fate Mapping in a Basal Metazoan, the

Ctenophore Mnemiopsis leidyi, Reveals the Origins of Mesoderm and the

Existence of Indeterminate Cell Lineages. Dev Biol. 1999;214:243–57.

http://dx.doi.org/10.1006/dbio.1999.9427

57.   Levin M, Anavy L, Cole AG, Winter E, Mostov N, Khair S, et al. The mid-

developmental transition and the evolution of animal body plans. Nature. 2016

Feb 17;531:637. https://doi.org/10.1038/nature16994

58.   Brockes JP, Kumar A. Comparative Aspects of Animal Regeneration. Annu Rev

Cell Dev Biol. 2008;24(1):525–49.

http://www.annualreviews.org/doi/10.1146/annurev.cellbio.24.110707.175336

59.   Owlarn S, Klenner F, Schmidt D, Rabert F, Tomasso A, Reuter H, et al. Generic

wound signals initiate regeneration in missing-tissue contexts. Nat Commun.

2017;8(1):1–13. http://dx.doi.org/10.1038/s41467-017-02338-x

60.   Tanaka EM, Drechsel DN, Brockes JP. Thrombin regulates S-phase re-entry by

cultured newt myotubes. Curr Biol. 1999;9(15):792–99.

http://dx.doi.org/10.1016/S0960-9822(99)80362-5

61.   Kato K, Orii H, Watanabe K, Agata K. The role of dorsoventral interaction in the

onset of planarian regeneration. Development. 1999;126(5):1031–40.

http://dx.doi.org/10.1242/dev.031682

62.   DuBuc TQ, Traylor-Knowles N, Martindale MQ. Initiating a regenerative

response; cellular and molecular features of wound healing in the cnidarian

Nematostella vectensis. BMC Biol. 2014;12(1):1–20.

http://dx.doi.org/10.1186/1741-7007-12-24

63.   Salinas-Saavedra M, Martindale MQ. Improved protocol for spawning and

immunostaining embryos and juvenile stages of the ctenophore Mnemiopsis

leidyi. Protocol exchange. 2018. http://dx.doi.org/10.1038/protex.2018.092

64.   Bhattacharjee T, Zehnder SM, Rowe KG, Jain S, Nixon RM, Sawyer WG, et al.

Writing in the granular gel medium. Sci Adv. 2015;1(8).

http://dx.doi.org/10.1126/sciadv.1500655

65.   O'Bryan CS, Kabb CP, Sumerlin BS, Angelini TE. Jammed Polyelectrolyte

Microgels for 3D Cell Culture Applications: Rheological Behavior with Added

Salts. ACS Appl Bio Mater. 2019;2(4):1509-17

http://dx.doi.org/10.1021/acsabm.8b00784

66.   Bhattacharjee T, Gil CJ, Marshall SL, Urueña JM, O'Bryan CS, Carstens M, et

al. Liquid-like Solids Support Cells in 3D. ACS Biomater Sci Eng.

2016;2(10):1787–95. http://dx.doi.org/10.1021/acsbiomaterials.6b00218

67.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et

al. Fiji: an open-source platform for biological-image analysis. Nat Methods.

2012;9(7):676–82. http://dx.doi.org/10.1038/nmeth.2019

1058

1059   **FIGURE LEGENDS**

1060

1061   **Figure 1. Cydippid stage of *Mnemiopsis leidyi* and animal surgeries.** (**A**) Lateral

1062   view of a *M. leidyi* cydippid. (**B**) Aboral view of a *M.leidyi* cydippid. Scale bars = 100μm.

1063   (**C**) Schematic representation of the body plan of a cydippid stage in a lateral and

1064   aboral views. (**D**) Diagrams showing the three types of animal surgeries performed in

1065   this study and the views presented for each one.

1066

Figure 2. Wound healing by filopodia-dependent cell crawling. (**A**) Schematic

representation of the puncture assay. (**B-E**) DIC images of the main phases of wound

closure (See **Additional file 1** for the time lapse video corresponding to these images).

(**B**) Cells from the mesoglea (yellow arrow caps) migrate upwards and adhere to the

edges of the wound. (**C**) Marginal cells of the wound gap extend filopodia to the

adjacent cells pulling the edges of the wound together. The inset shows a closer look

to the cells at the edge of the wound and yellow arrow caps point to the filopodia. (**D**)

When the diameter of the gap is considerably reduced, cells of the wound edge extend

filopodia towards the opposite edges of the gap. The inset shows a cell extending

multiple filopodia. (**E**) Network of filopodia connecting all the edges of the wound.

Timescale is in hours. Scale bars = 100μm.

1078

Figure 3. Half body regeneration in *Mnemiopsis leidyi* cydippid. (**A**) Schematic

representation of the time course of morphogenic events during cydippid half body

regeneration. All cartoons correspond to lateral views of the cydippid's body bisected

through the oral-aboral axis showing the cut site in the first plane. The apical organ is

located at the aboral end (top) and the mouth at the oral end (down). For simplicity,

tissues on the opposite body site are not depicted. (**B-F**) DIC images showing the cut

site of regenerating cydippids from 16 to 96 hab. Dotted line rectangles in (**B**), (**C**), (**D**)

and (**E**) show the area corresponding to higher magnifications on the right.

Magnifications show surface and deep planes. The vertical dotted line in (**F**) indicates

the approximate position of bisection, and all tissue in the left of the line is regenerated

tissue. N>100. Scale bars = 100 μm. Abbreviations: hours after bisection (hab), apical

organ (ao), pharynx (p), comb row (cr), ciliated furrow (cf), comb plate (cp), tentacle

bulb (tb), tentacle (t).

1092

Figure 4. Apical organ regeneration in *Mnemiopsis leidyi* cydippids. (**A**)
Schematic representation of the time course of morphogenic events during cydippid's
apical organ regeneration. Cartoons correspond to lateral views of the apical sensory
organ at different stages of regeneration. (**B-C'**) DIC images of the apical organ of an
intact cydippid in lateral (**B-B'**) and aboral (**C-C'**) views. (**D-D**) Schematic of the
components of the apical sensory organ in lateral (**D**) and aboral (**D'**) views. (**E-L'**) DIC
images showing the cut site of regenerating cydippids after apical organ amputation
from 12 to 60 hpa. Lateral and aboral views are included. Dotted line rectangles delimit
the area corresponding to higher magnifications on the right. Filled and empty white
arrow caps in (**G'**) point to aggregation of cells forming the primordia of the future
statolith and apical organ floor respectively. N>100. Scale bars = 100 μm.
Abbreviations: hours post amputation (hpa), wound epithelium (we), endodermal canal
(ec), dome cilia (dc), lithocyte (l), supporting cilia (sc), apical organ floor (ao fl), ciliated
furrows (cf).
1107
Figure 5. Cell proliferation in intact cydippids. (**A**) Schematic of the EdU pulse-
chase experiment and PH3 immunostaining in intact cydippids. (**B-D'**) Confocal stack
projections of whole intact cydippids oriented in a lateral view. White dotted rectangles
in (**B**) delimit the tentacle bulb (tb) (**C-C'''**) and apical organ (ao) (**D-D'''**) structures
showed in higher magnification at the bottom. Nuclei of S-phase cells are labeled with
EdU (magenta), M-phase cells are immunostained with anti-phospho-Histone 3 (PH3)
(green) and all nuclei are counterstained with DAPI (blue). Note that both markers of
cell proliferation (EdU and PH3) show the same pattern of distribution along the
cydippid's body. The inset in (**C'**) shows an aboral view of the tentacle bulb after EdU
staining. White asterisks in (**C'**) point to the symmetrical populations of intense cell
proliferation referred as aboral/external cell masses (aec). (**E-G'**) Confocal stack
projections of whole intact cydippids oriented in a lateral view. The time of the chase is
listed at the top of the columns, and the labeling corresponding to each panel is listed
to the left of the rows. Nuclei of S-phase cells are labeled with EdU (magenta) and all
nuclei are counterstained with DAPI (blue). Note that EdU+ cells migrate from the
tentacle bulb to the most distal end of the tentacle (yellow arrows). See **Additional file**
2 for further detail of EdU pulse-chase experiment in the tentacle bulb. Scale bars =
100 μm. Abbreviations: apical organ (ao), tentacle bulb (tb), aboral/external cell
masses (aec), lateral ridge (lr), medial ridge (mr), comb row (cr).
1127
Figure 6. Cell proliferation during half body regeneration. (**A**) Schematic of the
EdU incorporation experiment in cydippids bisected through the oral-aboral axis. (**B**)

Box plot showing the levels of cell proliferation at the wound site at different time points

after bisection. The thick horizontal bars indicate the median values. Each dot

represents one individual. (**C-F'**) Confocal stack projections of bisected cydippids

through the oral-aboral axis oriented in a lateral view showing the cut site in the first

plane. The time following bisection is listed to the left of the rows. Nuclei of S-phase

cells are labeled with EdU (magenta) and all nuclei are counterstained with DAPI

(blue). The pattern of EdU labeling corresponding to each time-point is shown in a

cartoon on the right of the rows. Dotted line rectangles in (**C**), (**D**) and (**E**) show the

area corresponding to higher magnifications on the right. White dotted lines in (**E''**) and

(**F'**) delimit the area corresponding to the regenerating comb rows and tentacle bulb.

White asterisks point to tentacle bulbs of the uncut site. Note that EdU+ cells at 72 hpa

(**F'**) are located at the oral end of the regenerating comb rows and no EdU+ cells are

detected at the aboral end where cells are already differentiated. Scale bars = 100 μm.

Abbreviations: hours after bisection (hab) comb row (cr), tentacle bulb (tb).

1144

Figure 7. Cell proliferation during apical organ regeneration. (**A**) Schematic of the

EdU incorporation experiment in cydippids in which the apical organ was amputated.

(**B**) Box plot showing the levels of cell proliferation at the wound site at different time

points post amputation. The thick horizontal bars indicate the median values. Each dot

represents one individual. (**C-H''**) Confocal stack projections of cydippids in which the

apical organ was amputated at different time points post amputation. Aboral and lateral

views are shown. The labeling corresponding to each panel is listed at the top of the

columns, and the time following amputation is listed to the left of the rows. Dotted line

rectangles in (**C**), (**D**), (**E**), (**F**), (**G**) and (**H**) show the area corresponding to higher

magnifications on the right. White rectangles in (**C''**) and (**D''**) delimit the area of apical

organ regeneration. Nuclei of S-phase cells are labeled with EdU (magenta) and all

nuclei are counterstained with DAPI (blue). DIC images of the tissue are shown in (**F'**)

and (**H'**). Scale bars = 100 μm. Abbreviations: tentacle bulb (tb), apical organ floor (ao

fl), hours post amputation (hpa).

1159

Figure 8. S-phase cells derived from the main regions of cell proliferation do not

contribute to the formation of new structures. (**A**) Schematic of the EdU pulse-

chase experiments and PH3 immunostaining in regenerating cydippids after oral-aboral

bisection and apical organ amputation. (**B-C''**) Confocal stack projections of bisected

cydippids through the oral-aboral axis oriented in a lateral view showing the cut site in

the first plane. (**D-E''**) Confocal stack projections of cydippids in which the apical organ

was amputated oriented in an aboral view. The labeling corresponding to each panel is

listed at the top of the columns, and the time of chase is listed to the left of the rows.

Nuclei of S-phase cells are labeled with EdU (magenta), M-phase cells are stained with

anti-PH3 (green) and all nuclei are counterstained with DAPI (blue). Dotted line

rectangles in (**B**), (**C**), (**D**) and (**E**) show the area corresponding to higher

magnifications on the right. The pattern of EdU and PH3 staining is shown in cartoons

on the left. Scale bars = 100 µm. Abbreviations: hours after bisection (hab), hours post

amputation (hpa), tentacle bulb (tb).

1174

Figure 9. Regeneration occurs after removal of the main regions of active cell

proliferation. (**A-C'**) DIC images of an uncut cydippid (**A**) and cydippids after tentacle

bulbs amputation (**B-C'**). Black dotted line rectangle in (**C**) show the area

corresponding to higher magnification on the right. (**D**) Schematic of the EdU

incorporation experiment in cydippids in which both tentacle bulbs were amputated. (**E-**

G'') Confocal stack projections of cydippids in which the tentacle bulbs were

amputated oriented in a lateral view at different time points post amputation. The

labeling corresponding to each panel is listed at the top of the columns, and the time

following amputation is listed to the left of the rows. Dotted line rectangles in (**E**), (**F**)

and (**G**) show the area corresponding to higher magnifications on the right. Dotted lines

in (**E'-E''**) and (**F-F''**) delimit the area corresponding to the endodermal canal. Nuclei of

S-phase cells are labeled with EdU (magenta) and all nuclei are counterstained with

DAPI (blue). Scale bars = 100µm. Abbreviations: tentacle sheath (ts), tentacle (t),

endodermal canal (ec), tentacle bulb (tb), hours post amputation (hpa).

1189

Figure 10. There is recruitment of slowly-dividing cells at the regenerating

structures. (**A**) Schematic of the EdU long pulse-chase experiment. (**B-I'**) Confocal

stack projections of whole intact cydippids oriented in an aboral (**B-E'**) and lateral (**F-I'**)

views after 2-hour EdU pulse. White dotted rectangles in (**B**) and (**F**) delimit the apical

organ (ao) (**C-C'** and **G-G'**), tentacle bulb (tb) (**D-D'** and **H-H'**) and comb row (cr) (**E-E'**

and **I-I'**) structures showed in higher magnification at the bottom. (**J-O'**) Confocal stack

projections of whole intact cydippids oriented in an aboral (**J-L'**) and lateral (**M-O'**)

views after the 2-hour EdU pulse and 5-day chase. White dotted rectangles in (**J**) and

(**M**) delimit the apical organ (ao) (**K-K'**), tentacle bulb (tb) (**L-L'** and **O-O'**) and pharynx

(p) (**N-N'**) structures showed in higher magnification at the bottom. White dotted lines in

(**L'**) and (**O'**) delimit the area corresponding to the tentacles. Note that no EdU+ cells

are detected at the tentacle bulbs nor the tentacles after the 5-day chase (**L-L'**, **O-O'**)

(see also **Additional files 10** and **11**). (**P-S'**) Confocal stack projections of cydippids in

which the apical organ was amputated after the 2-hour EdU pulse and 5-day chase at

24 hpa. Aboral (**P-Q'**) and lateral (**R-S'**) views are shown. Dotted line rectangles in (**P**)

and (**R**) show the area corresponding to higher magnifications on the bottom. Nuclei of

S-phase cells are labeled with EdU (magenta) and all nuclei are counterstained with

DAPI (blue). Scale bars = 100 μm. Abbreviations: hours post amputation (hpa), apical

organ (ao), tentacle bulb (tb), comb row (cr), pharynx (p).

1209

Figure 11. Ctenophore regeneration does not occur in the absence of cell

proliferation. (**A**) Schematic representation of the regeneration state of cydippids

shown in the panels on the right. Cartoons correspond to lateral views of the cydippid's

body bisected in half through the oral-aboral axis showing the cut site in the first plane.

(**B-E'**) DIC images of bisected cydippids in a lateral view at 72 hab. The type of

treatment corresponding to each panel is listed to the left of the rows. Dotted line

rectangles in (**B**) and (**D**) show the area corresponding to higher magnifications on the

right. Magnifications show surface (top) and deep (bottom) planes. Note that the wound

site in treated cydippids is covered by a continuous epithelium but there is no sign of

formation of missing structures. (**F**) Schematic representation of the apical sensory

organ in lateral (top) and aboral (bottom) views. Red dotted rectangle at the bottom

cartoon delimits the apical organ area shown in the images on the right. (**G-J'**) DIC

images of cydippids in which the apical organ was amputated at 72 hpa orientated in a

lateral (**G-H'**) and aboral (**I-J'**) views. The type of treatment corresponding to each

panel is listed to the top of the columns. Dotted line rectangles in (**G**), (**H**), (**I**) and (**J**)

show the area corresponding to higher magnifications on the bottom. Note that treated

cydippids show aggregation of cells around the surface of the wounded area although

none of the missing apical organ structures are formed. Scale bars = 100 μm.

Abbreviations: hours after bisection (hab), hours post amputation (hpa), apical organ

(ao), comb row (cr), tentacle bulb (tb), statolith (s), cupula (c), apical organ floor (ao fl),

endodermal canal (ec), lithocyte (l), ciliated furrow (cf).

1231

Figure 12. Regenerative ability is recovered after HU treatment ends. (**A**)

Reversible S-phase arrest after HU treatment as detected by EdU labeling performed

at the indicated time-points (red arrows). Nuclei of S-phase cells are labeled with EdU

(magenta) and all nuclei are counterstained with DAPI (blue). Note that cells have

already resumed cell cycle progression 1 day after HU wash. (**B** and **E**) Schematic

representation of HU treatment and wash experiment in bisected cydippids through the

oral-aboral axis (**B**) and cydippids in which the apical organ was amputated (**E**). (**C-C'**)

DIC images of bisected cydippids in lateral view showing the wound site at 48 hab.

Note that there is no sign of regeneration of the missing structures. (**D-D''**) DIC images

of bisected cydippids in lateral view showing the wound site at 96 hab. Note that all
missing structures (comb rows and tentacle bulb) are regenerated. The dotted line
rectangle in (**C**) and (**D**) show the area corresponding to higher magnification at the
bottom. (**F-G'**) DIC images of amputated cydippids oriented in a lateral (top panels)
and aboral (bottom panels) view showing the wound site at 48 hpa. Note that there is
no sign of regeneration of the missing structures. (**H-I'**) DIC images of amputated
cydippids oriented in a lateral (top panels) and aboral (bottom panels) view showing
the wound site at 96 hpa. Note that all components of the apical organ are
regenerated. Dotted line rectangles in (**F**), (**G**), (**H**) and (**I**) show the area corresponding
to higher magnification on the right. Scale bars = 100 µm. Abbreviations: hours after
bisection (hab), hours post amputation (hpa), apical organ (ao), comb row (cr), tentacle
bulb (tb), dome cilia (dc), lithocyte (l), supporting cilia (sc), apical organ floor (ao fl),
endodermal canal (ec), ciliated furrows (cf).

1254

Figure 13. Working model of ctenophore regeneration. Timeline of the
morphological and cellular events underlying ctenophore regeneration. Apical organ
regeneration is used as example. Proliferating cells (EdU+) are colored in magenta.
Wound closure is initiated immediately after surgery with the edges of the wound
forming a round circumference that reduces in diameter until meeting and is completed
within 1-2 hours after amputation. Reorganization of tissue including aggregation of
round-shaped cells at the wound epithelium – potentially derived from dedifferentiation
events – takes place during the first 6 hours after injury (Cell-proliferation independent
phase). Cell proliferation is activated at the wound site between 6-12 hours after
amputation and it reaches a maximum at 24 hours, when the primordia of the missing
structures are formed. After this peak of cell proliferation, the number of proliferating
cells at the wound site decreases while cells start to specify and differentiate into the
final structures.

1268

1269 **ADDITIONAL FILE LEGENDS**

1270

1271 **Additional file 1: Movie 1. Wound healing time-lapse.** DIC microscopy time-lapse
1272 movie of wound closure after puncture of the wound pictured in Figure 2. Note the
1273 migration of mesogleal cells to the edges of the wound during the wound closure
1274 process. Scale bar = 100 µm. [AVI, 13.6MB]

1275

Additional file 2: Figure S5. EdU pulse and chase experiment in the tentacle bulb.
(**A-D'**) Confocal stack projections of tentacle bulbs and tentacles in lateral view. The

time of the chase is listed to the left of the rows, and the labeling corresponding to each

panel is listed at the top of the columns. Nuclei of S-phase cells are labeled with EdU

(magenta) combined with a differential interference contrast image of the tissue (DIC).

Scale bars = 50 μm. [TIFF, 8.2MB]

1282

Additional file 3: Figure S6 and S7. EdU staining at 96 hab and 72 hpa. (**A-A''**)

Confocal stack projections of a bisected cydippid through the oral-aboral axis at 96 hab

oriented in a lateral view showing the cut site in the first plane. White asterisks point to

tentacle bulbs of the uncut site. (**B-C''**) Confocal stack projections of cydippids in which

the apical organ was amputated at 72 hpa. Examples of the two types of EdU pattern

observed at this time-point are shown. Nuclei of S-phase cells are labeled with EdU

(magenta) and all nuclei are counterstained with DAPI (blue). The pattern of EdU

labeling corresponding to each time-point is shown in a cartoon on the left of the rows.

Scale bars = 100 μm. Abbreviations: hours after bisection (hab), hours post amputation

(hpa), polar field (pf). [TIFF, 5.1MB]

1293

Additional file 4: Figure S6 and S7. PH3 immunostaining after oral-aboral

bisection and apical organ amputation. (**A-B''**) Confocal stack projections of a

bisected cydippid through the oral-aboral axis oriented in a lateral view showing the cut

site in the first plane. The time following surgery is listed to the left of the rows. M-

phase cells are stained with anti-PH3 (green) and all nuclei are counterstained with

DAPI (blue). (**C-D'**) Confocal stack projections of regenerating apical organs in a lateral

view. The time following surgery is listed to the left of the rows. M-phase cells are

stained with anti-PH3 (green) combined with DIC image of the tissue. DIC images of

surface and deep planes are shown for 24 hpa. Scale bars = 100 μm. Abbreviations:

hours after bisection (hab), hours post amputation (hpa). [TIFF, 3.1MB]

1304

Additional file 5: Figure S8. Controls of the EdU pulse-chase experiments in

regenerating cydippids. (**A**) Diagram of the experimental setup. (**B-D'**) Confocal stack

projections of an uncut cydippid just after EdU incubation (control 1). (**E-F'**) Confocal

stack projections of an amputated cydippid just after the cut (control 2). Nuclei of S-

phase cells are labeled with EdU (magenta) and all nuclei are counterstained with

DAPI (blue). White dotted rectangles in (**B**) and (**E**) show the area corresponding to

higher magnifications on the right. White arrows in (**E'**) point to the wound site. Note

that no EdU+ cells are detected at the wound site just after the cut while EdU staining

is found in the normal areas of cell proliferation (tentacle bulbs). Scale bars = 100μm.

[TIFF, 3.6MB]

1315

1316   **Additional file 6: Movie S10. Z-stack of whole body intact cydippid oriented in an**

1317   **aboral view after a 2-hour EdU incubation pulse.** Nuclei of S-phase cells are labeled

1318   with EdU (magenta) combined with a differential interference contrast image of the

1319   tissue (DIC). Note that EdU+ cells are located at the surface epidermis but not at the

1320   mesenchymal cells of the deep mesoglea. Scale bar = 100μm. [AVI, 36.4MB]

1321

1322   **Additional file 7: Movie S10. Z-stack of whole body intact cydippid oriented in a**

1323   **lateral view after a 2-hour EdU incubation pulse.** Nuclei of S-phase cells are labeled

1324   with EdU (magenta) combined with a differential interference contrast image of the

1325   tissue (DIC). Scale bar = 100μm. [AVI, 38.9MB]

1326

Additional file 8: Movie S10. Z-stack of the apical organ area of a cydippid

oriented in an aboral view after a 2-hour incubation pulse. Nuclei of S-phase cells

are labeled with EdU (magenta) combined with a differential interference contrast

image of the tissue (DIC). Note that most of the EdU+ cells are found at the epidermis

surrounding the apical organ area and some EdU+ cells are located at the apical organ

floor. Scale bar = 100μm. [AVI, 12.8MB]

1333

Additional file 9: Movie S10. Z-stack of the comb row area of a cydippid after a 2-

hour incubation pulse. Nuclei of S-phase cells are labeled with EdU (magenta) and

all nuclei are counterstained with DAPI (blue), combined with a differential interference

contrast image of the tissue (DIC). Note that EdU+ cells are found at the endodermal

canals that connect the comb rows with the digestive system. Scale bar = 100μm. [AVI,

17,7MB]

1340

1341   **Additional file 10: Movie S10. Z-stack of the tentacle apparatus of a cydippid**

1342   **oriented in an aboral view exposed to a 2-hour EdU pulse followed by a 5-day**

1343   **chase.** Nuclei of S-phase cells are labeled with EdU (magenta) combined with a

1344   differential interference contrast image of the tissue (DIC). Note that no EdU+ cells are

1345   found at the tentacle bulb nor the tentacle. Scale bar = 100μm. [AVI, 5.4MB]

1346

1347   **Additional file 11: Movie S10. Z-stack of the tentacle apparatus of a cydippid**

1348   **oriented in a lateral view exposed to a 2-hour EdU pulse followed by a 5-day**

1349   **chase.** Nuclei of S-phase cells are labeled with EdU (magenta) combined with a
1350   differential interference contrast image of the tissue (DIC). Note that no EdU+ cells are
1351   found at the tentacle bulb nor the tentacle. Scale bar = 100µm. [AVI, 1.5MB]

1352

Additional file 12: Figure S11. 5mM Hydroxyurea (HU) blocks cell proliferation in
intact and bisected cydippids. (**A**) Schematic of cell proliferation inhibitor
experiments with HU in intact cydippids. Confocal stack projections of intact cydippids
oriented in an aboral view at 24 hours (**B-C'**) and 72 hours (**D-E'**) after HU incubation.
The type of treatment is listed at the top of the columns, and the labeling corresponding
to each panel is listed to the left of the rows. Nuclei of S-phase cells are labeled with
EdU (magenta) and all nuclei are counterstained with DAPI (blue). Note that no EdU+
nuclei were detected in treated cydippids after HU incubation. (**F**) Schematic of cell
proliferation inhibitor experiments with HU in dissected cydippids. (**G-H'**) Confocal
stack projections of an untreated bisected cydippid oriented in a lateral view at 72 hab.
(**I-I'**) Confocal stack projections of a bisected cydippids treated with HU at 72 hab. (**J-**
J') Confocal stack projections of cydippids amputated from the apical organ treated
with HU at 72 hpa. The type of treatment and the time following bisection are listed at
the top of the columns, and the labeling corresponding to each panel is listed to the left
of the rows. The dotted line rectangle in (**G**) shows the area corresponding to higher
magnifications on the right. The white asterisks in (**G**), (**G'**) and (**I**) point to the tentacle
bulb of the uncut site. The white rectangle in (**J**) indicates the area of apical organ
regeneration. Nuclei of S-phase cells are labeled with EdU (magenta) and all nuclei are
counterstained with DAPI (blue). Note that no EdU+ nuclei were detected and none of
the missing structures had regenerated in treated cydippids. Scale bars = 100 µm.
Abbreviations: hours after bisection (hab), hours post amputation (hpa). [TIFF, 4.3MB]

A

Puncture

B 00:08

100 μm

C 00:53

100 μm

D 01:32

100 μm

E 02:00

100 μm

A Time course of cydippid half body regeneration

Lateral view

0   2   16   24   36   48   96 hab

Wound healing

Regeneration

Growing

B 16 hab
ao
cr
p
Surface
cf   ao   cf
Deep
ao
Apical organ — Ciliated furrows — Comb row — Pharynx

C 24 hab
Surface
cp   ao
Deep
ao
Comb plate

D 40 hab
ao
Surface
cr   cr
cr   cr
tb
cp
cp
Deep
tb
tb
Comb row — Comb plate — Tentacle bulb

E 48 hab
a.o.
Surface
cr   cr   tb   cr   cr
cp   cp
Deep
tb
tb
t
Tentacle

F 96 hab
c.r.
t.b.   a.o.   t.b.
Surface
cr   cr   cr   cr
t
Deep
cr   tb   cr   cr
cr
Aboral
Lateral

A

Time course of cydippid apical organ regeneration

0   2   12   24   48   72 hpa

Wound healing

Regeneration

Growing

Intact cydippid

B   B'   Lateral view

C   C'   Aboral view

D
Statolith
Cupula/Dome cilia
Lithocyte
Apical organ floor
Balancing cilia
Endodermal canal

D'
Polar field
Ciliated furrows
Anal pore
Statolith
Lithocyte

12 hpa

E   E'   we   ec   Lateral view

F   F'   Aboral view

48 hpa

I   I'   dc   l   sc   ao fl   ec   Lateral view

J   J'   cf   l   Aboral view

24 hpa

G   G'   Lateral view

H   H'   Aboral view

60 hpa

K   K'   dc   l   ao fl   sc   ec   Lateral view

L   L'   cf   l   Aboral view

A EdU pulse-chase experiment

EdU incubation
15 minutes

Washes thymidine +
1x FSW

Chase
t = 0, 6, 12, 24, 48h

EdU detection

PH3
immunostaining

DAPI EdU PH3

EdU

PH3

DAPI EdU

EdU

Control (no chase)

B ao / tb

B'

B''

C tb

C' tb
aec
lr
mr

C'' tb

D ao

D' ao

D'' ao

E t = 0h
tb / tb

E'

F t = 24h

F'

G t = 48h

G' ao / cr / p

A Apical organ regeneration

Apical organ amputation → Recovery to 6, 12, 24, 48 hpa → EdU incubation 15 minutes → EdU detection

A EdU pulse-chase experiment

EdU incubation 15 minutes → Washes Thymidine + FSW 1x → Oral-aboral bisection / Apical organ amputation → Recovery to 24, 48 hab/hpa → EdU detection (migrating cells) → PH3 immunostaining (local proliferation)

DAPI EdU PH3

B 24 hab — tb
B'
B''

C 48 hab
C'
C''

DAPI EdU (PH3)

D 24 hpa — tb, tb
D'
D''

E 48 hpa
E'
E''

A Uncut cydipppid

B 1hpa

C 48 hpa

C' Regenerated tentacle bulb
t
ts

D Tentacle bulb regeneration

Recovery to 18, 24, 48 hpa

EdU incubation 15 minutes

EdU detection

EdU DAPI

EdU DAPI

EdU

18 hpa

E

E'
ec

E''

24 hpa

F

F'
ec

F''

tb

48 hpa

G

G'

tb

G''

A   EdU long pulse-chase experiment

CTRL:
2 hrs pulse, no chase

EdU detection

EdU incubation
2 hours

Ctrl (t=0)

Chase

Washes thymidine
+ 1x FSW

EXP 1:
2 hrs pulse, 5 day chase

5 day chase

EdU detection

Apical organ
amputation

Recovery
for 24 hrs

EXP 2:
2 hrs pulse, 5 day chase, 24 hpa

EdU detection

CTRL: 2 hrs pulse, no chase

Aboral view

DAPI EdU | EdU

B | **B'**
cr | ao | tb

C ao | **C'** ao

D tb | **D'** tb

E cr | **E'** cr

Lateral view

DAPI EdU | EdU

F ao | **F'**
cr | tb

G ao | **G'** ao

H tb | **H'** tb

I cr | **I'** cr

EXP 1: 2 hrs pulse, 5 day chase

Aboral view

DAPI EdU | EdU

J tb | ao | **J'**

K ao | **K'** ao

L tb | **L'** tb

Lateral view

DAPI EdU | EdU

M tb | p | **M'**

N p | **N'** p

O tb | **O'** tb

EXP 2: 2 hrs pulse, 5 day chase, 24 hpa

Aboral view

DAPI EdU | EdU

P | **P'**

Q | **Q'**

Lateral view

DAPI EdU | EdU

R | **R'**

S | **S'**

A Half body regeneration

Regeneration of missing structures

No sign of regeneration

72 hab

B 1x FSW — ao

C Surface — ao

C' Deep — tb — cr cr cr

D 5mM HU — ao

E Surface — ao

E' Deep

F Apical organ regeneration

Statolith — Cupula/Dome cilia — Lithocyte — Balancing cilia — Apical organ floor — Endodermal canal

Comb row — Apical organ — Statolith — Lithocyte

G 1x FSW – 72 hpa — ao — Lateral view

H 5mM HU – 72 hpa

G' c — s — ao fl — ec

H'

I ao — Aboral view

J

I' l — cf

J'