Molecular Landscape of Anti-Drug Antibodies Reveals the Mechanism of the

Immune Response Following Treatment with TNFα Antagonists

Running title: Molecular landscape of anti-drug antibodies

[1,†]**Anna Vaisman-Mentesh,** [1,†]**Shai Rosenstein,** [2]**Miri Yavzori,** [1]**Yael Dror,** [2]**Ella**

Fudim, [2]**Bella Ungar,** [2]**Uri Kopylov,** [2]**Orit Picard,** [1]**Aya Kigel,** [2]**Shomron Ben-**

Horin, [1]**Itai Benhar,** [1,*]**Yariv Wine**

[1] George S. Wise Faculty of Life Sciences, School of Molecular Cell Biology and

Biotechnology, Tel Aviv University, Ramat Aviv, Israel

[2] Gastroenterology Department, Sheba Medical Center and Sackler School of Medicine,

Tel-Aviv University, Tel Hashomer, Israel

[†] These authors have contributed equally to this work.

*Author for correspondence: Yariv Wine

yarivwine@tauex.tau.ac.il

Keywords: Immunogenicity, anti-drug antibodies, next generation sequencing, Ig-Seq,

BCR-Seq, immune repertoire, antibody repertoire, proteomics, high-throughput

sequencing, monoclonal antibody, biologics, therapeutic antibodies

Molecular Landscape of Anti-Drug Antibodies Reveals the Mechanism of the

Immune Response Following Treatment with TNFα Antagonist

Abstract

Drugs formulated from monoclonal antibodies (mAbs) are clinically effective in various

diseases. Repeated administration of mAbs, however, elicits an immune response in the

form of anti-drug-antibodies (ADAs), thereby reducing the drug's efficacy.

Notwithstanding their importance, little is known about the molecular landscape of ADAs

and the mechanisms involved in their formation. Using a newly developed quantitative

bioimmunoassay, we found that ADA concentrations specific to TNFα antagonists can

exceed extreme concentrations of 1 mg/ml with a wide range of neutralization capacity.

Our data further suggest a preferential use of the λ light chain in a subset of neutralizing

ADAs. Moreover, we show that administration of TNFα antagonists result in a vaccine-

like response whereby ADA formation is governed by the extrafollicular T cell-

independent immune response. Our bioimmunoassay coupled with insights on the nature

of the immune response can be leveraged to improve mAb immunogenicity assessment

and facilitate improvement in therapeutic intervention strategies.

 Molecular Landscape of Anti-Drug Antibodies Reveals the Mechanism of the

 Immune Response Following Treatment with TNFα Antagonist

 Introduction

More than 30 years since the approval of the first therapeutic monoclonal antibody

(mAb) for clinical use, the therapeutic mAb market has expanded exponentially,

establishing mAbs as one of the leading biopharmaceutical therapeutic modalities (*1*).

Although mAbs hold significant promise for improving human health, their repeated

administration is often highly immunogenic and can elicit an undesirable antidrug

antibody (ADA) response (*2*). The formation of an ADA response interferes with the

effect of the drug or neutralizes it thereby altering the drug's pharmacokinetic (PK) and

pharmacodynamic (PD) properties and reducing its efficacy (*3*), which can lead to a

severe adverse immune reaction in humans (*4*).

Immunogenicity of mAbs and the formation of an ADA response has been suggested to

be dependent on the interplay between factors related to the drug itself (e.g., non-human

sequence, glycosylation, impurities, aggregation), to the patient (e.g., disease type,

genetic factors, concomitant immunomodulators), or to the drug's route and frequency of

administration (*5, 6*). However, the molecular mechanisms that lead to induction of ADA

are not well understood and were initially thought to be related to the murine origin of the

first mAbs because they were recognized as "non-self" by the human immune system.

This idea propelled the mAb discovery field to focus on engineering refined mAbs by

reducing the nonhuman portions and developing chimeric, humanized, and fully human

mAbs by using human libraries or humanized mice at the mAb discovery phase (*7*).

Unfortunately, this strategy did not abolish the immunogenicity potential of mAbs and

the associated induction of ADA. The question of why and how ADAs develop is further

complicated by data indicating that some patients develop ADAs, and some do not, and

by the observation that the extent of immunogenicity may differ among patients receiving

the same mAb (*8*). ADAs that develop in patients treated with an mAb can be stratified

into two main categories: 1) neutralizing ADAs (*nt*ADA) that directly block and interfere with the drug's ability to bind its target, and 2) nonneutralizing ADAs (i.e., binding ADA *b*ADA) that recognize other epitopes on the drug while still retaining the mAb binding activity (*9*). Neutralizing ADAs are generally considered to be more important in the clinical setting than *b*ADAs because they directly reduce a drug's efficacy. However, *b*ADAs may indirectly reduce the therapeutic efficacy of an mAb by compromising bioavailability or accelerating drug clearance from the circulation. In both cases, *nt*ADAs and *b*ADAs substantially alter the PK and PD of the mAb being administered (*10*).

Originator and biosimilar tumor necrosis factor alpha (TNFα) antagonistic mAbs are used extensively in clinical settings to treat inflammatory bowel disease (IBD; e.g., Crohn's disease and ulcerative colitis), rheumatoid arthritis, and other chronic inflammatory associated disorders such as psoriasis, psoriatic arthritis, and ankylosing spondylitis (*11*). TNFα antagonists help reduce inflammatory responses by targeting both membrane-bound and soluble TNFα. Neutralizing soluble TNFα prevents its binding to its receptor and impedes the secretion and upregulation of the signal cascade, thereby inhibiting its biological activity. The binding of TNFα antagonists to transmembrane TNFα on immune effector cells causes their destruction by inducing cell apoptosis or cell lysis through reverse signaling (*12*).

Currently, five TNFα antagonists have been approved by both the U.S. Food and Drug Administration and the European Medicines Agency: infliximab (IFX), adalimumab (ADL), etanercept, golimumab, and certolizumab pegol (*2*). Additionally, several biosimilars have already been approved or are in various stages of development (*13*). Both IFX and ADL belong to the group of TNFα antagonists and are routinely used in clinical settings to treat inflammatory diseases. IFX is a chimeric mAb (75% human and 25% murine), whereas ADL is fully human. The reported immunogenicity extent of these drugs is inconsistent. Whereas pharmaceutical companies report 10–15% and 2.6–26% immunogenicity for IFX and ADL, respectively (*14*), clinical data suggest higher immunogenicity rates for these drugs (*15*). Patients treated with IFX and ADL can be stratified based on the characteristics of their response to treatment or lack thereof. Primary nonresponders are patients whose disease does not respond to the drug at all, and

a certain subset of these may be mediated via early formation of an ADA (*15, 16*).

Secondary nonresponders are patients who initially respond to the drug but later fail

treatment, often due to development of ADA (for IFX, this was reported to develop

mostly within 12 months of treatment initiation) (*16*).

Historically, studies reporting immunogenicity following mAb administration and ADA

prevalence have been inconsistent due in part to the various assay formats used to

monitor immunogenicity in clinical trials (*17*). Current limitations of each available

format might reduce utility in clinical and research settings and complicate data

interpretation. Some assays have a poor dynamic range and may generate false negative

results because of interfering interaction with another circulating drug, or conversely,

false positive results due to the presence of other antibodies such as rheumatoid factor

(*18*). The pros and cons of available ADA detection assays were previously elaborated,

and the formation of ADA following treatment with IFX, ADL, and other TNFα

antagonists, including newly developed biosimilars, have been extensively studied and

reviewed elsewhere (*5, 19-21*).

Notwithstanding the effort invested in understanding the reasons that mAb

immunogenicity occurs and strategies to increase mAb efficacy, little is known about the

molecular mechanism that governs the formation of ADA following treatment with an

mAb. In this study, we investigated the molecular landscape of ADA following treatment

with TNFα antagonists. First, we developed a simple bioimmunoassay that accurately

quantifies ADA levels in patient sera. We further modified the bioimmunoassay to

evaluate the neutralization capacity of the ADA. Next, we aimed to profile the immune

response following mAb administration. We used flow cytometry to determine the

frequency of B cells in the circulation and whether the dynamics of the immune response

was akin to vaccine response. Finally, we used next-generation sequencing (NGS) and

high-resolution shotgun tandem mass spectrometry (LC-MS/MS) to elucidate the

molecular composition of the generated ADA. Using our bioimmunoassay we found that

ADA levels in sera from 55 patients ranged between 2.7 and 1,268.5 μg/ml. The

modified bioimmunoassay enabled us to differentiate between patients who have high

and low neutralization capacity. More interestingly, we found that patients with a high

neutralization capacity showed a strong bias in the λ/κ light chain ratio thereby

suggesting that *nt*ADA exhibits a preference for λ light chains.

To elucidate the nature of the immune response following drug administration we chose

to study a patient with IBD who was treated with IFX and who had high ADA levels and

neutralization capacity. At 10 days (D10) following IFX infusion, the patient exhibited an

approximately 16-fold and 4-fold increase in the frequency of plasmablasts (PB) and

memory B cells (mBC), respectively, compared with the preinfusion time point (D0).

Comparative NGS analysis from before and after the infusion of the antibody heavy

chain variable region (V_H) from isolated B cells showed a significant decrease in the level

of somatic hypermutation (SHM) and an increase in the length of the complementary

determining region 3 of the antibody heavy chain (CDRH3). Moreover, a proteomic

analysis of drug-specific ADA supports the observation that a preference for using λ light

chains exists and most serum antibodies map to non-class-switched IgM^+ B cells. These

data suggest a possible mechanism whereby the humoral immune response in patients

treated with TNFα antagonists is governed by a T cell-independent response that occurs

in the marginal zone (MZ). Moreover, the data support the hypothesis that the

immunocomplex that results from the drug-TNFα-ADA complex is a strong driver of

immunogenicity because these immune complexes may divert the immune response to

the MZ were B cells are activated by B cell receptor (BCR) cross-linking.

Results

Production of mAb-F(ab')$_2$ to be used in the immunoassay

To investigate the molecular landscape of ADA following mAb administration we first

aimed to develop an accurate, sensitive, robust bio-immunoassay to determine ADA

levels in sera. The working hypothesis was that anti-idiotypic antibodies dominate the

neutralizing ADA compartment (*21*) thus, the developed immunoassay was based on the

drugs' F(ab')$_2$ portion to be used as the antigen (i.e. coating agent).

To achieve this, we used the immunoglobulin G (IgG)-cleaving enzyme (IdeS), a cysteine

proteinase enzyme that proteolytically cleaves immunoglobulins below the hinge region

(*22*) (Fig. 1A). IFX was digested using IdeS by incubating 10 mg of clinical grade mAb

with IdeS to reach near complete digestion. Next, IFX-F(ab')$_2$ was purified from Fc

regions and undigested full IFX by consecutive affinity chromatography steps comprising

protein A and kappa-select columns.

Recovered IFX-F(ab')$_2$ purity was evaluated by SDS-PAGE and ELISA to ensure that the

IFX-F(ab')$_2$ exhibits no traces of IFX-Fc/undigested IFX that will contribute to the

background level when using anti-Fc HRP conjugate at the detection phase (Fig. 1B, C).

Recovered IFX-F(ab')$_2$ samples were found to be highly pure with basal anti-Fc signal

levels similar to the signal observed in the control samples (Fig. 1C). The produced IFX-

F(ab')$_2$ was tested for functionality by measuring its TNFα binding capacity, using

ELISA with TNFα as the coating agent, and was found to show similar functionality as

that of the intact IFX (Fig. 1D). ADL was subjected to the same preparative pipeline and

demonstrated similar results (Fig. S1).

ADA Standard curve

Quantification of total ADA in serum requires a standard reference. Thus, we generated a

standard ADA pool that facilitates the quantification of ADA levels in sera of patient

treated with IFX. ADA were pooled from several serum samples collected from patients

treated with IFX and purified by consecutive affinity chromatography steps comprising

protein G and a custom-made IFX-F(ab')$_2$ affinity columns. We confirmed the affinity

enrichment of ADA by applying the affinity chromatography elution fraction to ELISA

with IFX-F(ab')$_2$ as the antigen (Fig. S2A). The purity and concentration of the purified

ADA was determined by SDS-PAGE (Fig. S2B) and nanodrop.

Maximal serum concentration used in a bio-immunoassay (e.g. serum diluted 1:100 or

1:200) is a major factor that may contribute to high background signal levels due to non-

specific binding. Screening several maximal serum dilutions showed that 1:400 initial

serum dilution demonstrates the lowest background signal (data not shown). To evaluate

whether 1:400 dilution will affect the signal obtained by purified ADAs, we spiked-in

ADAs into negative control serum that was diluted 1:400.   Serial dilution of spiked-in

ADA and purified ADA showed similar signal in ELISA (Fig. S2C) indicating that serum

does not bias the ADA detection in our developed bio-immunoassay.

Quantitative measurement of ADA in serum

ADA detection is technically challenging as both the analyte and antigen are antibodies

which may result in the inability to differentiate between the mAb and ADA. To

overcome this challenge, many assays were previously developed (*5*). One of these

immunoassays is the anti-human λ chain (AHLC) immunoassay that is used in clinical

setups for monitoring the formation of ADA (*23*). The principle of this assay is to detect

ADA comprising λ light chain, thus avoiding cross reactivity with the drug that

comprises the κ light chain (Fig. 2A).

While AHLC is suitable for monitoring the development of ADA in clinical setups, when

one aims to study the molecular composition of ADA there is a need to provide

quantitative measures of total ADA in serum and to provide a solution for detecting ADA

that have kappa light chains. Thus, we developed a new bio-immunoassay based on the

$F(ab')_2$ portion of the mAb. The bio-immunoassay setup is described in Fig. 2B and is

based on mAb-$F(ab')_2$ as the coating antigen and anti-Fc HRP conjugate used as the

detection antibody. Each of the experimental setups to test ADA in serum included an

ADA negative serum as a control and ADA standard for the quantitation of total ADA.

First, we applied the newly developed bio-immunoassay on two serum samples:, one

negative and one positive for ADA as determined by the AHLC assay (AHLC[(-)] and

AHLC[(+)], respectively). As shown in Figure 2C-D, the ELISA signals obtained when

utilizing the new bio-immunoassay were higher compared to the signal obtained with the

AHLC assay. Moreover, applying the new bio-immunoassay on the AHLC[(-)] serum (no

detected ADA with the AHLC assay) detected relatively high levels of ADA. These

results indicate that not all ADA were detected with the AHLC assay as this assay is

based on the detection of ADA that comprise the λ light chain only.

Next, to extend and generalize the above results, sera from 55 patients treated with IFX were collected at the Chaim Sheba Medical Center and tested for drug levels and ADA using the AHLC assay. The established cohort showed very low drug trough levels and based on the AHLC results, sera were stratified into two groups: 26 serum samples were identified as AHLC$^{(-)}$ and 29 as AHLC$^{(+)}$. Using our newly developed quantitative bio-immunoassay, we found that ADA levels in tested sera ranged between 2.7 to 1068.5 μg/ml. Serum ADA levels using AHLC compared to the new bio-immunoassay are summarized in Table S1. More importantly, the new bio-immunoassay demonstrated improved sensitivity compared to AHLC assay manifested in the detection of higher concentrations of ADA in 40 out of the 55 serum samples, of which 12 belong to AHLC$^{(-)}$ group (Table S1). Overall, the average increase in ADA detection using the new bio-immunoassay compared to the AHLC assay was 17.3 and 60.6 fold in the AHLC$^{(+)}$ and AHLC$^{(-)}$, respectively.

Neutralization index of ADA

Due to high clinical relevance and different mechanism of action of *b*ADA and *nt*ADA, identifying their relative abundances in serum can provide valuable insights regarding the nature of the immune response following mAb administration. We therefore modified our newly developed mAb-F(ab')$_2$ based bio-immunoassay by blocking the coated IFX-F(ab')$_2$ binding site with TNFα in order to obtain a differential signal compared to the unblocked assay (Figure 3A). In order to block the binding site of IFX-F(ab')$_2$ towards TNFα and prevent the binding of anti-idiotypic ADA (i.e. *nt*ADA) to the drug, recombinant human TNFα (rhTNFα) fused to a His-tag was cloned and expressed (see materials and methods). In-house production of rhTNFα was essential, as the N terminal His-tag was used for monitoring the presence of the rhTNFα throughout the bio-immunoassay.

First, we evaluated the ability of rhTNFα to inhibit the binding of ADA to the coated IFX-F(ab')$_2$ by setting up a competitive ELISA where a series of ADA standard concentrations were incubated with a series of fixed rhTNFα concentrations (data no shown). Indeed, we observed a competitive effect while rhTNFα was fixed at the

concentratn of 5nM. This step was important as it enabled us to determine the ADA
equimolar concentration of rhTNFα to be used that will fully occupy the IFX-F(ab')$_2$
binding site and will prevent the binding of *nt*ADA. We monitored the presence of
rhTNFα using an HRP-conjugated anti-His tag antibody and observed that the ADA
concentration of 200mM completely blocks the binding of rhTNFα to the coated IFX-
F(ab')$_2$ (Fig. S3).

These data show a competitive effect between the ADA standard and the rhTNFα thus, to
ensure that the active site of the IFX-F(ab')$_2$ is occupied throughout the bio-
immunoassay, all steps were carried out while maintaining saturating concentrations of
rhTNFα (200nM) which was the equimolar concentration of ADA the blocks the binding
of rhTNFα to the drug.

In practice, IFX-F(ab')$_2$ binding site was blocked with rhTNFα by prior incubation of
serum with the coated IFX-F(ab')$_2$ hence, the differential signal w/ and w/o the presence
of rhTNFα represent the portion of ADA that could not bind the IFX-F(ab')$_2$ TNFα
binding site thus, reflects the neutralization capacity (hereby named neutralization index)
of the ADA in the tested serum. Using this assay, we evaluated the neutralization index
of the 55 sera from patients treated with IFX and 7 sera from patients treated with ADL.
In sera from patients treated with IFX, we noticed that there are two main neutralization
index patterns: those with high differential signal (Fig. 3B) and low differential signal
(Fig. 3C). All sera from patients treated with ADL were shown to have high
neutralization indexes (Fig. S4). More interestingly, we found that patients that were
stratified as AHLC$^{(+)}$ have a significantly higher neutralization index compared to those
that belong to the AHLC$^{(-)}$ group (Fig. 3D). This suggests that there is a preferential
usage of the λ light chain in *nt*ADA as the AHLC$^{(+)}$ group is *a priori* defined by the
presence of chain ADA with λ light chains.

mAb infusion induces a vaccine like immune response

To further investigate the molecular landscape of ADA we explored the dynamics of the
B cell response following mAb administration. When investigating well-controlled

clinical scenarios such as samples from post-vaccination individuals, it is convenient to
isolate the antigen-specific B cell as they peak at a well-defined time window (*24, 25*).
However, the characteristics of the humoral response and ADA encoding B cell dynamics
following mAb administration is unknown. Our working hypothesis assumed that the
immune response following mAb administration is a vaccine-like response thus, we
expected to observe a wave of PB and mBC peaking several days after mAb infusion. To
test this, we collected blood samples from a patient that was found to be positive to ADA
at two time points: prior to IFX infusion (D0) and 10 days after IFX infusion (D10).
Peripheral blood mononuclear cells (PBMCs) were sorted by FACS and the frequency of
B cell subsets including PBs (CD3$^-$CD19$^+$CD20$^-$CD27$^+$CD38^{high}) and mBCs (CD3$^-$
CD19$^+$CD20$^+$CD27$^+$) were determined. We identified a 16-fold and 4-fold increase in the
frequency of PB and mBC at D10, respectively. This data suggests that the B cell
dynamics following IFX infusion exhibits vaccine-like characteristics in accordance with
our working hypothesis (Table S2, Fig. S5).

Antibody repertoire of ADA encoding B-Cells

The waves of PB and mBC following challenge is enriched with antigen-specific B cells
(*24-26*). Based on this, a major fraction of B cells at D10 post mAb infusion is expected
to comprise B cell clones responding to the current antigen challenge. Thus, the
repertoire of B cells at two time points (pre- and post-infusion) is predicted to represent
the overall differences in the ongoing ADA encoding B cell response.

This diversity of antibodies is accomplished by several unique molecular mechanisms,
including chromosomal V(D)J rearrangement, somatic hypermutations (SHM) and class
switch recombination (*27*), processes that are mediated by recombination-activating gene
(RAG) and activation-induced cytidine deaminase (AID), respectively. The AID enzyme
functions mainly in secondary lymph nodes named germinal centers. Next-generation
sequencing (NGS) of the antibody variable regions (V genes) coupled with advanced
bioinformatics tools provides the means to elucidate the antigen-specific antibody
repertoire's immense diversity (*28*). To deep sequences antibodies' V genes, recovered
RNA from sorted PB and mBC is used as the template for first-strand cDNA synthesis,

followed by PCR amplification steps to produce barcoded amplicons of the V genes of

the antibody heavy, λ and κ light chains (V_H, V_λ and V_κ, respectively) as described

previously (*29*). While NGS of antibodies is a powerful tool for immune repertoire

analysis, relatively high rates of errors accumulate during the experimental procedure.

To overcome this challenge, we generated duplicates of the antibody V gene amplicons

and sequenced them using the Ilumina MiSeq platform (2x300bp). The resultant variable

region sequences were processed using our recently reported ASAP webserver that was

specifically developed to analyze NGS of antibody V gene sequences derived from

replicates (*27*).

In our analysis, we concentrated on several repertoire measures that collectively provide

a comprehensive molecular level characterization of the ADAs: i) V(D)J family usage; ii)

CDR3 length distribution; iii) SHM levels, and, iv) isotype distribution. Our data

revealed several interesting antibody repertoire features that shed light on the molecular

mechanism involved in the formation of ADA.

<u>V(D)J gene family usage is stable</u>

Examining the V(D)J family usage is important to determine whether the basal gene

frequency is similar to the expected frequency and if the B cell response following IFX

infusion drives B cells to exhibit a preferential V(D)J gene usage. Therefore, we

examined the frequency of family usage at two time points (D0 and D10), within two B

cell subsets (PB and mBC) and isotypes (IgG and IgM). The V(D)J family usage showed

no marked difference between the two time points, B cell subsets and isotypes. The

frequency of V-gene family usage was also found to have similar frequency profile as

previously described (*30, 31*). For example, the V gene λ light family frequency showed

that the V3, V4 and V1 have the most prevalent representation followed by V2, V5 and

V6 that had significantly lower frequencies (Fig. 4A). The same pattern trends were

identified for the D and J family usage (Fig. S6).

<u>CDRH3 length increases following IFX infusion</u>

Composed of the V(D)J join with its inherent junctional diversity, the CDRH3 specifies

the antibody V_H clonotype. The V_H clonotype is defined as the group of V_H sequences

that share germ-line V, D and J segments, have identical CDRH3 lengths, and exhibit

high amino acid identity in the CDRH3 sequences. The V_H clonotype is an important

immunological concept because it accounts for antibodies that likely originate from a

single B-cell lineage and may provide insight on the evolution of the antigen-specific

response (32).

By examining the length distribution of CDRH3 from the PB and mBC compartments,

across isotypes and time point we observed a shift towards longer CDRH3 at D10

regardless of B cell compartment and isotype (Fig. 4B). The median CDRH3 length

increased following IFX infusion in the PB and mBC compartments from 15 aa to 16 aa.

The only compartment showing no significant CDR3 length increase was within the mBC

compartment with IgG isotype that was stable with 15 aa across the two time points (Fig.

S7). Interestingly, this observation is in contrast to previous studies that reported a

decrease in the CDRH3 length post immunization with pneumococcal (33) and hepatitis

B vaccines (34) and when comparing antigen experienced B cell to naïve B cells (35).

<u>Somatic hypermutation levels decreases following IFX infusion</u>

Examining the level of SHM following vaccination provides insights regarding the extent

of the affinity maturation that antigen-stimulated B cell undergo in germinal centers. It

was previously reported that boost vaccination induces a substantial increase of the SHM

levels when comparing post- to pre-vaccination (34). Despite the vaccine like response

following IFX infusion, we observed a significant decrease on the SHM levels post-

infusion. This phenomenon was observed in the mBC as well as the PB compartments

(Fig. 4C) regardless if the mutations were synonymous and non-synonymous (Fig. S8).

Proteomic analysis of ADA

Analysis of serum antibodies provides a comprehensive profile of the humoral immune

response and is complementary to the genetic analysis derived from NGS of antibody

encoding B cell. Applying an approach that integrates NGS and tandem mass

spectrometry (LC-MS/MS) has been shown to provide valuable data regarding the
composition of antigen-specific serum antibodies and their relationship to B cells and
generates new insights regarding the development of the humoral immune response in
disease and following vaccination (*24, 25*). Here we utilized the previously developed
omics approach (*32, 36*) to elucidate the serum ADA composition following IFX
infusion. ADA from 10 ml of serum collected at D0 and D10 were subjected to protein G
affinity chromatography and total of 9 mg of recovered IgG was digested by IdeS to
remove the Fc regions that may mask the MS/MS signal obtained from low abundant
peptides. Following 5 hours of digestion, the reaction mixture was subjected to custom
made affinity column where the IFX-F(ab')$_2$ was coupled to agarose beads and served as
the antigen to isolate ADA. Recovered 48.57µg polyclonal ADA-F(ab')$_2$ (i.e., IFX-
F(ab')$_2$-specific IgG's) in the elution fraction and total F(ab')$_2$ (depleted from ADA-
F(ab')$_2$) in the flow through fraction were digested with trypsin and injected to high-
resolution tandem mass spectrometer analyzer in triplicates. LC-MS/MS raw data files
were analyzed using MaxQuant using label free quantitation mode (LFQ) and searched
against the custom antibody V gene database derived from the NGS data of B cells
isolated from the same donor. Identified peptide from the interpretation of the proteomic
spectra were stratified into three types of peptides: informative peptides (*i*Peptide) that
map uniquely to one antibody clonotype in a region that is upstream to the CDRH3, non-
informative CDRH3 peptides (*ni*CDRH3) that map to the CDRH3 region of the antibody
but do not map uniquely to a single antibody clonotype and informative CDRH3 peptides
(*i*CDRH3) that map uniquely to a single antibody clonotype. Summary of identified
peptides in LC-MS/MS are shown in Table S4. The *i*CDRH3 peptides enabled the
identification of 62 unique ADA CDRH3 clonotypes with 205 associated full-length V
gene sequences. The resulting V gene sequences were analyzed to determine their V(D)J
family usage and the B cell subset they are mapped to, based on our NGS data (Fig. 5).

The V(D)J family usage of the antibody variable region sequences that were identified by
LC-MS/MS (Fig. 5A) showed a similar distribution as observed in the NGS data (Fig.
4A, Fig. S6A). V family frequency analysis showed that the V1, V3, and V4 are the most
dominant V families followed by V2 and V5 that had significantly lower frequencies. D

family frequency analysis showed that the D6, D3, D2 and D1 have the most prevalent
representation, and J family frequency showed that the J4, J5 and J6 have the most
prevalent representation.

Next, we examined the distribution of the proteomically identified V gene sequences to B
cell subsets (Fig. 5B) and found that the V genes predominantly map to mBC from D0
(46.83%), followed by mBC from D10 (27.8%). Moreover, we found that 23.9% of V
genes map to D10 PB. Based on the dynamics of antibodies in serum (*24*), the majority of
antibodies produced following a boost challenge are the product of pre-existing mBC
cells that were re-activated following drug infusion, much like a response to a vaccine
boost (*25*).

As mentioned above, flow cytometry of B cells following IFX administration allowed us
to identify a substantial increase in the frequency of PB and mBC at D10, which suggests
that the B cell dynamics following IFX infusion exhibits vaccine-like characteristics.
Therefore, we expected to find a majority of V gene sequences mapping to IgG^+ B cells
that underwent class switch recombination in the germinal center. Surprisingly, the
majority of V gene sequences were mapped to IgM^+ B cells (Fig. 5C) indicating that the
class switch recombination process is impaired.

Next, we performed an analysis of the relative abundances of peptides that map to either
the κ and/or λ light chain constant regions. Here, we aimed to provide support to the
observation that *nt*ADA preferably use the λ light chain. By quantifying the accumulative
intensities of peptides derived from the constant region of both light chains we calculate
the change in the ratio between peptide that were derived from the affinity
chromatography elution fraction (ADA-specific) and the flow through fraction (non
ADA-specific). The expected κ/λ ratio of total IgG in serum is 2 (66% κ and 33% λ).
Indeed, proteomic analysis of the affinity chromatography flow through fractions (D0 and
D10) that represent the total IgG population depleted from ADA, resulted in average κ/λ
ratio of 2.1. The same analysis of the elution fraction showed a significant shift of the κ/λ
ratio to 1.19. These data support our observation that IFX specific ADA that were
positive in the AHLC assay as well as show neutralization index within the top %60

contain *nt*ADA that preferably use the λ light chain and thus, contribute to the shift in the

κ/λ ratio.

Discussion

The use of therapeutic mAbs in treating a wide range of diseases and disorders is growing exponentially. Nonetheless, a major shortcoming of their use is the development of ADA in patients receiving the mAb. Advances in mAb engineering have enabled the development of fully human mAbs with reduced immunogenicity without abolishing it completely. Thus an mAb administered to a patient can still induce an immune sensitization as reflected by the production of ADA, which is associated with low trough drug levels and can mediate loss of clinical response to the drug (*20*).

The precise mechanism underlying ADA production is unknown, and many questions related to its development remain unaddressed, including determining precise concentrations of ADA in serum, which portion of the ADA exhibits neutralizing capacity, the immune pathway governing the production of ADA, and ultimately, the molecular composition of ADA at the sequence level. To address these questions, we chose the chimeric TNFα antagonist IFX as the model system. First, we aimed to quantify the ADA level in patient sera. Many methods were previously reported to evaluate serum ADA levels; however, those assays provide mostly qualitative measures to assist physicians in deciding the most appropriate intervention when treating patients, and many (if not all) studies underestimated actual ADA levels (*19*). To provide quantitative measures quantitative measures describing the molecular landscape of ADA, we first developed a bioimmunoassay that would allow quantify ADA levels based on the F(ab′)$_2$ region of the mAb because previous reports indicated that the ADAs generated from mAb administration are mostly anti-idiotypic (*21*). Indeed, the bioimmunoassay demonstrated higher sensitivity compared with the AHLC assay used initially to detect ADA and was able to detect ADA when the AHLC assay could not. Leveraging its superior sensitivity, we applied our proprietary assay on sera from 55 patients treated with IFX and found that patients designated as AHLC[(+)] showed significantly higher levels of ADA (median 148.9 µg/ml) compared to the AHLC[(-)] group (median 0 µg/ml). These results support the clinical use of AHLC because overall, patients were correctly stratified leading to clinical decision-making that was based on a valid indicative assay. Notwithstanding, the applicability of the AHLC assay, the newly developed F(ab′)$_2$-based

bioimmunoassay demonstrated that ADA levels can reach extreme concentrations that

were not detected using the AHLC assay.

Some patients who develop ADA in response to IFX present a prolonged remission with

maintenance therapy despite repeated indications of high ADA and low IFX trough levels

(*20*). The mechanism of action of these ADA has significant influence on drug efficacy.

For example, *b*ADAs are most likely to enhance the clearance of a drug whereas *nt*ADAs

will prevent a drug from binding to its target. Hence, it is important to differentiate

between *b*ADAs and *nt*ADAs, or in other words, a need exists to identify sera with high

levels of *nt*ADAs that may predict the likelihood of a patient losing a favorable response

to an administrated mAb. To achieve this, we further revised our bioimmunoassay to

qualitatively measure the neutralization index of ADAs in the serum of patients treated

with IFX. Using this assay on sera from the 55 patients in our study revealed that patients

who tested positive for the AHLC antibody exhibit a significantly higher neutralization

index than patients testing negatively for it (i.e., AHLC$^{(-)}$). Noteworthy, the AHLC assay

is based on the anti-λ light chain antibody at the detection stage, suggesting that ADAs

with high neutralization index preferably use the λ light chain. This phenomenon

received additional support from our proteomic analysis in which we compared the

changes in the ratio of peptides derived from κ and λ constant light chains in mAb-

specific IgGs and in total IgGs. This analysis demonstrated that the κ/λ ratio in the total

IgG compartment is as expected and is decreased in the mAb-specific compartment (κ/λ

ratio 2.1 and 1.19 for total IgG and ADA IgG, respectively). The preferential use of the λ

light chain in neutralizing antibodies has been previously reported (*21, 37*), however, the

authors of those studies did not provide an explanation beyond the structural adaptability

of the light chain toward the target. The relevance of the reported cases showing λ chain

bias is not clear.

Here, specifically in the context of ADA, we hypothesize that the strong λ light chain

bias is associated with central tolerance processes that occur during development of B

cells in the bone marrow. B cells that encode antibodies that show autoreactivity may

avoid apoptosis by a mechanism termed receptor editing, in which the κ light chain is

substituted by the new recombined λ light chain, which results in the abolishment of

autoreactivity. These B cells may migrate to peripheral lymph nodes and engage

exogenous antigens. These central tolerance-associated processes also apply to endogenic

antibodies because they are considered to be self-molecules. Hence, administration of an

exogenous mAb may promote induction of B cells that undergo receptor editing to

replace the light chain, which eventually results in a preferable use of λ light chains by

neutralizing antibodies.

Repetitive administration of mAbs may induce a strong humoral response manifested in

the production of ADA. We hypothesized that mAb administration is similar to the

response that occurs following a boost vaccine. We and others have demonstrated that

boost vaccines induce a strong proliferation of PBs and mBCs that can be detected in

blood circulation several days after the boost. The "wave" of B cells after the boost

vaccine are dominated by class-switched antigen-specific B cells (*38*); thus repertoire

analysis of these cells can provide invaluable data about the antigen-specific antibody

repertoires. Use of flow cytometry showed an order of magnitude increase in PBs and, to

a lesser extent, mBCs 10 days after IFX infusion, suggesting that the immune response

following mAb administration is indeed similar to a vaccine response. This strong

humoral response is undesirable as opposed to a regular boost vaccination, which allows

evaluation of the efficacy of the vaccine by its ability to generate a strong humoral

response. So, mAb administration to patients' incudes a strong humoral response with the

production of high levels of ADA that as opposed to vaccines, decreases the efficacy of

the drug. To the best of our knowledge, this is the first report to identify a vaccine like

response following therapeutic mAb administration.

Next, we aimed to provide a comprehensive repertoire profile of the B cells induced after

mAb administration. To achieve this, we applied an "omics" approach as previously

described (*25, 32, 36*) that is based on the integration of high-throughput sequencing of

the V genes and proteomic analysis of serum IgGs. High-throughput sequencing of V

genes revealed no bias in the V(D)J usage across isotypes, cell types, and time point.

These data suggest that the original repertoire that existed before mAb administration and

antigen-specific repertoire that existed after mAb administration is formed by random

recombination processes without preferential use of any particular V(D)J segment.

Comparative repertoire analysis of the V genes between time points (before and after

mAb administration) revealed that after mAb administration, B cells exhibit longer

CDRH3s, lower SHM and class switch recombination (CSR) rates. Although the B cell

dynamics after mAb administration are similar to those that occur after a boost vaccine,

the repertoire measures show a different profile. It was previously reported that the

antibodies generated after a boost have shorter CDRH3, high SHM, and high CSR rates

(*33-35*).

This phenomenon can be explained by revisiting the study in which TNFα-deficient mice

exhibited a humoral immune response following vaccination that was "diverted" to the

marginal zone instead of to the germinal center (*39*). The characteristics of the immune

response in the marginal zone is directly affected by low levels of the activation-induced

cytidine deaminase (AID) enzyme that in turn is reflected in lower SHM and CSR levels.

On this basis we propose a mechanistic model according to which administration of

TNFα antagonist blocks the TNFα on one hand and induces a vaccine-like response on

the other. Due to the TNFα blockade, immune response of B cells occurs at the marginal

zone where AID is downregulated, thus the encoded ADAs exhibit lower SHM and CSR

rates. This idea is further corroborated by examining the relative high frequency of IgM$^+$

B cells after mAb administration and that serum IgGs are mapped mainly to IgM$^+$ B cells.

Additional support for this idea can be found in a previous *in vitro* study that reported

similar immune response characteristics (*40*).

Another possible mechanism that should be further explored is the strong T cell-

independent immune response in the MZ that is also induced by a drug/ADA

immunocomplex (IC). It was previously suggested that many of the immune-mediated

adverse effects attributed to ADA require the formation of an IC intermediate that can

have a variety of downstream effects (*6*). In the context of the system we investigated,

administration of a TNFα antagonist will divert the immune response to the MZ either by

TNFα blockade or by the formation of an IC carrying multiple mAbs that can induce the

cross-linking of cognate BCR. The BCR of ADA-encoding B cells will undergo co-

clustering leading to their activation in the MZ in the T-cell independent pathway. The
role of an IC in activation of the B cell response following mAb administration should be
further explored in the future.

Collectively, our data provide a comprehensive map of the molecular landscape of ADA
and addresses immunological questions related to the development of ADA following
treatment with an mAb. We believe that these data are invaluable and may assist
clinicians to better asses immunogenicity by identifying ADA at earlier stages of
treatment and provide new insights into the development of the humoral immune
response following treatment with mAbs.

Materials and Methods

Over expression and purification of rhTNFα

The sequence-encoding residues Val77–Leu233 of human TNFα was cloned and fused to
the N-terminal 6xHis tag in pET-28a+ vector (Novagen) and transformed into
Escherichia coli Rosetta (DE3) cells (Novagen). A single colony was inoculated into 2ml
LB supplemented with Kanamycin at final concentration of 100µg/ml and incubated over
night (O.N.) at 37°C, 250 RPM. The culture was next re-inoculated into a 0.5L
Erlenmeyer containing LB supplemented with Kanamycin, and grown at 37°C 250RPM
until O.D.$_{600}$~0.6-0.8 was reached. Induction was carried out by supplementing bacterial
culture with IPTG (0.1mM final concentration) and incubating the culture for 3 hours at
37°C, 250RPM. Bacterial cells were harvested by centrifugation at 8000 RPM, 15
minutes, at 10°C (SORVALL RC6 Plus, Thermo Fisher Scientific) and cell pellet was
stored O.N. at -20°C. Next, pellet was re-suspended in 30ml of binding buffer (50mM
sodium phosphate buffer pH 8.0, 300mM NaCl, 10mM imidazole) and sonicated on ice
for 8 cycles of 30 seconds pulse with 2-minute pause (W-385 sonicator, Heat Systems
Ultrasonics). Following sonication, cells were centrifuged at 12000 RPM, 30 minutes,
4°C (SORVALL RC 6+) and supernatant was applied to a HisTrap affinity column (GE
Healthcare) that was pre-equilibrated with binding buffer. All affinity purification steps
were carried out by connecting the affinity column to a peristaltic pump with flow rate of
1/ml/min. Column was washed with 5 column volumes (CV) of wash buffer (50mM
Sodium phosphate, pH 8.0, 300mM NaCl, 10% glycerol, 20mM imidazole) followed by
elution of rhTNFα with 5CV of elution buffer (50mM Sodium phosphate, pH 6.0,
300mM NaCl, 10% glycerol, 500mM imidazole). Elution was collected in 1ml fractions
and were analyzed by 12% SDS–PAGE. Fractions containing clean rhTNFα were merged
and dialyzed using Amicon Ultra (Mercury) cutoff 3K against PBS (pH 7.4). Dialysis
products were analyzed by 12% SDS–PAGE for purity and concentration was measured
using Take-5 (BioTek Instruments). To test functionality of the produced rhTNFα, 96
well plate (Nunc MaxiSorp™ flat-bottom, Thermo Fisher Scientific) was coated with
1µg/ml (in PBS) of purified rhTNFα and commercial hTNFα (PHC3011, Thermo Fisher
Scientific) and incubated at 4°C O.N. ELISA plates were then washed three times with

PBST (0.1% v/v Tween 20 in PBS) and blocked with 300µl of 2% w/v BSA in PBS for 1

hour at 37□C. Next, ELISA plates were washed three time with PBST, and incubated for

1 hour, room temperature (RT) in triplicates with anti-TNFα mAb (Infliximab or

Adalimumab) in 2% w/v BSA, PBS at the starting concentration of 50nM with 3-fold

dilution series. Plates were then washed three times with PBST with 30 second

incubation time at each washing cycle. For detection, 50µl of anti-human H+L HRP

conjugated antibody (Jackson) was added to each well (1:5000 ratio in 2% w/v BSA in

PBS) and incubated for 1 hour at RT, followed by three washing cycles with PBST.

Developing was carried out by adding 50µl of 3,3',5,5'-Tetramethylbenzidine (TMB,

Southern Biotech) and reaction was quenched by adding 50µl 0.1M sulfuric acid. Plates

were read using the Epoch Microplate Spectrophotometer ELISA plate reader (BioTek

Instruments).

582   Over expression and purification of IdeS

The coding sequence corresponding to amino acid residues 38–339 of *S. pyogenes* IdeS

(numbered from the start of the signal sequence) was sub-cloned into the expression

vector pET28b (Novagen). The coding sequencing was sub-cloned at the 3' end of

Thioredoxin 6xHis-TEV. The complete construct was sub-cloned as previously described

(*41*) and was kindly donated by Dr. Ulrich von Pawel-Rammingen from the Department

of Molecular Biology, Umea University. The transformation of pET-TRX_b plasmid

harboring the IdeS encoding gene (pET-IdeS) was carried out as follows: 200µl of

chemical-competent *E. coli* BL21-DE3 cells were thawed on ice for 20 minutes. 50ng of

the plasmid pET-IdeS was added to the thawed competent cells and incubated on ice for

20 minutes with gentle mixing every 5 minutes. Next, heat shock was applied by

incubating the cells at 42°C for 2 minutes followed by incubation on ice water for 2

minutes. For phenotypic expression, 800µl of LB was added, and cells were incubated at

37°C, 250 RPM for 1 hour in a horizontal position. Cells were plated on LB agar

supplemented with Kanamycin and incubated at 37°C overnight (O.N). Single colony

was inoculated into 2ml LB supplemented with Kanamycin and incubated O.N. at 37°C,

250 RPM. Next day, 2ml from the grown cultures were inoculated into two 2liter flasks,

each containing each 500ml LB supplemented with Kanamycin. Over expression and

purification of IdeS was carried out as described for rhTNFα with a minor modification

as follow: Ides was eluted with imidazole gradient (50, 150, 500mM imidazole), total of

20ml. 20 fractions of 1ml were collected from each elution step and evaluated for their

purity using 12% SDS–PAGE. All fractions containing clean IdeS were merged and

dialyzed O.N. at 4°C against 4L of PBS (pH 7.4), using SnakeSkin dialysis tubing with

10 kDa cutoff (Thermo Fisher Scientific). Dialysis products were analyzed by 12%

SDS–PAGE.

Production of mAb-F(ab')$_2$

Intact clinical grade IFX or ADL (designated here as mAb) were digested using in-house

produced IdeS. 10mg of mAb was incubated with 300µg of IdeS in the final volume of

500µl PBS for 2.5 hours at 37□C, followed by a spike-in of additional 300µg of IdeS to

achieve full digestion of the Fc fragments. IdeS inactivation was carried out by adding

0.1M of citric acid pH 3 and incubation for 1 minute at RT followed by the addition of

PBS (pH 7.4) to neutralized acidic pH. Next, reaction mixture was applied to a 1 mL

HiTrap KappaSelect affinity column (GE Healthcare Life Sciences). All affinity

purification steps were carried out by connecting the affinity column to a peristaltic pump

with flow rate of 1ml/min. The reaction mixture was recycled three times through the

KappaSelect column to maximize the capture of intact mAb and mAb-F(ab')$_2$.

KappaSelect column was subsequently washed with 5CV of PBS and eluted with 10CV

of 100mM glycine·HCl (pH 2.7). Collected 1ml elution fractions were immediately

neutralized with 100µl of 1.5M Tris·HCl (pH 8.8). Next, the recovered intact mAb and

mAb-F(ab')$_2$ fragments were applied to a custom packed 1ml Protein-G agarose column

(GenScript). The reaction mixture was recycled three times through the column, which

was subsequently washed with 5CV of PBS and eluted with 10CV of 100mM

glycine·HCl (pH 2.7). The 10ml elution fraction was immediately neutralized with 1ml of

1.5M Tris·HCl (pH 8.8). The recovered 10ml mAb-F(ab')$_2$ fragments were dialyzed

overnight at 4°C against 4L of PBS (pH 7.4) using SnakeSkin dialysis tubing with 10kDa

cutoff (Thermo Fisher Scientific). Recovered mAb-F(ab')$_2$ sample were evaluated for

purity by SDS-PAGE and their concentration measured by Take5 (BioTek instruments).

To test the functionality of the produced mAb-F(ab')$_2$, 96 ELISA plates (Nunc

MaxiSorp™ flat-bottom, Thermo Fisher Scientific) were coated with 1µg/ml of rhTNFα

in PBS and incubated at 4°C O.N. ELISA plates were then washed three times with

PBST and blocked with 300µl of 2% w/v BSA in PBS for 1 hour at 37^{o}C. Next, 50nM of

intact mAb and mAb-F(ab')$_2$ (IFX or ADL) in blocking solution was added to each well

in triplicates in a 3 fold dilution series, and plates were incubated at RT for 1 hour. Next,

plates were washed three times with PBST with 30 second incubation time at each

washing cycle. For detection, HRP conjugated anti-human kappa light chain (Jackson)

was added to each well (50µl, 1:5000 ratio in 2% w/v BSA in PBS) and incubated for 1

hour at RT, followed by three washing cycles with PBST. Developing was carried out by

adding 50µl of TMB and reaction was quenched by adding 0.1M sulfuric acid. Plates

were read using the Epoch Microplate Spectrophotometer ELISA plate reader. To

evaluate the purity of the mAb-Fa(b')$_2$ samples (i.e. to make sure there are no traces of

intact antibody or Fc fragment in the sample) , 96 ELISA plate (Nunc MaxiSorp™ flat-

bottom, Thermo Fisher Scientific) were coated with 5µg/ml of intact mAb and mAb-

F(ab')$_2$ in PBS and incubated at 4°C O.N. Next, plates were washed three time with

PBST and blocked with 300µl 2% w/v BSA in PBS for 1 hour at 37^{o}C. Next, plates were

washed three times with 300 µl PBST, followed by the incubation with HRP conjugated

anti-human IgG Fc antibody (Jackson) diluted 1:5000 in PBST. Developing was carried

out by adding 50µl of TMB and reaction was quenched by adding 0.1M sulfuric acid.

Plates were read using the Epoch Microplate Spectrophotometer ELISA plate reader

(BioTek Instruments).

Generation of ADA standard

A pool of 17 ADA to IFX positive sera were collected at Sheba Medical Center, and

passed through a 2ml custom packed protein G agarose column (GenScript). The pooled

sera was recycled three times over the column, which was subsequently washed with

5CV of PBS and eluted with 10CV of 100mM glycine·HCl (pH 2.7). The 10ml elution

fraction was immediately neutralized with 1ml of 1.5M Tris·HCl (pH 8.8). The purified

mAbs were immediately passed over a custom made rhTNFα affinity column (NHS-

activated agarose beads, Thermo Fisher Scientific) in gravity mode. The purified mAbs

were recycled three times over the column, which was subsequently washed with 5CV of

PBS and eluted with 10CV of 100mM glycine·HCl (pH 2.7). The 10ml elution fraction

was immediately neutralized with 1ml of 1.5M Tris·HCl (pH 8.8). The purified mAbs

were dialyzed overnight at 4°C against 4L of PBS (pH 7.4) using SnakeSkin dialysis

tubing with 10kDa cutoff (Thermo Fisher Scientific). Purified mAbs were analyzed for

purity using 12% SDS-PAGE and concentration was determined by Take3 (BioTek

instruments).

To test functionality, 96 ELISA plate were coated with 5µg/ml of mAb-F(ab')$_2$ in PBS

(pH 7.4) and incubated at 4°C O.N. ELISA plates were then washed three times with

PBST and blocked with 300µl of 2% w/v BSA in PBS for 1 hour at 37°C. Next, 50nM of

the purified ADA in blocking solution were added to each well in triplicates with 3-fold

dilution series and plates were incubated at RT for 1 hour. Next, plates were washed three

times with PBST with 30 second incubation time at each washing cycle. Next, anti-

human Fc HRP conjugate (Jackson) was added to each well at the detection phase (50µl,

1:5000 ratio in 2% w/v BSA in PBS) and incubated for 1 hour at RT, followed by three

washing cycles with PBST. Developing was carried out by adding 50µl of TMB and

reaction was quenched by adding 0.1M sulfuric acid. Plates were read using the Epoch

Microplate Spectrophotometer ELISA plate reader.

Quantitative measurement of ADA in serum

The schematic configuration of the bio-immunoassay for the quantitative measurement of

ADA in serum is described in Fig. 2B and was carried out as follows: ELISA plates that

were coated overnight at 4°C with 5µg/ml produced IFX-F(ab')$_2$ in PBS (pH 7.4). ELISA

plates were then washed three times with PBST and blocked with 300µl of 2% w/v BSA

in PBS for 1 hour at 37°C. Next, triplicates of 1:400 diluted serum samples were added at

triplicates and serially diluted 2 fold in 2% w/v BSA in PBS, 10% horse serum

(Biological Industries) and 1% Tween 20 in PBS (1:400– 1:51,200 serum dilution factor).

Plates were incubated for 1 hour at RT. On the same plate, serial dilutions of 10nM ADA

standard were incubated in triplicate and serially diluted 2 fold in 2% w/v BSA in PBS,

10% horse serum (Biological Industries) and 1% Tween 20 in PBS, to allow the

conversion of the tested serum to units per milliliter. ELISA plates were washed three

times with PBST and 50µl of HRP conjugated anti-human IgG Fc was added to each well

(50µl, 1:5000 ratio in 2% w/v BSA in PBS) and incubated for 1 hour at RT. ELISA plate

was then washed three times with PBST and developed by adding 50µl of TMB followed

by quenching with 50µl 0.1M sulfuric acid. Plates were read using the Epoch Microplate

Spectrophotometer ELISA plate reader.

Neutralization index of ADA

Neutralization capacity was determined using ELISA plates that were coated overnight at

4°C with 5µg/ml IFX-F(ab')$_2$ in PBS (pH 7.4 ELISA plates were then washed three times

with PBST and blocked with 300µl of 2% w/v BSA in PBS for 1 hr at 37□C. Next,

triplicates of 1:400 diluted serum samples were added to the negative TNFα wells and

serially diluted 2 fold in 2% w/v BSA in PBS, 10% horse serum (Biological Industries)

and 1% Tween 20 in PBS. Triplicates of 1:400 diluted serum samples with 200nM

rhTNFα were also added to the remaining wells and serially diluted 2 fold in 2% w/v

BSA in PBS, 10% horse serum (Biological Industries) and 1% Tween 20 in PBS (1:400–

1:51,200 serum dilution factor). Plates were incubated for 1 hour at RT. ELISA plates

were washed three times with PBST and 50µl of HRP conjugated anti-human IgG Fc

antibody or anti HRP conjugated His-tag antibody were added at the detection phase

(50µl, 1:5000 ratio in 2% w/v BSA in PBS) and incubated for 1 hour at RT, followed by

three washing cycles with PBST. Developing was carried out by adding 50µl of TMB

and reaction was quenched by adding 0.1M sulfuric acid. Plates were read using the

Epoch Microplate Spectrophotometer ELISA plate reader. Neutralization index was

calculated by the triplicate average differences between the ELISA equation curve

without rhTNFα and in the presence of rhTNFα where the value of the control

background is above 3 x standard deviation.

Blood processing

IFX treated patients with IBD cared for in the Department of Gastroenterology at the
Sheba medical center were included in the study. All subjects signed an informed
consent, and the study was approved by the Ethics Committee of the medical center. All
patients received IFX on a scheduled regimen and blood samples were drawn
immediately before their scheduled IFX infusion. Blood was collected into a single
Vacutainer Lithium Heparin collection tube (BD Bioscience).

For NGS analysis, blood was collected from a male donor treated with IFX, before IFX
administration and 10 days after administration. 30ml of peripheral blood were collected
into 3 single Vacutainer K-EDTA collection tubes (BD Biosciences). Collection of
peripheral blood mono-nuclear cells (PBMCs) was performed by density gradient
centrifugation, using Uni-SepMAXI+ lymphocyte separation tubes (Novamed) according
to the manufacturer's protocol.

Fluorescence-Activated Cell Sorting Analysis and sorting of B cell populations

PBMCs were stained for 15 minutes in cell staining buffer (BioLegend) at RT in the dark
using the following antibodies: anti-CD3–PerCP (clone OKT3; BioLegend), anti-CD19–
Brilliant Violet 510 (clone HIB19; BioLegend), anti-CD27–APC (clone O323;
BioLegend), anti-CD38–APC-Cy7 (clone HB-7; BioLegend), and anti-CD20–FITC
(clone 2H7; BioLegend).

The following B cell populations were sorted using a FACSAria cell sorter (BD
Bioscience):   CD3−CD19+CD20-CD27++CD38^{high} PB,   CD3−CD19+CD20-
CD27++CD38^{high}IFX-F(ab')$_2$+ IFX-F(ab')$_2$ positive PB, CD3−CD19+CD20+CD27^{var}
mBC, and CD3−CD19+CD20+CD27^{var}IFX-F(ab')$_2$+ IFX-F(ab')$_2$ positive mBC.

B cell subpopulations were sorted and collected into TRI Reagent solution (Sigma
Aldrich) and frozen at −80°C.

Amplification of V$_H$ and V$_L$ repertoires from B cells

Total RNA was isolated using RNeasy micro Kit (Qiagen), according to manufacturer's protocol. First-strand cDNA generation was performed with 100ng of isolated total RNA using a SuperScript RT II kit (Invitrogen) and oligo-dT primer, according to manufacturer's protocol. After cDNA synthesis, PCR amplification was performed to amplify the V_H and V_L genes using a primer set described previously (*25*) with overhang nucleotides to facilitate Illumina adaptor addition during the second PCR (Table S3). PCR reactions were carried out using FastStart™ High Fidelity PCR System (Roche) with the following cycling conditions: 95°C denaturation for 3 min; 95°C for 30 sec, 50°C for 30 sec, and 68°C for 1 min for four cycles; 95°C for 30 sec, 55°C for 30 sec, and 68°C for 1 min for four cycles; 95°C for 30 sec, 63°C for 30 sec, and 68°C for 1 min for 20 cycles; and a final extension at 68°C for 7 min. PCR products were purified using AMPure XP beads (Beckman Coulter), according to manufacturer's protocol (ratio x 1.8 in favor of the beads). Recovered DNA products from the first PCR was applied to a second PCR amplification to attach Illumina adaptors to the amplified V_H and V_L genes using the primer extension method as described previously (*29*). PCR reactions were carried out using FastStart™ High Fidelity PCR System (Roche) with the following cycling conditions: 95°C denaturation for 3 min; 95°C for 30 sec, 40°C for 30 sec, and 68°C for 1 min for two cycles; 95°C for 30 sec, 65°C for 30 sec, and 68°C for 1 min for 7 cycles; and a final extension at 68°C for 7 min. PCR products were applied to 1% agarose DNA gel electrophoresis and gel-purified with Zymoclean™ Gel DNA Recovery Kit (Zymo Research) according to the manufacturer's instructions. V_H and V_L libraries concentration were measured using Qubit system (Thermo Fisher Scientific) and library quality was assessed using the Bioanalyzer 2100 system (Agilent) or the 4200 TapeStation system (Agilent). All V_H libraries were produced in duplicates starting with RNA as the common source template. The V_L were produced with one replicate.

V_H and V_L libraries from sorted B cell were subjected to NGS on the MiSeq platform with the reagent kit V3 2x300 bp paired-end (Illumina), using an input concentration of 16pM with 5% PhiX.

Raw fastq files were processed using our recently reported ASAP webserver (*27*). ASAP analysis resulted in a unique, full-length V_H and V_L gene sequences database for each

769   time point. The resultant database was used as a reference database to search the LC-
770   MS/MS spectra.

Proteomic Analysis of the Serum ADA to IFX

Total IgG from each time point (D0, D10) were purified from 7-10ml of serum by protein
G enrichment. Serum was diluted 2 fold and passed through a 5ml Protein G agarose
column (GeneScript). The diluted serum was recycled three times over the column, which
was subsequently washed with 10CV of PBS and eluted with 7CV of 100mM
glycine·HCl (pH 2.7). A total of 35 fractions of 1ml were collected and immediately
neutralized with 100µl of 1.5M Tris·HCl (pH 8.8). All elution fractions were evaluated
for their purity using 12% SDS–PAGE and 11 purified 1 ml IgG fractions were combined
and dialyzed overnight at 4°C against 4L of PBS (pH 7.4) using SnakeSkin dialysis
tubing with 1 kDa cutoff (Thermo Fisher Scientific).

Next, 9mg of total IgG were digested with 100µg of IdeS in the final volume of 2ml PBS
for 5 hour at 37□C. IdeS inactivation was carried out by adding 0.1M of citric acid pH 3
and incubation for 1 minute at RT followed by the addition of PBS (pH 7.4) to neutralize
the low pH. Total serum $F(ab')_2$ was then applied to a one ml custom made affinity
column comprised of IFX-$F(ab')_2$ coupled to NHS-activated agarose beads (Thermo
Fisher Scientific). The purified serum $F(ab')_2$ were recycled three times over the affinity
column, which was subsequently washed with 5CV of PBS and eluted with 15CV of
100mM glycine·HCl (pH 2.7) and collected into Maxymum Recovery Eppendorf
(Axygen Scientific). A total of 30x0.5ml elution fractions and 1x50ml flow-through were
immediately neutralized with 50 and 100µl (respectively) of 1.5M Tris·HCl (pH 8.8). The
purified antigen-specific $F(ab')_2$ were dialyzed overnight at 4°C against 4L of PBS (pH
7.4) using SnakeSkin dialysis tubing with 10kDa cutoff (Thermo Fisher Scientific).
Elution and flow-through fractions were trypsin-digested, and resulting peptides were
fractionated and sequenced by nanoflow LC-electrospray MS/MS on an Orbitrap Velos
Pro hybrid mass spectrometer (Thermo Scientific), in the UT Austin mass spectrometry
core facility as described previously (*36*). MS/MS raw files were analyzed by MaxQuant
software version 1.6.0.16 (*42*) using the MaxLFQ algorithm (*43*) and peptide lists were

searched against the common contaminants database by the Andromeda search engine

(*44*) and a custom protein sequence database consisting of the donor-specific V_H and V_L

sequences derived from NGS of individual donor B cells. All searches were carried out

with cysteine carbamidomethylation as a fixed modification and methionine oxidations as

variable modifications. The false discovery rate was set to 0.01 for peptides with a

minimum length of seven amino acids and was determined by searching a reverse decoy

database. Enzyme specificity was set as C-terminal to arginine and lysine as expected

using trypsin as protease, and a maximum of two missed cleavages were allowed in the

database search. Peptide identification was performed with an allowed initial precursor

mass deviation up to 7ppm and an allowed fragment mass deviation of 20ppm. For LFQ

quantification the minimal ratio count was set to 2, and match between runs was

performed with three mass-spec injections originating from the same sample. MaxQunat

output analysis file, "peptides.txt", was used for further processing. Total peptides that

were identified in the elution samples were filtered using the following criteria: (a) were

not identified as contaminates; (b) did not match to the reversed decoy database; (c) were

identified as peptides derived from the region comprising the CDRH3, J region, FR4 and

the ASTK motif (derived from the N-terminal of the C_H1 region). The CDRH3 derived

peptides were further characterized as informative CDRH3 peptides (*i*CDRH3 peptides)

only if they map exclusively to a single antibody clonotype. A clonotype was defined as

all sequences that comprise CDRH3 with the same length and identity tolerating one

amino acid mismatch, and same V, J family. The intensities of high confidence *i*CDRH3

peptides were averaged between replicates while including only peptides that were

observed in at least two out of the three replicates. Clonotype frequencies within each

sample were calculated using only *i*CDRH3 peptides and were determined to be antigen-

specific if their frequency in the elution fraction was at least 5 fold greater than their

frequency in the flow-through fraction. The CDRH3 sequences identified by the mapping

of high confidence MS/MS peptides were used to generate a complete list of full length

VH sequences. These VH sequences were used to analyze the repertoire measures of the

antibodies that were identified in the donors' serum.

Statistical analysis

All curves were fitted on a sigmoidal dose–response curve and EC50 of each was

calculated. Mann-Whitney test was used to compare continuous variables. All reported P

values were two-tailed, and a P value less than 0.05 were considered statistically

significant. All statistics were performed with GraphPad Prism software (version 7, San

Diego, California).

Acknowledgments

General: We are grateful to George Georgiou for assisting with the LC-MS/MS

measurements at UT Austin, for Ulrich von Pawel-Rammingen from the Department of

Molecular Biology, Umea University who kindly donated the plasmid with the gene

encoding the IdeS.

Funding: The work was partially supported by BSF grant 2017359 (Y.W.)

Author contributions: A.V.M, S.B.H, I.B. and Y.W. conceived the research. A.V.M.

S.R. and Y.W. designed the experiments, A.V.M., S.R., M.Y., E.F. and Y.D. performed

experiments, A.V.M., S.R., S.B.H, M.Y., E.F., B.U. and U.K. collected and processed

clinical samples, A.V.M, A.K. and Y.W. carried out data analysis, A.V.M. and Y.W.

wrote the manuscript.

Competing interests: The authors declare no competing financial interests.

References
849
850
1.   A. L. Grilo, A. Mantalaris, The Increasingly Human and Profitable Monoclonal
Antibody Market. *Trends in biotechnology* **37**, 9-16 (2019).
2.   P. A. van Schouwenburg, T. Rispens, G. J. Wolbink, Immunogenicity of anti-
TNF biologic therapies for rheumatoid arthritis. *Nat Rev Rheumatol* **9**, 164-172
(2013).
3.   A. S. De Groot, D. W. Scott, Immunogenicity of protein therapeutics. *Trends in*
immunology **28**, 482-490 (2007).
4.   T. T. Hansel, H. Kropshofer, T. Singer, J. A. Mitchell, A. J. George, The safety
and side effects of monoclonal antibodies. *Nat Rev Drug Discov* **9**, 325-338
(2010).
5.   K. Bloem, B. Hernandez-Breijo, A. Martinez-Feito, T. Rispens, Immunogenicity
of Therapeutic Antibodies: Monitoring Antidrug Antibodies in a Clinical Context.
Ther Drug Monit **39**, 327-332 (2017).
6.   M. Krishna, S. G. Nadler, Immunogenicity to Biotherapeutics - The Role of Anti-
drug Immune Complexes. *Frontiers in Immunology* **7**, (2016).
7.   A. L. Nelson, E. Dhimolea, J. M. Reichert, Development trends for human
monoclonal antibody therapeutics. *Nat Rev Drug Discov* **9**, 767-774 (2010).
8.   S. Ben-Horin *et al.*, The immunogenicity of biosimilar infliximab: can we
extrapolate the data across indications? *Expert Rev Gastroenterol Hepatol* **9**
Suppl 1, 27-34 (2015).
9.   K. Bendtzen, Immunogenicity of Anti-TNF-α Biotherapies: II. Clinical Relevance
of Methods Used for Anti-Drug Antibody Detection. *Frontiers in Immunology* **6**,
109 (2015).
10. W. S. Putnam, S. Prabhu, Y. Zheng, M. Subramanyam, Y.-M. C. Wang,
Pharmacokinetic, pharmacodynamic and immunogenicity comparability
assessment strategies for monoclonal antibodies. *Trends in biotechnology* **28**,
509-516 (2010).
11. B. E. Sands *et al.*, IInfliximab maintenance therapy for fistulizing Crohn's disease.
New England Journal of Medicine **350**, 876-885 (2004).
12. H. Mitoma *et al.*, Infliximab induces potent anti-inflammatory responses by
outside-to-inside signals through transmembrane TNF-alpha. *Gastroenterology*
128, 376-392 (2005).
13. S. Ben-Horin, N. Vande Casteele, S. Schreiber, P. L. Lakatos, Biosimilars in
Inflammatory Bowel Disease: Facts and Fears of Extrapolation. *Clin*
Gastroenterol Hepatol **14**, 1685-1696 (2016).
14. M. P. Baker, H. M. Reynolds, B. Lumicisi, C. J. Bryson, Immunogenicity of
protein therapeutics: The key causes, consequences and challenges. *Self Nonself*
1, 314-322 (2010).
15. B. Ungar *et al.*, Prospective Observational Evaluation of Time-Dependency of
Adalimumab Immunogenicity and Drug Concentrations: The Poetic Study. *The*
American journal of gastroenterology **113**, 890-898 (2018).

16.   B. Ungar *et al.*, The temporal evolution of antidrug antibodies in patients with
inflammatory bowel disease treated with infliximab. *Gut* **63**, 1258-1264 (2014).
17.   F. B. Vincent *et al.*, Antidrug antibodies (ADAb) to tumour necrosis factor
(TNF)-specific neutralising agents in chronic inflammatory diseases: a real issue,
a clinical perspective. *Ann Rheum Dis* **72**, 165-178 (2013).
18.   S. Tatarewicz, J. M. Miller, S. J. Swanson, M. S. Moxness, Rheumatoid factor
interference in immunogenicity assays for human monoclonal antibody
therapeutics. *J Immunol Methods* **357**, 10-16 (2010).
19.   K. Bloem *et al.*, Systematic comparison of drug-tolerant assays for anti-drug
antibodies in a cohort of adalimumab-treated rheumatoid arthritis patients. *J
Immunol Methods* **418**, 29-38 (2015).
20.   S. Ben-Horin, Y. Chowers, Tailoring anti-TNF therapy in IBD: drug levels and
disease activity. *Nature reviews. Gastroenterology & hepatology* **11**, 243-
255 (2014).
21.   S. Ben-Horin *et al.*, The immunogenic part of infliximab is the F(ab′)2, but
measuring antibodies to the intact infliximab molecule is more clinically useful.
Gut **60**, gut.2009.201533-201548 (2010).
22.   U. von Pawel-Rammingen, B. P. Johansson, L. Björck, IdeS, a novel
streptococcal cysteine proteinase with unique specificity for immunoglobulin G.
The EMBO Journal **21**, 1607-1615 (2002).
23.   U. Kopylov *et al.*, Clinical utility of antihuman lambda chain-based
enzyme-linked immunosorbent assay (ELISA) versus double antigen ELISA for
the detection of anti-infliximab antibodies. *Inflammatory bowel diseases* **18**,
1628-1633 (2012).
24.   J. Lee *et al.*, Molecular-level analysis of the serum antibody repertoire in young
adults before and after seasonal influenza vaccination. *Nature Medicine* **22**, 1456-
+ (2016).
25.   J. J. Lavinder *et al.*, Identification and characterization of the constituent human
serum antibodies elicited by vaccination. *Proceedings of the National Academy of
Sciences of the United States of America* **111**, 2259-2264 (2014).
26.   G. Blanchard-Rohner, A. S. Pulickal, C. M. J.-v. der Zijde, M. D. Snape, A. J.
Pollard, Appearance of peripheral blood plasma cells and memory B cells in a
primary and secondary immune response in humans. *Blood* **114**, 4998-5002
(2009).
27.   O. Avram *et al.*, ASAP, A Webserver for Immunoglobulin-Sequencing Analysis
Pipeline. *Frontiers in Immunology* **9**, (2018).
28.   V. Greiff *et al.*, Quantitative assessment of the robustness of next-generation
sequencing of antibody variable gene repertoires from immunized mice. *BMC
immunology* **15**, 40 (2014).
29.   U. Menzel *et al.*, Comprehensive Evaluation and Optimization of Amplicon
Library Preparation Methods for High-Throughput Antibody Sequencing. *PLOS
ONE* **9**, (2014).
30.   E. S. Mroczek *et al.*, Differences in the composition of the human antibody
repertoire by B cell subsets in the blood. *Frontiers in immunology* **5**, 96-96
(2014).

31. J. M. Volpe, T. B. Kepler, Large-scale analysis of human heavy chain V(D)J
recombination patterns. *Immunome research* **4**, 3-3 (2008).
32. Y. Wine *et al.*, Molecular deconvolution of the monoclonal antibodies that
comprise the polyclonal serum response. *Proc Natl Acad Sci USA* **110**, 2993-2998
(2013).
33. A. Ademokun *et al.*, Vaccination-induced changes in human B-cell repertoire and
pneumococcal IgM and IgA antibody at different ages. *Aging cell* **10**, 922-930
(2011).
34. J. D. Galson *et al.*, Analysis of B Cell Repertoire Dynamics Following Hepatitis
B Vaccination in Humans, and Enrichment of Vaccine-specific Antibody
Sequences. *EBioMedicine* **2**, 2070-2079 (2015).
35. B. J. DeKosky *et al.*, Large-scale sequence and structural comparisons of human
naive and antigen-experienced antibody repertoires. *Proc Natl Acad Sci U S A*
113, E2636-2645 (2016).
36. D. R. Boutz *et al.*, Proteomic Identification of Monoclonal Antibodies from
Serum. *Analytical Chemistry* **86**, 4758-4766 (2014).
37. J. E. Robinson *et al.*, Most neutralizing human monoclonal antibodies target novel
epitopes requiring both Lassa virus glycoprotein subunits. *Nature*
communications **7**, 11544-11544 (2016).
38. A. N. Davydov *et al.*, Comparative Analysis of B-Cell Receptor Repertoires
Induced by Live Yellow Fever Vaccine in Young and Middle-Age Donors. *Front*
Immunol **9**, 2309 (2018).
39. M. Pasparakis, L. Alexopoulou, V. Episkopou, G. Kollias, Immune and
inflammatory responses in TNF alpha-deficient mice: a critical requirement for
TNF alpha in the formation of primary B cell follicles, follicular dendritic cell
networks and germinal centers, and in the maturation of the humoral immune
response. *Journal of Experimental Medicine* **184**, 1397-1411 (1996).
40. C. A. Marin, P. G. Gisbert, M. Chaparro, Immunogenicity and mechanisms
impairing the response to vaccines in inflammatory bowel disease. *World Journal*
of Gastroenterology **21**, 11273-11281 (2015).
41. K. Wenig *et al.*, Structure of the streptococcal endopeptidase IdeS, a cysteine
proteinase with strict specificity for IgG. *Proceedings of the National Academy of*
Sciences of the United States of America **101**, 17371-17376 (2004).
42. J. Cox, M. Mann, MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein
quantification. *Nature biotechnology* **26**, 1367-1372 (2008).
43. J. Cox *et al.*, Accurate proteome-wide label-free quantification by delayed
normalization and maximal peptide ratio extraction, termed MaxLFQ. *Molecular*
& cellular proteomics : MCP **13**, 2513-2526 (2014).
44. J. Cox *et al.*, Andromeda: a peptide search engine integrated into the MaxQuant
environment. *Journal of proteome research* **10**, 1794-1805 (2011).
978

Fig. 1: IFX digestion and IFX-F(ab')$_2$ purification. (A) Schematic representation of IgG digestion with IdeS. IdeS is a highly specific immunoglobulin-degrading enzyme that cleaves below the disulfide bonds in the IgG hinge region. The cleavage results in the production of one IFX-F(ab')$_2$ fragment and two ½ Fc fragments. (B) SDS-PAGE analysis of intact IgG (lane 2), following IdeS digestion (lane 3) and purified IFX-F(ab')$_2$ following a 2-step affinity chromatography purification including protein A and kappa-select columns (lane 4). (C) Presence of Fc and intact IgG traces was measured by direct ELISA where intact IFX IgG and purified IFX-F(ab')$_2$ were compared to a control antigen (streptavidin) as coating agents followed by direct incubation with an anti-Fc HRP conjugate at the detection phase. (D) The functionality of the recovered IFX-F(ab')$_2$ was confirmed by testing it for TNF binding by ELISA and compared to intact IFX. The ELISA setup included TNFα as the coating agent and anti-⬜ HRP conjugate at the detection phase. For panel C–D, triplicate averages were calculated as mean, with error bars indicating s.d.

Fig. 2: AHLC and the newly developed mAb-F(ab')₂ based bio-immunoassay configuration and their application on serum samples from patients treated with IFX. (A) AHLC assay configuration.

AHLC assay is based on an ELISA where TNFα is used as the coating agent, incubation with the mAb drug followed by differential dilutions of tested sera. anti-λ HRP conjugate is used at the detection phase. (B) The newly developed mAb-F(ab')$_2$ based bio-immunoassay configuration. The assay is based on an ELISA where mAb-F(ab')$_2$ is used as the coating agent followed by differential dilutions of tested sera. anti-Fc HRP conjugate is used at the detection phase. (C) Graph representing the ELISA results obtained utilizing the AHLC assay on two serum samples. Using this assay, one of the tested sera showed detectable levels of ADA (AHLC$^{(+)}$) and one hadno detectable levels of ADA (AHLC$^{(-)}$). (D) Both serum samples were tested by the newly developed mAb-F(ab')$_2$ based bio-immunoassay. This assay was able to detect ADA in both sera. For C–D, averages were calculated as mean from triplicates, with error bars indicating s.d.

Fig. 3: Configuration of the assay for determining the neutralization index in patient sera and ELISA results obtained from patient sera. (A)   The newly developed mAb-F(ab')$_2$ based bio-immunoassay configuration (left) and the modified configuration where mAb-F(ab')$_2$ binding site is blocked by saturating the assay with 200nM of rhTNFα (right). (B) Graph representing the ELISA results obtained utilizing the neutralization assay on serum sample that show detectable levels of ADA (AHLC$^{(+)}$) and (C) serum sample with no detectable levels of ADA (AHLC$^{(-)}$). (D) Scatter plot consolidating the neutralization index obtained by applying the immunoassay on sera from 55 patients treated with IFX (***P = 0.0003, Mann-Whitney U test). For B–C, averages were calculated as mean, with error bars indicating s.d.

A

B

C

Fig. 4: Changes in B cell repertoires following IFX infusion. mBC and PB from a patient treated with IFX were collected at two time points (D0, D10) and processed for NGS analysis. (A) The V family usage showed no difference between D0 and D10, different B cell subsets and isotypes. (B) An increase in antibody CDRH3 length and (C) a decrease in the number of bp mutations was observed at D10 in both PB and mBC.

(B) (For PB:****P<0.0001, For mBC: ****P<0.0001, Mann-Whitney U test)

(C) (For PB:****P<0.0001, For mBC: ****P<0.0001, Mann-Whitney U test)

Fig. 5: V gene and circulating antibody repertoire characteristics. (A) The V(D)J family usage of V gene sequences that were identified by LC-MS/MS. (B) Mapping of V gene sequences to B cell subsets and (C) isotypes, based on NGS data.