The Molecular Landscape of Anti-Drug Antibodies Reveals the Mechanism of the Immune Response Following Treatment with TNFα Antagonist

Running title: The molecular landscape of anti-drug antibodies

[1,†]**Anna Vaisman-Mentesh,** [1,†]**Shai Rosenstein,** [2]**Miri Yavzori,** [1]**Yael Dror,** [2]**Ella Fudim,** [2]**Bella Ungar,** [2]**Uri Kopylov,** [2]**Orit Picard,** [1]**Aya Kigel,** [2]**Shomron Ben-Horin,** [1]**Itai Benhar,** [1,*]**Yariv Wine**

[1]George S. Wise Faculty of Life Sciences, School of Molecular Cell Biology and Biotechnology, Tel Aviv University, Ramat Aviv, Israel

[2]Gastroenterology Department, Sheba Medical Center and Sackler School of Medicine, Tel-Aviv University, Tel Hashomer, Israel

[†] These authors have contributed equally to this work.
*Author for correspondence: Yariv Wine
 yarivwine@tauex.tau.ac.il

Keywords: Immunogenicity, Anti-Drug Antibodies, Next Generation Sequencing, Ig-Seq, BCR-Seq, Immune repertoire, Antibody Repertoire, Proteomics, High-throughput Sequencing, Monoclonal Antibody, Biologics, Therapeutic Antibodies

33 **Abstract**
34
Monoclonal antibodies (mAb) govern the biologics drug market with tremendous impact as clinical

therapeutics. However, repeated administration of mAbs may induce a strong immune response

manifested in the formation of anti-drug antibodies (ADA) that reduce the drug efficacy.

Notwithstanding their importance, little is known about the molecular landscape of ADA. Here we

utilized a newly developed quantitative bio-immunoassay and found that the concentration of ADA

specific to TNFα antagonists, can exceed the extreme concentrations of 1 mg/ml with a wide range of

neutralization capacity. More interestingly, our data suggest that there is a preferential usage of the λ

light chain in the subset of neutralizing ADA. Moreover, utilizing advanced omics tools we found that

TNFα antagonists administration results in a vaccine like response and ADA formation is governed by

the extrafollicular T cell independent immune response. Our developed bio-immunoassay coupled with

the insights regarding the nature of the immune response can be readily leveraged for mAb

immunogenicity assessment and assist clinicians in proposing better therapeutic intervention strategies.

MAIN TEXT

Introduction

More than 30 years since the approval of the first therapeutic monoclonal antibody (mAb) for clinical use (Muromonab-CD3, OKT3), the therapeutic mAb market has expanded exponentially bringing mAbs to establish themselves as the leading biopharmaceutical therapeutic modalities (*1, 2*). There are currently over 50 licensed mAb against a diverse panel of disease phenotypes with another ~540 in ongoing clinical development (*3*). It is predicted that in the current approval rate, more than 70 mAb products will be on the market by 2020 (*4*).

The clinical application of mAb to treat a wide range of diseases started to flourish during the 1970s following the development of the mouse hybridoma technology by Kohler and Milstein (*5*) and today, robust manufacturing and engineering platforms drive drug discovery efforts to seamlessly translate into clinical and commercial successes.

While mAbs hold significant promise for improving human health, repeated administration of mAbs is often highly immunogenic and is mostly manifested by the elicitation of an undesirable Anti-Drug Antibody (ADA) response (*6*). The formation of ADA interfere with, or neutralize the effect of the drug thus, altering the drugs' pharmacokinetics/pharmacodynamics properties, and overall reduces the drug efficacy (*7*). In a clinical context, ADA have significant impact on the safety of the administrated mAb as they may lead to serious adverse immune reactions in humans (*1, 8*).

Immunogenicity of mAbs and the formation of ADA is a multifactorial phenomenon and was suggested to be dependent on the interplay between factors related to the drug itself (e.g. non-human sequence, glycosylation, impurities, aggregation), patient related (e.g. disease type, genetic factors) as well as drugs' route and frequency of administration (*9, 10*).

The molecular mechanisms that lead to the induction of ADA are not well understood and were initially related to the murine origin of the first mAbs as they were recognized as "non-self" by the human immune system. This idea propelled the mAbs discovery field to focus on engineering refined mAbs by reducing the non-human portions while increasing its human ones. This effort led to the development of the first FDA approved chimeric mAb (Rituximab) where mouse immunoglobulin variable domains were fused to human immunoglobulin constant regions. Later in an attempt to further reduce immunogenicity, humanized mAbs were engineered by grafting the complementary determining regions (CDRs) into a human scaffold followed by the development of fully human mAbs by utilizing human

libraries or humanized mice at the mAb discovery phase (*1*). Unfortunately, these engineered mAbs did not totally diminish their immunogenicity and the associated induction of ADA. The question of why and how ADA are generated is further complicated by the fact that some patients develop ADA and some do not, and the extent of their immunogenicity may differ among patients receiving the same mAb (*11*).

ADA that are generated in patients treated with mAbs can be stratified into two main categories: i) neutralizing ADAs (*nt*ADA) are antibodies that directly block and/or interfere with the drug ability to bind its target, and ii) non-neutralizing ADA (i.e. binding ADA *b*ADA), that recognize other epitopes on the drug while still retaining the mAb binding activity (*12*).

*nt*ADAs are generally considered to be more important in the clinical setting than *b*ADA as they directly reduce drugs' efficacy. However, *b*ADA may indirectly reduce therapeutic efficacy by compromising bioavailability and/or accelerating the drug clearance from the circulation. In both cases, *nt*ADA and *b*ADA substantially alter the pharmacokinetics (PK) and pharmacodynamics (PD) of the administrated mAb (*13*).

TNFα antagonistic mAbs, originator and biosimilars, are probably the largest group of clinically used mAbs. These mAbs are used to treat Inflammatory Bowel Disease (IBD, e.g. Crohn's disease and Ulcerative Colitis), Rheumatoid arthritis, and other chronic inflammatory associated disorders such as psoriasis, psoriatic arthritis and ankylosing spondylitis (*14, 15*). TNFα antagonists help to reduce the inflammatory responses by targeting both membranes bound and soluble TNFα. Neutralizing soluble TNFα prevents its ability to bind to its receptor and impedes the secretion and up-regulation of the signal cascade thus inhibiting its biological activity. The binding of TNFα antagonists to TNFα anchored to the membrane of immune effector cells causes their destruction by inducing cell apoptosis or cell lysis through complement-dependent cytotoxicity (*16*).

Currently, there are five TNFα antagonists approved by both the FDA and EMA: infliximab, adalimumab, etanercept, golimumab and certolizumab pegol (*6*) with several biosimilars already approved or in different stages of development (*17*). Both infliximab and adalimumab belong to the TNFα antagonists' group and are routinely used in the clinic to treat inflammatory diseases. Infliximab (IFX) is a chimeric mAb (75% human and 25% murine) while Adalimumab (ADL) is fully human. The reported immunogenicity extent of these drugs is inconsistent where pharmaceutical companies report 10-15% and 2.6-26% immunogenicity for IFX and ADL, respectively (*18*). On the other hand, clinical

data suggests higher immunogenicity rates for these drugs (*19*). Patients treated with IFX and ADL can

be stratified based on the characteristics of the reaction towards the treatment. Primary non-responders

develop an immediate ADA response following infusion while secondary non-responders were reported

to develop ADA within 12 months of the initiation of treatment (*20*).

Historically, studies reporting on the response following mAb administration and ADA prevalence have

been inconsistent due, in part, to the various assay formats used to monitor immunogenicity in clinical

trials (*21, 22*). Each of the available formats has its limitations that can reduce its utility in clinical and

research settings and complicate interpretation of the data. Some assays have poor dynamic range and

may generate false-negative results because of interfering interaction with active drug or false-positive

results due to other antibodies such as rheumatoid factor (*23*). The pros and cons of the available assays

for the detection of ADA were previously elaborated and the formation of ADA following treatment

with IFX, ADL, and other TNFα antagonists including newly developed biosimilars were extensively

studied and reviewed elsewhere (*9, 24-27*).

Notwithstanding the effort invested in understanding the reasons for mAb immunogenicity and strategies

to increase their efficacy, little is known about the molecular mechanism that governs the formation of

ADA following treatment of mAbs.

In this study we investigated the molecular landscape of ADA following treatment with

TNFα antagonists. First, we developed a simple accurate bio-immunoassay to quantify ADA in patient

sera. We further modified the bio-immunoassay to evaluate the neutralization capacity of the ADA.

Next, we aimed to profile the immune response following mAb administration. We utilized flow

cytometry to determine the frequency of B cells in the circulation to determine if the immune response

is a vaccine like response with similar dynamics. Lastly, we used Next Generation Sequencing (NGS)

and high-resolution shotgun tandem mass spectrometry (LC-MS/MS) to elucidate the molecular

composition of the generated ADA.

Using our developed bio-immunoassay we found that the ADA levels in serums from 55 patients ranges

between 2.7 to 1268.5μg/ml. The modified bio-immunoassay enabled to differentiate between patients

that have high and low neutralization capacity. More interestingly, we found that patients with a high

neutralization capacity show a strong bias in the λ/κ light chain ratio thus, our data suggests that there

is a preferential usage of λ light chains by *n*tADA.

We chose an IBD patient treated with IFX with high ADA levels and neutralization capacity to elucidate
the nature of the immune response following drug administration. We found that 10 days (D10)
following IFX infusion there is a 16 fold and approximately 4 fold increase in the frequency of
plasmablast (PB) and memory B cells (mBC) respectively, compared to the pre-infusion time point (D0).
Next Generation Sequencing (NGS) analysis of the antibody heavy chain variable region (V_H) from
isolated B cells show a significant decrease in the somatic hypermutation (SHM) level and increase in
the length of the complementary determining region 3 of the antibody heavy chain (CDRH3) post-
infusion compared to pre-infusion time point. Moreover, proteomic analysis of drug-specific ADA
supports the observation there is a preferential usage of λ light chains and most serum antibodies map
to non-class switched IgM$^+$ B cells. These data suggest a possible mechanism where the humoral
immune response in patients treated with TNFα antagonists is governed by a T cell independent (TI)
response that occurs in the marginal zone (MZ). Moreover, the data supports the hypothesis that
immunocomplex comprised from drug-TNFα-ADA is a strong driver of immunogenicity as these
immune complexes may divert the immune response to the marginal zone were B cells are activated by
B cell receptor (BCR) crosslinking.
159
160

Results

Production of mAb-F(ab')$_2$ to be used in the immunoassay

To investigate the molecular landscape of ADA following mAb administration we first aimed to develop an accurate, sensitive, robust bio-immunoassay to determine ADA levels in sera. The working hypothesis was that anti-idiotypic antibodies dominate the neutralizing ADA compartment (*26, 28*) thus, the developed immunoassay was based on the drugs' F(ab')$_2$ portion to be used as the antigen (i.e. coating agent).

To achieve this, we used the immunoglobulin G (IgG)-cleaving enzyme (IdeS), a cysteine proteinase enzyme that proteolytically cleaves immunoglobulins below the hinge region (*29*) (Fig. 1A). IFX was digested using IdeS by incubating 10 mg of clinical grade mAb with IdeS to reach near complete digestion. Next, IFX-F(ab')$_2$ was purified from Fc regions and undigested full IFX by consecutive affinity chromatography steps comprising protein A and kappa-select columns.

Recovered IFX-F(ab')$_2$ purity was evaluated by SDS-PAGE and ELISA to ensure that the IFX-F(ab')$_2$ exhibits no traces of IFX-Fc/undigested IFX that will contribute to the background level when using anti-Fc HRP conjugate at the detection phase (Fig. 1B, C). Recovered IFX-F(ab')$_2$ samples were found to be highly pure with basal signal levels similar to the signal observed in the control samples (Fig. 1C). The produced IFX-F(ab')$_2$ was tested for functionality by ELISA with TNFα as the coating agent and was found to show similar functionality as observed with intact IFX (Fig. 1D). ADL was subjected to the same preparative pipeline and demonstrated similar results (Fig. S1).

ADA Standard curve

Quantification of total ADA in serum requires a standard reference thus, we generated a standard ADA pool that facilitates the quantification of ADA levels in sera of patient treated with IFX. ADA were pooled from several serum samples collected from patients treated with IFX and purified by consecutive affinity chromatography steps comprising protein G and a custom-made IFX-F(ab')$_2$ affinity columns. We confirmed the affinity enrichment of ADA by applying the affinity chromatography elution fraction to ELISA with IFX-F(ab')$_2$ as the antigen (Fig. S2A). The purity and concentration of the purified ADA was determined by SDS-PAGE (Fig. S2B) and nanodrop. Moreover, we evaluated the effect of serum on ADA detection by spiking-in varying concentrations

194   of ADA in a negative control serum and found that serum does not affect the detection of ADA

195   (Fig. S2C).

Quantitative measurement of ADA in serum

ADA detection is technically challenging as both the analyte and antigen are antibodies which may

result in the inability to differentiate between the mAb and ADA. To overcome this challenge, many

assays were previously developed (*9*). One of these immunoassays is the anti-human λ chain

(AHLC) immunoassay that is used in clinical setups for monitoring the formation of ADA (*28, 30*).

The principle of this assay is to detect ADA comprising λ light chain, thus avoiding cross reactivity

with the drug that comprises the κ light chain (Fig. 2A).

While AHLC is suitable for monitoring the development of ADA in clinical setups, when one aims

to study the molecular composition of ADA there is a need to provide quantitative measures of total

ADA in serum and to provide a solution for detecting ADA that have kappa light chains. Thus, we

developed a new bio-immunoassay based on the $F(ab')_2$ portion of the mAb. The bio-immunoassay

setup is described in Fig. 2B and is based on mAb-$F(ab')_2$ as the coating antigen and anti-Fc HRP

conjugate used as the detection antibody. Each of the experimental setups to test ADA in serum

included an ADA negative serum as a control and ADA standard for the quantitation of total ADA.

First, we applied the newly developed bio-immunoassay on two types of sera, negative and positive

for ADA as determined by the AHLC assay (AHLC[-] and AHLC[+], respectively). As shown in

Figure 2C-D, the ELISA signals obtained when utilizing the new bio-immunoassay were higher

compared to the signal obtained with the AHLC assay. Moreover, applying the new bio-

immunoassay on AHLC[-] serum (no detected ADA with the AHLC assay) detected relatively high

levels of ADA. These results indicated that not all ADA were detected with the AHLC assay as this

assay is based on the detection of ADA that comprise the λ light chain only.

Next, sera from 55 patients treated with IFX were collected at the Chaim Sheba Medical Center

and tested for drug levels and ADA using the AHLC assay. The established cohort showed very

low drug trough levels and based on the AHLC results, sera were stratified into two groups: 26

serum samples were identified as AHLC[-] and 29 as AHLC[+]. Using our newly developed

quantitative bio-immunoassay, we found that ADA levels in tested sera ranged between 2.7 to

1068.5 μg/ml. Serum ADA levels using AHLC compared to the new bio-immunoassay are

summarized in Table S1. More importantly, the new bio-immunoassay demonstrated improved

sensitivity compared to AHLC assay manifested in the detection of higher concentrations of ADA

in 40 out of the 55 serum samples of which 12 were from the 26 AHLC[(-)] samples.

Neutralization index of ADA

Due to high clinical relevance and different mechanism of action of *b*ADA and *nt*ADA, identifying

their relative abundances in serum can provide invaluable insights regarding the nature of the

immune response following mAb administration. We modified our newly developed mAb-F(ab')$_2$

based bio-immunoassay by blocking the coated IFX-F(ab')$_2$ binding site with TNFα in order to

obtain a differential signal compared to the unblocked assay (Figure 3A). In order to block the

binding site of IFX-F(ab')$_2$ towards TNFα and prevent the binding of anti-idiotypic ADA (i.e.

*nt*ADA) to the drug, recombinant human TNFα (rhTNFα) fused to a His-tag was cloned and

expressed (see materials and methods). In-house production of rhTNFα was essential, as the N

terminal His-tag was used for monitoring the presence of the rhTNFα throughout the bio-

immunoassay.

First, we evaluated the ability of rhTNFα to inhibit the binding of ADA to the coated IFX-F(ab')$_2$

by setting up a competitive ELISA where a series of ADA standard concentrations were incubated

with a fixed concentration (5nM) of rhTNFα. We monitored the presence of rhTNFα using an HRP-

conjugated anti-His tag antibody and observed that an increase in ADA concentrations inhibits the

binding of rhTNFα to the coated IFX-F(ab')$_2$ (Fig. S3).

These data show a competitive effect between the ADA standard and the rhTNFα thus, to ensure

that the active site of the IFX-F(ab')$_2$ is occupied throughout the bio-immunoassay, all steps were

carried out while maintaining saturating concentrations of rhTNFα (200nM) which is above the

maximum concentration of ADA standard applied in the assay.

In practice, IFX-F(ab')$_2$ binding site was blocked with rhTNFα by prior incubation of serum with

the coated IFX-F(ab')$_2$ hence, the differential signal w/ and w/o the presence of rhTNFα reflects the

neutralization capacity (hereby named neutralization index) of the ADA in the tested serum. Using

this assay, we evaluated the neutralization index of the 55 sera from patients treated with IFX and

7 sera from patients treated with ADL. In sera from patients treated with IFX, we noticed that there

are two main neutralization index patterns: those with high differential signal (Fig. 3B) and low

differential signal (Fig. 3C). All sera from patients treated with ADL were shown to have high

neutralization indexes (Fig. S4). More interestingly, we found that patients that were stratified as

AHLC[(+)] have a significantly higher neutralization index compared to those that belong to the

AHLC$^{(-)}$ group (Fig. 3D). This suggests that there is a preferential usage of the λ light chain in

*nt*ADA as the AHLC$^{(+)}$ group is *a priori* defined by the presence of chain ADA with λ light chains.

Flow cytometry of B cells following mAb administration

To further investigate the molecular landscape of ADA we explored the dynamics of the B cell

response following mAb administration. When investigating well-controlled clinical scenarios such

as samples from post-vaccination individuals, it is convenient to isolate the antigen-specific B cell

as they peak at a well-defined time window (*31-33*). However, the characteristics of the humoral

response and ADA encoding B cell dynamics following mAb administration is unknown. Our

working hypothesis postulated that the immune response following mAb administration is a

vaccine-like response thus, we expected to observe a wave of PB and mBC peaking several days

post mAb infusion. To confirm this, we collected blood samples from a patient that was found to

exhibit ADA (20.22µg/ml at D0, 32.33µg/ml at D10) at two time points: prior to IFX infusion (D0)

and 10 days post IFX infusion (D10). Total peripheral blood mononuclear cells (PBMCs) were

analyzed by FACS and the frequency of B cell subsets were determined. We identified a substantial

increase in the frequency of PB and mBC at D10, suggesting that the B cell dynamics following

IFX infusion exhibits vaccine-like characteristics in accordance with our working hypothesis (Table

S2, Fig. S5).

BCR-seq following IFX infusion

A well-functioning humoral response relies on its ability to generate an exceptional high diversity

in the genes encoding the variable region of B cell receptors and their secreted form, the antibodies.

This diversity of antibodies is accomplished by several unique molecular mechanisms, including

chromosomal V(D)J rearrangement during B cell maturation in the bone marrow, N-P

addition/deletion in the ligated V(D)J genes and somatic hypermutations (SHM) following antigen

stimuli in the peripheral lymph nodes (*34*). Further chromosomal rearrangement processes facilitate

class switch recombination generating different antibody isotypes (i.e. IgG, IgA etc.). The processes

following antigen engagement are collectively termed affinity maturation are mediated by the

enzyme activation-induced cytidine deaminase (AID) and take place in secondary lymph nodes

named germinal centers. Collectively, the underline processes can generate antibodies with the

theoretical diversity estimated to be 10^{11}–10^{13} in humans (*35*).

NGS of the antibody variable regions coupled with advanced bioinformatics tools provides the

means to elucidate the antibody repertoire's immense diversity (*36*). Due to the recombination and

non-templated diversification mechanisms that generates the CDRH3, it is considered the most

diverse determinant in terms of length and sequence and pivotal for antibody specificity hence, is

the focus of our repertoire analysis.

Isolating antigen-specific B cells at steady state is challenging as the frequency of these cells in the

circulation is very low. However, once the immune system is challenged with an antigen, the wave

of PB and mBC is dominated by antigen-specific B cells (*31, 32, 37*). Based on this, a major fraction

of B cells at D10 post mAb infusion should comprise B cell clones responding to the current antigen

challenge. Thus, the repertoire of B cells at two time points (pre- and post-infusion) will underline

the overall differences in the ongoing ADA encoding B cell response.

PBs ($CD3^-CD19^+CD20^-CD27^+CD38^{high}$) and mBCs ($CD3^-CD19^+CD20^+CD27^+$) were collected

using FACS followed by total RNA isolation. The recovered RNA was used as the template for

first-strand cDNA synthesis, followed by PCR amplification steps to produce barcoded amplicons

of the variable regions of the antibody heavy, λ and κ light chains (V_H, V_λ and V_κ, respectively) as

described previously (*38*). While NGS of antibodies is a powerful tool for immune repertoire

analysis, relatively high rates of errors accumulate during the experimental procedure. To

overcome this challenge, we generated duplicates of the antibody V gene amplicons and sequenced

them using the Ilumina MiSeq platform (2x300bp). The resultant variable region sequences were

processed using our recently reported ASAP webserver that was specifically developed to analyze

NGS of antibody V gene sequences derived from replicates (*34*).

In our analysis, we concentrated on several repertoire measures that collectively provide a

comprehensive molecular level characterization of the ADAs: i) V(D)J family usage; ii) CDR3

length distribution; iii) SHM levels, and, iv) isotype distribution.

Our data revealed a number of interesting antibody repertoire features that may shed light on the

molecular mechanism involved in the formation of ADA.

<u>V(D)J gene family usage is stable</u>

Examining the V(D)J family usage is important to determine whether the basal gene frequency is

similar to the expected frequency and if the B cell response following IFX infusion drives B cells

to exhibit a preferential V(D)J gene usage. Therefore, we examined the frequency of family usage

at two time point (D0 and D10), within two B cell subsets (PB and mBC) and isotypes (IgG and

IgM).

The V(D)J family usage showed no marked difference between the two time points, B cell subsets

and isotypes. The frequency of V-gene family usage was also found to have similar frequency

profile as previously described (*39, 40*). For example, the V gene λ light family frequency showed

that the V3, V4 and V1 have the most prevalent representation followed by V2, V5 and V6 that had

significantly lower frequencies (Fig. 4A). The same trends were identified for the D and J family

usage (Fig. S6).

CDRH3 length increases following IFX infusion

Composed of the V(D)J join with its inherent junctional diversity, the CDRH3 specifies the

antibody V_H clonotype. The V_H clonotype is defined as the group of V_H sequences that share germ-

line V, D and J segments, have identical CDRH3 lengths, and exhibit high amino acid identity in

the CDRH3 sequences. The V_H clonotype is an important immunological concept because it

accounts for antibodies that likely originate from a single B-cell lineage and may provide insight

on the evolution of the antigen-specific response (*41*).

By examining the length distribution of CDRH3 from the PB and mBC compartments, across

isotypes and time point we observed a shift towards longer CDRH3 at D10 regardless of B cell

compartment and isotype (Fig. 4B). The median CDRH3 length increased following IFX infusion

in the PB and mBC compartments from 15 aa to 16 aa. The only compartment showing no

significant CDR3 length increase was within the mBC compartment with IgG isotype that was

stable with 15 aa across the two time points (Fig. S7). Interestingly, this observation is in contrast

to previous studies that reported a decrease in the CDRH3 length post immunization with

pneumococcal (*42*) and hepatitis B vaccines (*43*) and when comparing antigen experienced B cell

to naïve B cells (*44*).

Somatic hypermutation levels decreases following IFX infusion

Examining the level of SHM following vaccination provides insights regarding the extent of the

affinity maturation that antigen-stimulated B cell undergo in germinal centers. It was previously

reported that boost vaccination induces a substantial increase of the SHM levels when comparing

post- to pre-vaccination (*43*). Despite the vaccine like response following IFX infusion, we

observed a significant decrease on the SHM levels post-infusion. This phenomenon was observed

in the mBC as well as the PB compartments (Fig. 4C) regardless if the mutations were synonymous

and non-synonymous (Fig. S8).

Proteomic analysis of ADA

Analysis of serum antibodies provides a comprehensive profile of the humoral immune response

and is complementary to the genetic analysis derived from NGS of antibody encoding B cell.

Applying an approach that integrates NGS and tandem mass spectrometry (LC-MS/MS) has been

shown to provide invaluable data regarding the composition of antigen-specific serum antibodies
and their relationship to B cells and generates new insights regarding the development of the
humoral immune response in disease (*45*) and following vaccination (*31, 32*). Here we utilized the
previously developed omics approach (*41, 46*) to elucidate the serum ADA composition following
IFX infusion. ADA from 10 ml of serum collected at D0 and D10 were subjected to protein G
affinity chromatography and total of 9 mg of recovered IgG was digested by IdeS to remove the Fc
regions that may mask the MS/MS signal obtained from low abundant peptides. Following 5 hours
of digestion, the reaction mixture was subjected to custom made affinity column where the IFX-
F(ab')$_2$ was coupled to agarose beads and served as the antigen to isolate ADA. Recovered 48.57μg
polyclonal ADA-F(ab')$_2$ (IFX-F(ab')$_2$-specific IgG's) in the elution fraction and total F(ab')$_2$
(depleted from ADA-F(ab')$_2$) in the flow through fraction were digested with trypsin and injected
to high-resolution tandem mass spectrometer analyzer in triplicates. LC-MS/MS raw data files were
analyzed using MaxQuant using label free quantitation mode (LFQ) and searched against the
custom antibody V gene database derived from the NGS data of B cells isolated from the same
donor. Identified peptide from the interpretation of the proteomic spectra were stratified into three
types of peptides: informative peptides (*i*Peptide) that map uniquely to one antibody clonotype in a
region that is upstream to the CDRH3, non-informative CDRH3 peptides (*ni*CDRH3) that map to
the CDRH3 region of the antibody but do not map uniquely to a single antibody clonotype and
informative CDRH3 peptides (*i*CDRH3) that map uniquely to a single antibody clonotype.
Summary of identified peptides in LC-MS/MS are shown in Table S4. The *i*CDRH3 peptides
enabled the identification of 62 unique ADA CDRH3 clonotypes with 205 associated full-length V
gene sequences. The resulting V gene sequences were analyzed to determine their V(D)J family
usage and the B cell subset they are mapped to, based on our NGS data (Fig. 5).
The V(D)J family usage of the antibody variable region sequences that were identifies by LC-
MS/MS (Fig. 5A) showed a similar distribution as observed in the NGS data (Fig. 4A, Fig. S6A).
V family frequency analysis showed that the V1, V3, and V4 are the most dominant V families
followed by V2 and V5 that had significantly lower frequencies. D family frequency analysis
showed that the D6, D3, D2 and D1 have the most prevalent representation, and J family frequency
showed that the J4, J5 and J6 have the most prevalent representation.
Next, we examined the distribution of the proteomically identified V gene sequences to B cell
subsets (Fig. 5B) and found that the of V genes predominantly map to mBC from D0 (46.83%),
followed by mBC from D10 (27.8%). Moreover, we found that 23.9% of V genes map to D10 PB.
Based on the dynamics of antibodies in serum (*31*), the majority of antibodies produced following

a boost challenge are the product of pre-existing mBC cells that were re-activated following drug
infusion, much like a response to a vaccine boost (*32*).
As mentioned above, flow cytometry of B cells following IFX administration allowed us to identify
a substantial increase in the frequency of PB and mBC at D10, which suggests that the B cell
dynamics following IFX infusion exhibits vaccine-like characteristics. Therefore, we expected to
find a majority of V gene sequences mapping to IgG$^+$ B cells that underwent class switch
recombination in the germinal center. Surprisingly, the majority of V gene sequences were mapped
to IgM$^+$ B cells (Fig. 5C) indicating that the class switch recombination process is impaired.
Next, we performed an analysis of the relative abundances of peptides that map to either the κ
and/or λ light chain constant regions. Here, we aimed to provide support to the observation that
*nt*ADA preferably use the λ light chain. By quantifying the accumulative intensities of peptides
derived from the constant region of both light chains we calculate the change in the ratio between
peptide that were derived from the affinity chromatography elution fraction (ADA-specific) and the
flow through fraction (non ADA-specific). The expected κ/λ ratio of total IgG in serum is 2 (*47*)
(66% κ and 33% λ). Indeed, proteomic analysis of the affinity chromatography flow through
fractions (D0 and D10) that represent the total IgG population depleted from ADA, resulted in
average κ/λ ratio of 2.1. The same analysis of the elution fraction showed a significant shift of the
κ/λ ratio to 1.19. These data support our observation that IFX specific ADA that were positive in
the AHLC assay as well as show neutralization index within the top 60% contain *nt*ADA preferably
use the λ light chain and thus, contribute to the shift in the κ/λ ratio.
415
416
417

Discussion

The administration of therapeutic mAbs to treat a wide range of diseases and disorders is growing exponentially. Along with the increase in their applicability, mAb still suffer from the serious shortcoming reflected mainly in the induction of ADA.

Advancements in mAb engineering enabled the development of fully human mAbs with reduced immunogenicity however, in various autoimmune and chronic inflammatory disorders, mAb administrated to patients still induce a strong immune response reflected in the production of ADA that drive the treated patients to lose response to the drug.

The development of ADA to mAbs is associated with low trough drug levels and have been implicated as predisposing factors to treatment failure (*48-52*).

The exact mechanism underlying ADA production is unknown and many questions related to the development of ADA remain un-addressed including, what are the exact concentration of ADA in serum, what portion of the ADA exhibit neutralizing capacity, what is the immune pathway the governess the production of ADA and ultimately, what is the molecular composition of ADA at the sequence level. To address these questions, we chose the chimeric TNFα antagonist - IFX, as the model system. First, we aimed to quantify the ADA level in patient's serum. Many approaches were previously reported to evaluate the serum ADA levels however, these assays mainly provide qualitative measures assisting physicians in decision making for the appropriate intervention strategy when treating the patients and many (if not all) result in underestimation of the actual ADA levels. Moreover, when one aims to investigate the molecular landscape of ADA, it is imperative to provide quantitative measures, thus we first developed a bio-immunoassay to quantify the ADA levels. The developed bio-immunoassay was based on the $F(ab')_2$ region of the mAb, as it was previously reported that the generated ADA are mostly anti-idiotypic (*26*). The developed bio-immunoassay demonstrated higher sensitivity compared to the AHLC assay used initially to detect ADA, and was able to detect ADA where AHLC assay could not. Leveraging its superior sensitivity, we applied the developed assay on sera from 55 patients treated with IFX, and found that patients that were stratified as AHLC[(+)], showed significantly higher levels of ADA (median 148.9μg/ml) compared to the AHLC[(-)] (median 0μg/ml). These results support the usage of AHLC in the clinic as overall the patients were stratified correctly and clinical decision making was based on a valid indicative assay. Notwithstanding the applicability of the AHLC assay, the new develop

F(ab')$_2$ based bio-immunoassay demonstrated that ADA levels can reach extreme concentration
that were not detected using the AHLC assay.

The mechanism of action of ADA have great influence on drug efficacy. For example, *b*ADA are
most likely to enhance the clearance of the drug while *nt*ADA will prevent the drug to bind to its
target. Noteworthy, some patients who develop ADA to IFX present a prolonged remission with
maintenance therapy, despite repeated indications of high ADA and low IFX trough levels (*25*).
Hence, there is great importance to differentiate between *b*ADA and *nt*ADA or in other words, there
is a need to identify serum with high levels of *nt*ADA that may predict the likelihood of the patient
to lose response to the administrated mAb. To achieve this, we further revised our developed bio-
immunoassay to qualitatively measure the neutralization index of the ADA in the serum of patients
treated with IFX. Applying this assay on sera from the 55 patients revealed that AHLC[(+)] patients
exhibit significantly higher neutralization index than those that were stratified as AHLC[(-)].
Noteworthy, AHLC assay is based on the anti-λ light chain antibody at the detection stage,
suggesting that ADA with high neutralization index preferably use the λ light chain. This
phenomena got additional support from the proteomic analysis comparing the changes in the ratio
of peptides derived from κ and λ constant light chain in mAb-specific IgGs and total IgGs. This
analysis demonstrated that the κ/λ in the total IgG compartment is as expected and is decreased in
the mAb-specific compartment (κ/λ ratio 2.1 and 1.19 for the total IgG and mAb-specific IgG,
respectively). The preferential usage of λ light chain in neutralizing antibodies has been previously
reported (*26, 53*) however, the authors in these cases did not provide an explanation beyond the
structural adaptability of the light chain towards the target. However, the relevance of the reported
cases showing λ chain bias is not clear.

Here, specifically in the context of ADA, we hypothesize that the strong λ light chain bias is related
to processes that are associated with the central tolerance during the development of the B cell in
the bone marrow. B cells that encode antibodies that show autoreactivity may avoid apoptosis by a
mechanism named receptor editing where the κ light chain is substituted by the new recombined λ
light chain that may results in the abolishment of autoreactivity. These B cells may migrate to the
peripheral lymph nodes and engage exogenous antigens. The central tolerance associated processes
apply also to endogenic antibodies because they are considered to be self-molecules. Hence, the
administration of exogenous mAbs may promote the induction of B cells that undergo receptor
editing to replace the light chain that eventually results in a preferable usage of λ light chains by
neutralizing antibodies.

The repetitive administration of mAbs may induce a strong humoral response manifested in the production of ADA. We hypothesized that mAb administration is similar to the response following a boost vaccine. We and others demonstrated that boost vaccines induce a strong proliferation of PB and mBC that can be detected several days post boost. The "wave" of B cells post boost are dominated by class switched antigen-specific B cells (*54*) thus, repertoire analysis of these cells can provide invaluable data regarding the antigen-specific antibody repertoires. Utilizing flow cytometry showed that there is an order of magnitude increase of PB and to the lesser extent mBC 10 days post IFX infusion, suggesting that the immune response following mAb administration is a vaccine like response. Needless to mention, this strong humoral response is undesirable as opposed to regular boost vaccination where the efficacy of the vaccine is evaluated by its ability to generate a strong humoral response.

Next, we aimed to provide a comprehensive repertoire profile of the B cells induced post mAb administration. To achieve this, we applied an omics approach as previously described (*32, 41, 46*) that is based on the integration of high throughput sequencing of the V genes and proteomic analysis of serum IgGs. High throughput sequencing of V genes revealed that there is no bias in the V(D)J usage across isotypes, cell types and time point. This data suggests that the original repertoire pre-mAb administration and antigen-specific post-mAb administration is formed by random recombination processes without preferential usage of any particular V(D)J segment. Comparative repertoire analysis of the V genes between time points (pre- and post mAb administration), revealed that the B cells post mAb administration exhibit longer CDRH3s, lower SHM levels and lower-class switch recombination (CSR) rates. While the B cell dynamics post mAb administration are similar to the dynamics post boost vaccine, the repertoire measures show a different profile. It was previously reported that the antibodies generated post boost have shorter CDRH3, high SHM and CSR rates (*42-44*).

This phenomena can be explained by revisiting the study of TNFα deficient mice were the humoral immune response following vaccination was "diverted" to the marginal zone instead to the germinal center (*55*). The characteristics of the immune response in the marginal zone is directly affected by low levels of the activation-induced cytidine deaminase enzyme (AID) that in turn is reflected in lower SHM and CSR levels. Based on this we propose the following mechanistic model: administration of TNFα antagonists block the TNFα on one hand and induces a vaccine like response on the other. Due to TNFα blockade, B cells immune response occurs at the marginal zone where AID is down regulated thus, the encoded ADA exhibit lower SHM and CSR rates. This idea is further corroborated by examining the relative high frequency of IgM$^+$ B cells post mAb

administration and that serum IgGs are mapped mainly to IgM$^+$ B cells. Additional support for this

idea can be found in a previous *in vitro* study that reported similar immune response characteristics

(*56*).

Another possible mechanism that should be further explored is the strong TI immune response in

the marginal zone is also induced by drug/ADA immunocomplexes (IC). It was previously

suggested that many of the immune-mediated adverse effects attributed to ADA require the

formation of IC intermediate that can have a variety of downstream effects (*10*). In the context of

our investigated system, the administration of TNFα antagonist will divert the immune response to

the marginal zone either by TNFα blockade or by the formation of ICs carrying multiple mAbs that

can induce the crosslinking of cognate BCR. The BCR of ADA encoding B cells will undergo co-

clustering leading to their activation in the marginal zone in the T-independent pathway. The role

of IC in the activation of the B cell response following mAb administration should be further

explored in the future.

Collectively, our data provide a comprehensive map of the molecular landscape of ADA and

addresses immunological question related to the development of ADA following treatment of

mAbs. We believe that these data are invaluable and could assist clinician to better asses

immunogenicity by identifying ADA at earlier stages of treatment as well as provide new insight

regarding the development of the humoral immune response following treatment with mAbs.

Materials and Methods

Over expression and purification of rhTNFα

The sequence-encoding residues Val77–Leu233 of human TNFα was cloned and fused to the N-terminal 6xHis tag in pET-28a+ vector (Novagen) and transformed into *Escherichia coli* Rosetta (DE3) cells (Novagen). A single colony was inoculated into 2ml LB supplemented with Kanamycin at final concentration of 100µg/ml and incubated over night (O.N.) at 37°C, 250 RPM. The culture was next re-inoculated into a 0.5L Erlenmeyer containing LB supplemented with Kanamycin, and grown at 37°C 250RPM until $O.D._{600}$~0.6-0.8 was reached. Induction was carried out by supplementing bacterial culture with IPTG (0.1mM final concentration) and incubating the culture for 3 hours at 37°C, 250RPM. Bacterial cells were harvested by centrifugation at 8000 RPM, 15 minutes, at 10°C (SORVALL RC6 Plus, Thermo Fisher Scientific) and cell pellet was stored O.N. at -20°C. Next, pellet was re-suspended in 30ml of binding buffer (50mM sodium phosphate buffer pH 8.0, 300mM NaCl, 10mM imidazole) and sonicated on ice for 8 cycles of 30 seconds pulse with 2 minute pause (W-385 sonicator, Heat Systems Ultrasonics). Following sonication, cells were centrifuged at 12000 RPM, 30 minutes, 4°C (SORVALL RC 6+) and supernatant was applied to a HisTrap affinity column (GE Healthcare) that was pre-equilibrated with binding buffer. All affinity purification steps were carried out by connecting the affinity column to a peristaltic pump with flow rate of 1/ml/min. Column was washed with 5 column volumes (CV) of wash buffer (50mM Sodium phosphate, pH 8.0, 300mM NaCl, 10% glycerol, 20mM imidazole) followed by elution of rhTNFα with 5CV of elution buffer (50mM Sodium phosphate, pH 6.0, 300mM NaCl, 10% glycerol, 500mM imidazole). Elution was collected in 1ml fractions and were analyzed by 12% SDS–PAGE. Fractions containing clean rhTNFα were merged and dialyzed using Amicon Ultra (Mercury) cutoff 3K against PBS (pH 7.4). Dialysis products were analyzed by 12% SDS–PAGE for purity and concentration was measured using Take-5 (BioTek Instruments). To test functionality of the produced rhTNFα, 96 well plate (Nunc MaxiSorp™ flat-bottom, Thermo Fisher Scientific) was coated with 1µg/ml (in PBS) of purified rhTNFα and commercial hTNFα (PHC3011, Thermo Fisher Scientific) and incubated at 4°C O.N. ELISA plates were then washed three times with PBST (0.1% v/v Tween 20 in PBS) and blocked with 300µl of 2% w/v BSA in PBS for 1 hour at 37^{0}C. Next, ELISA plates were washed three time with PBST, and incubated for 1 hour, room temperature (RT) in triplicates with anti-TNFα mAb (Infliximab or Adalimumab) in 2% w/v BSA, PBS at the starting concentration of 50nM with 3 fold dilution series. Plates were then washed three times with PBST with 30 second incubation time at each washing cycle. For detection , 50µl of anti-human H+L HRP conjugated antibody (Jackson) was added to each well (1:5000 ratio in 2% w/v

579   BSA in PBS) and incubated for 1 hour at RT, followed by three washing cycles with PBST.

580   Developing was carried out by adding 50µl of 3,3',5,5'-Tetramethylbenzidine (TMB, Southern

581   Biotech) and reaction was quenched by adding 50µl 0.1M sulfuric acid. Plates were read using the

582   Epoch Microplate Spectrophotometer ELISA plate reader (BioTek Instruments).

Over expression and purification of IdeS

The coding sequence corresponding to amino acid residues 38–339 of *S. pyogenes* IdeS (numbered

from the start of the signal sequence) was sub-cloned into the expression vector pET28b (Novagen).

The coding sequencing was sub-cloned at the 3' end of Thioredoxin 6xHis-TEV. The complete

construct was sub-cloned as previously described (*57*) and was kindly donated by Dr. Ulrich von

Pawel-Rammingen from the Department of Molecular Biology, Umea University. The

transformation of pET-TRX_b plasmid harboring the IdeS encoding gene (pET-IdeS) was carried

out as follows: 200µl of chemical-competent *E. coli* BL21-DE3 cells were thawed on ice for 20

minutes. 50ng of the plasmid pET-IdeS was added to the thawed competent cells and incubated on

ice for 20 minutes with gentle mixing every 5 minutes. Next, heat shock was applied by incubating

the cells at 42°C for 2 minutes followed by incubation on ice water for 2 minutes. For phenotypic

expression, 800µl of LB was added, and cells were incubated at 37°C, 250 RPM for 1 hour in a

horizontal position. Cells were plated on LB agar supplemented with Kanamycin and incubated at

37°C overnight (O.N). Single colony was inoculated into 2ml LB supplemented with Kanamycin

and incubated O.N. at 37°C, 250 RPM. Next day, 2ml from the grown cultures were inoculated into

two 2liter flasks, each containing each 500ml LB supplemented with Kanamycin. Over expression

and purification of IdeS was carried out as described for rhTNFα with a minor modification as

follow: Ides was eluted with imidazole gradient (50, 150, 500mM imidazole), total of 20ml. 20

fractions of 1ml were collected from each elution step and evaluated for their purity using 12%

SDS–PAGE. All fractions containing clean IdeS were merged and dialyzed O.N. at 4°C against 4L

of PBS (pH 7.4), using SnakeSkin dialysis tubing with 10 kDa cutoff (Thermo Fisher Scientific).

Dialysis products were analyzed by 12% SDS–PAGE.

Production of mAb-F(ab')₂

608   Intact clinical grade IFX or ADL (designated here as mAb) were digested using in-house produced

609   IdeS. 10mg of mAb was incubated with 300µg of IdeS in the final volume of 500µl PBS for 2.5

610   hours at 37⁰C, followed by a spike-in of additional 300µg of IdeS to achieve full digestion of the

611   Fc fragments. IdeS inactivation was carried out by adding 0.1M of citric acid pH 3 and incubation

612   for 1 minute at RT followed by the addition of PBS (pH 7.4) to neutralized acidic pH. Next, reaction

mixture was applied to a 1 mL HiTrap KappaSelect affinity column (GE Healthcare Life Sciences). All affinity purification steps were carried out by connecting the affinity column to a peristaltic pump with flow rate of 1ml/min. The reaction mixture was recycled three times through the KappaSelect column to maximize the capture of intact mAb and mAb-F(ab')$_2$. KappaSelect column was subsequently washed with 5CV of PBS and eluted with 10CV of 100mM glycine·HCl (pH 2.7). Collected 1ml elution fractions were immediately neutralized with 100µl of 1.5M Tris·HCl (pH 8.8). Next, the recovered intact mAb and mAb-F(ab')$_2$ fragments were applied to a custom packed 1ml Protein-G agarose column (GenScript). The reaction mixture was recycled three times through the column, which was subsequently washed with 5CV of PBS and eluted with 10CV of 100mM glycine·HCl (pH 2.7). The 10ml elution fraction was immediately neutralized with 1ml of 1.5M Tris·HCl (pH 8.8). The recovered 10ml mAb-F(ab')$_2$ fragments were dialyzed overnight at 4°C against 4L of PBS (pH 7.4) using SnakeSkin dialysis tubing with 10kDa cutoff (Thermo Fisher Scientific). Recovered mAb-F(ab')$_2$ sample were evaluated for purity by SDS-PAGE and their concentration measured by Take5 (BioTek instruments).

To test the functionality of the produced mAb-F(ab')$_2$, 96 ELISA plates (Nunc MaxiSorp™ flat-bottom, Thermo Fisher Scientific) were coated with 1µg/ml of rhTNFα in PBS and incubated at 4°C O.N. ELISA plates were then washed three times with PBST and blocked with 300µl of 2% w/v BSA in PBS for 1 hour at 37°C. Next, 50nM of intact mAb and mAb-F(ab')$_2$ (IFX or ADL) in blocking solution was added to each well in triplicates in a 3 fold dilution series, and plates were incubated at RT for 1 hour. Next, plates were washed three times with PBST with 30 second incubation time at each washing cycle. For detection, HRP conjugated anti-human kappa light chain (Jackson) was added to each well (50µl, 1:5000 ratio in 2% w/v BSA in PBS) and incubated for 1 hour at RT, followed by three washing cycles with PBST. Developing was carried out by adding 50µl of TMB and reaction was quenched by adding 0.1M sulfuric acid. Plates were read using the Epoch Microplate Spectrophotometer ELISA plate reader. To evaluate the purity of the mAb-Fa(b')$_2$ samples (i.e. to make sure there are no traces of intact antibody or Fc fragment in the sample) , 96 ELISA plate (Nunc MaxiSorp™ flat-bottom, Thermo Fisher Scientific) were coated with 5µg/ml of intact mAb and mAb-F(ab')$_2$ in PBS and incubated at 4°C O.N. Next, plates were washed three time with PBST and blocked with 300µl 2% w/v BSA in PBS for 1 hour at 37°C. Next, plates were washed three times with 300 µl PBST, followed by the incubation with HRP conjugated anti-human IgG Fc antibody (Jackson) diluted 1:5000 in PBST. Developing was carried out by adding 50µl of TMB and reaction was quenched by adding 0.1M sulfuric acid. Plates were read using the Epoch Microplate Spectrophotometer ELISA plate reader (BioTek Instruments).

Generation of ADA standard

A pool of 17 ADA to IFX positive sera were collected at Sheba Medical Center, and passed through a 2ml custom packed protein G agarose column (GenScript). The pooled sera was recycled three times over the column, which was subsequently washed with 5CV of PBS and eluted with 10CV of 100mM glycine·HCl (pH 2.7). The 10ml elution fraction was immediately neutralized with 1ml of 1.5M Tris·HCl (pH 8.8). The purified mAbs were immediately passed over a custom made rhTNFα affinity column (NHS-activated agarose beads, Thermo Fisher Scientific) in gravity mode. The purified mAbs were recycled three times over the column, which was subsequently washed with 5CV of PBS and eluted with 10CV of 100mM glycine·HCl (pH 2.7). The 10ml elution fraction was immediately neutralized with 1ml of 1.5M Tris·HCl (pH 8.8). The purified mAbs were dialyzed overnight at 4°C against 4L of PBS (pH 7.4) using SnakeSkin dialysis tubing with 10kDa cutoff (Thermo Fisher Scientific). Purified mAbs were analyzed for purity using 12% SDS-PAGE and concentration was determined by Take3 (BioTek instruments).

To test functionality, 96 ELISA plate were coated with 5µg/ml of mAb-F(ab')$_2$ in PBS (pH 7.4) and incubated at 4°C O.N. ELISA plates were then washed three times with PBST and blocked with 300µl of 2% w/v BSA in PBS for 1 hour at 37°C. Next, 50nM of the purified ADA in blocking solution were added to each well in triplicates with 3 fold dilution series and plates were incubated at RT for 1 hour. Next, plates were washed three times with PBST with 30 second incubation time at each washing cycle. Next, anti-human Fc HRP conjugate (Jackson) was added to each well at the detection phase (50µl, 1:5000 ratio in 2% w/v BSA in PBS) and incubated for 1 hour at RT, followed by three washing cycles with PBST. Developing was carried out by adding 50µl of TMB and reaction was quenched by adding 0.1M sulfuric acid. Plates were read using the Epoch Microplate Spectrophotometer ELISA plate reader.

Quantitative measurement of ADA in serum

The schematic configuration of the bio-immunoassay for the quantitative measurement of ADA in serum is described in Fig. 2B and was carried out as follows: ELISA plates that were coated overnight at 4°C with 5µg/ml produced IFX-F(ab')$_2$ in PBS (pH 7.4). ELISA plates were then washed three times with PBST and blocked with 300µl of 2% w/v BSA in PBS for 1 hour at 37°C. Next, triplicates of 1:400 diluted serum samples were added at triplicates and serially diluted 2 fold in 2% w/v BSA in PBS, 10% horse serum (Biological Industries) and 1% Tween 20 in PBS (1:400–1:51,200 serum dilution factor). Plates were incubated for 1 hour at RT. On the same plate, serial dilutions of 10nM ADA standard were incubated in triplicate and serially diluted 2 fold in 2% w/v BSA in PBS, 10% horse serum (Biological Industries) and 1% Tween 20 in PBS, to allow the

681 conversion of the tested serum to units per milliliter. ELISA plates were washed three times with

682 PBST and 50µl of HRP conjugated anti-human IgG Fc was added to each well (50µl, 1:5000 ratio

683 in 2% w/v BSA in PBS) and incubated for 1 hour at RT. ELISA plate was then washed three times

684 with PBST and developed by adding 50µl of TMB followed by quenching with 50µl 0.1M sulfuric

685 acid. Plates were read using the Epoch Microplate Spectrophotometer ELISA plate reader.

Neutralization index of ADA

Neutralization capacity was determined using ELISA plates that were coated overnight at 4°C with

5µg/ml IFX-F(ab')$_2$ in PBS (pH 7.4 ELISA plates were then washed three times with PBST and

blocked with 300µl of 2% w/v BSA in PBS for 1 hr at 37^{0}C. Next, triplicates of 1:400 diluted serum

samples were added to the negative TNFα wells and serially diluted 2 fold in 2% w/v BSA in PBS,

10% horse serum (Biological Industries) and 1% Tween 20 in PBS. Triplicates of 1:400 diluted

serum samples with 200nM rhTNFα were also added to the remaining wells and serially diluted 2

fold in 2% w/v BSA in PBS, 10% horse serum (Biological Industries) and 1% Tween 20 in PBS

(1:400– 1:51,200 serum dilution factor). Plates were incubated for 1 hour at RT. ELISA plates were

washed three times with PBST and 50µl of HRP conjugated anti-human IgG Fc antibody or anti

HRP conjugated His-tag antibody were added at the detection phase (50µl, 1:5000 ratio in 2% w/v

BSA in PBS) and incubated for 1 hour at RT, followed by three washing cycles with PBST.

Developing was carried out by adding 50µl of TMB and reaction was quenched by adding 0.1M

sulfuric acid. Plates were read using the Epoch Microplate Spectrophotometer ELISA plate reader.

Neutralization index was calculated by the triplicate average differences between the ELISA

equation curve without rhTNFα and in the presence of rhTNFα where the value of the control

background is above 3 x standard deviation.

Blood processing

IFX treated patients with IBD cared for in the Department of Gastroenterology at the Sheba medical

center were included in the study. All subjects signed an informed consent, and the study was

approved by the Ethics Committee of the medical center. All patients received IFX on a scheduled

regimen and blood samples were drawn immediately before their scheduled IFX infusion. Blood

was collected into a single Vacutainer Lithium Heparin collection tube (BD Bioscience).

For NGS analysis, blood was collected from a male donor treated with IFX, before IFX

administration and 10 days post administration. 30ml of peripheral blood were collected into 3

single Vacutainer K-EDTA collection tubes (BD Biosciences). Collection of peripheral blood

mono-nuclear cells (PBMCs) was performed by density gradient centrifugation, using Uni-

SepMAXI+ lymphocyte separation tubes (Novamed) according to the manufacturer's protocol.

Fluorescence-Activated Cell Sorting Analysis and sorting of B cell populations

PBMCs were stained for 15 minutes in cell staining buffer (BioLegend) at RT in the dark using the

following antibodies: anti-CD3–PerCP (clone OKT3; BioLegend), anti-CD19– Brilliant Violet 510

(clone HIB19; BioLegend), anti-CD27–APC (clone O323; BioLegend), anti-CD38–APC-Cy7

(clone HB-7; BioLegend), and anti-CD20–FITC (clone 2H7; BioLegend).

The following B cell populations were sorted using a FACSAria cell sorter (BD Bioscience):

CD3−CD19+CD20-CD27++CD38^{high} PB, CD3−CD19+CD20- CD27++CD38^{high}IFX-F(ab')$_2$+

IFX-F(ab')$_2$ positive PB, CD3−CD19+CD20+CD27^{var} mBC, and CD3−CD19+CD20+CD27^{var}IFX-

F(ab')$_2$+ IFX-F(ab')$_2$ positive mBC.

B cell subpopulations were sorted and collected into TRI Reagent solution (Sigma Aldrich) and

frozen at −80°C.

Amplification of V$_H$ and V$_L$ repertoires from B cells

Total RNA was isolated using RNeasy micro Kit (Qiagen), according to manufacturer's protocol.

First-strand cDNA generation was performed with 100ng of isolated total RNA using a SuperScript

RT II kit (Invitrogen) and oligo-dT primer, according to manufacturer's protocol. After cDNA

synthesis, PCR amplification was performed to amplify the V$_H$ and V$_L$ genes using a primer set

described previously (*32*) with overhang nucleotides to facilitate Illumina adaptor addition during

the second PCR (Table S3). PCR reactions were carried out using FastStart™ High Fidelity PCR

System (Roche) with the following cycling conditions: 95°C denaturation for 3 min; 95°C for 30

sec, 50°C for 30 sec, and 68°C for 1 min for four cycles; 95°C for 30 sec, 55°C for 30 sec, and 68°C

for 1 min for four cycles; 95°C for 30 sec, 63°C for 30 sec, and 68°C for 1 min for 20 cycles; and

a final extension at 68°C for 7 min. PCR products were purified using AMPure XP beads (Beckman

Coulter), according to manufacturer's protocol (ratio x 1.8 in favor of the beads). Recovered DNA

products from the first PCR was applied to a second PCR amplification to attach Illumina adaptors

to the amplified V$_H$ and V$_L$ genes using the primer extension method as described previously (*38*).

PCR reactions were carried out using FastStart™ High Fidelity PCR System (Roche) with the

following cycling conditions: 95°C denaturation for 3 min; 95°C for 30 sec, 40°C for 30 sec, and

68°C for 1 min for two cycles; 95°C for 30 sec, 65°C for 30 sec, and 68°C for 1 min for 7 cycles;

and a final extension at 68°C for 7 min. PCR products were applied to 1% agarose DNA gel

electrophoresis and gel-purified with Zymoclean™ Gel DNA Recovery Kit (Zymo Research)

according to the manufacturer's instructions. V_H and V_L libraries concentration were measured

using Qubit system (Thermo Fisher Scientific) and library quality was assessed using the

Bioanalyzer 2100 system (Agilent) or the 4200 TapeStation system (Agilent). All V_H libraries were

produced in duplicates starting with RNA as the common source template. The V_L were produced

with one replicate.

V_H and V_L libraries from sorted B cell were subjected to NGS on the MiSeq platform with the

reagent kit V3 2x300 bp paired-end (Illumina), using an input concentration of 16pM with 5%

PhiX.

Raw fastq files were processed using our recently reported ASAP webserver (*34*). ASAP analysis

resulted in a unique, full-length V_H and V_L gene sequences database for each time point. The

resultant database was used as a reference database to search the LC-MS/MS spectra.

Proteomic Analysis of the Serum ADA to IFX

Total IgG from each time point (D0, D10) were purified from 7-10ml of serum by protein G

enrichment. Serum was diluted 2 fold and passed through a 5ml Protein G agarose column

(GeneScript). The diluted serum was recycled three times over the column, which was subsequently

washed with 10CV of PBS and eluted with 7CV of 100mM glycine·HCl (pH 2.7). A total of 35

fractions of 1ml were collected and immediately neutralized with 100µl of 1.5M Tris·HCl (pH 8.8).

All elution fractions were evaluated for their purity using 12% SDS–PAGE and 11 purified 1 ml

IgG fractions were combined and dialyzed overnight at 4°C against 4L of PBS (pH 7.4) using

SnakeSkin dialysis tubing with 1 kDa cutoff (Thermo Fisher Scientific).

Next, 9mg of total IgG were digested with 100µg of IdeS in the final volume of 2ml PBS for 5 hour

at 37^0C. IdeS inactivation was carried out by adding 0.1M of citric acid pH 3 and incubation for 1

minute at RT followed by the addition of PBS (pH 7.4) to neutralize the low pH. Total serum F(ab')$_2$

was then applied to a one ml custom made affinity column comprised of IFX-F(ab')$_2$ coupled to

NHS-activated agarose beads (Thermo Fisher Scientific). The purified serum F(ab')$_2$ were recycled

three times over the affinity column, which was subsequently washed with 5CV of PBS and eluted

with 15CV of 100mM glycine·HCl (pH 2.7) and collected into Maxymum Recovery Eppendorf

(Axygen Scientific). A total of 30x0.5ml elution fractions and 1x50ml flow-through were

immediately neutralized with 50 and 100µl (respectively) of 1.5M Tris·HCl (pH 8.8). The purified

antigen-specific F(ab')$_2$ were dialyzed overnight at 4°C against 4L of PBS (pH 7.4) using SnakeSkin

dialysis tubing with 10kDa cutoff (Thermo Fisher Scientific). Elution and flow-through fractions

were trypsin-digested, and resulting peptides were fractionated and sequenced by nanoflow LC-

electrospray MS/MS on an Orbitrap Velos Pro hybrid mass spectrometer (Thermo Scientific), in

the UT Austin mass spectrometry core facility as described previously (*46*). MS/MS raw files were
analyzed by MaxQuant software version 1.6.0.16 (*58*) using the MaxLFQ algorithm (*59*) and
peptide lists were searched against the common contaminants database by the Andromeda search
engine (*60*) and a custom protein sequence database consisting of the donor-specific V_H and V_L
sequences derived from NGS of individual donor B cells. All searches were carried out with
cysteine carbamidomethylation as a fixed modification and methionine oxidations as variable
modifications. The false discovery rate was set to 0.01 for peptides with a minimum length of seven
amino acids and was determined by searching a reverse decoy database. Enzyme specificity was
set as C-terminal to arginine and lysine as expected using trypsin as protease, and a maximum of
two missed cleavages were allowed in the database search. Peptide identification was performed
with an allowed initial precursor mass deviation up to 7ppm and an allowed fragment mass
deviation of 20ppm. For LFQ quantification the minimal ratio count was set to 2, and match
between runs was performed with three mass-spec injections originating from the same sample.
MaxQunat output analysis file, "peptides.txt", was used for further processing. Total peptides that
were identified in the elution samples were filtered using the following criteria: i) were not
identified as contaminates; ii) did not match to the reversed decoy database; iii) were identified as
peptides derived from the region comprising the CDRH3, J region, FR4 and the ASTK motif
(derived from the N-terminal of the C_H1 region). The CDRH3 derived peptides were further
characterized as informative CDRH3 peptides (*i*CDRH3 peptides) only if they map exclusively to
a single antibody clonotype. A clonotype was defined as all sequences that comprise CDRH3 with
the same length and identity tolerating one amino acid mismatch, and same V, J family. The
intensities of high confidence *i*CDRH3 peptides were averaged between replicates while including
only peptides that were observed in at least 2 out of the 3 replicates. Clonotype frequencies within
each sample were calculated using only *i*CDRH3 peptides and were determined to be antigen-
specific if their frequency in the elution fraction was at least 5 fold greater than their frequency in
the flow-through fraction. The CDRH3 sequences identified by the mapping of high confidence
MS/MS peptides were used to generate a complete list of full length VH sequences. These VH
sequences were used to analyze the repertoire measures of the antibodies that were identified in the
donors' serum.
811
812 **Statistical analysis**
813 All curves were fitted on a sigmoidal dose–response curve and EC50 of each was calculated.
814 Mann-Whitney test was used to compare continuous variables. All reported P values were two-

815   tailed, and a P value less than 0.05 were considered statistically significant. All statistics were

816   performed with GraphPad Prism software (version 7, San Diego, California).

818
819

Acknowledgments

General: We are grateful to George Georgiou for assisting in the LC-MS/MS measurements at UT Austin, for Ulrich von Pawel-Rammingen from the Department of Molecular Biology, Umea University who kindly donated the plasmid with the gene encoding the IdeS.

Funding: The work was partially supported by BSF grant 2017359 (Y.W.)

Author contributions: A.V.M, S.B.H, I.B. and Y.W. conceived the research. A.V.M. S.R. and Y.W. designed the experiments, A.V.M., S.R., M.Y., E.F. and Y.D. performed experiments, A.V.M., S.R., S.B.H, M.Y., E.F., B.U. and U.K. collected and processed clinical samples, A.V.M, A.K. and Y.W. carried out data analysis, A.V.M. and Y.W. wrote the manuscript.

Competing interests: The authors declare no competing financial interests

References

1. A. L. Nelson, E. Dhimolea, J. M. Reichert, Development trends for human monoclonal antibody therapeutics. *Nat Rev Drug Discov* **9**, 767-774 (2010).
2. A. A. Shukla, L. S. Wolfe, S. S. Mostafa, C. Norman, Evolving trends in mAb production processes. *Bioeng Transl Med* **2**, 58-69 (2017).
3. C. Parola, D. Neumeier, S. T. Reddy, Integrating high-throughput screening and sequencing for monoclonal antibody discovery and engineering. *Immunology* **153**, 31-41 (2018).
4. D. M. Ecker, S. D. Jones, H. L. Levine, The therapeutic monoclonal antibody market. *MAbs* **7**, 9-14 (2015).
5. G. Kohler, C. Milstein, Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* **256**, 495-497 (1975).
6. P. A. van Schouwenburg, T. Rispens, G. J. Wolbink, Immunogenicity of anti-TNF biologic therapies for rheumatoid arthritis. *Nat Rev Rheumatol* **9**, 164-172 (2013).
7. A. S. De Groot, D. W. Scott, Immunogenicity of protein therapeutics. *Trends in immunology* **28**, 482-490 (2007).
8. T. T. Hansel, H. Kropshofer, T. Singer, J. A. Mitchell, A. J. George, The safety and side effects of monoclonal antibodies. *Nat Rev Drug Discov* **9**, 325-338 (2010).
9. K. Bloem, B. Hernandez-Breijo, A. Martinez-Feito, T. Rispens, Immunogenicity of Therapeutic Antibodies: Monitoring Antidrug Antibodies in a Clinical Context. *Ther Drug Monit* **39**, 327-332 (2017).
10. M. Krishna, S. G. Nadler, Immunogenicity to Biotherapeutics - The Role of Anti-drug Immune Complexes. *Frontiers in Immunology* **7**, (2016).
11. S. Ben-Horin *et al.*, The immunogenicity of biosimilar infliximab: can we extrapolate the data across indications? *Expert Rev Gastroenterol Hepatol* **9 Suppl 1**, 27-34 (2015).
12. K. Bendtzen, Immunogenicity of Anti-TNF-α Biotherapies: II. Clinical Relevance of Methods Used for Anti-Drug Antibody Detection. *Frontiers in Immunology* **6**, 109 (2015).

13. W. S. Putnam, S. Prabhu, Y. Zheng, M. Subramanyam, Y.-M. C. Wang, Pharmacokinetic,
pharmacodynamic and immunogenicity comparability assessment strategies for monoclonal
antibodies. *Trends in biotechnology* **28**, 509-516 (2010).
14. P. Rutgeerts *et al.*, Infliximab for Induction and Maintenance Therapy for Ulcerative Colitis.
dx.doi.org, (2009).
15. B. E. Sands *et al.*, Infliximab Maintenance Therapy for Fistulizing Crohn's Disease.
New England Journal of Medicine **350**, 876-885 (2004).
16. T. Horiuchi, H. Mitoma, S. Harashima, H. Tsukamoto, T. Shimoda, Transmembrane TNF-
alpha: structure, function and interaction with anti-TNF agents. *Rheumatology* **49**, 1215-
1228 (2010).
17. S. Ben-Horin, N. Vande Casteele, S. Schreiber, P. L. Lakatos, Biosimilars in Inflammatory
Bowel Disease: Facts and Fears of Extrapolation. *Clin Gastroenterol Hepatol* **14**, 1685-
1696 (2016).
18. M. P. Baker, H. M. Reynolds, B. Lumicisi, C. J. Bryson, Immunogenicity of protein
therapeutics: The key causes, consequences and challenges. *Self Nonself* **1**, 314-322 (2010).
19. T. R. D. J. Radstake *et al.*, Formation of antibodies against infliximab and adalimumab
strongly correlates with functional drug levels and clinical responses in rheumatoid arthritis.
Annals of the Rheumatic Diseases **68**, 1739-1745 (2009).
20. B. Ungar *et al.*, The temporal evolution of antidrug antibodies in patients with inflammatory
bowel disease treated with infliximab. *Gut* **63**, 1258-1264 (2014).
21. F. B. Vincent *et al.*, Antidrug antibodies (ADAb) to tumour necrosis factor (TNF)-specific
neutralising agents in chronic inflammatory diseases: a real issue, a clinical perspective.
Ann Rheum Dis **72**, 165-178 (2013).
22. S. S. Thomas *et al.*, Comparative Immunogenicity of TNF Inhibitors: Impact on Clinical
Efficacy and Tolerability in the Management of Autoimmune Diseases. A Systematic
Review and Meta-Analysis. *BioDrugs* **29**, 241-258 (2015).
23. S. Tatarewicz, J. M. Miller, S. J. Swanson, M. S. Moxness, Rheumatoid factor interference
in immunogenicity assays for human monoclonal antibody therapeutics. *J Immunol*
Methods **357**, 10-16 (2010).
24. K. Bloem *et al.*, Systematic comparison of drug-tolerant assays for anti-drug antibodies in
a cohort of adalimumab-treated rheumatoid arthritis patients. *J Immunol Methods* **418**, 29-
38 (2015).
25. S. Ben-Horin, Y. Chowers, Tailoring anti-TNF therapy in IBD: drug levels and disease
activity. *Nature reviews. Gastroenterology & hepatology* **11**, 243-255 (2014).
26. S. Ben-Horin *et al.*, The immunogenic part of infliximab is the F(ab′)2, but measuring
antibodies to the intact infliximab molecule is more clinically useful. *Gut* **60**,
gut.2009.201533-201548 (2010).
27. J. Goncalves *et al.*, Antigenic response to CT-P13 and infliximab originator in inflammatory
bowel disease patients shows similar epitope recognition. *Alimentary pharmacology &*
therapeutics **48**, 507-522 (2018).
28. M. Svenson, P. Geborek, T. Saxne, K. Bendtzen, Monitoring patients treated with anti-TNF-
α biopharmaceuticals: assessing serum infliximab and anti-infliximab antibodies.
Rheumatology (Oxford, England) **46**, 1828-1834 (2007).
29. U. von Pawel-Rammingen, B. P. Johansson, L. Björck, IdeS, a novel streptococcal cysteine
proteinase with unique specificity for immunoglobulin G. *The EMBO Journal* **21**, 1607-
1615 (2002).
30. U. Kopylov *et al.*, Clinical utility of antihuman lambda chain-based enzyme-linked
immunosorbent assay (ELISA) versus double antigen ELISA for the detection of anti-
infliximab antibodies. *Inflammatory bowel diseases* **18**, 1628-1633 (2012).

31. J. Lee *et al.*, Molecular-level analysis of the serum antibody repertoire in young adults
before and after seasonal influenza vaccination. *Nature Medicine* **22**, 1456-+ (2016).
32. J. J. Lavinder *et al.*, Identification and characterization of the constituent human serum
antibodies elicited by vaccination. *Proceedings of the National Academy of Sciences of the*
United States of America **111**, 2259-2264 (2014).
33. C. Giesecke *et al.*, Tissue Distribution and Dependence of Responsiveness of Human
Antigen-Specific Memory B Cells. *The Journal of Immunology* **192**, 3091-3100 (2014).
34. O. Avram *et al.*, ASAP, A Webserver for Immunoglobulin-Sequencing Analysis Pipeline.
Frontiers in Immunology **9**, (2018).
35. E. Miho *et al.*, Computational strategies for dissecting the high-dimensional complexity of
adaptive immune repertoires. *Frontiers in Immunology* **9**, (2018).
36. V. Greiff *et al.*, Quantitative assessment of the robustness of next-generation sequencing of
antibody variable gene repertoires from immunized mice. *BMC immunology* **15**, 40 (2014).
37. G. Blanchard-Rohner, A. S. Pulickal, C. M. J.-v. der Zijde, M. D. Snape, A. J. Pollard,
Appearance of peripheral blood plasma cells and memory B cells in a primary and
secondary immune response in humans. *Blood* **114**, 4998-5002 (2009).
38. U. Menzel *et al.*, Comprehensive Evaluation and Optimization of Amplicon Library
Preparation Methods for High-Throughput Antibody Sequencing. *PLOS ONE* **9**, (2014).
39. E. S. Mroczek *et al.*, Differences in the composition of the human antibody repertoire by B
cell subsets in the blood. *Frontiers in immunology* **5**, 96-96 (2014).
40. J. M. Volpe, T. B. Kepler, Large-scale analysis of human heavy chain V(D)J recombination
patterns. *Immunome research* **4**, 3-3 (2008).
41. Y. Wine *et al.*, Molecular deconvolution of the monoclonal antibodies that comprise the
polyclonal serum response. *Proc Natl Acad Sci USA* **110**, 2993-2998 (2013).
42. A. Ademokun *et al.*, Vaccination-induced changes in human B-cell repertoire and
pneumococcal IgM and IgA antibody at different ages. *Aging cell* **10**, 922-930 (2011).
43. J. D. Galson *et al.*, Analysis of B Cell Repertoire Dynamics Following Hepatitis B
Vaccination in Humans, and Enrichment of Vaccine-specific Antibody Sequences.
EBioMedicine **2**, 2070-2079 (2015).
44. B. J. DeKosky *et al.*, Large-scale sequence and structural comparisons of human naive and
antigen-experienced antibody repertoires. *Proc Natl Acad Sci U S A* **113**, E2636-2645
(2016).
45. C. M. Tipton *et al.*, Diversity, cellular origin and autoreactivity of antibody-secreting cell
population expansions in acute systemic lupus erythematosus. *Nature Immunology* **16**, 755
(2015).
46. D. R. Boutz *et al.*, Proteomic Identification of Monoclonal Antibodies from Serum.
Analytical Chemistry **86**, 4758-4766 (2014).
47. K. Murphy, P. Travers, M. Walport, C. Janeway, *Janeway's immunobiology*. (Garland
Science, New York, ed. 7th, 2008), pp. xxi, 887 p.
48. H. Nakase *et al.*, Significance of measurement of serum trough level and anti-drug antibody
of adalimumab as personalised pharmacokinetics in patients with Crohn's disease: a
subanalysis of the DIAMOND trial. *Alimentary pharmacology & therapeutics* **46**, 873-882
(2017).
49. E. A. Maser, R. Villela, M. S. Silverberg, G. R. Greenberg, Association of trough serum
infliximab to clinical outcome after scheduled maintenance treatment for Crohn's disease.
Clin Gastroenterol Hepatol **4**, 1248-1254 (2006).
50. K. Karmiris *et al.*, Influence of trough serum levels and immunogenicity on long-term
outcome of adalimumab therapy in Crohn's disease. *Gastroenterology* **137**, 1628-1640
(2009).

51. S. B. Hanauer *et al.*, Incidence and importance of antibody responses to infliximab after
maintenance or episodic treatment in Crohn's disease. *Clin Gastroenterol Hepatol* **2**, 542-
553 (2004).
52. C. H. Seow *et al.*, Trough serum infliximab: a predictive factor of clinical outcome for
infliximab treatment in acute ulcerative colitis. *Gut* **59**, 49-54 (2010).
53. J. E. Robinson *et al.*, Most neutralizing human monoclonal antibodies target novel epitopes
requiring both Lassa virus glycoprotein subunits. *Nature communications* **7**, 11544-11544
(2016).
54. A. N. Davydov *et al.*, Comparative Analysis of B-Cell Receptor Repertoires Induced by
Live Yellow Fever Vaccine in Young and Middle-Age Donors. *Front Immunol* **9**, 2309
(2018).
55. M. Pasparakis, L. Alexopoulou, V. Episkopou, G. Kollias, Immune and inflammatory
responses in TNF alpha-deficient mice: a critical requirement for TNF alpha in the
formation of primary B cell follicles, follicular dendritic cell networks and germinal centers,
and in the maturation of the humoral immune response. *Journal of Experimental Medicine*
184, 1397-1411 (1996).
56. C. A. Marin, P. G. Gisbert, M. Chaparro, Immunogenicity and mechanisms impairing the
response to vaccines in inflammatory bowel disease. *World Journal of Gastroenterology*
21, 11273-11281 (2015).
57. K. Wenig *et al.*, Structure of the streptococcal endopeptidase IdeS, a cysteine proteinase
with strict specificity for IgG. *Proceedings of the National Academy of Sciences of the*
United States of America **101**, 17371-17376 (2004).
58. J. Cox, M. Mann, MaxQuant enables high peptide identification rates, individualized p.p.b.-
range mass accuracies and proteome-wide protein quantification. *Nature biotechnology* **26**,
1367-1372 (2008).
59. J. Cox *et al.*, Accurate proteome-wide label-free quantification by delayed normalization
and maximal peptide ratio extraction, termed MaxLFQ. *Molecular & cellular proteomics :*
MCP **13**, 2513-2526 (2014).
60. J. Cox *et al.*, Andromeda: a peptide search engine integrated into the MaxQuant
environment. *Journal of proteome research* **10**, 1794-1805 (2011).
998

Fig. 1: IFX digestion and IFX-F(ab')$_2$ purification. (A) Schematic representation of IgG digestion with IdeS. IdeS is a highly specific immunoglobulin-degrading enzyme that cleaves below the disulfide bonds in the IgG hinge region. The cleavage results in the production of one IFX-F(ab')$_2$ fragment and two ½ Fc fragments. (B) SDS-PAGE analysis of intact IgG (lane 2), following IdeS digestion (lane 3) and purified IFX-F(ab')$_2$ following a 2-step affinity chromatography purification including protein A and kappa-select columns (lane 4). (C) Presence of Fc and intact IgG traces was measured by direct ELISA were intact IgG and purified IFX-F(ab')$_2$ were compared to a control antigen (streptavidin) as coating agents followed by direct incubation with an anti-Fc HRP conjugate at the detection phase. (D) The functionality of the recovered IFX-F(ab')$_2$ was confirmed by ELISA and compared to intact IFX. The ELISA setup included TNFα as the coating agent and anti-κ HRP conjugate at the detection phase. For panel C–D, triplicate averages were calculated as mean, with error bars indicating s.d.

Fig. 2: AHLC and the newly developed mAb-F(ab')₂ based bio-immunoassay configuration and their application on serum samples from patients treated with IFX. (A) AHLC assay configuration. AHLC

assay is based on an ELISA where TNFα is used as the coating agent, incubation with the mAb drug followed by differential dilutions of tested sera. anti-λ HRP conjugate is used at the detection phase. (B) The newly developed mAb-F(ab')$_2$ based bio-immunoassay configuration. The assay is based on an ELISA where mAb-F(ab')$_2$ is used as the coating agent followed by differential dilutions of tested sera. anti-Fc HRP conjugate is used at the detection phase. (C) Graph representing the ELISA results obtained utilizing the AHLC assay on two serum sample. This assay enabled the stratification of the tested sera to those that show detectable levels of ADA (AHLC[(+)]) and those with no detectable levels of ADA (AHLC[(-)]). (D) Both serum samples were applied to the newly developed mAb-F(ab')$_2$ based bio-immunoassay. This assay was able to detect ADA in both sera. For C–D, averages were calculated as mean, with error bars indicating s.d.

Fig. 3: Configuration of the assay for determining the neutralization index in patient sera and ELISA results obtained from patient sera. (A) The newly developed mAb-F(ab')$_2$ based bio-immunoassay configuration (left) and the modified configuration where mAb-F(ab')$_2$ binding site is blocked by saturating the assay with 200nM of rhTNFα (right). (B) Graph representing the ELISA results obtained utilizing the neutralization assay on serum sample that show detectable levels of ADA (AHLC$^{(+)}$) and (C) serum sample with no detectable levels of ADA (AHLC$^{(-)}$). (D) Scatter plot consolidating the neutralization index obtained by applying the immunoassay on sera from 55 patients treated with IFX (***P = 0.0003, Mann-Whitney *U* test). For B–C, averages were calculated as mean, with error bars indicating s.d.

Fig. 4: Changes in B cell repertoires following IFX infusion. mBC and PB from a patient treated with IFX were collected at two time points (D0, D10) and processed for NGS analysis. (A) The V family usage showed no difference between D0 and D10, different B cell subsets and isotypes. (B) An increase in antibody CDRH3 length and (C) a decrease in the number of bp mutations was observed at D10 in both PB and mBC.

(B) (For PB:****P<0.0001, For mBC: ****P<0.0001, Mann-Whitney U test)

(C) (For PB:****P<0.0001, For mBC: ****P<0.0001, Mann-Whitney U test)

Fig. 5: V gene and circulating antibody repertoire characteristics. (A) The V(D)J family usage of V gene sequences that were identified by LC-MS/MS. (B) Mapping of V gene sequences to B cell subsets and (C) isotypes, based on NGS data.