Wheat *VRN1* and *FUL2* play critical and redundant roles in spikelet meristem identity and

spike determinacy

Chengxia Li[1, 2*], Huiqiong Lin[1, 2*], Andrew Chen[1, 3], Meiyee Lau[1], Judy Jernstedt[1] and Jorge

Dubcovsky[1, 2‡]

[*] These authors contributed equally to this work

[1]Department of Plant Sciences, University of California, Davis, CA 95616, USA.

[2]Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA.

[3]Current address: University of Queensland, Brisbane, QLD4072, Australia

[‡]Author for correspondence: jdubcovsky@ucdavis.edu. Phone: 530 752 5159

Running title: Spikelet meristem identity genes

Key words: wheat, spike development, spikelet, meristem identity, MADS-box,

Word count: 6190

SUMMARY STATEMENT

The wheat MADS-box proteins VRN1 and FUL2 play critical and overlapping roles in the

development of spikelets, which are the basic unit of all grass inflorescences.

ABSTRACT

The spikelet is the basic unit of the grass inflorescence. In this study, we show that wheat

MADS-box genes *VRN1* and *FUL2* play critical and redundant roles in the determination of

spikelet meristem identity. Combined loss-of-function mutations of these two genes (*vrn1ful2*-

null) were sufficient to revert lateral spikelet meristems into vegetative meristems in the spikes

of tetraploid wheat. These plants were also unable to form a terminal spikelet and had an

indeterminate inflorescence. The single *vrn1*-null and *ful2*-null mutants showed increased

number of spikelets per spike, likely due to a delayed formation of the terminal spikelet.

Mutations in these two genes and in the more distantly related paralog *FUL3*, also affected wheat

heading time and stem elongation. The *ful2*-null mutant also showed a higher number of florets

per spikelet, which together with a higher number of spikelets, resulted in a significant increase

in the number of grains per spike in the field. Our results suggest that a better understanding of

the basic mechanisms underlying wheat spike development can inform future strategies to

improve grain yield in wheat.

INTRODUCTION

The grass family (Poaceae) has approximately 10,000 species, including important food crops such as rice, maize, sorghum, barley, and wheat (Kellogg, 2001). The flowers of these species are organized in a unique and diagnostic structure called spikelet (literally "little spike"), which is a compact and indeterminate branching inflorescence developing within the larger inflorescence (Malcomber et al., 2006). A spikelet typically has two sterile bracts (called glumes) enclosing one or more florets. Each floret includes a carpel, 3 or 6 stamens and two modified petals (called lodicules) subtended by two bract-like organs, the palea and the lemma (Preston et al., 2009).

The spikelet adds a recursive level of complexity to grass inflorescences, in which different types of meristems are responsible for a large diversity of inflorescence architectures (Ciaffi et al., 2011). Development of the ancestral grass inflorescence, the panicle, begins when the shoot apical meristem (SAM) changes from a vegetative meristem (VM) that produces leaves to an inflorescence meristem (IM), which elongates and generates lateral primary branch meristems (PBMs). The PBMs generate secondary branches as axillary meristems (SBM), and both PBMs and SBMs terminate into spikelet meristems (SMs), resulting in a highly branched structure. The SMs then generate glumes and lateral floral meristems (FMs), which vary in number across grass species (Malcomber et al., 2006).

In wheat, a drastic shortening of the lateral branches results in sessile spikelets attached directly to the central axis or rachis and the formation of a derived inflorescence, a spike, in which the spikelets are arranged alternately in opposite vertical rows (a distichous pattern) (Kellogg et al., 2013). In the initial stage of wheat spike development, the IM generates a double-ridge structure in which the lower ridges arrest, while the upper ridges acquire SM identity and form spikelets. The number of spikelets per spike is determined by the number of lateral meristems formed before the transition of the IM into a SM to form the terminal spikelet. Although the growth of the wheat spike is determinate, the growth of each spikelet is indeterminate, with each SM initiating a variable number of FMs (Ciaffi et al., 2011). The numbers of spikelets per spike and florets per spikelet determine the maximum number of grains per spike and are, therefore, important components of wheat grain yield potential.

Studies in Arabidopsis, which has a simpler inflorescence than grasses (Malcomber et al., 2006),

have shown that MIKC-type MADS-box transcription factors *APETALA1* (*AP1*),

CAULIFLOWER (*CAL*) and *FRUITFULL* (*FUL*) are prominent players in the determination of

floral meristem identity. In the triple *ap1calful* mutant, the IM is not able to produce flowers and

reiterates the development of leafy shoots (Ferrándiz et al., 2000). MIKC-type MADS-box

proteins have a highly conserved MADS DNA-binding domain (also important for dimerization

and nuclear localization), an Intervening (I) domain, a Keratin-like (K) domain critical for

dimerization and multimeric complex formation, and a C-terminal domain involved in

transcriptional activation. These proteins bind as dimers to DNA sequences named 'CArG'

boxes, and can organize in tetrameric complexes that can recognize different CArG boxes. The

multimeric nature of these complexes generates a large number of combinatorial possibilities

with different targets and functions (Honma and Goto, 2001; Theissen et al., 2016).

The closest homologs to the Arabidopsis MADS-box genes *AP1*, *CAL* and *FUL* in the grass

lineage are *VERNALIZATION 1* (*VRN1*), *FUL2* and *FUL3*. A phylogenetic analysis of the

proteins encoded by these genes indicates that these Arabidopsis and grass groups have

independent subfunctionalization stories (Preston and Kellogg, 2006) and Fig. S1). In the grass

lineage, a duplication that occurred close to the divergence between monocots and eudicots

resulted in the separation of the *FUL3* clade. A later duplication that occurred near the base of the

radiation of the grasses resulted in the *VRN1* and *FUL2* clades (Preston and Kellogg, 2006).

Large truncation mutations in the two *VRN1* homeologs in tetraploid wheat delay heading time,

but do not alter spikelet morphology or the ability of flowers to form viable grains (Chen and

Dubcovsky, 2012). Since *FUL2* and *FUL3* are the closest paralogs of *VRN1*, we hypothesized

that they could have redundant spikelet and floral meristem identity functions.

In this study, we generated truncation mutants for the two homeologs of both *FUL2* and *FUL3* in

the same tetraploid background carrying the *vrn1* truncation mutations and intercrossed them to

generate double- and triple-null mutants. Characterization of these mutants revealed that *VRN1*,

FUL2 and *FUL3* have overlapping roles in the regulation of flowering time and stem elongation

and, more importantly, that *VRN1* and *FUL2* (but not *FUL3*) are critical and redundant in the

determination of spikelet meristem identity and spike determinacy. Individual *vrn1* and *ful2*

truncation mutants showed significant increases in the number of spikelets and grains per spike,

suggesting that manipulations of these genes may contribute to increasing wheat grain yield

potential.

RESULTS

Mutations of *FUL2* and *FUL3* affect stem elongation

We identified loss-of-function mutations in the A and B genome homeologs of *FUL2* and *FUL3*

(Fig. S2) in an EMS-mutagenized population of the tetraploid spring wheat variety Kronos

(Krasileva et al., 2017; Uauy et al., 2009) and backcrossed each individual mutant two to three

times to Kronos to reduce background mutations. Then, we intercrossed the A and B genome

homeologs for each gene and selected plants homozygous for both mutations. For simplicity,

mutants with large truncation mutations in both homeologs will hereafter be referred to as null

mutants (e.g. *vrn1*-null). The *ful2*-null and *ful3*-null mutants were crossed with *vrn1*-null (Chen

and Dubcovsky, 2012) to generate *vrn1ful2*-null and *vrn1ful3*-null mutants, which were

intercrossed to generate the *vrn1ful2ful3*-null mutant (Fig. S2). The wild type Kronos carries a

functional *VERNALIZATION 2* (*VRN2*) flowering repressor, which results in extremely late

flowering in the presence of the *vrn1*-null mutation (Chen and Dubcovsky, 2012). To avoid this

problem all mutants described in this study were developed in a Kronos *vrn2*-null background

(Distelfeld et al., 2009), unless indicated otherwise. This line is referred to in figures as the

control line.

Since some mutant combinations lack real spikes, we decided to determine final stem length

from the base of the plant to the base of the spike instead of total plant height. The individual

ful3-null and *vrn1*-null mutants showed marginal or not significant reductions in stem length (5

% and 3 %, respectively, Fig. S3A). However, a factorial experiment including all four *VRN1* by

FUL3 wild type and null allele combinations, revealed highly significant effects for both genes

($P = 0.002$) and a significant synergistic interaction ($P = 0.019$) that resulted in a 17 % reduction

in stem length in the *vrn1ful3*-null mutant (Fig. S3B). The stems of plants carrying only the *ful2*-

null mutation were 27% shorter than the control ($P < 0.0001$, Fig. S3C) and the differences were

magnified in the double-null (*vrn1ful2*-null, 40% reduction, $P < 0.0001$, Fig. S3D) and triple-

null mutants (*vrn1ful2ful3*-null, 63% reduction, $P < 0.0001$, Fig. S3E). Taken together, these

results suggest that *VRN1*, *FUL2* and *FUL3* have redundant roles in the regulation of stem

elongation, and that the effect of the individual genes is larger in the absence of the other

paralogs.

FUL2 and FUL3 affect flowering and heading time

Functional redundancy among *VRN1*, *FUL2* and *FUL3* was also observed for heading time, but

the effect of the latter two genes was detected only in the absence of *VRN1*. The *vrn1*-null mutant

headed 37.5 d later than the control (Fig. 1A), but the differences in heading time for the *ful2*-

null (Fig. 1B), *ful3*-null and *ful2ful3*-null mutants in the presence of the strong *Vrn-A1* allele

were non-significant (Fig. 1C). For the *vrn1ful2*-null and *vrn1ful2ful3*-null mutants, it was not

possible to determine heading times accurately because they have short stems and abnormal

spikes that interfere with normal ear emergence. To study the effect of these mutations on

flowering time we determined instead the timing of the transition between vegetative and

double-ridge stages (Fig. 1D). The SAM in *vrn1*-null and *vrn1vrn3*-null transitioned to the

double-ridge stage around 31 days, but this transition was delayed 3-6 d in *vrn1ful2*-null and 9 to

12 d in *vrn1ful2ful3*-null (Fig. 1D). The latter result indicates that *FUL3* has a residual ability to

accelerate flowering time in the absence of *VRN1* and *FUL2*.

Transgenic Kronos plants overexpressing the coding regions of *FUL2* fused with a C-terminal

3xHA tag (henceforth *Ubi::FUL2*, Fig. 1E, events #1 and #6) or *FUL3* fused with a C-terminal

4xMYC tag (henceforth *Ubi::FUL3*, Fig. 1F, events #4 and #5) headed two to four days earlier

than the non-transgenic sister lines ($P < 0.0001$). The effect of *Ubi::FUL2* on heading time was

further characterized in the presence of different *VRN1* and *VRN2* alleles in the F_2 progeny from

the cross between *Ubi::FUL2* (*Vrn1Vrn2*) and *vrn1vrn2*-null under greenhouse conditions. Using

a three-way factorial ANOVA, we detected highly significant effects in heading time for each

gene and for all two- and three-way interactions ($P < 0.0001$, Table S3). As an example of these

interactions, the differences in heading time between the *FUL2*-wt and *Ubi::FUL2* alleles in the

presence of a functional *VRN2* allele were small in lines homozygous for the functional *VRN1*

allele (2.6 d, Fig. 1E), intermediate in *VRN1* heterozygous lines (11.1 d, Fig. 1G) and large in

homozygous *vrn1*-null mutants (53 d, Fig. 1H). These results indicate that the effect of the

Ubi::FUL2 transgene on heading time depends on the particular *VRN1* and *VRN2* alleles present

in the genetic background (Fig. G-H).

In summary, the strong effect of *VRN1* in the acceleration of wheat flowering time can mask the

smaller effects of *FUL2* and *FUL3*, but in the absence of *VRN1*, both *FUL2* and *FUL3* have

redundant effects on accelerating wheat flowering time.

FUL2 and VRN1 play critical but redundant roles in the determination of spikelet

meristem identity

The *vrn1*-null and *ful2*-null individual mutants showed normal spikelets and flowers (Fig. 2A-E),

but the plants carrying both mutations showed striking developmental changes. The *vrn1ful2*-null

mutants had a spike-like structure (Fig. 2A) in which all the lateral spikelets were replaced by

leafy shoots. These shoots produced normal leaves and most of them had no floral organs (Fig.

2F-G). However, dissection of tillers located in the basal part of this spike-like structure revealed

that some of them had leafy bracts subtending abnormal floral organs in the expected position of

the first floret (Fig. S4A-E). Floral abnormalities included leafy lodicules, reduced number of

anthers or anthers fused to ovaries and multiple ovaries (Fig. S4C-E). Dissection and Scanning

Electron-Microscopy (SEM) images of the *vrn1*-null and *vrn1ful2*-null IM showed lateral

meristems organized in a distichous phyllotaxis in both mutants (Fig. 2D and H). However,

while these lateral meristems progressed to SMs that formed normal glume flower primordia in

the *vrn1*-null and *vrn1ful3*-null mutants (Fig. 2D-E), they reverted to VMs generating leaf

primordia in the *vrn1ful2*-null mutant (Fig. 2H-I). After producing a variable number of leaves,

those lateral VMs transitioned again to IMs that generated their own lateral VMs (Fig. S4A-B).

The *vrn1ful2ful3*-null mutant showed a similar spike-like structure as the *vrn1ful2*-null mutant,

with vegetative meristems replacing the lateral spikelet meristems and vegetative shoots

replacing the spikelets (Fig. 2J-K). Compared with the *vrn1ful2*-null mutant, the stems in the

triple-null mutants were shorter and frequently had difficulty emerging from leaf sheaths,

resulting in smaller spike-like structures with curly leaves (Fig. 2A and J). Both *vrn1ful3*-null

and *ful2ful3*-null plants showed normal spikelets and fertile flowers. Since *FUL3* played a

limited role in the determination of the SM identity, we focused subsequent studies on the

vrn1ful2-null mutant.

To compare the relative roles of *VRN1* and *FUL2* on SM identity, we examined their individual

ability to restore fertile flowers when present as a single functional copy in a heterozygous state

(underlined). Both *ful2*-null/*Vrn-A1 vrn-B1-null* (functional *Vrn1* allele for spring growth habit)

and *ful2*-null/*vrn-A1-null* <u>*vrn-B1*</u> (functional *vrn1* allele for winter growth habit) produced spike-

like structures with leafy lateral shoots (Fig. 3A-B) and normal floral organs (Fig. 3C) but no

viable seeds. The developing spikes of these plants showed lateral meristems with floral

primordia (Fig. 3D), some of which later showed elongated rachillas and leafy organs (Fig. 3E-

G). By contrast, the presence of a single heterozygous copy of *FUL2* in a *vrn1*-null background

(*vrn1*-null/*ful2-A* <u>*Ful2-B*</u>) resulted in spikelets that were less leafy than the previous mutants

(Fig. 3H). Some of these spikelets were more similar to the control and were able to set some

viable seeds. Abnormal spikelets (Fig. 3I) and basal branches with lateral spikelets and fertile

florets (Fig. 3J) were also observed in this mutant. Taken together, these results indicate that

VRN1 and *FUL2* have critical but redundant roles in the determination and maintenance of SM

identity and that *FUL3* has a limited effect on this trait.

194 Transcript levels of *SHORT VEGETATIVE PHASE* (*SVP*)-like MADS-box genes *VRT2*,

195 *BM1* and *BM10* are upregulated in the developing spikes of the *vrn1ful2*-null mutant

A partial reversion of basal spikelets to vegetative tillers, similar to the one described above for

the *vrn1ful2*-null mutant, has been described in barley lines overexpressing *SVP*-MADS-box

genes *BM1* or *BM10* (see Discussion). To test if the transcript levels of the *SVP*-like wheat genes

were affected in the *vrn1ful2*-null mutants, we first studied their expression during normal spike

development in Kronos. Transcript levels of three related paralogs *BM1*, *BM10* and *VRT2*

(RefSeq v1.0 gene designations in Fig. S6) decreased three- to five-fold from the initial stages of

spike development (W2, Waddington scale) to the floret primordium stage (W3.5, Fig. S6A-C).

Then, we compared the transcriptional levels of the *SVP*-like wheat genes in *vrn1ful2*-null and

vrn1-null mutants. Plants were grown for 53 days in the same growth chamber until the

developing spikes of *vrn1*-null were at the terminal spikelet stage and those from *vrn1ful2*-null

had a similar number of lateral meristems (Fig. 2D and H). The transcript levels of wheat *BM1*,

BM10 and *VRT2* in the developing spikes were roughly ten-fold higher in the *vrn1ful2*-null

mutant than in the *vrn1*-null and control lines ($P < 0.0001$, Fig. S6D-F). These results suggest

that *VRN1* and *FUL2* are either direct or indirect transcriptional repressors of the three wheat

SVP-like genes.

***FUL2* and *VRN1* have redundant roles on spike determinacy and regulate the number of spikelets per spike**

Normal wheat spikes are determinate, with the distal IM transitioning into a terminal spikelet after producing a relatively stable number of lateral meristems (Fig. 4A). In *vrn1ful2*-null, by contrast, the spike-like structures are indeterminate, with a distal IM that continues to produce lateral meristems while growing conditions are favorable (Fig. 4B). In the *ful2*-null background, one functional copy of *VRN1* in the heterozygous state was sufficient to generate a determinate spike (Fig. 3D, *ful2*-null/*vrn-A1*-null <u>vrn-B1</u>), and the same was true for a single functional copy of *FUL2* in a *vrn1*-null background (Fig. 3K, *vrn1*-null/*ful2-A* <u>Ful2-B</u>).

The individual *vrn1*-null and *ful2*-null mutants (in a homozygous state) showed a larger number of spikelets per spike than the control. This increase was 58% in the *vrn1*-null mutant ($P < 0.0001$, Fig. 4C) and 10% in the *ful2*-null mutant ($P = 0.0014$, Fig 4D). Although no significant increases in the number of spikelets per spike were detected in the individual *ful3*-null mutant ($P = 0.4096$, Fig. 4E), two independent transgenic lines overexpressing *FUL3* (*Ubi::FUL3*) showed an average reduction of 1.12 spikelet per spike relative to their non-transgenic sister lines ($P = 0.0132$ and $P < 0.0001$, Fig. S7A). This last result indicates that *FUL3* can still play a role on the timing of the transition of the IM to a terminal spikelet.

A similar reduction in the number of spikelets per spike was observed in two independent *Ubi::FUL2* transgenic lines (1.05 spikelets per spike reduction, $P < 0.03$, Fig. S7B). We then investigated the effect of this transgene in the presence of different *VRN1* and *VRN2* alleles in the same F$_2$ population described previously (*Ubi::FUL2* x *vrn1vrn2*-null). In the *vrn2*-null F$_2$ plants, the differences in spikelet number between *Ubi::FUL2* and wild type alleles were larger in *vrn1*-null than in the *Vrn1*-Het plants (interaction $P < 0.0001$, Fig. S7C). In a separate group of F$_2$ plants fixed for *Vrn1*-Het and segregating for *VRN2* and *FUL2*, we did not detect significant effects for *Ubi::FUL2* and the interaction was not significant (Fig. S7D). However, we observed 3.3 more spikelets per spike in *Vrn2*-wt than in *vrn2*-null plants ($P < 0.0001$, Fig. S7D). These results suggest that the strong *Vrn-A1* allele for spring growth habit can mask the effects of the *Ubi::FUL2* transgene but not that of *VRN2* on the number of spikelets per spike.

Increased transcript levels of *CEN2*, *CEN4* and *CEN5* in developing spikes of the *vrn1ful2*-null mutant

Based on the strong effect observed in the Arabidopsis *tfl1* mutant and the *Antirrhinum cen*

mutant on inflorescence determinacy (see Discussion), we decided to investigate the effect of the

vrn1ful2-null mutations on the expression levels of the *TFL1/CEN*-like wheat homologs in the

developing spike. Since no previous nomenclature was available for the wheat *CEN* paralogs, we

assigned them numbers to match their chromosome locations, and designated them as *CEN2*,

CEN4 and *CEN5* (RefSeq v1.0 designations can be found in the legend of Fig. S8). The

transcript levels of these three genes were downregulated as the developing spike progressed

from the double-ridge stage to the floret primordium stage (Waddington scale 2 to 3.5, Fig. S8A-

C).

Comparison of *vrn1ful2*-null and *vrn1*-null plants grown for 53 days in the same growth chamber

(Fig. 2D and H) showed that the transcript levels of *CEN2*, *CEN4* and *CEN5* were significantly

higher ($P < 0.0001$) in the developing spikes of the *vrn1ful2*-null mutant than in those of the

vrn1-null mutant or the Kronos control (all in *vrn2*-null background). These differences were

larger for *CEN2* and *CEN4* than for *CEN5* (Fig. S8D-F). Taken together, these results suggested

that *VRN1* and *FUL2* work as transcriptional repressors of the *TFL1/CEN*-like wheat homologs.

The *ful2*-null mutant produces a higher number of florets per spikelet and more grains per

spike in the field

In addition to the higher number of spikelets per spike, the *ful2*-null mutant showed a higher

number of florets per spikelet than the Kronos control, an effect that was not observed for *vrn1*-

null (Fig. 2B-C) or *ful3*-null (Fig. S5A). The average increase in floret number was similar in

ful2-null (1.3 florets) and *ful2ful3*-null (0.9 florets), suggesting that *FUL3* has a limited effect on

this trait. In spite of some heterogeneity in the distribution of spikelets with extra florets among

spikes, the differences between the control and the *ful2*-null mutants were significant at all spike

positions (Fig. S5B).

Similar increases in the number of florets per spikelet were reported before in Kronos plants

overexpressing *miRNA172* (*Ubi::miR172*, (Debernardi et al., 2017). To study the genetic

interactions between *Ubi::miR172* and *ful2*-null we crossed the transgenic and mutant lines and

studied their effects on floret number in the progeny using a two-way factorial ANOVA. We

detected significant differences in average floret number for both *ful2*-null and *Ubi::miR172* (P

< 0.01) and a marginally significant interaction ($P < 0.0435$) that can be visualized in the

interaction graph in Fig. S5C. The differences in average floret number between *ful2*-null and the

wild type control were larger (and more variable) in the *Ubi::miR172* than in the non-transgenic

background (Fig. S5C-F). Both the mutant and transgenic lines showed heterogeneity among

spikes in the location of spikelets with increased numbers of florets (Fig. S5D-F).

Based on its positive effect on the number of florets per spikelet and spikelets per spike (and its

small effect on heading time), we selected the *ful2*-null mutant for evaluation in a replicated field

experiment. Relative to the control, the *ful2*-null mutant produced 20% more spikelets per spike

($P = 0.0002$) and 9% more grains per spikelet ($P= 0.05$), which resulted in a 31% increase in the

number of grains per spike ($P = 0.0002$, Fig. 4F). In this experiment, part of the positive effect

on grain yield was offset by a 19% reduction in average kernel weight ($P = 0.0012$). In spite of

these opposite trends, we observed a slight net increase of 6% in total grain weight per spike ($P = $

0.09, Fig. 4F). This negative correlation between grain number and grain weight suggest that in

this particular genotype by environment combination grain yield was more limited by the

"source" (produced and transported starch) than by the "sink" (number and size of grains).

DISCUSSION

RNA *in situ* hybridization studies at the early double-ridge stage of spike development in wheat

(Preston and Kellogg, 2008) and *Lolium temulentum* (Gocal et al., 2001) have shown that both

VRN1 and *FUL2* are expressed in the spike apical meristem and in spikelet primordia. Based on

their shared localization it was hypothesized that these two genes might have overlapping roles

in spike development. This study provides experimental evidence for this hypothesis and

demonstrates that *FUL2* and *VRN1* play central but redundant roles in the determination of

spikelet meristem identity and spike determinacy. It also shows that mutations in these two

genes, and in the more distantly related *FUL3* paralog, affect wheat heading time and stem

elongation.

Mutations in *VRN1*, *FUL2* and *FUL3* reduce stem elongation

No effects on stem elongation had been reported previously for *VRN1*, *FUL2* or *FUL3* in wheat

or *AP1*, *CAL* or *FUL* mutants in Arabidopsis, so we initially paid little attention to this trait.

However, differences in stem elongation in the double- and triple-null mutants were too obvious

to be ignored. The small effects of the single *vrn1*-null and *ful3*-null mutants were magnified in

the *vrn1ful3*-null mutant, indicating a synergistic interaction (Fig. S3B). A similar synergistic

interaction was detected when the *ful2*-null mutant (Fig. S3C) was combined with the *vrn1*-null

and *ful3*-null mutations (Fig. S3D-E). These results suggest that *VRN1*, *FUL2* and *FUL3* have

redundant functions in the regulation of stem elongation, and that their individual effects are

magnified in the absence of the other paralogs.

Although the molecular mechanisms by which these genes affect stem elongation are currently

unknown, an indirect way by which *VRN1* may contribute to this trait is through its strong effect

on the regulation of *FT1*. The induction of *FT1* has been shown to result in the upregulation of

gibberellic acid (GA) biosynthetic genes in developing wheat spikes (Pearce et al., 2013). GA

biosynthesis and sensitivity are known to play critical roles in the determination of stem

elongation in wheat (Peng et al., 1999). Therefore, the downregulation of *FT1* in the *vrn1*-null

mutant (Chen and Dubcovsky, 2012) is expected to reduce GA levels, which would inhibit stem

elongation.

A loss-of-function mutation in a homolog of *AP1* in rapeseed resulted in increased plant height

by 7.6 cm (Shah et al., 2018). This effect is opposite to what we observed in the loss-of-function

mutations in the wheat homologs, which suggests that different mechanisms are involved in the

effect of this group of meristem identity genes on plant height in these two species.

Mutations in *VRN1*, *FUL2* and *FUL3* delay flowering initiation in wheat

VRN1 is one of the main genes controlling natural variation in wheat flowering time (Fu et al.,

2005; Kippes et al., 2016; Yan et al., 2003; Zhang et al., 2008), so it was not surprising that *vrn1*-

null delayed heading time more than *ful2*-null or *ful3*-null. Although the strong *Vrn-A1* allele for

spring growth habit masked the smaller effects of *FUL2* and *FUL3* (Fig. 1A-C), in the absence of

VRN1 the *FUL2* and *FUL3* mutants showed delayed flowering initiation(Fig. 1D), indicating that

FUL2 and *FUL3* have retained some residual functionality in the acceleration of wheat flowering

time. This was further confirmed by the accelerated flowering of the *Ubi::FUL2* and *Ubi::FUL3*

transgenic plants (Fig. 1 E-F). Similar results have been reported in *Brachypodium distachyon,* in

which overexpression of *VRN1* (Ream et al., 2014)*, FUL2* or *FUL3* (Li et al., 2016) accelerates

flowering, and downregulation of *VRN1* delays flowering relative to non-transgenic controls

(Woods et al., 2016). These results suggest a conserved role of these genes in the regulation of

flowering time in temperate grasses.

Previous studies have shown a significant interaction between wheat *VRN1* and *VRN2* in the

regulation of heading time (Tranquilli and Dubcovsky, 2000). This study shows that similar

interactions exist between *FUL2* and *VRN2* (Fig. 1G-H). A tetraploid wheat population

segregating for *VRN1*, *FUL2* and *VRN2* revealed highly significant two-way and three-way

interactions among these genes, indicating that the effect of each of these genes on heading time

is dependent on the particular combination of alleles present for the other two. Previous studies

have shown that part of the ability of *VRN1* to accelerate flowering depends on its ability to

repress *VRN2* (Chen and Dubcovsky, 2012). The larger effect on heading time of the *Ubi::FUL2*

transgene in the presence of the functional *Vrn2* than in the *vrn2*-null background (Fig. 1G-H)

suggests that *FUL2* repression of *VRN2* can also contribute to its ability to accelerate heading

time.

VRN1 and FUL2 play critical and redundant roles in spikelet and floral development

The most significant discovery from this study was the central and redundant roles of *VRN1* and

FUL2 in the specification of the spikelet meristem identity. Even though the VM to IM transition

was not impaired in the *vrn1ful2*-null mutant, these plants were unable to form normal spikelets.

The *vrn1ful2*-null and *vrn1ful2ful3*-null mutants showed an elongated stem terminating in a

spike-like structure (Fig. 2A) with lateral meristems arranged in a distichous phyllotaxis, similar

to normal spikes (Fig. 2H, I, K and Fig. 4A-B). However, the fate of the lateral meristems was

drastically different in the wild type and mutant genotypes. In the wild type, the lateral

meristems acquired SM identity and formed normal spikelets with fertile flowers, whereas in the

vrn1ful2-null and *vrn1ful2ful3*-null mutants the lateral meristems reverted to VMs that formed

leaves. Interestingly, after forming a few leaves these VMs transitioned again into IMs with

lateral VMs, generating a recurrent developmental cycle that ended with the death of the plant.

No normal spikelets were observed in the double *vrn1ful2*-null mutant, but some of the tillers in

the basal region of the spike-like structure showed leafy bracts subtending abnormal floral

organs in the position of the first floret. This result indicates that *VRN1* and *FUL2* are not

absolutely essential for the production of floral organs. This was also observed in the

Arabidopsis *ap1calful* mutants, which usually failed to produce flowers but were able to produce

some abnormal floral organs under high temperatures (Ferrándiz et al., 2000). In Arabidopsis,

AP1 orchestrates a network of intermediate transcription factors that promote FM identity genes

and repress VM or IM identity genes (Kaufmann et al., 2010). The escape of these intermediate

transcription factors from AP1 control in old plants or under particular stress conditions

occasionally resulted in the formation of floral organs in the Arabidopsis *ap1calful* mutant.

Among the transcription factors directly repressed by AP1, the MADS-box *SVP* and

AGAMOUS-LIKE 24 (*AGL24*) play important roles in the formation of floral meristems

(Kaufmann et al., 2010; Liu et al., 2007). In the absence of AP1, ectopic expression of *SVP* and

AGL24 transform floral meristems into shoot meristems (Liu et al., 2007). The three related *SVP*-

like genes in wheat and barley lineages (*VRT2*, *BM1* and *BM10*) originated from duplications

independent of the one that originated *SVP* and *AGL24* in Arabidopsis and, therefore, have

independent subfunctionalization stories (Trevaskis et al., 2007). Overexpression of *BM1* and

BM10 in barley plants grown under short days resulted in the formation of vegetative tillers in

the base of the spike (Trevaskis et al., 2007), which were similar to the tillers formed in the

spike-like structures described in the wheat *vrn1ful2*-null spikes in this study. Based on these

observations, we hypothesize that the 10-fold upregulation of *VRT2*, *BM1* and *BM10* transcripts

detected in the *vrn1ful2*-null mutants (Fig. S6) may have contributed to the reversion of the

lateral SMs into VMs.

FUL2 mutants increase the number of florets per spikelet

In wheat, the spikelet meristem is indeterminate and can initiate a variable number of floret

primordia. Most of these primordia are aborted during spikelet development, a process that is

affected by genetic differences (Guo et al., 2016; Sakuma et al., 2018), photoperiod (Gonzalez et

al., 2005), plant height (Miralles et al., 1998), and available resources (Ferrante et al., 2013). An

increase in the number of florets per spikelet was detected in *ful2*-null but not in *vrn1*-null and

ful3-null, which suggests that among these three paralogs only *FUL2* contributes to maintaining

a limited number of florets per spikelet (Fig. S5B). Since a higher number of florets per spikelet

can contribute to a higher grain yield potential, it would be interesting to explore the natural

variation in *FUL2* and its effect on this trait.

Overexpression of wheat *miR172* or loss of function mutations in its *AP2-5* target also result in

higher number of florets per spikelet (Debernardi et al., 2017). The synergistic interaction

detected in this study between miR172 and *ful2*-null for this trait (Fig. S5C) suggests that

miR172/AP2-5 and *FUL2* may control floret number through a common pathway. However, the

actual mechanism by which *FUL2* contributes to maintaining a low number of florets per

spikelets is currently unknown. Since loss-of-function mutations in *FT2* also affect the number of

florets per spike (Shaw et al., 2018), we have initiated crosses between *ft2*-null and *ful2*-null to

investigate their genetic interactions.

395 *VRN1* and *FUL2* have essential and redundant roles in wheat spike determinacy

The determinate growth of the wheat spike requires the transition of the distal IM into a SM and

the formation of a terminal spikelet. The *vrn1ful2*-null mutant was unable to form spikelets, so it

was not surprising that it was unable to form a terminal spikelet (Fig. 2H and 4B). However, the

fate of the terminal and lateral meristems in *vrn1ful2*-null was different, with the lateral

meristems reverting to VMs and the terminal IM remaining indeterminate. The IM seems to be

very sensitive to the activity of *FUL2* or *VRN1*, since a single functional copy of either of these

genes in a heterozygous state was sufficient to restore spike determinacy (Fig 3D and K).

Loss-of-function mutations in *TERMINAL FLOWER 1* (*TFL1*) in Arabidopsis or in the

CENTRORADIALIS (*CEN*) homolog in *Antirrhinum* result in the formation of a terminal flower

and the transformation of indeterminate into determinate inflorescences (Bradley et al., 1997;

Ratcliffe et al., 1999). In Arabidopsis, *TFL1* is a direct target of AP1 (Kaufmann et al., 2010),

and the *ap1cal* and *ap1calful* mutants show *TFL1* upregulation and ectopic expression in young

lateral meristems preventing the formation of flowers (Ferrándiz et al., 2000; Ratcliffe et al.,

1999). The rice *CEN* homologs (*RCN1-RCN4*) also repress floral fate and affect inflorescence

architecture (Kaneko-Suzuki et al., 2018; Nakagawa et al., 2002), which suggests conserved

functions across distantly divergent plant lineages. Based on this conservation, it is tempting to

speculate that the upregulation of the wheat *CEN2*, *CEN4* and *CEN5* homologs in the developing

spike of the *vrn1ful2*-null mutant might have contributed to its indeterminate growth.

414 The *vrn1*-null and *ful2*-null mutants have a higher number of spikelets per spike

In wheat, the timing of the transition between the IM and the terminal spikelet determines the

number of spikelets per spike. Since *VRN1* and *FUL2* play a central role in this transition, we

hypothesized that changes in their dosage could affect the number of spikelets per spike. This

hypothesis was confirmed in the single *vrn1*-null and *ful2*-null mutants, which showed increases

of nine and two spikelets per spike respectively (Fig. 4C-D). The stronger effect of *vrn1*-null

relative to *ful2*-null on spikelet number is likely associated with *VRN1*'s stronger effect on

heading time (Fig. 1A-C), which results in a longer period of spike development, thereby

providing more time for the formation of additional spikelets. A stop codon mutation in a

homolog of AP1 in rapeseed altered plant architecture and increased the number of seeds per

plant (Shah et al., 2018), suggesting that mutations in this group of meristem identity genes may

be useful to modulate seed number in other species.

In the field study, the *ful2*-null plants showed a 30.8% increase in the average number of grains

per spike compared with the control sister lines. This large increase was likely a result of the

simultaneous increase in the number of spikelets per spike and florets per spikelet in this mutant.

Although in this experiment the positive increase in grain number was partially offset by a

decrease in average grain weight, the total grain weight per spike was still slightly higher (6.3%)

in the *ful2*-null mutant. It would be interesting to test if the introgression of this mutation in

genotypes which higher biomass (increased "source") can reduce the negative correlation

between grain number and grain weight. In addition, optimum agronomic conditions will be

required to translate the *ful2*-null increase in grain number into larger increases in total grain

yield.

The results presented here indicate that *VRN1* and *FUL2* play key and redundant roles in the

development of the spikelet, the reproductive structure that defines the grass family. Moreover,

our results indicate that these two genes are required for the determinate growth of the wheat

spike, and that loss-of-function of either gene affect the number of spikelets per spike, which is

an important component of grain yield. In summary, our results suggest that a better

understanding of the processes that control the development of grass flowers and inflorescences

may contribute to improving the productivity of a group of cereal crop species that is critical for

the global food supply.

ACKNOWLEDGEMENTS

This project was supported by the Howard Hughes Medical Institute, NRI Competitive Grant

2016-67013-24617 from the USDA National Institute of Food and Agriculture (NIFA) and the

International Wheat Partnership Initiative (IWYP). We thank Dr. Alejandra Alvarez and Dr. Josh

Hegarty for their help with field experiments and Dr. Daniel Wood and Dr. Juan Debernardi for

their valuable comments and suggestions.

COMPETING FINANCIAL INTERESTS STATEMENT

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

CL and JD designed research; JD provided overall supervision to the project; HL, CL, AC, ML

and JJ performed research; CL, HL and JD analyzed data; JD provided statistical analyses; CL

wrote first draft and JD the final version. All authors reviewed the paper.

MATERIALS AND METHODS

Selected mutations and mutant combinations

An ethyl methane sulphonate (EMS) mutagenized population of the tetraploid wheat variety

Kronos was screened for mutations initially using *Cel*I assays (Uauy et al., 2009) and later using

BLAST searches in the database of sequenced mutations for the same population (Krasileva et

al., 2017). We identified loss-of-function mutations in the A and B genome homeologs of *FUL2*

and *FUL3*, which were confirmed using genome specific primers described in Table S1. Single

genome mutants were backcrossed two to three times to Kronos *vrn2*-null to reduce background

mutations. A Kronos *vrn2*-null line with no functional copies of the *VRN2* flowering repressor

(Distelfeld et al., 2009) was used as recurrent parent to avoid the extremely late flowering of

mutant combinations including the *vrn1*-null. All mutants described in this study are in a *vrn2*-

null background unless indicated otherwise.

For *FUL-A2*, we selected line T4-837 (henceforth *ful-A2*), which has a mutation in the splice

donor site of the fifth intron. RT-PCR and sequencing analysis of the *ful-A2* transcripts revealed

two incorrect splicing forms. The most abundant form skipped the fifth exon, which resulted in a

deletion of 14 amino acids in the middle of the K-box (Δ144-157). In the other alternative

splicing form, the retention and translation of the fifth intron generated a premature stop codon

that disrupted the K-box and removed the entire C-terminus (Fig. S2A). For *FUL-B2*, we

selected line T4-2911 that carries a C to T change at nucleotide 484 (henceforth *ful-B2*). The *ful-*

B2 mutation generates a premature stop codon at position 162 (Q162*) that removed the last 13

amino acids of the K-box and the entire C-terminus (Fig. S2A).

For *FUL-A3*, we selected line T4-2375 that carries a G to A mutation in the splice acceptor site

of the third intron. Sequencing of *ful-A3* transcripts revealed that this mutation generated a new

splice acceptor site that shifted the reading frame by one nucleotide. The alternative translation

generated a premature stop codon that truncated 72% of the K-box and the entire C-terminus

(Fig. S2B). For *FUL-B3*, we selected line T4-2139 that carries a C to T mutation at nucleotide

position 394 that generates a premature stop codon at amino acid position 132 (Q132*). This

premature stop removed half of the K-box and the complete C-terminus (Fig. S2B). Given the

critical roles of the K-domain in protein-protein interactions, and the C-terminal domain in

transcriptional activation, these selected mutations are expected to impair the normal function of

the FUL2 and FUL3 proteins.

The A and B-genome mutants for each gene were intercrossed to generate double mutants, which

for simplicity, are referred to hereafter as null mutants. The *ful2*-null and *ful3*-null were

intercrossed with a *vrn1vrn2*-null mutant (*vrn-A1*-null T4-2268 / *vrn-B1* T4-2619 / *vrn2*-null)

(Chen and Dubcovsky, 2012) to generate *vrn1ful2*-null and *vrn1ful3*-null, which were finally

intercrossed to generate *vrn1ful2ful3*-null (all in a *vrn2*-null background). The *vrn1ful2*-null and

vrn1ful2ful3-null mutants were sterile, so they were maintained and crossed by keeping the *ful-*

B2 mutation in heterozygous state.

Transgenic Kronos plants overexpressing *FUL2* and *FUL3* coding regions were generated at the

UC Davis Transformation facility using *Agrobacterium*-mediated transformation. The coding

regions of these two genes were cloned from Kronos into the binary vector pLC41 (Japan

Tobacco, Tokyo, Japan) downstream of the maize *UBIQUITIN* promoter. A C-terminal 3xHA

tag was added to FUL2 and a C-terminal 4xMYC tag was added to FUL3. Mutant and transgenic

wheat plants were grown in PGR15 CONVIRON chambers under LD (16h light/8h dark, light

intensity ~330 μM m^{-2} s^{-1}) at 22 ºC during the day and 18 ºC during the night.

To study the effect of *Ubi::FUL2* in different genetic backgrounds we crossed the Kronos

Ubi::FUL2 with Kronos-*vrn1vrn2*-null and analyzed the effect of the three genes in the F_2

progeny under greenhouse conditions. A field experiment comparing *ful2*-null and its control

line was performed at the University of California, Davis field station during the 2017-2018

growing season (sowed on 12/1/2017 and harvested on 06/25/2018). One meter rows (30 grains

per row) were used as experimental units and the experiment was organized in a randomized

complete block design with eight blocks. During the growing season plants received 200 units of

N as ammonium sulfate, three irrigations, one application of broad-leaf herbicides (2, 4D +

Buctril) and alternating applications of fungicides Quadris and Tilt every 2 weeks.

Effect of the splice site mutations in *ful-A2* and *ful-B3* mutants

To determine the effect of the splice site mutations in *ful-A2* and *ful-B3*, we extracted total RNA

from leaf samples using the Spectrum™ Plant Total RNA kit. cDNA was synthesized from 2 μg

of RNA using the High Capacity Reverse Transcription Kit according to the manufacturer's

instructions and used as RT-PCR template. For *ful-A2*, we used primers *FUL2-837-F* (5'-

CCATACAAAAATGTCACAAGC-3') and *FUL2-837-R* (5'-TTCTGC CTCTCCACCAGTT-3')

for RT-PCR. These primers, which are not genome specific, amplified three fragments of 303 bp,

220 bp and 178 bp. We gel-purified these fragments, cloned them into pGEM-T vector

(Promega), and sequenced them. The 220 bp fragment was from the wild type *FUL-B2* allele,

whereas the other two fragments corresponded to two alternative splicing forms of *ful-A2* that

either retained the fifth intron (303 bp) or skipped the fifth exon (178 bp).

For the *ful-B3* mutant, we performed RT-PCR using primers FUL3-2375-F (5'-

ATGGATGTGATTCTTGAAC-3') and FUL3-2375-R (5'-

TGTCCTGCAGAAGCACCTCGTAGAGA-3'). Sequencing analysis of the PCR products

showed that the G to A mutation generated a new splice acceptor site with an adjacent G that

shifted the reading frame by one nucleotide after 333 bp, and generated a premature stop codon.

Scanning Electron-Microscopy (SEM)

Apices from different developmental stages were dissected and fixed for a minimum of 24 h in

FAA (50% ethanol, 5% (v/v) acetic acid, 3.7% (v/v) formaldehyde), and then dehydrated

through a graded ethanol series to absolute ethanol. Samples were critical-point dried in liquid

CO_2 (tousimis ® 931 Series critical point drier), mounted on aluminum stubs, sputter-coated

with gold (Bio-Rad SEM Coating System Model E5100), and examined with a Philips XL30

scanning electron-microscope operating at 5KV. Images were recorded at slow scan 3 for high

definition and saved as TIFF files.

RNA extraction and Real-time qPCR analysis

RNAs from apices were extracted using the Trizol reagent (ThermoFisher Scientific, Cat. No.15596026). One µg of RNA was used for cDNA synthesis following the instructions of the "High-Capacity cDNA Reverse Transcription Kit" (ThermoFisher Scientific, Cat. No. 4368814). The cDNA was then diluted 20 times and 5 µl of the dilution was mixed with 2×VeriQuest Fast SYBR Green qPCR Master Mix (Affymetrix, Cat. No. 75690) and with primers for the real-time qPCR analysis. Primer sequences are listed in Table S2. *INNITIATION FACTOR 4A* (*IF4A*) was used as an endogenous control. Transcript levels for all genes are expressed as linearized fold-*IF4A* levels calculated by the formula $2^{(IF4A\ C_T - TARGET\ C_T)} \pm$ standard error (SE) of the mean. The resulting number indicates the ratio between the initial number of molecules of the target gene and the number of *IF4A* molecules.

REFERENCES

Bradley, D., Ratcliffe, O., Vincent, C., Carpenter, R. and Coen, E. (1997). Inflorescence commitment and architecture in Arabidopsis. *Science* **275**, 80-83.

Chen, A. and Dubcovsky, J. (2012). Wheat TILLING mutants show that the vernalization gene *VRN1* down-regulates the flowering repressor *VRN2* in leaves but is not essential for flowering. *PLoS Genetics* **8**, e1003134.

Ciaffi, M., Paolacci, A. R., Tanzarella, O. A. and Porceddu, E. (2011). Molecular aspects of flower development in grasses. *Sexual Plant Reproduction* **24**, 247-282.

Debernardi, J. M., Lin, H., Chuck, G., Faris, J. D. and Dubcovsky, J. (2017). microRNA172 plays a crucial role in wheat spike morphogenesis and grain threshability. *Development* **144**, 1966-1975.

Distelfeld, A., Tranquilli, G., Li, C., Yan, L. and Dubcovsky, J. (2009). Genetic and molecular characterization of the *VRN2* loci in tetraploid wheat. *Plant Physiology* **149**, 245-257.

Ferrándiz, C., Gu, Q., Martienssen, R. and Yanofsky, M. F. (2000). Redundant regulation of meristem identity and plant architecture by *FRUITFULL*, *APETALA1* and *CAULIFLOWER*. *Development* **127**, 725-734.

Ferrante, A., Savin, R. and Slafer, G. A. (2013). Floret development and grain setting differences between modern durum wheats under contrasting nitrogen availability. *Journal of Experimental Botany* **64**, 169-184.

Fu, D., Szűcs, P., Yan, L., Helguera, M., Skinner, J., Hayes, P. and Dubcovsky, J. (2005). Large deletions within the first intron in *VRN-1* are associated with spring growth habit in barley and wheat. *Molecular Genetics and Genomics* **273**, 54-65.

Gocal, G. F. W., King, R. W., Blundell, C. A., Schwartz, O. M., Andersen, C. H. and Weigel, D. (2001). Evolution of floral meristem identity genes. Analysis of *Lolium temulentum* genes related to *APETALA1* and *LEAFY* of Arabidopsis. *Plant Physiology* **125**, 1788-1801.

Gonzalez, F. G., Slafer, G. A. and Miralles, D. J. (2005). Floret development and survival in wheat plants exposed to contrasting photoperiod and radiation environments during stem elongation. *Functional Plant Biology* **32**, 189-197.

Guo, Z. F., Slafer, G. A. and Schnurbusch, T. (2016). Genotypic variation in spike fertility

traits and ovary size as determinants of floret and grain survival rate in wheat. *Journal of*

Experimental Botany **67**, 4221-4230.

Honma, T. and Goto, K. (2001). Complexes of MADS-box proteins are sufficient to convert

leaves into floral organs. *Nature* **409**, 525-9.

Kaneko-Suzuki, M., Kurihara-Ishikawa, R., Okushita-Terakawa, C., Kojima, C., Nagano-

Fujiwara, M., Ohki, I., Tsuji, H., Shimamoto, K. and Taoka, K. I. (2018). TFL1-Like

proteins in rice antagonize rice FT-like protein in inflorescence development by competition for

complex formation with 14-3-3 and FD. *Plant and Cell Physiology* **59**, 458-468.

Kaufmann, K., Wellmer, F., Muino, J. M., Ferrier, T., Wuest, S. E., Kumar, V., Serrano-

Mislata, A., Madueno, F., Krajewski, P., Meyerowitz, E. M. et al. (2010). Orchestration of

floral initiation by APETALA1. *Science* **328**, 85-89.

Kellogg, E. A. (2001). Evolutionary history of the grasses. *Plant Physiology* **125**, 198-1205.

Kellogg, E. A., Camara, P. E. A. S., Rudall, P. J., Ladd, P., Malcomber, S. T., Whipple, C.

J. and Doust, A. N. (2013). Early inflorescence development in the grasses (Poaceae). *Frontiers*

in Plant Science **4**.

Kippes, N., Chen, A., Zhang, X., Lukaszewski, A. J. and Dubcovsky, J. (2016). Development

and characterization of a spring hexaploid wheat line with no functional *VRN2* genes.

Theoretical and Applied Genetics **129**, 1417–1428.

Krasileva, K. V., Vasquez-Gross, H., Howell, T., Bailey, P., Paraiso, F., Clissold, L.,

Simmonds, J., Ramirez-Gonzalez, R. H., Wang, X., Borrill, P. et al. (2017). Uncovering

hidden variation in polyploid wheat. *Proceedings of the National Academy of Sciences of the*

United States of America **114**, E913–E921

Li, Q., Wang, Y., Wang, F. X., Guo, Y. Y., Duan, X. Q., Sun, J. H. and An, H. L. (2016).

Functional conservation and diversification of *APETALA1/FRUITFULL* genes in *Brachypodium*

distachyon. *Physiologia Plantarum* **157**, 507-518.

Liu, C., Zhou, J., Bracha-Drori, K., Yalovsky, S., Ito, T. and Yu, H. (2007). Specification of

Arabidopsis floral meristem identity by repression of flowering time genes. *Development* **134**,

1901-1910.

Malcomber, S. T., Preston, J. C., Reinheimer, R., Kossuth, J. and Kellogg, E. A. (2006).

Developmental gene evolution and the origin of grass inflorescence diversity. *Advances in*

Botanical Research **44**, 425-481.

Miralles, D. J., Katz, S. D., Colloca, A. and Slafer, G. A. (1998). Floret development in near

isogenic wheat lines differing in plant height. *Field Crops Research* **59**, 21-30.

Nakagawa, M., Shimamoto, K. and Kyozuka, J. (2002). Overexpression of *RCN1* and *RCN2*,

rice *TERMINAL FLOWER 1/CENTRORADIALIS* homologs, confers delay of phase transition

and altered panicle morphology in rice. *Plant Journal* **29**, 743-750.

Pearce, S., Vanzetti, L. S. and Dubcovsky, J. (2013). Exogenous gibberellins induce wheat

spike development under short days only in the presence of *VERNALIZATION 1. Plant*

Physiology **163**, 1433–1445.

Peng, J. R., Richards, D. E., Hartley, N. M., Murphy, G. P., Devos, K. M., Flintham, J. E.,

Beales, J., Fish, L. J., Worland, A. J., Pelica, F. et al. (1999). 'Green revolution' genes encode

mutant gibberellin response modulators. *Nature* **400**, 256-261.

Preston, J. C., Christensen, A., Malcomber, S. T. and Kellogg, E. A. (2009). MADS-box

gene expression and implications for developmental origins of the grass spikelet. *American*

Journal of Botany **96**, 1419-1429.

Preston, J. C. and Kellogg, E. A. (2006). Reconstructing the evolutionary history of paralogous

APETALA1/FRUITFULL-like genes in grasses (Poaceae). *Genetics* **174**, 421-437.

Preston, J. C. and Kellogg, E. A. (2008). Discrete developmental roles for temperate cereal

grass *VRN1/FUL*-like genes in flowering competency and the transition to flowering. *Plant*

Physiology **146**, 265-276.

Ratcliffe, O. J., Bradley, D. J. and Coen, E. S. (1999). Separation of shoot and floral identity in

Arabidopsis. *Development* **126**, 1109-1120.

Ream, T. S., Woods, D. P., Schwartz, C. J., Sanabria, C. P., Mahoy, J. A., Walters, E. M.,

Kaeppler, H. F. and Amasino, R. M. (2014). Interaction of photoperiod and vernalization

determines flowering time of *Brachypodium distachyon. Plant Physiology* **164**, 694-709.

Sakuma, S., Golan, G., Guo, Z., Ogawa, T., Tagiri, A., Sugimoto, K., Bernhardt, N.,

Brassac, J., Mascher, M., Hensel, G. et al. (2018). Unleashing floret fertility by a mutated

homeobox gene improved grain yield during wheat evolution under domestication. *bioRxiv,*
doi.org/10.1101/434985.

Shah, S., Karunarathna, N. L., Jung, C. and Emrani, N. (2018). An *APETALA1* ortholog
affects plant architecture and seed yield component in oilseed rape (*Brassica napus* L.). *BMC*
Plant Biol **18**, 380.

Shaw, L. M., Lyu, B., Turner, R., Li, C., Chen, F., Han, X., Fu, D. and Dubcovsky, J.
(2018). *FLOWERING LOCUS T2* (*FT2*) regulates spike development and fertility in temperate
cereals. *J Exp Bot*.

Theissen, G., Melzer, R. and Rumpler, F. (2016). MADS-domain transcription factors and the
floral quartet model of flower development: linking plant development and evolution.
Development **143**, 3259-3271.

Tranquilli, G. and Dubcovsky, J. (2000). Epistatic interactions between vernalization genes
Vrn-A^{m}1 and *Vrn-A^{m}2* in diploid wheat. *Journal of Heredity* **91**, 304-306.

Trevaskis, B., Tadege, M., Hemming, M. N., Peacock, W. J., Dennis, E. S. and Sheldon, C.
(2007). *Short Vegetative Phase-like* MADS-box genes inhibit floral meristem identity in barley.
Plant Physiology **143**, 225-235.

Uauy, C., Paraiso, F., Colasuonno, P., Tran, R. K., Tsai, H., Berardi, S., Comai, L. and
Dubcovsky, J. (2009). A modified TILLING approach to detect induced mutations in tetraploid
and hexaploid wheat. *BMC Plant Biology* **9**, 115-128.

Woods, D. P., McKeown, M. A., Dong, Y. X., Preston, J. C. and Amasino, R. M. (2016).
Evolution of *VRN2/Ghd7*-like genes in vernalization-mediated repression of grass flowering.
Plant Physiology **170**, 2124-2135.

Yan, L., Loukoianov, A., Tranquilli, G., Helguera, M., Fahima, T. and Dubcovsky, J.
(2003). Positional cloning of wheat vernalization gene *VRN1*. *Proceedings of the National*
Academy of Sciences of the United States of America **100**, 6263-6268.

Zhang, X. K., Xia, X. C., Xiao, Y. G., Dubcovsky, J. and He, Z. H. (2008). Allelic variation at
the vernalization genes *Vrn-A1*, *Vrn-B1*, *Vrn-D1* and *Vrn-B3* in Chinese common wheat cultivars
and their association with growth habit. *Crop Science* **48**, 458-470.

663
664

FIGURE LEGENDS

Fig. 1. Effect of *FUL2* and *FUL3* on heading time under long day photoperiod. (A-F) Kronos *vrn2*-null background. (A) *vrn1*-null (n= 6) versus control (n= 6). (B) *ful2*-null (n= 8) *vs.* control (n= 11) in a *Vrn1* background. (C) *ful3*-null (n= 8) and *ful2ful3*-null (n= 15) *vs.* control (n= 10) in a *Vrn1* background. (D) Shoot apical meristem length. The red dotted line indicates the transition of the SAM to the reproductive stage (n= 6 per time point). (E-F) Heading time of Kronos T_1 transgenic plants from two independent events segregating for (E) *FUL2* (*Ubi::FUL2*, n= 4-23) and (F) *FUL3* (*Ubi::FUL3*, n= 9-31). (G-H) Two way interactions for F_2 plants segregating for *VRN1, VRN2* and *FUL2*. (G) *VRN2 x FUL2* in a *Vrn1* heterozygous background. (H) *VRN2 x FUL2* in a *vrn1*-null background. *P* values correspond to a 2 x 2 factorial ANOVA (3-way ANOVA in Table S3). Error bars are SEM. *** = $P < 0.0001$, NS = $P > 0.05$.

Fig. 2. Phenotypical characterization of the *vrn1ful2* and *vrn1ful2ful3* mutants. (A) Stems and heads of *vrn1*-null, *vrn1ful2*-null and *vrn1ful2ful3*-null mutants (leaves were removed before photography). (B-E) Phenotypes of *vrn1*-null (B) spike, (C) spikelet, and (D) dissected spike at terminal spikelet stage. (E) Scanning Electro-Microscopy (SEM) of developing spike. (F-I) *vrn1ful2*-null mutant. (F) Spike-like structure with leafy shoots replacing spikelets. (G) Detached "spikelet (shoot)" (H) Dissection of the spike-like structure showing lateral vegetative meristems and indeterminate growth (I) SEM detail of the spike-like meristem. (J-K) *vrn1ful2ful3*-null mutant. (J) Spike-like structure. (K) SEM detail of the spike-like meristems. All these mutants are in a Kronos *vrn2*-null background.

Fig. 3. **Phenotypical characterization of heterozygous mutants containing one copy of *VRN1* or *FUL2*.** (A-C) *ful2*-null/<u>*Vrn-A1*</u> *vrn-B1*-null. (A) "Head" at early stage. (B) Detached spikelet. (C) Dissection of an individual spikelet showing floral organs including lodicules, ovaries and stamens. (D-G) *ful2*-null/*vrn-A1*-null <u>*vrn-B1*</u>. (D) Terminal spikelet (TS) confirming spike meristem determinacy. (E) Head at later stage. (F) Detached "spikelet" showing indeterminate growth. (G) Spikelet dissection showing rachilla elongation. (H-K) *vrn1*-null/*ful2-A* <u>*Ful2-B*</u> (produces viable grain). (H) Representative spikes showing formation of lateral branches in the basal region of the spike and extra florets in the terminal spikelet. (I) Detached lateral branch. (J) Dissection of the lateral branch. (K) Terminal spikelet (TS) confirming spike

meristem determinacy. Sp= spikelets, Glu= glumes, Pa= palea, Le= lemma, Ra= rachilla. All

these mutants are in a Kronos *vrn2*-null background.

Fig. 4. *VRN1* and *FUL2* play redundant roles in the control of spike determinacy and affect

spikelet number. (A) Scanning Electro-Microscopy of a normal wheat spike with a terminal

spikelet in the *vrn1*-null control. (B) *vrn1ful2*-null mutant spike with indeterminate apical

meristem. (C-E) Number of spikelets per spike in a growth chamber experiment (n=6). (C) *vrn1*-

null (58% increase), (D) *ful2*-null (10% increase) and (E) *ful3*-null (no significant increase). Bars

represent mean ± SEM and asterisks indicate statistically significant difference to the control line

(** = $P < 0.01$, *** = $P < 0.001$, NS = $P > 0.05$). (F) ANOVAs for spike traits in *ful2*-null and

sister control lines in the field (randomized complete block design with 8 blocks).

SUPPLEMENTARY FIGURES

Fig. S1. Phylogeny of duplicated Arabidopsis AP1/CAL/FUL and grasses VRN1/FUL2/FUL3 clades.

Fig. S2. Selected *ful2* and *ful3* mutations and their effect on the encoded proteins.

Fig. S3. Effect of *vrn1*-null, *ful2*-null and *ful3*-null mutations on stem length.

Fig. S4. Dissection of basal tillers from the *vrn1ful2*-null spike-like structure.

Fig. S5. Effect of *ful2*-null and *ful3*-null on the number of florets per spikelet.

Fig. S6. Transcript levels of wheat *SVP*-like MADS-box genes *VRT2*, *BM1* and *BM10*.

Fig. S7. Effect of the overexpression of *FUL2* and *FUL3* on the number of spikelets per spike.

Fig. S8. Transcript levels of wheat *TFL1/CEN*-like genes *CEN2*, *CEN4* and *CEN5*.

Fig. 1

Fig. 2.

Fig. 3

Field exp.	No. Spikelets /Spike	Grain No. /spike	Grain No. /spikelet	Avg. grain weight (mg)	Grain weight /spike (g)
Control	15.7	41.9	2.67	57.7	2.41
ful2-null	18.9	54.8	2.91	47.0	2.57
% diff. of control	20.5%	30.8%	8.7%	-18.6%	6.3%
ANOVA *P* value	0.0002	0.0002	0.0502	0.0012	0.0982

Fig. 4