Title: Divergence in gut microbial communities mirrors a social group fission event in

a black-and-white colobus monkey (*Colobus vellerosus*)

Running title: Microbiome divergence mirrors group fission

Authors: Claire K. Goodfellow[1,2], Tabor Whitney[2], Diana M. Christie[1,3], Pascale Sicotte[4], Eva

C. Wikberg[5,*] Nelson Ting[1,3,*]

Author Affiliations:

1. Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97405, USA

2. Department of Biology, University of Oregon, Eugene, OR 97405, USA

3. Department of Anthropology, University of Oregon, Eugene, OR 97405, USA

4. Department of Anthropology and Archaeology, University of Calgary, Calgary, AB

T2N1N4, Canada

5. Department of Anthropology, University of Texas San Antonio, San Antonio, TX 78249,

USA

*. Co-corresponding author

ABSTRACT

Host behavior and social factors have increasingly been implicated in structuring the

composition of gut microbial communities. In social animals, distinct microbial communities

characterize different social groups across a variety of taxa, although little longitudinal research

has been conducted that demonstrates how this divergence occurs. Our study addresses this

question by characterizing the gut microbial composition of an African Old World monkey, the

black-and-white colobus (*Colobus vellerosus*), prior to and after a social group fission event. Gut

microbial taxonomic composition of these monkeys was profiled using the V-4 hypervariable

region of the bacterial 16s rRNA gene, and pairwise-relatedness values were calculated for all

individuals using 17 STR loci and partial pedigree information. The two social groups in this

study were found to harbor distinct microbial signatures after the fission event from which they

emerged, while these communities were not divergent in the same individuals prior to this event.

Furthermore, although this fission happened partially along lines of relatedness, relatedness did

not structure the differences that we found. Taken together, this study suggests that distinct gut

microbial profiles can emerge in social groups in less than one year and recommends further

work into more finely mapping the timescales, causes, and potentially adaptive effects of this

recurring trend toward distinct group microbial signatures.

Keywords: gut microbiome, group fission, social groups, sociality

Research highlights:

• Distinct gut microbial profiles emerge in two social groups of *C. vellerosus* less than nine

months after a fission event.

• Three genera differ in abundance between the two new social groups.

• Relatedness does not structure differences in microbial composition between the groups.

INTRODUCTION

The mammalian gut harbors a dynamic microbial community which contributes to host

physiology, metabolism, and defense (Huttenhower *et al.*, 2012; Cho & Blaser, 2012;

Barbachano *et al.,* 2017). This community both shapes host phenotypes and is shaped by host

characteristics that can include phylogeny, genetic variation, environment, and spatial

distribution (Wu *et al.*, 2011; Leamy *et al.*, 2014; Barelli *et al.*, 2015; Amato, 2013; Amato *et al.*,

2016). Moreover, behavior and social context can contribute to gut microbiome composition, and

diet choice, habitat use, mate choice, and social networks have all been shown to modulate gut

diversity (Archie *et al.*, 2015; Ezenwa *et al.*, 2012). In fact, the social transmission of beneficial

microbes has been cited as one of the benefits associated with group living (Lombardo, 2008),

and it has been suggested that more similar gut communities between hosts may confer similar

"ecosystem services" to their hosts (Costello *et al.*, 2012). Given that conspecifics in the same

social group likely encounter highly comparable ecological challenges, the social transmission

and ultimate convergence of a group's gut microbiota into the consortia that provides the most

ideal ecosystem services for that particular group's set of demands could prove to be of

evolutionary benefit.

Distinguishing the complex and intertwined forces that shape this dynamic community,

however, is difficult. Studies of wild populations can help to address this difficulty, providing

insight into the forces at play in natural communities as well as how they change over time

(Amato, 2013). In particular, studies of wild primates and other highly social animals allow us to

answer important questions about how social forces shape these changes in some of our own

closest living relatives. For example, in a number of wild primate populations, more closely

associated individuals have more homogenous gut microbiome compositions (Perofsky *et al.*,

2017; Moeller et al., 2016; Amato *et al.,* 2017) and distinct gut microbiota characterize different

social groups in the same population (Tung *et al.,* 2015; McCord *et al.*, 2014; Bennett *et al.*,

2016; Springer *et al.* 2017; Degnan *et al.,* 2012). Taken together, these findings highlight the

importance of social context in gut microbiome composition. Furthermore, immigrant males

which have resided in a new social group for a longer period of time have more similar gut

microbiota to the resident males of that group, suggesting that the convergence of group member

microbiota may occur over a span of months to years (Grieneisen *et al.*, 2017). Because dietary

shifts typically alter the composition of microbial communities over a shorter time scale of days

to weeks, this finding suggests that distinct group communities are not solely the result of

changes in diet (Bonte *et al.*, 2012; Williams *et al.*, 2012; Turnbaugh *et al.*, 2009). However,

further studies are needed to more thoroughly examine the time scales over which these

convergences occur in natural communities.

Group fission events provide an ideal natural system for interrogating such questions.

These are a means of group proliferation in social animals that occur when the costs of living in a

certain group have grown to outweigh the benefits (Sueur & Maire, 2014). When this happens,

one or more social groups will break off from the original group, oftentimes splitting along lines

of relatedness (Widdig *et al.*, 2006; Snyder-Mackler *et al.*, 2014). This type of event provides us

the opportunity for unique insight into the physiological and behavioral changes in individuals

following such an event as well as the time scales over which they occur. For example, fission

events and variations in group size have enabled insight into the effects of social context on

grooming networks, fertility and cortisol levels in primates (Henzi *et al.*, 1997; Dunbar, 2018;

Markham *et al.*, 2015). Here, we report on the gut microbiome compositions of a group of ursine

colobus or white-thighed black-and-white colobus (*C. vellerosus*) before, and less than a year

after, a fission event. The aim of this study was to examine the plasticity of the gut microbiome

shortly following a fission event as a way of gaining insight into the time scale over which

microbiomes diverge into the distinct microbiomes that have been shown to characterize

different social groups.

METHODS

Study system

The Boabeng-Fiema Monkey Sanctuary (BFMS) is a 1.92 km^2 dry semi-deciduous forest (Hall & Swaine, 1981) located in central Ghana (7°43' N and 1°42'W). Ursine colobus (*Colobus vellerosus*) is one of two diurnal primate species that resides at BFMS (Saj *et al.*, 2005). This is an arboreal, folivorous monkey (Saj & Sicotte 2007a, 2007b) that lives in uni-male or multi-male multi-female groups of 9-38 animals (Kankam & Sicotte, 2013; Wong & Sicotte, 2006). Dispersal is male-biased in this species (Teichroeb *et al*. 2011), although females do show facultative dispersal (Teichroeb *et al*. 2009, Teichroeb *et al*. 2011, Wikberg *et al*. 2012, Sicotte *et al.*, 2017). Female networks are affected by the presence of infants, kinship, and immigration status, but not by dominance rank (Wikberg *et al*. 2013, Wikberg *et al*. 2014a, Wikberg *et al*. 2014b, Wikberg et a. 2015). Several groups of *C. vellerosus* have been followed systematically since 2000 for behavioral, demographic and ecological data (see for example Teichroeb & Sicotte, 2009; Bǎdescu *et al*. 2015). Fecal samples were collected on a regular basis from each focal female in our study groups between 2006 and 2009 (see for example Wikberg *et al*. 2015). This study follows the fission of one social group to two daughter groups (named NP and DA) over the course of one year (2006-2007).

Sample collection and STR genotyping

All fecal samples from the study period were collected using masks, fresh gloves, sterile sticks, and sterile tubes to minimize contamination. 1-2 grams of feces were mixed with 6 μL of RNAlater® (Thermo Fisher Scientific, Waltham MA) immediately upon collection and stored at

-20°C in the field. After shipment to the Ting lab, samples were again stored at -20°C until DNA

extraction. DNA was extracted from two or more samples of each individual using a QIAamp

DNA Stool Mini Kit with a slightly modified manufacturer protocol (Wikberg *et al.* 2012), and

negative controls were processed with each round of extraction. 17 short tandem repeat (STR)

loci were amplified using Qiagen's multiplex PCR kit with a modified protocol and analyzed on

an ABI 3730 DNA analyzer (Wikberg *et al.*, 2012). We determined how many replicates were

needed to confirm homozygote genotypes based on real-time PCR DNA quantification (Morin *et*

al., 2001). Two replicates were used to confirm heterozygote genotypes.

Gut microbial profiling

We collected metagenomic data from matched genotyped samples from female members

of the original social group collected from June to August, 2006 (n = 12 samples) and from

matched genotyped samples from the same female individuals residing in the two daughter

groups from July to August, 2007 (NP with n = 6 samples; DA with n = 6 samples). DNA was

extracted again as above and quantified using a Qubit dsDNA BR Assay Kit protocol using a

Qubit 2.0 Fluorometer (Thermo Fisher Scientific, Waltham MA). DNA extracts containing at

least 1.0 ng/µL were chosen for preparation and sequencing of the V-4 hypervariable region of

the bacterial 16S ribosomal RNA gene in the Genomics and Cell Characterization Core Facility

at the University of Oregon. 200 ng of DNA diluted in 10 µL of H2O were PCR amplified using

barcoded Illumina 515F and 806R primers. Targets were amplified in reactions of 1 µL DNA,

1.25 µL of 10 µM primer mix, 10.25 µL H2O, and 12.5 µL NEB Q5 hot start 2x Master Mix.

The thermal cycling profile was as follows: initial denaturing at 98°C for 0:30, 20-30 cycles of

98°C for 0:10, 61°C for 0:20, and 72°C for 0:20, and a final extension of 72°C for 2:00. PCR

products were cleaned using Ampure XP beads (Beckman Coulter, Brea, CA), quantified and

normalized. Barcoded amplicons were pooled and pair-end sequenced with 150 base pair reads

on a partial medium output run on the Illumina NextSeq platform. Sequences were then

demultiplexed and denoised using DADA2 (Bolger *et al.*, 2014). Taxonomic units were assigned

using the Qiime2 pipeline. An OTU table was generated for samples rarefied to an even

sampling depth of 46,040 reads per sample, retaining 1,104,960 sequences for 24 samples.

Negative controls were processed at both the extraction and library preparation (PCR) stages,

and they were sequenced and carried through the data processing pipelines. No evidence of

contamination was found via fluorometry or gel electrophoresis, nor was there any evidence of

contaminating sequences in the negative controls.

Data Analyses

Unless otherwise noted, all subsequent statistical analyses were run in R. To test whether

average gut microbial composition differed by social group, samples were analyzed using four

groups based on social group at the time of sample collection: original group which became NP

after the fission, original group which became DA after the fission, DA, and NP. Beta diversity

was calculated for the samples as Bray-Curtis dissimilarity using the *phyloseq* package for R

(McMurdie & Holmes, 2013). This metric was selected over metrics accounting for evolutionary

relatedness as it represents a quantitative measure of community dissimilarity based on relative

abundance without adjusting for the phylogenetic proximity of OTUs, thereby allowing insight

into structure without accounting for relatedness among microbial taxa. PERMANOVAs were

run using the *adonis* function in the *vegan* package. Linear effect size analysis (LefSe) was run

for the groups at a KW alpha value of 0.01 and an LDA score of 3.0 (Segata *et al.*, 2011).

To assess whether the fission occurred along lines of host relatedness, a Kruskal-Wallis

test was run using metrics of average group relatedness for the three social groups (original

group, DA, NP). Finally, we determined whether host relatedness explained differences in gut

microbial beta diversity seen between groups. We investigated the correlation between the Bray-

Curtis dissimilarity matrix and the host relatedness matrix for each group using Mantel tests, and

Spearman rank correlation statistics were computed with 999 permutations.

169   **RESULTS**

In the summer of 2006, one of our study groups (28 individuals) at BFMS began to show

high female aggression, and subgroups started to range 50 meters apart for periods of time

(Wikberg, unpublished data). *C. vellerosus* typically exhibit a smaller group spread, and 50

meters is used to define a between-group encounter in this species (Sicotte & MacIntosh, 2004).

The home range of the original group spanned approximately 0.20 km^2 through both woodland

and mature forest. By May of 2007, this group had fissioned into two daughter groups: NP (10

individuals) and DA (18 individuals) (Table 1). These two groups both remained on the original

home range, splitting the range after the fission (Fig. 1). Both daughter groups ranged in subsets

of the original range that included both woodland and mature forest, but DA ranged 0.15 km^2,

moreso in woodland areas, while NP's new range was 0.054 km^2, moreso in mature forest after

the fission event.

The original group was broken down into two groups for the sake of analysis—those

female individuals which eventually split into the DA group and those females that split into the

NP group. Average Bray-Curtis dissimilarity was 0.527 (SD +/- 0.046) for the two groups before

the fission event, while it was 0.538 (SD +/- 0.048) for the two social groups after the fission

event. Bray-Curtis dissimilarity between the two groups was not significant prior to the fission (p

= 0.085, n = 12, pseudo F = 1.23), while differences after the fission were significant (p = 0.02, n

= 12, pseudo F = 1.62). Within year, group membership explained 7.0% of the variation in gut

microbiome diversity. Significant differences were found across years for all groups sampled (p

< 0.01, n = 24, pseudo F = 2.19), and this difference was the largest predictor of variation,

explaining 31.4% of the variation in gut microbiome diversity (Fig. 2).

The average pairwise-relatedness among females in the social group prior to the fission

was 0.153, while it was 0.232 in the post-fission NP group and 0.125 in the post fission DA

group (Fig. 3). This average pairwise-relatedness increased by 0.079 in the NP social group from

the original DA group, suggesting that this fission event may have happened partially along lines

of female relatedness, with a group of more closely-related female individuals splitting off to

form the new NP group. However, the average pairwise-relatedness decreased by 0.028 between

the pre-fission group and the new DA group. The pairwise-relatedness was not statistically

different between any of the groups (p = 0.103, df = 2, χ^2 = 4.5518). The Mantel test comparing

Bray-Curtis dissimilarity and relatedness showed no statistically significant correlation between

these variables (r < 0.01, p = 0.494).

On the whole, individual gut samples were vastly dominated by Firmicutes (57-78%) and

Bacteriodes (2-13%). The next most prevalent phyla across samples were Tenericutes (3-10%)

and Verucomicrobia (<1-16%). Linear discriminant effect analysis (LefSe) found three genera to

differ between the two new social groups: *Parabacteroides, Coprococcus, and*

Porphyromonadaceae (Fig. 4).

207   **DISCUSSION**

Distinct gut microbiota characterize different social groups across a wide range of taxa

(Bennett *et al.*, 2016; Degnan *et al.,* 2012; Tung *et al.,* 2015; Amato *et al.,* 2017). Examining the

process and the time scale over which these divergences occur is important to understand the

influence that social context can exert on gut microbiome assembly. Some recent research has

focused on the time scale across which an individual's microbiome converges with that of a new

social group. Grieniesen *et al.* (2017) found that the longer an immigrant male baboon resided in

a new social group, the more closely his core and non-core microbiomes resembled those of the

adult members of that group, suggesting that the process of group convergence takes place over a

span of months to years. Amaral *et al.* (2017) also found that, when newly-weaned infants joined

new social groups, their gut microbiomes converged to resemble their new groups within two

weeks. Our study focuses on the time scale across which social groups diverge from one another.

We found distinct gut microbial signatures to characterize two daughter groups of colobus less

than nine months after the fission event that resulted in these groups. Prior to this fission, the

same individuals did not harbor distinct microbial communities.

Hosts can gain microbes through changes in social context, such as alterations in direct

and indirect interactions with conspecifics that provide access to different microbes, potentially

affecting gut microbiome composition (Lombardo *et al.*, 2008). In social animals, changes in

group composition, size, and social networks could all contribute to this type of shift. In this

study, group composition of females before and after the fission remained similar overall (Table

1); thus the number and age structure of females in each fission product are unlikely to be

driving our results. There were however other changes in social context in DA social group

during the post-fission field season, including two males immigrating to DA group, another

extra-group male chasing females and infants during an intergroup encounter, and one infant

dying. Further investigation into how changes in social environment and social stress affect the

gut microbiome are required to determine how these events may have influenced the observed

gut microbiome divergence.

Other factors potentially contributing to the observed microbial shifts are diet and/or

ranging patterns. While diet has been suggested as a primary driver in structuring the gut

microbiome (Muegge *et al.*, 2011; Amato *et al.,* 2014; Hale *et al.*, 2018), diet has not been found

to explain differences in gut microbial beta diversity between individuals and groups in our

population (Wikberg *et al.,* 2017; Wikberg unpublished data). Alternatively, despite the large

degree of range overlap that remained between the two groups after the fission (Fig. 1), we

cannot rule out the effects of habitat use on the observed shifts. While product groups ranged in

subsets of the original range that included both woodland and mature forest, DA ranged

primarily in woodland areas after the fission while NP's new range tended toward the mature

forest. Because even small changes in environment can expose animals to new reservoirs of

environmentally-derived microbes, it is possible that the divergence observed between the two

product groups is in part driven by spatial distribution and habitat use.

As has been reported in other folivorous species, individual gut samples were vastly

dominated by Firmicutes and Bacteriodes with low but consistent proportions of Tenericutes and

Verucomicrobia (Yildirim *et al.*, 2010; Amato *et al.*, 2016). Linear discriminant analysis (LefSe)

revealed three genera to differ between the two product groups of this study. The new DA group

was found to have relatively more *Porphyromonadaceae* and *Parabacteriodes* than the NP group

after the fission, while the genus *Coprococcus* was found to be at greater prevalence in the NP

group than the new DA group. The *Coprococcus* genus is in the order Clostridiales, which can

assist in the degradation of plant material and is likely to reflect the folivorous diet of these

animals (Barelli *et al.*, 2015). It has been found at differential abundances in different social

groups of baboons, suggesting that this might be a genus with a strong propensity for social

transmission (Grieneisen *et al.*, 2017). This genus is also commonly used to gauge individual gut

health, and decreased levels of *Coprococcus* have been shown to accompany a stress response

(Derrien *et al.*, 2015), which could be related to the changes in social context and/or increased

ranging in woodland habitat seen in the DA social group.

This fission event resulted in an increase in average pairwise-relatedness for the NP but

not DA group. This is a common phenomenon in animals that disperse by group fission, and an

increase in relatedness in fission product groups has been demonstrated widely across primate

species (Widdig *et al.*, 2006; Snyder-Mackler *et al.*, 2014). Although genetic variation can play a

significant role in shaping the diversity of the gut microbiome (Goodrich *et al.*, 2014), previous

studies have found little evidence for a strong role of host genetics in structuring the microbial

communities of wild primates (Degnan *et al.*, 2012; Amato *et al.*, 2017; Spor *et al.*, 2011). In this

particular data set, no correlation existed between beta diversity and relatedness. Taken together,

our results suggest that while this fission happened partially along lines of relatedness,

relatedness did not play a significant role in structuring the differences in beta diversity seen

between the two groups.

Finally, the largest proportion of variation between groups in this study was explained by

year, rather than group membership. Previous studies have found temporal variation in the gut

microbiomes of other folivorous primates to change in response to seasonal changes in food

availability (Amato *et al.*, 2015; Springer *et al.* 2017), which is consistent with past observations

in this study population (Wikberg et al. 2016, Wikberg unpublished data). However, because

longitudinally collected samples in this study were all from the wet season, the observed

differences would need to be explained by some aspect of interannual variation in food

availability during the same season. Further work is needed to clarify this possibility, including

more sampling points between years and seasons as well as detailed data on changes in diet and

food availability through time.

Overall, we used a longitudinal approach that provides a new perspective into how social

groups acquire distinct gut microbial communities and the time period over which these

divergent communities establish. This has significant consequences for understanding the role of

social context in shaping the unique microbial signatures associated with distinct social groups

across a wide variety of taxa. Further work is recommended into more finely mapping the

timescales and factors that result in this divergence, especially within the context of the

potentially adaptive effects of this recurrent, social-context dependent trend.

289   **ACKNOWLEDGEMENTS**

We confirm that all research protocols reported in this manuscript were reviewed and

approved by an appropriate institution and/or governmental agency that regulates research with

animals, all research reported in this manuscript adhered to the legal requirements of the country

in which the work took place, and that the research adhered to the American Society of

Primatologists (ASP) Principles for the Ethical Treatment of Non Human Primates. Specifically,

we gained permission from the Ghana Wildlife Division, the management committee at BFMS,

and the University of Calgary's Animal Care Committee to conduct this study. Funding was

granted by Alberta Ingenuity, American Society of Primatologists, International Primatological

Society, Leakey Foundation, Natural Sciences and Engineering Research Council of Canada,

Province of Alberta, Sweden-America Foundation, Wenner–Gren Foundation (8172), the

REFERENCES

Amaral, W. Z., Lubach, G. R., Proctor, A., Lyte, M., Phillips, G. J., & Coe, C. L. (2017). Social

influences on prevotella and the gut microbiome of young monkeys. *Psychosomatic*

medicine, 79(8), 888-897.

Amato, K. (2013). Co-evolution in context: The importance of studying gut microbiomes in wild

animals. *Microbiome Sci Med*, 1(1), 10-29.

Amato, K. R., Leigh, S. R., Kent, A., Mackie, R. I., Yeoman, C. J., Stumpf, R. M., … Garber, P.

A. (2014). The Gut Microbiota Appears to Compensate for Seasonal Diet Variation in the Wild

Black Howler Monkey (Alouatta pigra). *Microbial Ecology*, 69(2), 434–443.

Amato, K. R., Yeoman, C. J., Cerda, G., Schmitt, C. A., Cramer, J. D., Miller, M. E. B., …

Leigh, S. R. (2015). Variable responses of human and non-human primate gut microbiomes to a

Western diet. *Microbiome*, 3, 53.

Amato, K. R., Martinez-Mota, R., Righini, N., Raguet-Schofield, M., Corcione, F. P., Marini, E.,

... & Williams, L. (2016). Phylogenetic and ecological factors impact the gut microbiota of two

Neotropical primate species. *Oecologia*, 180(3), 717-733.

Amato, K. R., Van Belle, S., Di Fiore, A., Estrada, A., Stumpf, R., White, B., … & Leigh, S. R.

(2017). Patterns in Gut Microbiota Similarity Associated with Degree of Sociality among Sex

Classes of a Neotropical Primate. *Microbial Ecology*, 74(1), 250–258.

Archie, E. A., & Tung, J. (2015). Social behavior and the microbiome. *Current Opinion in*

Behavioral Sciences, 6, 28-34.

Barbáchano, A., Fernández-Barral, A., Ferrer-Mayorga, G., Costales-Carrera, A., Larriba, M. J.,

& Muñoz, A. (2017). The endocrine vitamin D system in the gut. *Molecular and cellular*

endocrinology, 453, 79-87.

Bădescu, I., Sicotte, P., Ting, N., & Wikberg, E. C. (2015). Female parity, maternal kinship,

infant age and sex influence natal attraction and infant handling in a wild colobine (Colobus

vellerosus). *American journal of primatology*, 77(4), 376-387.

Barelli, C., Albanese, D., Donati, C., Pindo, M., Dallago, C., Rovero, F., … De Filippo, C.

(2015). Habitat fragmentation is associated to gut microbiota diversity of an endangered primate:

Implications for conservation. *Scientific Reports*, 5, 1–12.

Bennett, G., Malone, M., Sauther, M. L., Cuozzo, F. P., White, B., Nelson, K. E., ... & Amato, K.

R. (2016). Host age, social group, and habitat type influence the gut microbiota of wild ring-

tailed lemurs (Lemur catta). *American journal of primatology*, 78(8), 883-892.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics*, 30(15), 2114-2120.

Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., ... & Schtickzelle,

N. (2012). Costs of dispersal. *Biological Reviews*, 87(2), 290-312.

Cho, I., & Blaser, M. J. (2012). The human microbiome: at the interface of health and

disease. *Nature Reviews Genetics*, 13(4), 260.

Costello, E. K., Stagaman, K., Dethlefsen, L., Bohannan, B. J., & Relman, D. A. (2012). The

application of ecological theory toward an understanding of the human

microbiome. *Science*, 336(6086), 1255-1262.

Degnan, P. H., Pusey, A. E., Lonsdorf, E. V, Goodall, J., Wroblewski, E. E., & Wilson, M. L.

(2012). Factors associated with the diversification of the gut microbial communities within

chimpanzees from Gombe National Park. *Proceedings of the National Academy of Sciences*

(USA), 109, 13034–13039.

Derrien, M., Johan, E.T., & Vileg, H. (2015). Fate, activity, and impact of ingested bacteria

within the human gut microbiota. *Trends in Microbiology,* 23(6), 354-66.

Dunbar, R. I. M., MacCarron, P., & Robertson, C. (2018). Trade-off between fertility and

predation risk drives a geometric sequence in the pattern of group sizes in baboons. *Biology*

letters, 14(3), 20170700.

Ezenwa, V. O., Gerardo, N. M., Inouye, D. W., Medina, M., & Xavier, J. B. (2012). Animal

behavior and the microbiome. *Science*, 338(6104), 198–199.

Goodrich, J. K., Waters, J. L., Poole, A. C., Sutter, J. L., Koren, O., Blekhman, R., ... & Spector,

T. D. (2014). Human genetics shape the gut microbiome. *Cell*, 159(4), 789-799.

Grieneisen, L. E., Livermore, J., Alberts, S., Tung, J., & Archie, E. A. (2017). Group Living and

Male Dispersal Predict the Core Gut Microbiome in Wild Baboons. *Integrative and Comparative*

Biology, 0(0), 1–16.

Hale, V. L., Tan, C. L., Niu, K., Yang, Y., Knight, R., Zhang, Q., ... & Amato, K. R. (2018). Diet

versus phylogeny: a comparison of gut microbiota in captive Colobine monkey

species. *Microbial ecology*, *75*(2), 515-527.

Hall, J. B., & Swaine, M. D. (2013). *Distribution and ecology of vascular plants in a tropical*

rain forest: forest vegetation in Ghana(Vol. 1). Springer Science & Business Media.

Henzi, S. P., Lycett, J. E., & Weingrill, T. (1997). Cohort size and the allocation of social effort

by female mountain baboons. *Animal Behaviour*, 54, 1235–1243

Huttenhower, C., Gevers, D., Knight, R., Abubucker, S., Badger, J. H., Chinwalla, A. T., … &

White, O. (2012). Structure, function and diversity of the healthy human microbiome. *Nature*,

486(7402), 207–214.

Kankam, B. O., & Sicotte, P. (2013). The effect of forest fragment characteristics on abundance

of Colobus vellerosus in the Forest-Savanna Transition Zone of Ghana. *Folia*

Primatologica, 84(2), 74-86.

Leamy, L. J., Kelly, S. A., Nietfeldt, J., Legge, R. M., Ma, F., Hua, K., ... & Pomp, D. (2014).

Host genetics and diet, but not immunoglobulin A expression, converge to shape compositional

features of the gut microbiome in an advanced intercross population of mice. *Genome*

biology, 15(12), 552.

Lombardo, M. P. (2008). Access to mutualistic endosymbiotic microbes: an underappreciated

benefit of group living. *Behavioral Ecology and Sociobiology*, 62(4), 479-497.

Markham, A. C., Gesquiere, L. R., Alberts, S. C., & Altmann, J. (2015). Optimal group size in a

highly social mammal. *Proceedings of the National Academy of Sciences*, 112(48), 14882-

14887.

McCord, A. I., Chapman, C. A., Weny, G., Tumukunde, A., Hyeroba, D., Klotz, K., ... & Leigh,

S. R. (2014). Fecal microbiomes of non-human primates in Western Uganda reveal species-

specific communities largely resistant to habitat perturbation. *American journal of*

primatology, 76(4), 347-354.

McMurdie, P. J., & Holmes, S. (2013). phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. *PloS one*, 8(4), e61217.

Moeller, A. H., Foerster, S., Wilson, M. L., Pusey, A. E., Hahn, B. H., & Ochman, H. (2016).

Social behavior shapes the chimpanzee pan-microbiome. *Science Advances*, 2(1), e1500997.

Morin, P. A., Chambers, K. E., Boesch, C., & Vigilant, L. (2001). Quantitative polymerase chain

reaction analysis of DNA from noninvasive samples for accurate microsatellite genotyping of

wild chimpanzees (Pan troglodytes verus). *Molecular ecology*, 10(7), 1835-1844.

Muegge, B. D., Kuczynski, J., Knights, D., Clemente, J. C., González, A., Fontana, L., ... &

Gordon, J. I. (2011). Diet drives convergence in gut microbiome functions across mammalian

phylogeny and within humans. *Science*, 332(6032), 970-974.

Perofsky, A. C., Lewis, R. J., Abondano, L. A., Di Fiore, A., & Meyers, L. A. (2017).
Hierarchical social networks shape gut microbial composition in wild Verreaux's sifaka. *Proc. R.*
Soc. B, 284(1868), 20172274.

Rollins, L.A., Browning, L.E., Holleley, C.E., Savage, J.L., Russell, A.F., Griffith, S.C. (2012).
Building genetic networks using relatedness information: a novel approach for the estimation of
dispersal and characterization of group structure in social animals. *Mol. Ecol.* 21, 1727–1740.

Saj, T. L., Teichroeb, J. A., & Sicotte, P. (2005). The population status of the ursine colobus
(Colobus vellerosus) at Boabeng-Fiema, Ghana. *Commensalism and conflict: the human primate*
interface (Paterson, JD & Wallis, J., eds). American Society of Primatologists, Norman, OK,
350-375.

Saj, T. L., & Sicotte, P. (2007). Scramble competition among Colobus vellerosus at Boabeng-
Fiema, Ghana. *International Journal of Primatology*, 28(2), 337-355.

Saj, T. L., & Sicotte, P. (2007). Predicting the competitive regime of female Colobus vellerosus
from the distribution of food resources. *International Journal of Primatology*, *28*(2), 315-336.

Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C.
(2011). Metagenomic biomarker discovery and explanation. *Genome biology*, 12(6), R60.

Sicotte, P., & Macintosh, A. J. (2004). Inter-group encounters and male incursions in Colobus
vellerosus in central Ghana. *Behaviour*, 141(5), 533-553.

Sicotte, P., Teichroeb, J. A., Vayro, J. V., Fox, S. A., Bădescu, I., & Wikberg, E. C. (2017). The

influence of male takeovers on female dispersal in Colobus vellerosus. *American journal of*

primatology, 79(7), e22436.

Snyder-Mackler, N., Alberts, S. C., & Bergman, T. J. (2014). The socio-genetics of a complex

society: female gelada relatedness patterns mirror association patterns in a multilevel

society. *Molecular ecology*, 23(24), 6179-6191.

Spor, A., Koren, O., & Ley, R. (2011). Unravelling the effects of the environment and host

genotype on the gut microbiome. *Nature Reviews Microbiology*, 9(4), 279.

Springer, A., Fichtel, C., Al-Ghalith, G. A., Koch, F., Amato, K. R., Clayton, J. B., ... &

Kappeler, P. M. (2017). Patterns of seasonality and group membership characterize the gut

microbiota in a longitudinal study of wild Verreaux's sifakas (Propithecus verreauxi). *Ecology*

and evolution, 7(15), 5732-5745.

Strum, S. C. (2012). Darwin's monkey: Why baboons can't become human. *American journal of*

physical anthropology, 149(S55), 3-23.

Sueur, C., & Maire, A. (2014). Modelling animal group fission using social network

dynamics. *PloS one*, 9(5), e97813.

Teichroeb, J. A., Wikberg, E. C., & Sicotte, P. (2009). Female dispersal patterns in six groups of

ursine colobus (Colobus vellerosus): infanticide avoidance is important. *Behaviour*, 146(4), 551-

582.

Teichroeb, J. A., Wikberg, E. C., & Sicotte, P. (2011). Dispersal in male ursine colobus monkeys

(Colobus vellerosus): influence of age, rank and contact with other groups on dispersal

decisions. *Behaviour*, 148(7), 765-793.

Teichroeb, J. A., & Sicotte, P. (2012). Cost-free vigilance during feeding in folivorous primates?

Examining the effect of predation risk, scramble competition, and infanticide threat on vigilance

in ursine colobus monkeys (Colobus vellerosus). *Behavioral Ecology and Sociobiology*, 66(3),

453–466.

Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J. C., Lynch, J., Grieneisen, L. E., ... & Archie,

E. A. (2015). Social networks predict gut microbiome composition in wild baboons. *Elife*, *4*,

e05224.

Turnbaugh, P. J., Hamady, M., Yatsunenko, T., Cantarel, B. L., Duncan, A., Ley, R. E., ... &

Egholm, M. (2009). A core gut microbiome in obese and lean twins. *Nature*, 457(7228), 480.

Wang, J. (2011). COANCESTRY: A program for simulating, estimating and analysing

relatedness and inbreeding coefficients. *Molecular Ecology Resources*, 11, 141–145.

Widdig, A., Nürnberg, P., Bercovitch, F. B., Trefilov, A., Berard, J. B., Kessler, M. J., ... &

Krawczak, M. (2006). Consequences of group fission for the patterns of relatedness among

rhesus macaques. *Molecular Ecology*, 15(12), 3825-3832.

Wikberg, E. C., Sicotte, P., Campos, F. A., & Ting, N. (2012). Between-Group Variation in

Female Dispersal, Kin Composition of Groups, and Proximity Patterns in a Black-and-White

Colobus Monkey (Colobus vellerosus). *PLoS ONE*, 7(11), 1–14.

Wikberg, E. C., Teichroeb, J. A., Bădescu, I., & Sicotte, P. (2013). Individualistic female

dominance hierarchies with varying strength in a highly folivorous population of black-and-

white colobus. *Behaviour*, 150(3-4), 295-320.

Wikberg, E. C., Ting, N., & Sicotte, P. (2014). Familiarity is more important than phenotypic

similarity in shaping social relationships in a facultative female dispersed primate, Colobus

vellerosus. *Behavioural processes*, 106, 27-35.

Wikberg, E. C., Ting, N., & Sicotte, P. (2014). Kinship and similarity in residency status

structure female social networks in black-and-white colobus monkeys (Colobus

vellerosus). *American Journal of Physical Anthropology*, 153(3), 365-376.

Wikberg, E. C., Ting, N., & Sicotte, P. (2015). Demographic Factors Are Associated with

Intergroup Variation in the Grooming Networks of Female Colobus (Colobus vellerosus).

International Journal of Primatology, 36(1), 124–142.

Wikberg, E.C., Christie, D.M., Sicotte, P., & Ting, N. (2017, August 21-27). Between and within

group variation in the gut microbiome of a black-and-white colobus monkey (*Colobus

vellerosus*). Paper presented at the *International Primatologcal Society Congress*.

Wikberg, E. C., Christie, D.M., Campos, F. A., Sicotte, P., & Ting, N. (2017). The link between

social networks and gut microbial composition in black-and-white colobus (Colobus

vellerosus). In *American Journal of Physical Anthropology*, 162(S64), 409-409.

Williams, B.L., Hornig, M., Parekh, T., Lipkin, W.I. (2012). Application of novel PCR-based

methods for detection, quantitation, and phylogenetic characterization of Sutterella species in

intestinal biopsy samples from children with autism and gastro-intestinal disturbances. mBio 3,

e00261–11.

Wong, S. N. P., & Sicotte, P. (2006). Population size and density of Colobus vellerosus at the

Boabeng-Fiema Monkey Sanctuary and surrounding forest fragments in Ghana. *American*

Journal of Primatology, 68, 465–476.

Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y. Y., Keilbaugh, S. A., ... & Sinha, R.

(2011). Linking long-term dietary patterns with gut microbial enterotypes. *Science*, 334(6052),

105-108.

Yildirim, S., Yeoman, C. J., Sipos, M., Torralba, M., Wilson, B. A., Goldberg, T. L., ... &

Nelson, K. E. (2010). Characterization of the fecal microbiome from non-human wild primates

reveals species specific microbial communities. *PloS one*, 5(11), e13963.

492   **TABLES**

	Males			Females			Total
	Adults + Subadults	Juveniles	Infants	Adults + Subadults	Juveniles	Infants	
Prefission DA	9	0	2	12	4	1	28
Postfission DA	5	2	1	7	3	0	18
Postfission NP	1	0	2	6	0	1	10

494   Table 1. Composition of the different social groups before and after the fission event. Although

495   this study only examined the gut microbial compositions of females, males are included in this

496   table for insight into group composition overall.

497   **FIGURES**

499   Figure 1. Home range distributions of the DA and NP groups prior to and after the fission event.

500   (A) The original DA group maintained a large home range in the summer of 2006. (B) The two

501   product groups split the original home range by the summer of 2007.

Figure 2. Postfission group membership predicts Bray-Curtis dissimilarity of the NP and DA

groups. No significant difference in gut microbial diversity was observed prior to the fission

event ($p = 0.085$, n = 12, pseudo F = 1.23), while less than nine months after the fission event

these groups showed unique microbial signatures ($p = 0.02$, n = 12, pseudo F = 1.62). Significant

differences were found in gut microbial diversity between all groups across the years sampled (p

< 0.01, n = 24, pseudo F = 2.19).

512   Figure 3. Pairwise-relatedness as calculated using 17 STR loci are represented by group. On

513   average, pairwise-relatedness increased by 0.079 in the NP social group from the original group.

514   However, the pairwise-relatedness between groups was not statistically different (p=0.103, df=2,

515   χ^2=4.55).

516
517

519   Figure 4. LDA scores for taxa differing significantly between product group. Three genera were

520   found in different relative abundance between these two groups. Linear effect size analysis

521   (LefSe) was run for the groups at a KW alpha value of 0.01 and an LDA score of 3.0.