Experimentally-constrained biophysical models of tonic and burst firing modes in

thalamocortical neurons

Elisabetta Iavarone[1*], Jane Yi[2], Ying Shi[1,2], Bas-Jan Zandt[1], Christian O'Reilly[1], Werner Van Geit[1],

Christian A. Rössert[1], Henry Markram[1,2], and Sean L. Hill[1,3]

[1]Blue Brain Project, École polytechnique fédérale de Lausanne (EPFL), Geneva, Switzerland

[2]Laboratory of Neural Microcircuitry (LNMC), École polytechnique fédérale de Lausanne (EPFL)

(EPFL), Lausanne, Switzerland

[3]Krembil Centre for Neuroinformatics, CAMH, Toronto, Canada

* Corresponding author

E-mail: elisabetta.iavarone@epfl.ch

Abstract

Somatosensory thalamocortical (TC) neurons from the ventrobasal (VB) thalamus are central components in the flow of sensory information between the periphery and the cerebral cortex, and participate in the dynamic regulation of thalamocortical states including wakefulness and sleep. This property is reflected at the cellular level by the ability to generate action potentials in two distinct firing modes, called tonic firing and low-threshold bursting. Although the general properties of TC neurons are known, we still lack a detailed characterization of their morphological and electrical properties in the VB thalamus. The aim of this study was to build biophysically-detailed models of VB TC neurons explicitly constrained with experimental data from rats. We recorded the electrical activity of VB neurons (N = 49) and reconstructed morphologies in 3D (N = 50) by applying standardized protocols. After identifying distinct electrical types, we used a multi-objective optimization to fit single neuron electrical models (e-models), which yielded multiple solutions consistent with the experimental data. The models were tested for generalization using electrical stimuli and neuron morphologies not used during fitting. A local sensitivity analysis revealed that the e-models are robust to small parameter changes and that all the parameters were constrained by one or more features. The e-models, when tested in combination with different morphologies, showed that the electrical behavior is substantially preserved when changing dendritic structure and that the e-models were not overfit to a specific morphology. The models and their analysis show that automatic parameter search can be applied to capture complex firing behavior, such as co-existence of tonic firing and

35 low-threshold bursting over a wide range of parameter sets and in combination with different

36 neuron morphologies.

Author summary

Thalamocortical neurons are one of the main components of the thalamocortical system,

which are implicated in key functions including sensory transmission and the transition between

brain states. These functions are reflected at the cellular level by the ability to generate action

potentials in two distinct modes, called burst and tonic firing. Biophysically-detailed

computational modeling of these cells can provide a tool to understand the role of these neurons

within thalamocortical circuitry. We started by collecting single cell experimental data by

applying standardized experimental procedures in brain slices of the rat. Prior work has

demonstrated that biological constraints can be integrated using multi-objective optimization to

build biologically realistic models of neuron. Here, we employ similar techniques as those

previously employed, but extend them to capture the multiple firing modes of thalamic neurons.

We compared the model results with additional experimental data test their generalization and

quantitatively reject those that deviated significantly from the experimental variability. These

models can be readily integrated in a data-driven pipeline to reconstruct and simulate circuit

activity in the thalamocortical system.

Introduction

Introduction

Thalamocortical (TC) neurons are one of the main components of the thalamus and have

been extensively studied *in vitro* and *in computo*, especially in first order thalamic nuclei in

different species (1). One of these nuclei, namely the ventral posterolateral nucleus (VPL), relays

somatosensory, proprioceptive, and nociceptive information from the whole body to the

somatosensory (non-barrel) cortex (2). The VPL is located close to ventral posteromedial nucleus

(VPM), which transmits information from the face to the barrel cortex. The VPL and VPM nuclei

constitute the ventrobasal (VB) complex of the thalamus (3).

Despite its key role in sensory functions, a systematic characterization of the cellular

properties of the VB complex is still missing. The morphologies of VPL neurons in adult rats were

described in early anatomical studies but were limited to two-dimensional drawings of Golgi-

impregnated cells (4). The general electrical properties of TC neurons maintained *in vitro* are

known and similar in different thalamic nuclei and species with respect to the generation of two

distinct firing modes, called tonic firing and low-threshold bursting (5–8). However, a systematic

description on the electrical types in the VB thalamus in the rodents is still missing.

Collecting morphological and electrophysiological data, by following standardized

experimental procedures, is essential for the definition of cells types and it is the first step to

constraining computational models of single neurons (9,10). Although models of TC neurons have

already been previously published, they typically were aimed at studying specific firing properties

and their parameters were hand tuned to achieve the desired result (11–15).

The purpose of our study is to systematically define the morphological and electrical types

by collecting *in vitro* experimental data and to constrain biophysically detailed models of VB TC

neurons of the juvenile rat. To the best of our knowledge, automatic parameter search has not

been applied, thus far, to capture complex firing behavior in thalamic neurons, in particular low-

threshold bursting and tonic firing. We defined the electrical and morphological types of TC

neurons through in vitro patch-clamp recordings and 3D morphological reconstructions. We then

extended an existing method (16) to account for their distinctive firing properties. These

electrical models (e-models) were constrained by the electrical features extracted from

experimental data (9,17,18). Other experimental data were used to assess the generalization of

the models to different stimuli and morphologies. We further performed a sensitivity analysis by

varying each parameter at a time by a small amount and recording the resulting electrical

features. This analysis provides an assessment of the robustness of the models and a verification

that the selected features provide sufficient constraints for the parameters.

Results

Physiological and morphological characterization

We characterized TC neurons in slices of the rat VB thalamus, by combining whole-cell patch-

clamp recordings, biocytin filling and 3D Neurolucida (MicroBrightField) reconstruction, along

with anatomical localization in a reference atlas (19) (Fig 1).

Figure 1: Simultaneous physiological and morphological characterization.

(A) View of a patched cell under optic microscope and anatomical localization of biocytin-filled

neurons (insets) in the rat Paxinos and Watson atlas (19). Letters D and E identify morphologies

in a slice. (B) Voltage responses of two different thalamocortical (TC) neurons to a standardized

battery of current stimuli. Each current amplitude was normalized by the threshold current of

each neuron (e.g. 150 % threshold, see Methods). Third row is a low-threshold burst response

from a hyperpolarized holding potential, V_{hold} = −84 mV (burst mode), the other responses are

elicited from a depolarized holding potential, V_{hold} = −64 mV (tonic mode). Two different holding

currents (I_{hold} - tonic, I_{hold} - burst) are injected to obtain the desired V_{hold}. The vertical scale bar

applies to all the traces, the first horizontal scale bar from the top refers to the first two rows,

the second applies to the last four rows. (C) Analysis of adaptation index (AI) from recordings in

tonic mode. Solid line is a non-parametric estimation of the distribution, dashed lines are two

Gaussian distributions fitted to the data (see Methods). The vertical line indicates the cut-off

value.

Visual inspection of 50 reconstructed morphologies (24 from the VPL, 26 from the VPM

nuclei) revealed variability in the number of principal dendritic trunks and their orientation, in

agreement with previous anatomical studies (4).

Figure 2: Morphological properties.

(A) Renderings of 3D reconstructed TC neurons along with their persistence barcode according

to (20). Grey: soma and dendrites, blue: axon only small sections available). The persistence

barcode is a topological description of the branching pattern of the neurons' dendrites. (B) Sholl

analysis of TC neuron dendrites. For each Sholl ring, the number of intersections is shown (mean

± standard deviation, N = 50). Each grey circle represents one morphology, colored lines

correspond to the morphologies in A. See Fig S1 for further analysis.

The maximum radial extent of the dendrites ranged between 120 and 200 μm and they

started to branch between 20 and 50 μm from the soma (Fig S1). We then analyzed the

morphologies with two methods in order to quantitavely classify different morphological types.

We used algebraic topology to extract the persistent homology of each morphology and to

visualize the persistence barcode (20) (Fig 2A, see Methods). Each horizontal bar in the

persistence barcode represents the start and end point of each dendritic component in terms of

its radial distance from the soma. The barcodes of all the morphologies followed a semi-

continuous distribution of decreasing length. To quantify the differences between the barcodes,

we computed the pairwise distances of the persistence images (see Methods and Fig S1). We

found that they were in general small (<0.4, values expected to vary between 0 and 1). These

findings indicate that the morphologies cannot be grouped in different classes based on the

topology of their dendrites. Furthermore, we performed Sholl Analysis (21) to compare the

complexity of the dendritic trees (Fig 2B). We observed that all the morphologies had dense

dendritic branches, with a maximum number of 50-100 intersections between 50-80 μm from

the soma. When comparing the Sholl profiles for each pair of neurons we could not find any

statistically significant difference (Fig S1C). Considering the results of topological and Sholl

analyses, we grouped all the morphologies in one morphological type (m-type) called

thalamocortical (TC) m-type.

We used an adaptive stimulation protocol, called e-code, consisting of a battery of current

stimuli (e- code, see Methods for details), where the stimulation amplitude was adapted to the

excitability of different neurons. This standardized protocol has previously been used to build

biophysically-accurate models of cortical electrical types (e-types) (16). However, TC neurons

from different thalamic nuclei and species fire action potentials in two distinct firing modes,

namely tonic firing, when stimulated from a relatively depolarized membrane potential or low-

threshold bursting, from a hyperpolarized membrane potential (5). We thus extended the e-code

to include two different holding currents. All the neurons recorded in this study displayed tonic

and burst firing, when stimulated with the appropriate holding current (Fig 1). Moreover, we

were able to classify different e-types by considering the voltage traces recorded in tonic mode

in response to step current injections (Fig 1). The majority of the cells (59.3 %) showed a non-

adapting tonic discharge (continuous non-adapting low-threshold bursting, cNAD_ltb e-type)

while others (40.7 %) had higher adaptation rates (continuous non-adapting low-threshold

bursting, cAD_ltb e-type), as reflected by the adaptation index (Fig 1C). We followed the Petilla

convention (22) for naming the tonic firing discharge (cNAD or cAD), extending it to include "_ltb"

for the low-threshold bursting property. In some rare examples, we noticed acceleration in the

firing rate with decreasing inter-spike intervals (ISIs) towards the end of the stimulus. Similar

adapting and accelerating responses have already been described in the VB thalamus of the cat

(7). We also observed stereotypical burst firing responses within the same cell, with variation of

the number of spikes per burst in different cells, but the burst firing responses alone were

insufficient to classify distinct e-types.

Constraining the models with experimental data

Multi-compartmental models comes with the need of tuning a large number of parameters

(23), therefore we constrained the models as much as possible from experimental data. We first

combined the morphology and the ionic currents models in the different morphological

compartments (soma, dendrites and axon). Given that the reconstruction of the axon was

limited, we replaced it with a stub representing the initial segment (16). We used previously

published ionic current models and selected those that best matched properties measured in rat

TC neurons (see Methods). The kinetics parameters were not part of the free parameters of the

models. The distribution of the different ionic currents and their conductances in the dendrites

of TC neurons is largely unknown. The current amplitudes of the fast sodium, persistent and

transient (A-type) potassium currents were measured, but only up to 40-50 µm from the soma

(24). Indirect measures of burst properties (15) or Ca^{2+} imaging studies (25) suggest that the low-

threshold calcium (T-type) channels are uniformly distributed in the somatodendritic

compartments. We thus assumed different peak conductance in the soma, dendrites and axon

for all the ionic currents, except for I_{CaT}, which had the same conductance value in the soma and

dendrites. We then extracted the mean and standard deviation (STD) of different electrical

features in order to capture the variability of firing responses from different cells of the same e-

type (9) (Fig 3).

Figure 3: Histograms of electrical features.

Each vertical line represents the mean feature value for a cell. Tonic and burst refer to the holding

voltage as in Fig 1. (A) Feature values extracted from recordings in tonic mode (N = 11 cAD_ltb

cells, N = 16 cNAD_ltb cells). The features highlighted by a black box show different distributions

for the cNAD_ltb and cAD_ltb electrical types (e-types) (p-value<0.05, two- sided Mann-Whitney

U test with Bonferroni correction for multiple comparisons). Passive properties (V_{rest}, R_{input}) and

spike shape features (AHP depth, AP amp., etc.) did not show clear differences between the two

e-types. (B) Features measuring burst firing properties (N = 22 cells).

We observed that some features extracted from tonic firing responses had distinct

distributions between the cAD_ltb and cNAD_ltb e-types (Fig 3A). The features were chosen in

order to quantify salient physiological properties of TC neurons and to constrain the parameters

197   of the model, namely the peak conductance of each ionic current. The average value and STD of

198   the features were used as optimization objective (multi-objective optimization, MOO). Twenty-

199   five parameters were allowed to vary between the upper and lower bounds shown in Fig 5. The

200   models were associated with a training error, i.e. a set of all the feature errors (measured as

201   absolute z-scores) (9,18,26).

202
Figure 4: Models of different TC e-types and their fitting errors.

(A) Single neuron modelling pipeline. (B) Experimental and model voltage responses to a variety

of stimuli pattern used during the optimization of cNAD_ltb and cAD_ltb e-types. (C) Feature

errors of the models shown in (B) reported as deviation from the experimental mean. The models

are compared with the mean of features shown in Fig 3. Note that the models shown in B are

fitted in order to reproduce the mean firing properties, not only a specific experimental

recording. See Fig S2 for a complete list of fitting errors. By applying this MOO procedure, we

generated multiple models with distinct parameter combinations that reproduced tonic and low-

threshold burst firing in cNAD_ltb and cAD_ltb e-types (Fig 4).

Model and experimental diversity

We found that different sets of parameter values reproduced the target firing behavior (Fig

5B). We further analyzed models that had all the feature errors below 3 STD. Models' voltage

responses reflected the characteristic firing properties of TC neurons (Fig S3), indicating that the

selected set of features were sufficient to capture the two firing modes, in both the adapting and

non-adapting e-types. The voltage traces from different models showed small differences in spike

amplitude, firing frequency, and depth of the after-hyperpolarization, as reflected by the

variability of features values (Fig 5C).

222

223 **Figure 5: Diversity of model parameters and experimental variability.**

224 (A) Example of model fitting errors (sum of all feature errors) during optimization. (B) Initial

225 parameter ranges and diversity of solutions. Each vertical line represents the range for the

226 parameters, when the horizontal lower bar is missing the bound is 0. The characters following ".".

227 in the parameter name specifies the morphological compartment for the parameter ("s": soma,

228 "d": dendrites, "a": axon). Black circles: parameter values for one of the models in Fig 4, grey

229 circles: parameter values of the models with all feature errors below 3 STD. (C) Features

230 variability in the models and experiments. Blue crosses: feature errors of a sample of 10 models.

231 Each grey circle is the z-scored feature value of one experimental cell, obtained from the feature

232 values shown in Fig 3. The protocol names are shown in parenthesis and corresponds to the

233 stimuli shown in Fig 1 and Fig 4, tonic and burst refer to the holding current as in Fig 1.

234

Spike-shape related features (e.g. AP. amplitude) in the different models covered the space

of the experimental variability, while for some features (e.g. input resistance, R_{input}), all models

tended to cluster on one of the tails of the experimental distribution. R_{input} relates to the neuron

passive properties and depends both on the number of channels open at rest (inverse of the leak

conductance in the model) and the size of the cell. Given that all the models were constrained

on a single morphology, this result is not surprising. The number of action potentials (Num. of

APs) in different conditions (No stim, I_{hold}) ensured that the models did not spike in the absence

of a stimulus or in response to the holding current. For this reason, all the experimental and

model feature values in 5C are equal to 0. Other features, such as latency to the first spike and

sag amplitude were less variable in the models compared to experiments. We hypothesized that

this depended on the variable stimulation amplitudes applied to different experimental cells,

while all the models were stimulated with the same current amplitudes.

We examined the diversity of the parameter values with respect to the initial parameter

range (Fig 5B). Most of the optimized parameter values spanned intervals larger than one order

of magnitude. On the other hand, some parameter values were restricted to one order of

magnitude, for example the permeability of the low-threshold calcium current P_{CaT}. This result is

in agreement with experiments showing a minimum value of I_{CaT} is critical to generate burst

activity and this critical value is reached only at a certain postnatal age (27). The value of P_{CaT} was

constrained by features measuring burst activity (such as number of spikes, frequency, etc.).

Assessment of model generalization

We used different stimuli for model fitting (current steps) and for generalization

assessment (current ramps and noise). We simulated the experimental ramp currents in-silico,

by stimulating the models with the appropriate holding currents for the two firing modes and a

linearly increasing current. We first compared visually the model responses with the

experimental recordings (Fig 6A).

Figure 6: Model generalization.

(A) Responses to a ramp current injection in burst mode (left) and tonic mode (center). (B)

Responses to a noise current generated according to an Ornstein-Uhlenbeck process and scaled

based on the excitability of the different experimental cells and models (see Methods). (C)

Generalization errors for all the models that passed the generalization test (all generalization

errors <3 STD). (D) Proportion of models that passed the generalization test (see Fig S4 for

examples of models that failed this test).

In burst mode, the models reproduced the different behaviors observed experimentally:

absence of a burst, small low-threshold spike, burst, burst followed by tonic firing (Fig S4).

Moreover, the latency of burst generation substantially overlapped with the experimental one.

However, a small fraction of models (1.2 %) generate repetitive burst that we have never

observed in the experimental recordings (Fig S4). These models were quantitatively rejected by

considering the number of spikes and the inter-spike intervals. In tonic mode, the latency to first

spike, the voltage threshold, the shape of the subsequent action potentials and the increase in

firing frequency were comparable with the experimental recordings (Fig 6A). In addition, we

quantified the generalization error to ramp stimuli (Fig 6C), by considering the latency to first

spike, firing frequency increase over time (tonic mode) or number of spikes (burst mode).

Although conductance-based models can be fit by using step and ramp currents (26), these

stimuli are different from synaptic inputs, which can be simulated by injecting noisy currents. To

test the response to such network-like input, we used a noisy current varying accordingly to an

Ornstein-Uhlenbeck (OU) process (28) to compare models' responses with the experimental

data. Each experimentally recorded cell was stimulated with the same OU input, scaled by a

factor *w*. Experimentally, *w* was calculated during the experiment by evaluating the responses to

previous stimuli. We developed a similar approach to generate the noise stimuli in silico (see

Methods). The noise current was injected on top of the holding currents used during the

optimization. We found that the models reproduced well the subthreshold potential, spike times

and the distribution of single spikes and bursts (Fig 6B). Moreover, we quantitatively evaluated

the generalization to the noise stimulus by extracting features (e.g. number of spikes) and

comparing them with the experimental mean.

We computed generalization errors for each model, which were calculated similarly to the

optimization errors (Fig 6C). We considered a model acceptable after generalization if it had all

generalization errors <3 STD and we found that the majority of the models (>90%) passed the

generalization test.

Sensitivity of electrical features to small parameter perturbations

We assessed the robustness of the models to small changes in their parameter values. To

that end, we varied each parameter at a time by a small amount (± 2.5 % of the optimized value)

and computed the values of the features. A sensitivity value of 2 between parameter p and

feature y means that a 3 % change in p caused a 6 % change in f. We ranked the parameters from

the most to the least influential and the features from the most sensitive to the least sensitive.

Figure 7: Local sensitivity analysis.

(A) Sensitivity of the feature values to small changes to the parameter values for the cAD_ltb

model in Fig 4. Sensitivities (Δy/Δp) are color coded as a heat map. Features are ranked from the

most to the least sensitive and parameters are ranked from the most to the least influential. The

last three rows are features that ensure that the models were not firing without input or in the

response to the holding current. Small changes to the parameter values are not expected to make

the model firing and thus the sensitivity of these features is 0. (B) Same sensitivity values as in

(a), with features and parameters clustered by similar sensitivity and influences.

The conductance of the leak current g_{leak} emerged as the most influential parameter (Fig 7A). An

increase in g_{leak} caused a decrease in firing frequency (inverse of inter-spike intervals, ISIs) in both

the tonic and burst firing modes. These results are easy to interpret when considering Ohm's law:

increasing g_{leak} means decreasing the input resistance of the model, so that for the same input

current the voltage response becomes smaller. The second most influential parameter was the

conductance of the persistent sodium current g_{NaP} in the dendrites, which increased the tonic

firing rate as expected from a depolarizing current and had an effect on the late phase of the low-

threshold burst (inverse last ISI - burst). An increase in the permeability of the low-threshold

calcium current P_{CaT}, known to be one the main currents underlying low threshold bursting,

enhanced burst firing responses (it decreased the inverse of ISIs) and had effects on some of the

tonic features. P_{CaT} was the third most influential parameter. These findings show that I_{CaT} is the

main driver of the low-threshold burst, but other currents, such as I_{NaP} contributes as well.

Increasing the dendritic permeability of the high threshold calcium current P_{CaL} decreased the

tonic firing rate, despite being a depolarizing current. Increasing P_{CaL} means higher Ca^{2+} influx and

higher amplitude of the Ca^{2+}-activated potassium current (I_{SK}). The parameter g_{SK} had indeed a

similar effect on the features and thus clustered together with P_{CaL} (Fig 7B). Increasing the

conductance of the transient sodium conductance g_{NaT} increased action potential amplitude and

decreased its duration. Sag amplitude, that is known to depend on the activity of I_H, was mainly

influenced by change in g_{leak}, P_{CaT} and g_H. In summary, each parameter influenced at least one

feature. Some features were weakly influenced by small parameter changes, e.g. baseline

voltage, which depend more on the holding current amplitude, than on the model parameters.

These results indicate that the model ability to generate tonic and burst firing is robust to small

changes in parameter values and that all the parameters were constrained during the

optimization by one or more features.

We then analyzed which features depended similarly on parameter changes, as they may

add superfluous degrees of freedom during parameters search. Fig 7B shows the same

sensitivities as in Fig 7A, clustered by their similarities (see Methods). Features clustered together

if they were sensitive to similar parameter combinations and parameters clustered based on their

similar influence on the features. Not surprisingly, the same tonic features measured at different

level of current stimulation clustered together (e.g. AP amplitude and half-width, AHP depth,

latency of the first ISI) and tonic firing features belonged to a cluster that was different from burst

features.

Preservation of model firing properties with different morphologies

We optimized the parameters for the adapting and non-adapting e-models in combination with

two different experimental morphologies and then tested them with the other 48 morphologies.

Considering that morphologies could not be classified in different m-types based on topological

analysis of their dendrites and that TC neurons have been shown to be electrically compact (15),

we expected the electrical behavior to be conserved when changing morphology. Nonetheless,

different neurons vary in their input resistance R_{input} and rheobase current I_{thr} due to variation in

the surface area. Variation in R_{input} and I_{thr} made the current amplitude applied during the

optimization inadequate to generate the appropriate voltage trajectories. We thus devised an

algorithm to search for the holding current to obtain the target holding voltage (for example −64

mV or −84 mV for tonic and burst firing, respectively) and I_{thr} from the desired holding voltage.

The different e-model/morphology combinations (me-combinations) were evaluated by

computing the same feature errors calculated during optimization. For each morphology, we

selected the e-model that generated the smallest maximum error. All me-combinations

reproduced burst and tonic firing (Fig 8C). However, two me-combinations generated responses

with a small number of features that deviated from the experimental mean. We chose the value

of 3 STD as a threshold to define which me-combinations were acceptable (29), yielding 48

acceptable me-combinations out of the 50 tested (Fig 8A). We analyzed more closely which

features were significantly different from the experimental mean. In Fig 8B we show that the

rejected me-combinations had too many action potentials in the burst.

Figure 8: Model generalization to different experimental morphologies.

(A) Feature errors from the best electrical models (e-model) showed in Fig 4 applied to 50

different TC cell morphologies. Each morphology is represented with a different color. E-

models/morphology combinations with at least one feature error > 3 STD (dashed line) were

rejected. (B) Example of voltage responses from an accepted and from a rejected e-

model/morphology combination. Feature errors for the rejected combination are shown in red

in (A) and are indicated on the voltage trace.

Discussion

Our objective was to apply and extend an existing data-driven pipeline to identify the cell

types and build models of VB thalamocortical neurons that reproduce the multiple firing modes

that have been experimentally observed. We successfully modelled these novel firing types, by

including additional stimulation protocols and features to constrain the low-threshold burst.

Our morphological and electrical data were used to define the properties of VB TC neurons

in the rat. We found two electrical types (e-types) of TC neurons, but no objectively different

morphological types (m-types) were revealed either using Sholl analysis (21) or topological

analysis of dendritic branching (20). We cannot exclude that refinements to these methods will

reveal different m-types similar to the ones described in the visual thalamus of the mouse (30).

We also showed that automatic parameter search can be applied to build biophysically and

morphologically detailed models. This method was already applied to model canonical firing

behavior in cortical, hippocampal and cerebellar granule neurons (9,10,16,17,31,32). To the best

of our knowledge, such an automatic parameter search has not previously been used to capture

different firing modes and complex firing behavior such as low-threshold bursting in thalamic

neurons. Standardized electrophysiological protocols allowed us to identify for the first time in

juvenile rat adapting and non-adapting e-types of TC VB neurons that were previously observed

in other species (7). This finding suggests that the intrinsic properties of TC neurons contribute to

adaptation, a key phenomenon for filtering out irrelevant stimuli, before sensory information

reaches the neocortex. Further experiments are needed to elucidate the relative contribution of

intrinsic mechanisms and network properties to adaptation in somatosensory systems. We

named the two main e-types continuous non-adapting low-threshold bursting (cNAD_ltb) and

continuous adapting low-threshold bursting (cAD_ltb) by following and extending existing

conventions (16,22,31).

In this study, we improved upon previous morphologically and biophysically detailed

models of tonic and burst firing in TC neurons (12,13,15) by explicitly constraining the parameters

with experimental data, without hand-tuning of parameter values. Unlike previous models, we

chose a multi-objective optimization for a methodological and a scientific reason: it is more time-

efficient, reproducible, and it approximates the variability in ionic channel expression of

biological neurons (31,33–35), as shown by the family of acceptable solutions we found.

However, experiments aimed at quantifying ion channel conductances are essential to assess if

these solutions fall between biological ranges. Furthermore, we tested the generalization

capability of the models and found that more than 90% of the models were comparable with the

experimental data.

Nonetheless, we noticed some inaccuracies when comparing the voltage traces with the

experimental data when assessing the generalization of some models. For instance, some models

tended to generate small transient oscillations in response to ramp stimuli in burst mode. This

result is not surprising, considering that the exact kinetics for all the ionic currents are not

available and that there are known limitations in models of ionic channels derived from the

literature or from other models (36,37). In particular, modifications of the kinetics of the low-

threshold calcium current was shown to explain the propensity to generate oscillatory bursts in

TC neurons of other nuclei and species (38).

TC neurons have been shown to be electrically compact (15) and could, in principle, be

modeled as a single compartment. However, active mechanisms need to be located in the

dendrites in order to ensure synaptic integration and amplification (39). Information regarding

specific conductances or firing properties in the dendrites of TC neurons is limited. For this

reason, dendritic parameters in our models may be underconstrained. However, the sensitivity

analysis (see below) revealed that dendritic parameters did not appear to be the least

constrained because they influenced different tonic and burst-related features.

We included in the model fitting and validation pipeline a sensitivity analysis, which is often

neglected in computational neuroscience (40). Although we cannot use our simple univariate

approach to explore multidimensional parameter correlations and principles of co-regulation of

ion channels expression, it is useful to find better constraints for parameters optimization. The

selection of the features is indeed a step that still requires care and experience by modelers.

Furthermore, this type of sensitivity analysis allows to identify parameters that can be traded-off

during the optimization and that can be removed in order to reduce the dimensionality of the

problem. In our study, four parameters related to the calcium dynamics were shown to influence

the features in a very similar fashion. This type of analysis is of particular importance in future

work aimed at using the full diversity of ion channels that can be inferred from gene expression

data. More in detail, we propose that sensitivity analysis should be a fundamental tool in

selecting which conductances are successfully optimized by the available experimental

constraints. The example we showed is a local approach, applied to a specific solution to the

optimization problem, which showed that our models are robust to small parameter changes.

This analysis can be extended to study how the sensitivities vary in the neighborhood of different

solutions.

In conclusion, we systematically studied the morphological and electrical properties of VB

TC neurons and used these experimental data to constrain single neuron models, test their

generalization capability and assess their robustness. Further work will validate these models in

response to synaptic activity, in order to include them in a large-scale model of thalamocortical

microcircuitry (16).

Methods

Experimental procedures

Experimental data were collected in conformity with the Swiss Welfare Act and the Swiss

National Institutional Guidelines on Animal Experimentation for the ethical use of animals. The

Swiss Cantonal Veterinary Office approved the project following an ethical review by the State

Committee for Animal Experimentation.

All the experiments were conducted on coronal or horizontal brain slices (300 μm thick-

ness) from the right hemisphere of male and female juvenile (P14-18) Wistar Han rats. The region

of interest was identified using the Paxinos and Watson rat brain atlas (19). After decapitation,

brains were quickly dissected and sliced (HR2 vibratome, Sigmann Elektronik, Germany) in ice-

cold standard ACSF (in mM: NaCl 125.0, KCl 2.50, $MgCl_2$ 1.00, NaH_2PO_4 1.25, $CaCl_2$ 2.00, D-(+)-

Glucose 50.00, $NaHCO_3$ 50.00; pH 7.40, aerated with 95% O_2 / 5% CO_2). Recordings of

thalamocortical neurons in the VB complex were performed at 34 °C in standard ACSF with an

Axon Instruments Axopatch 200B Amplifier (Molecular Devices, USA) using 5–7 MΩ borosilicate

pipettes, containing (in mM): K^+-gluconate 110.00, KCl 10.00, ATP-Mg^{2+} 4.00, Na_2-

phosphocreatine 10.00, GTP-Na^+ 0.30, HEPES 10.00, biocytin 13.00; pH adjusted to 7.20 with

KOH, osmolarity 270-300 mOsm. Cells were visualized using infrared differential interference

contrast video microscopy (VX55 camera, Till Photonics, Germany and BX51WI microscope,

Olympus, Japan).

Membrane potentials were sampled at 10 kHz using an ITC-18 digitizing board (InstruTECH,

USA) controlled by custom-written software operating within IGOR Pro (Wavemetrics, USA).

Voltage signals were low-pass filtered (Bessel, 10 kHz) and corrected after acquisition for the

liquid junction potential (LJP) of −14 mV. Only cells with a series resistance <25 MΩ were used.

After reaching the whole-cell configuration, a battery of current stimuli was injected into

the cells and repeated 2-4 times (e-code). During the entire protocol, we defined offset currents

in order to keep the cell at −50 mV (tonic firing) or −70 mV (burst firing) before LJP correction and

applied them during the entire protocol. The step and ramp currents were injected with a delay

of 250 ms in the experiment. In the models, the stimuli were injected with a delay of 800 ms, to

allow for the decay of transients due to initialization. Each stimulus was normalized to the

rheobase current of each cell, calculated on-line as the current that elicited one spike (stimulus

TestAmp, duration 1350 ms). The stimuli used for in the experiments, for fitting and testing the

models were:

• IDRest: current step of 1350 ms, injected at different amplitude levels in 25 % increments

(range 50-300 % threshold). IDRest was renamed to Step in the model.

• IDThresh: current step with duration of 270 ms, 4 % increments (range 50 - 130 %).

• IV: hyperpolarizing and depolarizing steps of 3000 ms injected in 20 % increments (range

−140 - 60%).

• SponNoHold: the first 10 seconds of this stimulus was used to calculate the resting

membrane potential. No holding or stimulation currents were applied.

• SponHold: the first 10 seconds of this stimulus was used to calculate the holding current

applied to keep the cells at the target potential.

• PosCheops: ramps of current from 0 to 300 % and from 300 to 0 % having progressively

shorter durations (4000 ms, 2000 ms, 1250 ms). To test the models in tonic mode we used

the first increasing ramp in the stimulus, while we used the last one in the bursting firing

mode. We chose the last one because the biological cells were more likely to generate a

burst.

• NOISEOU3: the original wave was scaled and offset for each cell based on the spike

frequency responses to IDRest responses. The scaling factor w was extracted from the

frequency-current curve and corresponded to the current value that made the cell fire at

7.5 Hz.

Neurons that were completely stained and those with high contrast were reconstructed in

3D and corrected for shrinkage as previously described (41). Reconstruction used the

Neurolucida system (MicroBrightField). The location of the stained cells was defined by

overlaying the stained slice and applying manually an affine transformation to the Paxinos and

Watson's rat atlas (19).

Electrical features extraction

Electrical features were extracted using the Electrophys Feature Extraction Library (eFEL)

(42). We calculated the adaptation index (AI) from recordings in tonic mode (Step 200 %

threshold) and classified TC VB neurons into adapting (AI>=0.029) and non-adapting (AI<0.029)

electrical types. AI was calculated using the eFEL feature *adaptation_index2* and corresponded

to the average of the difference between two consecutive inter-spike intervals (ISI) normalized

by their sum. The cut-off value was calculated after fitting a Gaussian mixture model to the

bimodal data, using available routines for R (43,44). In order to group data from different cells

and generate population features, we normalized all the stimuli by the rheobase current I_{thr} of

each cell. To calculate I_{thr}, we used IDRest and IDThresh and selected the minimal amplitude that

evoked a single spike. The extracted features quantified passive (input resistance, resting

membrane potential), burst and tonic firing properties (number of spikes, inverse of inter-spike

intervals, latency to first spike), action potentials shape (amplitude, half-width, depth of the fast

after-hyperpolarization). We aimed at finding the minimal set of features that capture the most

important properties. This set was a trade-off between comprehensively describing the

experimental data (i.e. extracting all possible features), which can lead to over-fitting and loss of

generalizability, and a too small set that would miss some important characteristics. For the tonic

firing responses, we used three stimulation amplitudes (150 %, 200 %, 250 % of firing threshold)

which have been shown to reproduce the complete input-output function of the neurons (17,41).

Responses to two hyperpolarizing steps of different amplitudes (−40 % and −140 % threshold)

constrained the input resistance and the conductance of currents activated in hyperpolarization

(*sag_amplitude* feature). We included baseline voltage values in the optimization objectives to

ensure that the model was in the right firing regime and spike count to penalize models that were

firing in response to the holding currents. Along with the voltage features, we extracted mean

holding and threshold current values for all the experimental stimuli. Description of the features

and the details on their calculation are available on-line [49]. Current stimuli applied during the

optimization and generalization were directly obtained from the experimental values or

automatically calculated by following the experimental procedures (e.g. noise stimulus).

Morphology analysis

Reconstructed morphologies were analyzed to objectively identify different morphological

types. The Sholl profiles of each pair of cells was statistically tested by using k-samples Anderson-

Darling statistics. This test was preferred to the most common Kolmogorov- Smirnov test,

because it does not assume that the samples are drawn from a continuous distribution. The

different Sholl profiles are indeed an analysis of the intersections with discrete spheres.

To compare the topological description of each morphology we transformed the

persistence barcodes into persistence images and calculated their distances as in (20). Briefly, we

converted the persistence barcode, which encodes the start and end radial distances of a branch

in the neuronal tree, into a persistence diagram. In the persistence diagram, each bar of the

barcode is converted into a point in a 2D space, where the X and Y coordinates are the start and

end radial distances of each bar. The persistence diagram was then converted in a persistence

image by applying a Gaussian kernel. We used the library NeuroM (45) to perform Sholl and

morphometrics analyses. The reconstructed morphologies will be made publicly available on

neuromorpho.org.

Ionic currents models

We used Hodgkin-Huxley types of ionic current models, starting from kinetics equations

already available in the neuroscientific literature. Along with kinetics of the ionic currents, we

stored information on the experimental conditions, such as temperature and LJP, by using the

software NeuroCurator (46). Whenever the data was available, we compared simulated voltage-

clamp experiments to experimental data from juvenile rats. Ionic currents I_i were defined as

functions of the membrane potential v, its maximal conductance density g_i and the constant

value of the reversal potential E_i:

$$I_i = g_i\, m_i^x\, h_i^y (v - E_i)$$

m_{ion} and h_{ion} represent activation and inactivation probability (varying between 0 and 1), with

integer exponents x and y. Each probability varied according to:

$$n^{'}(v) = (n_\infty(v) - n)/\tau_n(v)$$

where $n_\infty(v)$ is a function of voltage that represents the steady-state activation/inactivation

function (normally fitted with a Boltzmann curve) and $\tau_n(v)$ is a voltage-dependent time constant.

Exceptions to this formalism are ionic currents that do not inactivate (y = 0) and ionic currents

with (in)activation processes mediated by two or more time constants. Calcium currents (I_{CaT} and

I_{CaL}) were modeled according to the Goldman-Hodgkin-Katz constant field equation and had

permeability values instead of conductance (47).

Fast transient sodium current I_{NaT} and delayed potassium current I_{Kd}. I_{NaT} and I_{Kd} were

taken from a previous models of rat TC neurons from the VB nucleus (12), available on SenseLab

ModelDB (accession no. 279). I_{NaT} was compared with recordings of transient sodium currents in

P7-11 rat neurons from the dorsolateral geniculate (dLGN) nucleus (48).

Low-threshold activated (T-type) calcium current I_{CaT}. I_{CaT} model was taken from (12) and

available on-line (ModelDB, accession no. 279). This model was based on data recorded from VB

neurons of Sprague-Dawley rats (P7-12) at room temperature and corrected for −9 mV LJP (11).

Hyperpolarization-activated cationic current I_H. The steady-state activation for I_H was

derived from VB thalamic neurons in P10-20 Long-Evans rats and was already corrected for −10

mV LJP in the original publication (49). The equation used was:

573
$$m_\infty = 1/(1 + \exp[(v + 86.4)/11.2])$$

574   The time constant of activation was modeled as in (50), which derived a mathematical

575   description of I_H based on data from the dLGN in adult guinea pigs, recorded at 35.5 °C (51). The

576   equation describing the time dependence of activation was not corrected for simulations at

577   different temperatures and was:

578
$$\tau_m = 1/[\exp(-14.59 - 0.086v) + exp(-1.87 + 0.0701v)]$$

The equilibrium potential of the channel E_H was −43 mV. In silico voltage-clamp

experiments were compared with data in (49).

Persistent sodium current I_{NaP}. We modeled I_{NaP} as in (17) which based their model on

recordings from entorhinal neurons of Long-Evans rats (P25-P35) (52). The steady-state

activation was modified according to (48) and the steady-state inactivation according to (14). The

original steady-state activation data were recorded at room temperature (22-24°) and corrected

for −6/−7 mV LJP. Limited data on I_{NaP} are available from dissociated neurons from the dLGN

nucleus in Wistar rats (48).

Fast transient (A-type) potassium current I_{KA}. The mathematical formulation of I_{KA} was

based on data recorded from VB neurons in Sprague-Dawley rats (P7-15), recorded at room

temperature (22-24 °C) (53). A Q_{10} = 2.8 was experimentally determined and used for simulations

at different temperatures. In the original experiments a small LJP (<−4 mV) was measured and

not corrected. The current had a rapid and a slow component, represented by two activation and

two inactivation variables. The model of this current was provided by the authors of (14).

High-threshold (L-type) calcium current I_{CaL}. I_{CaL} model is the same as TC neurons model

previously published (14,50). The model was based on data from isolated guinea-pig

hippocampal neurons, recorded at room temperature (20-22 °C) with modifications to the

Boltzmann curve parameters of activation contained in the correction to the original models [59].

A small LJP (<3 mV) was not corrected (50). A Q_{10} = 3 was used for simulations at different

temperatures.

Calcium-activated potassium currents. TC neuron express genes for BK-type (54) and SK-

type calcium-activated potassium channels (55). Models of BK-type currents, similar to the I_C

current, have already been used to model TC neurons (14,50,54). However, data characterizing

this current in mammalian neurons are not available. We thus included only a model of I_{SK}

(available in ModelDB, accession no. 139653) based on rat mRNA expression data in Xenopus

oocytes (56).

Intracellular calcium dynamics. A simple exponential decay mechanism was used to model

the intracellular calcium dynamics (ModelDB, accession no. 139653). Both I_{CaT} and I_{CaL}

contributed to the intracellular calcium concentration.

In addition, we included a voltage-insensitive membrane current I_{leak}. The equilibrium

potential was −79 mV and corresponded to the average resting potential from our experimental

recordings.

Simulation and parameters optimization

NEURON 7.5 software was used for simulation (57). We used NEURON variable time step

method for all simulations. For the sake of spatial discretization, each section was divided into

segments of 40 μm length. The following global parameters were set: initial simulation voltage

(−79 mV), simulation temperature (34 °C), specific membrane capacitance (1 μF/cm^2), specific

intracellular resistivity 100 Ωcm for all the sections, equilibrium potentials for sodium and

potassium were 50 mV and −90 mV, respectively.

BluePyOpt (18) with Indicator Based Evolutionary Algorithm (IBEA) were used to fit the

models to the experimental data. Each optimization run was repeated with three different

random seeds and evaluated 100 individuals for 100 generations. The evaluation of these 300

individuals for 100 generations was parallelized using the *iPython ipyparallel* package and took

between 21 and 52 h on 48 CPU cores (Intel Xeon 2.60 GHz) on a computing cluster. Each

optimization run typically resulted in tens or hundreds of unique acceptable solutions, defined

as models having all feature errors below 3 STD from the experimental mean.

The models will be made publicly available at ModelDB (58). The configuration files for the

optimization and analysis will be made publicly available on Github, Bluepyopt page (59).

Sensitivity analysis

We performed a sensitivity analysis of an optimization solution by varying one parameter

value (p_m) at a time and calculating the electrical features from the voltage traces (y$^+$ and y$^-$). We

defined the sensitivity as the ratio between the normalized feature change and the parameter

change, which for smooth functions approximates a partial derivative (60,61). The features

changes were normalized by the optimized feature value. For small changes of parameter values,

we assumed that the features depend linearly on its parameters. We could thus linearize the

relationship between the features and the parameters around an optimized parameter set and

calculate the derivatives. The derivatives were calculated with a central difference scheme (60).

636
$$\frac{\partial y_n}{\partial p_m} \approx \frac{y_n^+ - y_n^-}{2\Delta p_m}$$

We collected the derivatives (sensitivities) in the $N \times M$ Jacobian matrix, with N

representing the number of features and M the number of parameters.

To rank parameters and features we computed their relative importance by calculating

their norms (the square root of the summed squared values) from the Jacobian columns and

rows, respectively. To cluster parameters based on similar influences on the features and to

cluster features that were similarly dependent on the parameters, we used angles between

columns (or rows) to compute distances D between parameters (or features):

644
$$D = 1 - |\cos \theta|$$

645 Features where thus considered similar if they depended in a similar manner on the

646 parameters, independent of sign or magnitude.

Acknowledgements

648 We thank the Blue Brain Project Visualization Team for generating the meshes and

649 visualizing the 3D morphologies.

Author Contributions

651 **Conceptualization**: E.I., S.L.H. **Data Acquisition and Curation:** J.Y., Y.S. **Formal Analysis**: E.I., J.Y.,

652 B.Z., C.O. **Funding Acquisition**: S.L.H., H.M. **Investigation**: E.I., J.Y., Y.S. **Methodology**: E.I., B.Z.,

653   C.R., W.V. **Project Administration:** C.O., S.L.H., H.M. **Resources and Software**: J.Y., Y.S., B.Z., W.V.,

654   C.R. **Supervision**: S.L.H., H.M. **Validation and Visualization**: E.I. **Writing – Original Draft**

655   **Preparation**: E.I. **Writing – Review & Editing**: E.I., J.Y., C.O., W.V., H.M., S.L.H.

Supporting Information captions

658   **S1 Features Dataset – Electrical features**

659   Spreadsheet containing experimental electrical figures, separated in three different sheets. The

660   features, their means and standard deviations are used in Fig 1C, Fig 2, Fig 3, Fig 4C, Fig 5C, Fig

661   8A.

662   **S2 Supporting figures**

References

References

1. Sherman SM. The thalamus is more than just a relay. Current opinion in neurobiology.
2007;17(4):417–422.

2. Francis JT, Xu S, Chapin JK. Proprioceptive and cutaneous representations in the rat ventral
posterolateral thalamus. Journal of neurophysiology. 2008;99(5):2291–2304.

3. Jones E. The thalamus 2nd ed. Cambridge University Press Cambridge, UK; 2007.

4. McAlliser J, Wells J. The structural organization of the ventral posterolateral nucleus in the
rat. Journal of Comparative Neurology. 1981;197(2):271–301.

5. Llinás R, Jahnsen H. Electrophysiology of mammalian thalamic neurones in vitro. Nature.
1982;297(5865):406.

6. Jahnsen H, Llinas R. Electrophysiological properties of guinea-pig thalamic neurones: an in
vitro study. The Journal of physiology. 1984;349(1):205–226.

7. Turner JP, Anderson CM, Williams SR, Crunelli V. Morphology and membrane properties of
neurones in the cat ventrobasal thalamus in vitro. The Journal of Physiology.
1997;505(3):707–726.

8. Connelly WM, Crunelli V, Errington AC. Variable action potential backpropagation during
tonic firing and low-threshold spike bursts in thalamocortical but not thalamic reticular
nucleus neurons. Journal of Neuroscience. 2017;0015–17.

9. Druckmann S, Banitt Y, Gidon AA, Schürmann F, Markram H, Segev I. A novel multiple

objective optimization framework for constraining conductance-based neuron models by

experimental data. Frontiers in neuroscience. 2007;1:1.

10. Gouwens NW, Berg J, Feng D, Sorensen SA, Zeng H, Hawrylycz MJ, et al. Systematic

generation of biophysically detailed models for diverse cortical neuron types. Nature

communications. 2018;9(1):710.

11. Huguenard JR, McCormick DA. Simulation of the currents involved in rhythmic oscillations in

thalamic relay neurons. Journal of neurophysiology. 1992;68(4):1373–1383.

12. Destexhe A, Neubig M, Ulrich D, Huguenard J. Dendritic low-threshold calcium currents in

thalamic relay cells. Journal of Neuroscience. 1998;18(10):3574–3588.

13. Rhodes PA, Llinás R. A model of thalamocortical relay cells. The Journal of physiology.

2005;565(3):765–781.

14. Amarillo Y, Zagha E, Mato G, Rudy B, Nadal MS. The interplay of seven subthreshold

conductances controls the resting membrane potential and the oscillatory behavior of

thalamocortical neurons. Journal of Neurophysiology. 2014;112(2):393–410.

15. Connelly WM, Crunelli V, Errington AC. The global spike: conserved dendritic properties

enable unique Ca2+ spike generation in low-threshold spiking neurons. Journal of

Neuroscience. 2015;35(47):15505–15522.

16. Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, et al.

Reconstruction and Simulation of Neocortical Microcircuitry. Cell. 2015;163(2):456–492.

17. Hay E, Hill S, Schürmann F, Markram H, Segev I. Models of neocortical layer 5b pyramidal cells

capturing a wide range of dendritic and perisomatic active properties. PLoS computational

biology. 2011;7(7):e1002107.

18. Van Geit W, Gevaert M, Chindemi G, Rössert C, Courcol J-D, Muller EB, et al. BluePyOpt:

leveraging open source software and cloud infrastructure to optimise model parameters in

neuroscience. Frontiers in neuroinformatics. 2016;10:17.

19. Paxinos G, Watson C. The rat brain atlas in stereotaxic coordinates. San Diego: Academic.

1998;

20. Kanari L, D\lotko P, Scolamiero M, Levi R, Shillcock J, Hess K, et al. A topological

representation of branching neuronal morphologies. Neuroinformatics. 2018;16(1):3–13.

21. Sholl DA. Dendritic organization in the neurons of the visual and motor cortices of the cat.

Journal of anatomy. 1953;87(Pt 4):387.

22. Ascoli G a, Alonso-Nanclares L, Anderson S a, Barrionuevo G, Benavides-Piccione R,

Burkhalter A, et al. Petilla terminology: nomenclature of features of GABAergic interneurons

of the cerebral cortex. Nature reviews Neuroscience. 2008;9(7):557–568.

23. Almog M, Korngreen A. Is realistic neuronal modeling realistic? Journal of neurophysiology.

2016;116(5):2180–2209.

24. Williams SR, Stuart GJ. Action potential backpropagation and somato-dendritic distribution

of ion channels in thalamocortical neurons. Journal of Neuroscience. 2000;20(4):1307–1317.

25. Errington AC, Renger JJ, Uebele VN, Crunelli V. State-dependent firing determines intrinsic

dendritic Ca2+ signaling in thalamocortical neurons. Journal of Neuroscience.

2010;30(44):14843–14853.

26. Druckmann S, Berger TK, Hill S, Schürmann F, Markram H, Segev I. Evaluating automated

parameter constraining procedures of neuron models by experimental and surrogate data.

Biological cybernetics. 2008;99(4–5):371.

27. Velazquez JLP, Carlen PL. Development of firing patterns and electrical properties in neurons

of the rat ventrobasal thalamus. Developmental brain research. 1996;91(2):164–170.

28. Pozzorini C, Mensi S, Hagens O, Naud R, Koch C, Gerstner W. Automated high-throughput

characterization of single neurons by means of simplified spiking models. PLoS computational

biology. 2015;11(6):e1004275.

29. Hay E, Schürmann F, Markram H, Segev I. Preserving axosomatic spiking features despite

diverse dendritic morphology. Journal of neurophysiology. 2013;109(12):2972–2981.

30. Krahe TE, El-Danaf RN, Dilger EK, Henderson SC, Guido W. Morphologically distinct classes of

relay cells exhibit regional preferences in the dorsal lateral geniculate nucleus of the mouse.

Journal of Neuroscience. 2011;31(48):17437–17448.

31. Migliore R, Lupascu CA, Bologna LL, Romani A, Courcol J-D, Antonel S, et al. The physiological

variability of channel density in hippocampal CA1 pyramidal cells and interneurons explored

using a unified data-driven modeling workflow. PLoS computational biology.

2018;14(9):e1006423.

32. Masoli S, Rizza MF, Sgritta M, Van Geit W, Schürmann F, D'Angelo E. Single neuron

optimization as a basis for accurate biophysical modeling: the case of cerebellar granule cells.

Frontiers in cellular neuroscience. 2017;11:71.

33. Schulz DJ, Goaillard J-M, Marder E. Variable channel expression in identified single and

electrically coupled neurons in different animals. Nature neuroscience. 2006;9(3):356.

34. Taylor AL, Goaillard J-M, Marder E. How multiple conductances determine

electrophysiological properties in a multicompartment model. Journal of Neuroscience.

2009;29(17):5573–5586.

35. O'Leary T, Williams AH, Caplan JS, Marder E. Correlations in ion channel expression emerge

from homeostatic tuning rules. Proceedings of the National Academy of Sciences.

2013;110(28):E2645–E2654.

36. Ranjan R, Khazen G, Gambazzi L, Ramaswamy S, Hill SL, Schürmann F, et al. Channelpedia: an

integrative and interactive database for ion channels. Frontiers in neuroinformatics.

2011;5:36.

37. Podlaski WF, Seeholzer A, Groschner LN, Miesenböck G, Ranjan R, Vogels TP. Mapping the

function of neuronal ion channels in model and experiment. Elife. 2017;6:e22152.

38. Wei H, Bonjean M, Petry HM, Sejnowski TJ, Bickford ME. Thalamic burst firing propensity: a

comparison of the dorsal lateral geniculate and pulvinar nuclei in the tree shrew. Journal of

Neuroscience. 2011;31(47):17287–17299.

39. Connelly WM, Crunelli V, Errington AC. Passive synaptic normalization and input synchrony-

dependent amplification of cortical feedback in thalamocortical neuron dendrites. Journal of

Neuroscience. 2016;36(13):3735–3754.

40. Tennøe S, Halnes G, Einevoll GT. Uncertainpy: A Python toolbox for uncertainty quantification

and sensitivity analysis in computational neuroscience. bioRxiv. 2018;274779.

41. Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, et al.

Reconstruction and simulation of neocortical microcircuitry. Cell. 2015;163(2):456–492.

42. BlueBrain. BlueBrain/eFEL [Internet]. 2018. Available from:

https://github.com/BlueBrain/eFEL

43. RStudio Team. RStudio: Integrated Development Environment for R [Internet]. Boston, MA:

RStudio, Inc.; 2016. Available from: http://www.rstudio.com/

44. Choisy M. Identifying a cutoff value from bimodal data [Internet]. 2018. Available from:

http://marcchoisy.free.fr/fmm/index.html

45. BlueBrain. BlueBrain/NeuroM [Internet]. 2018. Available from:

https://github.com/BlueBrain/NeuroM

46. O'Reilly C, Iavarone E, Hill SL. A Framework for Collaborative Curation of Neuroscientific

Literature. Frontiers in neuroinformatics. 2017;11:27.

47. Hille B. Ionic channels in excitable membranes. Current problems and biophysical

approaches. Biophysical Journal. 1978;22(2):283–294.

48. Parri HR, Crunelli V. Sodium current in rat and cat thalamocortical neurons: role of a non-

inactivating component in tonic and burst firing. Journal of Neuroscience. 1998;18(3):854–

867.

49. Budde T, Biella G, Munsch T, Pape H-C. Lack of Regulation by Intracellular Ca2+ of the Hyper

Polarization-Activated Cation Current in Rat Thalamic Neurones. The Journal of physiology.

1997;503(1):79–85.

50. McCormick D a, Huguenard JR. A model of the electrophysiological properties of

thalamocortical relay neurons. Journal of neurophysiology. 1992;68(4):1384–1400.

51. McCORMICK DA, Pape H-C. Properties of a hyperpolarization-activated cation current and its

role in rhythmic oscillation in thalamic relay neurones. The Journal of physiology.

1990;431(1):291–318.

52. Magistretti J, Alonso A. Biophysical properties and slow voltage-dependent inactivation of a

sustained sodium current in entorhinal cortex layer-II principal neurons: a whole-cell and

single-channel study. The Journal of general physiology. 1999;114(4):491–509.

53. Huguenard J, Coulter D, Prince D. A fast transient potassium current in thalamic relay

neurons: kinetics of activation and inactivation. Journal of neurophysiology.

1991;66(4):1304–1315.

54. Ehling P, Cerina M, Meuth P, Kanyshkova T, Bista P, Coulon P, et al. Ca2+-dependent large

conductance K+ currents in thalamocortical relay neurons of different rat strains. Pflügers

Archiv-European Journal of Physiology. 2013;465(4):469–480.

55. Gymnopoulos M, Cingolani LA, Pedarzani P, Stocker M. Developmental mapping of small-

conductance calcium-activated potassium channel expression in the rat nervous system.

Journal of Comparative Neurology. 2014;522(5):1072–1101.

56. Köhler M, Hirschberg B, Bond C, Kinzie JM, Marrion N, Maylie J, et al. Small-conductance,

calcium-activated potassium channels from mammalian brain. Science.

1996;273(5282):1709–1714.

57. Hines ML, Carnevale NT. The NEURON simulation environment. Neural computation.

1997;9(6):1179–1209.

58. McDougal RA, Morse TM, Carnevale T, Marenco L, Wang R, Migliore M, et al. Twenty years

of ModelDB and beyond: building essential modeling tools for the future of neuroscience. J

Comput Neurosci. 2017 Feb 1;42(1):1–10.

59. BlueBrain. BlueBrain/BluePyOpt [Internet]. 2018. Available from:

https://github.com/BlueBrain/BluePyOpt

60. Weaver CM, Wearne SL. Neuronal firing sensitivity to morphologic and active membrane

parameters. PLoS Computational Biology. 2008;4(1):e11.

61. Olifer AV. Neuronal Parameter Sensitivity. In: Jaeger D, Jung R, editors. Encyclopedia of

Computational Neuroscience [Internet]. New York, NY: Springer New York; 2013 [cited 2018

Dec 19]. p. 1–6. Available from: https://doi.org/10.1007/978-1-4614-7320-6_172-1