Divergent selection following speciation in two ectoparasitic honey bee mites

Maeva A. Techer[1*], Rahul V. Rane[2,3], Miguel L. Grau[1], John M. K. Roberts[4], Shawn T. Sullivan[5], Ivan Liachko[5], Anna K. Childers[6,7], Jay D. Evans[7] and Alexander S. Mikheyev[1,8*]

Okinawa Institute of Science and Technology, 1919-1 Tancha Onna-son, 904-0495 Okinawa, Japan

Commonwealth Scientific and Industrial Research Organisation, Clunies Ross St, (GPO Box 1700), Acton, ACT, 2601, Australia

Bio21 Institute, School of BioSciences, University of Melbourne, 30 Flemington Road, Parkville, 3010, Australia

CSIRO, Canberra, ACT, 2601, Australia

Phase Genomics Inc, Seattle, WA, 98195, USA

USDA-ARS Insect Genetics and Biochemistry Research Unit, Fargo, ND, USA

USDA-ARS Bee Research Lab, Beltsville, MD, USA

Australian National University, Canberra, ACT, 2600, Australia

* Corresponding authors: maeva.techer@oist.jp, alexander.mikheyev@oist.jp

Abstract

Multispecies host-parasite evolution is common, but how parasites evolve after speciating remains poorly understood. Shared evolutionary history and physiology may propel species along similar evolutionary trajectories whereas pursuing different strategies can reduce competition. We test these scenarios in the economically important association between honey bees and ectoparasitic mites by sequencing the genomes of the sister mite species *Varroa destructor* and *Varroa jacobsoni*. These genomes were closely related, with 99.7% sequence identity. Among the 9,628 orthologous genes, 4.8% showed signs of positive selection in at least one species. Divergent selective trajectories were discovered in conserved chemosensory gene families (IGR, SNMP), and Halloween genes (CYP) involved in moulting and reproduction. However, there was little overlap in these gene sets and associated GO terms, indicating different selective regimes operating on each of the parasites. Based on our findings, we suggest that species-specific strategies may be needed to combat evolving parasite communities.

Interactions between hosts and parasites shape biodiversity by driving coevolutionary

arms races and by regulating populations over ecological timescales. Parasitism is a remarkably

successful strategy, occurring in nearly half of recognized animal phyla [1,2]. Parasites depend on

their hosts which form their principal ecological niches and provide essential resources to

survive [3]. Commonly, multiple parasite generations elapse per host generation. Thus parasites

are prone to evolve rapidly and may circumvent any gained host advantage [4]. The similar

selective regimes acting on phylogenetically unrelated parasites resulted in reproducible

evolution such as genome size reduction, functional losses associated with reduced metabolism

and adaptive functional gains linked to parasite virulence [5]. Most models of parasite

coevolution are limited to interactions between single focal parasites and their hosts, in part

because multi-species dynamics are much more complex [6,7]. In nature, the reality is that

parasite's success and evolution will also depend on interactions with co-infecting conspecific

and heterospecific within its host niche [8–10]. Yet, coevolutionary dynamics involving more than

one parasitic species remain poorly understood [11,12].

For example, co-infecting parasites occupying the same host can exclude competitors

via direct, or interference competition [13,14]. One strategy to reduce interspecific competition

can be spatial separation within regions of the host gut as adopted by cestodes that naturally

co-infect sticklebacks [15]. Within the same host, spatial segregation that allows parasite co-

existence can ultimately lead to intrahost speciation, as found in gill parasites (*Dactylogyrus*) of

freshwater fish (*Cyprinidae*) [16]. Niches may also be partitioned temporally, which allows two

fungal sibling species (*Erysiphe quercicola* and *E. alphitoides*) to exploit the same host tree

(*Quercus robur*) by reducing direct interference [17]. Alternatively, parasites may shift their

strategies to exploit different aspects of the host niche. For example, fire ants (*Solenopsis spp.*)

are parasitized by a genus of decapitating phorid fly (*Pseudacteon*), which are highly host-

specific and have the same life cycle, but specialize on attacking ants in different contexts (*e.g.,*

while foraging, *vs.* at the nest.) [18]. However, almost nothing is known about evolutionary

trajectories of speciating parasites. On one hand, they share physiology and host-specific

adaptations, which may predispose them to evolve along common lines, particularly in

allopatry. On the other hand, selective regimes may be less predictable, or even disruptive, in

the case of character displacement.

We investigated the evolution of speciating parasites by sequencing genomes of two

economically important mite species that specialize on honey bees (*Apis sp.*): *Varroa jacobsoni*

and *Varroa destructor*. The honey bee colony, which is a densely packed community of

genetically similar individuals, can host many diseases and parasites. Varroa mites are

obligatory ectoparasites which feed on fat bodies of developing larvae/pupae and adult honey

bees (shown in *V. destructor*) [19] and reproduce in the brood [20]. Parasitic feeding injures vital the

honey bee host but leading to abnormal development [21], reduced worker lifespan [22], the

transmission of viruses [23] and pathogenic bacteria [24]. These combined negative effects on

individual honey bee decrease the fitness of the colony as a superorganism. Without acaricide

treatments, managed western honey bee colonies were reported to decline and die within two

to three years [20]. As its name indicates, *V. destructor* is considered as the main devastating

force behind colony collapses and impact deeply western honey bee (*A. mellifera*) beekeeping

[25,26].

The dramatic effects of *Varroa* mites on *A. mellifera* are due to their recent

coevolutionary history. Originally, brood parasitism by *Varroa* ectoparasitic mites has evolved in

all *Apis* species except for *Apis mellifera* [27]. Until the 2000s, *V. jacobsoni* and *V. destructor* were

considered a single species, based on their life cycle, morphological similarities (Figure 1A) [28]

and given that both originally parasitized the widespread eastern honey bee (*Apis cerana*)

(Figure 1B). However, as western honey bees were brought into contact with the eastern honey

bee in Asia and Oceania, both mites repeatedly colonized this novel host [29,30] (Figure 1C). In

contrast to what has happened to *A. mellifera*, the host-parasite interaction with *A. cerana* is a

balanced relationship in which a tolerance "equilibrium" is maintained by a range of parasite

adaptations and host counter-adaptations [31–36]. Specifically, the eastern honey bee has

impressive host defense traits, including i) self and collective grooming behaviors [37], ii) hygienic

behavior by removing mites from infested cell [38], iii) worker brood susceptibility stopping

parasite reproduction [33,34], iv) entombing of mites in drone brood and v) shorter worker

development cycle limiting mites offspring number [32]. By contrast, most western honey bee

populations lack these defensive mechanisms, and hives collapse quickly after *Varroa*

infestation. On the part of *Varroa* mites, they evolved highly specialized and adaptive strategies

to find, select, feed, reproduce and hide on its host [39]. While many mite adaptations have

already been discovered [20,39], the recent use of 'omics approaches has quickly broadened our

understanding on *V. destructor* hidden specialization to host [36,40–42]. For example,

transcriptomic and proteomic profiling of different *V. destructor* life stages burst the

identification of genomic regions related to chemosensory system [40,42,43], reproductive success

[41] and linking oogenesis initiation or failure to parasitic reduced metabolism and host defense

[44,45].

Here, we ask how both mite species have arrived at this tolerance equilibrium in *A.*

cerana – did they follow similar or different evolutionary paths? On one hand, the two species

naturally occur in parapatry (Figure 1B), which is expected to result from divergent local

adaptation with the distribution of different *A. cerana* lineages [46,47]. In actuality, what drives the

parapatric distribution is unclear. *V. destructor*, which was historically absent from Southeast
Asia, now occurs there on its novel host *A. mellifera* in sympatry with *V. jacobsoni*, which does
not parasitize *A. mellifera* in this region (Figure 1C). Thus, it is possible that the current
geographical distribution resulted from secondary contact following allopatric speciation. This
view is further supported by the fact that both species have the genetic capacity to shift to *A.*
mellifera as a novel host, suggesting a level of physiological similarity. We tested these
alternative hypotheses (parallel *vs.* divergent evolution) by generating high-quality hybrid *de*
novo genome assembly of both species and examining them for signatures of adaptive
evolution. We also compared them to the recently published genome of a distantly related
ectoparasitic mite *Tropilaelaps mercedesae*, to see whether the patterns appear universal at
different timescales.
127

Figure 1: ***V. destructor* and *V. jacobsoni* are morphologically similar sister species originally parasitizing the eastern honey bee (*A. cerana*).** These species were recognized upon the basis of quantitative morphometric and genetic data [28.] Morphological differences are restricted to differences in body size and shape, as shown in (A) 3D surface models comparison of two fully sclerotized females. (B) Both mites parasitize *A. cerana* as their original host and can coexist in sympatry, occasionally even at the same apiary [46,48], but are mainly parapatric [49]. (C) With the introduction of *A. mellifera*, *V. destructor* can be found on this novel host throughout the original *V. jacobsoni* range. *V. jacobsoni* has also extended its range into Papua New Guinea on *A. cerana* followed by a shift to *A. mellifera*, where *V. destructor* is currently absent [30].

Methods

Sequencing strategy and *Varroa* mite collection

We took advantage of the haplodiploid sex determination of both *V. destructor* and *V. jacobsoni* to generate data to reduce the ambiguity due to heterozygous sites for a high contiguous hybrid genome assembly. The sequencing strategy adopted in this study combined shorts reads (Illumina) for which input DNA can be obtained from a single mite individual, long reads of tens of kilobases (PacBio) which requires a large quantity of high-molecular-weight DNA and Hi-C to capture chromosome conformation and interactions.

Haploid males (~0.7 mm by 0.7 mm) are only found inside capped brood cells after initiation of reproduction while diploid females (~1 mm by 1.5 mm) can be found in the brood and on adult honey bees. All *V. destructor* samples were collected from *A. mellifera* colonies of Okinawa Institute of Science and Technology (OIST) experimental apiary in Japan (26.477 N 127.835 E) in August-September 2016. One mature male *V. destructor* was collected within a worker cell at the red-eye pupa developmental stage (day 15 to 16) using a brush and kept in absolute ethanol at -80° C. Mature adult females were collected from adult honey bee workers. In order to obtain a large number of mites infecting adult workers, we modified the standard powdered sugar method [50] for an entire managed colony as follows: one hive box containing three to four frames with honey bee workers and no queen was placed on top of a collecting white tray; approximately 500g of icing sugar was then applied using sieve filter powder strainer; the sugar and honey bees were shaken in the tray for 2 min and then separated using a sieve filter, placing the sugared-workers directly within the hive. This process was repeated on adjacent colonies. *Varroa* mites were separated from their hosts and trapped in the icing sugar. Following a water rinse on a gauze mesh, a total of 1,207 alive *Varroa* females were snap frozen at -80° C until laboratory processing (inactive, sluggish or dead mites were discarded). Samples of *V. jacobsoni* were obtained from its first detection survey where it was found reproducing in the western honey bee [30]. One mature male (VJ856) collected from *A. mellifera* drone brood cell in Papua New Guinea (between EHP and Henganofi border, 30/05/2008) was used to prepare a whole genome library.

One sclerotized mature female *V. destructor* (VDOKI-01) collected from OIST apiary in the reproductive phase and one sclerotized mature female *V. jacobsoni* (VJ347-11) collected from *A. cerana* originating in Java (Cililin central Java, 13/06/1998 by D. Anderson) were processed for X-ray microtomography (micro-CT) using a Zeiss Xradia 510 Versa scanner. 3D surface models were reconstructed from micro-CT outputs using Amira software to visualize

176 the slight morphological differences when focusing on idiosoma ventral and dorsal views and
177 whole body in profile.
178
179 **DNA extraction, genome sequencing, assembly, and annotation**
180
Genomic DNA was extracted from each single haploid male *V. jacobsoni* and *V.*
destructor using QIAamp DNA Micro Kit (Qiagen) following manufacturer's instructions. The
total amount of dsDNA was measured using Quant-iT™ PicoGreen™ dsDNA Assay Kit
(Invitrogen). Short-insert of 250 bp paired-end libraries were prepared for both individuals
using NEBNext® Ultra™ II DNA Library Prep Kit for Illumina® (Biolabs) and 10 PCR cycles. Size-
selection and cleanup was applied using SPRI beads and library was subsequently checked using
a Bioanalyzer High Sensitivity DNA kit (Agilent). Both paired-end libraries were each sequenced
on two lanes of Illumina HiSeq 2500 at the OIST sequencing center in paired-end 250 cycle
mode.
Prior to assembly paired-end reads were de-duplicated using ParDRe [51], and host and
microbial contamination was filtered using DeConSeq (parameters: -i 97 -c 95) [52], screening
against genomes of *Apis cerana*, *Apis mellifera*, as well as bacterial genomes and genomes of
bee diseases and parasites. The cleaned reads were then merged using PEAR (parameters: --
min-overlap 10 --memory 80G --threads 10 -n 200 -m 600 -p 0.0001) [53] and assembled using
Newbler (v. 2.6) (parameters: -large -m -cpu 24 -mi 98 -siom 390 -l 1000 -a 500 -urt -novs -a
1000 -minlen 45 -ml 100) [54]. The contigs were then scaffolded using publicly available RNA-seq
data assembled using Trinity [55,56]. For *V. destructor* further scaffolding was carried out using Hi-
C, performed by Phase Genomics. Hi-C read pairs were aligned to the initial assembly using
BWA-MEM with the "-5" option [57], keeping only primary properly paired alignments (samtools
with the "-F 2316" filter [58]). Scaffolding was then performed using Phase Genomics' Proximo Hi-
C scaffolding pipeline [59], which builds on the LACHESIS method [60].
To estimate genome characteristics, K-mers from cleaned reads were counted using
jellyfish version 2.2.7 [61] for the following values of K: 19, 25, 31 and 42 and coverage cut-off of
1,000. Frequencies were only computed on HiSeq 2500 reads from one individual haploid male
of each species. Computed and selected K-mer histograms for k=42 were analyzed with
GenomeScope to estimate genome size, kcoverage, number of duplicates [62]. We used the
bioinformatically derived genome sizes for genome coverage estimation.
Small batches of adult females *V. destructor* (1,207 mites total) were pooled and
crushed by pestle in 1.5 mL Eppendorf tubes. The powder was then lysed using proteinase K
and lysis buffer at 56 for 20 min. Sample lysate was thus transferred to a Maxwell Cartridge.

The remaining step was fully performed on the Maxwell 16 automated purification system
(Promega) with the LEV Blood DNA mode. Library preparation for PacBio RSII sequencing was
performed using SMRTbell Template Prep Kit 1.0 (Pacific Biosciences) following the
manufacturer's recommendations. SMRTbell libraries were prepared and sequenced at the
OIST sequencing center on 48 SMRT cells. These reads were trimmed using proovread [63] and
then used to gap-fill the Hi-C scaffolded *V. destructor* assembly using PBJelly (parameters: -
minMatch 8 -minPctIdentity 70 -bestn 1 -nCandidates 20 -maxScore -500 -nproc 24 -
noSplitSubreads) (v. 15.8.24) [64].
Annotation for both genomes Vdes_3.0 and Vjacob_1.0 was carried out by NCBI using
NCBI's automated genome annotation pipeline. This pipeline takes advantage of species-
specific RNA-seq data, as well as extensive protein homology data stored in GenBank [65].
222
223 **Identification of *Varroa* mite strains using mtDNA and sequence similarity**
224
Most previous taxonomic work relied on mitochondrial fragments. We used
mitochondrial analysis to determine and compare the lineages of *Varroa* species used for
genome sequencing to the previously described haplotypes. The whole mitogenome of *V.*
destructor (accession NC_004454.2 [66,67]) was available from the NCBI database and used as a
reference for read mapping. Raw Illumina® reads for both *V. destructor* and *V. jacobsoni* males
were mapped onto the entire reference sequence (16,476 bp) using NextGenMap 0.5.0 [68].
Mapped reads were sorted and PCR duplicates were removed using Samtools 1.3.1 [58].
Maximum coverage of 500x was subsampled on all mapped reads using VariantBam 1.4.3 [69].
Subsequently, variant calling was performed using FreeBayes v1.1.0-46 [70], with the following
parameters others than the default: minimum mapping quality of 20 and minimum base quality
of 5. Similarly, Illumina reads used to construct Vjacob_1.0 were mapped against the reference
Vdes_3.0 to detect the number of variants in the nuclear genome. Consensus mitochondrial
sequence was generated for each individual using vcf2fasta from Vcflib library [71].
One standard method to identify Varroa species and haplogroup is to sequence a partial
region of the *Cox1* gene (426 bp) from the mitochondrial DNA [50]. The same region was
extracted from both Vdes_3.0 and Vjacob_1.0 generated mitochondrial sequence and aligned
together with 26 *Varroa spp.* haplotypes previously described [28,72–74]. Given its worldwide bee
pest status, *V. destructor* haplogroups were better described than *V. jacobsoni* and additional
larger *Cox1, Cox3, Atp6* and *cytB* mitochondrial sequences were available on NCBI. Therefore, a
more resolutive comparison within *V. destructor* was done by aligning concatenated sequences
of the four genes with 22 haplotypes described previously [72,75]. Sequences were aligned using

ClustalW, checked by hand and a consensus maximum likelihood tree was constructed after
5000 bootstraps in Mega 7.0.18 [76].

A distribution map of the *Varroa* mites infecting *A. cerana* was built from reviewing
literature. Conservatively, only the presence of *Varroa spp.* confirmed by mtDNA barcoding
after the taxonomic revision was plotted [28,46,72–74,77–79] onto the potential native and introduced
range of the Asian honey bee host [47,80]. The map was produced using QGIS 2.16.1 (Open Source
Geospatial Foundation Project qgis.osgeo.org) and the public domain natural 10m earth raster
NE1_HR_LC github.com/nvkelso/natural-earth-raster). An additional interactive map showing
the reported distribution of the 26 mtDNA haplotypes is available at
maevatecher.github.io/varroa-denovo-genomes with layers control options.

Prediction of gene orthology, selection, and duplication in *Varroa* mites

The Orthonome pipeline (www.orthonome.com) [81] was applied to predict orthologues
(pairwise and 1:1 orthogroups), inparalogues and lineage-specific gene expansions from the
two parasitic *Varroa* mites' genomes sequenced and annotated here and four Acari annotated
reference genomes available from NCBI
(www.ncbi.nlm.nih.gov/genome/browse/#!/overview/Acari). The Orthonome pipeline uses
phylogenetic information and microsynteny to identify gene copy number changes and tandem
duplications between pairs of species using the MSOAR2.0 pipeline [82]. We included four
outgroups which had fully sequenced genomes, including the eastern honey bee ectoparasitic
mite *Tropilaelaps mercedesae* [83], the parasitic tick *Ixodes scapularis* [84]; the free-living Western
predatory mite *Metaseiulus occidentalis* [85] and the free-living two spotted mites *Tetranychus*
urticae [86]. Total and pairwise/lineage intersection of orthologs gene sets were represented
using Upset plots [87] using R. The 1:1 orthogroups with both *Varroa* species represented and > 3
species were used for positive selection hypothesis tested on phylogeny using HyPhy to
calculate the mean dN/dS and test using aBSREL (adaptive Branch-Site Random Effects
Likelihood) [88]. This test allowed estimation of selection along specific branches of the tree,
including ancestral branches. Only branches with a significant p-value (< 0.05) after FDR
corrections were considered as positively selected in each orthogroup. Additionally, tandem
duplications in the three parasitic mites were identified from inparalogues in immediate vicinity
of the orthologues. All orthologous genes in Varroa mites under positive selection and/or
duplicated in tandem were checked by hand by using BLAST. Duplication events with more than
two genes were systematically checked by aligning the annotated coding sequence (CDS) with

ClustalW alignment implemented in Geneious R8 (Biomatters, Ltd., Auckland, New Zealand) and
construct a tree with IQ-TREE [89].

Gene ontology (GO) enrichment analysis of the positively selected genes in honey bee
parasitic mites were carried out using the GOstats R package [90]. The same process was applied
for gene duplication in *Varroa* mites. Biological processes associated with these GO terms were
summarized and visualized using REVIGO [91] (http://revigo.irb.hr). The semantic similarity was
measured using the Resnik's measure (SimRel) [92] and the threshold used was C = 0.5 (i.e. small).
The results were then used to produce a scatter plot and treemap plot using the ggplot2
package. The same package was used to plot the localization of the genes under positive
selection and/or duplicated onto *V. destructor* chromosomal scaffolds. Significance of overlaps
between gene sets and associated GO terms were tested using Fisher's exact test with
GeneOverlap R package. Code and raw plots for exploratory analysis of positively selected
genes, duplication in tandem and GO term association are available online
(maevatecher.github.io/varroa-denovo-genomes).

Chemosensory and ecdysteroid pathway-related gene families in host-parasite interactions

Varroa mites are blind and rely on their chemosensory system to find a suitable host
within a densely packed honey bee colony [39]. Chemical cues are transported to the nervous
system by binding to soluble carrier proteins and membrane proteins in insects. Several
homologous chemosensory protein families involved in these pathways have been identified in
Acari and *Varroa* mites [40,42,43,93]. Here we used the available catalogue of protein sequences
reported from *V. destructor* forelegs transcripts [43,93] containing six Odorant Binding Proteins
(OBP), six Niemann-Pick disease protein, type C2 (NPC2), seven Sensory Neuron Membrane
Proteins (SNMPs), three Glutamate Receptors (GRs) and 17 Ionotropic Glutamate Receptor
(IGRs). We identified these proteins in both *V. destructor* and *V. jacobsoni* annotated genomes
using BLASTP and compiled predicted protein sequences (keeping one isoform if several exist
for the same gene). We then aligned each family of protein sequences with other sequences
from other Acari and insects using MAFFT version 7 [94] (parameters: E-INS-i, the BLOSUM62
matrix, and MAFFT-homolog option with 100 homologs, E threshold = 0.1). We used IQ-TREE [95]
to find the best substitution model for each family alignment, with state frequency estimated
by maximum likelihood, 1000 ultra-fast bootstraps, 1000 replicate for SH-aLRT branch test. All
generated phylogenetic trees were visualized and exported using iTOL v4 [96].

We ran the same process for the cytochrome P450 enzyme (CYP) genes (including the
Halloween genes), known to be involved in the steroid molting hormone ecdysone biosynthesis
pathway in Arthropods. Ecdysteroids derivatives are also important as they are involved in the
vitellogenin production and oogenesis initiation in *V. destructor* after feeding on its host[44,97]. To
assess which CYP genes and in particular the Halloween genes were present in both *V.*
destructor and *V. jacobsoni* genomes*,* we used sequences from the fruit fly *Drosophila*
melanogaster, the mosquito *Aedes aegypti*, both honey bee hosts *A. mellifera* and *A. cerana*, as
well as other Acari (same name code as in a previous study on *M. occidentalis* [98]).
324
325

Results

Two new reference genomes for *V. destructor* and *V. jacobsoni*

We generated 28.6Gb (333 $\pm$ S.D. 56 bp) and 56.2 Gb (409 $\pm$ S.D. 61 bp) of de-duplicated, decontaminated, and merged Illumina reads for *V. jacobsoni* and *V. destructor*, corresponding to a coverage of 80$\times$ and 152$\times$, respectively. PacBio sequencing yielded 20.3 Gb of *V. destructor* reads with an N50 of 13.5 kb (66× coverage), which was used to fill 84.4% of the gaps in the scaffolded HiC assembly. Final *de novo* assemblies were named Vdes_3.0 (GCA_002443255.1) and Vjacob_1.0 (GCA_002532875.1) and are available on NCBI.

Through the k-mer analysis, the genome size estimates ranged from 369.02 to 365.13 Mbp using k = 42 (Figure S1). Both Vdes_3.0 and Vjacob_1.0 final assembly size were close to the predicted estimates but *V. jacobsoni* genome size was slightly shorter than *V. destructor* with 365.59 Mbp < 368.94 Mbp, respectively (Table 1). Previous assemblies size and contiguity for *V. destructor* with Vdes_1.0 [99] and Vdes_2.0 were improved as detailed in Table 1. According to the N50 statistics, more than 50% of the genome was in scaffolds of 58.5 Mbp or more for Vdes_3.0 while the Vdes_2.0 N50 was around 0.1 Mbp. When comparing to invertebrate assemblies, Vdes_3.0 genome is among the most highly contiguous with to date the best-reported scaffold N50 and best contig N50 for Acari and Mesostigmata as reported on the i5k database (i5k.github.io/arthropod_genomes_at_ncbi) [100].

Using Hi-C data, seven major scaffolds were constructed for *V. destructor* ranging from 39.4 to 76.9 Mbp with additional of 1,418 nuclear minor scaffolds > 0.2 Mbp. The longest genomic scaffolds correspond to the seven haploid chromosomes (2n = 14) previously detected in *V. destructor* karyotype [101]. Contrary to Vdes_3.0 which benefited from long reads and chromosome capture sequencing, *V. jacobsoni* Vjacob_1.0 genome was assembled from 8,241 contigs into 4,881 scaffolds with an N50 of 233,810 bp. Similar GC content composed 40.9% were found for both sister species Vdes_3.0 and Vjacob_1.0. The automated NCBI Eukaryote annotation pipeline predicted 10,247 protein-coding genes (out of 12,849 genes and pseudo-genes) for Vdes_3.0, and 10,739 protein-coding genes (out of 15,486 genes and pseudo-genes) for Vjacob_1.0 [65] (based upon transcriptome data available on the database in October 2017).

When Vjacob_1.0 reads were mapped onto the Vdes_3.0 reference genome, a total of 1,125,230 biallelic, filtered SNPs were detected. Relative to Vdes_3.0 genome size, sequence divergence with Vjacob_1.0 amounted to 0.30% across the nuclear genome sequence.

Table 1: Statistics for the release of new genome assembly for honey bee *Varroa* mites and improvement from the first *V. destructor* draft genome.

		V. destructor			*V. jacobsoni*
Genome assembly		**BRL_Vdes_1.0**	**Vdes2.0**	**Vdes_3.0**	**Vjacob_1.0**
NCBI accession		GCA_000181155.1	GCA_000181155.2	GCA_002443255.1	GCA_002532875.1
	Assembly size (Mb)	294.13	331.92	368.94	365.59
	Coverage	5x	60x	119x	57x
	Gap size (bp)	0	2,821,918	271,335	408,908
	GC content (%)	-	40.1	40.9	40.9
Scaffold	Scaffolds (n)	-	20,448	1,426	4,881
	N50 scaffold length (bp)	-	128,078	58,536,683	233,810
	L50 scaffold length (bp)	-	703	3	482
Contig	Contigs (n)	184,190	52,152	4,498	8,241
	N50 contig length (bp)	2,262	15,568	201,886	96,009
	L50 contig length (bp)	40,912	6,465	556	1,168
Annotation	Gene number (n)	-	-	12,849	15,486
	Mean length (bp)	-	-	18,634	14,657
	mRNA number (n)	-	-	30,208	26,214
	Mean mRNA length (bp)	-	-	4,193	3,489
	CDS number (n)	-	-	30,208	26,214
	Mean CDS length (bp)	-	-	1,953	1,695
	Intron number (n)	-	-	94,154	93,749
	Mean intron length (bp)	-	-	3,299	2,898
	Exons number (n)	-	-	110,06	111,624
	Mean exons length (bp)	-	-	537	487
Source		from Cornman *et al.*(2010) BMC Genomics		This study	This study

Figure S1: Similar genome sizes between the two species of *Varroa* mites estimated from haploid male genome through the k-mer 42 frequency in GenomeScope.

Identification and confirmation of *Varroa* mite species and mtDNA lineages

Mapping and pairwise whole mitochondrial alignment (16,505 bp) showed that Vdes_3.0 diverged of only 0.1% to the reference *V. destructor* mitogenome (NC004454.2) [66,67] (Figure 2A). Despite presenting similar morphological phenotypes mainly distinguishable through female body size [28,50] (Figure 1A), the *V. jacobsoni* mitochondrial genome diverged from Vdes_3.0 mtDNA at 5.1% of all nucleotides.

Given that *Cox1* is a standard marker used to identify the *Varroa* mite "haplogroup", this region was extracted and realigned with 26 described sequences from native and introduced populations. Divergence within the complete *Cox1* sequences of the *V. destructor* and *V. jacobsoni* reference genomes rose up to 5.9%. The partial *Cox1* sequence (426 bp) of Vdes_3.0 clustered with *V. destructor* sequences and was clearly distinct from Vjacob_1.0 which clustered with *V. jacobsoni* haplotypes (Figure 2B). The concatenated *Cox1, Atp6, Cox3,* and *Cytb* sequence (2,696 bp) from the Vdes_3.0 mitochondrial genome was identical to that of the invasive Korean K1-1/K1-2 sequence (Figure 2C).

By contrast, the *V. jacobsoni* (Vjacob_1.0 i) partial *Cox1* region (426 bp) was a unique new haplotype with 99% nucleotide similarity to Java (one transition point mutation G/A) and Bali (two transitions) haplotypes and ii) the concatenate sequence (only *Cox1, Atp6,* and *Cox3*)

was also a distinct haplotype from the only referenced *V. jacobsoni* Laos 1 and 2. We named

Vjacob_1.0 mtDNA haplotype as Papua New Guinea, which has been formerly detected in field

surveys and reported in host switch events on *A. mellifera* (Roberts, unpublished data).

Figure 2: Inter and intraspecies mtDNA variability across the genus *Varroa* and host switched lineages.

Different mitochondrial markers used in the literature to discriminate the four *Varroa* species (and

unresolved one) based on partial *Cox1* [28] (A,B) and intraspecies lineages using larger *Cox1, ATP6-Cox3*

and *CytB* regions [72,75] (A, C). The two cryptic species *V. destructor* (red) and *V. jacobsoni* (blue) are

genetically divergent as shown by the unrooted phylogenetic tree of *Varroa* mite partial *Cox1* sequences

(B). Variability of *V. jacobsoni* is higher than that of any other *Varroa* species and the reference genome

(blue arrow) corresponded to one of the previously detected haplotypes switching on *A. mellifera* in

Papua New Guinea [30] (B). Several haplotypes from the Korean and Japanese lineages successfully

jumped on *A. mellifera* (bold) [72,75] but only K1-1/K1-2 is quasi-cosmopolitan and can even be retrieved in

the native range of the Japanese lineage as illustrated by its presence in Okinawa (red arrow) (C). Ac:

identified from *Apis cerana*, Am: identified from *Apis mellifera.*

Predicting orthology relationships between *Varroa* mites and other Parasitiformes mites

The arthropod genome database at NCBI presently contains only two other published

representative Mesostigmata genomes: the brood parasitic honey bee mite *Tropilaelaps*

mercedesae (GCA_002081605.1) [83] and the non-parasitic Western predatory mite *Metaseiulus*

(= Typhlodromus or Galendromus) occidentalis (GCA_000255335.1)[102]. Orthologue prediction
was performed by comparing annotated *Varroa* genes to those of the two other Mesostigmata
species gene sets and those of two additional Acari: the parasitic tick *Ixodes scapularis*
(GCA_000208615.1)   and   the   free-living   two-spotted   mite   *Tetranychus   urticae*
(GCA_000239435.1). The tick presents the largest set of genes (n = 20,486) and also possessing
a haploid genome more than four times larger 1.76 Gb [84]. Orthology prediction analysis using
the six genomes clustered the genes by domain into 7,502 super-orthologue groups consisting
of orthologues and in-paralogues [81]. A total of 11,123 orthologue groups were predicted and
42.8% were shared by the five parasitiformes and the outgroup spider mite *Tetranychus*
(Supplementary Table 1). The concatenated phylogenetic tree built on the genomic sets based
on 4,758 genes using IQ-TREE [89] confirmed the close relationship between the three mite
species compared to other Acari (Figure 3). The honey bee ectoparasitic mites, *Varroa* and
Tropilaelaps shared 813 lineage-specific orthogroups (akin to gene families) of which 65.9%
were Varroidae-lineage-specific. The two sibling *Varroa* species shared 9,628 orthologous genes
including 536 which were private to Varroidae.

Orthologous genes specific to the Parasitiformes ancestral branch (*i.e.*, before *T. urticae*
and other Acari species split) was enriched for molecular functions and biological processes
involved in several transporters and channel activities (Supplementary Table 2). When
considering the Mesostigmata ancestral branch (*i.e.*, before *M. occidentalis* and the three
honey bee parasites species split), GO terms related with structural constituents of the cuticle
were the most represented. At the Varroidae ancestral branch level (*i.e.*, before the two species
split), orthologous genes showed GO terms mainly related to endonuclease activity.

435
436

Figure 3: Positive selection and gene duplication of orthologous genes in parasitic and free-living Acari. For the honey bee mite parasites, host ranges are shown as checkboxes and individual records had to meet the following conditions: 1) Mites had to have been found reproducing or observed in bee colonies in independent surveys, and 2) the identity of cryptic *Tropilaelaps* and *Varroa* species was confirmed by molecular markers (e.g. *Cox1* barcoding [28,103]). The number of orthologous genes in each lineage is circled at the nodes. The number of positively selected genes for each branch is shown in green, with the number of species-specific genes shown in purple. Gene duplications within honey bee parasitic mites are shown in boxes: white squares = shared genes, black squares = *Varroa*-specific genes by, red and blue squares = *V. destructor* and V. *jacobsoni* specific genes, respectively. The two *Varroa* mites have similar host ranges, through *V. jacobsoni* also occurs on *A. nigrocincta*, a close relative of *A. cerana*, possibly as a result of a recent secondary host shift [28,50]. Despite sharing the same ancestral host, the *Varroa* sister species show different patterns of adaptive molecular evolution and exhibit different gene expansion patterns, suggesting different evolutionary trajectories.

450
451

Divergent patterns of selection and gene expansion among honey bee ectoparasitic mites

A total of 234 and 225 genes were detected under positive selection for *V. destructor* and *V. jacobsoni* branches, respectively, though only 13 were shared among this list (2.8%)

(Fisher's exact test p = 0.003) (Figure 3). Additionally, 40 genes were detected under positive
selection when looking at the Varroa ancestral branch, *i.e.*, before the two species split. On
these set combined, the sibling species *V. destructor* and *V. jacobsoni* presented 81.0% and
80.4% species-specific genes with positive selection signals, respectively (Figure S2,
Supplementary Table S3). The remaining proportion of positively selected genes shared
involved the other honey bee mite *T. mercedesae* rather than with any other Acari. The
genomic regions associated with genes under positive selection were found and distributed
along all major chromosomes of *V. destructor*, and, assuming preservation of synteny, in *V.*
jacobsoni as well (Figure 4A).

Semantic distribution of the GO terms for the genes under positive selection in each
Varroa species showed non-significant overlap (p = 0.385 > 0.050). The few terms that
overlapped involved biological processes in essential functions such as the molting cycle -
chitin-based, fatty acid, and lipid metabolism and responses to pH (Figure 4B). More
specifically, chitin is the main constituent of the mite cuticle hardening it against both bees and
the environment. Its metabolism varies throughout the mite life cycle, especially during
reproduction or molting [41]. Fatty acid metabolism could be related to the feeding on the fat
body on the host, as well as oogenesis. Although few overlaps exists when GO terms list were
reduced using Resnik's measure, the REVIGO scatterplot and treemap visually suggest
functionally different routes to host adaptation (Figure 4B, Supplementary Table S4). In *V.*
destructor, GO terms were mainly associated with 1) regulation of membrane depolarization, 2)
retinal cone cell development and myofibroblast differentiation, 3) mRNA cis splicing via
spliceosome, 4) cellular response to alcohol (i.e., any process that results in a change or activity
of a cell) whereas GO terms in *V. jacobsoni* related to 1) protein processing involved in targeting
to mitochondria, 2) vitamin K metabolic process, 3) response to pH, 4) gonad development l
and other processes detailed in (Figure S3AB).

Figure S2: UpSet plot of orthologous genes detected under positive selection detailing the shared number per species. Vd Vj Tm Mo ANC = genes shared with the ancestral and modern Mesostigmata (*V. destructor, V. jacobsoni, T. mercedesae,* and *M. occidentalis*), Vd Vj Tm ANC = set of genes shared with the ancestor of Asian mites and Vd Vj ANC = set of genes shared with the ancestral Varroidae.

488
489

Figure 4: Genes and pathways under positive selection in the *Varroa* sister species. (A) Red and blue
bars (in 5 kb windows) represent locations of genes in *V. destructor* and *V. jacobsoni*, respectively. *V.*
jacobsoni data are mapped to *V. destructor* scaffold positions for comparison. Black asterisks indicated
genes under positive selection, shared by both species (n = 12) and detected in the *Varroa* ancestral
lineage prior to the split of the two species (n = 40). (B) Semantic space analysis of significantly enriched
GO terms (BP, CC, and MF) over-represented in genes detected under positive selection for *Varroa* mites
Bubble color indicates the species for which GO terms were enriched (all p-value < 0.05) and size
indicates the frequency of the GO term found in the GOA database. There was little overlap in analyses
at both gene (2.8%) and functional levels (0.8%), suggesting different selective pressures on the two
sister species since they split.
500

501
502 **Figure S3: Treemap plots (REVIGO) analysis for the summary of GO terms in biological processes for**
503 **gene detected under positive selection when comparing *Varroa* species.** Related GO terms were joined
504 in clusters (color) and the size of the rectangles was adjusted to reflect the p-value and with a semantic
505 similarity (C = 0.5).
506
507

Tandem duplication was detected in 51 genes for *V. destructor*, 35 in *V. jacobsoni* and
70 in *T. mercedesae* (Figure 3). Numbers of duplication events and gene clusters within a
species varied from two to six units and generally involved adjacent chromosomal locations in
V. destructor (Figure 5A, Supplementary Table S5). In duplicated genes found only in *V.*
jacobsoni, 23 had no orthologs in the *V. destructor* genome. However, most duplications are on
the same scaffold with the exception of the ABC gene family where genes are present on two
scaffolds (online supplemental data). Manual checks of duplicated clusters with more than two
genes validated all of the duplications except for the Histone 2A family (vd.dup.gene_01) where
gene LOC111255102 located on the minor scaffold BEIS01000444.1 was removed since it is
100% identical to LOC111250948 located on chromosome 1 (BEIS01000001.1) and may be an
artefact of mis-assembly.

Only five duplicated gene clusters were shared by all three honey bee parasite species.
These amounted to 9.8% and 14.3% of detected tandem gene duplications for *V. destructor* and
V. jacobsoni, respectively (Figure 2, Supplementary Table S6). GO terms associated with these
duplicated gene clusters related to chitin, amino sugars, and drug and carbohydrate derivatives.
Conversely, the two *Varroa* sister species shared 14 gene clusters undergoing tandem
duplication, representing 27.5% and 40.0% of the total size for *V. destructor* and *V. jacobsoni*,
respectively. When summarizing the list of duplicated shared genes by both *Varroa* species, GO
terms were associated with biological processes in glycerophospholipid catabolism, body
development, regulation of DNA-templated transcription/elongation and amino-sugar
metabolism (Supplementary Table S7). Finally, 62.7% of the duplicated genes were exclusive to
V. destructor whereas 45.7% were only detected in *V. jacobsoni*. Genes duplicated in *V.*
destructor were associated with GO terms related to biological processes such as
mechanical/external stimulus, regulation of endoribonuclease activity and phagocytosis (Figure
5B, Supplementary Table S7-8). In contrast, genes duplicated in *V. jacobsoni* were involved in
biological processes of the skeletal myofibril assembly, Golgi calcium ion transport, and striated
muscle contraction (Figure 5B, Supplementary Table S7-8).

535
536
537
538

539
540

Figure 5: Duplicated genes in *V. destructor* are found throughout the genome and are involved in
different biological pathways. (A) Chromosomal location and similarity of duplicated genes (arrows) for
V. destructor. Different genes and chromosomal regions underwent duplication in the two species. (B)
Cluster analysis of significantly enriched GO terms for biological processes over-represented among
duplicating genes in *Varroa* mites. Not only were most of the GO terms species-specific, but they also
comprised non-overlapping categories of biological processes. Neither gene duplication nor selection
analysis suggests a substantial degree of parallel evolution in these mites.
548
549　**Chemosensory and cytochrome P450 proteins are conserved in honey bee parasites but**
550　***Varroa* species show different selective trajectories**
551
Lipid carrier proteins: Based on the comparison of six amino acid sequences of OBPs previously
described from *V. destructor* legs transcripts [93], we found seven homologs (with OBP conserved
domain IPR036728 or OBP-like) in *V. destructor* annotated genome (Figure S4). Each OBP gene
found in *V. destructor* presented an ortholog in *V. jacobsoni* coding for an identical OBP
sequence (Supplementary Table S9). An analogous patterns of strong conservation was found
for eight NCP2s (IPR003172) or similar with a Ganglioside GM2 activator conserved domain
(IPR028996) in both sibling Varroa mites (Figure S5). None of the genes encoding for OBP and
NCP2 proteins showed signatures of positive selection or tandem duplication in Varroa sister
species.
561
562

Figure S4: OBP repertoires are highly conserved between *V*arroa sibling species and other Acari. The phylogenetic tree was constructed using 42 amino acid sequences from Acari. *V. destructor* (Vdes = genome and Var = protein sequences from Eliash et al. 2018), *V. jacobsoni* (Vjac), *T. mercedesae* (Tmer), *M. occidentalis* (Mocc), *I. scapularis* (Isca), *T. urticae* (Turt) (see list in Table S9). Additionally, sequences from mosquitos Aedes aegypti (Aaeg), *A. albopictus* (Aaeg) and fruit flies *Drosophila* (D), were downloaded from NCBI accession and aligned with MAFFT. Best fit model computed for the tree is WAG+I+G4 using IQ-TREE. Bootstrap values were estimated using an SH-like aLRT with 1000 and bootstraps over 95% are shown by a purple circle.

Figure S5: *V. destructor* **and** *V. jacobsoni* **genomes showed eight NCP2 amino acid sequences mostly conserved with the other honey bee parasite** *T. mercedesae*. The phylogenetic tree was constructed using 86 amino acid sequences from Acari *V. destructor* (Vdes = genome and Var = protein sequences from Eliash et al. 2018), *V. jacobsoni* (Vjac), *T. mercedesae* (Tmer), *M. occidentalis* (Mocc), *I. scapularis* (Isca) and *T. urticae* (Turt) (see list in Table S9). Additionally, sequences from the fruit fly *Drosophila melanogaster* (Dmel), downloaded from NCBI and aligned with MAFFT. Best fit model computed for the tree is LG+R4 using IQ-TREE. Bootstrap values were estimated using an SH-like aLRT with 1000 and bootstraps over 95% are shown by a purple circle.

Membrane-bound receptors: A total of four genes per Varroa genome were found to encode
for GR proteins presenting either the 7m_7 or Trehalose receptor conserved domains
(IPR013604 and IPR009318, respectively). All four GR were closely related to GRs proteins
found either in the other honey bee parasitic mite *T. mercedesae* or in the parasitic tick *I.
scapularis* (Figure S6). As previously reported, IGR protein profiles are the largest in *V.
destructor.* According to the IGR phylogeny, this is also the case for *V. jacobsoni* with 35
identified genes (mainly presenting IPR001828, IPR019594 or IPR001320 signatures) (Figure 6).
Among these, 19 proteins found in Varroa genomes had closest homolog in *T. mercedesae*
indicating high conservation of the gene family within honey bee parasites. Interestingly, 3
genes encoding for IGR or IGR-like proteins out of the 37 identified in *V. destructor* had no
orthologous in *V. jacobson*i. The opposite situation was also true but only for one gene in *V.
jacobsoni* (LOC111272277). Additionally, divergent selective trajectories were found as a
glutamate [NMDA] receptor subunit 1-like gene (LOC111270396) was under positive selection
for *V. jacobsoni* while a metabotropic glutamate receptor 3-like gene (LOC111245193) show
positive selection signal for *V. destructor*. Finally, despite a small number of SNMPs (with
conserved domain IPR002159) detected in *V. jacobsoni*, we also found that one scavenger
receptor class B member 1-like gene (LOC111267160) was under positive selection but none for
V. destructor (Figure S7).
609
610

Figure S6: ***V. destructor* and *V. jacobsoni* genomes presented only four gustatory receptors**
amino acid sequences like. The phylogenetic tree was constructed using 45 amino acid
sequences from Acari *V. destructor* (Vdes = genome and Var = protein sequences from Eliash et
al. 2018), *V. jacobsoni* (Vjac), *T. mercedesae* (Tmer), *M. occidentalis* (Mocc), *I. scapularis* (Isca)
and *T. urticae* (Turt) (see list in Table S9). Additionally, sequences from the fruit fly *Drosophila*
melanogaster (Dmel) and the honey bee *A. mellifera (*Amel), sequences were downloaded from
NCBI and aligned with MAFFT. The best-fit model computed for the tree is LG+F+R4 using IQ-
TREE. Bootstrap values were estimated using an SH-like aLRT with 1000 and bootstraps over
95% are shown by a purple circle.

Figure 6: The largest IGR repertoire in honey bee parasites show divergent selective

trajectories for host and environment chemosensing between *V. destructor* and *V. jacobsoni*.

The phylogenetic tree was constructed using 153 amino acid sequences from Acari *V. destructor*

(Vdes = genome and Var = protein sequences from Eliash et al. 2018), *V. jacobsoni* (Vjac), *T.*

mercedesae (Tmer), *M. occidentalis* (Mocc), *I. scapularis* (Isca) and *T. urticae* (Turt) which were

aligned with MAFFT (see list in Table S9). Best fit model computed for the tree is VT+R5 using

IQ-TREE. Bootstrap values were estimated using an SH-like aLRT with 1000 and bootstraps over

95% are shown by a purple circle.

Figure S7: Sensory neuron membrane protein repertoires are conserved among honey bee parasites but sibling Varroa species show signs of differential selective regime. The phylogenetic tree was constructed using 67 amino acid sequences from Acari *V. destructor* (Vdes = genome and Var = protein sequences from Eliash et al. 2018), *V. jacobsoni* (Vjac), *T. mercedesae* (Tmer), *M. occidentalis* (Mocc), *I. scapularis* (Isca) (see list in Table S9). Additionally, sequences from the fruit fly *Drosophila melanogaster* (Dmel), were implemented with NCBI accession and aligned with MAFFT. Best fit model computed for the tree is LG+R5 using IQ-TREE. Bootstrap values were estimated using an SH-like aLRT with 1000 and bootstraps over 95% are shown by a purple circle.

Ecdysteroids and Halloween genes: All CYP enzymes sequences from *V. destructor* and *V.*
jacobsoni annotation were extracted by filtering gene description list containing "cytochrome
p450" or exhibiting the associated conserved domain cl12078 or pfam00067. A total of 36 and
38 CYP (one isoform per gene) were aligned together with the seven Halloween genes reported
from *D. melanogaster* and *A. aegypti* and other CYPs from Acari. Previously, only some
members of this pathway, namely orthologues of *spook*, *shade* and *disembodied* were found
from a genomic sequence generated by 454 pyrosequencing [44]. Homologs proteins of
CYP302A1 and CYP314A1 (~32% and ~28% sequence identity to *disembodied* and *shade* of *D.*
melanogaster, respectively) were found for both Varroa sibling species but surprisingly no
homolog *spook* (CYP307A1) was found for *V. jacobsoni*, though there was one in *V. destructor*
(Figure 7)*.* Phylogenetic relationships among Varroa mites, *T. urticae* and *M. occidentalis* and *D.*
melanogaster CYP315A1 sequences (~36% identity with the fruit fly) suggest that both *V.*
destructor and *V. jacobsoni* Varroa mites possess *shadow* homologs. Likely, *shroud* and
phantom are also presents in both Varroa species but not *neverland*. In the CYP family, we
detected three genes under positive selection only in *V. jacobsoni*, including *shadow*-like
(LOC111271433) and disembodied (LOC111265704) (Figure 7 and Table S9).
669

Figure 7: CYP repertoire in Varroa mites is broader than expected with two functionaly important Halloween genes *shade* and *disembodied* under positive selection in *V. jacobsoni*. The phylogenetic tree was constructed using 174 amino acid sequences from Acari *V. destructor* (Vdes = genome and Var = protein sequences from Eliash et al. 2018), *V. jacobsoni* (Vjac), *T. mercedesae* (Tmer), *M. occidentalis* (Mocc), *I. scapularis* (Isca) and *T. urticae* (Turt). Additionally, sequences from the fruit fly *Drosophila melanogaster* (Dmel), mosquito *Aedes aegypti* (Aaeg) and the honey bee *A. mellifera (*Amel) and *A. cerana* (Acer) were downloaded form NCBI and aligned with MAFFT. Best fit model computed for the tree is LG+R5 using IQ-TREE. Bootstrap values were estimated using a SH-like aLRT with 1000 and bootstraps over 95% are shown by a purple circle. *V. destructor* was previously believed to have only *disembodied* (CYP302a1)*, shade* (CYP314a1) and *spook* (CYP307a1) homologs of the seven Halloween genes (background in black)[44].

Other genes possibly related to nutrition and detoxification: Some annotated genes detected under positive selection or undergoing tandem duplication in *Varroa* have been linked to play a role in external stress tolerance (e.g. in-hive temperature environment), nutrition, molting, reproduction, and metabolizing toxic xenobiotics possibly leading to acaricide resistance (detailed in Table 2). On *V. destructor* chromosome 4 (NW_019211457.1), two duplicated genes were also under positive selection and were involved in adult-specific rigid cuticular proteins. These genes are involved in physical properties of the mite cuticle, which in addition to providing protection may constitute the first barrier to reduce the penetration of external chemical agents [104]. On the other hand, two duplicate clusters were also detected undergoing positive selection and annotated as part of the ABC family (ATP-Binding Cassette sub-family A) and to aminopeptidase M1-B-like, in *V. jacobsoni*. Finally, the most remarkable tandem duplication occurred on Vdes_3.0 chromosome 1 (NW_019211454.1) with six duplicated genes coding for heat shock 70 kDa protein. These were not detected for *V. jacobsoni* (Table 2). Both positive gene selection and duplication patterns in the *V. destructor* and *V. jacobsoni* genomes and related GO terms showed that the two sister species have undergone dissimilar evolutionary trajectories.

Table 2: Genes undergoing positive selection or duplication, that have been implicated in tolerance to external stressors and stimuli. Research on Acari and *V. destructor*, in particular, has identified a large number of genes involved in stress and detoxification (possibly leading to acaricide resistance), nutrition, reproduction and chemosensing in interaction with the host. Intriguingly, many of these genes appear to be under ancestral positive selection, as *V. jacobsoni* has not been extensively targeted by acaricides on *A. mellifera*, and most of the evolutionary history captured by positive selection analyses occurred in *A. cerana*, given that coevolution with *A. mellifera* is a relatively recent evolutionary phenomenon, particularly for *V. jacobsoni*.

	V. destructor (Vdes_3.0)		*V. jacobsoni* (Vjacob_1.0)	
Stress from in-hive temperature	**Annotated gene under positive selection**	**Annotated gene undergoing duplication**	**Annotated gene under positive selection**	**Annotated gene undergoing duplication**
Heat Shock 70 Kda protein-like [105]		6		

Nutrition				
Guanine metabolism [106]			1	
Detoxification				
Glutathione-S-transferase [107]			1	
Sodium channel related [108]	4		2	
Transmembrane proteins	3		4	2
G-protein coupled receptors	1			
Esterases [109]	2			
GABA-gated ion channels [110]	1			
ATP-binding cassette gene family [111]	1		2	2
Molting and reproduction				
Ecdysteroids pathway with cytochrome P450 [44]			3	
Cuticular protein	2	4		1
Nutrition and reproduction				
Fatty acid metabolism [97,112,113]	3			
Chemosensory receptors				
Sensory neuron membrane protein			1	
Ionotropic glutamate receptor	1		1	

Discussion

In this study, we developed high-quality *de novo* reference genomes and annotations for economically important honey bee parasites *V. destructor* and *V. jacobsoni*. Because *Varroa* mites are, to a large extent, responsible for the global honey bee crisis, and research efforts worldwide target these parasites, these genomic resources respond to a pressing need [36]. Improved genome assembly and annotation with transcriptomic data from different life stages produced the most comprehensive gene catalogue to date, as reflected by our discovery of previously missing Halloween genes homologs [44]. The supposed absence of such genes has led to speculation control measures development or host counter adaptation could target this pathway[45]. Our analysis also revealed that the two *Varroa* species share 99.7% sequence identity across the nuclear genome. Yet, after the two species have split, they have gone on largely dissimilar evolutionary trajectories, as evidenced by different patterns of positive selection and gene duplication. In particular, signals of positive selection were found in functionally important genes involved in chemosensing, molting, and reproduction. Thus, it seems likely that the two species underwent different patterns of adaptation while continuing to exploit the same host, the eastern honey bee (*A. cerana*). Differences between the two species most likely result from local adaptation to host geographic subspecies, the environment or their interaction. Though not enough data exist to distinguish between these alternatives, below we summarize what is already known and outline promising avenues for investigation.

Inferring what constitutes the ancestral ecological niche of the two mites is key to understanding what selective pressures they have faced and how they responded to them. *V. destructor* and *V. jacobsoni* are very similar morphologically, and until 20 years ago they were mistakenly considered the same species [28]. Their host specificity overlaps as they both parasitize the eastern honey bee *A. cerana* throughout Asia. The only exception reported for the Indonesian *A. nigrocincta* infested by *V. jacobsoni*, though possibly as a result of a recent host switch from *A. cerana* [28,114]. The most recent common ancestor of *V. destructor* and *V. jacobsoni* might have also been a parasite of the *A. cerana* lineage. This hypothesis is supported by the host range of *V. underwoodi*, another closely related mite that also parasitizes *A. cerana* in addition to *A. nigrocincta* and *A. nulensis* [115,116]. With no known varroid (Mesostigmata) fossils, no information exists about speciation timing in *Varroa* mites. Based on the *Cox1* gene, a recognized molecular clock for insects with a divergence rate of 2.3% per Myr [117], the divergence between *V. destructor* and *V. jacobsoni* of 5.9% suggests that the two species split ~2-3 Myr ago. This is after the split between *A. cerana* and *A. mellifera* which occurred 6-9 Myr

ago based on *Cox1* and *ND2* [118,119] to 17-19 Myr using a combination of mitochondrial,
ribosomal and nuclear *loci* [120]. Thus, host ranges and molecular estimates suggest that most
Varroa evolution occurred on ancestors of *A. cerana*.

The major ecological difference between these mites is their parapatric distribution
(Figure 1). Whether parapatry resulted from their respective biogeographic history, adaptation
to different subspecies of *A. cerana*, or is shaped by competition is not known. In fact, even the
extent of the geographical overlap between the mite distributions is uncertain, since few
surveys actually used molecular markers for identification and most have focused on
associations with *A. mellifera* [28,72,116,121]. Yet, parapatry is clearly not due to the inability of the
mites to complete their phoretic life stage outside the hive, as host-switched V. *destructor*
successfully expanded its range following the range extension of *A. mellifera*. While allopatry
could result from adaptation to local hosts/conditions or competitive exclusion, both mites
have been reported from the same Thai apiary on *A. cerana* [46,48]. Unfortunately, there are not
enough data on *A. cerana* infestation under natural conditions to determine the likelihood that
the two *Varroa* species interact and compete. One survey in southern Thailand found 98%
parasitism by *V. jacobsoni* under natural conditions, albeit at variable levels of infestation [122],
suggesting that encounters between the two species are indeed likely. If the divergent
evolutionary history of the two *Varroa* species resulted from competition, we predict that long-
term coexistence within a hive of *A. cerana* should be unlikely, a hypothesis that we hope will
be tested in the future in the zone of sympatry.

In addition, or as an alternative to interspecific competition, it is possible that the
divergence between *V. destructor* and *V. jacobsoni* was driven by local adaptation to different
A. cerana populations. For instance, *V. destructor* was found to infect the "Northern *cerana*",
"Himalayan *cerana*" and the "Indo-Chinese *cerana*" morpho-clusters identified by [47]
whereas *V. jacobsoni* was reported on the "Indo-Chinese *cerana*" and the "Indo-Malayan
cerana". These *A. cerana* populations differ in their nesting behaviour, such as the number of
combs or even in their body size [80]. In China alone, *A. cerana* populations have been shown to
be as genetically divergent for more than 300,000 to 500,000 years as recognized subspecies of
A. mellifera [123]. In addition to genetic differences between the hosts, the climactic
environments across Asia are highly variable and could result in local adaptation, though, as
mentioned earlier, this did not prevent range expansions after switching hosts (Figure 1).
Consequently, the dissimilar selective pressures observed in the mites may have resulted from
selection due to different host genotypes.

Understanding the dynamics of co-infestation by both mites will become important if

the host-switched *V. destructor* and *V. jacobsoni* are ever found to co-occur on *A. mellifera*.

Presently, host-switched *V. jacobsoni* is restricted to Papua New Guinea, where *V. destructor* is

absent [30]. However, if its range expands, it may impose an additional parasitic burden on *A.*

mellifera, but whether the effects of this extra parasite are cumulative or whether the parasites

will simply replace each other locally is unknown. Our data suggest that if the parasites are able

to tolerate each other within *A. mellifera* colonies, given their different suites of molecular

adaptations, they may in fact impose a greater cost than either parasite alone. Clearly, this is

another avenue where further experimental work is urgently needed.

Likewise, it is unknown whether chemical treatment options developed for *V. destructor*

will also be effective against *V. jacobsoni*. Interestingly, both mite genomes show signs of

selection on a range of genes previously found to be involved in stress and pesticide resistance

(Table 2). Since *V. destructor* has evolved resistance to acaricides, it cannot be ruled out that

positive selection resulted from recent acaricide pressures rather than co-evolution with their

host. By contrast, acaricides are not yet widely used on *V. jacobsoni*, though the number of

genes under selection is similar, suggesting that most selection is ancient. In the case of *V.*

jacobsoni, and possibly generally, these genes have the potential to become co-opted for

pesticide resistance. This has been the case with pesticide tolerance in the sheep blowfly

(*Lucilia cuprina*), where sequencing of museum specimens has shown that resistance variants

pre-dated pesticide use [124]. Similarly, the two-spotted mite (*Tetranychus urticae*) appears to

have co-opted existing detoxification machinery in a selective response to miticides [125].

However, more detailed population-level work comparing switched and unswitched

populations of mites will be necessary to test this hypothesis. Museum specimens or other

historical material would be particularly useful in understanding evolutionary trends [126].

Conclusion

Despite the economic importance of brood parasitic mites in *A. mellifera*, it is clear that

various pressing research needs still exist concerning their biology, cryptic diversity in the native

range, possible geographic ranges of host-switched species, and how co-infestations may

proceed. These new reference genomes will allow researchers to pursue such investigations,

develop better tools for *Varroa* control, and to better understand the invasion and evolution of

Varroa on *A. mellifera*. In particular, we hope that these new resources will greatly benefit

future studies to track the population genetic changes and demography of *V. destructor*'s

worldwide spread, which has been difficult due to the low polymorphism of previously available

markers [121]. More *Varroa spp.* are continuously testing new host [73,75] and may remain

undetected. Finally, the recent boom in -omics methods for the study of honey bee parasites

and diseases underlines the need to improve our knowledge of genome structure and

organization to investigate the genetic basis of newly reported acaricide resistance, virulence

level or even new host tolerance. Such knowledge combined with progress in novel

manipulation tools, such as RNAi [107,127], and more experimental tools like gene drive [128], may

provide ways of specifically controlling mite pests without harming honey bees or other insects.

Supplementary data

Figure S1: Similar genome sizes between the two species of Varroa mites estimated from haploid male genome through the k-mer 42 frequency in GenomeScope.

Figure S2: Vdes3.0 clusters with the worldwide invasive V. destructor K1-1/K1-2 haplogroup. Maximum likelihood tree was build on 5000 bootstraps using concatenate sequences of *Cox1, Cox3, Atp6* and *Cytb* partial genes.

Figure S3: UpSet plot of orthologous genes detected under positive selection detailing the shared number per species.

Figure S4: Treemap plots (REVIGO) analysis for the summary of GO terms in biological processes for gene detected under positive selection when comparing Varroa species.

Supplementary table 1: Summary output of orthologs genes and correspondence among the six species included in the phylogeny from ORTHONOME pipeline

Supplementary table 2: GO terms enrichment for ortholog clusters before species split between Varroa mites and with four other Acari

Supplementary table 3: Genomic details of the genes detected under positive either on *V. destructor, V. jacobsoni* or Varroa ancestral (before species split) branch

Supplementary table 04: GO terms enrichment analysis of genes detected under positive selection for *V. destructor, V. jacobsoni* and Varroa ancestral branch (before species split)

Supplementary table 05: Gene duplicated in tandem different between *V. destructo*r and *V. jacobsoni* from the MSOAR2 pipeline

Supplementary table 06: Gene duplicated in tandem shared by Varroa mites and Tropilaelaps

Supplementary table 07: GO terms enrichment analysis of duplicated in tandem genes for *V. destructor* and *V. jacobsoni*

Supplementary table 08: Gene set enrichment analysis of genes duplicated shared by Varroa

mites

Supplementary table 09: Details of the proteins and related genes used for phylogenetic

analysis and comparison of chemosensory proteins

Acknowledgments

We would like to acknowledge Yoann Portugal (Ecology and Evolution OIST) for helping with the

fieldwork collection of *V. destructor* mites, Jo Tan (Ecology and Evolution OIST) for processing in

laboratory *Varroa* males and Hiroki Goto (OIST DNA sequencing section – Onna, Okinawa) for

leading and processing the PacBio RSII sequencing. We would like also to thank Francisco Hita

Garcia (Biodiversity and Biocomplexity OIST) for his support in preparation and scanning of

specimens using the micro-CT. We thank Steven Aird (OIST technical editor) for carefully

checking and helping to correct our manuscript. This work was supported by the Okinawa

Institute of Science and Technology Graduate University, as well as grants from the Japan

Society for the Promotion of Science (Kakenhi grants 16KK0175, 16H06209 and 18H02216) and

the Australian Research Council (Future Fellowship Scheme FT160100178). Genome assembly

was partially supported by Driscoll's, Inc.

Data accessibility

V. destructor (Vdes_3.0) and *V. jacobsoni* (Vjacob_1.0) assembled reference genomes and

annotation are available on NCBI database respectively with the accession number

GCF_002443255 and GCF_002532875. Raw Illumina and PacBio reads are also available in the

Sequence Read Archive (SRA) under the Bioprojects PRJDB6279 and PRJNA391052. Online

supplemental codes and map are available with an R markdown hosted from on

github.com/MaevaTecher/varroa-denovo-genomes.

Author contributions

MAT collected samples for sequencing, carried out analyses, interpreted the data and wrote the

manuscript;

RVR carried out analyses and wrote the manuscript;

JMKR collected samples for sequencing;

MLG carried out the assembly and quality analyses;

SS carried out the computational analyses for Hi-C genome assembly;

IL carried out the experimental protocols for Hi-C genome assembly;

AKC conceived and designed experiments;

JDE conceived and designed experiments;

ASM conceived and designed experiments, carried out the assembly and analyses, interpreted the data

and wrote the manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

IL and STS are employees and shareholders of Phase Genomics, a company commercializing

proximity-ligation technology.

Materials & Correspondence

Maeva A. Techer: maeva.techer@oist.jp

ORCID number: 0000-0001-5417-5103

Alexander S. Mikheyev: alexander.mikheyev@oist.jp

References

1. de Meeûs, T. & Renaud, F. Parasites within the new phylogeny of eukaryotes. *Trends*

Parasitol. **18**, 247–251 (2002).

2. Weinstein, S. B. & Kuris, A. M. Independent origins of parasitism in Animalia. *Biol. Lett.* **12**,

(2016).

3. Poulin, R., Krasnov, B. R., Mouillot, D. & Thieltges, D. W. The comparative ecology and

biogeography of parasites. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **366**, 2379–2390 (2011).

4. Kochin, B. F., Bull, J. J. & Antia, R. Parasite evolution and life history theory. *PLoS Biol.* **8**,

e1000524 (2010).

5. Poulin, R. & Randhawa, H. S. Evolution of parasitism along convergent lines: from ecology

to genomics. *Parasitology* **142 Suppl 1**, S6–S15 (2015).

6. Rigaud, T., Perrot-Minnot, M.-J. & Brown, M. J. F. Parasite and host assemblages:

embracing the reality will improve our knowledge of parasite transmission and virulence.

Proc. Biol. Sci. **277**, 3693–3702 (2010).

7. Schmid Hempel, P. *Evolutionary parasitologythe integrated study of infections,*

immunology, ecology, and genetics. (sidalc.net, 2011).

8. Petney, T. N. & Andrews, R. H. Multiparasite communities in animals and humans:

frequency, structure and pathogenic significance. *Int. J. Parasitol.* **28**, 377–393 (1998).

9. Vaumourin, E., Vourc'h, G., Gasqui, P. & Vayssier-Taussat, M. The importance of

multiparasitism: examining the consequences of co-infections for human and animal

health. *Parasit. Vectors* **8**, 545 (2015).

10. Pouyaud, L., Desmarais, E., Deveney, M. & Pariselle, A. Phylogenetic relationships among

monogenean gill parasites (Dactylogyridea, Ancyrocephalidae) infesting tilapiine hosts

(Cichlidae): systematic and evolutionary implications. *Mol. Phylogenet. Evol.* **38**, 241–249

(2006).

11. Rabajante, J. F. *et al.* Red Queen dynamics in multi-host and multi-parasite interaction

system. *Sci. Rep.* **5**, 10004 (2015).

12. Betts, A., Gray, C., Zelek, M., MacLean, R. C. & King, K. C. High parasite diversity accelerates

host adaptation and diversification. *Science* **360**, 907–911 (2018).

13. Miller, R. S. Pattern and Process in Competition. in *Advances in Ecological Research* (ed.

958 Cragg, J. B.) **4**, 1–74 (Academic Press, 1967).

959 14. Bashey, F. Within-host competitive interactions as a mechanism for the maintenance of

960 parasite diversity. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, (2015).

961 15. Chappell, L. H. Competitive Exclusion between Two Intestinal Parasites of the Three-Spined

962 Stickleback, Gasterosteus aculeatus L. *J. Parasitol.* **55**, 775–778 (1969).

963 16. Simková, A., Morand, S., Jobet, E., Gelnar, M. & Verneau, O. Molecular phylogeny of

964 congeneric monogenean parasites (Dactylogyrus): a case of intrahost speciation. *Evolution*

965 **58**, 1001–1018 (2004).

966 17. Hamelin, F. M., Bisson, A., Desprez-Loustau, M.-L., Fabre, F. & Mailleret, L. Temporal niche

967 differentiation of parasites sharing the same plant host: oak powdery mildew as a case

968 study. *Ecosphere* **7**, e01517 (2016).

969 18. Orr, M., Seike, S. & Gilbert, L. Foraging ecology and patterns of diversification in dipteran

970 parasitoids of fire ants in south Brazil. *Ecol. Entomol.* **22**, 305–314 (1997).

971 19. Ramsey, S. D. *et al.* Varroa destructor feeds primarily on honey bee fat body tissue and not

972 hemolymph. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 1792–1801 (2019).

973 20. Rosenkranz, P., Aumeier, P. & Ziegelmann, B. Biology and control of Varroa destructor. *J.*

974 *Invertebr. Pathol.* **103 Suppl 1**, S96–119 (2010).

975 21. Le Conte, Y., Ellis, M. & Ritter, W. Varroa mites and honey bee health: can Varroa explain

976 part of the colony losses? *Apidologie* **41**, 353–363 (2010).

977 22. Annoscia, D., Del Piccolo, F., Covre, F. & Nazzi, F. Mite infestation during development

978 alters the in-hive behaviour of adult honeybees. *Apidologie* **46**, 306–314 (2015).

979 23. Carreck, N. L., Ball, B. V. & Martin, S. J. Honey bee colony collapse and changes in viral

prevalence associated with Varroa destructor. *J. Apic. Res.* **49**, 93–94 (2010).

24. Gliński, Z. & Jarosz, J. Varroa jacobsoni as a carrier of bacterial infections to a recipient bee

host. *Apidologie* **23**, 25–31 (1992).

25. Neumann, P. & Carreck, N. L. Honey bee colony losses. *J. Apic. Res.* **49**, 1–6 (2010).

26. Steinhauer, N. *et al.* Drivers of colony losses. *Curr Opin Insect Sci* **26**, 142–148 (2018).

27. Eickwort, G. C. Evolution and Life-History Patterns of Mites Associated with Bees. in *Mites:*

Ecological and Evolutionary Analyses of Life-History Patterns (ed. Houck, M. A.) 218–251

(Springer US, 1994).

28. Anderson, D. L. & Trueman, J. W. Varroa jacobsoni (Acari: Varroidae) is more than one

species. *Exp. Appl. Acarol.* **24**, 165–189 (2000).

29. Crane, E. The varroa mite. *Bee World* **59**, 12 (1978).

30. Roberts, J. M. K., Anderson, D. L. & Tay, W. T. Multiple host shifts by the emerging

honeybee parasite, Varroa jacobsoni. *Mol. Ecol.* **24**, 2379–2391 (2015).

31. Oldroyd, B. P. Coevolution while you wait: Varroa jacobsoni, a new parasite of western

honeybees. *Trends Ecol. Evol.* **14**, 312–315 (1999).

32. Rath, W. Co-adaptation of Apis cerana Fabr. and Varroa jacobsoni Oud. *Apidologie* **30**, 97–

110 (1999).

33. Page, P. *et al.* Social apoptosis in honey bee superorganisms. *Sci. Rep.* **6**, 27210 (2016).

34. Lin, Z. *et al.* Reproduction of parasitic mites Varroa destructor in original and new

honeybee hosts. *Ecol. Evol.* **8**, 2135–2145 (2018).

35. Zhang, Y. & Han, R. A Saliva Protein of Varroa Mites Contributes to the Toxicity toward Apis

cerana and the DWV Elevation in A. mellifera. *Sci. Rep.* **8**, 3387 (2018).

36. Evans, J. D. & Cook, S. C. Genetics and physiology of Varroa mites. *Curr Opin Insect Sci* **26**,

130–135 (2018).

37. Peng, Y.-S., Fang, Y., Xu, S. & Ge, L. The resistance mechanism of the Asian honey bee, Apis

cerana Fabr., to an ectoparasitic mite, Varroa jacobsoni Oudemans. *J. Invertebr. Pathol.* **49**,

54–60 (1987).

38. Lin, Z. *et al.* Go East for Better Honey Bee Health: Apis cerana Is Faster at Hygienic

Behavior than A. mellifera. *PLoS One* **11**, e0162647 (2016).

39. Nazzi, F. & Le Conte, Y. Ecology of Varroa destructor, the Major Ectoparasite of the

Western Honey Bee, Apis mellifera. *Annu. Rev. Entomol.* **61**, 417–432 (2016).

40. McAfee, A., Chan, Q. W. T., Evans, J. & Foster, L. J. A Varroa destructor protein atlas reveals

molecular underpinnings of developmental transitions and sexual differentiation. *Mol.*

Cell. Proteomics **16**, 2125–2137 (2017).

41. Mondet, F. *et al.* Transcriptome profiling of the honeybee parasite Varroa destructor

provides new biological insights into the mite adult life cycle. *BMC Genomics* **19**, 328

(2018).

42. Iovinella, I. *et al.* Proteomic analysis of chemosensory organs in the honey bee parasite

Varroa destructor: a comprehensive examination of the potential carriers for

semiochemicals. *bioRxiv* 260539 (2018). doi:10.1101/260539

43. Eliash, N. *et al.* Chemosensing of honeybee parasite, Varroa destructor: Transcriptomic

analysis. *Sci. Rep.* **7**, 13091 (2017).

44. Cabrera, A. R. *et al.* Three Halloween genes from the Varroa mite, Varroa destructor

(Anderson & Trueman) and their expression during reproduction. *Insect Mol. Biol.* **24**, 277–

1024  292 (2015).

1025 45. Conlon, B. H. *et al.* A gene for resistance to the Varroa mite (Acari) in honey bee (Apis

1026  mellifera) pupae. *Mol. Ecol.* (2019). doi:10.1111/mec.15080

1027 46. Warrit, N., Smith, D. R. & Lekprayoon, C. Genetic subpopulations of Varroa mites and their

1028  Apis cerana hosts in Thailand. *Apidologie* **37**, 19–30 (2006).

1029 47. Radloff, S. E. *et al.* Population structure and classification of Apis cerana. *Apidologie* **41**,

1030  589–601 (2010).

1031 48. Rueppell, O., Hayes, A. M., Warrit, N. & Smith, D. R. Population structure of Apis cerana in

1032  Thailand reflects biogeography and current gene flow rather than Varroa mite association.

1033  *Insectes Soc.* **58**, 445–452 (2011).

1034 49. Chantawannakul, P., de Guzman, L. I., Li, J. & Williams, G. R. Parasites, pathogens, and

1035  pests of honeybees in Asia. *Apidologie* **47**, 301–324 (2016).

1036 50. Dietemann, V. *et al.* Standard methods for varroa research. *J. Apic. Res.* **52**, 1–54 (2013).

1037 51. González-Domínguez, J. & Schmidt, B. ParDRe: faster parallel duplicated reads removal

1038  tool for sequencing studies. *Bioinformatics* **32**, 1562–1564 (2016).

1039 52. Schmieder, R. & Edwards, R. Fast identification and removal of sequence contamination

1040  from genomic and metagenomic datasets. *PLoS One* **6**, e17288 (2011).

1041 53. Zhang, J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: a fast and accurate Illumina Paired-

1042  End reAd mergeR. *Bioinformatics* **30**, 614–620 (2014).

1043 54. Margulies, M. *et al.* Genome sequencing in microfabricated high-density picolitre reactors.

1044  *Nature* **437**, 376–380 (2005).

1045 55. Xue, W. *et al.* L_RNA_scaffolder: scaffolding genomes with transcripts. *BMC Genomics* **14**,

604 (2013).

56. Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a

reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).

57. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

58. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

59. Bickhart, D. M. *et al.* Single-molecule sequencing and chromatin conformation capture

enable de novo reference assembly of the domestic goat genome. *Nat. Genet.* **49**, 643–650

(2017).

60. Burton, J. N. *et al.* Chromosome-scale scaffolding of de novo genome assemblies based on

chromatin interactions. *Nat. Biotechnol.* **31**, 1119–1125 (2013).

61. Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. *Bioinformatics* **27**, 764–770 (2011).

62. Vurture, G. W. *et al.* GenomeScope: fast reference-free genome profiling from short reads.

Bioinformatics **33**, 2202–2204 (2017).

63. Hackl, T., Hedrich, R., Schultz, J. & Förster, F. proovread: large-scale high-accuracy PacBio

correction through iterative short read consensus. *Bioinformatics* **30**, 3004–3011 (2014).

64. Worley, K. C. *et al.* Improving genomes using long reads and Pbjelly 2. in *International*

Plant and Animal Genome Conference Xxii. San Diego, CA 10–15 (2014).

65. Thibaud-Nissen, F., Souvorov, A., Murphy, T., DiCuccio, M. & Kitts, P. *Eukaryotic Genome*

Annotation Pipeline. (National Center for Biotechnology Information (US), 2013).

66. Navajas, M., Le Conte, Y., Solignac, M., Cros-Arteil, S. & Cornuet, J.-M. The complete

sequence of the mitochondrial genome of the honeybee ectoparasite mite Varroa

destructor (Acari: Mesostigmata). *Mol. Biol. Evol.* **19**, 2313–2317 (2002).

67. Evans, J. D. & Lopez, D. L. Complete mitochondrial DNA sequence of the important honey

bee pest, Varroa destructor (Acari: Varroidae). *Exp. Appl. Acarol.* **27**, 69–78 (2002).

68. Sedlazeck, F. J., Rescheneder, P. & von Haeseler, A. NextGenMap: fast and accurate read

mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790–2791 (2013).

69. Wala, J., Zhang, C.-Z., Meyerson, M. & Beroukhim, R. VariantBam: filtering and profiling of

next-generational sequencing data using region-specific rules. *Bioinformatics* **32**, 2029–

2031 (2016).

70. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

arXiv [q-bio.GN] (2012).

71. Garrison, E. Vcflib: A C++ library for parsing and manipulating VCF files. *GitHub*

https://github. com/ekg/vcflib (2012).

72. Navajas, M. *et al.* New Asian types of Varroa destructor: a potential new threat for world

apiculture. *Apidologie* **41**, 181–193 (2010).

73. Beaurepaire, A. L. *et al.* Host Specificity in the Honeybee Parasitic Mite, Varroa spp. in Apis

mellifera and Apis cerana. *PLoS One* **10**, e0135103 (2015).

74. Wang, S. *et al.* Ectoparasitic Mites Varroa underwoodi (Acarina: Varroidae) in Eastern

Honeybees, but not in Western Honeybees. *J. Econ. Entomol.* **112**, 25–32 (2019).

75. Gajic, B. *et al.* Variability of the honey bee mite Varroa destructor in Serbia, based on

mtDNA analysis. *Exp. Appl. Acarol.* **61**, 97–105 (2013).

76.   Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis

1091   Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874 (2016).

1092   77.   Anderson, D. L. & Fuchs, S. Two genetically distinct populations of Varroa jacobsoni with

1093   contrasting reproductive abilities on Apis mellifera. *J. Apic. Res.* **37**, 69–78 (1998).

1094   78.   Zhou, T. *et al.* Identification of Varroa mites (Acari: Varroidae) infesting Apis cerana and

1095   Apis mellifera in China. *Apidologie* **35**, 645–654 (2004).

1096   79.   Solignac, M. *et al.* The invasive Korea and Japan types of Varroa destructor, ectoparasitic

1097   mites of the Western honeybee (Apis mellifera), are two partly isolated clones. *Proc. Biol.*

1098   *Sci.* **272**, 411–419 (2005).

1099   80.   Koetz, A. H. Ecology, Behaviour and Control of Apis cerana with a Focus on Relevance to

1100   the Australian Incursion. *Insects* **4**, 558–592 (2013).

1101   81.   Rane, R. V., Oakeshott, J. G., Nguyen, T., Hoffmann, A. A. & Lee, S. F. Orthonome - a new

1102   pipeline for predicting high quality orthologue gene sets applicable to complete and draft

1103   genomes. *BMC Genomics* **18**, 673 (2017).

1104   82.   Shi, G., Zhang, L. & Jiang, T. MSOAR 2.0: Incorporating tandem duplications into ortholog

1105   assignment based on genome rearrangement. *BMC Bioinformatics* **11**, 10 (2010).

1106   83.   Dong, X. *et al.* Draft genome of the honey bee ectoparasitic mite, Tropilaelaps

1107   mercedesae, is shaped by the parasitic life history. *Gigascience* **6**, 1–17 (2017).

1108   84.   Gulia-Nuss, M. *et al.* Genomic insights into the Ixodes scapularis tick vector of Lyme

1109   disease. *Nat. Commun.* **7**, 10507 (2016).

1110   85.   Hoy, M. A. *et al.* Genome Sequencing of the Phytoseiid Predatory Mite Metaseiulus

1111   occidentalis Reveals Completely Atomized Hox Genes and Superdynamic Intron Evolution.

Genome Biol. Evol. **8**, 1762–1775 (2016).

86.   Grbić, M. *et al.* The genome of Tetranychus urticae reveals herbivorous pest adaptations.

Nature **479**, 487–492 (2011).

87.   Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R. & Pfister, H. UpSet: Visualization of

Intersecting Sets. *IEEE Trans. Vis. Comput. Graph.* **20**, 1983–1992 (2014).

88.   Pond, S. L. K. & Muse, S. V. HyPhy: Hypothesis Testing Using Phylogenies. in *Statistical*

Methods in Molecular Evolution (ed. Nielsen, R.) 125–181 (Springer New York, 2005).

89.   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**,

268–274 (2015).

90.   Falcon, S. & Gentleman, R. Using GOstats to test gene lists for GO term association.

Bioinformatics **23**, 257–258 (2007).

91.   Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and visualizes long lists

of gene ontology terms. *PLoS One* **6**, e21800 (2011).

92.   Resnik, P. Semantic Similarity in a Taxonomy: An Information-Based Measure and its

Application to Problems of Ambiguity in Natural Language. *arXiv [cs.AI]* (2011).

93.   Eliash, N. *et al.* Varroa chemosensory proteins: some are conserved across Arthropoda but

others are arachnid specific. *Insect Mol. Biol.* (2018). doi:10.1111/imb.12553

94.   Katoh, K., Rozewicki, J. & Yamada, K. D. MAFFT online service: multiple sequence

alignment, interactive sequence choice and visualization. *Brief. Bioinform.* (2017).

doi:10.1093/bib/bbx108

95.   Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A. & Minh, B. Q. W-IQ-TREE: a fast online

1134  phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* **44**, W232–5 (2016).

1135 96. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new

1136  developments. *Nucleic Acids Res.* (2019). doi:10.1093/nar/gkz239

1137 97. Frey, E., Odemer, R., Blum, T. & Rosenkranz, P. Activation and interruption of the

1138  reproduction of Varroa destructor is triggered by host signals (Apis mellifera). *J. Invertebr.*

1139  *Pathol.* **113**, 56–62 (2013).

1140 98. Wu, K. & Hoy, M. A. The Glutathione-S-Transferase, Cytochrome P450 and

1141  Carboxyl/Cholinesterase Gene Superfamilies in Predatory Mite Metaseiulus occidentalis.

1142  *PLoS One* **11**, e0160009 (2016).

1143 99. Cornman, S. R. *et al.* Genomic survey of the ectoparasitic mite Varroa destructor, a major

1144  pest of the honey bee Apis mellifera. *BMC Genomics* **11**, 602 (2010).

1145 100. Thomas, G. W. C. *et al.* The Genomic Basis of Arthropod Diversity. *bioRxiv* 382945 (2018).

1146  doi:10.1101/382945

1147 101. De Ruijter, A. & Pappas, N. Karyotype and sex determination of Varroa jacobsoni Oud.

1148  *Varroa jacobsoni Oud. affecting honey bees: present status and needs (ed. R. Cavalloro)*

1149  41–47 (1983).

1150 102. Jeyaprakash, A. & Hoy, M. A. The nuclear genome of the phytoseiid Metaseiulus

1151  occidentalis (Acari: Phytoseiidae) is among the smallest known in arthropods. *Exp. Appl.*

1152  *Acarol.* **47**, 263–273 (2009).

1153 103. Anderson, D. L. & Morgan, M. J. Genetic and morphological variation of bee-parasitic

1154  Tropilaelaps mites (Acari: Laelapidae): new and re-defined species. *Exp. Appl. Acarol.* **43**,

1155  1–24 (2007).

104. Martin, S. J. Acaricide (pyrethroid) resistance in Varroa destructor. **85**, 67–69 (2015).

105. Garrido, P. M. *et al.* Heat shock proteins in Varroa destructor exposed to heat stress and

in-hive acaricides. *Exp. Appl. Acarol.* **76**, 421–433 (2018).

106. Posada-Florez, F. *et al.* Insights into the metabolism and behaviour of Varroa destructor

mites from analysis of their waste excretions. *Parasitology* 1–6 (2018).

107. Campbell, E. M., Budge, G. E. & Bowman, A. S. Gene-knockdown in the honey bee mite

Varroa destructor by a non-invasive approach: studies on a glutathione S-transferase.

Parasit. Vectors **3**, 73 (2010).

108. González-Cabrera, J. *et al.* Novel Mutations in the Voltage-Gated Sodium Channel of

Pyrethroid-Resistant Varroa destructor Populations from the Southeastern USA. *PLoS One*

11, e0155332 (2016).

109. Sammataro, D., Untalan, P., Guerrero, F. & Finley, J. The resistance of varroa mites (Acari:

Varroidae) to acaricides and the presence of esterase. *Int. J. Acarology* **31**, 67–74 (2005).

110. Ménard, C. *et al.* Multiple combinations of RDL subunits diversify the repertoire of GABA

receptors in the honey bee parasite Varroa destructor. *J. Biol. Chem.* **293**, 19012–19024

(2018).

111. Dermauw, W. & Van Leeuwen, T. The ABC gene family in arthropods: comparative

genomics and role in insecticide transport and resistance. *Insect Biochem. Mol. Biol.* **45**,

89–110 (2014).

112. Dmitryjuk, M., Zalewski, K., Raczkowski, M. & Żółtowska, K. Composition of fatty acids in

the Varroa destructor mites and their hosts, Apis mellifera drone-prepupae. *Ann. Parasitol.*

61, 21–26 (2015).

113. Zalewski, K., Zaobidna, E. & Żółtowska, K. Fatty acid composition of the parasitic mite

Varroa destructor and its host the worker prepupae of Apis mellifera : Fatty acids of V.

destructor and its host. *Physiol. Entomol.* **41**, 31–37 (2016).

114. de Guzman, L. I. & Rinderer, T. E. Identification and comparison of Varroa species infesting

honey bees. *Apidologie* **30**, 85–95 (1999).

115. Delfinado-Baker, M. & Aggarwal, K. A new Varroa (Acari: Varroidae) from the nest of Apis

cerana (Apidae). *Int. J. Acarology* **13**, 233–237 (1987).

116. Warrit, N. & Lekprayoon, C. Asian Honeybee Mites. in *Honeybees of Asia* (eds. Hepburn, H.

R. & Radloff, S. E.) 347–368 (Springer Berlin Heidelberg, 2011).

117. Papadopoulou, A., Anastasiou, I. & Vogler, A. P. Revisiting the insect mitochondrial

molecular clock: the mid-Aegean trench calibration. *Mol. Biol. Evol.* **27**, 1659–1672 (2010).

118. Garnery, L., Vautrin, D., Cornuet, J. M. & Solignac, M. Phylogenetic relationships in the

genus Apis inferred from mitochondrial DNA sequence data. *Apidologie* **22**, 87–92 (1991).

119. Arias, M. C. & Sheppard, W. S. Molecular phylogenetics of honey bee subspecies (Apis

mellifera L.) inferred from mitochondrial DNA sequence. *Mol. Phylogenet. Evol.* **5**, 557–566

(1996).

120. Ramírez, S. R. *et al.* A molecular phylogeny of the stingless bee genus Melipona

(Hymenoptera: Apidae). *Mol. Phylogenet. Evol.* **56**, 519–525 (2010).

121. Solignac, M. *et al.* Characterization of microsatellite markers for the apicultural pest Varroa

destructor (Acari: Varroidae) and its relatives. *Mol. Ecol. Notes* **3**, 556–559 (2003).

122. Rath, W. & Drescher, W. Response of Apis cerana Fabr towards brood infested with Varroa

jacobsoni Oud and infestation rate of colonies in Thailand. *Apidologie* **21**, 311–321 (1990).

123. Chen, C. *et al.* Population genomics provide insights into the evolution and adaptation of

the eastern honey bee (Apis cerana). *Mol. Biol. Evol.* (2018). doi:10.1093/molbev/msy130

124. Hartley, C. J. *et al.* Amplification of DNA from preserved specimens shows blowflies were

preadapted for the rapid evolution of insecticide resistance. *Proc. Natl. Acad. Sci. U. S. A.*

103, 8757–8762 (2006).

125. Dermauw, W. *et al.* A link between host plant adaptation and pesticide resistance in the

polyphagous spider mite Tetranychus urticae. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E113–22

(2013).

126. Yeates, D. K., Zwick, A. & Mikheyev, A. S. Museums are biobanks: unlocking the genetic

potential of the three billion specimens in the world's biological collections. *Curr Opin*

Insect Sci **18**, 83–88 (2016).

127. Huang, Z. Y., Bian, G., Xi, Z. & Xie, X. Genes important for survival or reproduction in Varroa

destructor identified by RNAi. *Insect Sci.* **26**, 68–75 (2019).

128. McFarlane, G. R., Whitelaw, C. B. A. & Lillico, S. G. CRISPR-Based Gene Drives for Pest

Control. *Trends Biotechnol.* **36**, 130–133 (2018).

1215

A

Varroa destructor

Varroa jacobsoni

0.5 mm

B

Original host
A. cerana

native range

introduced range

C

New host
A. mellifera

0   1000 km

Varroa cryptic species complex infecting *A. cerana* colonies

● *V. destructor*

■ *V. destructor* / *V. jacobsoni* coexisting on the same site

◆ *V. jacobsoni*

Other Varroa species known to infect *A. cerana*

■ *V. underwoodi*

★ *Varroa sp.* (unresolved)

Varroa species only infecting *A. koschevnikovi*

▲ *V. rindereri*

Mesostigmata mite ectoparasites of *Apis* honey bees

214/234

80/98
204/225

+5 +14 +32   *V. destructor*
+5 +14 +16   *V. jacobsoni*
+5   *T. mercedesae*

21/31   *M. occidentalis*
(predatory mite)

6/10   *I. scapularis*
(blacklegged tick)

3/4   *T. urticae*
(red spider mite)

A. cerana
A. mellifera
A. nigrocincta
A. dorsata
A. laboriosa

0   0.25   0.50   0.75   substitutions / site

Ⓧ Lineage specific orthogroups   **X** Positively selected genes   **X** Species specific positively selected genes

X **X** **X** **X** Genes undergoing tandem duplication   ↔ Shared ortholog duplicated genes

Figure 5.

A

B

DNA-templated transcription elongation

regulation of cell killing

locomotory behavior

membrane permeability

response to external stimulus

cell-cell adhesion

behavior

glycerolipid metabolic process

cell killing

organophosphate catabolic process

rythmic process

nitrogen compound process

cation transport

lymphocyte mediated immunity

GO terms count

Semantic space y

Semantic space x

duplicated only in *V. destructor*

duplicated in both Varroa

chromosomal localization of the orthologs duplicated genes

V. destructor   *V. jacobsoni*   duplicated in both Varroa

Ionotropic glutamate receptors (IGRs)

Cytochromes P450 (CYPs)