The proteome of *Tetrasphaera elongata* is adapted to changing

conditions in wastewater treatment plants

Florian-Alexander Herbst[a], Morten S. Dueholm[a], Reinhard Wimmer[a], Per Halkjær Nielsen[a#]

[a]Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark

#Corresponding author

Per Halkjær Nielsen

Center for Microbial Communities, Department of Chemistry and Bioscience, Fredrik Bajers Vej
7H, Aalborg University, DK-9220, Aalborg, Denmark

Phone: +45 9940 8503

Fax: +45 9814 1808

Running title: Proteome of *Tetrasphaera elongata*

Keywords: Proteomics, Metabolomics, Enhanced biological phosphorus removal, Wastewater
treatment

Abstract

The activated sludge in wastewater treatment plants (WWTP) designed for enhanced biological phosphorus removal (EBPR) experiences periodically changing nutrient and oxygen availability. *Tetrasphaera* is the most abundant genus in Danish WWTP and represent up to 20-30% of the activated sludge community based on 16S rRNA amplicon sequencing and quantitative fluorescence *in situ* hybridization analyses, although it is in low abundance in the influent wastewater. Here we investigate how *Tetrasphaera* can successfully out-compete most other microorganisms in such highly dynamic ecosystems. To achive this, we analyze the physiological adaptations of the WWTP isolate *T. elongata* str. LP2 during an aerobic to anoxic shift by label-free quantitative proteomics and NMR-metabolomics. *Escherichia coli* was used as reference organism as it shares several metabolic capabilities and is regularly introduced to wastewater treatment plants, but without succeeding there. When compared to *E. coli,* only minor changes in the proteome of *T. elongata* were observed after the switch to anoxic conditions. This indicates that metabolic pathways for anaerobic energy harvest were already expressed during the aerobic growth. This allows continuous growth of *Tetrasphaera* immediately after the switch to anoxic conditions. Metabolomics furthermore revealed that the substrates provided were exploited far more efficiently by *Tetrasphaera* than by *E. coli*. These results suggest that *T. elongata* prosper in the dynamic WWTP environment due to adaptation to the changing environmental conditions.

Significance of the study

Members of the genus *Tetrasphaera* are widely distributed and highly abundant in most well-operating WWTPs with EBPR configuration. However, despite their high abundance *in situ*, little is known about their physiology and ecological role. Although the importance of *Tetrasphaera* in engineered wastewater treatment systems is slowly being recognized, additional information is needed to understand the full extent of functions the microorganisms have in many of the essential biological processes in the WWTP. Such information may improve available process models and ultimately lead to better wastewater treatment as well as resource recovery. This study supplies proteomic and metabolomic data on the aerobic/anoxic adaptation of *Tetrasphaera* and provides a hypothesis on how *Tetrasphaera* might compete in dynamic engineered systems.

Introduction

Engineered environments allow for the enrichment of beneficial microorganisms by applying specific environmental conditions that benefit them. In nature, the microorganisms may encounter changes in e.g. temperature or substrate as well as electron acceptor availability, and many of these changes follow irregular dynamic schedules. Microorganisms have developed different physiological adaptations to cope with such changes in carbon and electron donor/acceptor availability and they include use of storage compounds such as polyhydroxyalkanoate (PHA), lipids, glycogen, poly-phosphate (poly-P), elemental sulphur, and nitrate [1–3]. Very often adaptation also involves the expression of specific sets of genes [4], e.g., for acquisition and recycling pathways in case of substrate limitations [5], chaperones and proteases in case of heat [6], catalase or superoxide dismutase in case of oxidative stress [7], or expression of genes for alternative electron acceptors and fermentation in the absence of oxygen [8].

The adaptation of the microbial community to cope with dynamic conditions is exploited in modern wastewater treatment plants (WWTPs). By applying oxic/anoxic alternating phases and/or substrate rich and poor phases (feast-famine) as selective pressures, different microorganisms are enriched that efficiently remove carbon (C), nitrogen (N), and phosphorus (P) from domestic and industrial wastewater [2]. N is removed efficiently by nitrification/denitrification (oxic/anoxic) cycles which convert surplus N to inert N_2 or biomass [9]. P is removed by enhanced biological phosphorus removal (EBPR) by including an anaerobic tank in addition to tanks for nitrification/denitrification and these plants can reach a high effluent water quality in respect to P with no or minimal chemical supplementation. In EBPR plants, poly-P accumulating organisms (PAOs) are enriched [10] and they store excess amounts of intracellular poly-P [2]. The general model assumes that poly-P is used for energy production during an anaerobic phase to take up carbon substrates. These are then stored as polyhydroxyalkanoates (PHAs) which are later oxidized aerobically in carbon-limited oxic

phases for growth and poly-P regeneration. Over time P is depleted from the wastewater as the poly-P production is overall net-positive, and additional P is also incorporated into the biomass. Both can be removed as excess sludge and P can be recovered as raw material, e.g., before [11] or after anaerobic digestion [12].

The general EBPR model is primarily based on studies on the betaproteobacterial *Candidatus* Accumulibacter and the uptake of volatile fatty acids, mainly acetate, during an anaerobic phase, storage as PHA, and later oxidation in the aerobic phase [13,14]. Recently, another mechanism was revealed in PAOs belonging to the actinobacterial genus *Tetrasphaera*. Its members carry out aerobic heterotrophic growth and nitrate respiration, but they can also ferment. All these processes are important for a well-working EBPR process. Surprisingly, *Tetrasphaera* are not able to produce and store PHAs [15]. Instead, experiments using *T. elongata* str. Lp2 under dynamic anaerobic/aerobic conditions showed that they could accumulate free amino acids such as glycine for later oxidation in the aerobic phase [16]. The general model is undoubtedly appropriate for *Ca.* Accumulibacter and may explain many lab-scale and full-scale observations. However, microbial community analyses by fluorescence *in situ* hybridization (FISH) and 16S rRNA gene amplicon sequencing indicate that *Ca.* Accumulibacter accounts for only a small fraction of the PAO population in many EBPR plants and that *Tetrasphaera* is more abundant, occasionally reaching 20-30% of the biomass [17–19].

The aim of this study was to understand how *Tetrasphaera* can successfully compete in highly dynamic EBPR plants. Therefore, the proteome and extracellular metabolome (exometabolome) of aerobically grown *T. elongata* str. Lp2 cells before and after a 3 h anoxic phase were analyzed. *E. coli* str. K-12 was treated in the same way to serve as reference. *E. coli* was chosen as it can grow anaerobically by fermentation and nitrate reduction and it is being introduced in high amounts by the incoming wastewater without constituting any major fraction in WWTPs [18]. Also, *E. coli* is

generally very versatile and can survive under many different conditions. This specific strain is

further one of the best described microorganisms and good annotations as well as literature are

available. As for most bacteria, the expression of the necessary genes for anaerobic energy harvest

is strictly hierarchically controlled [8,20] and adaptation needs some time. The acquired data

indicated that this is not the case for *T. elongata* and that it does indeed show a high level of

metabolic robustness and readiness. This, together with poly-P as energy storage compound and a

metabolic diversity, could partly explain continuous growth and successful competition in the

dynamic WWTP environment.

Materials and Methods

Cultivation and sampling

T. elongata str. Lp2 and *E. coli* str. K-12 were cultivated in modified R2A (minimal) medium [16] to be comparable with previous studies. Inoculation was performed from liquid overnight cultures to an optical density (OD) at 600 nm of 0.01. Initial oxic cultivation was done in 50 ml medium within 250 ml conical flasks (25°C, 150rpm). For the 3 h of anoxic incubation, cultures were transferred to serum flasks, and oxygen was removed by repeatedly replacing the headspace with >99.9% pure N_2. Cultivations were performed in quadruplicates, and whole cultures were sacrificed at the end of the aerobic or anaerobic phase. Growth was assessed by measuring the change in OD at 600 nm of 1 ml culture broth in a cuvette and with total protein concentration (see below).

Metabolomics & Proteomics

All metabolomic and proteomic samples were obtained and measured as four biological replicates. Extracellular metabolites were extracted and analyzed by 600 MHZ NMR as previously described [21]. In short, 15 ml of culture supernatant were lyophilized, rehydrated in 600 µl D2O with TSP as standard, adjusted to a pH of 7, measured by NMR, and analyzed using ChenomX.

For protein extraction, cells were lysed (in 1 % sodium deoxycholate, 50 mM triethylammonium bicarbonate) using the FastPrep-96 Instrument (MP Biomedicals) for 1 min at 1600 rpm and the All FastDNA-96 (MP Biomedicals) lysis matrix. Protein concentrations were assessed by BCA method in triplicates, and approximately 20 µg of protein was subjected to a polyvinylidene fluoride membrane-based proteomic sample preparation [22]. The protocol was adjusted to 20 µg of protein, and the membranes were washed twice with 66% acetonitrile before equilibration with an 8 M urea solution. Tryptic peptides were measured by nLC-MS/MS (Ultimate 3000 coupled to a Q Exactive, Thermo Fisher Scientific, Waltham, USA) applying a 3 h method (~140 min elution window),

details can be found in detail elsewhere [21]. Mass spectra were analyzed by MaxQuant (v.

1.5.3.30) [23] as previously described [21], but with up to 4 allowed missed cleavages. Organism-

specific protein databases in FASTA file format were obtained from UniProt [24]. Proteins were

kept if they could be quantified (at least 2 peptides) in at least three replicates in one condition

(aerobic or anaerobic). Label-free quantification (LFQ) values were used to compare the relative

changes. The abundance data was log2 transformed and missing values were replaced by imputation

from the normal distribution before statistical analysis. Significant changes in abundance were

identified by t-test (two-tailed, permutation-based correction, 250 randomizations, FDR < 5%) in

Perseus [25]. Plots were created using R [26] and the ggplot2 package [27]. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium

(http://www.proteomexchange.org/) via the PRIDE partner repository [28] with the dataset

identifier PXD005211.

Results and Discussion

The genus *Tetrasphaera* is repeatedly observed as the most abundant genus in many EBPR plants

[18,19]. Since dynamic conditions with oxic/anoxic changes of roughly 3-4 h duration are

fundamental to these plants, the reaction to oxygen deprivation under controlled conditions was

investigated using *T. elongata* str. Lp2. This species has been used as the model organism for this

clade of PAOs as it has a relatively high similarity on 16S rRNA gene level to *in situ* abundant

phylotypes [17].

Efficient growth under anoxic conditation

Pure cultures of *E. coli* and *T. elongata* were grown in modified R2A medium (without starch) and

they showed as expected different growth patterns. *T. elongata* grew considerably slower than *E.*

coli and required approx. three times as long time to reach a similar amount of biomass before the

start of the anaerobic phase (24 vs. 7 hours). During the anoxic growth period, relative and absolute

growth were at least as high for *T. elongata* as for *E. coli* as determined by biomass differences as

assessed by the change in OD and total protein concentration (Table 1).

Cost-effective use of substrates

To obtain detailed information of the substrate usage and the production of fermentation products,

the medium was analyzed by NMR at different growth stages (Fig. 1). *T. elongata* was found to use

considerably less sugars and amino acids for growth compared to *E. coli*. Surprisingly, *T. elongata*

first depleted trehalose in the medium and subsequently exploited glucose and aspartate. Other

amino acids were not visibly used as substrates, although *T. elongata* can metabolize all the

measured amino acids in glucose-free medium. Previous genomic analyses suggested that *T.*

elongata might produce lactate as fermentation product [15]. This could not be observed in this

study and emphasizes the need for methods that provide direct information about phenotypic

behaviour, such as metabolomics. Under the conditions applied, the only fermentation product

observed in significant amounts was succinate. Both *E. coli* and *T. elongata* showed increased

succinate production during anaerobic incubation.

Substrates were not a limiting factor in the experiments (Fig. 1), although provided in low

concentrations, as they are in WWTPs and thus the oxic/anoxic switch can be seen as the primary

environmental variable.

Minor changes in the proteomes upon shift to anoxic conditions

It was possible to quantify 1318 and 1224 proteins or 42.7% and 28.5% of the theoretical proteome

for *T. elongata* and *E. coli*, respectively (up to 1703 identified proteins, see Dataset1.xlsx). Out of

these, 228 and 171 were significantly altered in abundance (FDR 5%), but with higher maximum

fold changes in *E. coli* (Fig. 2a). Classification of the differentially expressed proteins based on

COG class IDs [29] revealed considerable larger changes in the abundance of proteins related with

energy production and conversion, amino acid metabolism and transport, carbohydrate metabolism

and transport, and inorganic ion transport and metabolism for *E. coli* compared to for *T. elongata*

(Fig. 2b).

Looking at specific changes for proteins involved in energy production and stress response, the

proteomic similarities between the two organisms were primarily restricted to an aspartate

ammonia-lyase (AspA) and, interestingly, to an ATP-dependent RNA helicase, which was the most

significantly down-regulated protein in both proteomes: DeaD and HelY in *E. coli* and *T. elongata*,

respectively. These RNA helicases are required to adapt to different environmental situations [36].

AspA provides fumarate, presumably for reduction to succinate, from aspartate. For *E. coli*, it has

been shown that AspA is part of the NarL regulon [20]. These proteomic adaptations fit the

observed depletion of aspartate in the medium. The aspartate-succinate conversion has been

reported for several bacteria [32–34] and yeast [35] and might be a general anaerobic adaptation

and be preferred to the phosphoenolpyruvate to oxaloacetate conversion (slight, but significant up-

regulation in *T. elongata* of PckG). Although succinate was undoubtedly the dominant fermentative

product for *T. elongata*, no fumarate reductase [38] for fermentation is annotated in its genome. For

B. subtilis it was shown that the 'aerobic' succinate dehydrogenase could fulfil this role [39] and *T.*

elongata might do likewise. No other fermentation products were of importance to *T. elongata*.

Although it depleted lactate under aerobic conditions, it did not produce any during anaerobic

incubation. However, the lactate dehydrogenase (Ldh) could be identified in both conditions. *E. coli*

did produce high amounts of lactate in the absence of oxygen (and other terminal electron

acceptors), and the necessary lactate dehydrogenase (LdhA) was significantly more abundant. This

assumingly led to additional acidic stress which was compensated by the expression of a lysine

decarboxylase (CadA) which is regarded as a major acid stress enzyme [30]. Apart from the

increased abundance of AspA, the hierarchical control of anaerobic gene expression [20,37] was

evident only for *E. coli* (Fig. 3). Enzymes for fermentation or nitrate reduction were identified

already under aerobic conditions for *T. elongata* and did not change significantly in their relative

abundance during anaerobic incubation. In contrast to this, enzymes for nitrate reduction or

fermentation were not identified before or showed clear up-regulation during anaerobic incubation

for *E. coli* (Fig. 3). The presence of the anaerobic metabolic machinery for energy production in *T.*

elongata explains their ability to ignore the anaerobic shock and keep a steady growth. Under

relatively stable conditions like in the human gut this will probably be a costly disadvantage, but in

dynamic systems like wastewater treatment plants with regular environmental changes, this might

be a major cost-saving advantage.

PAO-metabolism in T. elongata

As a PAO, poly-P accumulation and degradation in *T. elongata* are of special interest. One central

protein, which might be indicative of a PAO physiology, is the low-affinity Pit phosphate

transporter [31]. In the well-described PAO *Ca.* Accumulibacter it seems to drive VFA uptake in

the anaerobic phase [32]. In accordance with previously described observations, the abundance of

Pit did not show any significant change between oxic and anaerobic conditions for *T. elongata*.

Neither did the high-affinity Pst system. A small (~1.3 fold), but significant, upregulation in the

anaerobic phase could be observed for the polyphosphate kinase Ppk2. In contrast to Ppk1, which is

widely conserved in bacteria [33] and did not show any significant change, Ppk2 regenerates GTP

[34] as well as ATP [35] by utilizing poly-P, potentially providing additional energy during

anaerobic conditions. Another poly-P utilizing enzyme which was slightly, but significantly, more

abundant was the polyphosphate glucokinase PpgK. PpgK serves as an alternative hexokinase,

which can phosphorylate glucose to glucose-6-phosphate by exploiting poly-P instead of ATP [36],

thus, providing glucose-6-phosphate for glycolysis while preserving ATP. Apart from poly-P,

glycogen is supposed to be the major element in *Tetraphaeras*'s PAO metabolism. Based on

genomic and biochemical analyses, free amino acids and glycogen are the major C storage under

anaerobic conditions [15,37]. A recent comprehensive *in situ* study using Raman spectroscopy

could not verify the accumulation of glycogen in *Tetrasphaera* [38]. In this study, several putative

glycogen related enzymes for synthesis (GlgB, GlgC, N0E1Q7, N0E176) and degradation (GlgX,

GlgP) could be identified. Only N0E1Q7, a 1,4-alpha-glucan branching enzyme, showed a

statistically significant, but very small (~ 1.2 fold) change in abundance under aerobic conditions.

This change was contrary to the assumption that glycogen synthesis is required during anaerobic

conditions and carbon storage in *Tetrasphaera* remains a question of interest.

Overall, the data suggest that *T. elongata*, in contrast to *E. coli*, is pre-adapted to anaerobic

conditions with the pathways necessary for energy production at the ready. If oxygen as a terminal

electron acceptor is unavailable, *E. coli* reacts by first using alternative electron acceptors like

nitrate or fumarate and ultimately relies on fermentation. As for most bacteria, the expression of the

necessary genes is strictly hierarchically controlled [8,20] and adaptation needs time. The acquired

data indicated that this is not the case for *T. elongata* and that it does indeed show a high level of

metabolic robustness and readiness [39]. The theoretical advantage of being able to keep growing

without major metabolic adaptations will be enhanced under *in situ* conditions as *T. elongata* can

store excessive amounts of poly-P for rapid supplementation of energy or accumulation of carbon

substrates under anaerobic conditions [16]. The proteomic and metabolomic data helps to

complement the existing metabolic model [15] (Fig. 4) and gives an explanation for the abundance

of *Tetrasphaera*. Although *Ca.* Accumulibacter and *Tetrasphaera* differ in central aspects of their

metabolism and substrate preferences, they might follow a similar strategy. In that sense, no marked

effects on relative protein abundances were observed across an EBPR cycle for *Ca.* Accumulibacter

[40,41]. On the other hand, clear indications of regulation could be found on transcriptome level for

Ca. Accumulibacter [42] which might not have been recognizable due to sample complexity and

technical challenges. Unfortunately, transcription and translation do not always correlate [43] or are

at least time-delayed [44] and conclusions can be challenging. Here, it was possible to observe the

small but significant effects on proteome level which might not be observable in complex *in situ*

experiments. A question that arises is, if the observations are entirely correct on the single-cell level

or if community level adaptations are responsible. *T. elongata* has a more pronounced tendency to

aggregate compared to *E. coli*. An advantage in WWTPs, but a disadvantage for cultivation in the

laboratory. This study was performed at low cell-densities, vigorous shaking in large flasks with

little culture volumen to reduce aggregation and its impact on physiology. Nevertheless, aggregates

cannot be excluded which would lead to pre-adaptation of some *T. elongata* cells which enable

constant growth in regularly changing environments. In favor of the results reflecting single-cell

adaptations is that, e.g., the nitrate reductase (alpha and beta subunits) did not change in abundance

which might be expected if all cells are now challenged with oxygen limitation (instead of a minor

fraction that is already expressing the enzyme) as was observed for *E. coli*. *T. elongata*'s behavior is

somewhat paradox. *T. elongata* is preventively expressing unnecessary genes (e.g., nitrate

reduction) while showing an economical metabolism with relatively low growth rates and without

strong reactions to environmental stimuli (at least oxygen limitation). Of course, predicting *in situ*

responses from *in vitro* data must be done carefully, but the observations and the concluded

hypothesis fit *in situ* observations and will guide future enrichment reactor and *in situ* studies.

Concluding remarks

As expected, *E. coli* grew initially faster than *T. elongata*, but *T. elongata* grew consistently during

the 3 h anaerobic switch. *T. elongata* behaved more economically to produce similar amounts of

biomass. Furthermore, *T. elongata* mainly used succinate fermentation as an electron sink, whereas

E. coli produced succinate, acetate, formate/CO_2, and high amounts of lactate. The proteomic data

lead to the identification of roughly 200 statistically significantly regulated proteins for both

organisms, but the regulation in *T. elongata* resembled more a fine-tuning of already present

pathways whereas *E. coli* underwent major rearrangements. Analyses of enriched pathways in *E.

coli* showed a clear down-regulation of pathways necessary for translation and aerobic respiration

as well as an up-regulation of anaerobic respiration, fermentation, and severe stress (probably due to

lactic acid production) when challenged with oxygen limitation. At the same time, *T. elongata*

already expressed necessary pathways under aerobic conditions, ignoring the classic hierarchical

control of anaerobic gene expression and just fine-tuned its metabolism. While this strategy might

not be suitable for many natural environments, it seems well suited for engineered habitats like

WWTPs and their scheduled dynamics.

Acknowledgments

289 This study was supported by the Danish Research Council for Independent Research (FNU), grant

290 no. 4002-00455B. The NMR laboratory at Aalborg University is supported by the Obelske Family,

291 SparNord, and Carlsberg Foundations. We thank the PRIDE team for providing data deposition

292 support.

Conflict of interest statement

295 The authors have declared no conflict of interest.

References

[1] van Loosdrecht, M.C.M., Pot, M.A., Heijnen, J.J., Importance of bacterial storage polymers in

bioprocesses. *Water Sci. Technol.* 1997, 35.

[2] Nielsen, P.H., Saunders, A.M., Hansen, A.A., Larsen, P., Nielsen, J.L., Microbial communities

involved in enhanced biological phosphorus removal from wastewater--a model system in

environmental biotechnology. *Curr. Opin. Biotechnol.* 2012, 23, 452–459.

[3] Salman, V., Bailey, J. V, Teske, A., Phylogenetic and morphologic complexity of giant sulphur

bacteria. *Antonie Van Leeuwenhoek* 2013, 104, 169–86.

[4] López-Maury, L., Marguerat, S., Bähler, J., Tuning gene expression to changing environments: from

rapid responses to evolutionary adaptation. *Nat. Rev. Genet.* 2008, 9, 583–593.

[5] Merchant, S.S., Helmann, J.D., Elemental economy: microbial strategies for optimizing growth in the

face of nutrient limitation. *Adv. Microb. Physiol.* 2012, 60, 91–210.

[6] Rosen, R., Ron, E.Z., Proteome analysis in the study of the bacterial heat-shock response. *Mass

Spectrom. Rev.* 2002, 21, 244–265.

[7] Cabiscol, E., Tamarit, J., Ros, J., Oxidative stress in bacteria and protein damage by reactive oxygen

species. *Int. Microbiol.* 2000, 3, 3–8.

[8] Unden, G., Becker, S., Bongaerts, J., Schirawski, J., Six, S., Oxygen regulated gene expression in

facultatively anaerobic bacteria. *Antonie Van Leeuwenhoek* 1994, 66, 3–22.

[9] Seviour, R.J., Nielsen, P.H., *Microbial ecology of activated sludge*, IWA Publishing, 2010.

[10] Nielsen, P.H., Mielczarek, A.T., Kragelund, C., Nielsen, J.L., et al., A conceptual ecosystem model of

microbial communities in enhanced biological phosphorus removal plants. *Water Res.* 2010, 44,

5070–88.

[11] Takiguchi, N., Kishino, M., Kuroda, A., Kato, J., Ohtake, H., A laboratory-scale test of anaerobic

digestion and methane production after phosphorus recovery from waste activated sludge. *J. Biosci.*

Bioeng. 2004, 97, 365–8.

[12] Nyberg, U., Aspegren, H., Andersson, B., Jorgensen, P.E., Jansen, J. la C., Circulation of phosphorus

in a system with biological p-removal and sludge digestion. *Water Sci. Technol.* 1994, 30, 293–302.

[13] García Martín, H., Ivanova, N., Kunin, V., Warnecke, F., et al., Metagenomic analysis of two

enhanced biological phosphorus removal (EBPR) sludge communities. *Nat. Biotechnol.* 2006, 24,

1263–1269.

[14] McMahon, K.D., Read, E.K., Microbial contributions to phosphorus cycling in eutrophic lakes and

wastewater. *Annu. Rev. Microbiol.* 2013, 67, 199–219.

[15] Kristiansen, R., Nguyen, H.T.T., Saunders, A.M., Nielsen, J.L., et al., A metabolic model for

members of the genus *Tetrasphaera* involved in enhanced biological phosphorus removal. *ISME J.*

2013, 7, 543–54.

[16] Nguyen, H.T.T., Kristiansen, R., Vestergaard, M., Wimmer, R., Nielsen, P.H., Intracellular

accumulation of glycine in polyphosphate-accumulating organisms in activated sludge, a novel

storage mechanism under dynamic anaerobic-aerobic conditions. *Appl. Environ. Microbiol.* 2015, 81,

4809–18.

[17] Nguyen, H.T.T., Le, V.Q., Hansen, A.A., Nielsen, J.L., Nielsen, P.H., High diversity and abundance

of putative polyphosphate-accumulating *Tetrasphaera*-related bacteria in activated sludge systems.

FEMS Microbiol. Ecol. 2011, 76, 256–67.

[18]   McIlroy, S.J., Saunders, A.M., Albertsen, M., Nierychlo, M., et al., MiDAS: the field guide to the
340   microbes of activated sludge. *Database* 2015, 2015, bav062.

341   [19]   Stokholm-Bjerregaard, M., McIlroy, S.J., Nierychlo, M., Karst, S.M., et al., A critical assessment of
342   the microorganisms proposed to be important to enhanced biological phosphorus removal in full-scale
343   wastewater treatment systems. *Front. Microbiol.* 2017, 8, 718.

344   [20]   Goh, E.-B., Bledsoe, P.J., Chen, L.-L., Gyaneshwar, P., et al., Hierarchical control of anaerobic gene
345   expression in *Escherichia coli* K-12: the nitrate-responsive NarX-NarL regulatory system represses
346   synthesis of the fumarate-responsive DcuS-DcuR regulatory system. *J. Bacteriol.* 2005, 187, 4890–9.

347   [21]   Herbst, F.-A., Danielsen, H.N., Wimmer, R., Nielsen, P.H., Dueholm, M.S., Label-free quantification
348   reveals major proteomic changes in *Pseudomonas putida* F1 during the exponential growth phase.
349   *Proteomics* 2015, 15, 3244–3252.

350   [22]   Berger, S.T., Ahmed, S., Muntel, J., Cuevas Polo, N., et al., MStern blotting–Hhgh throughput
351   polyvinylidene fluoride (PVDF) membrane-based proteomic sample preparation for 96-well plates.
352   *Mol. Cell. Proteomics* 2015, 14, 2814–2823.

353   [23]   Cox, J., Mann, M., MaxQuant enables high peptide identification rates, individualized p.p.b.-range
354   mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* 2008, 26, 1367–1372.

355   [24]   Uniprot Consortium, Activities at the Universal Protein Resource (UniProt). *Nucleic Acids Res.* 2014,
356   42, D191-8.

357   [25]   Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., et al., The Perseus computational platform for
358   comprehensive analysis of (prote)omics data. *Nat. Methods* 2016.

[26]   R Core Team, R: A Language and Environment for Statistical Computing 2014.

[27]   Wickham, H., *ggplot2: elegant graphics for data analysis*, Springer New York, 2009.

[28]   Vizcaíno, J.A., Côté, R.G., Csordas, A., Dianes, J.A., et al., The PRoteomics IDEntifications (PRIDE)

database and associated tools: status in 2013. *Nucleic Acids Res.* 2013, 41, D1063-9.

[29]   Tatusov, R.L., Galperin, M.Y., Natale, D.A., Koonin, E. V, The COG database: a tool for genome-

scale analysis of protein functions and evolution. *Nucleic Acids Res.* 2000, 28, 33–6.

[30]   Harden, M.M., He, A., Creamer, K., Clark, M.W., et al., Acid-adapted strains of *Escherichia coli* K-

12 obtained by experimental evolution. *Appl. Environ. Microbiol.* 2015, 81, 1932–41.

[31]   McIlroy, S.J., Albertsen, M., Andresen, E.K., Saunders, A.M., et al., 'Candidatus Competibacter'-

lineage genomes retrieved from metagenomes reveal functional metabolic diversity. *ISME J.* 2014, 8,

613–24.

[32]   Saunders, A.M., Mabbett, A.N., McEwan, A.G., Blackall, L.L., Proton motive force generation from

stored polymers for the uptake of acetate under anaerobic conditions. *FEMS Microbiol. Lett.* 2007,

274, 245–51.

[33]   Zhang, H., Ishige, K., Kornberg, A., A polyphosphate kinase (PPK2) widely conserved in bacteria.

Proc. Natl. Acad. Sci. 2002, 99, 16678–16683.

[34]   Ishige, K., Zhang, H., Kornberg, A., Polyphosphate kinase (PPK2), a potent, polyphosphate-driven

generator of GTP. *Proc. Natl. Acad. Sci.* 2002, 99, 16684–16688.

[35]   Nocek, B., Kochinyan, S., Proudfoot, M., Brown, G., et al., Polyphosphate-dependent synthesis of

ATP and ADP by the family-2 polyphosphate kinases in bacteria. *Proc. Natl. Acad. Sci.* 2008, 105,

17730–17735.

[36]   Szymona, M., Ostrowski, W., Inorganic polyphosphate glucokinase of Mycobacterium phlei.

Biochim. Biophys. Acta 1964, 85, 283–95.

[37]   Marques, R., Santos, J., Nguyen, H., Carvalho, G., et al., Metabolism and ecological niche of

Tetrasphaera and Ca. Accumulibacter in enhanced biological phosphorus removal. *Water Res.* 2017,

122, 159–171.

[38] Fernando, E.Y., McIlroy, S.J., Nierychlo, M., Herbst, F.-A., et al., Resolving the individual

contribution of key microbial populations to enhanced biological phosphorus removal with Raman-

FISH. *bioRxiv* 2018, 387795.

[39] Félix, M.-A., Barkoulas, M., Pervasive robustness in biological systems. *Nat. Rev. Genet.* 2015, 16,

483–496.

[40] Wilmes, P., Andersson, A.F., Lefsrud, M.G., Wexler, M., et al., Community proteogenomics

highlights microbial strain-variant protein expression within activated sludge performing enhanced

biological phosphorus removal. *ISME J.* 2008, 2, 853–864.

[41] Wexler, M., Richardson, D.J., Bond, P.L., Radiolabelled proteomics to determine differential

functioning of Accumulibacter during the anaerobic and aerobic phases of a bioreactor operating for

enhanced biological phosphorus removal. *Environ. Microbiol.* 2009, 11, 3029–3044.

[42] Oyserman, B.O., Noguera, D.R., del Rio, T.G., Tringe, S.G., McMahon, K.D., Metatranscriptomic

insights on gene expression and regulatory controls in *Candidatus* Accumulibacter phosphatis. *ISME

J.* 2016, 10, 810–22.

[43] Taniguchi, Y., Choi, P.J., Li, G.-W., Chen, H., et al., Quantifying *E. coli proteome* and transcriptome

with single-molecule sensitivity in single cells. *Science* 2010, 329, 533–538.

[44] Gedeon, T., Bokes, P., Delayed protein synthesis reduces the correlation between mRNA and protein

fluctuations. *Biophys. J.* 2012, 103, 377–85.

Figure 1

NMR-quantified exometabolites. (a) carbohydrates, (b) fermentative products, and (c) amino acids in the initial medium as well as for the end of the aerobic and anoxic cultivations of *E. coli* and *T. elongata*.

Figure 2

Overview of differentially expressed proteins for the end of the aerobic and anoxic cultivations of *E. coli* and *T. elongata*. (a) Volcano plot for quantified (after imputation) proteins from *T. elongata* and *E. coli*. Significance testing was corrected for multiple hypotheses testing at an FDR of 5%. Vertical red lines indicate a twofold change in abundance. The horizontal line reflects a p-value threshold of 0.05. (b) Classification of differentially expressed proteins based on COG functional classes. One-letter abbreviations for the functional categories: J, translation, including ribosome structure and biogenesis; L, replication, recombination and repair; K, transcription; O, molecular chaperones and related functions; M, cell wall structure and biogenesis and outer membrane; N, secretion, motility and chemotaxis; T, signal transduction; P, inorganic ion transport and metabolism; C, energy production and conversion; G, carbohydrate metabolism and transport; E, amino acid metabolism and transport; F, nucleotide metabolism and transport; H, coenzyme metabolism; I, lipid metabolism; D, cell division and chromosome partitioning; R, general functional prediction only; S and "-", no functional prediction.

Figure 3

Normalized abundance of selected proteins for the end of the aerobic and anoxic cultivations of *E. coli* and *T. elongata*. The estimated abundance of each protein in each organims based on LFQ values was nomalized based on the average abundance of that protein across samples.

Figure 4

Updated metabolic model of *Tetrasphaera elongata* str. Lp2. Red arrows indicate increased activity under anaerobic conditions. Pink ovals are storage compounds, lilac ovals are substrates, and blue ovals are fermentation products.

Table 1

Growth of *E. coli* and *T. elongata* after shift to anoxic conditions. Growth was determined by optical density (OD) at 600 nm and protein concentration after cell lysis. The mean and standard deviation of four measurement is presented.

	OD_{600nm}		Protein concentration	
	E. coli	*T. elongata*	*E. coli*	*T. elongata*
Aerobic phase (t = 0 hr)	0.183 ± 0.003	0.174 ± 0.026	107.8 ± 0.8 µg/mL	78.4 ± 3.1 µg/mL
Anoxic phase (t = 3 hr)	0.214 ± 0.009	0.209 ± 0.032	114.0 ± 5.1 µg/mL	88.7 ± 1.8 µg/mL
Absolute increase	0.031 ± 0.009	0.036 ± 0.015	6.2 ± 5.3 µg/µL	10.3 ± 3.8 µg/µL
Relative increase	16.8 ± 4.8 %	20.7 ± 9.2 %	5.8 ± 5.0 %	13.2 ± 5.3 %