Title :

Intrinsic plasticity of cerebellar Purkinje cells in motor learning circuits

Dong Cheol Jang[1,2,*], Hyun Geun Shim[2,3,*], Sang Jeong Kim[2,3,4,5]

[1] Department of Brain and Cognitive Science, College of Natural Science, Seoul National University, 1, Gwanak-

ro, Gwanak-gu, Seoul, Republic of Korea, 08826

[2] Department of Physiology, College of Medicine, Seoul National University, 103, Daehak-ro, Jongno-gu, Seoul,

Republic of Korea, 03080

[3] Department of Biomedical Science, College of Medicine, Seoul National University, 103, Daehak-ro, Jongno-

gu, Seoul, Republic of Korea, 03080

[4] Neuroscience Research Institute, College of Medicine, Seoul National University, 103, Daehak-ro, Jongno-gu,

Seoul, Republic of Korea, 03080

[5] Lead contact

[*] These authors contributed equally to this work

Corresponding author: Sang Jeong Kim (sangjkim@snu.ac.kr)

Abstract

Intrinsic plasticity of cerebellar Purkinje cells (PCs) is recently highlighted in the cerebellar local circuits, however,

its physiological impact on the cerebellar learning and memory remains elusive. Using a mouse model of memory

consolidation deficiency, we found that the intrinsic plasticity of PCs may be involved in motor memory

consolidation. Gain-up training of the vestibulo-ocular reflex produced a decrease in the synaptic weight of PCs

in both the wild-type and knockout groups. However, intrinsic plasticity was impaired only in the knockout mice.

Furthermore, the observed defects in the intrinsic plasticity of PCs led to the formation of improper neural

plasticity in the vestibular nucleus (VN) neurons. Our results suggest that the synergistic modulation of intrinsic

and synaptic plasticity in PCs is required for the changes in the local connectivity between the cerebellum and

VN that contribute to the long-term storage of motor memory.

Introduction

It is widely believed that the cellular basis of memory is derived from modifications of synaptic transmission, such as long-term potentiation (LTP) and long-term depression (LTD) (Kandel et al., 2014). In addition, changes in intrinsic neuronal excitability (intrinsic plasticity) may be an additional mechanism through which memory is stored in the brain (Daoudal and Debanne, 2003; Zhang and Linden, 2003). However, the impact of intrinsic plasticity on the information storage is unknown

For decades, since Ito proposed the flocculus hypothesis (Ito, 1982), numerous studies have demonstrated that synaptic plasticity between parallel fibers (PFs) and cerebellar Purkinje cells (PCs) is the key mechanism of VOR learning (De Zeeuw et al., 1998; Hansel et al., 2006; Schonewille et al., 2010). However, it has recently been proposed that the synaptic plasticity at the PF-PC synapse is not sufficient to explain motor learning (Ito, 2013; Ke et al., 2009; Schonewille et al., 2011; Wulff et al., 2009). Emerging evidence suggests that neural plasticity at multiple sites, including the cerebellar cortex and vestibular nucleus (VN), is required for motor learning (Boyden et al., 2004; Clopath et al., 2014; Porrill and Dean, 2007; Yamazaki et al., 2015). Furthermore, it has been suggested that motor memory is formed in cortical areas through PF-PC plasticity at the early phase of adaption and transferred to sub-cortical area, especially in the VN (Ito, 2013). In the motor learning circuits, cerebellar PCs integrate the information and then project their output signal to the VN. Interestingly, synaptic plasticity at mossy fibers to VN neurons show dependency on the activity of PCs (McElvain et al., 2010; Medina, 2010). This suggests that the activity-dependent modulation of the output of PCs might derive adequate memory transduction from the cerebellar cortex to the VN. Interestingly, we previously verified that the excitability of PCs shows bi-directionality in response to the different patterns of synaptic plasticity induction (Shim et al., 2017) and postulated that the intrinsic plasticity of PCs plays a role in VOR learning, based on former results (Ryu et al., 2017).

Here, we provide insight into the circuit mechanism through which the intrinsic plasticity of cerebellar PCs may be required for long-term memory storage. We used a memory consolidation deficit mouse model, in which stromal interaction molecule 1 (STIM1) was knocked out PC specifically (STIM1^{PKO}) (Ryu et al., 2017), and applied cerebellum-dependent eye movement learning. The VOR learning is concomitant with synaptic and intrinsic plasticity in the cerebellar PCs in the wild-type mice. Interestingly, the STIM1^{PKO} mice showed deficient intrinsic plasticity, although synaptic plasticity was induced after learning. To investigate the impact of intrinsic plasticity within the motor learning circuits, we assessed neural activity in the VN neurons following the gain-increase learning. Firing rate potentiation was gradually developed over time,

and excitatory synaptic transmission was also increased after learning. Notably, neither synaptic plasticity

nor intrinsic plasticity of the VN neurons was observed in STIM^{PKO} mice. This implies that the subsequent

increases in the neural activity in the VN neurons may be derived from changes in the cortical output, as

determined by the intrinsic plasticity of the cerebellar PCs.

Materials and Methods

Animal model

We crossed homozygous PCP2-Cre line (B6.129-Tg(Pcp2-cre)2Mpin/J line, Jackson Laboratory) with the

STIM1-floxed line (C57BL/6 background) to generate PC-specific STIM1 knockout line as our recent study (Ryu

et al., 2017). Only male mice were used in all experiments, and procedures were approved by the Institutional

Animal Care and Use Committee of Seoul National University College of Medicine.

Surgery

All surgical procedure was similar to our previous paper (see Ryu et al., 2017). Surgery for head fixation has been

done at 7- to 8- weeks-old. Zoletil (Zoletil 50, Virbac, 15mg/kg) + xylazine (Rompun, Bayer, 15mg/kg) mixture

was applied to anesthetize mice through intraperitoneal injection. After surgery, 24 to 48 hours of recovery time

were given to mice.

Behavior test

The installation and most of procedure were similar to our previous paper (see Ryu et al., 2017). To control

pupil dilatation, physostigmine salicylate solution (Eserine; Sigma Aldrich) was treated with brief isoflurane

anesthetization, and at least 20mins were given to washout the side-effect of anesthetization. The concentration

of eserine solution was constantly increased from 0.1%, 0.15% and 0.2% because of drug resistance. Two sessions

of acclimation were performed. At first, the mouse was fixed onto a restrainer for 15 minutes and experienced

light on and off, several brief visual and vestibular stimulation for checking surgical failure. Secondly, calibration

step was included. We measured three basal ocular-motor responses, which are OKR, dVOR and lVOR. Visual

stimulation via drum rotating was provided in sinusoidal rotation with ±5° for OKR. Vestibular stimulation was

delivered through turn table rotation for dVOR and lVOR with same rotation amplitude as OKR. The only

different between dVOR and lVOR was under light off and on, respectively. Each response was recorded at four

different rotating frequencies 0.1, 0.25, 0.5, 1.0Hz.   To increase 0.5Hz dVOR gain, associative visuo-vestibular

stimulation was applied. Drum and table simultaneously rotated with ±5° of amplitude out of phase. The protocol

contained three 10min training sessions and four check points (Figure 1A). Trained mice went back to its home

cage which placed in complete dark condition until next check point, and after examine memory retention level,

the mouse was sacrificed for *ex vivo* recording. Given stimulus and the response were fitted to sine curves. Gain

was calculated by the ratio of the response amplitude to stimulus amplitude, and the phase differences between

the two sine curves were determined as phase values. Custom built LabView (National Instrument) analysis tool

was used for all these procedures. The level of memory consolidation was calculated by percent ratio of remained

memory to learned memory.

Slice preparation

Coronal cerebellar (flocculus) and brainstem slices of 270 - 320 μm were dissected by vibratome (Leica, VT1200)

from behaviour tested 9 to 11 weeks old male mice in ice-cold NMDG cutting solution contained with the

following (in mM): 93 NMDG, 93 HCl, 2.5 KCl, 1.2 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 Glucose, 5 sodium

ascorbate, 2 Thiourea, 3 Sodium pyruvate, 10 $MgSO_4 \cdot 7H_2O$, 0.5 $CaCl_2 \cdot 2H_2O$ (pH 7.3). The brainstem slices

containing VN were obtained from more rostral part in which the brainstem was attached to the cerebellum. The

coronal plane of the cerebellar and brainstem slices were transferred into recovery chamber containing NMDG-

cutting solution at 32 ˚C for 10 minutes, and then incubated in standard artificial cerebrospinal fluid (aCSF)

contained with the following (in mM): 125 NaCl, 2.5 KCl, 1 $MgCl_2$, 2 $CaCl_2$, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 10

glucose at room temperature for an hour. NMDG-cutting solution and aCSF were oxygenated with 95% O_2-5%

CO_2 (pH 7.4).

Whole cell recording

Brain slices were put onto a submerged recording chamber on the stage of Olympus microscope (BX50WI,

Japan) and perfused with standard aCSF. We used EPC9 amplifier with PatchMaster software (HEKA Elektronik)

and multiclamp 700B amplifier with pClamp 10 (Molecular Device). Sampling frequency of 20 kHz and signals

were filtered at 2 kHz (1 kHz filter for sEPSC). Inhibitory synaptic inputs were totally blocked by 100 μM

picrotoxin (Sigma) in PC recording, and strychnine (1 μM) was added to block glycinergic input in VN recording.

Patch pipettes (3-4 MΩ) were borosilicate glass and filled with internal solution containing the following: 9 KCl,

10 KOH, 120 K-gluconate, 3.48 $MgCl_2$, 10 HEPES, 4 NaCl, 4 Na_2ATP, 0.4 Na_3GTP and 17.5 sucrose (pH 7.25)

for testing *in vitro* recordings, *ex vivo* PC excitability and *ex vivo* VN recordings; 140 CsCl, 4 NaCl, 0.5 $CaCl_2$,

10 HEPES, 2 MgATP and 5 EGTA (pH 7.3) for *ex vivo* sEPSC recording from PCs in the medial part of the

flocculus. To evaluate the excitability of the cerebellar Purkinje cells, we compared gain responses of the neurons.

We injected series of square-wised current steps ranging from +100 pA to +600 pA with increments of 100 pA for

500 ms. The membrane potential of the cerebellar Purkinje cells was held at -70 mV by injecting bias current in

current clamp mode. To measure the excitability of VN neurons, we injected from -150 pA to +150 pA with

increments of 50 pA for 1 s. There was no bias current in current clamp mode to hold membrane potential at these

recordings. All patch clamp data, except for sEPSC recordings, were imported and analyzed by Igor Pro (Wave

Metrics). The sEPSC data were analyzed using Mini Analysis (Synaptosoft). Other recording and analysis details

including plasticity induction protocol were similar to our previous paper (see Shim et al., 2017)

Statistical analysis

As we described above, behavior data was analyzed by custom built LabView (National Instrument) tool, and

electrophysiology data was analyzed by Igor Pro (Wave Matrics) and Mini Analysis (Synaptosoft). All statistical

analysis was performed using Graphpad Prism 7 and Microsoft Excel. One-way ANOVA with Fisher's LSD test,

One- or Two-way repeated measure ANOVA with Sidak's post-hoc test was used for several time groups analysis,

and unpaired *t*-test was performed to compare wild-type and knockout group. All graphs are shown as mean±SEM,

and asterisks *, ** and *** indicates $p<0.05$, $p<0.01$ and $p<0.001$, respectively. *n* for each experiments are written

in the figure legends.

Results

Consolidation deficit in the STIM1[PKO] mouse may occur between 1 to 4 hours after learning.

Accumulating evidence support the theory that eye movement memory is firstly formed in the cerebellar cortex

and is then transferred to the sub-cortical regions (Ito, 2013; Kassardjian et al., 2005; Matsuno et al., 2016;

Okamoto et al., 2011). This sequential memory acquisition within the cortical and sub-cortical areas has been

implicated in long-term memory storage. Interestingly, Okamoto et al., (2011) observed that memory transfer

occurs between 2.5 and 4 hours after learning in mice. This implies that this period of time is the critical period

for communication between the cerebellar cortex and sub-cortical areas (in this circumstance, the VN). Because

we have previously reported that an impairment in long-term memory storage, but not memory acquisition, is

observed in the STIM1[PKO] mice (Ryu et al., 2017), we expected that a dysfunctional memory transfer process

would manifest as a consolidation deficit in this model. To determine whether VOR memory was attenuated in

the STIM1[PKO] mice over the memory transfer period, we selected three time periods after learning task in which

measurements were made: 0.5 and 1 hours two points as short-term period, 4 hours as the mid-term period and 24

hours as the long-term period. The memory retention level was evaluated at each of these periods. Both groups

showed normal basal oculomotor performance (Figure S1) and no memory impairment at the short-term period

(Figure 1B). However, at the mid-term period, the memory retention level was significantly lower in the

STIM1[PKO] group than in the wild-type group. Furthermore, at the long-term period memory retention was

eliminated in the STIM1[PKO] group (Figure 1B). This temporal alteration in the memory retention level was re-

calculated as the ratio of the remaining memory at the test session to the acquired memory at the training session

(Figure 1C). This revealed a showing gradual reduction in memory retention over the studied periods. Given that

the slight decline in the level of memory retention began 1 hour after the learning task and developed further, we

speculated that the impaired motor memory consolidation that was observed in the STIM1[PKO] mice was based on

defect of memory transfer process, leading to inappropriate communication between the cerebellar cortex and the

VN.

Learning generally induces PF-PC synaptic plasticity in both groups

Many previous studies, using various LTD-deficient animal models, have reported that synaptic plasticity at the

PF-PC synapse is strongly correlated with motor learning (Boyden et al., 2006; De Zeeuw et al., 1998; Hansel et

al., 2006). Late-phase LTD has also been implicated in VOR memory consolidation (Ahn et al., 1999; Boyden et

al., 2006). This suggests that a learning-induced long-lasting reduction of cerebellar cortical activity drives the

transduction of memory to the sub-cortical region. Interestingly, because the time course of late-phase LTD is

similar to that of the mid-term period of study, we investigated whether PF-PC LTD is involved in memory

consolidation. To verify this in detail, electrophysiological *ex vivo* recordings were made from floccular PCs to

investigate the signs of synaptic plasticity at the short-, mid- and long-term periods after learning task. This

approach enabled us to monitor neuronal activity for periods of over an hour, overcoming the experimental

limitation of the whole-cell patch clamp technique. Due to the location of micro zone of the flocculus that regulates

horizontal VOR behavior, we recorded from PCs that were located in the medial part of the flocculus (Schonewille

et al., 2006). As a control group, we used sham animals that had undergone surgery and restraint without the

learning task. We firstly measured the PF-stimuli-evoked synaptic response (eEPSC) following the injection of

ranges of electrical stimuli intensities. In these experiments, the amplitude 99-101e of the eEPSC was decreased

at the short- (1hr) and mid-term (4hr) period than sham group. However, this alteration was recovered, with the

eEPSC returning to a level that was not significantly different to that of sham control before the long-term phase

in both group (Figure 2A and B). In addition to the PF-evoked synaptic events, the spontaneous excitatory

postsynaptic current (sEPSC) was recorded after the learning task. Between the sham groups of both genotypes,

there was no significant difference in the spontaneous glutamatergic synaptic transmission (Figure S2). The

distribution of the inter-event intervals (IEIs) of synaptic events was shifted to the right after the learning task and

was restored at the long-term period in both groups (Figure 2C and E). However, the mean frequency of sEPSC

in the STIM1^{PKO} mice was transiently reduced until 1 hour after the learning task, and was recovered 4 hours later,

while in the change of wild-type littermates, the change was maintained throughout all of the time periods after

the learning task (insets of Figure 2C and E). Conversely, the aspect of changes in the sEPSC amplitudes in the

STIM1^{PKO} mice were comparable to that of the wild-type littermates (Figure 2D and F). In both groups, the

distribution of the sEPSC amplitude was left-shifted after the learning task and this change was maintained until

the long-term period. Collectively, these results, as well as the results of a previous study (Boyden et al., 2006),

indicate that VOR gain-up training elicited synaptic weakening of the PF-PC synapses. Although the alterations

of sEPSC frequency that were observed in the STIM1^{PKO} mice were relatively transient compared to the results

observed in the wild-type group, overall, the aspects of plasticity that were induced by VOR learning were similar

between the groups. Therefore, we suggest that the learning-induced PF-PC LTD presumably did not contribute

to the long-term memory deficit observed in the STIM1^{PKO} mice.

Learning-induced intrinsic plasticity shows relevance in memory consolidation

Experience-dependent neural plasticity includes not only synaptic plasticity but also alterations in the intrinsic

excitability (Daoudal and Debanne, 2003; Zhang and Linden, 2003). Because PF-PC LTD was not sufficient to

account memory consolidation, we investigated whether if intrinsic plasticity, the other form of neural plasticity,

could be a considerable factor in memory consolidation. To investigate whether excitability changes of cerebellar

PCs are required for memory consolidation, we firstly performed whole-cell patch clamp recordings to compare

the long-term depression of intrinsic excitability (LTD-IE) in the STIM1^{PKO} group and the wild-type littermates.

A PF burst protocol (7 of 100 Hz PF burst followed by a single CF stimulation; Shim et al., 2017) was introduced

to induce PC synaptic and intrinsic plasticity in the presence of an inhibitory synaptic transmission inhibitor,

picrotoxin. As shown in Figure 2, both STIM1^{PKO} and the wild-type groups showed normal induction of PF-PC

LTD (Figure S3A) and LTD-IE. However, the magnitude of the intrinsic plasticity was weaker in the STIM1^{PKO}

group than in the wild-type group. Interestingly, the reduction in the excitability following LTD induction was

recovered 40 min after the induction in the STIM1^{PKO} mice, whereas, in the wild-type littermates LTD-IE was

elicited and developed further (Figure S3B).

Next, we examined the temporal alteration of PC excitability through *ex vivo* recordings after the learning task;

at short-, mid- and long-term time periods. In agreement with the results of *in vitro* experiments (Shim et al., 2017;

Figure S3A), the firing frequency was decreased 1 hour after training in the wild-type littermates (Figure 3A).

The AP firing frequency of PCs was measured in current clamp mode through the injection of brief current steps

from the membrane potential of approximately -70 mV (500 ms, from +100 pA to +500 pA with an increment of

100 pA, step interval 4.5 s). The learning-induced intrinsic plasticity was partially recovered at the mid-term time

period and fully recovered to the value from the sham control at the long-term period. However, the STIM1[PKO]

group showed a deficiency in the learning-induced intrinsic plasticity throughout the studied periods (Figure 3B).

Comparing the results from the different genotypes over the same time period, wild-type littermates has

significantly higher firing frequency in sham control, but the frequency reversed in the short-term period and

gradually recovered over the time period studied (Figure S3C-F). Given the notion that the intrinsic plasticity of

cerebellar PCs amplifies the modification of synaptic weight to properly project the learned signal from the PCs

to their relay neurons, our observation may imply that the impairment of intrinsic plasticity results in dysfunctional

memory transfer to the VN neurons. Because intrinsic plasticity was almost restored 40 min after plasticity

induction in the STIM1[PKO] group (Figure S3B), we expect that the learning-induced intrinsic plasticity may be

already abolished within 1 hour after training. Taken together, we conclude that the intrinsic plasticity of the

cerebellar PCs would be involved in long-lasting reduction of the cerebellar cortical activity, and thereby, may

contribute to VOR memory consolidation.

Appropriate synaptic and intrinsic plasticity of VN neurons require intrinsic plasticity of cerebellar PC

We observed an impairment of intrinsic plasticity in the cerebellar PCs of the STIM1[PKO] group through *in vitro*

and *ex vivo* recordings. This suggests that memory consolidation requires the transduction of a memory from the

cerebellar cortex into the sub-cortical area through the intrinsic plasticity of PCs. A large population of VN neurons

receive information from floccular PCs (Matsuno et al., 2016; Shin et al., 2011). Furthermore, the output of the

cerebellar cortex serves as an instructive signal in the control the aspect of neuronal plasticity between mossy

fibers (MFs) and VN neurons (Clopath et al., 2014; Dean et al., 2010; McElvain et al., 2010; Medina, 2010; Porrill

and Dean, 2007; Shutoh et al., 2006; Yamazaki et al., 2015). Thus, we hypothesized that the impairment of the

intrinsic plasticity of the cerebellar cortex that was observed in STIM1[PKO] mice would lead to an inadequate

alteration of the VN neuron activity following VOR learning. To address this, we performed *ex vivo* recordings to

investigate spontaneous synaptic transmission after the learning task during three distinct time periods, short-, mid- and long-term by *ex vivo* recordings. In the sham groups, the frequency of sEPSC in the STIM1[PKO] group showed remarkable augmentation compared to the wild-type group (Figure S4A). However, the sEPSC amplitude was not significantly different between the STIM1[PKO] group and wild-type group (Figure S4B). These results imply that the homeostatic scaling in the VN neurons are due to the reduction of PC excitability in the STIM1[PKO] group (Figure S3C). Intriguingly, synaptic transmission was found to be potentiated after VOR learning in the wild-type littermates throughout the periods of study (Figure 4A and 4B). Although the increase in the mean frequency of sEPSC after training seemed to be restored at the long-term period, the cumulative distribution of the IEIs was found to be left-shifted throughout the periods of study (Figure 4A). The cumulative fraction of the sEPSC amplitude was especially right shifted at the long-term period, indicating that the proportion of increased glutamatergic synaptic events was enhanced during this period (Figure 4B). However, the mean value was not significantly altered compared to that of the sham control (Figure 4B, inset). These results indicate that VOR gain-up learning induces LTP at the MF-VN synapse, in line with the previous expectation (Boyden et al., 2006). In contrast to the results presented from the wild-type littermates, the STIM1[PKO] group showed a slight depression of sEPSC frequency in cumulative distribution in the short- and mid-term time periods that continuously recovered to baseline (Figure 4C). However, the mean frequency was not significantly altered among the periods of study (Figure 4C, inset). The amplitude of sEPSC was slightly left-shifted in the short- and mid-term time periods, and mean amplitude in mid-term time period was significantly lower than that of the sham group (Figure 4D). In light of previous reports, which have suggested that cerebellar PC activity contribute to MF-VN plasticity (Dean et al., 2010; Matsuno et al., 2016; McElvain et al., 2010; Medina, 2010), we speculated that the synaptic plasticity at the MF-VN synapse is inappropriately induced in the STIM1[PKO] group due to the absence of PC intrinsic plasticity.

Furthermore, we asked whether the intrinsic plasticity of cerebellar PCs is also required for the adequate induction of intrinsic plasticity in the VN neurons, because VOR training involves a change in the excitability as well as synaptic transmission (Carcaud et al., 2017; Shutoh et al., 2006). To answer this, the gain responses were measured through the injection of square-wised somatic depolarizing current into the VN neurons at the three time periods after the learning task. The VN neurons of the STIM1[PKO] group showed higher firing frequency in response to the current injection than were observed in the wild-type littermates in the sham group (Figure S4C). Interestingly, VOR training elicited the intrinsic plasticity of the VN neurons in the wild-type littermates (Figure 5A), whereas, there was no alteration of the gain responses in the STIM1[PKO] group (Figure 5B). The excitability of the MVN neurons gradually increased over the studied time periods and the intrinsic plasticity was maintained

24 hours after training (Figure 5A, right). The difference between the sham groups of both genotypes was faded

during the short- and the mid-term time period, and finally reversed with statistical significance at the long-term

time period (Figure S4C-S4F). To clarify whether the neural plasticity in the VN neurons is affected by knockout

of STIM1 in the PC, we delivered the conventional protocol for induction of LTP in VN neurons. The VN neurons

from the wild-type and STIM1^{PKO} groups exerted potentiation of the synaptic weight and excitability (Figure S4G,

S4H). Taken together with modification of synaptic weight and intrinsic properties in VN neurons, we suggest

that the intrinsic plasticity of the cerebellar PCs following VOR learning could enable to induction of the proper

forms of neuronal plasticity in VN neurons, corresponding to the specific behavior, such as consolidation of

memory.

Discussion

Here, we demonstrate a role of the intrinsic plasticity of cerebellar Purkinje cells in motor learning through

which the VOR adaptive memory is transferred to the sub-cerebellar cortical region for long-term memory storage.

After gain-up training, the synaptic strength at PF-PC synapses is decreased and learning-induced PF-PC LTD

occurs concomitantly with a reduction of intrinsic excitability in PCs. Furthermore, VOR learning causes

potentiation of the synaptic weight and intrinsic excitability in VN neurons, as well as the plasticity in PCs. Mice

that were subjected to impaired memory consolidation, the STIM1^{PKO} group, show a normal learning curve and

synaptic plasticity, whereas the PC intrinsic plasticity is declined within an hour. In addition to the unstable

induction of intrinsic plasticity, there were no appropriate learning-induced alterations of synaptic transmission

and excitability in the VN neurons. Our observations indicate that experience-dependent modulation of the

neuronal excitability is required for long-term memory consolidation, in terms of the cerebellum-dependent motor

learning.

There have been two long-lasting hypotheses for VOR learning that hold many different points of views. Marr-

Albus-Ito proposed that plasticity in the cerebellar cortex may be the key player in the adaptive eye-movement

motor learning (Albus, 1971; Ito, 1982; Marr, 1969). During the VOR training, PF-PC LTD is established through

the conjunction of head movement-driven PF activation with visuo-vestibular mismatch-driven instructive CF

activation. For decades, the Marr-Albus-Ito theory has been confirmed and expanded by abundant experimental

evidence supporting the contribution of the PF-PC LTD to the adjustment of the VOR gain (De Zeeuw et al., 1998;

Hansel et al., 2006; Schonewille et al., 2010). In spite of several literatures corroborating this hypothesis, there has been an alternative perspective for VOR behavior. Miles and Lisberger, (1981) suggested that the cellular basis for the adaptive motor learning is activity-dependent neural plasticity of the VN neurons guided by an instructive signal from the cerebellar cortex. In parallel with this perspective, several experimental observations have insisted that cerebellar LTD might be not sufficient to be a central dogma of the cellular mechanism for the VOR learning (Alphen and Zeeuw, 2002; Ke et al., 2009; Schonewille et al., 2011). There is, however, accumulating evidence supporting a reconciliatory learning model. (Boyden et al., 2004) proposed multiple plasticity mechanisms, insisting that plasticity at the MF-VN synapse, as well as PF-PC synapses requires the motor memory formation. In addition, VOR gain-up learning has been found to be selectively engaged in the PF-PC LTD and thereby continue to MF-VN LTP (Boyden et al., 2006). The authors pointed out that the output of PCs may be responsible for the VN plasticity as Miles and Lisberger suggested. More recently, numerous computational modeling studies have supported the theory that synaptic plasticity in both regions is required for successful memory storage (Clopath et al., 2014; Dean et al., 2010; Porrill and Dean, 2007; Yamazaki et al., 2015). This theory highlights the importance of the communication between these brain regions. Despite many implications suggesting that long-term storage of motor memory requires the memory transfer process from cerebellar cortex to nuclei, the detailed mechanisms of this memory transfer have yet to be elucidated. The data we presented here provides experimental evidence that VOR training results in the alteration of synaptic weight and excitability at multiple sites, the cerebellar cortex and the VN. Moreover, this study elucidates an unrevealed role of the intrinsic plasticity of cerebellar PCs in the VOR memory circuit by using the memory consolidation deficit mice model (Figure 1). Learning-induced synaptic plasticity at the PF-PC synapse is observed in wild-type littermates (Figure 2), however, intrinsic plasticity is abolished within an hour of the learning task in the memory consolidation deficient mouse model (Figure 3). Furthermore, this impairment of the intrinsic plasticity of PCs is concomitant with the failure of VOR training-induced plasticity induction in the VN neurons (Figure 4), although the VN neurons in the STIM1[PKO] mice are endowed with neural plasticity *in vitro* (Figure S4G and S4H), implying that the learning-induced alteration of excitability in the PCs might serve as an instructive signal to induce the appropriate plasticity induction in VN neurons (Figure S5). These results support the previous expectations in which suggested that PC activity can affect the synaptic and intrinsic plasticity induction in VN neurons (McElvain et al., 2010; Medina, 2010). Collectively, our results reconcile two long-standing hypotheses by providing experimental evidence for the induction of multiple forms of plasticity through VOR learning in both the cerebellar cortex and sub-cortical regions.

It has been assumed that motor memory is firstly formed in the cerebellar cortex and that neurons in the VN are

involved in late phase adaption for VOR gain (Ito, 2013; Shutoh et al., 2006). This assumption implies that the

temporal order between PC and VN plasticity has to be considered in memory processing. In our results, the VN

plasticity is induced at a relatively later period than the plasticity in the PCs, and it indicates two major aspects.

One is that PF-PC LTD contributes to memory acquisition, and the other is that the consequent induction of

plasticity in VN neurons encodes long-term memory storage. Our data indicates that the impaired intrinsic

plasticity of the cerebellar PCs would impair memory transfer and disrupt long-term memory storage. This

supports the theory that the intrinsic plasticity of PCs connects two distinct brain regions and shapes the flow of

information flow from the cerebellar cortex to sub-cortical area. The temporal order of plasticity at multiple sites

may reflect the loci of memory storage. The *ex-vivo* recordings we presented here were executed at distinct time

points: short- (~1hr), mid- (~4hrs) and long-term (~24hrs) periods after learning. At the short-term period, the

VOR learning curve and synaptic plasticity were not impaired in the memory consolidation deficient mouse model,

although intrinsic plasticity was abolished (Figure S5). These results indicate that the memory acquisition may

require synaptic plasticity in the cerebellar cortex, but not intrinsic plasticity. Rather, the aspects of the memory

retention and deficiency of intrinsic plasticity in STIM1[PKO] lead us to assume that the learning-induced alteration

in PC excitability might be involved in the memory transfer process. Consistent with previous implications, our

results suggest that the memory transfer occurs within 4 hours after learning (Kassardjian et al., 2005; Okamoto

et al., 2011; Shutoh et al., 2006). Synergies between synaptic and intrinsic plasticity may provide an instructive

signal to convey the learned information into the sub-cortical area, the VN, at the mid-term (~4hrs) period.

Interestingly, the synaptic plasticity in the VN neurons is observed slightly later than the intrinsic plasticity of the

PC. Additionally, there is another slight delay in the VN intrinsic plasticity to reach peak (Figure S5). These results

indicate that sequential flow of information from the cerebellar cortex to the sub-cortical region is responsible for

memory processing. Taken together, we conclude that the acquired VOR memory might be located in the

cerebellar cortex and the VN at the short- and long-term period, respectively, and a guiding instructive signal,

driven by the intrinsic plasticity of the PCs may take part in the transfer of memory from the cortical area to the

sub-cortical area during the mid-term time period.

It is widely believed that the plasticity of neuronal excitability is involved in the cellular mechanism for memory

storage. In particular, the intrinsic plasticity of cerebellar PCs shows features in the cerebellar memory circuits

that are distinct from other types of neurons. In the neurons in the amygdala and hippocampal, learning-related

neurons show higher excitability (Zhou et al., 2009), and the depolarization of the membrane potential of these

cells enables the promotion of further synaptic plasticity (Ramakers and Storm, 2002; Watanabe et al., 2002).

Thus, these excitable neurons form a stable connection by strengthening the synaptic weight the given neural

network, thereby consolidating the memory. In contrast, one previous study suggested that the intrinsic plasticity

of PCs occludes the subsequent induction of PF-PC synaptic plasticity (Belmeguenai et al., 2010). Hence, the

plasticity of excitability may ensure that synaptic activity remains within a physiological limit by restricting

further synaptic plasticity and adjusting the impact of PF activation on the output of PCs. In addition, our data

show that there is no significant difference in the magnitude of synaptic plasticity at the PF-PC synapses between

the wild-type littermate group and STIM1^{PKO} group, although the excitability is lower in the STIM1^{PKO} group

than the wild-type group (Figure S3 and S4A). This suggests that the basal membrane excitability in PCs is not

correlated with the synaptic plasticity induction or the magnitude of synaptic plasticity.

Given that potentiation of the spontaneous firing rates in cerebellar PCs is not sufficient for affecting the firing

rates in the DCN neurons (Belmeguenai et al., 2010), this implies that potentiation and/or reduction of PC

excitability alone would be unable to significantly influence the neuronal activity in the sub-cortical area. Because

the intrinsic plasticity of PCs is modulated with the same polarity of the PF-PC synaptic plasticity, concurrence

of synaptic and intrinsic plasticity may synergistically produce a PC output in response to external inputs, such as

vestibular stimuli (Belmeguenai et al., 2010; Shim et al., 2017). Furthermore, the *ex-vivo* recordings in this study

reveal that VOR gain-up learning induces PF-PC LTD and LTD-IE in PCs, indicating that the intrinsic plasticity

is accompanied by synaptic plasticity, which corresponds to the activity-pattern (Figure 2 and 3). Collectively, we

suggest that learning-induced intrinsic plasticity may amplify the alteration of the synaptic transmission, resulting

in the synergistic modulation of the net output of PCs in order to maximize information storage.

Acknowledgments

We thanks to Seungha Kim and Yong Gyu Kim for experimental support, and Misun Moon, Geehoon Chung for

reviewing original draft. This study was supported by National Research Foundation of Korea (NRF) Grant funded

by Korean government (MSIP) (2012R1A5A2A44671346 and 2017M3C7A1029611 to S.J.K; Global Ph.D

fellowship programme, 2013H1A2A1034318 to D.C.J). H.G.S. received a scholarship from the BK21-plus

education program provided by the National Research Foundation of Korea (5262-20170100)

Author Contributions

Conceptualization, Investigation, Writing (Original Draft), D.C.J. and H.G.S.; Writing (Review & Editing), D.C.J.,

H.G.S. and S.J.K.; Funding Acquisition, D.C.J. and S.J.K.

Declaration of Interests

388 The authors declare no competing interests.

References

Ahn, S., Ginty, D.D., Linden, D.J., 1999. A late phase of cerebellar long-term depression requires activation of

CaMKIV and CREB. Neuron 23, 559–568.

Albus, J.S., 1971. A theory of cerebellar function. Math Biosci 10, 25–61. doi:10.1016/0025-5564(71)90051-4

Alphen, V.A., Zeeuw, D.C., 2002. Cerebellar LTD facilitates but is not essential for long-term adaptation of the

vestibulo-ocular reflex. European Journal of … 16. doi:10.1046/j.1460-9568.2002.02094.x

Belmeguenai, A., Hosy, E., Bengtsson, F., Pedroarena, C.M., Piochon, C., Teuling, E., He, Q., Ohtsuki, G., Jeu,

M.T., Elgersma, Y., Zeeuw, C.I., Jörntell, H., Hansel, C., 2010. Intrinsic Plasticity Complements Long-

Term Potentiation in Parallel Fiber Input Gain Control in Cerebellar Purkinje Cells. J. Neurosci. 30,

13630–13643. doi:10.1523/JNEUROSCI.3226-10.2010

Boyden, E.S., Katoh, A., Pyle, J.L., Chatila, T.A., Tsien, R.W., Raymond, J.L., 2006. Selective engagement of

plasticity mechanisms for motor memory storage. Neuron 51, 823–834. doi:10.1016/j.neuron.2006.08.026

Boyden, E.S., Katoh, A., Raymond, J.L., 2004. Cerebellum-dependent learning: the role of multiple plasticity

mechanisms. Annu. Rev. Neurosci. 27, 581–609. doi:10.1146/annurev.neuro.27.070203.144238

Carcaud, J., de Barros, F., Idoux, E., Eugène, D., Reveret, L., Moore, L.E., Vidal, P.-P., Beraneck, M., 2017.

Long-lasting visuo-vestibular mismatch in freely-behaving mouse reduces the vestibulo-ocular reflex and

leads to neural changes in the direct vestibular pathway 4, ENEURO. 0290–16.2017.

doi:10.1523/eneuro.0290-16.2017

Clopath, C., Badura, A., Zeeuw, C.I., Brunel, N., 2014. A Cerebellar Learning Model of Vestibulo-Ocular

Reflex Adaptation in Wild-Type and Mutant Mice. J. Neurosci. 34, 7203–7215.

doi:10.1523/JNEUROSCI.2791-13.2014

Daoudal, G., Debanne, D., 2003. Long-term plasticity of intrinsic excitability: learning rules and mechanisms.

Learn. Mem. 10, 456–465. doi:10.1101/lm.64103

De Zeeuw, C.I., Hansel, C., Bian, F., Koekkoek, S.K., van Alphen, A.M., Linden, D.J., Oberdick, J., 1998.

Expression of a protein kinase C inhibitor in Purkinje cells blocks cerebellar LTD and adaptation of the

vestibulo-ocular reflex. Neuron 20, 495–508. doi:10.1016/S0896-6273(00)80990-3

Dean, P., Porrill, J., Ekerot, C.-F.F., Jörntell, H., 2010. The cerebellar microcircuit as an adaptive filter:

experimental and computational evidence. Nat. Rev. Neurosci. 11, 30–43. doi:10.1038/nrn2756

Hansel, C., de Jeu, M., Belmeguenai, A., Houtman, S.H., Buitendijk, G.H.H., Andreev, D., De Zeeuw, C.I.,

Elgersma, Y., 2006. alphaCaMKII Is essential for cerebellar LTD and motor learning. Neuron 51, 835–

843. doi:10.1016/j.neuron.2006.08.013

Ito, M., 2013. Error detection and representation in the olivo-cerebellar system. Front Neural Circuits 7, 1.

doi:10.3389/fncir.2013.00001

Ito, M., 1982. Cerebellar control of the vestibulo-ocular reflex--around the flocculus hypothesis. Annu. Rev.

Neurosci. 5, 275–296. doi:10.1146/annurev.ne.05.030182.001423

Kandel, E.R., Dudai, Y., Mayford, M.R., 2014. The molecular and systems biology of memory. Cell 157, 163–

186. doi:10.1016/j.cell.2014.03.001

Kassardjian, C.D., Tan, Y.-F.F., Chung, J.-Y.J.Y., Heskin, R., Peterson, M.J., Broussard, D.M., 2005. The site

of a motor memory shifts with consolidation. J. Neurosci. 25, 7979–7985. doi:10.1523/JNEUROSCI.2215-

05.2005

Ke, M.C., Guo, C.C., Raymond, J.L., 2009. Elimination of climbing fiber instructive signals during motor

learning. Nat. Neurosci. 12, 1171–1179. doi:10.1038/nn.2366

Marr, D., 1969. A theory of cerebellar cortex 437–470. doi:10.1113/jphysiol.1969.sp008820

Matsuno, H., Kudoh, M., Watakabe, A., Yamamori, T., Shigemoto, R., Nagao, S., 2016. Distribution and

Structure of Synapses on Medial Vestibular Nuclear Neurons Targeted by Cerebellar Flocculus Purkinje

Cells and Vestibular Nerve in Mice: Light and Electron Microscopy Studies. PLoS ONE 11, e0164037.

doi:10.1371/journal.pone.0164037

McElvain, L.E., Bagnall, M.W., Sakatos, A., Lac, du, S., 2010. Bidirectional plasticity gated by

hyperpolarization controls the gain of postsynaptic firing responses at central vestibular nerve synapses.

Neuron 68, 763–775. doi:10.1016/j.neuron.2010.09.025

Medina, J.F., 2010. A Recipe for Bidirectional Motor Learning: Using Inhibition to Cook Plasticity in the

Vestibular Nuclei 68, 607–609. doi:10.1016/j.neuron.2010.11.011

Miles, F.A., Lisberger, S.G., 1981. Plasticity in the vestibulo-ocular reflex: a new hypothesis. Annual review of

neuroscience.

Okamoto, T., Endo, S., Shirao, T., Nagao, S., 2011. Role of cerebellar cortical protein synthesis in transfer of

memory trace of cerebellum-dependent motor learning. J. Neurosci. 31, 8958–8966.

doi:10.1523/JNEUROSCI.1151-11.2011

Porrill, J., Dean, P., 2007. Cerebellar Motor Learning: When Is Cortical Plasticity Not Enough? 3, 1935–1950.

doi:10.1371/journal.pcbi.0030197

Ramakers, G.M.J., Storm, J.F., 2002. A postsynaptic transient K(+) current modulated by arachidonic acid

regulates synaptic integration and threshold for LTP induction in hippocampal pyramidal cells. Proc. Natl.

Acad. Sci. U.S.A. 99, 10144–10149. doi:10.1073/pnas.152620399

Ryu, C., Jang, D.C., Jung, D., Kim, Y.G., Shim, H.G., Ryu, H.-H., Lee, Y.-S., Linden, D.J., Worley, P.F., Kim,

S.J., 2017. STIM1 Regulates Somatic Ca2+Signals and Intrinsic Firing Properties of Cerebellar Purkinje

Neurons. J. Neurosci. 37, 8876–8894. doi:10.1523/JNEUROSCI.3973-16.2017

Schonewille, M., Luo, C., Ruigrok, T.J., Voogd, J., Schmolesky, M.T., Rutteman, M., Hoebeek, F.E., De Jeu,

M.T., De Zeeuw, C.I., 2006. Zonal organization of the mouse flocculus: physiology, input, and output. J.

Comp. Neurol. 497, 670–682. doi:10.1002/cne.21036

Schonewille, M., Belmeguenai, A., Koekkoek, S.K., Houtman, S.H., Boele, H.J., van Beugen, B.J., Gao, Z.,

Badura, A., Ohtsuki, G., Amerika, W.E., Hosy, E., Hoebeek, F.E., Elgersma, Y., Hansel, C., De Zeeuw,

C.I., 2010. Purkinje cell-specific knockout of the protein phosphatase PP2B impairs potentiation and

cerebellar motor learning. Neuron 67, 618–628. doi:10.1016/j.neuron.2010.07.009

Schonewille, M., Gao, Z., Boele, H.-J.J., Veloz, M.F., Amerika, W.E., Simek, A.A., De Jeu, M.T., Steinberg,

J.P., Takamiya, K., Hoebeek, F.E., Linden, D.J., Huganir, R.L., De Zeeuw, C.I., 2011. Reevaluating the

role of LTD in cerebellar motor learning. Neuron 70, 43–50. doi:10.1016/j.neuron.2011.02.044

Shim, H.G., Jang, D.C., Lee, J., Chung, G., Lee, S., Kim, Y.G., Jeon, D.E., Kim, S.J., 2017. Long-Term

Depression of Intrinsic Excitability Accompanied by Synaptic Depression in Cerebellar Purkinje Cells. J.

Neurosci. 37, 5659–5669. doi:10.1523/JNEUROSCI.3464-16.2017

Shin, M., Moghadam, S.H., Sekirnjak, C., Bagnall, M.W., Kolkman, K.E., Jacobs, R., Faulstich, M., Lac, du, S.,

2011. Multiple types of cerebellar target neurons and their circuitry in the vestibulo-ocular reflex. J.

Neurosci. 31, 10776–10786. doi:10.1523/JNEUROSCI.0768-11.2011

Shutoh, F., Ohki, M., Kitazawa, H., Itohara, S., Nagao, S., 2006. Memory trace of motor learning shifts

transsynaptically from cerebellar cortex to nuclei for consolidation. Neuroscience 139, 767–777.

doi:10.1016/j.neuroscience.2005.12.035

Watanabe, S., Hoffman, D.A., Migliore, M., Johnston, D., 2002. Dendritic K+ channels contribute to spike-

timing dependent long-term potentiation in hippocampal pyramidal neurons. Proc. Natl. Acad. Sci. U.S.A.

99, 8366–8371. doi:10.1073/pnas.122210599

Wulff, P., Schonewille, M., Renzi, M., Viltono, L., Sassoè-Pognetto, M., Badura, A., Gao, Z., Hoebeek, F.E.,

van Dorp, S., Wisden, W., Farrant, M., De Zeeuw, C.I., 2009. Synaptic inhibition of Purkinje cells

mediates consolidation of vestibulo-cerebellar motor learning. Nat. Neurosci. 12, 1042–1049.

doi:10.1038/nn.2348

Yamazaki, T., Nagao, S., Lennon, W., Tanaka, S., 2015. Modeling memory consolidation during posttraining

periods in cerebellovestibular learning. Proc. Natl. Acad. Sci. U.S.A. 112, 3541–3546.

doi:10.1073/pnas.1413798112

Zhang, W., Linden, D.J., 2003. The other side of the engram: experience-driven changes in neuronal intrinsic

excitability. Nat. Rev. Neurosci. 4, 885–900. doi:10.1038/nrn1248

Zhou, Y., Won, J., Karlsson, M.G., Zhou, M., Rogerson, T., Balaji, J., Neve, R., Poirazi, P., Silva, A.J., 2009.

CREB regulates excitability and the allocation of memory to subsets of neurons in the amygdala. Nat.

Neurosci. 12, 1438–1443. doi:10.1038/nn.2405

Figure legends

Figure 1. Long-term memory storage was impaired in STIM^{PKO} mice.

(A) Illustration of the VOR test and training (upper) and experimental scheme (below). For training, visuo-vestibular stimulation was delivered through drum and turntable, 10 min each session, three times; dVOR at 0.5 Hz was recorded in before the learning and after each session of training. After whole training session, animals were put in the completely dark condition.

(B) Normalized gain of the eye movement in learning. Note there is no significant differences between wild-type and STIM1^{PKO} mice in learning (points on white-colored background; wild-type n=36, STIM1^{PKO} n=39). We measured memory retention level at 0.5 and 1 hour (described as short-term; wild-type n=21, STIM1^{PKO} n=21), 4 hours (mid-term; wild-type n=10, STIM1^{PKO} n=11) and 24 hours (long-term period; wild-type n=13, STIM1^{PKO} n=17) after training (points on grey-dotted background). STIM1^{PKO} showed significantly lower memory retention level from the mid-term period compared to the wild-type littermates (4hr, p=0.037; 24hr; p=0.004).

(C) Calculated consolidation level. Memory retention level was obtained by calculating the ratio remained to learned memory (4hr, p <0.001; 24hr, p<0.001).

Two-way repeated measure ANOVA was used for panel B, and asterisks in the graph were marked by post-hoc Sidak test for pairwise comparison. Unpaired *t*-test was used for panel C. Error bars denote SEM, *p<0.05, **p<0.01, ***p<0.001.

Figure 2. Long-term depression at the PF-PC synapses after gain-up learning.

(A) Representative eEPSCs traces of both genotype groups in each time point. Scale bars, 200pA (vertical) and 30ms (horizontal).

(B) Amplitude of eEPSC by serial PF stimulation in wild-type littermates. In comparison to sham group (n=9), the amplitude was significantly reduced at 1 hour and 4 hours after learning (at 50 μA injection; 1hr, n=19, p<0.001; 4hr, n=9, p=0.002), and depressed amplitude was recovered 24 hours after (at 50μA injection; 24hr, n=8, p=0.922).

(C) Amplitude of eEPSC by serial PF stimulation in STIM1^{PKO}. Same as wild-type littermates, the amplitude was considerably decreased at 1 and 4 hours after, and restored at 24 hours after learning (at 50 μA injection;

sham, n=14; 1hr, n=14, p<0.001; 4hr, n=14, p=0.013; 24hr, n=7, p=0.232).

(D) Representative sEPSC traces of wild-type group in each time point. Scale bars, 25pA (vertical) and 1s

(horizontal).

(E) Cumulative plots of inter-events-interval (IEI) of sEPSC in wild-type littermates. The cumulative fraction

was right-shifted after learning, implying reduction of frequency (sham n=15, 1hr n=15, 4hr n=8, 24hr n=8).

Inset bar graph is mean frequencies of sEPSC indicating depression of frequency was maintained until 24

hours after learning in comparison to sham group (1hr, p<0.001; 4hr, p<0.001; 24hr, p=0.029).

(F) Cumulative plots of amplitude of sEPSC in wild-type littermates. The cumulative fraction was left-shifted

after learning, implying reduction of amplitude. Inset bar graph is mean amplitudes of sEPSC indicating that

depression of amplitude was maintained for 24 hours after learning (1hr, p <0.001; 4hr, p<0.001; 24hr,

p<0.001).

(G) Representative sEPSC traces of STIM^{PKO} group in each time point. Scale bars, 25pA (vertical) and 1s

(horizontal).

(H) Cumulative plots of inter-events-interval (IEI) of sEPSC in STIM^{PKO} mice. The cumulative fraction was right-

shifted 1 hour after learning, but most of changes returned to sham level from 4 hours after training (sham

n=15, 1hr n=13, 4hr n=14, 24hr n=10). Inset shows summarizing bar graph of sEPSC frequency, implying

that frequency of sEPSC was transiently depressed at 1 hour and recovered from 4 hours after learning (1hr,

p <0.001; 4hr, p<0.001; 24hr, p<0.001).

(I) Cumulative plots of amplitude of sEPSC in STIM^{PKO} mice. The cumulative fraction was left-shifted after

learning. Inset bar graph shows that amplitude of sEPSC was depressed and maintained for 24 hours after

learning (1hr, p =0.001; 4hr, p<0.001; 24hr, p=0.016).

Two-way repeated measure ANOVA was used for panel A and B, and asterisks in the graph were marked by post-

hoc Sidak test for pairwise comparison. One-way ANOVA with Fisher LSD test was used for insets in panel C,

D, E and F. Asterisks in each time points were calculated by comparing to sham groups. Error bars denote SEM.

*p<0.05, **p<0.01, ***p<0.001.

Figure 3. VOR learning also induces plastic change of intrinsic excitability in PCs.

(A) Representative traces from whole-cell recording in wild-type group. Scale bars (upper), 20mV (vertical) and

200ms (horizontal). Scale bars (lower), 20mV (vertical) and 50ms (horizontal).

(B) PC excitability of wild-type littermates over several time points. VOR learning decreased gain responses of

the cerebellar PCs in response to square-wised current injection ranging from 100 pA to 600 pA for 500 ms

(sham vs 1hr, p<0.001; sham vs 4hr, p=0.024; sham vs 24hr, p=0.717, left; sham, n=20; 1hr, n=11; 4hr, n=16;

24hr, n=20). Excitability in 600 pA injection was significantly decreased at short-term (1hr, p=0.002) and

mostly recovered at mid-term (4hr, p=0.196) and fully recovered at long-term (24hr, p=0.914).

(C) Representative traces from whole-cell recording in STIM1[PKO] group. Scale bars (upper), 20mV (vertical) and

200ms (horizontal). Scale bars (lower), 20mV (vertical) and 50ms (horizontal).

(D) Excitability of PC in STIM1[PKO]. Different from wild-type littermates, STIM1[PKO] showed no alteration of

excitability after learning (sham vs 1hr, p=370; sham vs 4hr, p=0.343; sham vs 24hr, p=0.768, left; sham,

n=17; 1hr, n=13; 4hr, n=17; 24hr, n=13). There were no significant changes in 600 pA injection (1hr, p=0.371.;

4hr, p=0.184; 24hr, p=0.814)

Two-way repeated measure ANOVA was used for injected current-frequency graphs. One-way ANOVA with

Fisher LSD test was used for bar graphs. Asterisks in each time points were calculated by comparing to sham

groups. Error bars denote SEM. **p<0.01.

Figure 4. VOR gain-up learning induced long-term depression of the excitatory input in VN neurons.

(A) Representative sEPSC traces of wild-type group in each time point. Scale bars, 25pA (vertical) and 1s

(horizontal).

(B) Inter-events-interval (IEI) of sEPSC in wild-type mice. The cumulative distributions of IEI were left-shifted

after learning (sham, n=23; 1hr, n=15; 4hr, n=16; 24hr, n=16). The frequency of sEPSC was potentiated at

short- (1hr) and mid-term (4hr) after training (inset bar graph; 1hr, p=0.046; 4hr, p=0.013; 24hr, p=0.242).

(C) Amplitude of sEPSC in wild-type mice. The cumulative distribution was right-shifted at 24 hours after

learning. There was trend of potentiation at long-term (24hr) after training, but overall, the amplitude of

sEPSC was not significantly affected by learning (inset bar graph; 1hr, p=0.850; 4hr, p=0.874; 24hr, p=0.122).

(D) Representative sEPSC traces of STIM1[PKO] group in each time point. Scale bars, 25pA (vertical) and 1s

(horizontal).

(E) Inter-events-interval (IEI) of sEPSC in STIM[PKO] mice. In contrast to wild-type littermates, the cumulative

distribution was slightly right-shifted after learning (sham, n=14; 1hr, n=21; 4hr, n=111; 24hr, n=15). The

mean frequency of sEPSC was not significantly altered after learning (inset bar graph; 1hr, p=0.613; 4hr,

p=0.066; 24hr, p=0.314).

(F) Cumulative plots of amplitude of sEPSC in STIM[PKO] mice. The cumulative distribution was slightly left-

shifted at 4 hours after learning. The amplitude of sEPSC was significantly reduced at mid-term (4hr) after

learning (inset bar graph; 1hr, p=0.111; 4hr, p=0.030; 24hr, p=0.420).

One-way ANOVA with Fisher LSD test was used for bar graphs in all panels. Asterisks in each time points were

calculated by comparing to sham groups. Error bars denote SEM. *p<0.01

Figure 5. VOR gain-up learning induced potentiation of intrinsic excitability in VN neurons.

(A) Representative traces from whole-cell recordings of wild-type group in each time point. Scale bars (upper),

20mV (vertical) and 500ms (horizontal). Scale bars (lower), 20mV (vertical) and 125ms (horizontal).

(B) The excitability of VN neurons in wild-type littermates. VOR learning significantly potentiated gain

responses of the VN neurons in response to square-wised current injection ranging from -150 pA to 150 pA

for 1 s (sham vs 1hr, p=0.319; sham vs 4hr, p=0.072; sham vs 24hr, p=0.002, left; sham, n=23; 1hr, n=16;

4hr, n=38; 24hr, n=19). Excitability in 150 pA injection was significantly increased at mid- (4hr, p=0.032)

and long-term (24hr, p<0.001) after training.

(C) Representative traces from whole-cell recordings of wild-type group in each time point. Scale bars (upper),

20mV (vertical) and 500ms (horizontal). Scale bars (lower), 20mV (vertical) and 125ms (horizontal).

(D) Excitability of VN neurons in STIM1^{PKO}. There was no alteration of excitability after learning (sham vs 1hr,

p=0.422; sham vs 4hr, p=0.801; sham vs 24hr, p=0.493, left; sham, n=17; 1hr, n=25; 4hr, n=15; 24hr, n=15),

and no significant changes in 150 pA injection as well (1hr, p=0.530.; 4hr, p=0.908; 24hr, p=0.371)

Two-way repeated measure ANOVA was used for injected current-frequency graphs. One-way ANOVA with

Fisher LSD test was used for bar graphs. Asterisks in each time points were calculated by comparing to sham

groups. Error bars denote SEM. *p<0.05 ***p<0.001.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary Information

Intrinsic plasticity of cerebellar Purkinje cells in motor learning circuits

Dong Cheol Jang, Hyun Geun Shim and Sang Jeong Kim

Figure S1 (Related to Figure 1). The basal ocular-motor performance of STIM1^{PKO} mice was not altered
compared to wild-type littermates.

(A) OKR responses in various frequency of drum rotating. OKR gain and phase values were not different between

wild-type littermates (n=20) and STIM1^{PKO} (n=17).

(B)　VOR responses under dark (dVOR) in various frequency of drum rotating. dVOR gain and phase values were

not different between wild-type littermates and STIM1^{PKO}.

(C)　VOR responses under light (lVOR) in various frequency of drum rotating. lVOR gain and phase values were

not different between wild-type littermates and STIM1^{PKO}.

Figure S2. Synaptic transmission in STIM1WT and STIM1PKO was not significantly different (Related to Figure

2).

(A) Cumulative plots of IEI of sEPSC in wild-type (blue) and STIM1PKO mice (red). The cumulative fraction of IEI

and bar graph (inset) of sEPSC frequency indicated that frequency of sEPSC was not changed in STIM1PKO

compared to wild-type littermates (wild-type, n=15; STIM1PKO, n=15, p=0.145).

(B) Cumulative plots of amplitude of sEPSC. The cumulative fraction of amplitude and bar graph (inset) of sEPSC

frequency indicated that amplitude of sEPSC was not changed in STIM1PKO compared to wild-type littermates

(p=0.587).

Unpaired *t*-test was used for bar graphs. Error bar denotes SEM.

Figure S3. Intrinsic plasticity was impaired in STIM1^{PKO} (Related to Figure 3).

(A) Plots of the normalized EPSC change after application of LTD induction protocol in wild-type littermates (n=8, n=4 after 40min, blue) and STIM1^{PKO} mice (n=7, n=6 after 40min, red). Synaptic plasticity was normally induced in STIM1^{PKO} mice. Scale bar, 200pA

(B) Bar graphs showed the comparison of excitability changes after LTD induction. In STIM1^{PKO} mice, down-

regulation of excitability in PCs was shown 20 min after induction (n=7, p=0.010), however, fully recovered 40

min and 50 min after induction (40 min, n=7, p=0.331; 50 min, n=6, p=0.748). On the contrary, the intrinsic

plasticity was induced and slightly further developed in time from wild-type littermates (20 min, n=8, p<0.001;

40 min, n=8, p<0.001; 50 min, n=4, p=0.001). Scale bars (upper), 20mV (vertical) and 200ms (horizontal)

(C-F) Comparing PC excitability of wild-type littermates (blue) and STIM1[PKO] mice (red) in each time points. (C)

STIM1[PKO] group (n=17) showed significantly lower excitability than wild-type littermates (n=20, p<0.001). (D) While

the excitability of STIM1[PKO] (n=13) was unchanged, excitability of wild-type littermates (n=11) was plunged

(p=0.067). (E) At 4 hours after learning, the altered excitability of wild-type littermates (n=16) partly restored, and

overlapped to excitability of STIM1[PKO] (n=17), which was still unchanged (p=0.644). (F) As the excitability of wild-

type littermates (n=20) was fully recovered, significant difference from STIM1[PKO] group (n=13) was also restored

(p=0.022).

One-way repeated measure ANOVA with post hoc Sidak correction was used for panel B. Two-way repeated measure

ANOVA was used for panel C to F. Error bars denote SEM. *p<0.05, **p<0.01, ***p<0.001.

Figure S4. Basal synaptic transmission and excitability in VN neurons before VOR gain up training (Related

to Figure 4 and 5).

(A) Frequency of synaptic transmission in VN neurons from wild-type littermates (n=23, blue) and STIM1^{PKO} mice

(n=14, red). The cumulative fraction of IEI and bar graph (inset) of sEPSC frequency indicated that frequency

of sEPSC was higher in STIM1^{PKO} compared to wild-type littermates (p=0.032).

(B) Amplitude of sEPSC in VN neurons from wild-type littermates (blue) and STIM1^{PKO} mice (red). The cumulative

fraction of amplitude and bar graph (inset) of sEPSC frequency indicated that amplitude of sEPSC was not

changed in STIM1^{PKO} compared to wild-type littermates (p=0.161).

(C-F) Excitability in VN neurons from STIM1^{WT} (blue) and STIM1^{PKO} (red) mice in each time points. Square-wised

somatic current steps were injected from membrane potential with various ranges from -150 pA to 150 pA with

increment of 50 pA for 1 s. (C) Overall, the gain responses of VN neurons from STIM1^{PKO} (n=16) were not

significantly different from wild-type littermates (n=23, p=0.423), but at 150 pA injection, STIM1^{PKO} showed higher

excitability than wild-type littermates (p=0.003). (D) At the short-term period, the excitability of VN neurons of the

wild-type (n=16) has slightly increased, while the excitability of STIM1^{PKO} (n=25) was unchanged. Statistical

difference has disappeared (p=0.428). (E) At 4 hours after learning, the excitability of VN neurons of the wild-type

(n=38) increased more, and that of STIM1^{PKO} (n=15) was unchanged again. IO curve of both groups are overlapped

(p=0.759). (F) As the excitability of wild-type littermates (n=19) became much higher at the long-term period, the

wild-type group showed significant higher frequency than STIM1^{PKO} group (n=15) (p=0.016). At 100pA and 150pA

injection, statistical significance was 0.005 (100pA) and <0.001 (150pA) by post-hoc Sidak test.

(G) Synaptic LTP of VN neuron. Both wild-type (n=6) and STIM1^{PKO} (n=7) groups showed intact synaptic plasticity

through *in vitro* induction protocol.

(H) Intrinsic plasticity of VN neuron was intact in both genotypes. Through LTP induction protocol, the excitability

of VN neurons was significantly increased in both genotypes (wild-type, n=6, p=0.031; STIM1^{PKO}, n=6, p=0.031).

There was no significant difference between post-induction groups of both genotypes (p=0.386)

Unpaired *t*-test was used for bar graphs in panel A and B. Two-way repeated measure ANOVA with Sidak test was

used for panel C-F. Mann-Whitney U test was used for panel H. Error bars denote SEM. *p<0.05, ***p<0.001.

Figure S5. Schematic illustration for memory trace of VOR motor memory.

(A) Summary of behavioral test and *in vitro*, *ex vivo* recording showing Temporal order of the memory retention and

neural plasticity in PCs and VN neurons. Differences between wild-type and STIM1[PKO] mice are indicated as red

shade of each plots). VOR memory retention level is maintained over a day whereas the motor memory is declined

at mid-term period (+1Hr - +4Hr) in STIM1[PKO]. Alterations of the neural activity corresponding to each period

are presented below. There is no difference in PF-PC synaptic plasticity between wild-type and STIM1[PKO] mice.

However, learning-induced LTD-IE is abolished within an hour in STIM1[PKO] and the difference in PC intrinsic

plasticity between groups may lead to MF-VN synaptic plasticity and intrinsic plasticity of VN neurons (see the

red shade). Furthermore, peak difference of each plot seems to move from 1 to 24 hours after learning, indicating

that the plasticity in cerebellar PCs and VN neurons is connected in order. Thus, we speculate the behavioral

difference between wild-type littermates and STIM1^{PKO} (red shade in top) underlies PC intrinsic plasticity, which

plays a role in memory transfer process, but not PF-PC synaptic plasticity.

(B) Schematic illustration of neural circuit for VOR memory storage shown in wild-type (top) and STIM1^{PKO} mice

(bottom). For successful memory acquisition and storage, four different types of neural plasticity are necessary

(synaptic and intrinsic plasticity in the PCs and VN neurons). Especially, intrinsic plasticity of PC has important

role in transfer acquired memory to sub-cortical area in this circuitry (top). When PC intrinsic plasticity is

abolished, synaptic and intrinsic plasticity in VN neurons are impaired thereby failure to long-term memory

storage (bottom)