Integrated multi-omics analysis to study the effects of simulated weightlessness on rhesus macaques (*Macaca mulatta)*

Peng Zhang[1,2, *], Libin Shao[3,8, *], Jie Zhang[3,*], Wenjiong Li[1,*], Guangyi Fan[3,7], Ying Zhou[6], Guanghan Kan[1], Hongju Liu[1], Weidong Li[6], Fei Wang[2], Xixia Chu[6], Peng Han[3], Ling Peng[3], Xingmin Liu[3],Jianwei Chen[3], Xinming Liang[4],Jingkai Ji[4],Shiyi Du[4], Zhanlong Mei [4],Ronghui Li[3], Xun Xu[3,4,5,†], Shanguang Chen[2,†], Xin Liu[3,4,5,†], Xiaoping Chen[1,2,†]

[1]State Key Laboratory of Space Medicine Fundamentals and Application, China Astronaut Research and Training Center, Beijing 100094, China.

[2]National Key Laboratory of Human Factors Engineering, China Astronaut Research and Training Center, Beijing 100094, China.

[3]BGI-Qingdao, BGI-Shenzhen, Qingdao, 266555, China

[4]BGI-Shenzhen, Shenzhen 518083, China

[5]China National GeneBank, BGI-Shenzhen, Shenzhen 518120, China

[6]Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders, Ministry of Education, Shanghai Jiao Tong University, Shanghai 200240, China

[7]State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Macao, China

[8]School of Biology and Biological Engineering, South China University of Technology, Guangzhou, 510006, China

*These authors contributed equally to this work.

[†]Correspondence authors: Xun Xu(xuxun@genomics.cn),Shanguang Chen (tigercsg@163.com) , Xin Liu (liuxin@genomics.cn) and Xiaoping Chen (xpchen2009@163.com).

Abstract

Safety and health of astronauts in space is one of the most important aspects of space exploration, however, the genomic research about how a weightless space can affect astronaut's health was limited. In this study, we sequenced 25 transcriptomic, 42 metabolomic and 35 metagenomic data of 15 rhesus macaques (*Macaca mulatta)* spanning seven simulated weightlessness experiment stages. We identified 84 genes, 1911 features and 55 genera which are significantly changed in blood and muscle, hippocampal region, dorsomedial prefrontal cortex as well as fecal,

respectively. Furthermore, performing the integrated analysis of three omics data, we found several pathways which were related to regulation of immune system process, glucose uptake, reaction to threatens, neurotoxic and bone or joints damage, such as tyrosine metabolism and tryptophan metabolism. Our results provided an initial attempt of "multi-omics" approaches which combined transcriptomics, metabolomics and metagenomics to illustrate some molecular clues for simulated weightlessness effect on the rhesus macaques and potential sight of microgravity's effect on astronauts' health.

Keywords

weightlessness, transcriptomics, metabolomics, metagenomics, immune

Introduction

Since the first traveling to space, the frequency of long-term spaceflight has increased rapidly. However, health issues are threatening mankind's space exploring and the normal life of retired astronauts. Because research under real condition of spaceflight is limited, researchers often use ground-based analogs to study spaceflight's effect on organisms. One such popular model is the head down-tilt bed rest (HDBR) which mimics the headward fluid shift and axial body unloading of spaceflight [1], and in which organisms remain either horizontal or -6 degree HDBR for days to months [1, 2]. Several previous studies suggested that long duration space flight can affect intracranial pressure [3], brain structure and function [4, 5], osteoclastogenesis [6], blood pressure [7], visual impairment [8] and the immune system [9, 10], indicating that solutions for maintain astronaut's health. Multiple studies on space traveling astronauts, animals and plants, as well as microgravity analogues on animal models have been taken to study the biology changes during and after the spaceflight [11, 14].

The characteristics of space environment have two main factors, space radiation and microgravity [15]. It has been reported that the space radiation can damage DNA, leading to potentially harmful health consequences [15]. Space flight, especially microgravity, is one of the most extreme conditions that humans encounter [16]. However, little is known about the changes of gene expression, gut microbiota and metabolites of astronaut under microgravity. Addition of omics profiling to microgravity space flight experiments will understand key areas of variance in the molecular landscape [17]. The genomic and transcriptome studies can find gene expressions under difference environmental [18]. Space flight affects the community behavior of bacteria that harmful and beneficial human microbial interactions change during space flight [19]. About microgravity of systemic immune dysfunction may make the host more susceptible to pathogen infection [16]. Besides, metabolomics, the comprehensive study of metabolic reactions, was applied to study the metabolite profile in the studies of spaceflight effects on astronauts [2, 7], which help to understand the mechanism of physiological changes during and after space life. Thus, the results of integrated omics analysis could be used for aerospace medicine and research [17].

In this study, we recruited 15 rhesus macaques in different stages of ground-based HDBR analog using transcriptomics, metabolomics and metagenomics to understand the effects of weightless space flight analogues (HDBR) in rhesus macaques. As far as we know, our study will be the first multi-omics sights into the molecular study of space flight analogues.

Results

Metabolism significance of rhesus macaques in HDBR study

To investigate the effect of spaceflight on metabolism, we established a head down-tilt bed rest (HDBR) model using rhesus macaques. We then used ultra-performance liquid chromatography / mass spectrometry (UPLC/MS)-based metabolomics approach to quantify the metabolism signatures of the muscle, hippocampal region (HIP), dorsomedial prefrontal cortex (dmPFC) of

15 rhesus macaques in three different time points: pre-HDBR (control, T1), HDBR (T4) and

recovery states (T7) (Figure 1, Table S1). We firstly conducted pair-wised comparison among

these three time points using univariate and multivariable analysis. 356, 287 and 100 significantly

different abundance features (DAFs) were identified in muscle by comparing HDBR to control,

recovery to HDBR, and recovery to control, respectively (P < 0.05, fold change > 1.2 or fold

change < 1/1.2, VIP>1, Figure S1). Besides, we also identified 328, 430 and 177 DAFs in HIP;

368, 472 and 109 DAFs in dmPFC. For the further analysis, we mainly focused on the 154, 198

and 207 DAFs that were significantly changed in HDBR comparing to control and recovery in

muscle, HIP and dmPFC, respectively (P<0.05, Figure 2a, Figures S2a and S2b). Based on

KEGG pathways analysis (Table S5), we found that the tryptophan metabolism pathway emerged

in all comparisons of three tissues and tyrosine metabolism pathway contained 8 metabolites

emerged in comparisons of muscle and dmPFC. Out of these eight metabolites, L-dopa

(3,4-dihydroxy-L-phenylalanine,   C00355,   HMDB00609),   L-Norepinephrine   (C00547,

HMDB00216), epinephrine (C00788, HMDB00068), L-Metanephrine (C05588, HMDB04063)

were reported to be involved in neurotransmitter precursor, blood pressure control, stress reaction,

glucose uptake and energy metabolism [20, 24].

In addition, to observe the interaction of these DAFs, we performed the weighted correlation

network analysis (WGCNA) for three tissues, respectively [34]. In details, we classified 552

DAFs into six modules in muscle, 668 DAFs into seven modules in HIP and 691 DAFs into six

modules in dmPFC, respectively (Figure 2b, Figures S3a and S3b). Particularly, we found the

DAFs of tyrosine metabolism pathway (mcc00350) mainly distributed in three modules (M1, M5

and M6).   Interestingly, the DAFs of M1 showed a good co-abundant pattern in muscle

that all of these features decreased in HDBR and increased in recovery (Figure S3c and Figure

2c). Among them, the abundance of L-Alanine (C00041, HMDB00161), L-Carnitine (C00318,

HMDB00062) and calcitroic acid (HMDB06472) were decreased, which might be related to the

muscle weakness and bone loss (Figures S3d-S3f). L-Alanine (C00041, HMDB00161) is an

important amino acid in glucose-alanine cycle, which moves gluconeogenesis from muscle to

liver to produce ATP stored in muscle for muscle contraction [26]. L-Carnitine ((C00041,

HMDB00161) is most abundant in skeletal muscle and cardiac muscle, and it actives and

transports fatty acids into the mitochondria for energy generation in skeletal muscle [27].

Calcitroic acid (HMDB06472) is a major metabolite from 1,25-Dihydroxyvitamin D3 which

plays an important role on bone homeostasis and resorption, was found decreasing after long-term

bed rest or space flight [38] [28]. Besides, we found that epinephrine (C00788, HMDB00068)

was increased the metabolic rate in skeletal muscle (C00788, HMDB00068) [30], and it was the

hub metabolite of M1 (Figure 2d). Pyridoxamine (C00534, HMDB01431) was the hub metabolite

of M6, which was one form of vitamin B_6 taking part in metabolism of L-Alanine.

Gut microbiome changes of rhesus macaque in HDBR study

The gastrointestinal tract harbors most of microbes, which play important roles in organisms'

health, disease, immunity, and even behavior [31, 32]. To investigate the effect of HDBR on the

microbiome, we sequenced 35 metagenomics gut samples from five rhesus macaque individuals

of seven time points (T1~T7) of HDBR analog (Figure 1). We generated a total of 286.86 Gb raw

data and 275.99 Gb high-quality data after removing low quality data and the host, the average

clean data was 7.89 Gb for each sample (Table S2). Then, we constructed a reference gene

catalogue of rhesus macaque's gut metagenome using all samples (Table S3). Among of these

genes, 63.10% of gene catalogue could be annotated to phylum level, 21.00% to genus level, and

2.43% to species level. And 45.51% could be annotated to 6,631 KEGG Orthologs (KOs).

From the annotation of microbiome species, we found that Firmicutes and Bacteroidetes were

dominated on the phylum level, as well as Prevotella and Clostridium were the dominated genera

on the genus level (Figure S4). From the annotation of KEGG function, K03088, K03327,

K06147, K00599 and K00754 were most dominated KOs with a higher abundance which

annotated to transcription machinery, ion-coupled transporters, transporters, histidine

metabolism/tyrosine metabolism and fructose and mannose metabolism. In all, 49.48% of genera

and 95.58% of KO functions were shared by monkey, human gut gene catalogues (Figure S5).

We identified 55 genera with a significant change of abundance from T1 to T7 (Table S6).

Interestingly, out of these 55 genera, 43% of them had decreased in their abundance in the

experimental stages (T2~T4) and increase during the recovery phases (T5-T7). These genera

include such as *Acinetobacter* and *Lactococcus,* which is involved regulation of inflammation [33,

34, 35] and protection of infections [36], indicating that HDBR caused a disorder in the rhesus

macaque's intestinal flora (Figure 3a and 3b). The abundance of two genera had been decrease

continuously throughout the study, such as *Bifidobacterium,* which could modulate host immune

responses, inhibit infection by pathogens, and regulate intestinal microbial homeostasis [37, 38,

39] (Figure 3c). In addition, we found 537 KOs with significantly changed abundance were

assigned into 139 genera, of which 27 genera were significantly changed during T1 to T7

(P<0.05). Correlation analysis showed that 44.76% of pair-wise correlation between KOs and

genera were positive, whereas only 3.38% correlation were negative (R2>0.3, P<0.05) (Figure

3d). This results indicate that the changes of genera abundance positively affect the gut microbial

function, such as *Myroides* and *Acinetobacter*, which could help to improve the K00121, K00151,

K00276, K00451 and K01555 which participate in tyrosine metabolism pathway. In summary,

our findings revealed that HDBR affects the abundance of gut microbiome, which might be

related to the host immune response and metabolism.

Transcription features of rhesus macaque in HDBR study

To investigate the effect of HDBR on gene expression in immune cells, we collected 25 blood

samples from five rhesus macaque individuals of six time points (T1, T3~T7) (Figure 1). We

generated 779.09 M reads in six time points of HDBR analog (Table S4). Using pairwise

comparison of each time point, we detected 84 significantly differential expressed genes (DEGs,

fold change>2 and P<0.05) in at least two time points. We firstly focused on 65 DEGs in HDBR

(T3 and T4) comparing to the control (T1) and the recovery phases (T5~T7, Figure 4a). These

DEGs were significantly enriched in 44 biological processes (P<0.05), and 41 of them were

related to the regulation of immune system, including regulation of leukocyte activation

(GO:0002694), regulation of T cell differentiation (GO:0045580) and regulation of interleukin-2

production (GO:0032663) (Figure 4b and Table S7). DEGs in these immune response biological

processes were mainly down-regulated in the HDBR, which were consistent with the previous

researches about the immune system of animal and human in simulated long-term microgravity

environment [36][40]. In addition, we also found 71 DEGs in recovery status (T5~T7) comparing

to the control (T1, Figure 4a), which were also mainly enriched in the immune system. Besides,

WGCNA [25] method was employed to construct genes co-expression networks in

transcriptomes, 1941 genes with max median absolute deviation (MAD) were clustered into

eleven modules. In accordance with the differential expression analysis, genes in the purple

module were mainly related to immune response (Table S8) with the expression dropped in

HDBR.

Integrated analysis of omics data in rhesus macaque

Short-chain fatty acids (SCFA) is a dominant metabolite produced from bacteria, related to the

immune response, such as butyrate regulates the size and function of the regulatory T cell

network by promoting the induction and fitness of regulatory T cells in the colonic environment

[41, 42, 43]. In our study, DEGs in lymphatic cells were related to reduce immune response and

dysregulation of muscle butyrate metabolism. In the gut microbiome, the relative abundance of

butyrate-producing colon bacteria *Eubacterium*, *Roseburia* and their cross-feeding bacteria

Bifidobacteria were reduced in the HDBR. Thus, our results revealed that the weak immune

response and the reduced abundance of butyrate during HDBR might be related to the abundance

change of gut microbe (Figure 5).

Besides, previous study showed that 3-hydroxyphenylacetate is a by-product of the tyrosine

metabolism mediated by *Clostridium spp* [28]. Enzyme 4-hydroxyphenylacetate

3-monooxygenase (EC: 1.14.14.9), which was not found in rhesus macaque, could transform

3-hydroxyphenylacetate (C00642, HMDB00020) to 3,4-Dihydroxybenzeneacetic acid

(3,4DPHAA, C01161, HMDB01336) in tyrosine metabolism. Coincidently, enzyme

4-hydroxyphenylacetate 3-monooxygenase and 3,4DPHAA both increased in during HDBR and

decreased during the recovery phase, consistent with the abundance changes of a gut microbe

Providencia rettgeri which produces enzyme 4-hydroxyphenylacetate 3-monooxygenase (EC:

1.14.14.9), indicating *Providencia rettgeri* influence tyrosine metabolism by producing

3,4DPHAA (Figure 5). Besides, we also found enzyme tryptophan 2,3-dioxygenase (EC:

1.13.11.11) could produce N'-Formylkynurenine (C02700, HMDB60485), which then

transformed to Formylanthranilic acid (C05653, HMDB04089) by enzyme kynureninase (EC:

3.7.1.3) in tryptophan metabolism. Enzyme tryptophan 2,3-dioxygenase and kynureninase could

be produced by the gut microbes with abundance change during HDBR, such as *Myroides* and

Comamonas. Furthermore, we found *Myroide* were positively and significantly correlated with

Leucodopachrome (C05604, HMDB04067) and Dopaquinone (C00822, HMDB01229) in the

tyrosine metabolism pathway (mcc00350) and *Providencia* were found positively and

significantly correlated with p-Cresol (C01468, HMDB01858) which is a metabolite of tyrosine

(P<0.05). Meanwhile, *Lactococcus* were positively correlated with L-Carnitine (C00318,

HMDB00062) which is crucial in providing energy to muscles [27] (Figure S6). Therefore, the

significantly changed abundance of gut microbes were involved in tyrosine metabolism

(mcc00350) and tryptophan metabolism (mcc00380) pathways during the HDBR analog

(P<0.05).

Discussion

Living in a space environment with microgravity and motionless for long periods of time may

have adverse effects on immunity, metabolism and health. The first report from the Soviet

immunologist, Konstantinova and coworkers [44], found that lymphocyte responsiveness to

mitogens was remarkably reduced after astronauts after a longtime spaceflight. In this study, we

applied transcriptomic, metabolomic and metagenomic analysis in a head down-tilt bed rest

model to elucidate the effect of spaceflight and analogue microgravity environment on biology

functions. In transcriptome data, DEGs were enriched in multiple biological processes mainly

related to immune response, indicating that gene expression has plausible functional connections

with spaceflight and microgravity. For the first time, we also found the effects of an analogue

microgravity environment on immunity by combining metabolomics and metagenomics. We

found that tyrosine metabolism, which plays an important role in muscle function, was affected

by HDBR. The abundance of L-Dopa, norepinephrine, epinephrine and L-Metanephrine in

catecholamines biosynthesis and metabolism pathway, dopaquinone and leucodopachrome in

L-Dopachrome biosynthesis pathway and 4-Hydroxyphenylpyruvate and homogentisate in

homogentisate biosynthesis pathway were significantly affected by HDBR (Figure 2e). We also

found that calcitroic acid (related to bone balance), L-Alanine and L-Carnitine (related to muscle

contraction and muscle weakness) were affected by HDBR.

To our knowledge, this was the first study successfully combining metabolomics and

metagenomics in studying simulated microgravity on health and metabolism. The findings

provided valuable insights into how long-term spaceflight could affect metabolism and health.

More importantly, our findings on the link between gut microbiome with metabolism and

immunity provided the possible solutions to combat dysregulated immunity and metabolism in

spaceflight. This study was conducted with animal model in a simulated environment, we

supposed a solid combination of omics in the study of astronauts could be a useful tool for future

study of astronauts in extremely environment. Further study with more astronauts would provide

more information during long term space flight.

Materials and Method

Animal experiments

We sampled 20 healthy male rhesus macaques, aged 4 to 6 years and weighing 4 to 8 kg from

Beijing Institute of Xie'erxin Biology Resource (Beijing, China). All of these rhesus macaques

received 3 months of domestication (involving preliminary caretaker handling, confinement

jacket fitting, and tilt-table acclimation training) at the Laboratory Animal Center of China

Astronaut Research and Training Center prior to the start of the experiments. Only 15

well-domesticated rhesus macaques were selected and separated into 3 groups: 1) CON:

ground-based controls, 2) HDBR: 42 days of HDBR, 3) REC: 42 days of HDBR plus 32 days of

recovery.

A six-week head-down tilted bed rest (HDBR) experiment was performed on rhesus macaques

to simulate weightlessness as described previously [45]. Briefly, rhesus macaques laid on beds,

which were tilted backward 10 °C from the horizontal. The head-down monkeys wore the

confinement jacket, which enabled them to be fixed to the bed. These rhesus macaques were

housed one per bed in rooms with air temperature maintained at $23 \pm 2°C$ and a standard 12:12 h

dark–light cycle (lights were turned on at 8:00 a.m. and off at 8:00 p.m.). After six weeks of

HDBR, each rhesus macaque was solely removed into stainless steel mesh cages to recovery for

32 days. Throughout the duration of the experiment, rhesus macaques received an intensive

humanistic care. For instance, the rhesus macaques always had free access to food and water.

Toys (such as the drum-shaped rattle, a Chinese traditional toy) were available all the time except

during experimental procedures. The caretaker accompanied the rhesus macaques during the day

time to help relieve anxiety. The general health condition of the rhesus macaques was also

carefully monitored. All procedures were performed in accordance with the principles of the

Association for Assessment and Accreditation of Laboratory Animal Care International

(AAALAC), approved by Institutional Animal Care and Use Committee of China Astronaut

Research and Training Center (ACC-IACUC-2014-001).

Sample collection

All samples provided consent forms which were approved by BGI genomics Committee of Ethics.

Under light ketamine sedation, sterile heparinized peripheral blood samples were obtained from

femoral vein of the five rhesus macaques in REC group before (Pre-3, T1), during (HB-12,

HB-25, HB-40, T2~T4) and after (R-12, R-24, R-32, T5~T7) the HDBR at 10:00 a.m. Peripheral

blood mononuclear cells (PBMCs) were then collected by Ficoll-Hypaque density-gradient

centrifugation. Fecal samples were also collected from the five rhesus macaques in REC group

before (Pre-2, T1), during (HB-16, HB-30, HB-42, T2~T4) and after (R-13, R-17, R-28, T5~T7)

the HDBR. Skeletal muscle sample, dmPFC sample and HIP sample were collected from all of

the selected rhesus macaques in each group.

Transcriptome analysis

mRNA of the PBMCs was isolated by oligo(dT) and sequenced by Complete Genomics(CG)

SE50, we then use SOAPnuke to filter out the reads of low quality, at least 20 million reads were

generated for each sample. HISAT2 was applied to map sequence reads to genome and RSEM to

calculate the FPKM. Spearman correlation between all samples were calculated using FPKM, the

correlation between all samples was very high (>0.95) and the biological replicates did not cluster

well together. We used DESeq2 package in R to find different expressed genes [46], and GO was

used to annotated function information.

Metabolomics analysis

UPLC-MS technology was implemented for metabolites detection. The experimental quality was

evaluated by quality control (QC) samples. Features alignment, picking and identification were

performed by Progenesis QI (Waters, Nonlinear Dynamics, Newcastle, UK). MetaX software was

used for data cleaning and statistical analysis [47]. Low quality features were removed within

data cleaning. By combining the univariate and multivariate statistical analysis, significantly

changed features (P value<0.05, fold change <1/1.2 or fold change >1.2, VIP >1) were acquired.

Those features were further annotated by Progenesis QI with Human Metabolome databases

(version   3.6),   and   online   Kyoto   Encyclopedia   of   Genes   and   Genomes   database

(www.genome.jp/kegg/).

Clusters of co-abundant metabolites of muscle, HIP and dmPFC were performed by R package

WGCNA [25], respectively. Soft threshold β=9 for muscle features, soft threshold β=13 for

dmPFC features, and soft threshold β=12 for HIP features were chosen by scale free topology

analysis, for signed, weighted features co-abundance correlation network construction. Dynamic

hybrid tree-cutting algorithm by deepSplit of 4 and a minimum cluster size of 30 were applied for

clusters identification. If the biweight mid-correlation between the cluster's eigenvectors

exceeded 0.8, the similar clusters would be subsequently merged. The muscle, dmPFC and HIP

features clusters, were labelled by M1-M6, D1-D7 and H1-H6, respectively.

Metagenomics analysis

Raw reads sequenced on the Illumina Hiseq 2000 platform (Expression Analysis Inc., San Diego,

CA, USA) at BGI were filtered to remove the adaptor contamination, low-quality reads and host

genomic DNA (Rhesus macaques, assembly Mmul_8.0.1, NCBI). The remaining high-quality

reads were assembled by metaSPAdes (v3.10.1). Open Reading Frames (ORFs) in contigs of each

sample were obtained using GeneMark (v2.7). The non-redundant gene set of all ORFs was

clustered using CD-HIT (v4.5.7) based on nucleotide sequence and the identity is 95% at the

coverage 90%. Taxonomic annotation of gene set was made with CARMA3 on the basis of

BLASTP alignment with bacteria and archaea from NCBI-NR database. And the gene set was

also annotated against KEGG (v59) databases with BLAST (v2.2.23).

Gene abundance profiling was calculated based on the alignment of SOAP2 and the species

abundant profile and functional profile were summarized from their respective genes [48]. The

differential alpha diversity, species and KOs of different time points were calculated using R

kruskal.test. Spearman coefficient was used to calculate the relationship between different genus

and different Kos, between different genus and different metabolic features according to their

abundance.

Acknowledgments

This work was supported by grants from the National Basic Research Program of China (2011CB711003), the National Natural Science Foundation of China (81772016, 81272177) and the 1226 major project (AWS16J018) to X.C.

Competing interests

Competing interest statement: The author denies that he has any intention to obtain any financial interests.

Figure 1. Overview of the study integrating metabolome, metagenome and transcriptome data and rhesus macaques HDBR experiments. Experimental Design for 15 rhesus macaques HDBR experiments. Each group of 5 rhesus macaques participated in this experiment. 5 rhesus macaques before HDBR were chosen for muscle, HIP and dmPFC tissues obtaining. 5 of the others took part in following HDBR experiment, then muscle, HIP and dmPFC tissues of which at HDBR 42 days were obtained. The 5 remaining rhesus macaques got into recovery experiments, and muscle, HIP and dmPFC tissues were obtained at 32 days after recovery experiment. Among all the experiments, blood samples 3 days before HDBR, 12 days, 25, and 40 days during HDBR, 12 days, 24, 32 days during recovery, and fecal samples 2 days before HDBR, 16 days, 30, and 42 days during HDBR, and 13 days, 17 and 28 days during recovery were all collected from the last 5 rhesus macaques.

Figure 2. Meaningful metabolites and tyrosine metabolism pathway were found in muscle by metabolomics analysis.(a)A heatmap of 154 DAFs which only changing significantly in HDBR comparing to control and recovery in muscle.(b)Dendrogram of 552 muscle DAFs mainly clustering into 6 modules by WGCNA analysis. 6 modules colored with 'turquoise', 'yellow', 'red', 'green', 'blue' and 'brown' were labeled by 'M1' - 'M6', respectively. And 'grey' module included the remaining DAFs that did not fit clustering criteria.(c)A heatmap of DAFs in M1

showing that all DAFs down-regulated during HDBR and up-regulated during recovery in

muscle.(d)Epinephrine was the hub metabolite of M1 network in muscle. Correlation of M1

DAFs were calculated by WGCNA. And the bigger size of the node means the more neighbor of

this node. The adjacent lines of epinephrine colored with 'red', and the others colored with 'light

gray'.(e)11 significantly changed muscle metabolites showing in tyrosine metabolism pathway. 5

metabolites in M1 decreasing during HDBR and increasing during recovery were colored with

'turquoise', 3 metabolites in M2 showing opposite changes to M1 were colored with blue. And 5

metabolites in M3 showing similar trend with M1 were colored with brown. The 5 remaining

metabolites colored with grey weren't found in differential metabolites.

Figure 3. Several important intestinal genera alterations in seven time points with 95%

confidence interval for the medians, and associations of differential gut microbial genera with

differential KOs.(a-c) Box plot of the differential genera abundance of Acinetobacter, Lactococcus

and Bifidobacterium. The x coordinate represents T1 ～ T7 time points, y coordinate represents   the

genera abundance (log10). For each interquartile ranges (IQRs), the first and third quartiles were

showed as the boxes, and the line inside the box represents the median. The circles data points

represent the outside of the whiskers which marked as the lowest or highest values within 1.5 times

IQR boxes.(d) Spearman correlation coefficient heatmap (P<0.05) between 27 genera differed in

abundance (x coordinate) and 442 Kos differed in abundance (y coordinate). The red color represented

significantly positive correlation, the blue color represented significantly negative correlation, and

blank areas indicate no significant correlation.

Figure 4. Interesting DEGs and biological processes found in transcriptomics

analysis.(a)Histograms of DEGs between any two time points. Up-regulated and down-regulated

DEGs were colored with 'orange' and 'blue', respectively. (b)GO analyses of the 65 DEGs showing a

significant enrichment of several biological processes.

Figure 5. Reducing gut microbes produced lesser butyrate to weaken the host immune response, and the gut microbes might take part in tyrosine metabolism (mcc00350) and tryptophan metabolism pathway (mcc00380).

References

[1] Hargens, A. R., & Vico, L. (2016). Long-duration bed rest as an analog to microgravity. Journal of applied physiology, 120(8), 891-903.

[2] Smith, S. M., Heer, M., Wang, Z., Huntoon, C. L., & Zwart, S. R. (2012). Long-duration space flight and bed rest effects on testosterone and other steroids. The Journal of Clinical Endocrinology & Metabolism, 97(1), 270-278.

[3] Marshall-Bowman, K., Barratt, M. R., & Gibson, C. R. (2013). Ophthalmic changes and increased intracranial pressure associated with long duration spaceflight: an emerging understanding. Acta Astronautica, 87, 77-87.

[4] Pompeiano, O., d'Ascanio, P., Balaban, E., Centini, C., & Pompeiano, M. (2004). Gene expression in autonomic areas of the medulla and the central nucleus of the amygdala in rats during and after space flight. Neuroscience, 124(1), 53-69.

[5] Demertzi, A., Van Ombergen, A., Tomilovskaya, E., Jeurissen, B., Pechenkova, E., Di Perri, C., ... & Sijbers, J. (2016). Cortical reorganization in an astronaut's brain after long-duration spaceflight. Brain Structure and function, 221(5), 2873-2876.

[6] Rucci, N., Rufo, A., Alamanou, M., & Teti, A. (2007). Modeled microgravity stimulates osteoclastogenesis and bone resorption by increasing osteoblast RANKL/OPG ratio. Journal of cellular biochemistry, 100(2), 464-473.

[7] Fritsch-Yelle, J. M., Charles, J. B., Jones, M. M., Beightol, L. A., & Eckberg, D. L. (1994). Spaceflight alters autonomic regulation of arterial pressure in humans. Journal of Applied Physiology, 77(4), 1776-1783.

[8] Mader, T. H., Gibson, C. R., Pass, A. F., Kramer, L. A., Lee, A. G., Fogarty, J., ... & Phillips, J. L. (2011). Optic disc edema, globe flattening, choroidal folds, and hyperopic shifts observed in astronauts after long-duration space flight. Ophthalmology, 118(10), 2058-2069.

[9] Sonnenfeld, G., & Shearer, W. T. (2002). Immune function during space flight. Nutrition, 18(10), 899-903.

[10] Taylor, G. R., & Janney, R. P. (1992). In vivo testing confirms a blunting of the human cell□mediated immune mechanism during space flight. Journal of Leukocyte Biology, 51(2), 129-132.

[11] Smith, S. M., Abrams, S. A., Davis-Street, J. E., Heer, M., O'Brien, K. O., Wastney, M. E., & Zwart, S. R. (2014). Fifty years of human space travel: implications for bone and calcium research. Annual review of nutrition, 34, 377-400.

[12] Taylor, K., Kleinhesselink, K., George, M. D., Morgan, R., Smallwood, T., Hammonds, A. S., ... & Hoshizaki, D. K. (2014). Toll mediated infection response is altered by gravity and spaceflight in Drosophila. PloS one, 9(1), e86485.

[13] Zhang, Y., Wang, L., Xie, J., & Zheng, H. (2015). Differential protein expression profiling of Arabidopsis thaliana callus under microgravity on board the Chinese SZ-8 spacecraft. Planta, 241(2), 475-488.

[14] van Oosterhout, W. P. J., Terwindt, G. M., Vein, A. A., & Ferrari, M. D. (2015). Space headache on Earth: Head-down-tilted bed rest studies simulating outer-space microgravity. Cephalalgia, 35(4), 335-343.

[15] Moreno-Villanueva M1,2, Wong M1,3, Lu T1,4, Zhang Y5, Wu H1.Interplay of space radiation and microgravity in DNA damage and DNA damage response.NPJ Microgravity. 2017 May 10;3:14.

[16] Li P1, Shi J1, Zhang P1, Wang K1, Li J1, Liu H1, Zhou Y1, Xu X1, Hao J1, Sun X1, Pang X1, Li Y1, Wu H1, Chen X2, Ge Q2.Simulated microgravity disrupts intestinal homeostasis and increases colitis susceptibility.FASEB J. 2015 Aug;29(8):3263-73

[17] Schmidt MA1, Goodwin TJ2.Personalized medicine in human space flight: using Omics based analyses to develop individualized countermeasures that enhance astronaut safety and performance.Metabolomics. 2013;9(6):1134-1156

[18] Srinivasan Shanmugarajan,1,2 Ye Zhang,3 Maria Moreno-Villanueva,1,4 Ryan Clanton,5 Larry H. Rohde,2 Govindarajan T. Ramesh,6 Jean D. Sibonga,1 and Honglu Wu1,Combined Effects of Simulated Microgravity and Radiation Exposure on Osteoclast Cell Fusion.Int J Mol Sci. 2017 Nov; 18(11): 2443

[19] Ronald Anderson,1,* Gregory Tintinger,1,2 Riana Cockeran,1 Moliehi Potjo,1 and Charles Feldman3.Beneficial and Harmful Interactions of Antibiotics with Microbial Pathogens and the Host Innate Immune System.Pharmaceuticals (Basel). 2010 May; 3(5): 1694–1710.

[20] Lopez, V. M., Decatur, C. L., Stamer, W. D., Lynch, R. M., & McKay, B. S. (2008). L-DOPA is an

endogenous ligand for OA1. PLoS biology, 6(9), e236.

[21] Hiroshima, Y., Miyamoto, H., Nakamura, F., Masukawa, D., Yamamoto, T., Muraoka, H., ... &

Endo, I. (2014). The protein Ocular albinism 1 is the orphan GPCR GPR 143 and mediates

depressor and bradycardic responses to DOPA in the nucleus tractus solitarii. British journal of

pharmacology, 171(2), 403-414.

[22] Chrousos, G. P. (2009). Stress and disorders of the stress system. Nature reviews endocrinology,

5(7), 374.

[23] Ratheiser, K. M., Brillon, D. J., Campbell, R. G., & Matthews, D. E. (1998). Epinephrine produces a

prolonged elevation in metabolic rate in humans. The American journal of clinical nutrition, 68(5),

1046-1052.

[24] Eisenhofer, G., Kopin, I. J., & Goldstein, D. S. (2004). Catecholamine metabolism: a contemporary

view with implications for physiology and medicine. Pharmacological reviews, 56(3), 331-349.

[25] Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network

analysis. BMC bioinformatics, 9(1), 559.

[26] Nelson, David L.; Cox, Michael M. (2005). Principles of Biochemistry (4th ed.). New York: W. H.

Freeman, pp. 684–85, ISBN 0-7167-4339-6.

[27] Alesci, S., Manoli, I., Costello, R., Coates, P., Gold, P. W., Chrousos, G. P., & Blackman, M. R.

(2004). Carnitine: Lessons from one hundred years of research. Annals of the New York Academy

of Sciences, 1033.

[28] Yu, O. B., & Arnold, L. A. (2016). Calcitroic Acid–A Review. ACS chemical biology, 11(10),

2665-2672.

[29] Smith, S. M., McCoy, T., Gazda, D., Morgan, J. L., Heer, M., & Zwart, S. R. (2012). Space flight

calcium: implications for astronaut health, spacecraft operations, and Earth. Nutrients, 4(12),

2047-2068.

[30] Ratheiser, K. M., Brillon, D. J., Campbell, R. G., & Matthews, D. E. (1998). Epinephrine produces a

prolonged elevation in metabolic rate in humans. The American journal of clinical nutrition, 68(5),

1046-1052. .

[31] Schmidt, T. S., Raes, J., & Bork, P. (2018). The human gut microbiome: from association to

modulation. Cell, 172(6), 1198-1215.

[32] Johnson, K. V. A., & Foster, K. R. (2018). Why does the microbiome affect behaviour?. Nature

Reviews Microbiology, 1.

[33] Cekanaviciute, E., Yoo, B. B., Runia, T. F., Debelius, J. W., Singh, S., Nelson, C. A., ... &

Crabtree-Hartman, E. (2017). Gut bacteria from multiple sclerosis patients modulate human T cells

and exacerbate symptoms in mouse models. Proceedings of the National Academy of Sciences, 114(40), 10713-10718.

[34] Visca, P., Seifert, H., & Towner, K. J. (2011). Acinetobacter infection–an emerging threat to human health. IUBMB life, 63(12), 1048-1054.

[35] Hughes, L. E., Bonell, S., Natt, R. S., Wilson, C., Tiwana, H., Ebringer, A., ... & Vowles, J. (2001). Antibody Responses to Acinetobacterspp. and Pseudomonas aeruginosa in Multiple Sclerosis: Prospects for Diagnosis Using the Myelin-Acinetobacter-Neurofilament Antibody Index. Clinical and diagnostic laboratory immunology, 8(6), 1181-1188.

[36] Nakamura, S., Morimoto, Y. V., & Kudo, S. (2015). A lactose fermentation product produced by Lactococcus lactis subsp. lactis, acetate, inhibits the motility of flagellated pathogenic bacteria. Microbiology, 161(4), 701-707.

[37] Mayo, B., & Van Sinderen, D. (Eds.). (2010). Bifidobacteria: genomics and molecular aspects. Horizon Scientific Press.

[38] Pinzone, M. R., Celesia, B. M., Di Rosa, M., Cacopardo, B., & Nunnari, G. (2012). Microbial translocation in chronic liver diseases. International journal of microbiology, 2012.

[39] Lievin, V., Peiffer, I., Hudault, S., Rochat, F., Brassart, D., Neeser, J. R., & Servin, A. L. (2000). Bifidobacterium strains from resident infant human gastrointestinal microflora exert antimicrobial activity. Gut, 47(5), 646-652.

[40] Luo, H., Wang, C., Feng, M., & Zhao, Y. (2014). Microgravity inhibits resting T cell immunity in an exposure time-dependent manner. International journal of medical sciences, 11(1), 87.

[41] Smith, P.M., Howitt, M.R., Panikov, N., Michaud, M., Gallini, C.A., Bohlooly-Y, M., Glickman, J.N., and Garrett, W.S. (2013). The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. Science 341, 569–573.

[42] Furusawa, Y., Obata, Y., Fukuda, S., Endo, T.A., Nakato, G., Takahashi, D., Nakanishi, Y., Uetake, C., Kato, K., Kato, T., et al. (2013). Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. Nature 504, 446–450.

[43] Arpaia, N., Campbell, C., Fan, X., Dikiy, S., van der Veeken, J., deRoos, P., Liu, H., Cross, J.R., Pfeffer, K., Coffer, P.J., and Rudensky, A.Y. (2013). Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. Nature 504, 451–455.

[44] Gmünder, F. K., & Cogoli, A. (1996). Effect of space flight on lymphocyte function and immunity. Handbook of Physiology, 2, 799-813.

[45] Koenig SC1, Ewert DL, Ludwig DA, Fanton JF, Convertino VA.Bed rest affects ventricular and arterial elastances in monkeys: implications for humans.Aviat Space Environ Med. 2004 Jan;75(1):7-15.

[46] Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology, 15(12), 550.

503   [47] Wen, B., Mei, Z., Zeng, C., & Liu, S. (2017). metaX: a flexible and comprehensive software for
504   processing metabolomics data. BMC bioinformatics, 18(1), 183.
505   [48] Xie, H., Guo, R., Zhong, H., Feng, Q., Lan, Z., Qin, B., ... & Chen, B. (2016). Shotgun
506   metagenomics of 250 adult twins reveals genetic and environmental impacts on the gut
507   microbiome. Cell systems, 3(6), 572-584.

a

b

c

d

e

Epinephrine

Tyrosine

Dopaquinone → Leucodopachrome → Dopachrome

3-Methoxytyramine

L-DOPA → Dopamine → Norepinephrine → Epinephrine → L-Metanephrine

4-Hydroxy-phenylpyruvic acid

Tyramine → N-Methyltyramine → Hordenine

4-Hydroxy-phenylacetaldehyde → Tyrosol → Salidroside

Identified in M1
Identified in M5
Identified in M6
Not identified in differential metabolites

a

Acinetobacter

b

Lactococcus

c

Bifidobacterium

d

a

b