Bacterioplankton dynamics driven by inter-annual variation in phytoplankton spring bloom

communities in the Baltic Sea

María Teresa Camarena-Gómez*[1,2], Clara Ruiz-González[3], Jonna Piiparinen[2], Tobias

Lipsewers[2], Cristina Sobrino[4], Ramiro Logares[3], Kristian Spilling[2]

[1]Tvärminne Zoological Station, University of Helsinki, J.A. Palménin tie 260, 10900 Hanko,

Finland

[2]Finnish Environment Institute, Marine Research Center, Agnes Sjöbergin katu 2, 00790,

Helsinki, Finland

[3]Department of Marine Biology and Oceanography, Institut de Ciències del Mar (ICM-CSIC).

Pg Marítim de la Barceloneta 37–49, E08003 Barcelona, Catalunya, Spain

[4] Faculty of Science, University of Vigo, Campus Lagoas-Marcosence s/n, 36310 Vigo, Spain

Running title: Bacterioplankton dynamics and phytoplankton composition in the Baltic Sea

* Corresponding author:

Phone: +358 401823179

Email: maria.camarena@environment.fi

Abstract

In the Baltic Sea, climate change has caused shifts in the phytoplankton spring bloom

communities with co-occurrence of diatoms and dinoflagellates. Such changes likely affect the

composition and function of associated bacterioplankton, key members of the carbon cycling,

although the actual effects are unknown. To understand how changes in phytoplankton impact

on bacterioplankton composition and function, we analysed bacterioplankton communities and

their production during different phases of the spring bloom in four consecutive years across

the Baltic Sea, and related them to environmental variables. Phytoplankton communities varied

largely in composition, modifying the taxonomic structure and richness of the associated

bacterioplankton assemblages. In presence of certain diatoms (*Achnanthes taeniata,*

Skeletonema costatum and *Chaetoceros* spp.), bacterial production and diversity were high and

with more relative abundance of Flavobacteriia, Gammaproteobacteria and Betaproteobacteria.

This bacterial community structure correlated positively with high diatom biomass and with

high bacterial production rates. In contrast, during dinoflagellate-dominated blooms or when

the diatom *Thalassiosira baltica* was abundant, both bacterial production rates and diversity

were low, with bacterial communities dominated by SAR11 and Rhodobacteraceae. Our results

demonstrate that, changes in the phytoplankton spring bloom will have profound consequences

on bacterial community structure and their role in carbon cycling.

Introduction

The Baltic Sea is suffering from a progressive increase in surface water temperature, which has been suggested to alter the food web structure, favoring small flagellates and dinoflagellates [1]. During the last decades, shifting phytoplankton spring bloom communities from diatom-dominated blooms towards higher abundances of dinoflagellates have been reported in some subbasins of the Baltic Sea [2-4], likely due to the increase in frequency of milder winters [1,5]. However, the consequences of these changes in spring blooms on the structure and functioning of the associated bacterioplankton communities are still poorly understood.

The nature and the timing of the spring bloom in the Baltic Sea vary between sub-basins [6]. The bloom typically starts with the light promoting net primary production in the southernmost Baltic Sea in February/March, reaching the Gulf of Finland in April, and the Gulf of Bothnia in May (Map in Fig. 1). The bloom reaches the peak at the time when inorganic nutrients have been depleted; N-limitation prevails in most of the Baltic Sea except for the P-limited Bay of Bothnia [7-9]. The subsequent decline phase is characterized by rapid sinking of the phytoplankton cells [10].

Diatoms and dinoflagellates have different ecological features, as diatoms grow faster than co-occurring dinoflagellates [11], they also differ in terms of quality and quantity of dissolved organic matter (DOM) release [12-14], and have different sedimentation patterns [10,15]. Since bacteria are the main consumers of the DOM produced by phytoplankton, also in the Baltic Sea [1,16], understanding how the predicted changes in the spring bloom may impact bacterial community composition and activity is essential to anticipate the biogeochemical consequences of this global warming driven process.

Assessing these issues in the Baltic Sea is complex due to the large environmental and spatial-temporal heterogeneity between subbasins, i.e. the pronounced salinity (2 - 20) and temperature gradients (0 - 20°C) [8]. Thus, the taxonomic composition of bacteria varies along

the salinity gradient with Alphaproteobacteria and Gammaproteobacteria dominating in more

saline waters whereas Actinobacteria, Betaproteobacteria and Planctomycetes are favored by

lower salinity [17-19]. The strong seasonality in the Baltic Sea also shapes bacterial community

structure and dynamics: Alphaproteobacteria are present throughout the year [16,19],

Bacteroidetes, Beta- and Gammaproteobacteria are associated with phytoplankton spring

blooms [17,20,21] and Verrucomicrobia, Actinobacteria and Planctomycetes appear later in the

year [18,22]. Although it is clear that both physical and chemical properties affect bacterial

communities, most of these studies have been restricted to specific sub-basins or did not capture

the full development of the spring bloom. The potential of diatoms to shape the bacterial

communities towards the dominance of Gamma-, Betaproteobacteria and Bacteriodetes and to

increase heterotrophic production in the Baltic Sea was recently demonstrated a in mesocosm

study [23]. However, so far no study has addressed this issue in natural conditions or across the

naturally occurring environmental gradients in the Baltic Sea.

The aim of this study was to investigate how differences in the taxonomic composition of

phytoplankton spring bloom communities (diatom- or dinoflagellate-dominated), affect the

community structure and dynamics of the associated bacterioplankton across different areas of

the Baltic Sea. We collected samples from four consecutive years (2013–2016) during and after

the spring bloom, from the northern Gulf of Bothnia to the southern Baltic Proper. Based on

our previous observations from laboratory studies [23], we hypothesized that the predicted

increase in dinoflagellates may lead to less productive bacterial communities dominated by less

copiotrophic bacterial groups.

Material and Methods

Study area, sampling design and environmental variables

The water samples were collected during four cruises (2013–2016) on board of the R/V Aranda.

In total, 127 stations were sampled and the area spanned from the southern Baltic Proper (BP,

55°22'N) to the northern Bay of Bothnia (BoB, 65°53'N, Fig 1). At all stations, the water was

collected from 3 m depth either using an oceanographic rosette with Niskin bottles of 5 L (n =

122) or from the flow-through system on board (n = 5). Salinity, temperature, and depth, were

recorded *in situ*. Inorganic nutrients, measured as nitrogen (NO_2+NO_3-N, NH_4-N), phosphorus

(PO_4-P), and dissolved silica (DSi), particulate organic carbon (POC), biogenic silicate (BSi),

dissolved organic carbon (DOC), dissolved organic nitrogen (DON) and chlorophyll *a* (Chl *a*)

concentrations were measured as described by Camarena-Gómez, et al. [23]. The particulate

organic carbon to chlorophyll *a* (POC:Chl *a*) ratio, based on weight ($\mu g\ L^{-1}$), was calculated as

an indicator of the trophic status (autotrophic or heterotrophic) of the system, considering

heterotrophic status those values larger than 200 [24].

Plankton community composition, carbon biomass and productivity

Samples for the determination of nano- and microplankton, including phytoplankton, were

determined as described by Lipsewers and Spilling [25]. A modified (carbon biomass instead of

biovolume) diatom/dinoflagellate (Dia/Dino) index = diatom biomass/diatom + dinoflagellate

biomass, was calculated for each station according to Wasmund, et al. [26]. The values higher

than 0.5 indicate diatom dominance whereas values lower than 0.5 indicate dinoflagellate

dominance. Primary production (PP) was estimated by measuring the incorporation of [14]C-

labelled sodium bicarbonate (DHI, Denmark) according to Gargas [27] and following the method

described in Camarena-Gómez, et al. [23].

The percent extracellular release of [14]C (PER) was calculated based on the dissolved

organic fraction ($DO^{14}C < 0.2\ \mu m$) of the total [14]C fixation after 24 h incubation. Since

substantial heterotrophic consumption of DOC (on average, 30–50 %) might take place during

the incubation period [28], the results are regarded as net DOC production rates.

Four different stages of the spring bloom were defined based on the concentration of

inorganic nutrients (PO_4^{3-} and $NO_3^-+NO_2^-$) and Chl *a* (Table S1): the bloom growth phase

(Growth), the peak of the bloom phase (Peak), the declining bloom phase (Decline) and after

the bloom phase (Postbloom). The $NO_3^-+NO_2^-$ was used for the N-limited sub-basins Gulf of

Finland (GoF), BP, Archipelago Sea (ArS), Åland Sea (ÅS) and Bothnian Sea (BS); and PO_4^{3-}

for the P-limited BoB subbasin. High Chl *a* was defined as within 20 % of the average peak

concentration during spring, which is different in the different sub-basins.

Bacterial heterotrophic production, abundance and community composition

Bacterial cell production (BPT) and protein production (BPL) were measured in triplicates by

using the dual labeling with [metil-^{3}H]-thymidine and [^{14}C (U)]-leucine (PerkinElmer Inc.,

Waltham, MA, USA) of samples processed by the centrifugation method as described in

Camarena-Gómez, et al. [23]. Bacterial abundances (BA) were determined in duplicates by flow

cytometry according to Gasol and Del Giorgio [29], applying some modifications of the method

described in Camarena-Gómez, et al. [23]. Bacterial community analysis including DNA

extraction and *Illumina* sequencing were done as described by Camarena-Gómez, et al. [23]. The

post-processing of the sequences was done according to the pipeline described by Logares [30].

The Operational Taxonomic Units (OTUs) were clustered at 99 % of similarity. After the

removal of singletons as well as the OTUs classified as chloroplasts and mitochondria, based

on the classification with SILVA v123, 2 247 OTUs and 2 250 000 reads were retained.

Samples with less than 7 000 reads (4 samples) were removed. The remaining samples (122)

were rarefied using *rrarefy* in R to the lowest number of reads (n = 7 024 reads). Downstream

analyses were carried out in the rarefied table, including 122 samples and 2 128 OTUs. Raw

sequence data will be deposited in the European Nucleotide Archive (acc. nun. XXXXX).

Statistical analyses

To assess whether there were differences in the diatom and dinoflagellate carbon biomass,

between the different phytoplankton bloom phases a non-parametric Wilcoxon rank sum test

was applied. This test was also applied to the bacterial diversity in order to test whether there

were differences between the two main bacterial community clusters observed. Non-metric

multidimensional scaling (NMDS) analysis was done on the Bray-Curtis dissimilarity matrix.

Mantel tests with Spearman correlation were applied to study the correlation between the

composition of bacterial and phytoplankton communities. Redundancy analysis (RDA) with

forward selection was applied to obtain the most important environmental variables that

explain bacterial community variance. This analysis included an ANOVA test in each step

(permutation test for RDA under reduced model, permutations: free, number of permutations:

9 999) to identify the significant environmental variables ($p < 0.05$). Spearman correlation was

used to obtain the correlation between the main bacterial groups and the significant

environmental variables as well as the bacterial activity. All figures and analyses were

performed with the software RStudio (RStudio Team 2015) with the packages *colorRamps,*

plotrix, rworldmap, rworldxtra, *vegan, ggplot2*, *ade4*, *ggpubr* and *Hmisc* and

ComplexHeatmap.

Results

Inter-annual and spatial variability in environmental variables and bloom dynamics

The environmental conditions varied largely within and between the years due to the location

of the different subbasins and time of the sampling (Table 1, Fig. 1A). For instance, salinity

ranged from ~ 8 in the southernmost station of the BP to ~2 in the northernmost part of the

BoB (Table 1). In general, the $NO_2^-+NO_3^-$ was depleted after the bloom with the exception of

the P-limited BoB, where the $NO_2^-+NO_3^-$ concentration was similar to the stations in the

Growth phase and phosphate was low. The lowest dissolved DSi concentration was measured

in the GoF in 2014 while the maximum DSi concentration was detected in the Kemi river plume

in the northernmost BoB. The DOC and DON were highest in the ArS in 2014, and higher in

the GoF and the Gulf of Bothnia compared with the BP.

Overall, diatom carbon biomass was significantly higher in the Growth and Peak

phytoplankton bloom phases compared to the Decline and Postbloom phases (Wilcoxon test, p

< 0.05), whereas dinoflagellate carbon biomass was also significantly high in the Decline phase

(Wilcoxon test, $p < 0.05$). However, no clear increasing trend in the biomass of dinoflagellates

compared to diatoms was observed across the studied years. In 2013, we sampled right after

the ice melt in the Gulf of Finland and the water temperature was low in the entire sampling

area (Table 1). The spring bloom was in the Growth to Decline phase in the 2013 cruise (Table

1), and was largely dominated by diatoms, displaying much higher carbon contribution to total

biomass than in the other three cruises (Fig. 1B). In 2014, the temperature was 4-5.5 °C and

the spring bloom was in a more advanced phase with low phytoplankton biomass (Fig. 1B). In

2015, the conditions were similar to 2014, with the exception of the Gulf of Bothnia where the

spring bloom was in the Peak to Decline phase at the time of the sampling. In 2016, we sampled

the same areas and time of the year as 2013, but the temperature was warmer in the BP

subbassin and the spring bloom was more advanced in the latest year than in the first sampling

cruise (Table 1). In these warmer cruises, dinoflagellates contributed > 300 µg C L^{-1} to the

carbon biomass at some stations (Fig. 1B).

High PP was observed during the Growth and Peak phases (Fig. 1C), following the

tendency of the Chl *a* and coinciding with the stations that presented high carbon biomass (Fig.

2A, Table S2). The PER, measured after 24 h, was high in the Decline and Postbloom phase in the BP and the BoB, in contrast to PP (Fig. 1C, Table S2). The trophic status, estimated as the POC:Chl *a* ratio, was in general low, except at some stations in the BP and the BoB (Table 1).

Changes in plankton community composition

We observed pronounced differences in taxonomic composition of the studied plankton communities (mainly phytoplankton and nano-microzooplankton here) across the different subbasins and years, which appeared mainly shaped by the inorganic nutrient concentrations and the water temperature (Fig. 2A, B). Diatoms largely dominated in the GoF and ÅS (Table 1), with high contributions of *Achnanthes taeniata*, *Skeletonema marinoi*, *Thalassiosira levanderi*, and *Chaetoceros* spp., whereas in the Bothnian Sea, *T. baltica* dominated during the Peak bloom phase (Fig. 2A). Dinoflagellates were dominated by the dinoflagellate complex, mainly formed by *Biecheleria baltica* and/or *Gymnodinium corollarium* in the GoF and BP. The chain forming *Peridiniella catenata* was present in all the subbasins except at the stations with clearly low carbon biomass (Fig. 2A). Other autotrophic dinoflagellates (*Heterocapsa* spp.) and heterotrophic/mixotrophic dinoflagellates (e.g. *Protoperidinium* spp. and *Dinophysis* spp., respectively) were also present, but their contribution to the total plankton carbon biomass was < 5 %. The contribution of other autotrophic plankton organisms, such as Crytophyceae, Chrysophyceae, Prasinophyceae, Prymnesiophyceae, Chlorophyceae, to the total carbon biomass was also generally low (< 2 µg C L^{-1}).

The mixotrophic ciliate, *Mesodinium rubrum,* was present across all sub-basins having highest contribution in the BS and being in general more abundant in 2016 compared with other years (Fig. 2A). Heterotrophic ciliates were most abundant during the late phase of the bloom in the GoF and the northern BP. The silicoflagellate *Ebria tripartita* and the heterotrophic nanoflagellates (HNF) had generally low abundance but contributed more than 10 µg C L^{-1} to

the total carbon biomass during the Peak and Decline phases of the bloom also at some stations

in the GoF and the BP.

Bacterial community composition and its environmental and biological drivers

The taxonomic composition of the studied bacterial communities varied largely across the

stations sampled (Fig. 3). Overall, Alphaproteobacteria, mainly SAR11, dominated in all the

sampled years (Fig. 3A). Bacteroidetes were dominated by the class Flavobacteriia, which

comprised on average ~ 25 % of the total relative abundance. Within Flavobacteriia, the genus

Flavobacterium contributed 8–12 % to the total relative abundance during the Peak/Decline

phase in 2013 and 2016, together with *Fluviicola*, *Polaribacter* and NS3 marine group, whereas

Owenweeksia was found in the Growth phase. Gammaproteobacteria (*Crenothrix*) had higher

relative abundance (> 10 %) in the GoF and the ÅS in 2013 and 2016. The classes

Actinobacteria (hgcl clade) and Acidimicrobiia (CL500-29 marine group) had higher relative

abundance in 2014 and 2015. The phylum Phycisphaerae (CL500-3) was scarce but reached

its highest values (~ 8 %) in the Gulf of Bothnia in 2015.

Based on their overall taxonomic structure, the communities were clustered into two main

groups (Wilcoxon; $p < 0.001$) differing largely in taxonomic richness (Fig. 3B) and diversity

(Shannon index; table S2) as well as in the bacteria abundances of the major bacterial groups.

The group with high taxonomic diversity comprised communities sampled during 2013 and

2014, and showed higher relative abundances of Bacteroidetes, Gammaproteobacteria and

Actinobacteria, whereas the low taxonomic richness group (2015–2016) showed a dominance

of Alphaproteobacteria (SAR11 and Rhodobacteraceae) and Cyanobacteria (Fig. 3B).

Although the bacterial community structure was strongly influenced by the salinity gradient

(Fig. 3C), the shift in community structure observed between 2013–2014 and 2015–2016

seemed to be largely driven by certain phytoplankton groups such as the diatoms *A. taeniata,*

S. marinoi, T. levanderi and *Chaetoceros* spp*.,* as well as temperature and the POC:Chl *a* ratio

(Fig. 3C). The role of the phytoplankton community composition in shaping bacterial

assemblages was further supported by the strong correlations between bacterial and

phytoplankton taxonomic dissimilarity matrices found across the four years of study

(Spearman r values 0.6–0.8, Mantel test *p* = 0.0001; Table S3).

When exploring the individual correlations between different environmental and biological

factors and the abundances of the main bacterial groups (Fig. 4), we observed two groups of

bacterial taxa showing fundamentally contrasting relationships. Bacteroidetes (Flavobacteriia,

Cytophagia and Sphingobacteriia), Gamma- and Betaproteobacteria were strongly correlated

with diatoms and *E. tripartita,* and negatively correlated with temperature and POC:Chl *a* ratios

(Fig. 4). In contrast, Rhodobacteraceae, SAR11 and Cyanobacteria correlated positively with

temperature, POC:Chl *a* ratios, HNFs and ciliates, whereas the dominance of certain diatoms

or dinoflagellates caused decreases in their abundances (Fig. 4). Phycisphaerae correlated

strongly with the diatom *T. baltica*. Within dinoflagellates, the dinoflagellate complex was the

only group that correlated positively with all the bacterial taxa dominant in 2013–2014, but

also with Rhodobacteraceae (Fig. 4), which was more prevalent in 2015–2016 (Fig. 3B).

The observed changes in composition of the phytoplankton communities were

accompanied by a large spatial-temporal variation of bacterial abundances and production (Fig.

5). Overall, the bacterial production rates measured with leucine (BPL) were highest in 2013

(Fig. 5A, Table S2), having similar pattern of the PP (Fig. 2A), while when measured with

thymidine (BPT), the rates were highest in 2014 (Fig. 5A). In contrast, the highest BA values

were recorded at the stations sampled during the 2015 cruise (Fig. 5B; Table S2).

The differences in BCC had an impact on community functioning. Most of the bacterial

groups that were prevalent in the high diversity assemblages from 2013 and 2014, i.e. Beta-,

Gammaproteobacteria, Actinobacteria, Bacteroidetes, and more specifically Flavobacteriia

showed strong positive correlations with BPL and BPT (Fig. 5C−D; Table S4). In contrast, the

abundance of Alphaproteobacteria (SAR11 in particular), was negatively correlated with

bacterial production rates (Fig. 5E, Table S4).

Discussion

Our results demonstrate that the bacterioplankton community structure was strongly affected

by the different phytoplankton communities, either dominated by diatoms or dinoflagellates,

and by the different phases of the phytoplankton bloom. Although our results do not evidence

a clear trend of increasing dominance of dinoflagellates from 2013 to 2016, we did capture

much higher carbon biomass of diatoms during the 2013 cruise than in the other years. This

heterogeneity allowed us to explore how changes in the dominant phytoplankton groups are

translated into changes in the structure and functioning of the associated bacterioplankton

communities.

Dynamics of environmental variables and phytoplankton biomass

We found that the different environmental conditions, primarily temperature and inorganic

nutrient concentrations such as NO_3^- and PO_4^{3-}, shaped the phytoplankton community structure

and bloom dynamics in terms of Chl *a* concentration, PP and dominant species, in agreement

with previous studies in the Baltic Sea [9,31]. The phytoplankton community during the early

bloom development, with low temperature and high inorganic nutrient concentrations, was

clearly dominated by diatoms, in particular in the GoF. Diatom dominance was also associated

with high primary productivity suggesting an actively growing community. The prevailing

diatoms were the taxa that typically occur during the cold water season [32-34]. In the BS, the

Peak phase was dominated by the single diatom species *Thalassiosira baltica* along with high

carbon biomass of the mixotrophic ciliate *Mesodinium rubrum,* which both seem to be common

in this sub-basin [7]. The diatom-dominated communities co-occurred to a varying extent with

several dinoflagellates, most likely *Gymnodinium corollarium* and *Biecheleria baltica* [35,36] in

the GoF and the BP, similarly to the earlier observations [2,3]. The heterotrophic *Ebria tripartita*

was more abundant during diatom dominance, likely because it grazed on diatom species such

as *Skeletonema* sp. and *Thalassiosira* spp. [37].

During the declining phytoplankton bloom, the water was warmer, the Chl *a* concentration

and the total carbon biomass were lower and dinoflagellates contributed more to the biomass

than in the Growth and Peak phases. This is the typical situation in the GoF with diatoms being

more abundant in early spring and dinoflagellates gradually becoming more dominant as the

nutrients are depleted [10,38]. For instance, the stations sampled during 2014 presented low

phytoplankton biomass and also low DSi concentration in the GoF, suggesting that a diatom-

dominated bloom had settled out of the surface layer before the time of sampling. This probably

explained the similarity in structure between the bacterial communities and productivity found

in 2014 and those in the diatom-dominated stations of 2013. In the BoB, the low phytoplankton

carbon biomass was limited by phosphorus with excess NO_3^- and DSi and high DOC and DON

concentrations. This sub-basin is P-limited [9] and is more turbid due to terrestrial DOM inputs,

which limits light penetration resulting in light-limited primary production [39-41].

The abundance of nano- and microzooplankton (HNF, ciliates and *M. rubrum*) were also

higher during the Decline phase of the bloom, which, together with higher POC:Chl *a* ratios (

> 175), points to a more heterotrophic plankton community [24]. A decrease in phytoplankton

biomass and an increase in the proportion of dinoflagellates and heterotrophic organisms have

been projected for a future, warmer Baltic Sea [1], in addition to the earlier onset of the spring

bloom [42]. Our results represent different temperature scenarios, for example the 2013 and 2016

cruises sampled the same subbasins during varying environmental conditions with 2016 being

warmer and the bloom in a more advance stage, in line with these predictions.

Bacterial community dynamics and their environmental drivers

The changes in the phytoplankton community composition between the bloom phases resulted in pronounced shifts in the structure of the associated bacterial communities and likely related to the availability of the released DOM, largely produced by phytoplankton [43,44]. The availability of this DOM can lead to changes in bacterial community dynamics [45-47], which have been also reported in several field studies during phytoplankton spring blooms [16,17,48].

The studied bacterial communities clustered into two clearly different groups of high and low taxonomic richness and diversity, which corresponded to the communities sampled during 2013−2014, and those sampled during 2015−2016, respectively. Salinity and temperature appeared as relevant factors in shaping the bacterial communities in the Baltic Sea, similar to previous studies [20,22,49]. However, in our study, the salinity was not responsible for the segregation observed in the bacterial communities. Instead, changes in the carbon biomass of specific phytoplankton groups such as *Achnanthes taeniata*, *Chaetoceros* spp., *Skeletonema marinoi*, and *T. levanderi* emerged as some of the most important drivers in the configuration of bacterial communities. Thus, the diatom bloom observed in 2013 and the diatom bloom that had likely recently settled out of the surface layer in 2014 promoted a higher diversity of associated bacterial communities with more Gammaproteobacteria, Betaproteobacteria, and Bacteroidetes. This conclusion was further strengthened by the strong and positive correlations between the diatom species and these copiotrophic bacteria, in accordance with their preference for productive or bloom-like conditions. All these groups are known to be favored during diatom blooms [23,50]. In contrast, the lower diatom carbon biomass observed during 2015 and 2016 were associated with less diverse communities dominated by the SAR11 group in 2015 and also by Rhodobacteraceae (genus *Gemmobacter*) in 2016, supported by the strong negative correlations between this bacterial community structure and the phytoplankton carbon biomass. The conditions within the less productive stations sampled during 2015 and 2016 likely

promoted the dominance of SAR11, which typically occurs in nutrient-poor environments [51,52].

Other groups like Phycisphaerae, and in particular CL500-3, appeared to follow more clearly

the salinity gradient and was strongly associated to the presence of *T. baltica*. The presence of

Phycisphaerae has been associated to the lower salinity of the Gulf of Bothnia and to a higher

influence of allochthonous DOM [19]. Interestingly, temperature emerged as a one of the main

drivers for the structure of both, the phytoplankton and bacterioplankton communities,

suggesting that the predicted increases in water temperature along the Baltic Sea [53,54] may lead

to dramatic changes in the planktonic assemblages inhabiting this vulnerable system.

Within the high-diversity bacterial communities found in 2013 and 2014, we observed that

the dynamics of Gammaproteobacteria and Bacteroidetes differed between bloom phases and

subbasins. Bacteroidetes were always present with fluctuating relative abundance, whereas

Gammaproteobacteria, dominated by the genus *Crenothrix,* peaked shortly and were restricted

to those stations with complex diatom communities. These bacterial groups are known to

occupy different niches in terms of metabolic affinities: Flavobacteriia are capable of degrading

high molecular weight compounds, such as proteins, chitin and polysaccharides [48,55], whereas

Gammaproteobacteria are more specialized on breaking down low molecular weight

substrates, peaking shortly during increased organic and inorganic substrate availability [56,57].

Thus, their different patterns were likely related to the DOM availability and niche partitioning

during the highly dynamic phytoplankton blooms. This was supported by the lower net PER

during diatom dominance compared with dinoflagellate dominance, suggesting that diatoms

release more labile DOC, and therefore more rapidly consumed, than dinoflagellates (Lignell

1990).

The succession of taxa within Bacteroidetes in the different phases of the bloom was similar

to that observed in a mesocosm experiment simulating diatom and dinoflagellate blooms with

Baltic Sea bacterial communities [23]. In the Growth phase, *Owenweeksia*, *Flabovacterium* and

Fluviicola were more dominant, whereas the genera *Polaribacter* and NS3 marine group were

favored in the declining phase. Within Gammaproteobacteria, the presence of *Crenothrix* in

surface waters was surprising since this taxon is a methane-oxidizer specialized in the uptake

of single-carbon substrate, such as methane and methanol, but also acetate and glucose [58]. This

filamentous bacterium may have benefited by the increase of the suitable substrates, either by

fresh and labile DOM released or as a product from a metabolic cascade after the degradation

process carried out by other bacterial groups.

A lower diversity of Bacteroidetes was observed in the presence of the diatom *T. baltica* in

the BS. A previous study has reported the production of certain polyunsaturated aldehydes

(PUAs) by *Thalassiosira spp.* [59], which can affect the bacterial protein production of

Bacteroidetes negatively [60]. This could have played a role in the reduced growth of this bacterial

group. Any anti-bacterial effects of phytoplankton are not well studied, but there are indications

that some species inhibit bacterial growth [50].

Links between bacterial community composition and ecosystem functioning

The observed changes in bacterial community structure resulted in pronounced changes in

the bacterial production of the studied communities. Most stations sampled during 2013 and

2014 showed much higher rates of BPL compared to the stations sampled in 2015 and 2016

and were similar or, in some cases, larger than those reported by previous studies conducted at

similar temperatures during the spring in the Baltic Sea [16,61]. The larger values of BPL

coincided with the higher abundances of copiotrophic groups such as Betaproteobacteria,

Gammaproteobacteria and Bacteroidetes associated to diatom dominance. The variation in

BPL between studies conducted at similar temperatures may be due to the different dominating

phytoplankton groups that produce DOM of different lability. This emphasizes the importance

of the phytoplankton taxonomical composition as the driver of the bacterial carbon cycling

mediated by the presence of specific bacterial taxa. Large peaks in BPT were also observed in 2014 indicating an actively dividing community and mostly related to the increases in relative abundance of Actinobacteria, which is known to occur after phytoplankton blooms [20] and correlate positively with thymidine incorporation [62]. In contrast, SAR11 correlated negatively with BPL, which agrees with the small and slow-growing pattern of this clade typically occurring in oligotrophic ecosystems [51,52]. Similar bacterial responses were observed in a mesocosm study, where the growth of diatoms resulted in higher bacterial production and higher dominance of Flavobacteriia, Gammaproteobacteria and Cytophagia than in the presence of dinoflagellates [23]. The robustness of the observed patterns in our study, linking BCC and functioning, are essential in microbial ecology since the studies that explore these relationship are scarce in the Baltic Sea [63].

Gammaproteobacteria and Bacteroidetes are known to be quickly grazed by HNF [51], which prefer to graze on actively growing bacteria. This may explain the low bacterial abundance of the active community observed in 2013. The dinoflagellate complex also correlated positively with the relative abundances of these bacterial groups. However, previous experimental observations reported a decrease in bacterial productivity upon the addition of *Biecheleria baltica* [23], one of the species forming the dinoflagellate complex.

Conclusions

Our results during and after phytoplankton spring blooms highlight the differences in BCC, diversity and productivity under distinct phytoplankton communities dominated by diatoms and/or dinoflagellates. Diatoms like *Skeletonema marinoi, Achnanthes taeniata* and *Chaetoceros* spp. likely release high-quality DOM that boosts the bacterial production and enhances the development of highly diverse bacterial communities dominated by Gamma-, Betaproteobacteria and Bacteroidetes. In contrast, less diverse and less productive

bacterioplankton communities dominated by SAR11 and Cyanobacteria are observed in

presence of more dinoflagellates. Based on our results, it is not the overall dominance of

dinoflagellates or diatoms what drives the largest changes in bacterial community structure,

but rather the presence of particular taxonomic groups. Finally, although temperature appears

as a main factor shaping both bacterial and phytoplankton communities, higher temperatures

are not necessarily related to higher bacterial production rates. Thus, it is needed to incorporate

knowledge on the changes in plankton community composition to be able to understand and

predict the biogeochemical consequences of expected temperature increases and associated

changes in phytoplankton blooms in Baltic Sea waters.

Conflict of interest

The authors declare no conflict of interest

Acknowledgements

This study was funded by the Walter and Andrée de Nottbeck Foundation and the Academy of

Finland (decision numbers 259164 and 292711). We would like to thank the staff at Tvärminne

Zoological Station and the staff on board the Aranda vessel for help with the analytical

measurements and sampling during the experiments.

References

1   Legrand, C. *et al.* Interannual variability of phyto-bacterioplankton biomass and

production in coastal and offshore waters of the Baltic Sea. *Ambio* 2015; **44:** 427–438.

2   Klais, R., Tamminen, T., Kremp, A., Spilling, K. & Olli, K. Decadal-scale changes of

dinoflagellates and diatoms in the anomalous Baltic Sea spring bloom. *PLoS One* 2011;

6: e21567

3   Wasmund, N., Tuimala, J., Suikkanen, S., Vandepitte, L. & Kraberg, A. Long-term

trends in phytoplankton composition in the western and central Baltic Sea *J Mar Syst*

2011; **87**: 145–159.

4   Wasmund, N. & Uhlig, S. Phytoplankton trends in the Baltic Sea. *ICES J Mar Sci* 2003;

60: 177–186.

5   Klais, R. *et al.* Spring phytoplankton communities shaped by interannual weather

variability and dispersal limitation: Mechanisms of climate change effects on key

coastal primary producers. *Limno Oceanogr* 2013; **58**: 753–762.

6   Kahru, M. & Nômmann, S. The phytoplankton spring bloom in the Baltic Sea in 1985,

1986: multitude of spatio-temporal scales. *Cont Shelf Res* 1990; **10**. 329–354.

7   Andersson, A., Hajdu, S., Haecky, P., Kuparinen, J. & Wikner, J. Succession and

growth limitation of phytoplankton in the Gulf of Bothnia (Baltic Sea). *Mar Biol* 1996;

126: 791–801.

8   Hagström, Å., Azam, F., Kuparinen, J. & Zweifel, U.-L. in *A systems analysis of the*

Baltic Sea. Springer 2001: 177–210.

9   Tamminen, T. & Andersen, T. Seasonal phytoplankton nutrient limitation patterns as

revealed by bioassays over Baltic Sea gradients of salinity and eutrophication. *Mar Ecol*

Prog Ser 2007; **340**: 121–138.

10   Heiskanen, A. in *Monographs of the Boreal Environment Research 8*. Finnish

Environment Institute, 1998.

11   Spilling, K. & Markager, S. Ecophysiological growth characteristics and modeling of

the onset of the spring bloom in the Baltic Sea. *J Mar Syst* 2008; **73**: 323–337.

12   Castillo, C. R., Sarmento, H., Alvarez-Salgado, X. A., Gasol, J. M. & Marraséa, C.

Production of chromophoric dissolved organic matter by marine phytoplankton. *Limno*

Oceanogr 2010; **55**: 446–454.

13 Myklestad, S. M. Release of extracellular products by phytoplankton with special

emphasis on polysaccharides. *Sci Total Environ* 1995; **165**: 155–164.

14 Spilling, K., Kremp, A., Klais, R., Olli, K. & Tamminen, T. Spring bloom community

change modifies carbon pathways and C: N: P: Chl a stoichiometry of coastal material

fluxes. *Biogeosciences* 2014; **11**: 7275-7289.

15 Spilling, K. & Lindström, M. Phytoplankton life cycle transformations lead to species-

specific effects on sediment processes in the Baltic Sea. *Con Shelf Res* 2008; **28**: 2488–

2495.

16 Lindh, M. V. *et al.* Disentangling seasonal bacterioplankton population dynamics by

high-frequency sampling. *Environ Microbiol* 2015; **17**: 2459–2476.

17 Bunse, C. *et al.* Spatio-temporal interdependence of bacteria and phytoplankton during

a Baltic Sea spring bloom. *Front Microbiol* 2016; **7**: 517.

18 Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity

gradient of the Baltic Sea. *ISME J* 2011; **5**: 1571–1579.

19 Rieck, A., Herlemann, D. P., Jürgens, K. & Grossart, H.-P. Particle-associated differ

from free-living bacteria in surface waters of the Baltic Sea. *Front Microbiol* 2015; **6**:

1297.

20 Hugerth, L. W. *et al.* Metagenome-assembled genomes uncover a global brackish

microbiome. *Genome Biol* 2015; **16**: 279.

21 Laas, P. *et al.* Redox-specialized bacterioplankton metacommunity in a temperate

estuary. *PloS One* 2015; **10**: e0122304.

22 Andersson, A. F., Riemann, L. & Bertilsson, S. Pyrosequencing reveals contrasting

seasonal dynamics of taxa within Baltic Sea bacterioplankton communities. *ISME J*

2010; **4**: 171–181.

23   Camarena-Gómez, M. T. *et al.* Shifts in phytoplankton community structure modify

bacterial production, abundance and community composition. *Aquat Microb Ecol*

2018; **81**: 149–170.

24   Vargas, C. A. *et al.* The relative importance of microbial and classical food webs in a

highly productive coastal upwelling area. *Limnol Oceanogr* 2007; **52**,:1495–1510.

25   Lipsewers, T. & Spilling, K. Microzooplankton, the missing link in Finnish plankton

monitoring programs. *Boreal Environ Res* 2018; **23**, 127–137.

26   Wasmund, N. *et al.* The diatom/dinoflagellate index as an indicator of ecosystem

changes in the Baltic Sea. 1. Principle and handling instruction. *Front Mar Sci* 2017; **4**:

22.

27   Gargas, E. A manual for phytoplankton primary production studies in the Baltic. *Baltic*

Mar Biologists 1975; **2**: 1–88. (Copies available from: Water Quality Institute, 11

Agern Allé, DK-2970 Hørsholm, Denmark).

28   Morán, X. A. G. & Estrada, M. Phytoplanktonic DOC and POC production in the

Bransfield and Gerlache Straits as derived from kinetic experiments of 14C

incorporation. *Deep Sea Research Part II: Topical Studies in Oceanography* 2002; **49**:

769–786.

29   Gasol, J. M. & Del Giorgio, P. A. Using flow cytometry for counting natural planktonic

bacteria and understanding the structure of planktonic bacterial communities. *Sci Mar*

2000; **64**: 197–224.

30   Logares, R. Workflow for analysing MiSeq amplicons based on Uparse v1. 5. 2017

Zenodo10 **5281**.

31   Fleming, V. & Kaitala, S. Phytoplankton spring bloom intensity index for the Baltic

Sea estimated for the years 1992 to 2004. *Hydrobiologia* 2006; **554**: 57–65.

Ikävalko, J. & Thomsen, H. A. The Baltic Sea ice biota (March 1994): a study of the protistan community. *Eur J Protistol* 1997; **33**: 229–243.

Lignell, R. Excretion of organic carbon by phytoplankton: its relation to algal biomass, primary productivity and bacterial secondary productivity in the Baltic Sea. *Mar Ecol Prog Ser* 1990; **68**: 85–99 *Oldendorf.*

Spilling, K. On the ecology of cold-water phytoplankton in the Baltic Sea. 2007.

Olli, K. & Trunov, K. Abundance and distribution of vernal bloom dinoflagellate cysts in the Gulf of Finland and Gulf of Riga (the Baltic Sea). *Deep Sea Research Part II: Topical Studies in Oceanography* 2010; **57**: 235–242.

Sundström, A. M. *et al. Gymnodinium corollarium* sp. nov.(dinophyceae)—a new cold-water dinoflagellate responsible for cyst sedimentation events in the baltic sea 1. *J Phycol* 2009; **45**: 938–952.

Hargraves, P. E. The ebridian flagellates Ebria and Hermesinum. *Plankton Biol Ecol* 2002; **49**: 9–16.

Spilling, K. *et al.* Shifting Diatom—Dinoflagellate Dominance During Spring Bloom in the Baltic Sea and its Potential Effects on Biogeochemical Cycling. *Front Mar Sci* 2018; **19**.

Andersson, A. *et al.* Projected future climate change and Baltic Sea ecosystem management. *Ambio* 2015; **44**: 34–356.

Rowe, O. F. *et al.* Major differences in dissolved organic matter characteristics and bacterial processing over an extensive brackish water gradient, the Baltic Sea. *Mar Chem* 2018; **202**: 27–36.

Sandberg, J., Andersson, A., Johansson, S. & Wikner, J. Pelagic food web structure and carbon budget in the northern Baltic Sea: potential importance of terrigenous carbon. *Mar Ecol Prog Ser* 2004; **268**: 13–29.

42   Groetsch, P. M., Simis, S. G., Eleveld, M. A. & Peters, S. W. Spring blooms in the

Baltic Sea have weakened but lengthened from 2000 to 2014. *Biogeosciences* 2016; **13:**

4959–4973.

43   Ducklow, H. W. Production and fate of bacteria in the oceans. *Bioscience* 1983; **33**:

494–501.

44   Kirchman, D. Measuring bacterial biomass production and growth rates from leucine

incorporation in natural aquatic environments. *Methods in microbiology* 2001; **30**: 227–

237.

45   Pinhassi, J. *et al.* Changes in bacterioplankton composition under different

phytoplankton regimens. *Appl Environ Microbiol* 2004; **70**: 6753–6766.

46   Riemann, L., Steward, G. F. & Azam, F. Dynamics of bacterial community composition

and activity during a mesocosm diatom bloom. *Appl Environ Microbiol* 2000; **66**: 578–

587.

47   Sarmento, H. *et al.* Phytoplankton species-specific release of dissolved free amino acids

and their selective consumption by bacteria. *Limnol Oceanogr* 2013; **58**: 1123–1135.

48   Teeling, H. *et al.* Substrate-controlled succession of marine bacterioplankton

populations induced by a phytoplankton bloom. *Science* 2012; **336**: 608–611.

49   Herlemann, D. P., Lundin, D., Andersson, A. F., Labrenz, M. & Jürgens, K.

Phylogenetic signals of salinity and season in bacterial community composition across

the salinity gradient of the Baltic Sea. *Front Microbiol* 2016; **7**.

50   Amin, S. A., Parker, M. S. & Armbrust, E. V. Interactions between diatoms and

bacteria. *Microbiol Mol Biol Rev* 2012; **76**: 667–684.

51   Alonso-Sáez, L. *et al.* Bacterial assemblage structure and carbon metabolism along a

productivity gradient in the NE Atlantic Ocean. *Aquat Microb Ecol* 2007; **46**: 43–53.

52 Giovannoni, S. J. *et al.* Genome streamlining in a cosmopolitan oceanic bacterium.

Science 2005; **309**: 1242–1245.

53 Meier, H. *et al.* Modeling the combined impact of changing climate and changing

nutrient loads on the Baltic Sea environment in an ensemble of transient simulations

for 1961–2099. *Clim Dynam* 2012; **39**: 2421–2441.

54 Thomas, D. N. *et al.* in *Biological oceanography of the Baltic Sea*. Springer 2017; 333–

357.

55 Cottrell, M. T. & Kirchman, D. L. Natural assemblages of marine proteobacteria and

members of the Cytophaga-Flavobacter cluster consuming low-and high-molecular-

weight dissolved organic matter. *Appl Environ Microbiol* 2000; **66**: 1692–1697.

56 Gómez-Consarnau, L., Lindh, M. V., Gasol, J. M. & Pinhassi, J. Structuring of

bacterioplankton communities by specific dissolved organic carbon compounds.

Environ Microbiol 2012; **14**: 2361–2378.

57 Pinhassi, J. & Berman, T. Differential growth response of colony-forming α-and γ-

proteobacteria in dilution culture and nutrient addition experiments from Lake Kinneret

(Israel), the eastern Mediterranean Sea, and the Gulf of Eilat. *Appl Environ Microbiol*

2003; **69**: 199–211.

58 Stoecker, K. *et al.* Cohn's Crenothrix is a filamentous methane oxidizer with an unusual

methane monooxygenase. *Proceedings of the National Academy of Sciences* 2006; **103**:

2363–2367.

59 Wichard, T., Poulet, S. A. & Pohnert, G. Determination and quantification of α, β, γ, δ-

unsaturated aldehydes as pentafluorobenzyl-oxime derivates in diatom cultures and

natural phytoplankton populations: application in marine field studies. *J Chromatogr B*

2005; **814**: 155–161.

60   Balestra, C., Alonso-Sáez, L., Gasol, J. M. & Casotti, R. Group-specific effects on

coastal bacterioplankton of polyunsaturated aldehydes produced by diatoms. *Aquat*

Microb Ecol 2011; **63**: 123–131.

61   von Scheibner, M., Herlemann, D. P., Lewandowska, A. M. & Jürgens, K. Phyto-and

bacterioplankton during early spring conditions in the Baltic Sea and response to short-

term experimental warming. *Front Mar Sci* 2018; **5**: 231.

62   Pérez, M. T., Hörtnagl, P. & Sommaruga, R. Contrasting ability to take up leucine and

thymidine among freshwater bacterial groups: implications for bacterial production

measurements. *Environ Microbiol* 2010; **12**: 74–82.

63   Lindh, M. V. & Pinhassi, J. Sensitivity of Bacterioplankton to Environmental

Disturbance: a Review of Baltic Sea Field Studies and Experiments. *Front Mar Sci*

2018; **5**: 361.

 Figure legends

Figure 1. Sampling transects and phytoplankton dynamics across the studied period. A) Map

with the sampling transects during the four cruises: 2013 (red), 2014 (dark blue), 2015 (light

blue) and 2016 (green). B) Total carbon biomass of dinoflagellates (black) and diatoms (blue)

during the cruises and bloom phases. C) Percentage of extracellular release (PER, black) and

primary production (PP, blue) in the different sub-basins. See Table 1 for more details of the

x-axis abbreviations.

Figure 2. Plankton dynamics in A) total carbon biomass of the main plankton groups and the

chlorophyll *a* (Chl *a*) concentration at each station. The years are indicated on top of the figure

and the bloom phases and the sampled subbasins are at the bottom of the figure. See Table 1

for more details of x-axis abbreviations. B) Non-metric multidimensional scaling (NMDS) plot

based on Bray-Curtis distances between plankton community taxonomic composition (N =

126, stress = 0.1602). The vectors indicate the environmental factors most strongly explaining

the observed distribution based on RDA forward selection (NO_2^--NO_3^{-1}; nitrate-nitrite, Tem;

temperature, PO_4^{3-}; phosphate, POC:Chl *a;* particulate organic carbon to Chl *a* ratio, diatoms

and dinos; total carbon of diatoms and dinoflagellates). The colour gradient indicates the water

temperature, shape indicates the different bloom phases and size indicates the log-transformed

total carbon biomass.

Figure 3. Bacterial community dynamics and their environmental drivers. A) Relative

abundance of the bacterial groups (class-family and genus level) that contribute more than 0.5

% of the total OTUs. Abbreviations as in figure 1. B) Non-metric multidimensional scaling

(NMDS) plot based on Bray-Curtis distances of bacterial community structure (N = 122, stress

= 0.134). Colour-code indicates the years (see Fig. 1) and size indicates the OTU richness in

each community. Vectors indicate the main bacterial groups found in the four years. The

abbreviations in the legend of Fig. A) and in Fig. B) indicate the main clades found during the

sampling: Acidi; Acidimicrobiia, Actino; Actinobacteria, Flavo; Flavobacteriia; Sphingo;

Sphingobacteriia; Cyano; Cyanobacterias, Alpha; Alphaproteobacteria; Beta;

Betaproteobacteria; Gamma; Gammaproteobacteria; Phycis; Phycisphaerae (Phy). C) Non-

metric multidimensional scaling (NMDS) plot linking bacterial community structure with

environmental variables and plankton community. Colour-code indicates the salinity gradient

and shape indicates the phytoplankton bloom phase. The vectors indicate the significant

environmental variables as follow: *Achnanthes taeniata*, *Skeletonema marinoi*, *Thalassiosira*

levanderi, *Chaetoceros* spp., Dinos (dinoflagellate complex), *Ebria tripartite*, *T. baltica*, HNF

(heterotrophic nanoflagellates), temperature, POC Chl *a*, *Mesodinium rubrum*, ciliates, other

autotrophs and salinity.

Figure 4. Heatmap with Spearman correlations between significant environmental variables,

phytoplankton or heterotrophic plankton groups and the main bacterial groups. Colour-code

indicates the correlation coefficients (rho values). The environmental factors (columns) are as

in Fig. 3. In addition, BA = bacterial abundance; BSi = biogenic silicate; BPT = bacterial

production thymidine and BPL = bacterial production leucine. N = 122.

Figure 5. Bacterioplankton productivity and abundance during the sampled spring blooms and

the link to bacterial community. A) Bacterial heterotrophic production based on leucine

incorporation (BPL, black) and thymidine incorporation (BPT, blue), and B) bacterial

abundance. The x-axis indicates the sub-basins across the sampling. See Fig. 1 for more detail

of legend abbreviation. The dashed lines separate the different years. C-E) Spearman

correlations between bacterial production measured as leucine uptake (BPL) or thymidine

uptake (BPT) and the relative abundance (%) of Flavobacteriia (C), Actinobacteria (D) and

SAR11 (E), during the study. Rho values and p values are presented on the figure. N = 122; p

< 0.05.

Tables

Table 1. Environmental and biological variables measured along the four sampling transects

(2013−2016). Values represent the range between the minimum and maximum values, in bold,

within the four years. The different sub-basins covered are indicated: Gulf of Finland (GoF),

Baltic Proper (BP), Åland Sea (ÅS), Archipelago Sea (ArS), Bothnian Sea (BS), Kvark (Kv)

and Bay of Bothnia (BoB). The different phytoplankton bloom phases are indicated: Growth

(G), Peak (P), Decline (D) and Post-bloom (PB), defined based on the Chl *a*, nitrate (NO_3^-) and

phosphate (PO_4^{3-}) concentrations during the sampling at each subbasin (see Methods and Table

S1). The inorganic nutrients and dissolved organic nitrogen are presented in µmol L^{-1} and the

dissolved organic carbon is presented in mmol L^{-1}. The particulate organic carbon to Chl *a*

(POC:Chl *a*) ratio indicates the trophic status (considering heterotrophic status those values

larger than 200) and the diatoms:dinoflagellates index (Dia/Dino index) indicates diatom

dominance or dinoflagellates dominance with values higher or lower than 0.5, respectively.

Figure 1

A

B

Figure 3

Figure 4

Table 1

	#of Stations	Subbasins	Bloom phase	Temp (° C)	Saliniy (psu)	NO_2^--NO_3^-	NH_4^+	PO_4^{3-}	DSi	DOC	DON	POC:Chl a	Dia/Dino index
2013 April 10th-26th	29	GoF	G, P, D	0.76-1.98	5.09-6.38	0-5.85	0.10-0.39	0.19-0.48	7.84-19.75	0.30-1.08	10.06-45.57	33.36-52.47	0.30-0.86
	11	BP	G, P	1.62-2.72	5.97-7.24	0-2.08	0.04-0.19	0.06-0.33	8.27-14.24	0.41-1.01	13.49-30.78	38.96-100.35	0.09-0.94
	4	ÅS	G, P	1.29-1.59	5.45-6.16	0-2.59	0.07-0.13	0.02-0.35	8.92-13.10	0.35-0.51	16.07-17.85	44.54-56.39	0.36-0.74
2014 May 5th-10th	9	GoF	D, PB	4.25-5.18	5.03-6.01	0-0.26	0.5-0.24	0.12-0.33	2.27-9.10	0.50-1.05	16.07-28.70	49.43-80.00	0.11-0.29
	2	BP	D, PB	4.62-5.69	6.56-6.63	0	0.05	0.28	7.60-11.60	0.44-0.52	15.16-16.06	106.82-176.30	0.01-0.36
	4	ArS	D, PB	4.58-5.17	5.90-6.29	0-0.4	0.08-0.11	0.20-0.33	6.01-8.10	0.60-1.12	16.02-67.86	88.28-208.84	0.05.0.49
2015 May 4th-15th	5	GoF	D, PB	3.91-5.35	4.96-5.86	0-0.02	0.08-0.44	0.18-0.36	5.50-11.20	0.46-1.07	16.36-25.40	75.28-119.28	0.12-0.40
	6	ArS	P, PB	4.41-6.09	5.75-6.38	0-0.23	0.06-0.13	0.06-0.33	5.60-11.10	0.32-0.64	11.33-18.58	58.81-190.20	0.04-0.32
	4	ÅS	D	5.03-6.22	5.41-6.06	0	0-0.06	0.05-0.22	6.70-8.70	0.37-0.47	12.73-13.97	76.75-108.31	0.17-0.68
	4	BP	D, PB	4.52-5.69	5.72-6.56	0-0.44	0.05-0.34	0.27-0.40	7.6-11.8	0.37-0.55	15.16-18.95	83.49-149.96	0.09-0.33
	10	BS	P	3.59-4.16	5.12-5.54	0-0.38	0.05-0.13	0-0.21	8.10-12.70	0.33-1.07	12.40-18.66	31.93-82.12	0.48-0.90
	9	Kv, BoB	P, D	1.35-4.73	2.20-5.09	0-7.47	0.04-0.35	0-0.02	9.40-55.70	0.35-0.68	15.73-55.53	70.71-223.37	0.36-0.68
2016 April 4th-15th	4	GoF	G, P, D	1.56-2.89	4.95-5.20	0-6.64	0.14-0.22	0.34-0.61	12.21-19.09	0.44-0.58	14.71-31.52	25.85-57.10	0.13-0.77
	20	BP	G, P, D, PB	2.47-5.90	5.92-7.77	0-1.89	0-1.19	0.16-0.57	11.77-17.65	0.32-0.62	10.73-22.10	44.49-260.31	0.01-0.58
	4	ÅS	P	2.84-3.28	5.51-5.68	0-0.17	0.11-0.18	0.12-0.18	15.57-18.17	034-0.99	12.40-14.54	58.41-67.04	0.61-0.80