Joint seasonality in geographic and ecological spaces,
illustrated with a partially migratory bird

Mathieu Basille iD

Department of Wildlife Ecology and Conservation, Fort Lauderdale Research and Education Center, University of Florida, Davie, FL 33314, USA.
Corresponding author: basille@ufl.edu

James Watling iD

Department of Biology, John Carroll University, University Heights, OH 44118, USA.

Stephanie Romañach iD

Wetland and Aquatic Research Center, U.S. Geological Survey, Fort Lauderdale, FL 33314, USA.

Rena Borkhataria

Department of Wildlife Ecology and Conservation, Everglades Research and Education Center, University of Florida, Belle Glade, FL 33430, USA.

Abstract.   As most species live in seasonal environments, considering varying

conditions is essential to understand species dynamics in both geographic and ecological

spaces. Both resident and migratory species need to contend with seasonality, and balance

settling in favorable areas with tracking favorable environmental conditions during the

year. We present an exploratory framework to jointly investigate a species' niche in

geographic and ecological spaces, applied to wood storks (*Mycteria americana*), which are

partially migratory wading birds, in the southeastern U.S. We concurrently described

monthly geographic distributions and climatic niches based on temperature and

precipitation. Geographic distributions of wood storks were more similar throughout the

year than climatic niches were, suggesting that birds stay within specific areas seasonally,

rather than tracking areas of similar climate. However, wood storks expressed consistent

selection of warm areas during the winter, and wet areas during the summer, indicating

that the selection of seasonal ranges may be directly related to environmental conditions

across the entire range. Our flexible framework, which simultaneously considered

geographic and ecological spaces, suggested that tracking climate alone did not explain

seasonal distributions of wood storks in breeding and non-breeding areas.

Key words: climatic niche, ecological niche, kernel overlap, range, wood stork

₂₂   Introduction

₂₃ At the core of ecology is the question of where do animals live? Early on, ecologists

₂₄ acknowledged the dual nature of this issue, by investigating species ranges (i.e., their

₂₅ location in geographic space, Candolle 1855, Darwin 1859, Wallace 1876), and species

₂₆ ecological niches (i.e., their location in ecological space, Grinnell 1917, Hutchinson 1957),

₂₇ actually reflecting what has been termed "Hutchinson's duality" (Colwell and Rangel

₂₈ 2009). More recent is the joint investigation of the effect of a change in one space on the

₂₉ other—e.g., how range constraints caused by migration or breeding affect a species' niche

₃₀ (Kearney and Porter 2009), or how climate change may affect species' range limits, (Chen

₃₁ et al. 2011). However, the dynamic nature of both spaces in time—i.e., their

₃₂ *seasonality*—often goes unacknowledged, despite the fact that most species live in seasonal

₃₃ environments (Fretwell 1972), with strongly cyclic variations in resource availability and

₃₄ conditions throughout the year. Here, we illustrate the use of a generalized framework for

₃₅ joint seasonality in geographic and ecological (specifically climatic) spaces, with the case of

₃₆ wood storks (*Mycteria americana*) in the southeastern United States.

₃₇   The direct correspondence between environmental conditions (i.e., ecological niche) and

₃₈ a species range at equilibrium is the fundamental assumption of species distribution

₃₉ modeling (Araújo and Pearson 2005, Pearman et al. 2008, Elith and Leathwick 2009).

₄₀ However, variations of environmental conditions during the year has led to the recognition

₄₁ of seasonal variation in ecological niches, with species responses varying on a continuum

₄₂ from "niche trackers" to "niche switchers" (Martínez–Meyer et al. 2004, Nakazawa et al.

₄₃ 2004). Broennimann et al. (2011) proposed a framework to quantify niche overlap using

₄₄ essentially kernel overlap metrics in a simplified 2-dimensional niche. Relying on this

₄₅ framework, several studies investigated seasonality in ecological niches: for instance, Laube

₄₆ et al. (2015) estimated overlap between breeding and non-breeding niches of warblers,

₄₇ while Gómez et al. (2016) investigated seasonal niche overlap of passerine birds.

₄₈   Movement has evolved as the primary means to manage heterogeneous environments in

space or in time (Nathan et al. 2008). Motile animal species can be placed on a gradient from sedentary to migratory, through nomadic (Chapman et al. 2011, Cagnacci et al. 2011, Jonzén et al. 2011), with direct implications for seasonal overlap, both in geographic and ecological spaces. Sedentary species should exhibit a high similarity in their range, but will experience environmental seasonality, resulting in low ecological similarity throughout the year. Alternatively, nomadism can be thought of as a response to highly variable and unpredictable environments in both time and space, which should lead to variable ranges through time, as species track their ecological niche closely (Jonzén et al. 2011). At the end of the continuum, migration is a dramatic response to seasonal environmental change (Dingle and Drake 2007), which leads to two distinct seasonal ranges. As a consequence, occupied space should show high similarity within seasons in both geographic and ecological spaces, but not necessarily between seasons, with migrations potentially resulting in shifts in both spaces. Some species also migrate "in place", e.g., through altitudinal migration (i.e., migratory range shifts along an elevation gradient, Zweifel-Schielly et al. 2009). These species mostly stay within their global range year-round, while locally buffering for seasonally varying conditions to some extent. As a result, it is expected that their occupied space should vary throughout the year, but with intermediate patterns compared to the high expected similarity of residents in geographic space, and of migrants in ecological space.

Finally, it has been recognized that many species express a form of partial migration (a fraction of the population is migratory, while the other part remains resident, Chapman et al. 2011), or facultative migration (individuals that may or may not migrate in a given year, Newton 2012). Partial or facultative migrants pose a conundrum, as the global geographic range and environmental conditions experienced may encompass distinct and contrasted subsets (for residents and migrants) during part of the year or in different years. This plasticity allows facultative migrants to pick and choose when to migrate in response to environmental conditions or internal state, as opposed to presumably more

hard-wired obligate migrants. In the case of partial migration, we thus expect the mix

between the two strategies (resident or migrant) to be reflected in both geographic and

ecological spaces, resulting, at the population level, in similarity being in between that of

resident and migrant species during the migratory season.

Here we describe the seasonality in the distribution of wood storks, which are partially

migratory birds, in both geographic and ecological (specifically climatic) spaces. We built

on the approaches advocated by Fieberg and Kochanny (2005) in the geographic space

only, and by Broennimann et al. (2011) in the ecological space only, to propose a general

framework that allows joint investigation in geographic and ecological spaces. To facilitate

combined interpretation in both spaces, we directly used a simplified 2-D niche space with

two critical climatic components varying throughout the year: temperature and

precipitation. We use home range overlap metrics (Fieberg and Kochanny 2005) to

simultaneously investigate similarity of monthly ranges in geographic space and monthly

niches in climatic space. Because wood storks are partial migrants breeding in South

Florida during winter (dry season, Coulter et al. 1999), we expect both monthly ranges

and niches to be more restricted and highly similar during this season. On the other hand,

as a fraction of the population migrates to northern foraging grounds after breeding, we

then expect both monthly ranges and niches to show low similarity during the rest of the

year.

Methods

Wood Storks in southeastern U.S.

Sixty-one adult wood storks were captured between 2005 and 2011 from 11 sites spread

throughout their southeastern U.S. range in Florida, Georgia, and South Carolina. Wood

storks were equipped with solar-powered Argos/GPS tags set up to take a fix every 1h or

2h. Tagged birds were monitored for an average of 649 ± 563 d from July 2005 to

December 2011, and the tags collected a total of 445,638 GPS locations, which is more

than 7,000 locations per individual on average $(7,306 \pm 6,036)$. We defined the study area

by the coastline to the South (i.e., all land masses), and by the limit of a convex hull

around all GPS locations to the North (see the study area limits in Fig. 1).

We first grouped all GPS data by month across all years to define monthly ranges. To

account for variability in number of locations among individuals, and the fact that

individual wood storks are a random sample of the population, we sampled 5,000 GPS

locations in each month by 1) removing individuals with fewer than 100 locations in a

given month, and 2) sampling 5,000 random individual wood storks (with replacement)

and 1 random location for each of the 5,000 sampled individuals. We thus compiled 60,000

random locations for the population, evenly distributed throughout the year.

In climatic space, the wood stork niche was described by average monthly temperature

and precipitation using climate rasters from the Climate Research Unit (CRU) data set,

available at a resolution of approximately 10 arc-min (New et al. 2002). A previous

analysis (Watling et al. 2012) showed that variables describing monthly climate (e.g., mean

monthly temperature and precipitation) performed equally well as the more ecologically

sensible "bioclimate" variables derived from seasonal relationships between precipitation

and temperature (e.g., maximum temperature of the warmest month, or mean

precipitation of the driest month) to define climatic niches of twelve terrestrial vertebrate

species in the southeastern U.S. Remarkably, the only exception was wood storks, which

can be explained by the global range of the species spanning both hemispheres, hence

effectively preventing monthly variables from be meaningful (Watling et al. 2012).

Focusing on the southeastern U.S. range only circumvents this issue, and actually supports

the use of monthly variables as the simplest approach.

Global layers were clipped to the study area, and temperature and precipitation at

sampled locations defined a 2D-climatic niche, similar to the range defined in the

2D-geographic space (longitude and latitude) by GPS locations. To allow meaningful

comparison between temperature/precipitation and longitude/latitude, the former were

standardized as to be on the same scale (subtraction of the mean followed by division by

the standard deviation for all locations).

131   *A note on ecological niches*

At this stage, it is important to note two things: First of all, as in every species

distribution model, what we observe by means of occurrence data does only reflect realized

niches (Colwell and Rangel 2009, Soberón and Nakamura 2009), i.e., realizations of the

fundamental niche of a species in a given context, restricted by biotic interactions and

movement or dispersal constraints (Soberón 2007). Several studies have tried to

circumvent this limitation (Broennimann and Guisan 2008, Panzacchi et al. 2015), but

they require several populations in different ranges to approach the fundamental niche. In

this manuscript, we thus only consider the realized niche through the year. Niches can

change through time, and such changes observed in the realized niche also include any

potential change of the fundamental niche, although it is impossible to disentangle both

(Guisan et al. 2014).

Second, the seminal work of Hutchinson (1957) provided at the same time a conceptual

framework to study both fundamental and realized niches, and a clear geometrical model

of the niche, defined as a "n-dimensional hypervolume", i.e., a multivariate cloud of points

in the ecological space. This model can be useful to investigate niches, whether we

consider fundamental, potential, or realized niches (Soberón and Nakamura 2009). As a

matter of fact, we rely on this model to evaluate how the niche change through time, and

we extend it to define the concept of monthly niche: just as we can measure "monthly

ranges", defined from all geographical locations in a given month, we define "monthly

niches", as the realization of the fundamental niche in a given month—i.e., a realized niche

at a monthly scale.

154   *Geographic and climatic similarity*

Our framework relies on the observation that the problems of measuring home range

overlap in the geographic space (Fieberg and Kochanny 2005), and measuring niche

overlap in the ecological space reduced to a 2D Cartesian plane (Broennimann et al. 2011)

are actually mathematically identical problems, which reduce to defining what metric of

overlap to use. In the context of home ranges, Fieberg and Kochanny (2005) recommend

Bhattacharyya's affinity (BA), a synthetic measure (i.e., symmetric) appropriate to

quantify the overall similarity of two bivariate probability densities, and ranges from zero

(no overlap) to 100% (complete overlap). In the context of niche overlap, Broennimann

et al. (2011) presented a method to compute overlap between two ecological niches (i.e., in

the ecological space) in three steps: 1) reducing the multidimensional space of the niche to

two dimensions; 2) compute kernel densities to circumvent potential sampling biases; 3)

estimate overlap using Schoener's D. Warren et al. (2008) however recommend the use of

a derivative of Hellinger's distance over Schoener's D. It can actually be shown that their

newly presented metric ($1 - \frac{1}{2}\sqrt{\sum_i \left(\sqrt{p_{X,i}} - \sqrt{p_{Y,i}}\right)^2}$) corresponds mathematically to BA,

which provides fundamental justification for the use of BA as an appropriate metric of

overlap.

We thus computed utilization distributions (UDs), i.e., the bivariate probability density

associated with wood storks in both geographic (defined by longitude and latitude) and

climatic (defined by standardized temperature and precipitation) spaces using the kernel

method (Worton 1989) with standard parameterization (*ad-hoc* method for the smoothing

parameter, which supposes a bivariate normal UD). We then measured overlap in both

geographic and climatic spaces throughout the year with Bhattacharyya's affinity, which

defines seasonal similarity (Cagnacci et al. 2016).

<center>*Marginality*</center>

Different constraints acting in the climatic and geographic spaces render quantitative

comparison of overlap in both spaces difficult. Instead, as the geographic range bounds

what is available to a species, it is possible to better understand the dynamics at play in

climatic space by further investigating monthly selection of climatic conditions by means

of marginality. Marginality, the vector connecting the centroid of environmental conditions

in the geographic area and the centroid of conditions in the realized niche, measures the

difference between available and used conditions (Hirzel et al. 2002, Basille et al. 2008). In

our case, we used marginality to measure the difference between climatic conditions in the

study area and climatic conditions at wood stork GPS locations as temperature and

precipitation change month by month throughout a year. Note here that the study area, in

climatic space, is completely defined throughout the year as the union of the 12 monthly

climate layers.

All analyses were performed in R (R Core Team 2017), with the help of the

"adehabitatLT" package (Calenge 2006) to process GPS locations. All scripts are available

in a GitHub repository (`https://basille.github.io/wost-seasonality/`).

194   RESULTS

There was greater similarity (as measured by overlap) in geographical space than in

climatic space (Fig. 2), despite large geographic variation during the year (see the full

video in Metadata S1). Overall, similarity in the geographic range between any two

months was always greater than 50 %, with minima reached between the months of

January–February and July–September (consistently between 51–55 %). On the other

hand, similarity in the climatic space was much lower, ranging from 0 % between many

months throughout the year, to a maximum of 83 % between July and August. In

particular, similarity from successive months ranges from 6 % to 83 % (mean±SD:

42.0±23.5) in the climatic space vs. 89 % to 99 % (mean±SD: 94.8±3.4) in the geographic

space.

The different patterns of seasonal change in geographic and climatic space confound

interpretation of the overlap in both spaces. For instance, the use of a high threshold in

the geographic space (e.g. 80 %) delineates two very consistent seasons, Summer from

April to October included, and Winter from November to March included (Fig. 2).

However, transitions from each season are very smooth, with high values of overlap (89 %)

in both cases. In other words, the change of the geographic distribution is very progressive

from month to month. Conversely, using thresholds as low as 35 % in the climatic space

still delineates two seasons, Summer from June to September included, and Winter from

November to February included, which are actually subset of the seasons previously

defined in the geographic space (Fig. 2).

Within-season similarity is greater in geographic space than in climatic space: the

lowest similarity between any two months for a given season is 84 % (between April and

September, and between May and September), while similarity drops to 1 % in the

climatic space, even in the subset seasons (between July and October) (Fig. 2). This

highlights both the high similarity of monthly ranges as compared to monthly climatic

niches, but also the higher discriminatory power of the latter, due to larger contrasts.

Finally, monthly marginality in climatic space highlighted a very clear picture of an

annual cycle (Fig. 3): wood storks consistently used areas wetter and slightly warmer than

generally available in their range during the summer (July–August), and consistently used

warmer and drier areas during the winter (November–April). Transitional months between

these two seasons appeared clearly to be May and October, with areas used by wood

storks in these two months warmer and slightly wetter than generally available in their

entire range. In fact, marginality in May and October were very similar, and much more

similar to each other than to any other month of the year (Fig. 3).

229   DISCUSSION

In this study, we investigated seasonal similarity in both geographic space (range) and

ecological space (climatic niche) of the partially migratory wood storks in the southeastern

U.S. Using home range metrics of overlap, we highlighted seasonal consistency in both

spaces: a summer season (wet season: June–September *a minima*) and a winter season

(dry season: November–February *a minima*) have been identified in both geographic and

climatic spaces, punctuated by intermediate seasons (April–May and October). This

resulted in wood storks using drier areas than generally available in the study area during

the winter, but wetter areas during the summer, while using warmer areas year-round.

²³⁸ However, higher overall similarity in the geographic than in the climatic space indicated

²³⁹ the primacy of the former to lead to seasonal space use, with climate alone unable to fully

²⁴⁰ explain temporal patterns.

²⁴¹   The lowest value of geographic overlap occurred between the months of January and

²⁴² July/August, and is still greater than most values of climatic overlap. This can be in part

²⁴³ explained by their migratory strategy: wood storks are partially migratory birds (Coulter

²⁴⁴ et al. 1999), which results in some individual migrating while others stay resident

²⁴⁵ year-round. As a consequence, the overall (monthly) distributions at the population level

²⁴⁶ are not as contrasted as would be for a fully migratory species, especially between

²⁴⁷ migration phases. This result is highly similar to Golden Eagle (*Aquila chrysaetos*), which

²⁴⁸ expressed high overlap (BA $= 87 \pm 9$) between breeding and non-breeding home ranges in

²⁴⁹ western U.S. (Watson et al. 2014), but can also be compared with very low values (< 10)

²⁵⁰ of overlap at the individual level for large migratory vertebrates, with minimum overlap as

²⁵¹ low as 3 (reindeer *Rangifer tarandus*) or 8 (red deer *Cervus elaphus*) documented in

²⁵² Norway (Cagnacci et al. 2016). Despite the very high geographic overlap in our study

²⁵³ system, BA (Fieberg and Kochanny 2005) showed enough fluctuation to define clear

²⁵⁴ seasons in the geographic space as well.

²⁵⁵   On the other hand, similarity in the climatic space was surprisingly low, and defined

²⁵⁶ narrower seasons. While BA delineated clear seasons in the climatic space, their values

²⁵⁷ were consistently lower than the corresponding ones in the geographic space, indicating

²⁵⁸ higher fidelity in space than for climate—or, in other words, that seasonal ranges (breeding

²⁵⁹ and non-breeding) are sought after for different features than their climatic characteristics,

²⁶⁰ suggesting that wood storks could be consider "niche switchers" (Nakazawa et al. 2004).

²⁶¹ Over the long term, birds have been shown to somewhat track climate change, however at

²⁶² a slower pace than the actual climatic shift (Devictor et al. 2008). Altogether this seems to

²⁶³ indicate that birds' fidelity to specific areas could counteract their need to track climatic

²⁶⁴ conditions at both fine (within year, this study) and large (across years, Devictor et al.

2008) temporal scales. However, and contrary to our expectations, climatic similarity was
greater during the wet season than during the dry season, as shown by larger BA values in
between June–September than during the rest of the year. This is potentially a
consequence of the low climatic variability during summer throughout the southeastern
U.S.: it is wet and hot everywhere in the range, and despite wood storks occupying a
larger area, available conditions are narrower than during the winter.

Selection of climatic conditions—in terms of temperature and precipitation—was
remarkably stable within each season (especially during the summer), even though average
climatic conditions still vary greatly from month to month in the study area. This is
another indication of spatial constraint: instead of tracking climate conditions directly,
wood storks settle in specific geographic areas, which directly defines and constrains the
conditions available there in a given season. These areas are in turn associated with
climatic conditions different than the rest of the range, however still varying in each
season. Wood storks are heavily tied to prey availability, especially during the breeding
season, which is directly regulated by local hydrology of the wetland (Kahl 1964, Gawlik
2002). As a result, wood storks are potentially more sensitive to local variations in
weather (especially rain) than average monthly climatic conditions (Borkhataria et al.
2012). As wood storks are constrained to their nests during the breeding season (generally
February–May, Coulter et al. 1999), their winter range corresponds to the driest areas of
the year-round distribution, which are associated with the best foraging opportunities
(Coulter et al. 1987, Herring and Gawlik 2011).

Historically, South Florida was host to large breeding populations of wading birds,
including wood storks, although their number have sharply declined with the drainage of
South Florida's wetlands (Frederick and Ogden 2003). The date of nest initiation has been
delayed by several months (February–March vs. historical November–December initiations)
in response to deteriorating habitat conditions (Frederick et al. 2009). With sufficient
sample size (i.e., enough individuals monitored simultaneously and over several years to

₂₉₂ bring a good generalization power), our framework could be used to highlight fine-scale

₂₉₃ dynamics within single years, and investigate temporal trends (e.g., directly related to the

₂₉₄ timing of migration) over time. Wading birds (white ibises *Eudocimus albus* and wood

₂₉₅ storks) are considered an indicator of Everglades restoration success (Frederick et al.

₂₉₆ 2009), so understanding dynamics for their migration is important to be able to detect

₂₉₇ changes from restoration vs. climatic conditions.

₂₉₈   We used a simplified definition of the niche in two dimensions only—arguably,

₂₉₉ temperature and precipitation are amongst the most critical climatic variables that define

₃₀₀ niche dynamics (Luoto et al. 2007, Bucklin et al. 2015). However, ecological niches, even

₃₀₁ restricted to their climatic form, are highly dimensional constructs, and by extension, the

₃₀₂ ecological space too. Our approach measuring overlap is by no means restricted to a plane.

₃₀₃ While measures of multi-dimensional overlap exist (see for instance an alternative

₃₀₄ approach of a multivariate index of niche overlap based on Tukey depth in Cerdeira et al.

₃₀₅ 2018), it is unclear how the resulting indexes can be compared to a 2-dimensional index of

₃₀₆ geographic overlap, as overlap mathematically decreases with higher dimensionality. A

₃₀₇ first step is thus to reduce the ecological space to two dimensions, for instance focusing on

₃₀₈ the first two axes of a multivariate analysis, which typically account for most of the

₃₀₉ information (see e.g., Dallas et al. 2017), at the expense of a more difficult interpretation.

₃₁₀ In this particular case of temporally-varying niches, the K-select (Calenge et al. 2005)

₃₁₁ provides a natural way to find commonalities between monthly climatic niches, which are

₃₁₂ essentially similar to individual habitat selection with varying availability. Such an

₃₁₃ approach, given relevant data over the entire range, would allow to include other

₃₁₄ temporally-varying factors potentially critical to the dynamics of wood storks, like water

₃₁₅ levels, which are largely managed in South Florida (Gawlik 2002).

₃₁₆   Finally, it can be noted that not all environments that are varying temporally are also

₃₁₇ seasonal—both short- and long-term changes are present in nature. Many arid regions are

₃₁₈ subject to unpredictable rainfall with no temporal correlation within the year, leading to

nomadism (Jonzén et al. 2011). On the other extreme, global changes such as climate

change (Harvey 2016) or land-use change (Song et al. 2018) potentially affect the entire

planet, with direct impact on species ranges (Di Marco and Santini 2015). While we

focused on seasonality in this manuscript, the presented framework is perfectly suited to

handle such cases, as there is no particular assumption as to which movement strategy

animals use, and it is not necessary that the temporal trend follows a yearly cycle as is the

case for seasonal patterns. Given suitable data at the right temporal scale, investigating

any temporal pattern will allow to highlight how the niche evolves in both geographic and

ecological spaces, and identify either short- or long-term shifts.

While seasonality can be addressed in both spaces separately (home range dynamics in

the geographic space, e.g., Lesage et al. 2000, dynamic habitat use in the ecological space,

e.g., Basille et al. 2013), little attention has been devoted to a joint approach. At the

individual level, Peters et al. (2017) present an elaborate approach to 1) identify migratory

tactics in individual roe deer (*Capreolus capreolus*) in partially migrating populations, and

2) describe environmental factors which are different in both groups—and would explain

the different tactics. We aimed to fill the gap at the population or species level by

providing a simple, yet very general approach to explore seasonal variability in geographic

and ecological spaces, which should prove very useful in dynamic ecosystem (La Sorte

et al. 2018). For instance, the correspondence between the geographic and the ecological

space is at the core of species distribution modeling, which infers potential range maps by

considering all suitable areas, and predicts future potential ranges under changing

conditions, such as climate change (Elith and Leathwick 2009). While in a stable,

saturated system, ecological niches provide a direct equivalence to the geographic range

(Soberón and Peterson 2005), it becomes increasingly critical to account for variation

(seasonality, long-term trends) in natural ecosystems(Pearman et al. 2008).

Acknowledgments

We thank David Bucklin for extracting temperature and precipitation data from the
Climate Research Unit dataset. Clément Calenge provided useful feedback on a previous
version of this manuscript. Financial support for this work was provided by the U.S.
Geological Survey's Southeast Region, the U.S. Fish and Wildlife Service, the Army Corps
of Engineers, the National Park Service, the Environmental Protection Agency (STAR
Fellowship to R.B.), and the U.S.D.A. National Institute of Food and Agriculture (Hatch
project 1009101). Handling of Wood Storks was conducted under U.S. Fish and Wildlife
Service Threatened and Endangered Species Permits TE051552, TE801914, and
TE37498B-0; Florida Fish and Wildlife Conservation Commission Special Purpose Permit
WX07646; Georgia Department of Natural Resources Scientific Collecting Permit
29-WJH-05-79; U.S. Geological Survey Bird Banding Permit 22598-E; and University of
Florida IACUC Protocol #E013 and ARC Protocol 002-11BGL. Any use of trade, firm, or
product names is for descriptive purposes only and does not imply endorsement by the
U.S. Government.

Literature Cited

Araújo, M. B., and R. G. Pearson. 2005. Equilibrium of species' distributions with climate.
    Ecography **28**:693–695.

Basille, M., C. Calenge, E. Marboutin, R. Andersen, and J.-M. Gaillard. 2008. Assessing habitat
    selection using multivariate statistics: Some refinements of the Ecological-Niche Factor
    Analysis. Ecological Modelling **211**:233–240.

Basille, M., D. Fortin, C. Dussault, J.-P. Ouellet, and R. Courtois. 2013. Ecologically based
    definition of seasons clarifies predator-prey interactions. Ecography **36**:220–229.

Borkhataria, R. R., P. C. Frederick, R. A. Keller, and J. A. Collazo. 2012. Temporal variation in
    local wetland hydrology influences postdispersal survival of juvenile wood storks (*Mycteria
    americana*). The Auk **129**:517–528.

Broennimann, O., M. C. Fitzpatrick, P. B. Pearman, B. Petitpierre, L. Pellissier, N. G. Yoccoz,

W. Thuiller, M.-J. Fortin, C. Randin, N. E. Zimmermann, and et al. 2011. Measuring

ecological niche overlap from occurrence and spatial environmental data. Global Ecology and

Biogeography **21**:481–497.

Broennimann, O., and A. Guisan. 2008. Predicting current and future biological invasions: both

native and invaded ranges matter. Biology Letters **4**:585–589.

Bucklin, D. N., M. Basille, A. M. Benscoter, L. A. Brandt, F. J. Mazzotti, S. S. Romañach,

C. Speroterra, and J. I. Watling. 2015. Comparing species distribution models constructed

with different subsets of environmental predictors. Diversity and Distributions **21**:23–35.

Cagnacci, F., S. Focardi, A. Ghisla, B. Van Moorter, E. Merrill, E. Gurarie, M. Heurich,

A. Mysterud, J. Linnell, M. Panzacchi, R. May, T. Nygård, C. Rolandsen, and M. Hebblewhite.

2016. How many routes lead to migration? Comparison of methods to assess and characterise

migratory movements. Journal of Animal Ecology **85**:54–68.

Cagnacci, F., S. Focardi, M. Heurich, A. Stache, A. J. M. Hewison, N. Morellet, P. Kjellander,

J. D. C. Linnell, A. Mysterud, M. Neteler, L. Delucchi, F. Ossi, and F. Urbano. 2011. Partial

migration in roe deer: migratory and resident tactics are end points of a behavioural gradient

determined by ecological factors. Oikos **120**:1790–1802.

Calenge, C. 2006. The package "adehabitat" for the R software: A tool for the analysis of space

and habitat use by animals. Ecological Modelling **197**:516–519.

Calenge, C., A. Dufour, and D. Maillard. 2005. K-select analysis: A new method to analyse

habitat selection in radio-tracking studies. Ecological Modelling **186**:143–153.

Candolle, A. d. 1855. Géographie botanique raisonnée. Masson, Paris.

Cerdeira, J. O., T. Monteiro-Henriques, M. J. Martins, P. C. Silva, D. Alagador, A. M. A. Franco,

M. L. Campagnolo, P. Arsénio, F. C. Aguiar, and M. Cabeza. 2018. Revisiting niche

fundamentals with Tukey depth. Methods in Ecology and Evolution **9**:2349–2361.

Chapman, B. B., C. Brönmark, J.-Å. Nilsson, and L.-A. Hansson. 2011. The ecology and

evolution of partial migration. Oikos **120**:1764–1775.

Chen, I.-C., J. K. Hill, R. Ohlemüller, D. B. Roy, and C. D. Thomas. 2011. Rapid range shifts of

species associated with high levels of climate warming. Science **333**:1024–1026.

Colwell, R. K., and T. F. Rangel. 2009. Hutchinson's duality: The once and future niche.

Proceedings of the National Academy of Sciences **106**:19651–19658.

Coulter, M. C., A. L. Bryan, H. E. Mackey, J. R. Jensen, and M. E. Hodgson. 1987. Mapping of

Wood Stork Foraging Habitat with Satellite Data. Colonial Waterbirds **10**:178.

Coulter, M. C., J. A. J. Rodgers, J. C. Ogden, and F. C. Depkin, 1999. Wood Stork (*Mycteria*

americana), version 2.0. Page 409 *in* A. F. Poole and F. B. Gill, editors. The Birds of North

America. Cornell Lab of Ornithology, Ithaca, NY, USA.

Dallas, T., R. R. Decker, and A. Hastings. 2017. Species are not most abundant in the centre of

their geographic range or climatic niche. Ecology Letters **20**:1526–1533.

Darwin, C. 1859. On the Origin of Species. Gramercy.

Devictor, V., R. Julliard, D. Couvet, and F. Jiguet. 2008. Birds are tracking climate warming,

but not fast enough. Proceedings of the Royal Society B: Biological Sciences **275**:2743–2748.

Di Marco, M., and L. Santini. 2015. Human pressures predict species' geographic range size

better than biological traits. Global Change Biology **21**:2169–2178.

Dingle, H., and V. A. Drake. 2007. What Is Migration? BioScience **57**:113–121.

Elith, J., and J. R. Leathwick. 2009. Species Distribution Models: Ecological Explanation and

Prediction Across Space and Time. Annual Review of Ecology and Systematics **40**:677–697.

Fieberg, J., and C. O. Kochanny. 2005. Quantifying home-range overlap: the importance of the

utilization distribution. Journal of Wildlife Management **69**:1346–1359.

Frederick, P., D. E. Gawlik, J. C. Ogden, M. I. Cook, and M. Lusk. 2009. The white ibis and

wood stork as indicators for restoration of the Everglades ecosystem. Ecological Indicators

9:S83–S95.

Frederick, P., and J. Ogden, 2003. Monitoring wetland ecosystems using avian populations:

seventy years of surveys in the Everglades. Pages p. 321–350 *in* D. Bush and J. Trexler,

editors. Monitoring ecosystems: interdisciplinary approaches for evaluating ecoregional

initiatives. Island Press, Washington, D.C.

Fretwell, S. D. 1972. Populations in a seasonal environment. Princeton University Press,

Princeton, NJ, USA.

Gawlik, D. E. 2002. The effects of prey availability on the numerical response of wading birds.

Ecological Monographs **72**:329–346.

Grinnell, J. 1917. The niche-relations of the California Thrasher. Auk **34**:427–433.

Guisan, A., B. Petitpierre, O. Broennimann, C. Daehler, and C. Kueffer. 2014. Unifying niche

shift studies: insights from biological invasions. Trends in Ecology & Evolution **29**:260–269.

Gómez, C., E. A. Tenorio, P. Montoya, and C. D. Cadena. 2016. Niche-tracking migrants and

niche-switching residents: evolution of climatic niches in New World warblers (Parulidae).

Proceedings of the Royal Society B: Biological Sciences **283**:20152458.

Harvey, L. D. D. 2016. Climate and Global Environmental Change. Taylor & Francis Group.

Herring, H. K., and D. E. Gawlik. 2011. Resource Selection Functions for Wood Stork Foraging

Habitat in the Southern Everglades. Waterbirds **34**:133–142.

Hirzel, A. H., J. Hausser, D. Chessel, and N. Perrin. 2002. Ecological-Niche Factor Analysis: How

to compute habitat-suitability maps without absence data? Ecology **83**:2027–2036.

Hutchinson, G. E., 1957. Concluding remarks. Pages 415–427 *in* Cold Spring Harbour

Symposium on Quantitative Biology, volume 22.

Jonzén, N., E. Knudsen, R. D. Holt, and B.-E. Sæther, 2011. Uncertainty and predictability: the

niches of migrants and nomads. Pages 91–109 *in* E. Milner-Gulland, J. M. Fryxell, and A. R.

Sinclair, editors. Animal Migration: A Synthesis. Oxford University Press.

Kahl, M. P. 1964. Food Ecology of the Wood Stork (Mycteria americana) in Florida. Ecological

Monographs **34**:97–117.

Kearney, M., and W. Porter. 2009. Mechanistic niche modelling: combining physiological and

spatial data to predict species' ranges. Ecology Letters **12**:334–350.

La Sorte, F. A., D. Fink, and A. Johnston. 2018. Seasonal associations with novel climates for

North American migratory bird populations. Ecology Letters **21**:845–856.

Laube, I., C. H. Graham, and K. Böhning-Gaese. 2015. Niche availability in space and time:

migration in Sylvia warblers. Journal of Biogeography **42**:1896–1906.

Lesage, L., M. Crête, J. Huot, A. Dumont, and J.-P. Ouellet. 2000. Seasonal home range size and

philopatry in two northern white-tailed deer populations. Canadian Journal of Zoology

78:1930–1940.

Luoto, M., R. Virkkala, and R. K. Heikkinen. 2007. The role of land cover in bioclimatic models

depends on spatial resolution. Global Ecology and Biogeography **16**:34–42.

Martínez–Meyer, E., A. Townsend Peterson, and A. G. Navarro–Sigüenza. 2004. Evolution of

seasonal ecological niches in the Passerina buntings (Aves: Cardinalidae). Proceedings of the

Royal Society of London. Series B: Biological Sciences **271**:1151–1157.

Nakazawa, Y., A. T. Peterson, E. Martínez-Meyer, and A. G. Navarro-Sigüenza. 2004. Seasonal

Niches of Nearctic-Neotropical Migratory Birds: Implications for the Evolution of Migration.

The Auk **121**:610.

Nathan, R., W. M. Getz, E. Revilla, M. Holyoak, R. Kadmon, D. Saltz, and P. E. Smouse. 2008.

A movement ecology paradigm for unifying organismal movement research. Proceedings of the

National academy of Sciences of the United States of America **105**:19052–19059.

New, M., D. Lister, M. Hulme, and I. Makin. 2002. A high-resolution data set of surface climate

over global land areas. Climate Research **21**:1–25.

Newton, I. 2012. Obligate and facultative migration in birds: ecological aspects. Journal of

Ornithology **153**:171–180.

Panzacchi, M., B. Van Moorter, O. Strand, L. E. Loe, and E. Reimers. 2015. Searching for the

fundamental niche using individual-based habitat selection modelling across populations.

Ecography **38**:659–669.

Pearman, P. B., A. Guisan, O. Broennimann, and C. F. Randin. 2008. Niche dynamics in space

and time. Trends in Ecology & Evolution **23**:149–158.

Peters, W., M. Hebblewhite, A. Mysterud, D. Spitz, S. Focardi, F. Urbano, N. Morellet,

M. Heurich, P. Kjellander, J. D. C. Linnell, and F. Cagnacci. 2017. Migration in geographic

and ecological space by a large herbivore. Ecological Monographs **87**:297–320.

R Core Team, 2017. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria. URL `https://www.R-project.org/`.

Soberón, J. 2007. Grinnellian and Eltonian niches and geographic distributions of species.

Ecology Letters **10**:1115–1123.

Soberón, J., and A. T. Peterson. 2005. Interpretation of models of fundamental ecological niches

and species' distributional areas. Biodiversity Informatics **2**:1–10.

Soberón, J., and M. Nakamura. 2009. Niches and distributional areas: Concepts, methods, and

assumptions. Proceedings of the National Academy of Sciences **106**:19644–19650.

Song, X.-P., M. C. Hansen, S. V. Stehman, P. V. Potapov, A. Tyukavina, E. F. Vermote, and

J. R. Townshend. 2018. Global land change from 1982 to 2016. Nature **560**:639–643.

Wallace, A. R. 1876. The geographical distribution of animals: With a study of the relations of

living and extinct faunas as elucidating the past changes of the Earth's surface. Harper and

brothers, New York.

Warren, D. L., R. E. Glor, and M. Turelli. 2008. Environmental niche equivalency versus

conservatism: quantitative approaches to niche evolution. Evolution **62**:2868–2883.

Watling, J. I., S. S. Romañach, D. N. Bucklin, C. Speroterra, L. A. Brandt, L. G. Pearlstine, and

F. J. Mazzotti. 2012. Do bioclimate variables improve performance of climate envelope models?

Ecological Modelling **246**:79–85.

Watson, J. W., A. A. Duff, and R. W. Davies. 2014. Home range and resource selection by

GPS-monitored adult golden eagles in the Columbia Plateau Ecoregion: Implications for wind

power development. The Journal of Wildlife Management **78**:1012–1021.

Worton, B. J. 1989. Kernel methods for estimating the utilization distribution in home-range

studies. Ecology **70**:164–168.

Zweifel-Schielly, B., M. Kreuzer, K. C. Ewald, and W. Suter. 2009. Habitat selection by an

Alpine ungulate: the significance of forage characteristics varies with scale and season.

Ecography **32**:103–113.

505   FIGURES

Figure 1: Wood stork monthly range (left and center) and climatic niche (right) during

January, May, August and October. Range maps show current GPS locations in a given

month in black with kernel contour lines, and all locations from the entire dataset shown

as a reference in gray, on a background of temperature (left, from blue for colder

temperatures to red for warmer temperatures) or precipitations (center, from yellow for

dry conditions to blue for wet conditions). Climatic niches show current conditions used in

a given month (precipitation on the X-axis, and temperature on the Y-axis) in black, in

comparison to conditions available in the study area in a given month in gray. The

marginality arrow defines the difference between average available conditions ("A") and

average used conditions ("U").

Figure 2: Matrices of A) geographic and B) climatic overlap during the year. Each

matrix provides Bhattacharyya's affinity as a percentage (between 0 and 100, higher values

corresponding to darker color), as a measure of overlap between two months i (rows) and j

(columns). The main diagonal, where $i = j$ is grayed out as all values equal 1. Consistent

seasons in a given space are highlighted with a bold line, while seasons identified in the

other space are highlighted with a dotted line for reference.

Figure 3: Seasonal variation in marginality, as measured by the difference between

climatic conditions at wood stork GPS locations vs. the entire study area for each month.

Each arrow represents selection in a given month for both precipitation (X-axis) and

temperature (Y-axis): Wood storks select drier and warmer areas during the winter

(November–April in fuchsia), wetter areas during the summer (June–September in dark

blue), and intermediate conditions in transitional seasons (May and October in orange).