Transcriptomic basis and evolution of ant nurse-larval social regulatory interactions

Michael R. Warner[1*], Alexander S. Mikheyev[2,3], Timothy A. Linksvayer[1]

[1]Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania, USA

[2]Ecology and Evolution Unit, Okinawa Institute of Science and Technology, Onna, Okinawa, Japan

[3]Research School of Biology, Australian National University, Canberra, ACT, Australia

* Corresponding Author

E-mail: michael.ryan.warner@gmail.com

Abstract

Development is often strongly regulated by interactions among close relatives, but the underlying

molecular mechanisms are largely unknown. In eusocial insects, interactions between caregiving worker

nurses and larvae regulate larval development and resultant adult phenotypes. Here, we begin to

characterize the social interactome regulating ant larval development by collecting and sequencing the

transcriptomes of interacting nurses and larvae across time. We find that the majority of nurse and larval

transcriptomes exhibit parallel expression dynamics across larval development. We leverage this

widespread nurse-larva gene co-expression to infer putative social gene regulatory networks acting

between nurses and larvae. Genes with the strongest inferred social effects tend to be peripheral elements

of within-tissue regulatory networks and are often known to encode secreted proteins. This includes

interesting candidates such as the nurse-expressed *giant-lens*, which may influence larval epidermal

growth factor signaling, a pathway known to influence various aspects of insect development. Finally, we

find that genes with the strongest signatures of social regulation tend to experience relaxed selective

constraint and are evolutionarily young. Overall, our study provides a first glimpse into the molecular and

evolutionary features of the social mechanisms that regulate all aspects of social life.

Author Summary

Social interactions are fundamental to all forms of life, from single-celled bacteria to complex plants and

animals. Despite their obvious importance, little is known about the molecular causes and consequences

of social interactions. In this paper, we study the molecular basis of nurse-larva social interactions that

regulate larval development in the pharaoh ant *Monomorium pharaonis*. We infer the effects of social

interactions on gene expression from samples of nurses and larvae collected in the act of interaction

across a developmental time series. Gene expression appears to be closely tied to these interactions, such

that we can identify genes expressed in nurses with putative regulatory effects on larval gene expression.

Genes which we infer to have strong social regulatory effects tend to have weak regulatory effects within

individuals, and highly social genes tend to experience relatively weaker natural selection in comparison

to less social genes. This study represents a novel approach and foundation upon which future studies at

the intersection of genetics, behavior, and evolution can build.

Introduction

Social interactions play a prominent role in the lives of nearly all organisms [1] and strongly affect trait

expression as well as fitness [2–4]. Social interactions in the context of development (e.g. parental care)

often strongly regulate developmental trajectories and resultant adult phenotypes, for example via

transferred compounds such as milk in mammals [5,6], milk-like secretions in arthropods [7,8], and other

forms of nutritional provisioning [9,10]. In many taxa including certain birds, mammals, and insects, care

for offspring and the regulation of offspring development has shifted at least in part from parents to adult

siblings, who perform alloparental care [11]. In eusocial insect societies, sterile nurse workers regulate the

development of their larval siblings by modulating the quantity and quality of nourishment larvae receive

[12–14], as well as through the direct transfer of growth-regulating hormones and proteins [15,16]. At the

same time, larvae influence nurse provisioning behavior via pheromones [17–20] and begging behavior

[21,22].

In general, traits such as caregiving behavior that are defined or influenced by social interactions

are the property of the genomes of multiple interacting social partners [2,14]. This has implications for

both the mechanistic (e.g., molecular) underpinnings of development and trait expression as well as the

genetic basis of trait variation at the population level -- i.e. how allelic variation in the genomes of

interacting social partners affects trait variation [2,14]. Furthermore, because social traits are expressed in

one individual but impact the fitness of other individuals, social behavior and socially-influenced traits

experience distinct forms of selection, including kin selection and social selection [23,24]. Altogether,

these distinct genetic features and patterns of selection are often thought to lead to distinct evolutionary

features, such as rapid evolutionary dynamics in comparison to other traits [25–27]. In eusocial insects,

previous studies show that variation in larval developmental trajectories and ultimate adult phenotypes

(including reproductive caste, body size, etc.) depends on the combination of larval and nurse genotypes

[28–34]. However, the identity of specific genes and molecular pathways that are functionally involved in

the expression of social interactions (e.g., genes underlying nurse and larval traits affecting nurse-larva

interactions) and the patterns of molecular evolution for these genes have remained less well studied

[15,16,35,36].

Transcriptomic studies are often used to identify sets of genes underlying the expression of

particular traits by performing RNA-sequencing on individuals that vary in the expression of such traits.

For example, in social insects, recent studies have compared the transcriptomes of workers that perform

nursing versus foraging tasks [37–39], or nurses feeding larvae of different stages or castes [35,40].

However, given the phenotypic co-regulation known to occur between interacting social partners (here,

nurses and larvae), it is likely that genes expressed in one social partner affect the expression of genes in

the other social partner, and vise-versa, such that interacting social partners are connected by "social"

gene regulatory networks [14,32,41,42]. Thus, identifying the genes important for *social interactions* such

as nurse-larva interactions is only possible by studying the transcriptomic dynamics of both interacting

social partners across a time series of interactions.

To understand the transcriptomic basis of host-symbiont interactions, recent studies have

reconstructed gene regulatory networks acting between hosts and symbionts by collecting and profiling

the transcriptomes of each social partner across a time series of interactions [43–47]. Here, we use

analogous methodology to study transcriptomic signatures of nurse-larva interactions in the pharaoh ant,

Monomorium pharaonis. We sample a developmental time series of larvae as well as the nurses that feed

each larval stage in this series, collecting individuals at the moment of interaction in order to identify

genes involved in the expression of nurse-larva interactions, as well as genes affected by these

interactions (i.e. the full "social interactome" [14]). Pharaoh ant nurses tend to specialize on feeding

young versus old larvae, and nurses feeding young versus old larvae show different transcriptomic

profiles [40]. Larval transcriptomic profiles also change over development [48,49]. Given these results,

we predicted that we would observe concerted changes in broad-scale gene expression in larvae and their

nurses across larval development (Fig 1), reflective of the functional importance of nurse-larva

interactions. Based on our dual RNA-seq data, we infer social gene regulatory networks acting between

nurses and larvae to identify candidate genes predicted to have important social regulatory effects.

Finally, we combine our measures of social regulatory effects with available population genomic data

[48] to characterize the patterns of molecular evolution of genes underlying nurse-larva interactions.

Results

Transcriptome-wide signatures of nurse-larva co-expression across larval development

To elucidate transcriptomic signatures of nurse-larva interactions, we performed RNA-sequencing on

worker-destined larvae across five developmental stages and nurses that fed larvae of each developmental

stage (termed "stage-specific" nurses; see S1 Fig for sampling scheme, S1 Table for list of samples),

building upon a previously published dataset focused on caste development in *M. pharaonis* [48]. We

hypothesized that if genes expressed in larvae regulate the expression of genes in nurse and vice versa, we

would observe correlated expression profiles across larval development in larvae and nurses (Fig 1). As a

biological control, we collected "random nurses" that we observed feeding any stage of larvae in the

colony, and hence would not be expected to show correlated expression dynamics with larvae across the

five larval developmental stages. We also collected reproductive-destined larvae, but unless clearly stated

otherwise, all analyses were performed on only worker-destined larvae. We collected ten individuals of

each sample type to pool into one sample, and we sequenced whole bodies of larvae but separated nurse

heads and abdomens prior to sequencing.

We grouped genes into co-expression profiles or "modules" using an algorithm designed to

characterize gene co-expression dynamics across a short time series [50], known as Short Time-Series

Expression Mining (STEM) [51]. Each module represents a standardized pre-defined expression profile,

consisting of five values that each represent the $\log_2$ fold-change between the given developmental stage

and the initial (L1) stage (see S2 Fig; this results in a total of 81 possible modules). We sorted genes into

the module that most closely represented their expression profile by Pearson correlation. We identified

modules containing a greater than expected number of genes, where we formed null expectations using

permutation tests across developmental stages [50]. We identified such significantly-enriched modules

separately for larvae, stage-specific nurse heads, stage-specific nurse abdomens, random nurse heads, and

random nurse abdomens. We focused on both parallel (i.e. positive regulation or activation) and anti-

parallel (i.e. inhibitory) correlated expression patterns by identifying significantly-enriched modules that

were shared in both larvae and nurses (parallel), as well as significantly-enriched modules for which the

inverse of the module was identified as significantly-enriched in the social partner (anti-parallel).

Larvae and stage-specific nurses shared many significantly-enriched modules (S2 Table). These

shared modules contained the majority of genes expressed in nurses (65% of genes in stage-specific nurse

heads and 76% in abdomens). A substantial proportion of the larval transcriptome was also shared with

stage-specific nurse heads (22% of larval genes) and abdomens (60% of larval genes). Overall there was a

widespread signature of correlated transcriptional patterns between stage-specific nurses and larvae across

larval development (Fig 2A-D). These coordinated dynamics were dominated by parallel associations in

nurse abdomens (possibly reflecting shared metabolic pathways) but anti-parallel associations in nurse

heads (possibly reflecting the social regulation of larval growth). In contrast to stage-specific nurses,

random nurses (our biological control) shared few significantly-enriched modules with larvae (S2 Table),

and modules shared between random nurses and larvae contained significantly fewer genes than modules

shared between stage-specific nurses and larvae (Fig 2E; Wilcoxon test, P < 0.001 for all comparisons).

Specifically, 2% of genes expressed in random nurse heads and 13% of genes expressed in random nurse

abdomens were in modules shared with larvae; 3% of genes expressed in larvae were in modules shared

with random nurse heads, and 2% of genes expressed in larvae were in modules shared with random nurse

abdomens.

Identification of genes putatively involved in social interactions

Given that we observed transcriptome-wide patterns consistent with nurse-larva transcriptional co-

regulation across larval development, we next identified the genes that might be driving these patterns

(see S3 Fig). We performed differential expression analysis to identify genes that varied in larval

expression according to larval developmental stage, as well as genes that varied in nurse expression

according to the developmental stage of larvae they fed. We identified 8125 differentially expressed

genes (DEGs) in larvae (78% of 10446 total genes). We identified 2057 and 1408 DEGs in stage-specific

nurse heads and abdomens, respectively, compared to 599 and 520 DEGs in random nurse heads and

abdomens, respectively. We removed genes differentially expressed in both stage-specific and random

nurses (N = 272 DEGs in heads, N = 140 DEGs in abdomens), which might differ among our colony

replicates due to random colony-specific effects that were not consistently associated with social

regulation of larval development. After this removal, we retained the top 1000 DEGs, sorted by P-value,

for each sample type other than random nurses (larvae, stage-specific nurse heads, stage-specific nurse

abdomens) for social gene regulatory network reconstruction, reasoning that these genes were the most

likely to be involved in the regulation of larval development.

Reconstruction of social gene regulatory networks

To infer putative gene-by-gene social regulatory relationships between nurses and larvae, we

reconstructed gene regulatory networks acting within and between nurses and larvae (S3 Fig). The output

of regulatory network reconstruction is a matrix of connection strengths, which indicate the regulatory

effect (positive or negative) one gene has on another, separated according to the tissue the gene is

expressed in. To identify the most highly connected (i.e. centrally located, upstream) genes of regulatory

networks, we calculated within-tissue connectivity and social connectivity by averaging the strength of

connections across each connection a gene made, differentiating between within-tissue (nurse-nurse or

larva-larva) and social connections (nurse-larva) (Fig 1B). On average, within-tissue connectivity was

higher than social connectivity (Wilcoxon test; P < 0.001 in all tissues), and within-tissue connectivity

was negatively correlated with social connectivity in each tissue (S4 Fig). The top enriched gene ontology

terms based on social connectivity in nurses were entirely dominated by metabolism (S3,S4 Tables; see

also S5 Table for the top 20 genes by nurse social connectivity).

Secreted proteins and social gene regulation

While based on our data it is not possible to distinguish between genes that code for protein products that

are actually exchanged between nurses and larvae versus genes that affect behavior or physiology within

organisms (Fig 1A), proteins that are known to be cellularly secreted represent promising candidates for

the social regulation of larval development [40]. We downloaded the list of proteins that are known to be

cellularly secreted from FlyBase [52] and used a previously-generated orthology map to identify ant

orthologs of secreted proteins [40]. Genes coding for proteins with orthologs that are cellularly secreted in

Drosophila melanogaster had higher social connectivity than genes coding for non-secreted orthologs in

nurse heads (Fig 3A; Wilcoxon test; P = 0.025), though not for nurse abdomens (P = 0.067).

For the most part, we have focused on broad patterns of nurse-larva gene coregulation. In this

paragraph, we will highlight the potential social role of one of the genes with the highest social

connectivity within nurse heads, *giant-lens* (S6 Table; *giant-lens* is the 7[th] highest gene coding for

secreted proteins by social connectivity in nurse heads). *Giant-lens* is an inhibitor of epidermal growth

factor receptor (EGFR) signaling [53], and g*iant-lens* expression in nurse heads was negatively associated

with the expression of the homolog of *eps8*, human EGFR substrate 8 in larvae, most prominently seen in

the spike in nurse *giant-lens* expression accompanied by a drop in larval *eps8* expression at the end of

larval development (Fig 3B). *Giant-lens* was also used in regulatory network reconstruction in larvae (i.e.

it was one of the top 1000 DEGs), and *giant-lens* expression in larvae drops steadily throughout

development (S5 Fig; in contrast to the pattern of *giant-lens* expression in nurse heads). Interestingly,

eps8 does not exhibit a similar peak and drop in expression level in reproductive-destined larvae in

comparison to worker-destined larvae (S6 Fig). It is important to note that these patterns were not seen for

all genes in the EGFR pathway, and the results presented here cannot be taken as concrete evidence of

EGFR regulation via social processes. Nonetheless, the mechanism illustrated here represents a tangible

example of how nurse-larva interactions could function at the molecular level.

Molecular evolution of social gene regulatory networks

To investigate the selective pressures shaping social regulatory networks, we used population genomic

data from 22 resequenced *M. pharaonis* workers, using one sequenced *M. chinense* worker as an outgroup

[48]. Using polymorphism and divergence data, we estimated gene-specific values of selective constraint,

which represents the intensity of purifying selection that genes experience [54]. To identify genes

disproportionately recruited to the core of social regulatory networks, we calculated "sociality index" as

the difference between social connectivity and within-tissue connectivity for each gene. Sociality index

was negatively correlated to selective constraint due to a positive correlation between within-tissue

connectivity and constraint and a negative correlation between social connectivity and constraint (Fig 4A-

C). Additionally, genes differed in sociality index according to their estimated evolutionary age, with

ancient genes exhibiting lower sociality indices than genes in younger age categories (Fig 4D). Finally,

while evolutionary age and evolutionary rate appear to be somewhat empirically confounded [55],

selective constraint and evolutionary age were each independently associated with sociality index, based

on a model including both variables as well as tissue (GLM; LRT; evolutionary age: $\chi^2 = 21.536$, P <

0.001; selective constraint: $\chi^2 = 22.191$, P < 0.001).

Discussion

In organisms with extended offspring care, developmental programs are controlled in part by socially-

acting gene regulatory networks that operate between caregivers and developing offspring [14,42]. In this

study, we sequenced the transcriptomes of ant nurses and larvae as they interacted across larval

development to assess the effects of social interactions on gene expression dynamics. We found that large

sets of genes (i.e. modules) expressed in ant larvae and their caregiving adult nurses show correlated

changes in expression across development (Fig 2). The majority of nurse and larval transcriptomes was

represented in these correlated modules, suggesting that the tight phenotypic co-regulation characterizing

nurse-larva interactions over the course of larval development is also reflected at the molecular level.

To characterize the overall network and evolutionary patterns of genes involved in nurse-larva

interactions, we reverse engineered nurse-larva gene regulatory networks and calculated the "social

connectivity" for each gene, defined as the sum of inferred social regulatory effects on all genes

expressed in social partners. We found that genes with high social connectivity tended to have low

within-individual connectivity (S4 Fig; where within-individual connectivity is defined as the sum of

inferred regulatory effects acting within a given tissue). Nurse-expressed genes with higher sociality

indices (i.e disproportionately higher social connectivity than within-individual connectivity) tended to be

evolutionarily young and rapidly evolving due to relaxed selective constraint (Fig 4). Genes with high

social connectivity were enriched for a number of Gene Ontology (GO) categories associated with

metabolism (S3,S4 Tables), consistent with the idea that molecular pathways associated with metabolism

are involved in the expression of social behavior [56,57]. Previously, many of the proteins found to be

widely present in social insect trophallactic fluid transferred from nurses to larvae were involved in sugar

metabolism (e.g. Glucose Dehydrogenase, several types of sugar processing proteins) [15]. Along the

same lines, many of the genes with with high social connectivity in our study are also annotated with

terms associated with sugar metabolism (S5 Table; e.g. Glycerol-3-phosphate dehydrogenase, Glucose

dehydrogenase FAD quinone, Pyruvate dehydrogenase). Finally, we found that genes encoding for

orthologs of cellularly-secreted proteins in *Drosophila melanogaster* (possibly important for intercellular

signaling) tended to exhibit higher levels of social connectivity than their non-secreted counterparts (Fig

3A).

One gene that stands out in terms of being cellularly secreted and exhibiting a relatively high

social connectivity is *giant-lens*, which inhibits EGFR signaling [53]. EGFR signaling affects eye and

wing development [58] as well as body size in *D. melanogaster* [59], caste development in the honey bee

Apis mellifera [59,60] via the transfer of royalactin from nurses to larvae [59], and worker body size

variation in the ant *Camponotus floridanus* [61]. Further experimental work is necessary to ascertain

whether *giant-lens* is actually orally secreted by nurses and transferred to larvae, but gene expression

dynamics are consistent with the social transfer of *giant-lens* from nurses to larvae, followed by the

inhibition of EGFR signaling at the end of larval development in worker-destined larvae (Fig 3B).

Importantly, this inhibition is not seen in reproductive-destined larvae (S6 Fig). While caste in *M.*

pharaonis is socially regulated in the first larval stage [49], social inhibition of EGFR signaling could

play a role in the regulation of worker body size [61] or secondary caste phenotypes such as wings

[62,63].

In terms of broad evolutionary patterns, our study complements previous results suggesting genes

with worker-biased expression tend to be rapidly evolving, evolutionarily young, and loosely connected

in regulatory networks in comparison to genes with queen-biased expression [38,48,64–66]. Because

pharaoh ant workers are obligately sterile, their traits are shaped indirectly by kin selection, based on how

they affect the reproductive success of fertile relatives (i.e. queens and males) [23,67]. As a result, all-

else-equal, genes associated with worker traits are expected to evolve under relaxed selection relative to

genes associated with queen traits [68,69].

In general, the suite of genic characteristics commonly associated with worker-biased genes

(rapidly evolving, evolutionarily young, loosely connected) are all consistent with relaxed selection acting

on genes associated with workers [49]. Here, we show that within the worker caste, genes that appear to

be functionally involved in the expression of social behavior (i.e. nursing) experience relaxed selective

constraint relative to genes important for within-worker processes. Therefore, the combination of kin

selection as well rapid evolution thought to be characteristic of social traits [25] likely act in concert to

shape the labile evolutionary patterns commonly associated with worker-biased genes. Finally, it has also

been suggested that plastic phenotypes such as caste recruit genes which were evolving under relaxed

selection prior to the evolution of such plastic phenotypes [70–72]. Our results could also be consistent

with this hypothesis, though the population genomic patterns we observe show that relaxed selective

constraint is ongoing.

In this study, we sought to reconstruct regulatory networks acting between nurses and larvae,

beginning with the assumption that nurse gene expression changes as a function of the larval stage fed.

This is more likely to be the case when nurses are specialized on feeding particular larval stages.

According to a previous study, about 50% of feeding events are performed by specialists (though note

specialization is likely a continuous trait, and the 50% figure is the result of a binomial test) [40].

Therefore, we expect our stage-specific nurse samples to comprise about 50% specialists. We also expect

random nurse samples to contain 50% specialist nurses, but, crucially, the specialists should be relatively

evenly divided among larval stages since random nurses were collected regardless of which larval stage

they were observed feeding. Because our stage-specific nurse samples did not consist of 100% specialists,

we expect that the signal of nurse-larva co-expression in our analysis is effectively diluted. In order to

maximize the signal of nurse-larval co-expression dynamics, future studies would ideally focus entirely

on specialists, as well as on tissues such as brains and the specific exocrine glands [73] known to be

important for social behavior and communication. Despite these limitations, we were still able to observe

transcriptomic signatures consistent with the social regulation of larval development.

Conclusions

In this study, we uncovered putative transcriptomic signatures of social regulation and identified distinct

evolutionary features of genes that underlie "social physiology", the communication between individuals

that regulates division of labor within social insect colonies [74,75]. Because we simultaneously collected

nurses and larvae over a time series of interactions, we were able to elucidate the putative molecular

underpinnings of nurse-larval social interactions. This is a promising approach that could be readily

extended to study the molecular underpinnings of all forms of social regulation in social insect colonies,

including regulation of foraging, regulation of reproduction, etc.. Furthermore, by adapting the

methodology presented here (i.e. simultaneous collection over the course of interactions followed by

sequencing), the molecular mechanisms and evolutionary features of genes underlying a diverse array of

social interactions, including courtship behavior, dominance hierarchy formation, and regulation of

biofilm production could all be investigated. Overall, this study provides a foundation upon which future

research can build to elucidate the genetic underpinnings and evolution of interacting phenotypes.

Methods

This study builds on previous work investigating genomic signatures of kin selection in which we

characterized transcriptomic profiles from adult queens and workers, as well as queen- and worker-

destined larvae [48]. While stage-specific nurses were used in the previous analysis, the knowledge of the

developmental stage of larvae they fed was not, as they were simply treated as adult workers. This study

also complements the past dataset with new data from random nurses, which were collected concurrently

with previous samples.

Study Design

To construct experimental colonies, we began by creating a homogenous mixture of approximately fifteen

large source colonies of the ant *Monomorium pharaonis*. From this mixture, we created thirty total

replicate experimental colonies of approximately equal sizes (~300-400 workers, ~300-400 larvae). We

removed queens from ½ the study colonies to promote the production of reproductive-destined larvae.

Reproductive caste is determined in *M. pharaonis* by the end of the first larval instar, likely in the egg

stage [76], and queen presence promotes culling of reproductive-destined L1 larvae. Removing queens

halts this culling, but it is unknown which colony members actually perform such culling [76]. While we

initially expected the presence of queens to impact the gene expression profiles of nurses, we detected 0

DEGs (FDR < 0.1) between queen-present and queen-absent colonies for every sample type. This could

indicate that nurses don't perform culling and that worker developmental trajectories (and nutritional

needs) are not appreciably different between queen-present and queen-absent colonies. Because queen

presence did not substantially impact gene expression, in this study we pooled samples across queen-

present and queen-absent colonies for all analyses.

We pre-assigned colonies to one of five larval developmental stages (labeled L1-L5, where L1

and L2 refer to 1st-instar and 2nd-instar larvae and L3, L4, and L5 refer to small, medium, and large 3rd-

instar larvae [77]). We identified larval stage through a combination of hair morphology and body size.

L1 larvae are nearly hairless, L2 larvae have straight hairs and are twice the length of L1 larvae, and L3-

L5 larvae have dense, branched hairs [78]. We separated 3rd-instar larvae into three separate stages based

on body size [77] because the vast majority of larval growth occurs during these stages. We sampled

individuals (larvae as well as nurses) across larval development time: beginning at the L1 stage, we

sampled colonies assigned to each subsequent stage at intervals of 3-4 days, by the time the youngest

larvae in colonies lacking queens were of the assigned developmental stage (note that in colonies lacking

queens, no new eggs are laid so the age class of the youngest individuals progressively ages). We sampled

each colony once, according to the developmental stage we had previously assigned the colony (e.g. for

colonies that we labeled 'L4', we waited until it was time to sample L4 larvae and nurses and sampled

individuals from that colony at that time). From each colony, we sampled stage-specific nurses and

worker-destined larvae, as well as random nurses from colonies with queens and reproductive-destined

larvae from colonies without queens (starting at the L2 stage, because at L1 caste cannot be distinguished

[76,77]. Reproductive-destined larvae include both males and queens (which cannot be readily

distinguished), though samples are expected to be largely made up of queen-destined individuals given

the typically skewed sex ratio of *M. pharaonis* [48]. See S1 Table for full sample list.

For each time point in each assigned colony, we collected stage-specific nurses, nurses feeding

larvae of the specified developmental stage (L1, L2, etc). Concurrently, we collected random nurses,

nurses we observed feeding a larva of any developmental stage. Rather than paint-marking nurses, we

collected them with forceps as soon as we saw them feeding larvae. We collected random nurses as soon

as we observed them feeding a larva of any developmental stage in the course of visually scanning the

colony. We did not make an attempt to systematically collect nurses from different areas of the nest but

did so haphazardly, such that the distribution of larval stages fed resembled overall colony demography.

Nurses feed L1 and L2 larvae exclusively via trophallaxis (i.e. liquid exchange of fluid), while nurses

feed L3-L5 larvae both via trophallaxis and by placing solid food in larval mouthparts [79]. To get a

representative sample of all types of nurses, we did not distinguish between nurses feeding liquid and

solid food, though all L3-L5 samples contained a mixture of the two. After collecting nurses, we

anaesthetized the colony using carbon dioxide and collected larvae of the specified developmental stage.

All samples were flash-frozen in liquid nitrogen immediately upon sample collection. Note that workers

in *M. pharaonis* are monomorphic [80].

We performed mRNA-sequencing on all samples concurrently using Illumina HiSeq 2000 at

Okinawa Institute of Science and Technology Sequencing Center. Reads were mapped to the NCBI

version 2.0 *M. pharaonis* assembly [38], and we used RSEM [81] to estimate counts per locus and

fragments per kilobase mapped (FPKM) for each locus. For further details on RNA extraction and library

preparation, see [48].

Transcriptome-wide signatures of nurse-larva co-expression across larval development

We used an algorithm that categorizes genes based on their expression dynamics over time into a number

of modules represented by pre-defined expression profiles [50]; see S2 Fig for workflow). To create

modules, we started at 0 and either doubled, halved, or kept the expression level the same at each

subsequent stage, resulting in 81 possible modules ($3*3*3*3 = 81$; four stages after L1). To generate

gene-specific expression profiles based on real results, we calculated the average $\log_2$ fold change in

expression (FPKM) of the gene at each developmental stage compared to the initial expression level at

stage L1. We then assigned each gene to the closest module by Pearson correlation between gene

expression profile and module expression profile [50]. To identify significantly-enriched modules, we

generated null distributions of the number of genes present in each module (based on permutation of

expression over time), and retained modules with a significantly greater than expected number of genes

based on these null distributions (FDR < 0.05 after Bonferroni multiple correction [50]).

Identification of genes putatively involved in social interactions

We used the package EdgeR [82] to construct models including larval developmental stage and replicate

and performed differential expression analysis for each sample type separately. We retained genes

differentially expressed according to a nominal P-value of less than 0.05 (i.e. no false discovery

correction), as the purpose of this step was simply to identify genes that could be involved in interactions

that shape larval development (rather than spurious interactions arising from replicate-specific effects).

See S1 Dataset for a list of all stage-specific nurse and larval differentially expressed genes.

Social regulatory network reconstruction

We normalized expression for each gene using the inverse hyperbolic sine transformation of FPKM. As

input to the algorithm, we constructed "meta-samples" by combining expression data within the same

replicate and time point from nurses and larvae and labeling genes according to the tissue they were

expressed in, along the lines of host-symbiont studies [43,45]. We utilized the program GENIE3 [83,84]

to construct two types of networks: those acting between larvae and nurse heads, and those acting

between larvae and nurse abdomens.

GENIE3 uses a random forest method to reconstruct regulatory connections between genes, in

which a separate random forest model is constructed to predict the expression of each gene, with the

expression of all other genes as predictor variables. The output of GENIE3 is a matrix of pairwise

directional regulatory effects, where the regulatory effect of gene i on gene j is estimated as the feature

importance of the expression of gene i for the random forest model predicting the expression of gene j

(i.e. regulatory effect is how important the expression of gene i is for determining the expression of gene

j). These regulatory effects (or strengths) include both positive and negative as well as non-linear effects,

though these different effect types are not distinguished.

As a side note, a version of GENIE3 that was developed for time series data, dynGENIE3 [85],

does exist. However, we opted to utilize the original GENIE3 algorithm because we reasoned that the

temporal spacing of developmental stages was likely too sparse for regulatory network reconstruction to

incorporate time (note also that the co-expression algorithm we used, STEM, was explicitly designed for

short time series such as ours). While our method therefore does not explicitly incorporate temporal

dynamics, we purposefully biased our results to emphasize larval development over differences between

replicates by only utilizing genes differentially expressed across larval development (or based on larval

stage fed in the case of nurses).

We repeated the entire regulatory reconstruction reconstruction process 1000 times and averaged

pairwise connection strengths across runs, as the algorithm is non-deterministic. To capture the total

effect of each gene on the transcriptome dynamics within tissues, we averaged the regulatory effects each

gene had on all other 999 genes expressed in the same tissue ("within-individual connectivity").

Similarly, to capture the effect each gene had on the transcriptome of social partners, we averaged

regulatory effects each gene had on the 1000 genes expressed in social partners ("social connectivity").

Estimation of selective constraint, and evolutionary rate

Previously, we performed whole-genome resequencing on 22 diploid *M. pharaonis* workers as well as

one diploid *M. chinense* worker to serve as an outgroup [48]. We estimated selective constraint using

MKtest2.0 [86], assuming an equal value of alpha (an estimate of the proportion of nonsynonymous

substitutions fixed by positive selection) across all genes. Selective constraint is the estimate of the

proportion of nonsynonymous mutations that are strongly deleterious and thereby do not contribute to

polymorphism or divergence [86]. Selective constraint is estimated using polymorphism data, so it

represents the strength of purifying selection genes experience within the study population [54].

Phylostratigraphic Analysis

Phylostrata are hierarchical taxonomic categories, reflecting the most inclusive taxonomic grouping for

which an ortholog of the given gene can be found [87–90]. We focused on distinguishing between genes

that were evolutionarily "ancient", present in non-insect animals, versus genes present in only insects,

hymenopterans, or ants [49]. We constructed a database containing 48 hymenopteran available genomes,

10 insect non-hymenopteran genomes, and 10 non-insect animal genomes (S2 Dataset). For outgroup

genomes, we focused on well-annotated genomes which spanned as many insect orders and animal phyla

as possible. Using this database, we estimated evolutionary age of genes based on the most evolutionarily

distant identified BLASTp hit (E-value 10^{-10}).

Gene Set Enrichment Analysis

We performed gene set enrichment analysis based on social connectivity for each gene in each tissue

separately using the R package topGO [91]. We identified enriched gene ontology terms using

Kolmogorov-Smirnov tests (P < 0.05).

General Analyses

We performed all statistical analyses and generated all plots using R version in R version 3.4.0 [92], aided

by the packages "reshape2" [93], "plyr" [94], and "ggplot2" [95].

Data Availability

All raw reads are available at DDBJ bioproject PRJDB3164. All source data for generating figures is

included as S3 Dataset. All scripts and processed data (e.g. expression matrices, evolutionary measures)

are available at https://github.com/warnerm/MonomoriumNurseLarva.

Acknowledgments

We would like to thank the following: Mandy Tin for constructing RNA-sequencing libraries and

performing RNA-sequencing, Luigi Pontieri for images of pharaoh ants, Chao Tong for compiling

hymenopteran genomes for use in phylostratigraphy, and Rohini Singh for comments on the manuscript.

References

1.   Frank SA. All of life is social. Curr Biol. 2007 Aug 21;17(16):R648–50.

2.   Moore AJ, Brodie ED III, Wolf JB. Interacting phenotypes and the evolutionary process: I. Direct

and indirect genetic effects of social interactions. Evolution. 1997;51(5):1352–62.

3.   Wolf JB, Brodie ED III, Cheverud JM, Moore AJ, Wade MJ. Evolutionary consequences of indirect
genetic effects. Trends Ecol Evol. 1998 Feb 1;13(2):64–9.

4.   Bleakley BH, Wolf JB, Moore AJ. The quantitative genetics of social behaviour. Social behaviour:
genes, ecology and evolution. 2010;29–54.

5.   Bijma P. Estimating maternal genetic effects in livestock. J Anim Sci. 2006 Apr;84(4):800–6.

6.   Bouwman AC, Bergsma R, Duijvesteijn N, Bijma P. Maternal and social genetic effects on average
daily gain of piglets from birth until weaning 1. J Anim Sci. 2010;88(9):2883–92.

7.   Stay B, Coop AC. "Milk" secretion for embryogenesis in a viviparous cockroach. Tissue and Cell.
1974;6(4):669–93.

8.   Chen Z, Corlett RT, Jiao X, Liu S-J, Charles-Dominique T, Zhang S, et al. Prolonged milk
provisioning in a jumping spider. Science. 2018 Nov 30;362(6418):1052–5.

9.   Moczek AP. Horn polyphenism in the beetle *Onthophagus taurus*: larval diet quality and plasticity in
parental investment determine adult body size and male horn morphology. Behav Ecol. 1998 Jan
1;9(6):636–41.

10. Lindström J. Early development and fitness in birds and mammals. Trends Ecol Evol. 1999
Sep;14(9):343–8.

11. Wilson Edward O. Sociobiology: the new synthesis. Cambridge, MA: Belknap; 1975.

12. Wheeler DE. Developmental and physiological determinants of caste in social Hymenoptera:
evolutionary implications. Am Nat. 1986;128(1):13–34.

13. Linksvayer TA, Kaftanoglu O, Akyol E, Blatch S, Amdam GV, Page RE. Larval and nurse worker
control of developmental plasticity and the evolution of honey bee queen–worker dimorphism. J
Evol Biol. 2011 Sep 1;24(9):1939–48.

14. Linksvayer TA. The molecular and evolutionary genetic implications of being truly social for the
social insects. In: Zayed A, Kent CF, editors. Advances in Insect Physiology. Academic Press; 2015.
p. 271–92.

15. LeBoeuf AC, Waridel P, Brent CS, Gonçalves AN, Menin L, Ortiz D, et al. Oral transfer of chemical
cues, growth proteins and hormones in social insects. eLife Sciences. 2016 Nov 29;5:e20375.

16. LeBoeuf AC, Cohanim AB, Stoffel C, Brent CS. Molecular evolution of juvenile hormone esterase-
like proteins in a socially exchanged fluid. Sci Rep. 2018 Dec 13;8:17830.

17. Brian MV. Larval recognition by workers of the ant *Myrmica*. Anim Behav. 1975 Nov;23(4):745–
56.

18. Le Conte Y, Sreng L, Poitout SH. Brood pheromone can modulate the feeding behavior of *Apis
mellifera* workers (Hymenoptera: Apidae). J Econ Entomol. 1995 Aug 1;88(4):798–804.

19. Slessor KN, Winston ML, Le Conte Y. Pheromone communication in the honeybee (*Apis mellifera*
L.). J Chem Ecol. 2005 Nov;31(11):2731–45.

20.   Penick CA, Liebig J. A larval "princess pheromone"identifies future ant queens based on their
500   juvenile hormone content. Anim Behav. 2017 June;128:33-40.

501   21.   Creemers B, Billen J, Gobin B. Larval begging behaviour in the ant *Myrmica rubra*. Ethol Ecol
502   Evol. 2003 Jul 1;15(3):261–72.

503   22.   Kaptein N, Billen J, Gobin B. Larval begging for food enhances reproductive options in the ponerine
504   ant *Gnamptogenys striatula*. Anim Behav. 2005;69(2):293–9.

505   23.   Hamilton WD. The genetical evolution of social behaviour. I. J Theor Biol. 1964 Jul;7(1):1–16.

506   24.   Wolf JB, Brodie ED III, Moore AJ. Interacting phenotypes and the evolutionary process. II.
507   Selection resulting from social interactions. Am Nat. 1999;153(3):254–66.

508   25.   West-Eberhard MJ. Sexual selection, social competition, and speciation. Q Rev Biol.
509   1983;58(2):155–83.

510   26.   McGlothlin JW, Moore AJ, Wolf JB, Brodie ED III. Interacting phenotypes and the evolutionary
511   process. III. Social evolution. Evolution. 2010 Sep;64(9):2558–74.

512   27.   Bailey NW, Marie-Orleach L, Moore AJ, Simmons L. Indirect genetic effects in behavioral ecology:
513   does behavior play a special role in evolution? Behav Ecol. 2018 Jan 13;29(1):1–11.

514   28.   Osborne KE, Oldroyd BP. Possible causes of reproductive dominance during emergency queen
515   rearing by honeybees. Anim Behav. 1999 Aug;58(2):267–72.

516   29.   Beekman M, Oldroyd BP. Effects of cross-feeding anarchistic and wild type honey bees: anarchistic
517   workers are not queen-like. Naturwissenschaften. 2003 Apr;90(4):189–92.

518   30.   Linksvayer TA. Direct, maternal, and sibsocial genetic effects on individual and colony traits in an
519   ant. Evolution. 2006 Dec;60(12):2552–61.

520   31.   Linksvayer TA. Ant species differences determined by epistasis between brood and worker genomes.
521   PLoS One. 2007 Oct 3;2(10):e994.

522   32.   Linksvayer TA, Fondrk MK, Page RE Jr. Honeybee social regulatory networks are shaped by
523   colony-level selection. Am Nat. 2009 Mar;173(3):E99–107.

524   33.   Teseo S, Châline N, Jaisson P, Kronauer DJC. Epistasis between adults and larvae underlies caste
525   fate and fitness in a clonal ant. Nat Commun. 2014;5:3363.

526   34.   Villalta I, Blight O, Angulo E, Cerdá X, Boulay R. Early developmental processes limit socially
527   mediated phenotypic plasticity in an ant. Behav Ecol Sociobiol. 2016 Feb 1;70(2):285–91.

528   35.   Vojvodic S, Johnson BR, Harpur BA, Kent CF, Zayed A, Anderson KE, et al. The transcriptomic
529   and evolutionary signature of social interactions regulating honey bee caste development. Ecol Evol.
530   2015;5(21):4795–807.

531   36.   Benowitz KM, McKinney EC, Cunningham CB, Moore AJ. Predictable gene expression related to
532   behavioral variation in parenting. Behav Ecol;ary179;doi:10.1093/beheco/ary179.

533   37.   Manfredini F, Lucas C, Nicolas M, Keller L, Shoemaker D, Grozinger CM. Molecular and social
534   regulation of worker division of labour in fire ants. Mol Ecol. 2014 Feb;23(3):660–72.

535   38.   Mikheyev AS, Linksvayer TA. Genes associated with ant social behavior show distinct
536   transcriptional and evolutionary patterns. Elife. 2015 Jan 26;4:e04775.

537   39.   Kohlmeier P, Alleman AR, Libbrecht R, Foitzik S, Feldmeyer B. Gene expression is more strongly
538   associated with behavioural specialisation than with age or fertility in ant workers. Mol Ecol. 2018
539   Dec 7;doi:10.1111/mec.14971.

540   40.   Walsh JT, Warner MR, Kase A, Cushing BJ, Linksvayer TA. Ant nurse workers exhibit behavioural
541   and transcriptomic signatures of specialization on larval stage. Anim Behav. 2018 Jul 1;141:161–9.

542   41.   Bloch G, Grozinger CM. Social molecular pathways and the evolution of bee societies. Philos Trans
543   R Soc Lond B Biol Sci. 2011 Jul 27;366(1574):2155–70.

544   42.   Linksvayer TA, Fewell JH, Gadau J, Laubichler MD. Developmental evolution in social insects:
545   regulatory networks from genes to societies. J Exp Zool B Mol Dev Evol. 2012 May;318(3):159–69.

546   43.   Tierney L, Linde J, Müller S, Brunke S, Molina JC, Hube B, et al. An interspecies regulatory
547   network inferred from simultaneous RNA-seq of *Candida albicans* invading innate immune cells.
548   Front Microbiol. 2012 Mar 12;3:85.

549   44.   Westermann AJ, Gorski SA, Vogel J. Dual RNA-seq of pathogen and host. Nat Rev Microbiol. 2012
550   Sep;10(9):618–30.

551   45.   Schulze S, Henkel SG, Driesch D, Guthke R, Linde J. Computational prediction of molecular
552   pathogen-host interactions based on dual transcriptome data. Front Microbiol. 2015 Feb 6;6:65.

553   46.   Westermann AJ, Förstner KU, Amman F, Barquist L, Chao Y, Schulte LN, et al. Dual RNA-seq
554   unveils noncoding RNA functions in host–pathogen interactions. Nature. 2016 Jan
555   20;529(7587):496–501.

556   47.   Burns JA, Zhang H, Hill E, Kim E, Kerney R. Transcriptome analysis illuminates the nature of the
557   intracellular interaction in a vertebrate-algal symbiosis. ELife. 2017;6:e22054.

558   48.   Warner MR, Mikheyev AS, Linksvayer TA. Genomic signature of kin selection in an ant with
559   obligately sterile workers. Mol Biol Evol. 2017 Jul 1;34(7):1780–7.

560   49.   Warner MR, Qiu L, Holmes MJ, Mikheyev AS. Convergent eusocial evolution is based on a shared
561   reproductive groundplan plus lineage-specific plastic genes. Preprint. Available from:
562   https://www.biorxiv.org/content/early/2018/10/26/454645.abstract
563

564   50.   Ernst J, Nau GJ, Bar-Joseph Z. Clustering short time series gene expression data. Bioinformatics.
565   2005 Jun;21 Suppl 1:i159–68.

566   51.   Ernst J, Bar-Joseph Z. STEM: a tool for the analysis of short time series gene expression data. BMC
567   Bioinformatics. 2006 Apr 5;7:191.

568   52.   Gramates LS, Marygold SJ, Santos G dos, Urbano J-M, Antonazzo G, Matthews BB, et al. FlyBase
569   at 25: looking to the future. Nucleic Acids Res. 2016;gkw1016.

570   53.   Schweitzer R, Howes R, Smith R, Shilo BZ, Freeman M. Inhibition of *Drosophila* EGF receptor
571   activation by the secreted protein *Argos*. Nature. 1995 Aug 24;376(6542):699–702.

54. Kreitman M, Hudson RR. Inferring the evolutionary histories of the *Adh* and *Adh-dup* loci in
Drosophila melanogaster from patterns of polymorphism and divergence. Genetics. 1991
Mar;127(3):565–82.

55. Moyers BA, Zhang J. Evaluating phylostratigraphic evidence for widespread de novo gene birth in
genome evolution. Mol Biol Evol. 2016 May;33(5):1245–56.

56. Toth AL, Robinson GE. Evo-devo and the evolution of social behavior. Trends Genet. 2007
Jul;23(7):334–41.

57. Toth AL, Varala K, Henshaw MT, Rodriguez-Zas SL, Hudson ME, Robinson GE. Brain
transcriptomic analysis in paper wasps identifies genes associated with behaviour across social insect
lineages. Proc Biol Sci. 2010 Jul 22;277(1691):2139–48.

58. Shilo BZ. Signaling by the *Drosophila* epidermal growth factor receptor pathway during
development. Exp Cell Res. 2003 Mar 10;284(1):140–9.

59. Kamakura M. Royalactin induces queen differentiation in honeybees. Nature. 2011 May
26;473(7348):478–83.

60. Mutti NS, Dolezal AG, Wolschin F, Mutti JS, Gill KS, Amdam GV. IRS and TOR nutrient-signaling
pathways act via juvenile hormone to influence honey bee caste fate. J Exp Biol. 2011 Dec
1;214(23):3977–84.

61. Alvarado S, Rajakumar R, Abouheif E, Szyf M. Epigenetic variation in the *Egfr* gene generates
quantitative variation in a complex trait in ants. Nat Commun. 2015 Mar 11;6:6513.

62. Abouheif E, Wray GA. Evolution of the gene network underlying wing polyphenism in ants.
Science. 2002 Jul 12;297(5579):249–52.

63. Trible W, Kronauer DJC. Caste development and evolution in ants: it's all about size. J Exp Biol.
2017 Jan 1;220(Pt 1):53–62.

64. Johnson BR, Tsutsui ND. Taxonomically restricted genes are associated with the evolution of
sociality in the honey bee. BMC Genomics. 2011 Mar 29;12:164.

65. Harpur BA, Kent CF, Molodtsova D, Lebon JMD, Alqarni AS, Owayss AA, et al. Population
genomics of the honey bee reveals strong signatures of positive selection on worker traits. Proc Natl
Acad Sci U S A. 2014 Feb 18;111(7):2614–9.

66. Morandin C, Tin MMY, Abril S, Gómez C, Pontieri L, Schiøtt M, et al. Comparative transcriptomics
reveals the conserved building blocks involved in parallel evolution of diverse phenotypic traits in
ants. Genome Biol. 2016 Mar 7;17:43.

67. Bourke AFG, Franks NR. Social evolution in ants. Princeton University Press; 1995.

68. Linksvayer TA, Wade MJ. Genes with social effects are expected to harbor more sequence variation
within and between species. Evolution. 2009 Jul;63(7):1685–96.

69. Linksvayer TA, Wade MJ. Theoretical predictions for sociogenomic data: the effects of kin selection
and sex-limited expression on the evolution of social insect genomes. Front Ecol Evol; 2016
June;4:65.

70. Hunt BG, Ometto L, Wurm Y, Shoemaker D, Yi SV, Keller L, et al. Relaxed selection is a precursor
to the evolution of phenotypic plasticity. Proc Natl Acad Sci U S A. 2011 Sep 20;108(38):15936–41.

71. Leichty AR, Pfennig DW, Jones CD, Pfennig KS. Relaxed genetic constraint is ancestral to the
evolution of phenotypic plasticity. Integr Comp Biol. 2012 Jul;52(1):16–30.

72. Helanterä H, Uller T. Neutral and adaptive explanations for an association between caste-biased
gene expression and rate of sequence evolution. Front Genet. 2014 Aug 29;5:297.

73. Eelen D, Børgesen L, Billen J. Functional morphology of the postpharyngeal gland of queens and
workers of the ant *Monomorium pharaonis* (L.). Acta Zool. 2006;87(2):101–11.

74. Seeley TD. The Wisdom of the Hive: the social physiology of honey bee colonies. Harvard
University Press; 2009. 318 p.

75. Johnson BR, Linksvayer TA. Deconstructing the superorganism: social physiology, groundplans,
and sociogenomics. Q Rev Biol. 2010 Mar;85(1):57–79.

76. Warner MR, Lipponen J, Linksvayer TA. Pharaoh ant colonies dynamically regulate reproductive
allocation based on colony demography. Behav Ecol Sociobiol. 2018 Mar 1;72(3):31.

77. Warner MR, Kovaka K, Linksvayer TA. Late-instar ant worker larvae play a prominent role in
colony-level caste regulation. Insectes Soc. 2016 Nov 1;63(4):575–83.

78. Berndt KP, Kremer G. Larvenmorphologie der Pharaoameise *Monomorium pharaonis* (L.)
(Hymenoptera, Formicidae). Zool Anz. 1986;216.

79. Cassill DL, Butler J, Vinson SB, Wheeler DE. Cooperation during prey digestion between workers
and larvae in the ant, *Pheidole spadonia*. Insectes Soc. 2005 Nov 1;52(4):339–43.

80. Frumhoff PC, Ward PS. Individual-level selection, colony-level selection, and the association
between polygyny and worker monomorphism in ants. Am Nat. 1992 Mar 1;139(3):559–90.

81. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a
reference genome. BMC Bioinformatics. 2011 Aug 4;12:323.

82. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression
analysis of digital gene expression data. Bioinformatics. 2010 Jan 1;26(1):139–40.

83. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring regulatory networks from expression
data using tree-based methods. PLoS One. 2010 Sep 28;5(9).
http://dx.doi.org/10.1371/journal.pone.0012776

84. Marbach D, Costello JC, Küffner R, Vega NM, Prill RJ, Camacho DM, et al. Wisdom of crowds for
robust gene network inference. Nat Methods. 2012 Jul 15;9(8):796–804.

85. Huynh-Thu VA, Geurts P. dynGENIE3: dynamical GENIE3 for the inference of gene networks from
time series expression data. Sci Rep. 2018 Feb 21;8(1):3384.

86. Welch JJ. Estimating the genomewide rate of adaptive protein evolution in *Drosophila*. Genetics.
2006 Jun;173(2):821–37.

87. Domazet-Lošo T, Tautz D. A phylogenetically based transcriptome age index mirrors ontogenetic

divergence patterns. Nature. 2010 Dec 9;468(7325):815–8.

88.   Quint M, Drost H-G, Gabel A, Ullrich KK, Bönn M, Grosse I. A transcriptomic hourglass in plant
embryogenesis. Nature. 2012 Oct 4;490(7418):98–101.

89.   Drost H-G, Gabel A, Grosse I, Quint M. Evidence for active maintenance of phylotranscriptomic
hourglass patterns in animal and plant embryogenesis. Mol Biol Evol. 2015 May;32(5):1221–31.

90.   Domazet-Lošo T, Brajković J, Tautz D. A phylostratigraphy approach to uncover the genomic
history of major adaptations in metazoan lineages. Trends Genet. 2007;23(11):533–9.

91.   Alexa A, Rahnenfuhrer J. topGO: enrichment analysis for gene ontology. R package version 2.28.0

92.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2017.

93.   Wickham H. Reshaping data with the reshape package. J Stat Soft. 2007;21(12):1-20.

94.   Wickham H. The split-apply-combine strategy for data analysis. J Stat Soft. 2011;40(1):1-29.

95.   Wickham H. ggplot2: elegant graphics for data analysis. New York: Springer-Verlag; 2016.

Figure Captions

Fig 1. Social regulation of gene expression between ant nurses and larvae.

(A) Cartoon depicting positive gene regulation (i.e. activation) between larvae and nurses, where gene 1 is

expressed in nurses and genes 2 and 3 are expressed in larvae. After the expression of gene 1 increases,

the expression of gene 2 increases as a result of the social interaction of nursing (depicted in [B]). This

can occur if gene 1 itself codes for a protein passed to larvae, if the mRNA transcript is passed directly, or

if gene 1 activates the expression of some other gene in nurses, which in turn is passed as mRNA (or

codes for a protein that is passed) to larvae. Following the increase in expression of gene 2, the expression

of gene 3, which is shown to be activated by gene 2, also increases. While we have depicted a time-lag in

this social regulation of gene expression, the time lags are likely too short to observe in our data, as larvae

were collected every 3-4 days across development. Therefore, correlated transcriptome dynamics over

development (see Fig 2) would reflect mechanisms shown here. (B) Gene regulatory networks act

between and within individuals engaged in social interactions. Blue boxes are genes expressed in larvae,

and red boxes are genes expressed in nurses. Solid lines depict regulatory interactions within tissues

(here, within larvae or within nurses), while dashed lines represent social connections (nurse-larva or vice

versa).

Fig 2. Nurse and larval transcriptomes show strong signatures of gene co-expression across larval

development.

Plots (A-D) depict the expression profiles of individual genes (light lines) as expressed in (A) nurse head,

and (B) nurse abdomens, as well as (C) larvae, shared with nurse heads, and (D) larvae, shared with nurse

abdomens. Dark lines indicate the median expression values of all genes sorted into modules, with pre-

defined expression profiles of modules depicted in plot insets. Colors indicate the pre-defined expression

profile (i.e. module) that genes have been sorted into. Only the five shared modules containing the most

nurse-expressed genes are shown for clarity. Larval expression profiles are divided by the nurse tissue

they are shared with, such that (C) depicts larval gene expression shared with nurse heads (A), while (D)

depicts larval gene expression shared with nurse abdomens (B). Note that nurse heads and larvae shared

inversely-related expression profiles, and that this algorithm does not reveal the direction of regulation as

it is simply correlation-based. (E) Stage-specific nurses have more genes than random nurses in modules

shared with larvae than do random nurses, reflecting more broad-scale co-expression across development.

"Connection type" refers to the tissue that the number of genes was calculated in (i.e. larva $\rightarrow$ nurse head

indicates the number of genes expressed in larvae that are in modules shared with nurse heads), though

directionality is not determined in this algorithm. Error bars indicate 95% confidence intervals derived

from systematic drop-1 jackknifing of nurse samples. N = 10944 genes total.

Fig 3. Genes encoding secreted proteins such as *giant-lens* are important for social gene regulation.

(A) Genes encoding for proteins that are secreted in *Drosophila melanogaster* exhibit higher social

connectivity (i.e. more strongly socially regulate larval expression) in nurse heads than genes encoding

for non-secreted proteins (P-values from Wilcoxon test). (B) The protein *giant-lens* is one of the genes

coding for secreted proteins with the highest social connectivity in nurse heads. Based on our data, *giant-*

lens expressed in stage-specific nurse heads (red) appears to inhibit the expression of the homolog of

human EGFR substrate 8 (*eps8*) expressed in worker-destined larvae (blue). The expression of *giant-lens*

in nurses of a given colony was negatively correlated to the expression of *eps8* in larvae of the same

sampled colony (rho = -0.270, P < 0.001, N = 25 colony/stage pairings after removing missing samples).

Expression at stage i is equal to $\log_2(\text{expression}_i/\text{expression}_1)$, i.e. the ratio of expression at the given

stage to expression at L1.

Fig 4. Highly social genes tend to be less evolutionarily constrained.

Selective constraint, estimated from whole-genome polymorphism data, is (A) positively correlated with

within-tissue connectivity (Spearman correlation; head: rho = 0.122, P < 0.001; abdomen: rho = 0.217, P

< 0.001), but negatively correlated with (B) social connectivity (head: rho = -0.090, P = 0.009; abdomen:

rho = -0.150, P < 0.001) and (C) sociality index (head: rho = -0.132, P < 0.001; abdomen: rho = -0.223, P

< 0.001), where sociality index is the difference between social and within-tissue connectivity per gene.

Each point in (A-C) indicates a single gene, as expressed in nurse heads or abdomens. Lines are trendlines

from linear model. (D) Sociality index differs according to estimated evolutionary age (GLM; LRT; χ^2 =

57.357, P < 0.001), as ancient genes tended to have lower sociality indices than all other categories

(Tukey's post-hoc test; ancient - insect: P < 0.001, ancient - hymenoptera: P < 0.001, ancient - ant: P <

0.001, all other comparisons P > 0.05). Individual points depict average values across nurse heads and

abdomens for all genes within each estimated evolutionary age class, indicated by labels on points. Error

bars depict 95% confidence intervals from bootstrapping. Numbers in parentheses indicate number of

genes in each age class.

S1 Fig. Diagram of sampling scheme.

We collected ten worker-destined larvae, ten stage-specific nurses, and ten random nurses from each

colony (six colonies per time point, where time points represent larval developmental stages L1, L2, etc).

We collected stage-specific nurses when we observed them feeding larvae of the given developmental

stage. We collected random nurses when we observed them feeding any stage of larvae.

S2 Fig. Identification of significantly-enriched modules shared between larvae and nurses.

Inset tables depict pre-defined expression profiles of modules genes can be assigned to. First, we

construct modules using all possible expression profiles (top left bubble). Expression profiles consist of

five values, starting at zero, that indicate the $\log_2$ fold-change in expression from the initial value (at stage

L1). At each subsequent stage, we either double, halve, or keep the expression level the same. This

process is repeated to produce 81 (four stages after L1; $3*3*3*3 = 81$) total modules. Next, for each tissue

separately (here we depict workflow in larvae with yellow bubbles), we calculate individual gene

expression profiles as the $\log_2$ fold-change in expression from the initial value at stage L1 and assign

genes to the closest related module by Pearson correlation. Concurrently, we permute the developmental

stage labels for each gene and assign the stage-permuted genes to modules (repeated 1000 times). From

these stage-permuted results, we calculate the mean number of genes assigned to each module and treat

this number as a null expectation (as each expression profile is not equally likely to occur by chance). We

then identify significantly-enriched modules using a one-way binomial test (with the calculated mean as

the null), with a Bonferroni-corrected false discovery rate of 0.05. This entire process is repeated in a

nurse tissue and significantly-enriched modules are found (blue bubble). Finally, we compare

significantly-enriched modules between larvae and nurses and retain identical and inverse modules as

shared profiles. An example of an inversely related profile is shown in red, where larvae exhibit the

enriched module [0, 0, -1, -2, -3] and nurses exhibit the inverse module, [0, 0, 1, 2, 3].

S3 Fig. Workflow of preliminary differential expression analysis and gene regulatory network

reconstruction.

On the left, we identify putatively socially-acting genes through differential expression analysis. First, for

nurse heads and abdomens separately, we perform differential expression analysis in stage-specific and

random nurses to identify genes differentially expressed according to larval stage fed, using a nominal P-

value of 0.05. We remove genes differentially expressed in random nurses, as these correspond to colony-

specific environmental effects unrelated to social regulation of larval development. Next, we select the

top 1000 differentially expressed genes by P-value in stage-specific nurses (after removing those DE in

random nurses) as well as the top 1000 differentially expressed genes in larvae. From these genes, we

create "meta-samples" by combining gene expression of larvae and stage-specific nurses collected from

the same colony (separately for heads and abdomens), and labeling genes by the tissue they are expressed

in. Using these meta-samples, we perform gene regulatory reconstruction (right) to identify genes

expressed in nurses that regulate larval gene expression, and vise-versa. We repeat gene regulatory

reconstruction 1000 times and average connection strength across runs, as the algorithm is non-

deterministic. The output of gene regulatory reconstruction is a matrix of regulatory connections acting

between genes. From this matrix, we calculate the average connectivity for each gene, separating within-

tissue (larva-larva or nurse head-nurse head) from social (nurse-larva) connections. Genes with high

connectivity are predicted to interact with many genes, i.e. are central to the network. Finally, we

calculate each genes' sociality index as the difference between social connectivity and within-tissue

connectivity.

S4 Fig. Genes highly connected in social regulatory networks are loosely connected in within-tissue

regulatory networks

Connectivity is representative of the number and strength of regulatory connections each gene makes.

Points indicate the average connectivity for a given gene, as measured within-tissue (x-axis; i.e. larva-

larva or nurse-nurse) or socially (y-axis; i.e. larva-nurse). Points are colored by tissue the connectivity is

measured in (e.g., dark blue indicates genes expressed in larvae, with connectivity measured in networks

constructed with nurse abdomens). Spearman rho = -0.166, -0.374, -0.276, -0.342 for the four tissues as

ordered in legend; P < 0.001 in all cases.

S5 Fig. Expression of *giant-lens* in nurse heads and worker-destined larvae. Expression at stage *i* is

equal to $\log_2(\text{expression}_i/\text{expression}_1)$, i.e. the ratio of expression at the given stage to expression at the

initial (L1) stage. **: $P < 0.01$, ns: $P > 0.05$ (Wilcoxon test at each stage).

S6 Fig. Expression of *eps8* (epidermal growth factor receptor substrate 8) in worker-destined and

reproductive-destined larvae. Expression at stage *i* is equal to $\log_2(\text{expression}_i/\text{expression}_1)$, i.e. the ratio

of expression at the given stage to expression at the initial (L1) stage. Expression of *eps8* changed

differently over time in worker-destined versus reproductive-destined larvae (linear model with

developmental stage treated as an ordinal variable; LRT; $\chi^2 = 12.574$, $P = 0.014$ for the interaction term

stage*caste).

S1 Table. Description of samples included in study. Worker-destined larvae are indicated by larva (W),

and reproductive-destined larvae are indicated by larva (R). Larval caste cannot be distinguished at the L1

stage, so L1 larvae are labeled larva (W/R). For network reconstruction, "meta" samples were used as

input for network reconstruction, in which genes were labeled by sample type and grouped such that each

gene contained a measurement of expression in worker-destined larvae, nurse heads, and nurse abdomens.

After sample collection and RNA extraction, some samples exhibited clearly degraded RNA according to

an Agilent Bioanalyzer assay. Removing these samples caused sampling to be uneven, so we used the

minimum number of samples contained across tissues at a given stage for stage-specific nurse heads and

abdomens, and randomly dropped excess samples. Overall, 25 "aggregate" samples were used as input for

gene regulatory network reconstruction.

S2 Table. Number of nurse significantly-enriched modules shared with larvae.

Significantly-enriched modules are defined as modules with a statistically significant number of genes

assigned, as determined by a permutation test (FDR < 0.05). Left column is the total number of

significant modules for each tissue, while the second and third columns indicate number shared with

larvae (out of 24 larval significantly-enriched modules). The last column indicates the total number of

genes in these shared modules.

S3 Table. Nurse head social connectivity GO terms based on GSEA of social connectivity.

P-value (unadjusted) is from Kolmogorov Smirnov (K-S) test. Enriched terms have higher than expected

social connectivity in nurse heads.

S4 Table. Nurse abdomen GO terms based on GSEA of social connectivity.

P-value (unadjusted) is from Kolmogorov Smirnov (K-S) test. Enriched terms have higher than expected

social connectivity in nurse abdomens.

S5 Table. Top 20 genes by social connectivity in nurses.

SwissProt ID is listed from automated annotation where a term was found.

S6 Table. SwissProt annotations for the top genes coding for secreted proteins, sorted by social

connectivity.

Only genes with SwissProt annotations are included. All genes listed encode for secreted proteins in *D.*

melanogaster.

S1 Dataset. Complete list of all differentially expressed genes.

Each gene can be differentially expressed in three tissues: worker larva, nurse head, or nurse abdomen. P-

values are listed for each tissue. The top 1000 differentially expressed genes (by P-value) were used for

regulatory network reconstruction. Social connectivity is the sum of all regulatory interactions in the

direction specified by "estimated regulatory direction".

S2 Dataset. List of species used for phylostratigraphy.

Each species listed, with their NCBI taxonomic ID, was used in the construction of the phylostratigraphic

database to estimate evolutionary ages of genes.

S3 Dataset. Data files used to construct all figures.

All data are organized in text files, with the relevant figure listed in the title.

L1
larvae x10
stage-specific nurses x10
? random nurses x10

L2
larvae x10
stage-specific nurses x10
? random nurses x10

L3
larvae x10
stage-specific nurses x10
? random nurses x10

L4
larvae x10
stage-specific nurses x10
? random nurses x10

L5
larvae x10
stage-specific nurses x10
? random nurses x10

Differential Expression Analysis

Gene Regulatory Network Reconstruction

development stage	sample type	number of samples
L1	larva (W/R)	5
	stage-specific nurse head	5
	stage-specific nurse abdomen	6
	random nurse head	3
	random nurse abdomen	2
L2	larva (W)	6
	larva (R)	3
	stage-specific nurse head	6
	stage-specific nurse abdomen	5
	random nurse head	3
	random nurse abdomen	3
L3	larva (W)	6
	larva (R)	3
	stage-specific nurse head	5
	stage-specific nurse abdomen	6
	random nurse head	3
	random nurse abdomen	3
L4	larva (W)	6
	larva (R)	3
	stage-specific nurse head	4
	stage-specific nurse abdomen	5
	random nurse head	2
	random nurse abdomen	2
L5	larva (W)	6
	larva (R)	3
	stage-specific nurse head	5
	stage-specific nurse abdomen	5
	random nurse head	3
	random nurse abdomen	3

	number of significantly-enriched modules	modules positively shared with larvae	modules negatively shared with larvae	number of genes in shared modules
stage-specific nurse head	9	0	5	6838
random nurse head	9	0	2	209
stage-specific nurse abdomen	21	13	4	7943
random nurse abdomen	10	0	1	1400

GO.ID	Term	Annotated	Significant	Expected	P-value
GO:0044710	single-organism metabolic process	81	14	7.82	0.00011
GO:0055114	oxidation-reduction process	45	11	4.35	0.00230
GO:0019637	organophosphate metabolic process	18	1	1.74	0.00506
GO:0044711	single-organism biosynthetic process	18	0	1.74	0.00506
GO:0006629	lipid metabolic process	12	2	1.16	0.00565
GO:0009117	nucleotide metabolic process	14	0	1.35	0.00755
GO:0006812	cation transport	18	2	1.74	0.00781
GO:0015672	monovalent inorganic cation transport	11	1	1.06	0.00898
GO:0090407	organophosphate biosynthetic process	11	0	1.06	0.01033
GO:0055086	nucleobase-containing small molecule metabolic process	16	0	1.55	0.01156

GO.ID	Term	Annotated	Significant	Expected	P-value
GO:0055114	oxidation-reduction process	52	7	4.96	0.022
GO:0008152	metabolic process	246	25	23.46	0.033

Gene	estimated regulatory direction	social connectivity	SwissProt ID
LOC105833299	nurse abdomen → larva	0.464	5'-nucleotidase domain-containing protein 1
LOC105838526	nurse head → larva	0.456	1 5-anhydro-D-fructose reductase
LOC105828583	nurse head → larva	0.433	Glycerol-3-phosphate dehydrogenase NAD( ) cytoplasmic
LOC105835488	nurse head → larva	0.433	Myosin regulatory light chain 2
LOC105837148	nurse head → larva	0.419	Short-chain dehydrogenase/reductase family 16C member 6
LOC105828656	nurse head → larva	0.413	N-acetyltransferase 6
LOC105837075	nurse head → larva	0.409	Hemolymph lipopolysaccharide-binding protein
LOC105834708	nurse head → larva	0.401	Glucose dehydrogenase FAD quinone
LOC105836140	nurse head → larva	0.399	Ankyrin-2
LOC105832040	nurse abdomen → larva	0.393	-
LOC105834910	nurse abdomen → larva	0.376	Pyruvate dehydrogenase E1 component subunit beta mitochondrial
LOC105835497	nurse abdomen → larva	0.373	Phenoloxidase 2
LOC105836690	nurse abdomen → larva	0.366	-
LOC105836193	nurse head → larva	0.365	-
LOC105833826	nurse head → larva	0.360	Probable tubulin polyglutamylase TTLL2
LOC105833938	nurse head → larva	0.357	-
LOC105836189	nurse abdomen → larva	0.348	Venom metalloproteinase 3
LOC105829511	nurse head → larva	0.341	La-related protein 1
LOC105840715	nurse abdomen → larva	0.336	-
LOC105833984	nurse abdomen → larva	0.331	Procollagen-lysine 2-oxoglutarate 5-dioxygenase 3

nurse head	nurse abdomen
Basement membrane-specific heparan sulfate proteoglycan core protein	Procollagen-lysine 2-oxoglutarate 5-dioxygenase 3
Collagen alpha-1(IV) chain	Collagen alpha-1(IV) chain
Spondin-1	Tubulointerstitial nephritis antigen-like
Serine proteinase stubble	Papilin
Angiotensin-converting enzyme (Fragment)	Semaphorin-2A
Thrombospondin-4	Transferrin
Protein giant-lens	Basement membrane-specific heparan sulfate proteoglycan core protein
Protein lev-9	Protein NPC2 homolog
Papilin	Testican-2
Glypican-6	Lysozyme