Calorie restriction intervention induces enterotype-associated BMI loss in nonobese individuals

Hua Zou[1,2, 6*], Dan Wang[2, 6*], Huahui Ren[2,3,5*], Peishan Chen[2,3], Chao Fang[2,3,5], Zhun Shi[2,3], Pengfan Zhang[1,2], Jian Wang[2,4], Huanming Yang[2,4], Kaiye Cai[2,3,6#], Huanzi Zhong[2,3,5 #]

[1] BGI Education Center, University of Chinese Academy of Sciences, Shenzhen 518083, China

[2] BGI-Shenzhen, Shenzhen 518083, China

[3] China National Genebank, BGI-Shenzhen, Shenzhen 518120, China

[4] James D. Watson Institute of Genome Sciences, Hangzhou 310058, China

[5] Laboratory of Genomics and Molecular Biomedicine, Department of Biology, University of Copenhagen, 2100 Copenhagen Ø, Denmark

[6] Shenzhen Key Laboratory of Human commensal microorganisms and Health Research

[*]These authors contributed equally to this work

[#]**Corresponding authors:**

Kaiye Cai. E-mail: caikaiye@genomics.cn;

Huanzi Zhong. E-mail: zhonghuanzi@genomics.cn

Abstract

Calorie restriction (CR), which has the potential effect to weight loss and blood amino acids, has been demonstrated to associate with gut microbiota in human, especially in obese individuals. However, studies for simultaneously evaluating enterotype-dependent impacts of CR on the gut microbiota and blood amino acids in nonobese individuals are still limited.

Here, 41 nonobese individuals received a 3-week CR diet with approximately 50% fewer calories than normal diet. We measured their BMI and blood amino acid concentration, along with the gut microbiota before and after the intervention. In this trial, 28 Enterotype *Bacteroides* (ETB) subjects and 13 Enterotype *Prevotella* (ETP) subjects were identified before the intervention. Short-term CR dietary intervention decreased the body mass index (BMI) in most subjects but varied in subjects with different enterotypes. ETP subjects exhibited significantly higher BMI loss ratio than the ETB subjects. CR additionally induced substantial enterotype-independent changes in blood amino acids, but only minor changes in gut microbial composition.

We further built a prediction model based on baseline relative abundances of 7 gut microbial species showing high performance in predicting CR-associated BMI loss ratio. Among them, the relative abundance of ETB-enriched *Clostridium bolteae* and *C. ramosum* were negatively correlated with BMI loss ratio while the relative abundance of *Dorea longicatena* which was slightly enriched in ETP subjects, was positively correlated with BMI loss ratio.

Together, our work points out that the individual variation of BMI loss after CR could be partially correlated with different microbial composition and highlights the potential application for microbiome stratification in personalized nutrition intervention.

Keywords: Calorie restriction, gut microbiota, enterotype, amino acids, body mass index.

Introduction

Calorie restriction, a nutritional intervention of reduced energy intake, has been demonstrated to reduce body weight and modulate serum metabolic in human population in many studies [1, 2]. Observational studies with long-term CR found that CR resulted in weight loss and decreasing chronic disease risk factors in nonobese persons [3-6]. For instance, in a 2-year nonobese human trail, CR significantly decreased body weight and

cardiometabolic risk factors, such as triglycerides and total cholesterol [4]. Another study [5] also provided evidence for reduced BMI and improved mood and sleep duration with CR in healthy nonobese adults. However, studies about potential negative impacts of CR on human health remain limited.

CR intervention affecting gut microbiota and blood amino acids has been explored in many studies, particularly among overweight and obese individuals [7-10]. A 4-week CR intervention improved gut barrier integrity, reduced systemic inflammation on gut microbial diversity and BMI loss in obese women, suggesting a potential association among gut microbiota, CR and BMI [8]. Increasing studies have demonstrated that the gut microbiota has been implicated in modulating host energy and nutrient metabolism [11] and regulating the production of gastrointestinal hormones and host appetite [12, 13]. Enterotype, a concept for stratifying individuals based on the gut microbiota, was first described in 2011 and was closely linked to long-term dietary patterns [14, 15]. Plenty of studies have reported that individuals have shown microbial-dependent (enterotypes, *Bacteroides* to *Prevotella* ratio) metabolic responses to the same intervention, including changes of BMI and glycemic indices [12, 15-20]. As for the association between CR and blood amino acids, Biolo *et.al* reported changes in blood amino acids of 9 healthy nonobese men after 2-week CR diet [10]. Despite the plenty of studies on CR as we mentioned above, to our knowledge, there are no studies for simultaneously evaluating impacts of CR on BMI, gut microbiota and blood amino acids in nonobese individuals in a single study.

Here, we conducted a 3-week CR intervention on 41 nonobese subjects with two enterotypes, including enterotype *Bacteroides* (ETB) and enterotype *Prevotella* (ETP). We found that ETP subjects had higher BMI loss ratio than ETB subjects after CR intervention. On the other hand, we found that there were no obvious changes in the gut microbiota composition of two enterotype groups after the CR intervention and the two enterotype groups showed consistent changes in levels of multiple blood amino acids. We further demonstrated that baseline relative abundances of 7 gut microbial species including 3 enterotype-specific species (*C. bolteae*, *C. ramosum*, and *D. longicatena*), rather than baseline BMI and blood amino acids levels, could well predict CR-associated

84 BMI loss ratio. Additionally, we found significantly increased levels of 3-methylhistidine

85 after the intervention, a potential biomarker for muscle protein turnover, suggesting

86 possible muscle protein loss under a CR diet with lower than normal protein availability.

Materials and methods

Volunteer Recruitment

Volunteer-wanted posters were propagated at the China National Gene Bank in Shenzhen

from March to April 2017. A non-obese healthy volunteer was considered if his/her BMI

was less than 28 kg/m^2 [21]. In addition, recruited volunteers should meet all the

following criteria: 1) without antibiotics in the recent 2 months; 2) without prebiotic or

probiotic supplements in the recent 2 months; 3) without hypertension, diabetes mellitus,

gastrointestinal disease and other severe auto-immune disease; 4) regular eating and

lifestyle patterns; 5) no international travel in the recent 3 month. 50 individuals met all

the criteria and were recruited in this study, and 41 individuals (24 females and 17 males

aged 30 ± 6 years old) completed the whole intervention (**Table 1**). The study was

approved by the institutional review board on bioethics and biosafety of BGI-Shenzhen,

Shenzhen (NO. BGI-IRB 17020). All participates were fully informed of the design and

purpose of this intervention study and signed a written informed consent letter.

Study design and low-calorie food preparation

The study included a one-week run-in period (baseline) and a three-week CR dietary

intervention period. During the first week (run-in period), all healthy volunteers

consumed their usual diet and were encouraged to avoid yoghurt, high-fat foods and

alcohol. The CR diet was comprised of ~50% calories of a normal-calorie diet (female,

1000kcal/day; male, 1200kcal/day). It was designed with carbohydrate, fat and protein as

approximately 55%, 30% and 15% of the total energy intake respectively, according to

the Dietary Guidelines for Chinese Residents (2016) and nutritionally balanced [22] and a

recent large nutritional study in China [23]. Common foods in low-calorie diets such as

rice, vegetables, eggs, pork and beef were prepared in our study center to control

experimental variables introduced by different foods and calorie estimation errors.

Traditional Chinese cooking style - boiled, stir-fried and stewed, were applied for our

foods. For each meal, digital scales were used to measure the nutritional and caloric

values of different foods and total meal for male and female respectively.

BMI, blood and fecal samples of each volunteer were collected at our study center at

baseline and after the 3-week CR intervention (**Figure 1**). To avoid intra-individual

variations, BMI was collected multiple times of each volunteer during the last week of

the CR intervention, and the averaged BMI value was used as his/her after-intervention

BMI (**Figure 1**).

Fecal sampling and shotgun metagenomic sequencing

Fecal samples were self-collected and then transferred to the laboratory on dry ice and

kept frozen at -80°C before and after the CR intervention. Fecal DNA was extracted

following a manual protocol as described previously [24]. The DNA concentration was

estimated by Qubit (Invitrogen). Library construction and shotgun metagenomics

sequencing were performed on qualified DNA samples based on the BGISEQ-500

protocol in the single-end 100bp mode [25].

Metagenomic analysis

Raw reads of BGISEQ-500 with SE100 mode were trimmed by an overall accuracy (OA)

control strategy to control quality [25]. After trimming, averagely, 98.15% of the raw

reads still remained as high-quality reads (**Supplemental Table 1**). By using SOAP2.22

software, the high-quality reads were aligned to hg19 to remove reads from host DNA

(identity $\geq$ 0.9). The retained clean reads were aligned to the integrated non-redundant

gene catalog (IGC) using SOAP2.22 [26] and the average mapping rate and unique

mapping rate were 80.18% and 65.76% respectively (identity $\geq$ 0.95, **Supplemental**

Table 1). The relative abundance profiles of genes, genera, species and Kyoto

Encyclopedia of Genes and Genomes orthologous groups (KEGG, KOs) of each sample

were calculated by summing the relative abundances of their assigned IGC genes [26].

For   enterotyping,   we   applied   a   recently   published   universal   classifier

(http://enterotypes.org/),   which   circumvents   major   shortcomings   in   enterotyping

methodology such as lack of standard and small sample size [27].

At baseline, 41 individuals were clustered into two groups: 28 ETB (*Bacteroides* enriched) and 13 ETP (*Prevotella* enriched) individuals 87.8% (36 of 41) individuals were clustered to the same enterotype after the 3-week CR intervention. Detailed enterotype information for each individual is provided in **Supplemental Table 2**.

Genus or species with an occurrence rate > 80% and a median relative abundance > 1e-6 in all samples were defined as common genus or species and used for further intra- and inter-enterotype comparison analyses **Supplemental Table 3-4**.

Differentially enriched KEGG pathways were identified between enterotypes and between different time points, based on the distribution of Z-scores of all KOs belonging to a given pathway [28, 29]. A reporter score $|Z| > 1.96$ (95% confidence interval according to a normal distribution) was used as a detection threshold for significantly differentiating pathways.

Alpha diversity of each individual was calculated on the gene and species relative abundance profiles using the Shannon index. Beta-diversity on the gene and species relative abundance profiles was calculated using the Bray-Curtis distance.

Blood sample collection and amino acids profiling

Fasting blood samples were collected before and after the intervention for amino acids analysis. These blood samples were then centrifuged, and serum samples were collected and stored at -80 □. The concentrations of 31 amino acids and derivatives in the serum samples were then measured via ultra-high pressure liquid chromatography (UHPLC) coupled to an AB Sciex Qtrap 5500 mass spectrometry (AB Sciex, US) as described previously [30].

Statistical methods

Pearson's chi-square test

Pearson's chi-square test was performed to assess sex distribution between individuals of two enterotypes.

Wilcoxon rank-sum test & Wilcoxon Signed-rank test

Wilcoxon rank-sum test was used to detect the significant differences on phenotypes, the

concentrations of blood amino acids and the relative abundances of genera and species

between enterotypes.

Wilcoxon signed-rank test was used to detect the significant differences on phenotypes,

the concentrations of blood amino acids and the relative abundances of genera and

species in paired samples before and after the intervention.

BMI loss ratio

BMI loss ratio of a given individual was calculated using the following equation:

$$BMI\ loss\ ratio = \frac{Before_{BMI} - After_{BMI}}{Before_{BMI}} * 100\%$$

Where $Before_{BMI}$ and $After_{BMI}$ are the BMI value of the same individual before and

after the CR intervention, respectively.

PERMANOVA

The association between enterotypes and the overall blood amino acid profile at baseline

was assessed using permutational multivariate analysis of variance (PERMANOVA) with

9,999 permutations on enterotypes (R *vegan* package, *adonis* function, method="bray").

PCoA

Principal coordinate analysis (PCoA) of fecal samples was performed based on the

relative abundances of common species using Bray-Curtis distance (R *ape* package).

PCA

Principal component analysis (PCA) was performed based on the blood amino acid

profiles to visual overall amino acid composition between enterotypes, and that before

and after the intervention.

Feature selection of gut species and serum amino acids

To investigate whether we could predict BMI loss ratio using omics features, we

performed a Lasso (Least absolute shrinkage and selection operator) regression analysis

between baseline relative abundances of gut common species and the concentrations of

blood amino acids (independent variables), and BMI loss ratio (dependent variables).

We first normalized values of both independent and dependent variables (R, *scale*

function). We then used the R function cv.glmnet to choose the most appropriate value

for λ in the Lasso model (R *glmnet* package, alpha =1, family="gaussian", nfolds=10,

alpha=1, nlambda=100). Here, λ is the tuning parameter (λ > 0) which controls the

strength of the shrinkage of the variables [31]. We then applied the Lasso feature

selection process by shrinking the Lasso regression coefficients of non-informative

variables to zero and selecting the variables of non-zero coefficients. Seven gut microbial

species including *Clostridium bolteae*, *Clostridium ramosum*, *Dorea longicatena*,

Coprococcus eutactus, *Streptococcus mitis, Clostridiales genomosp. BVAB3* and

Mobiluncus curtisii were selected at this step.

Performance estimation of BMI loss ratio prediction model

To reduce overfitting with a limited sample size (n=41), we applied leave-one-out cross

validation (LOOCV) to estimate the prediction performance of BMI loss ratio using a

generalized linear model (GLM) of the seven selected features (creatFolds function in R

caret package and the glm function in R *base* package). Likewise, we also used baseline

BMI values for LOOCV to estimate its prediction performance for CR-associated BMI

loss ratio. Spearman's rho values were calculated between actual BMI loss ratios and the

predicted values.

Significance cutoff

P-value adjustment was applied for multiple hypothesis testing on the concentrations of

blood amino acids, the relative abundances of gut microbial genera and species used

Benjamini-Hochberg (BH) method. BH-adjusted P value less than 0.05 was considered

as statistical significance. The significance for α-diversity, β-diversity and phenotypes

(age, female to male ratio, BMI and BMI loss ratio) was set at $p < 0.05$.

All statistical analyses were conducted using R (version 3.5.0).

Result

BMI loss of ETB and ETP subjects responded differentially to CR Intervention

Based on the baseline genera abundance profile, individuals can be robustly clustered into two enterotypes: enterotype *Bacteroides* (ETB, n = 28) and enterotype *Prevotella* (ETP, n = 13) (See Materials and Methods, **Figure 2A**).

Comparisons of the baseline phenotypes between two enterotype groups revealed that all collected phenotypes, including sex distribution (female/male ratio), age, BMI and weight, showed no significant differences between ETB and ETP subjects **(Table 1, Figure 2B)**. By contrast, the baseline compositional and functional characteristics of the gut microbiota showed marked differences between two groups, in agreement with the previous studies [12, 15, 32]. For instance, genera *Prevotella* and *Paraprevotella* and four species from the two genera were significantly enriched in ETP subjects whereas 19 common species, from such as genera *Bacteroides* and *Clostridium* including *C. bolteae* and *C. ramosum*, were significantly enriched in ETB subjects (Wilcoxon rank-sum test, BH-adjusted P < 0.05; **Supplemental Figure 1A, Supplemental Table 5-6**). At the functional level, multiple pathways in carbohydrate metabolism and membrane transport were highly enriched in ETB subjects whereas four pathways including biosynthesis of phenylalanine, tyrosine and tryptophan (map00400), peptidoglycan (map00550), terpenoid backbone (map00900); and methane metabolism (map00680) were enriched in ETP subjects (|Z-score| > 1.96, **Supplemental Figure 1B**, **Supplemental Table 7**).

After the 3-week CR intervention, BMI values were decreased significantly in both ETB and ETP subjects **(Figure 2C**; Wilcoxon Signed-rank test, p < 0.05). Interestingly, subsequent analysis revealed that ETP subjects showed significantly greater BMI loss ratio than ETB subjects (Wilcoxon rank-sum test, p < 0.05; mean BMI loss ratio 3.27% versus 1.84%; **Figure 2D**; see Materials and Methods).

Overall gut microbiome composition is stable to CR Intervention

We next investigated the extent and impacts of the CR intervention on gut microbial composition in subjects of different enterotypes. Principal coordinate analysis (PCoA) based on species abundance profile of all samples showed that the projected coordinates

of each enterotype group did not change significantly before and after the intervention
(**Figure 3A**, p > 0.05). Furthermore, 23 of 28 ETB subjects and 12 of 13 ETP subjects
were assigned to the same enterotypes after the intervention (**Supplemental Table 2**). In
addition, α-diversity **(Figure 3B-C)** and β-diversity **(Figure 3D-E)** at the genera and
species level of fecal samples also showed no significant changes before and after the
intervention in two enterotype groups, respectively (Wilcoxon Signed-rank test, P **>**
0.05). We further revealed that no common species differed significantly in abundance
before and after the intervention in each enterotype group (**Supplemental Table 8**, BH-
adjusted P > 0.05). All these findings suggest overall stable gut microbial composition in
response to a 3-week 50% energy deficit CR intervention.

On the other hand, we observed that the abundances of several functional pathways were
changed after the intervention. For instance, abundances of pathways for pentose and
glucuronate interconversions (map00040) and galactose metabolism (map00052) were
significantly decreased after the CR intervention in both enterotype groups. ETB
individuals additionally exhibited significantly reduced levels of pathways involved in
cofactors metabolism (pantothenate and CoA biosynthesis, map00770; biotin metabolism,
map00780) and increased levels of pathways for amino acids metabolism (phenylalanine
metabolism, map00360; cysteine and methionine metabolism, map00270) after the
intervention (**Supplemental Table 9**).

Enterotype-independent alterations of blood amino acids to CR Intervention

Reflecting differential gut microbial functional potentials including amino acid
metabolism between two enterotype groups, we asked whether blood amino acid
composition was associated with enterotype. PCA of baseline amino acid profiles showed
no separation of subjects of two enterotypes (**Figure 4A,** PERMANOVA, p > 0.05, See
Material and Methods). In line with this result, we found no significant differences of
baseline levels of blood amino acids between two enterotype groups (Wilcoxon rank-sum
test, BH-adjusted P > 0.05, **Figure 4B, Supplemental Table 10**).

We next examined the potential impacts of the CR diet on blood amino acids. Notably,
we observed similar changes in multiple blood amino acid concentrations in subjects of

both enterotypes in response to the CR intervention (**Figure 4C, Supplemental Table**
11). We therefore combined all samples and found that levels of 13 blood amino acids or
amino acid derivatives such as α-aminoisobutyric acid, β-alanine, glycine, serine, lysine
and its degradation product 2-aminoadipic acid, and 3-methyl-histidine were significantly
increased whereas only one measured amino acid tyrosine was significantly decreased
after the intervention (Wilcoxon Signed-rank test, BH-adjusted $P < 0.05$, **Figure 4D**). In
addition, we observed no significant differences in levels of blood amino acids between
ETB and ETP subjects after the CR intervention (Wilcoxon rank-sum test, BH-adjusted
$P > 0.05$, **Supplemental Table 12**), suggesting the effects of the CR intervention on
blood amino acids were enterotype independent.

Prediction of BMI loss ratio induced by CR intervention using gut microbial species

Considering the differential response in the BMI loss ratio to the CR intervention in two
enterotypes, we next asked whether we could predict BMI loss ratio from the baseline
omics measures. We thus built a Lasso shrinkage model between baseline levels of gut
microbial species and blood amino acids and BMI loss ratio (**See Materials and**
Methods). We successfully selected 7 gut microbial species showing associations with
BMI loss ratio (Coefficient estimate > 0, **Figure 5A**). Interestingly, the relative
abundances of 2 selected species *C. bolteae* and *C. ramosum*, which were enriched in
ETB (Wilcoxon rank-sum test, BH-adjusted $P < 0.05$, **Supplemental Figure 1A,**
Supplemental Table 8), were negatively correlated with BMI loss ratio. On the other
hand, the relative abundance of *D. longicatena* which was slightly enriched in ETP
(Wilcoxon rank-sum test, BH-adjusted $P = 0.06$) was positively correlated with BMI loss
ratio (**Figure 5A**). Baseline abundances of the other 4 Lasso selected species, however,
showed no significant differences between two enterotype groups (BH-adjusted $P > 0.05$,
Supplemental Table 6). Among them, the abundances of *Coprococcus eutactus*,
Streptococcus mitis and *Clostridiales genomosp. BVAB3* were positively associated with
BMI loss ratio whereas the abundance of *Mobiluncus curtisii* was negatively associated
with BMI loss ratio (**Figure 5A**).

To estimate the performance of 7 gut microbial species on the prediction of BMI loss
ratio, we applied a general linear model between predicted BMI loss ratios and true

values using leave-one-out cross validation (LOOCV). Notably, the result showed the predicted BMI loss ratio was significantly correlated with the true BMI loss ratio with a Spearman's rho of 0.646 (**Figure 5B**, See Materials and Methods). By contrast, we found that the individual baseline BMI could hardly predict their BMI loss after the CR intervention (Spearman's rho = -0.016, **Supplemental Figure2**).

Discussion

Here we evaluated changes in BMI, blood amino acids and the gut microbiota of 41 nonobese subjects stratified by enterotype after a 3-week 50% energy deficit CR intervention. We revealed that CR had significant effects on BMI and ETP subjects had higher BMI loss ratio than ETB subjects. Additionally, CR showed only minor impacts on gut microbial composition but significantly impacted blood amino acids in an enterotype-independent manner. We further selected 7 gut bacterial species whose baseline relative abundances could well predict BMI loss ratio induced by the CR intervention.

In this study, nonobese ETP individuals showed greater BMI loss than ETB individuals after our 3-week intervention. Similar findings were also reported in overweight individuals that high *Prevotella/Bacteroides* ratio group had higher weight loss than low *Prevotella/Bacteroides* ratio group when receiving the New Nordic Diet [17] and a 500 kcal/day energy deficit diet [9]. Importantly, we successfully constructed a BMI loss ratio prediction model based on baseline relative abundance of 7 gut microbial species. Among them, *C. ramosum* and *C. bolteae* (negatively associated with BMI loss ratio in our model) were significantly more enriched in ETB subjects while *D. longicatena* (positively associated with BMI loss ratio in our model) was slightly more enriched in ETP subjects. These three enterotype-specific species in our prediction model also have been reported by several other studies. A previous study showed that *C. ramosum* could enhance high-fat-diet-induced obesity in gnotobiotic mice, probably by enhancing nutrient absorption [33]. Two human studies have also reported a positive link between the abundance of *C. ramosum* and several obese-related metabolic parameters [34, 35]. In an obese

postmenopausal women study, the fecal abundance of *C. bolteae* was positively correlated with multiple metabolic markers such as fasting glucose and insulin resistance, whereas *D. longicatena* was negatively correlated [36].

Combining the results of our prediction model and all the above-mentioned studies, the different enrichment of *C. ramosum, C. bolteae* and *D. longicatena* between ETB and ETP subjects may help explain why ETP subjects showed significantly larger BMI loss ratio than ETB subjects. Moreover, all selected 7 species annotated to genera *Coprococcus*, *Streptococcus*, *Clostridium*, *Dorea* and *Mobiluncus* were not the main contributors to enterotypes, suggesting the need for human gut microbial stratification in a higher resolution than enterotypes in further microbiome-based personalized intervention studies.

Attention should also be paid that levels of multiple amino acids and their derivatives were significantly increased in healthy non-obese subjects after the CR intervention, including 3-methylhistidine, a well-established biomarker of skeletal muscle protein turnover [37, 38]. Previous studies have reported that skeletal muscle acted as a fuel reserve to provide glucose disposal for other organs during fasting in human [38, 39]. Since the CR in our study was 50% of normal energy-intake and contained much lower protein than a normal diet, it can be speculated that skeletal muscle proteins might be partially broken down for supplying energy. By contrast, another previous study which provided a high-soy-protein diet induced weight loss without losing muscle mass in obese and pre-obese subjects [40]. All these findings suggest that the availability of adequate dietary proteins in further diet interventions is important to preserve healthy muscle mass during weight loss.

Acknowledgements

We are very grateful to acknowledge colleagues from the China National Genebank for preparing calorie restriction diets, and for DNA extraction, library construction and sequencing.

Author Contributions Statement

Z.H.Z and K.Y.C designed the project. P.S.C and K.Y.C oversaw the collection of fecal and blood samples and phenotypic data. H.Z, D.W, H.H.R and Z.H.Z established the concept and analysis framework of the study. H.Z, H.H.R and D.W performed the bioinformatic analysis. H.Z wrote the first version of the manuscript. D.W and Z.H.Z substantially revised the manuscript. All authors contributed to and approved the final manuscript.

Funding source and disclosure of conflicts of interests

The authors declare that they have no competing interests. This work was funded by National Key Research and Development Program of China (No.2017YFC0909703) and support of Shenzhen Municipal Government of China (No. CXB201108250098A).

Ethics approval and consent to participate

The study protocol was approved by the institutional review board on bioethics and biosafety of BGI-Shenzhen, Shenzhen (NO. BGI-IRB 17020). Written informed consent was obtained from all participants before the CR intervention.

Availability of data and material

The metagenomic sequencing data of this study have been deposited in the China National Genebank Nucleotide Sequence Archive (CNSA) (https://db.cngb.org/cnsa/) under the BioProject number CNP0000247. The scripts and gut microbial profiles used to perform statistical analysis and figure plotting are available at https://github.com/HuaZou/NonobeseBP.

Reference

1. Zheng X, Wang S, Jia W: **Calorie restriction and its impact on gut microbial composition and global metabolism**. *Front Med* 2018, **12**(6):634-644.
2. Schmedes MS, Yde CC, Svensson U, Håkansson J, Baby S, Bertram HC: **Impact of a 6-week very low-calorie diet and weight reduction on the serum and fecal metabolome of overweight subjects**. *European Food Research and Technology* 2015, **240**(3):583-594.
3. Redman LM, Huffman KM, Landerman LR, Pieper CF, Bain JR, Muehlbauer MJ, Stevens RD, Wenner BR, Kraus VB, Newgard CB *et al*: **Effect of caloric restriction with and without exercise on metabolic intermediates in nonobese men and women**. *J Clin Endocrinol Metab* 2011, **96**(2):E312-321.
4. Ravussin E, Redman LM, Rochon J, Das SK, Fontana L, Kraus WE, Romashkan S, Williamson DA, Meydani SN, Villareal DT *et al*: **A 2-Year Randomized Controlled Trial of Human Caloric Restriction: Feasibility and Effects on Predictors of Health Span and Longevity**. *J Gerontol A Biol Sci Med Sci* 2015, **70**(9):1097-1104.
5. Martin CK, Bhapkar M, Pittas AG, Pieper CF, Das SK, Williamson DA, Scott T, Redman LM, Stein R, Gilhooly CH *et al*: **Effect of Calorie Restriction on Mood, Quality of Life, Sleep, and Sexual Function in Healthy Nonobese Adults: The CALERIE 2 Randomized Clinical Trial**. *JAMA Intern Med* 2016, **176**(6):743-752.
6. Lefevre M, Redman LM, Heilbronn LK, Smith JV, Martin CK, Rood JC, Greenway FL, Williamson DA, Smith SR, Ravussin E *et al*: **Caloric restriction alone and with exercise improves CVD risk in healthy non-obese individuals**. *Atherosclerosis* 2009, **203**(1):206-213.
7. Santacruz A, Marcos A, Warnberg J, Marti A, Martin-Matillas M, Campoy C, Moreno LA, Veiga O, Redondo-Figuero C, Garagorri JM *et al*: **Interplay between weight loss and gut microbiota composition in overweight adolescents**. *Obesity (Silver Spring)* 2009, **17**(10):1906-1915.
8. Ott B, Skurk T, Hastreiter L, Lagkouvardos I, Fischer S, Buttner J, Kellerer T, Clavel T, Rychlik M, Haller D *et al*: **Effect of caloric restriction on gut permeability, inflammation markers, and fecal microbiota in obese women**. *Sci Rep* 2017, **7**(1):11955.
9. Hjorth MF, Blaedel T, Bendtsen LQ, Lorenzen JK, Holm JB, Kiilerich P, Roager HM, Kristiansen K, Larsen LH, Astrup A: **Prevotella-to-Bacteroides ratio predicts body weight and fat loss success on 24-week diets varying in macronutrient composition and dietary fiber: results from a post-hoc analysis**. *Int J Obes (Lond)* 2019, **43**(1):149-157.
10. Biolo G, Ciocchi B, Stulle M, Bosutti A, Barazzoni R, Zanetti M, Antonione R, Lebenstedt M, Platen P, Heer M: **Calorie restriction accelerates the catabolism of lean body mass during 2 wk of bed rest**. *The American journal of clinical nutrition* 2007, **86**(2):366-372.
11. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI: **An obesity-associated gut microbiome with increased capacity for energy harvest**. *Nature* 2006, **444**(7122):1027-1031.
12. Kang C, Zhang Y, Zhu X, Liu K, Wang X, Chen M, Wang J, Chen H, Hui S, Huang L *et al*: **Healthy Subjects Differentially Respond to Dietary Capsaicin Correlating with Specific Gut Enterotypes**. *J Clin Endocrinol Metab* 2016, **101**(12):4681-4689.
13. Tolhurst G, Heffron H, Lam YS, Parker HE, Habib AM, Diakogiannaki E, Cameron J, Grosse J, Reimann F, Gribble FM: **Short-chain fatty acids stimulate glucagon-like peptide-1 secretion via the G-protein–coupled receptor FFAR2**. *Diabetes* 2012, **61**(2):364-371.

437   14.   Arumugam M, Raes J, Pelletier E, Le Paslier D, Yamada T, Mende DR, Fernandes GR, Tap
438   J, Bruls T, Batto JM *et al*: **Enterotypes of the human gut microbiome**. *Nature* 2011,
439   **473**(7346):174-180.
440   15.   Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, Bewtra M, Knights D,
441   Walters WA, Knight R *et al*: **Linking long-term dietary patterns with gut microbial
442   enterotypes**. *Science* 2011, **334**(6052):105-108.
443   16.   Vandeputte D, Falony G, Vieira-Silva S, Tito RY, Joossens M, Raes J: **Stool consistency is
444   strongly associated with gut microbiota richness and composition, enterotypes and
445   bacterial growth rates**. *Gut* 2016, **65**(1):57-62.
446   17.   Hjorth MF, Roager HM, Larsen TM, Poulsen SK, Licht TR, Bahl MI, Zohar Y, Astrup A: **Pre-
447   treatment microbial Prevotella-to-Bacteroides ratio, determines body fat loss success
448   during a 6-month randomized controlled diet intervention**. *Int J Obes (Lond)* 2018,
449   **42**(3):580-583.
450   18.   Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, Britt EB, Fu X, Wu Y, Li L *et al*:
451   **Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes
452   atherosclerosis**. *Nat Med* 2013, **19**(5):576-585.
453   19.   Zeevi D, Korem T, Zmora N, Israeli D, Rothschild D, Weinberger A, Ben-Yacov O, Lador D,
454   Avnit-Sagi T, Lotan-Pompan M *et al*: **Personalized Nutrition by Prediction of Glycemic
455   Responses**. *Cell* 2015, **163**(5):1079-1094.
456   20.   Korem T, Zeevi D, Zmora N, Weissbrod O, Bar N, Lotan-Pompan M, Avnit-Sagi T,
457   Kosower N, Malka G, Rein M *et al*: **Bread Affects Clinical Parameters and Induces Gut
458   Microbiome-Associated Personal Glycemic Responses**. *Cell Metab* 2017, **25**(6):1243-
459   1253 e1245.
460   21.   Potts M: **China's one child policy**. *BMJ* 2006, **333**(7564):361-362.
461   22.   Wang SS, Lay S, Yu HN, Shen SR: **Dietary Guidelines for Chinese Residents (2016):
462   comments and comparisons**. *J Zhejiang Univ Sci B* 2016, **17**(9):649-656.
463   23.   Wan Y, Wang F, Yuan J, Li D: **Optimal dietary macronutrient distribution in China
464   (ODMDC): a randomised controlled-feeding trial protocol**. *Asia Pac J Clin Nutr* 2017,
465   **26**(5):972-980.
466   24.   Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, Liang S, Zhang W, Guan Y, Shen D *et al*: **A
467   metagenome-wide association study of gut microbiota in type 2 diabetes**. *Nature* 2012,
468   **490**(7418):55-60.
469   25.   Fang C, Zhong H, Lin Y, Chen B, Han M, Ren H, Lu H, Luber JM, Xia M, Li W *et al*:
470   **Assessment of the cPAS-based BGISEQ-500 platform for metagenomic sequencing**.
471   *Gigascience* 2018, **7**(3):1-8.
472   26.   Li J, Jia H, Cai X, Zhong H, Feng Q, Sunagawa S, Arumugam M, Kultima JR, Prifti E, Nielsen
473   T *et al*: **An integrated catalog of reference genes in the human gut microbiome**. *Nat
474   Biotechnol* 2014, **32**(8):834-841.
475   27.   Costea PI, Hildebrand F, Arumugam M, Bäckhed F, Blaser MJ, Bushman FD, de Vos WM,
476   Ehrlich SD, Fraser CM, Hattori M *et al*: **Enterotypes in the landscape of gut microbial
477   community composition**. *Nature Microbiology* 2017, **3**(1):8-16.
478   28.   Patil KR, Nielsen J: **Uncovering transcriptional regulation of metabolism by using
479   metabolic network topology**. *Proceedings of the National Academy of Sciences of the
480   United States of America* 2005, **102**(8):2685-2689.
481   29.   Kanehisa M, Goto S: **KEGG: kyoto encyclopedia of genes and genomes**. *Nucleic acids
482   research* 2000, **28**(1):27-30.

30.   Jie Z, Liang S, Ding Q, Tang S, Li F, Wang D, Lin Y, Chen P, Cai K, Rao W *et al*: **A multi-omic**
cohort as a reference point for promoting a healthy human gut microbiome. *bioRxiv*
2019:585893.
31.   Tibshirani R: **Regression shrinkage and selection via the lasso: a retrospective**. *Journal*
of the Royal Statistical Society: Series B (Statistical Methodology) 2011, **73**(3):273-282.
32.   Lim MY, Rho M, Song YM, Lee K, Sung J, Ko G: **Stability of gut enterotypes in Korean**
monozygotic twins and their association with biomarkers and diet. *Sci Rep* 2014,
4:7348.
33.   Woting A, Pfeiffer N, Loh G, Klaus S, Blaut M: **Clostridium ramosum promotes high-fat**
diet-induced obesity in gnotobiotic mouse models. *MBio* 2014, **5**(5):e01530-01514.
34.   Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, Almeida M, Arumugam M,
Batto JM, Kennedy S *et al*: **Richness of human gut microbiome correlates with**
metabolic markers. *Nature* 2013, **500**(7464):541-546.
35.   Karlsson FH, Tremaroli V, Nookaew I, Bergstrom G, Behre CJ, Fagerberg B, Nielsen J,
Backhed F: **Gut metagenome in European women with normal, impaired and diabetic**
glucose control. *Nature* 2013, **498**(7452):99-103.
36.   Brahe LK, Le Chatelier E, Prifti E, Pons N, Kennedy S, Hansen T, Pedersen O, Astrup A,
Ehrlich SD, Larsen LH: **Specific gut microbiota features and metabolic markers in**
postmenopausal women with obesity. *Nutr Diabetes* 2015, **5**:e159.
37.   Ballard F, Tomas F: **3-Methylhistidine as a measure of skeletal muscle protein**
breakdown in human subjects: the case for its continued use. *Clinical science* 1983,
65(3):209-215.
38.   Tipton KD, Hamilton DL, Gallagher IJ: **Assessing the Role of Muscle Protein Breakdown**
in Response to Nutrition and Exercise in Humans. *Sports Med* 2018, **48**(Suppl 1):53-64.
39.   Wolfe RR: **The underappreciated role of muscle in health and disease**. *The American*
journal of clinical nutrition 2006, **84**(3):475-482.
40.   Deibert P, Konig D, Schmidt-Trucksaess A, Zaenker KS, Frey I, Landmann U, Berg A:
Weight loss without losing muscle mass in pre-obese and obese subjects induced by a
high-soy-protein diet. *Int J Obes Relat Metab Disord* 2004, **28**(10):1349-1352.

Figures, tables and additional files

Figure 1. Overview of the experimental design. Illustration of our experimental design, including a 1-week run-in period (baseline) and a 3-week calorie restriction (CR) dietary intervention trial with 50% energy deficit diet (male, ~1200Kcal/day; female, ~1000Kcal/day). BMI, fasting blood samples and fecal samples of 41 enrolled healthy subjects were collected before and after the CR intervention to assess its effects on BMI, blood amino acids and gut microbiome in two enterotype groups.

Figure 2. A short-term CR intervention altered BMI. (A) Principal coordinates analysis (PCoA) based on genera-level Bray-Curtis distance between all baseline fecal samples. Orange, subjects of enterotype *Bacteroides* (ETB) and blue, subjects of enterotype *Prevotella* (ETP). **(B)** Baseline BMI between ETB and ETP subjects. **(C)** Changes in BMI before and after intervention in individuals of each enterotype. **(D)** Boxplot showing BMI loss ratio between ETB subjects and ETP subjects. *, $P < 0.05$; **, $P < 0.01$.

Figure 3. Overall gut microbial composition of two enterotypes before and after the CR. (A) Species-based principal coordinates analysis (PCoA) of subjects before and after the CR trial. Trangers, samples of ETB; Circles, samples of ETP. Arrows indicate paried samples from the same indivudul. Boxplot showing the projected coordinate 1 (PCo1) and PCo2 of samples before and after the intervention. **(B-C)** α-diversity (Shannon index) at the gene and species levels before (blue) and after (red) intervention in each enterotype group. **(D-E)** β-diversity (Bray-Curtis distance) at the gene and species levels before (blue) and after (red) the intervention in each enterotype group.-ns, no significance, $P > 0.05$, Wilcoxon signed-rank test.

Figure 4. A short-term CR intervention altered blood amino acids. (A) Principal component analysis (PCA) of 41 subjects using baseline blood amino acid profiles. Orange, ETB; blue, ETP. **(B)** Comparison of baseline blood amino acid concentrations between ETB subjects (orange) and ETP subjects (blue). Wilcoxon rank-sum test, P values are transformed to Z-scores to represent enrichment directions. **(C)** Amino acid based PCA of samples before and after the intervention. Trangers, samples of ETB

subjects; Circles, samples of ETP subjects. Arrows indicate paried samples from the same indivudul. **(D)** Changes in blood amino acid concentrations of ETB subjects, ETP subjects and all subjects before and after the intervention. Wilcoxon rank-sum test, P values are transformed to Z-scores to represent enrichment directions. Dashlied line indicates the absolute Z score of 1.96 (P = 0.05). Asterisk (*) indicates significance of Benjamini-Hochberg (BH) adjusted P < 0.05.

Figure5. Prediction of BMI loss ratio using baseline abundances of gut microbial species. (A) Bar plot showing the 7 gut microbial species selected by least absolute shrinkage and selection operator (Lasso). Bar length indicates regression coefficient of each species estimated by Lasso. Orange, species significantly enriched in ETB subjects (BH-adjusted P < 0.05); blue, species slightly enriched in ETP subjects (P < 0.05 and BH-adjusted P = 0.06); grey, species with no significant enrichment between two enterotypes (P > 0.05).

(B) Scatter plot showing prediction performance of BMI loss ratio based on the 7 selected species. Leave-one-out cross validation (LOOCV) was applied to evaluate the performance of generalized liner model (GLM), showing a strong Spearman's rho between actual BMI loss ratios and predicted BMI loss ratios of 0.646. Red circles, ETB individuals; blue circles, ETP individuals.

Supplemental Figure 1. Baseline gut microbial differences between two enterotype groups. **(A)** Differentially enriched species between two enterotype groups. Orange and blue represent species overrepresented in ETB and ETP subjects, respectively. **(B)** Differential enrichment of KEGG pathways between ETB and ETP subjects. Dashed lines indicate a reporter score of 1.96, corresponding to 95% confidence in a normal distribution. Orange and blue bars indicate reporter scores of selected KEGG pathway overrepresented in ETB and ETP subjects, respectively.

Supplemental Figure 2. Performance of baseline BMI for prediction of BMI loss ratio.

Scatter plot showing prediction performance of BMI loss ratio using baseline BMI values.
A Spearman's rho between actual BMI loss ratios and predicted BMI loss ratios was -
0.016. Red circles, ETB individuals; blue circles, ETP individuals.

Table 1. Comparison of baseline phenotypes between ETB and ETP subjects

Table 2. Cohort description

Supplemental Table 1. Statistics for metagenomic sequencing data of fecal samples

Supplemental Table 2. Enterotype classification of individual fecal samples before and
after the intervention

Supplemental Table 3. List of common genera

Supplemental Table 4. List of common species

Supplemental Table 5. Comparison of baseline genera relative abundance between ETB
and ETP subjects

Supplemental Table 6. Comparison of baseline species relative abundance between ETB
and ETP subjects

Supplemental Table 7. Differential enrichment of KEGG pathway between ETB and
ETP subjects at baseline

Supplemental Table 8. Comparison of species relative abundance before and after the
intervention in each enterotype

Supplemental Table 9. Differential enrichment of KEGG pathway before and after the
intervention in each enterotype

Supplemental Table 10. Comparison of baseline blood amino acid levels between ETB
and ETP subjects

Supplemental Table 11. Comparison of blood amino acid levels before and after the
intervention

Supplemental Table 12. Comparison of blood amino acid levels between ETB and ETP

subjects after the intervention

Table 1. Comparison of baseline phenotypes of ETB and ETP subjects

	ETB Group (Mean ± SD)	ETP Group (Mean ± SD)	P value ETB vs ETP
Number of subjects	28	13	
Sex (female/male)	16/12	8/5	1
Age	29 ± 6	30 ± 7	0.44
BMI (kg/m^2)	23.50 ± 2.81	24.21 ± 2.85	0.42
Weight (kg)	64.40 ± 11.37	65.81 ± 12.39	0.62

Table 2. Cohort Description

	Cohort (Mean ± SD)
Number of subjects	41
Sex (female/male)	24/17
Age	30 ± 6
BMI (kg/m^2)	23.72 ± 2.81
Weight(kg)	64.84 ± 11.57