Community size affects the signals of ecological drift and niche selection on biodiversity

Tadeu Siqueira[1], Victor S. Saito[2], Luis M. Bini[3], Adriano S. Melo[3], Danielle K. Petsch[3], Victor L.

Landeiro[4], Kimmo T. Tolonen[5], Jenny Jyrkänkallio-Mikkola[6], Janne Soininen[6], Jani Heino[7]

[1]Institute of Biosciences, São Paulo State University (UNESP), Av. 24 A 1515, Rio Claro, SP,

13506-900, Brazil

[2]Universidade Federal de São Carlos (UFSCar), Departamento de Ciências Ambientais, São Carlos,

Brazil

[3]Departamento de Ecologia, ICB, Universidade Federal de Goiás, Goiânia, GO, Brazil

[4]Departamento de Botânica e Ecologia, IB, Universidade Federal de Mato Grosso, Cuiabá, MT,

Brazil

[5]University of Jyväskylä, Department of Biological and Environmental Sciences, Jyväskylä,

Finland

[6]University of Helsinki, Department of Geosciences and Geography, Helsinki, Finland

[7]Finnish Environment Institute, Freshwater Centre, Oulu, Finland

Keywords: aquatic insects; beta diversity deviation; community assembly; demographic

stochasticity; dispersal; metacommunities; null models

Abstract. Ecological drift can override the effects of deterministic niche selection on small populations and drive the assembly of small communities. We tested the hypothesis that smaller local communities are more dissimilar among each other because of ecological drift than larger communities, which are mainly structured by niche selection. We used a unique, comprehensive dataset on insect communities sampled identically in a total of 200 streams in climatically different regions (Brazil and Finland) that differ in community size by fivefold. Null models allowed us to estimate the magnitude to which beta diversity deviates from the expectation under a random assembly process while taking differences in species richness and relative abundance into account, i.e., beta deviation. Beta diversity of small tropical communities was consistently higher but closer to null expectations than β-diversity of large communities. However, although β-deviation and community size were strongly related in both regions, the direction of the relationship varied according to dissimilarity metrics. While incidence-based β-diversity was lower than expected (communities were less dissimilar than null expectations) and negatively related to community size in Brazil, abundance-based β-diversity was higher than expected (communities were more dissimilar than null expectations) and positively related to community size in both regions. We suggest that ecological drift drives variation in small communities by increasing the chances of species with low abundance and narrow distribution to occur within the metacommunity. Also, while weak niche selection and high dispersal rates likely reduced variation in community structure among large tropical streams, niche selection was likely sufficient to cause non-random variations in the relative abundances of genera among large communities in both regions. Habitat destruction, overexploitation, pollution, and reductions in connectivity have been reducing the size of biological communities. These environmental pressures will make smaller communities more vulnerable to novel conditions and render community dynamics more unpredictable, as random demographic processes should prevail under these conditions. Incorporation of community size into ecological models should provide conceptual, empirical and applied insights into a better understanding of the processes driving changes in biodiversity.

Introduction

Recent syntheses in community ecology propose that four main processes drive the dynamics of metacommunities – deterministic niche selection, ecological drift, dispersal and speciation (Vellend 2010, 2016, Leibold and Chase 2018). At broad spatial scales, dispersal, colonization history, and spatial heterogeneity are thought to produce major biodiversity patterns (Leibold and Chase 2018), while speciation also plays a role by altering the composition of regional species pools at longer time frames (Vellend 2010). Within localities, niche selection determines community structure mainly through species interactions, differential use of resources and species responses to environmental gradients (Leibold and Chase 2018). However, stochasticity may also play a major role in driving local community dynamics, for example, when demographic events occur at random with respect to species identities (Vellend et al. 2014). Indeed, mechanistic models (Orrock and Fletcher 2005, Orrock and Watling 2010) and recent empirical evidence (Gilbert and Levine 2017) suggest that ecological drift can even override the effects of niche selection under certain circumstances, such as when species populations in local communities are small and isolated from other populations.

Small communities have few individuals per unit area, and thus random birth and death events are likely to have a high impact on their structure (species composition and relative abundances). For example, the species composition of a local community would change if all individuals of a species die before reproducing. This is likely to happen in nature especially among small populations or in those large populations in which only a reduced fraction of adults successfully reproduce (Bunn and Hughes 1997). Some models indicate that ecological drift can even reduce competition asymmetries in small communities to a level that strong and weak competitors become effectively neutral (Orrock and Watling 2010) – i.e., the negative effect of a superior competitor on other species is relatively small compared to the effects of demographic stochasticity. Indeed, in an experiment with annual plants, Gilbert and Levine (2017) found that larger communities converged to a state in which a strong competitor dominated after three years,

whereas in smaller communities the strong competitor co-occurred with other species at different

densities. Thus, both theory and experimental evidence suggest that ecological drift can change the

structure of local communities by altering species relative abundances as well as species

occurrences. Investigating the relationship between variation in community structure and

community size would help us better understanding the relative importance of deterministic and

stochastic processes on beta diversity. This should also be relevant from an applied perspective, as

many types of environmental disturbances, such as deforestation and floods, tend to reduce the size

of local communities (Barnes et al. 2014, Petsch et al. 2015).

87   Most methods commonly used to estimate β-diversity are affected by differences in species

richness or species abundance distributions (Chase and Myers 2011). This is undesirable for

understanding the beta diversity-community size relationship for two main reasons. First, estimates

of β-diversity can be influenced by random sampling effects that are neutral with respect to species

identity (Chase and Myers 2011, Myers et al. 2013, 2015). For example, let us assume that the

species composition of 10 local communities embedded in a larger metacommunity represents each

a random subset (say 5 species) of the regional species pool with, for instance, 100 species. This

would likely result in high β-diversity as, by chance, many pairs of local communities would not

share species. Second, studies employing incidence-based estimates of β-diversity (e.g., Jaccard or

Sørensen dissimilarity indices) do not capture variation in species relative abundances (Anderson et

al. 2011). In this case, if the same species occur at two sites but with different relative abundance,

β-diversity should not be equal. A solution to the first issue is to use a null model to produce

expected values under random assembly from a large species pool, contrast observed and expected

values and use the difference between them as estimates of β-deviation; i.e., the extent to which β-

diversity differs from null expectations (Kraft et al. 2011, Myers et al. 2013, 2015, Catano et al.

2017). In this case, positive and negative values of β-deviation indicate that communities are more

dissimilar and less dissimilar than expected by chance, respectively. Beta deviation values close to

zero indicate communities are as dissimilar as expected by chance (Kraft et al. 2011, Chase et al.

2011, Catano et al. 2017). Beta deviation provides a means to determine whether β-diversity

deviates from the expectations of stochastic assembly and if the magnitude of the deviation is

related to community size. A solution to the second issue is to analyze the data with dissimilarity

coefficients that take into account species composition and species relative abundance (e.g., Bray-

Curtis; Anderson et al. 2011). Such dissimilarity coefficients provide complementary information

regarding the main mechanisms responsible for β-diversity patterns (e.g., Siqueira et al. 2015).

In this study, we tested the hypothesis that ecological drift is a major process causing

variation among small communities. To reach our goal, we used a unique, comprehensive dataset

on aquatic insect communities sampled using identical methods in a total of 200 streams in two

climatically highly different regions (100 in Brazil and 100 in Finland). The sampling design

included 5 streams (communities) per watershed (20 in each region) and provided us replicates of

metacommunities (watersheds; Fig. S1). This allowed us to make specific predictions considering a

community size gradient. Our previous study showed that local community size is, on average, five-

fold larger in boreal than in tropical streams, with the smallest boreal stream communities being as

large as the largest tropical communities (Heino et al. 2018). Thus, we expected that β-diversity

would be high (E1) and close (E2) to null expectations (random assembly from the regional pool) in

watersheds with the smallest communities (some watersheds in Brazil only; Fig. 1). This would

indicate that ecological drift plays a major role in structuring these small subtropical communities.

Second, all else being equal, we expected that (E2a) β-deviation would be positive in all

watersheds, but higher in the largest communities (Fig. 1). This would indicate that niche selection

and sufficient dispersal rates are the main processes resulting in large communities to be more

dissimilar than expected by chance, as species sorting occurs when dispersal is sufficient to allow

individuals to reach sites that match their ecological requirements (Winegardner et al. 2012, Leibold

and Chase 2018). Although dispersal limitation can also cause positive values of β-deviation, this

was unlikely in our study system as the watersheds we studied are not large enough to lead to strong

dispersal limitation (see details below). Taken together, these two expectations would lead to a

negative relationship between β-diversity (before controlling for sampling effects) and community

size, but to a positive relationship between β-deviation and community size (Fig. 1). Finally,

because environmental heterogeneity and spatial extent are also expected to increase β-diversity

(Heino et al. 2015a), we included them as co-variates in our models (Fig. S1).

Material and methods

Study area and sampling

The streams sampled in Brazil and Finland were distributed across a strong gradient of land

cover among watersheds. It is well established that watershed land cover influences the structure of

stream communities (Hynes 1975, Allan 2004, Roque et al. 2010, Siqueira et al. 2015). In Brazil,

we sampled 100 streams distributed in 20 watersheds located in the southeastern region of the

country – i.e., five streams per watershed, with spatial extent of 70 km in north-south and 120 km in

east-west directions (Fig. S2). These streams drain through three major Atlantic Forest protected

areas ('Carlos Botelho', 'Intervales' and 'Alto Ribeira' State Parks; São Paulo State). The

watersheds are mainly dominated by pastures and planted forests (*Eucalyptus* and *Pinus*). The

region has a dry season from April to August (average rainfall from 45 to 80 mm per month;

average temperature from 16 to 20 $^{\circ}$C) and a wet season from September to March (average rainfall

from 105 mm to 180 mm per month; average temperature from 20 to 23 $^{\circ}$C). Sampling was done

between September and November in 2015.

Figure 1. Graphical representation describing *a priori* expectations (E) about the relationship

between beta diversity and beta deviation with community size in Brazil (vermilion) and Finland

(blue).

The study sites in Finland were situated in the western part of the country. We sampled 100

streams that were distributed in 20 watersheds (as described above) with spatial extents of ca. 500

km and 300 km in north-south and in east-west directions (Fig. S2). The streams drain watersheds

covered with agriculture and boreal forests. Western Finland has four typical seasons: a long winter

that lasts from November to March, a short spring from April to May, a short summer from June to

August, and an autumn period from September to October. Sampling was done in September 2014.

Streams were generally of the same order within each watershed, but varied among watersheds,

including 2nd and 3rd order streams in Brazil, and a few 4th order streams in Finland. In Brazil,

maximum distances between pairs of streams within watersheds varied from 2.48 to 8.86 km,

whereas in Finland it varied from 12.77 to 109.5 km. Most in-stream abiotic variables varied within

a similar range and had similar mean values between regions, except conductivity, total nitrogen

and total phosphorus, which were much higher in Finland. A more detailed description of both

regions can be found in Heino et al. (2018).

At each of the 100 stream sites in both regions, we took a 2-minute kick-net sample (net

mesh size: 0.5 mm), which was composed of four 30-seconds sample units obtained in the main

microhabitats at a riffle site (e.g., which considered differences in current velocity, depth, benthic

particle size and macrophyte cover). The four sample units were pooled, preserved in alcohol in the

field and taken to the laboratory for further processing and identification. All insects were separated

from debris and the following taxonomic orders were identified to genus level: Ephemeroptera,

Odonata, Plecoptera, Megaloptera, Trichoptera and Coleoptera. We sampled 16,113 individuals,

distributed among 83 genera in Brazil, and 86,048 individuals (77 genera) in Finland. The mean

number of genera per stream was 17.84 (standard deviation = 7.46) in Brazil and 14.01 (sd = 5.07)

in Finland, while the mean number of individuals per stream was 181.50 (sd = 111.38) and 886.57

(sd = 700.73), respectively (Heino et al. 2018).

We adopted the definition of community size provided by Orrock and Watling (2010) and

estimated local community size as the number of individuals sampled in a stream site. However, as

β-diversity was estimated for each watershed (i.e., considering 5 stream sites; see below), we

averaged the number of individuals across five streams within each watershed. This resulted in 20

values of community size, one per watershed (Fig. S1).

Beta diversity and beta deviations

We first estimated β-diversity by using the Sørensen incidence-based coefficient and the

Bray-Curtis abundance-based coefficient. To do that, we calculated dissimilarity values between all

pairs of five streams within each of the 20 watersheds, separately for the tropical and boreal

datasets. Our measure of β-diversity was represented by the mean of these values in each watershed.

Thus, we estimated 20 values of Sørensen β-diversity and 20 values of Bray-Curtis β-diversity per

region (Fig. S1).

We estimated β-deviations by using two procedures based on null models. To estimate

incidence-based β-deviations that accounted for random sampling effects, we used a modified

version of the Raup-Crick coefficient following Chase et al. (2011) and the procedures described in

the package vegan (Oksanen et al. 2018): (i) we defined the genus pool as all genera occurring in

each region and their observed occupancy across all 100 streams; (ii) an algorithm calculated the

number of genera that every pair of streams share (SG_{obs}); (iii) and assembled local stream

communities by randomly sampling genus from the pool until reaching the local (observed) genus

richness and by using the observed genus' occupancy frequency to determine the probability to

sample a genus; (iv) step (iii) was repeated 10,000 times to generate 10,000 random matrices of

genus composition and, posteriorly, the number of genera shared between each pair of streams

within each watershed (SG_{exp}); (v) β-deviations were calculated as the number of random draws in

which $SG_{exp} \geq SG_{obs}$ divided by the total number of random draws. This index was rescaled to range

between -1 and 1, "indicating whether local communities are more dissimilar (approaching 1), as

dissimilar (approaching 0), or less dissimilar (approaching -1), than expected by random chance"

within each watershed (Chase et al. 2011). As with β-diversity, this procedure resulted in 20 values

of incidence-based β-deviation per region, with one value per watershed (Fig. S1).

As selection and drift act on individuals, but not adding or removing entire populations, the

use of abundance-based metrics should provide complementary information about assembly

mechanisms. Thus, to estimate abundance-based β-deviations, we followed the procedure described

by Kraft et al. (2011): (i) we defined the genus pool as all genera and their abundances across all

100 streams in each region; (ii) an algorithm then randomly assigned individuals to stream sites

while preserving the overall genus-abundance distribution in the region and the total number of

individuals in each stream site (local community size); (iii) step (ii) was repeated 10,000 to generate

a distribution of pairwise dissimilarities within each watershed assuming random colonization of

individuals from the genera pool; (iv) β-deviations were calculated as the difference between the

observed Bray-Curtis dissimilarity and the mean expected (i.e. simulated) dissimilarity, divided by

the standard deviation of the (null) simulated distribution within each watershed. Positive values

and negative values indicate greater and lower dissimilarity than expected from random assembly,

respectively. This null model allows one to analyze how β-diversity differs from patterns generated

without the processes that cause clumping of species across the landscape (Kraft et al. 2011, Myers

et al. 2015). These procedures also resulted in 20 values of abundance-based β-deviation per region,

one value per watershed (Fig. S1).

We used linear regression models to test whether β-diversity (and β-deviation) was related

to community size. Environmental heterogeneity within watersheds was estimated using

environmental data and a Permutational analysis of multivariate dispersions (PERMDISP), as

described by Anderson et al. (2006). This analysis was based on a Euclidean distance calculated

with the standardized values (zero mean and unit variance) of the following variables: current

velocity (m/s), depth (cm), stream width (cm), % of sand (0.25-2 mm), gravel (2-16 mm), pebble

(16-64 mm), cobble (64-256 mm), and boulder (256-1024 mm), % of canopy cover by riparian

vegetation, pH, conductivity, total nitrogen, and total phosphorus. These were all measured at the

stream riffle scale (for details, see Heino et al. 2018). We also included the following watershed-

scale variables estimated through satellite images within a 400-m buffer along tracts of the sampled

streams: average slope, % of native forest cover, pasture, agriculture, planted forests, urban areas,

mining, water bodies, bare soil, secondary forest cover, and mixed land uses. We mapped land use

and land cover of Brazilian and Finnish watersheds using 5-m resolution RapidEye multispectral

imagery and the CORINE database (https://land.copernicus.eu/pan-european/corine-land-cover),

respectively. Geographical coordinates of the sampling sites were transformed to a Euclidean

distance matrix and then submitted to a PERMDISP procedure (Anderson et al. 2006), using

watershed as a grouping variable, to estimate mean spatial extent. Thus, we regressed our response

variables (β-diversity and β-deviation; with 20 values; one per watershed) against community size,

environmental heterogeneity, and spatial extent separately for each region (Fig. S1). We used the

vegan package (Oksanen et al. 2018) in R (version 3.5.0; R Core Team 2018) to estimate β-

diversity, Raup-Crick β-deviation, environmental heterogeneity, and spatial extent. In the tropical

data, the strongest correlation between explanatory variables was between environmental

heterogeneity and spatial extent (r = 0.47), whereas in the boreal data, it was between community

size and spatial extent (r = 0.36). All other correlations were lower than 0.07. Thus,

multicollinearity was not an issue in our study. Code and data that fully reproduce the analyses are

archived on Zenodo (https://doi.org/10.5281/zenodo.2620550).

Results

There was a strong negative relationship between β-diversity (both with incidence- and

abundance-based data) and community size in Brazil ($b = -0.87$, $t = -7.47$, $p < 0.001$, $R^2 = 0.74$; and

$b = -0.78$, $t = -5.32$, $p < 0.001$, $R^2 = 0.59$, respectively; Fig. 2), indicating that small communities

were more dissimilar among each other than larger communities, as we expected (E1). However,

we found that neither incidence-based (Sørensen) nor abundance-based (Bray-Curtis) β-diversity

were related to community size in Finland ($b = -0.14$, $t = -0.59$, $p = 0.563$; and $b = -0.22$, $t = -0.94$, p

$= 0.357$, respectively).

Figure 2. (A) Incidence-based (Sørensen) and (B) abundance-based (Bray-Curtis) beta diversity-community size (average number of individuals per watershed) relationships within tropical (vermilion) and boreal (blue) stream watersheds ($n = 20$ for each region). The average number of individuals per watershed was calculated with a sample size of 5 streams.

We found strong relationships between β-deviation (i.e., β-diversity apart from null expectations) and community size in both regions, but, contrary to our expectations, they varied according to the type of dissimilarity coefficient (incidence- and abundance-based) and the direction of relationship (negative and positive; Table 1, Fig. 3). As we expected (E2), beta diversity of tropical smaller communities was closer to null expectations than those of larger communities (Fig. 3). However, differently from what we expected, mean Raup-Crick β-deviations were negative in Brazil, varying from -0.98 to -0.02, indicating that tropical communities were less dissimilar in genus composition than expected by random sampling from the species pool. Neither environmental heterogeneity nor spatial extent were related to Raup-Crick β-deviation in Brazil. Raup-Crick β-deviation in boreal watersheds (ranging from -0.99 to 0.30) was not related to community size (Fig. 3A) but it was to environmental heterogeneity (Table 1).

Table 1. Relationship between beta deviation (incidence-based [Raup-Crick] and abundance-based [Bray-Curtis]) and community size, environmental heterogeneity and spatial extent ($n = 20$ watersheds in each region). CS = Community size; EH = Environmental heterogeneity; SE = Spatial extent. R^2 and adj. R^2 = coefficient of determination and adjusted coefficient of determination of the full model, respectively. b = standardized regression coefficient.

			b	se	t	P	R^2	adj. R^2
Brazil	Raup-Crick	CS	**-0.737**	**0.164**	**-4.505**	**<0.001**	0.510	0.419
		EH	-0.114	0.200	-0.572	0.575		
		SE	0.057	0.200	0.289	0.776		
	Bray-Curtis	CS	**0.777**	**0.147**	**5.279**	**<0.001**	0.658	0.594
		EH	0.211	0.167	1.267	0.223		
		SE	-0.087	0.167	-0.522	0.609		

Finland	Raup-Crick	CS	-0.252	0.221	-1.137	0.271	0.319	0.192
		EH	**0.461**	**0.207**	**2.229**	**0.040**		
		SE	0.338	0.222	1.527	0.146		
	Bray-Curtis	CS	**0.728**	**0.168**	**4.328**	**<0.001**	0.607	0.534
		EH	-0.115	0.157	-0.731	0.475		
		SE	0.087	0.168	0.518	0.611		

Bray-Curtis (abundance-based) β-deviation was positively related to community size in both regions (Table 1; Fig. 3B), supporting our expectations (E2a). In tropical watersheds, mean Bray-Curtis β-deviations ranged from 3.08 to 16.66, indicating that communities were more dissimilar than expected by random changes in species abundances and genus composition. Again, β-deviations of smaller communities were closer to zero than those of larger tropical communities, supporting our expectation E1. In boreal watersheds, mean Bray-Curtis β-deviation varied from 17.40 to 102.51, also indicating that communities were more dissimilar than expected by random changes in genus abundances and composition. Neither environmental heterogeneity nor spatial extent were related to Bray-Curtis β-deviation in any region.

Figure 3. (A) Incidence-based (Raup-Crick) and (B) abundance-based (Bray-Curtis) beta deviation-community size (average number of individuals per watershed) relationships within tropical

(vermilion) and boreal (blue) stream watersheds ($n = 20$ for each region). The average number of

individuals per watershed was calculated with a sample size of 5 streams. The dashed grey line

indicates expected beta diversity under null assembly.

Although there was a negative relationship between Raup-Crick β-deviation and community

size in tropical watersheds only, mean incidence-based β-deviation in Brazil was not different from

mean incidence-based β-deviation in Finland (Fig. 3A). This suggests that the departure from null

expectations related to compositional change among streams is similar in tropical and boreal

regions. On the other hand, mean abundance-based β-deviation was five times higher in Finland

than in Brazil, suggesting that departures from null expectations are much higher in boreal than in

tropical streams (Fig. 3B).

Discussion

The role of stochastic community assembly processes in arranging species within and

among communities has gained support from models (Mouquet and Loreau 2003, Orrock and

Fletcher 2005, Durães et al. 2016) and data (Cottenie 2005, Lancaster and Downes 2017, Germain

et al. 2017, Swan and Brown 2017, Valente-Neto et al. 2017). Here, we provide empirical evidence

that community size, a simple characteristic of communities, may mediate the interplay between

niche selection and ecological drift as drivers of β-diversity in tropical and boreal

metacommunities. Based on null models, our estimates of variation in community structure

accounted for both differences in species richness and species relative abundance that deviate from

random assembly (i.e., β-deviations). We showed that β-diversity of smaller tropical communities

deviates less from null expectations than larger communities. This means that the high β-diversity

we observed among smaller communities was, to some extent, indistinguishable from patterns

generated by random changes in local genus richness and community size. As findings generated by

null models are our best approximation of patterns generated by stochastic processes (Kraft et al.

2011, Chase et al. 2011), we suggest that demographic stochasticity plays a major role in small

ecological communities (Orrock and Fletcher 2005, Orrock and Watling 2010, Gilbert and Levine

2017).

Explanations for the major role of demographic stochasticity in small communities involve

the alteration of the occupancy frequency and relative abundance of species with different fitness

(Orrock and Watling 2010). When local communities are small, even species with high fitness are

at a high risk of extinction due to demographic stochasticity in comparison to a situation when

communities have large populations. Consequently, species with low relative abundances have a

chance to increase their populations in small communities (Orrock and Watling 2010, Gilbert and

Levine 2017). As this would be the result of a local demographic random process, and because

dispersal ability is highly variable among species in riverine systems (Heino et al. 2015b, Tonkin et

al. 2018a), the outcome of the assembly would likely differ among local communities, increasing β-

diversity within metacommunities. Indeed, in Brazil, the three most abundant and widespread

genera (Heino et al. 2018) were not dominant in smaller communities: the mayfly *Farrodes* was not

among the most abundant in any of the smallest communities; the beetle *Heterelmis* was among the

most abundant in two communities only; and the caddisfly *Smicridea* was among the most abundant

in three of the five smallest communities. These three genera were dominant in four of the five

largest communities in Brazil. The smallest communities in Brazil were dominated by genera with

intermediate regional abundance and occupancy. For example, *Gripopteryx* (stonefly), *Cloeodes*

(mayfly) and *Callibaetis* (mayfly) were the sixth, eleventh and sixteenth most abundant genera in

Brazil, respectively. In Finland, these differences were less evident, as the three most abundant and

widespread genera dominated small and large communities: the beetle *Elmis* (two of the smallest

vs. three of the largest communities); the mayfly *Baetis* (three vs. three); and stonefly *Nemoura*

(one vs. one). This is likely because the smallest boreal stream communities were as large as the

largest tropical communities (Heino et al. 2018).

The positive relationship between abundance-based β-deviation and community size in both regions is in line with our expectations. The positive slope and values of β-deviation indicate that the variation in community structure among streams increased with community size more than what was expected under random assembly. Thus, in terms of which genera were more abundant or less abundant, and more aggregated or less aggregated, communities within the same watershed differed from each other more than expected by chance, especially in mid to large communities. As dispersal within watersheds was likely not limited, this positive relationship indicates that niche selection was sufficient to cause non-random variations in genera relative abundance and aggregation patterns among large communities. We suggest that as community size increases, demographic stochasticity becomes less important and niche selection determines which species are more abundant locally and widely distributed within the metacommunity. In this case, small random variations in the number of individuals of relatively abundant genera occurring in larger communities, which can only be detected with abundance-based β-diversity metrics, do not result in major changes in genus occurrence.

Although small communities had incidence-based β-deviations values close to random assembly expectations, as we expected, the relationship between β-deviation and community size was negative in Brazil. This result contradicts our expectations and indicates that the genus composition of streams harboring mid to large communities in tropical watersheds is less dissimilar (negative values of incidence-based β-deviation) than patterns predicted by random assembly – i.e., these communities share more genera than expected. In general, communities can be less dissimilar than random expectations in at least two cases. First, dissimilarity should be low when niche selection is spatially constant, as the environment maximizes the fitness of a few species (Vellend 2016). Within watershed environmental heterogeneity was weakly and positively related with incidence-based β-deviation in Finland only, where community sizes are on average five-fold larger than in tropical streams (Heino et al. 2018). This suggests a tendency towards environmental determinism; the higher the environmental variation within watersheds, the higher their β-deviation.

Thus, it is likely that even the large size of boreal communities was not sufficient to allow niche

selection to be the main driver of spatial variation in genus composition among those communities.

Alternatively, we might have failed to measure key abiotic and biotic variables (e.g., biological

interactions) underlying community variation, or despite including key correlates of beta diversity,

our snapshot analysis might not have been enough to represent the complexity of all mechanisms

involved in niche selection in watersheds.

A second reason for communities to be less dissimilar than random expectations is when

dispersal rates are high within the metacommunity (Mouquet and Loreau 2003, Leibold and Chase

2018). By distributing organisms among communities, dispersal can reduce β-diversity if it is

excessive and is combined with a source-sink system, where populations with high growth rates

supply individuals to other localities where they would otherwise be excluded by niche selection

(Mouquet and Loreau 2003, Siqueira et al. 2014). Although dispersal across watersheds tends to be

limited for many aquatic species, those with an adult flight stage may have higher dispersal rates

especially along the stream channel (Hughes 2007, Lancaster and Downes 2017). A meta-analysis

by Muehlbauer et al. (2014) showed that most adult aquatic insects tend to fly ca. 1.5 m around

their natal stream, but that a few individuals can fly 550 m away from the stream, with some

caddisflies being able to reach sites distant more than 650 m. Also, while Macneale et al. (2005)

found that the stonefly *Leuctra ferruginea* was able to fly across headwater forested catchments,

Flenner and Sahlin (2008) estimated annual range expansions of up to 88 km in non-migratory

dragonflies in a boreal region of Sweden. The northern faunas of Europe should also be composed

of species with good dispersal ability, as they have reached these areas since the Last Glacial

Period. On the other hand, many tropical aquatic insects have multiple reproduction events per year,

i.e., many are multivoltine species (Wallace and Anderson 1996, Vásquez et al. 2009). Unless local

population growth rate is low, multivoltine species should have many opportunities per year to

disperse from their natal streams. If a few of these multivoltine species can fly 250 m (a

conservative estimate considering the results by Muehlbauer et al. 2014), after some generations,

some individuals of these species could reach distant sites, at a velocity of one kilometer per year.

We thus suggest that long-term availability for colonization and multivoltinism allowed some

genera to reach widespread distribution within tropical watersheds (Saito et al. 2016). Similarly,

good dispersal abilities of northern species may have masked the role of within-watershed

environmental heterogeneity in the boreal region, making these communities less dissimilar than

expected. Thus, as communities become larger, demographic stochasticity plays a less important

role, while dispersal surplus plays a major role in homogenizing the genus composition of streams

within tropical watersheds. However, this, as almost everything in ecology, should be scale

dependent (Chase et al. 2018). If our watersheds were larger (in extent), dispersal limitation would

likely have played a role and niche selection could have been the major driver of community

structure. Our results reinforce previous suggestions that dispersal should be viewed as a process

occurring over time and not only considering the distance individuals can move in one dispersal

event during one generation (Saito et al. 2015).

Predicting community assembly dynamics has been a major challenge for ecology in a

changing world (Mouquet et al. 2015). Many drivers of environmental change, such as habitat

destruction, overexploitation, pollution, and reductions in landscape connectivity, are likely to cause

reductions in community size. For example, Hallmann et al. (2017) estimated a decline of more than

70% in insect biomass between 1989 and 2016 in Germany and Lister and Garcia (2018) found that

arthropod biomass has fallen 10 to 60 times since 1970 in Puerto Rico. It is possible, then, that

results from previous studies associating high β-diversity with environmental changes were related

to small communities being more variable. For example, Hawkins et al. (2015) showed that

disturbed sites had high β-diversity due to a decreased prevalence of more common taxa. Although

they did not analyze community size per se, the mechanism they evoked to explain this pattern –

i.e., the relative abundance of species with lower fitness become progressively higher – is what has

been suggested as the reason for a major role of ecological drift in small communities (Orrock and

Watling 2010, Gilbert and Levine 2017). Additionally, the effects of environmental changes on

naturally small communities might be combined with the effects of ecological drift in non-obvious

ways, making community dynamics even harder to predict (e.g., Bini et al. 2014) and smaller

communities more vulnerable to novel environmental conditions, such as altered flow regimes

(Tonkin et al. 2018b, Ruhi et al. 2018). This should also be relevant for the conservation and

management of ecosystems. For example, stream restoration efforts tend to be directed towards

sites with low habitat quality and with reduced number of species and low-density populations.

Even if restoration efforts overcome dispersal constraints and barriers to recolonization, which are a

major cause for unsuccessful stream restoration (Bond and Lake 2003, Sundermann et al. 2011,

Tonkin et al. 2014), restored communities will likely be small at the earlier stages of community

assembly and, thus, more prone to ecological drift.

Our study provides empirical evidence of the role that community size can play in mediating

stochastic and deterministic process as drivers of metacommunity dynamics in tropical and boreal

streams, providing one possible solution to a long-standing debate about why some communities

are apparently more influenced by stochastic processes than others. The magnitude of β-diversity

deviation from the null models (negative and positive values) and how it related with community

size (negative and positive slopes) indicate that stochastic and deterministic processes affect species

occurrences and their abundances differently. While incidence-based β-deviation was negative and

decreased with community size in Brazil, abundance-based β-deviation was positive and increased

with community size in both regions. These results indicate that: (1) as communities get larger,

demographic stochasticity plays a less important role and excessive dispersal combined with niche

selection tend to homogenize the genus composition of larger communities; and (2) niche selection

was likely strong enough to produce variations in genera relative abundances, but not to cause

compositional changes among communities within the same watershed. However, we acknowledge

that these are, at least partially, *a posteriori* explanations as we expected that the results would be

similar (i.e., a positive relationship between beta deviation and community size) independently of

the type of coefficient. On the other hand, these contrasting results appear interesting avenues for

future research. Our findings complement previous empirical and conceptual efforts that indicated

that ecological drift can drive biodiversity patterns even in communities where species have clear

differences in life-history traits, resource use and competitive abilities (i.e., those communities

assumed to be non-neutral). Incorporation of community size into ecological models should provide

conceptual, empirical and applied insights towards better understanding of the processes driving

changes in biodiversity.

Acknowledgements. This research originated during a joint research project funded by grant

#2013/50424-1 (São Paulo Research Foundation, FAPESP) to TS, and by grants #273557 and

#273560 to JH and JS (Academy of Finland). ASM and LMB are supported by research grants from

the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq no. 307587/2017-7

and 304314/2014-5, respectively). This study was financed in part by the Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 (scholarship

to DKP). This work was also developed in the context of the National Institutes for Science and

Technology (INCT) in Ecology, Evolution and Biodiversity Conservation, supported by

MCTIC/CNPq (proc. 465610/2014-5) and FAPEG. We thank Christopher Catano and Jonathan

Myers for gently providing R code to generate abundance-based beta deviations, and Mauricio

Vancine for preparing a map of the regions. We thank Eric Harvey, Kevin Cazelles, and Romain

Bertrand for their comments and suggestions on earlier versions of this manuscript, which was

submitted to PCI Ecology.

Author contributions. TS conceived the idea, ran the analyses, and wrote the manuscript with

substantial support from VSS, LMB, ASM, and JH. TS, LMB, VLL, ASM, JS, and JH planned the

sampling design. DKP and JJ-M sampled the data. DKP, VSS and KTT identified the insects. DKP,

JJ-M, KT, VLL and JS commented on the manuscript.

References

Allan, J. D. 2004. Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems. Annual Review of Ecology, Evolution, and Systematics 35:257–284.

Anderson, M. J., T. O. Crist, J. M. Chase, M. Vellend, B. D. Inouye, and A. L. Freestone. 2011. Navigating the multiple meanings of b diversity: a roadmap for the practicing ecologist. Ecology Letters 14:19–28.

Anderson, M. J., K. E. Ellingsen, and B. H. McArdle. 2006. Multivariate dispersion as a measure of beta diversity. Ecology Letters 9:683–693.

Barnes, A. D., M. Jochum, S. Mumme, N. F. Haneda, A. Farajallah, T. H. Widarto, and U. Brose. 2014. Consequences of tropical land use for multitrophic biodiversity and ecosystem functioning. Nature Communications 5:5351.

Bini, L. M., V. L. Landeiro, A. A. Padial, T. Siqueira, and J. Heino. 2014. Nutrient enrichment is related to two facets of beta diversity for stream invertebrates across the United States. Ecology 95:1569–1578.

Bond, N. R., and P. S. Lake. 2003. Local habitat restoration in streams: Constraints on the effectiveness of restoration for stream biota. Ecological Management & Restoration 4:193–198.

Bunn, S. E., and J. M. Hughes. 1997. Dispersal and Recruitment in Streams: Evidence from Genetic Studies. Journal of the North American Benthological Society 16:338–346.

Catano, C. P., T. L. Dickson, and J. A. Myers. 2017. Dispersal and neutral sampling mediate contingent effects of disturbance on plant beta-diversity: a meta-analysis. Ecology Letters 20:347–356.

Chase, J. M., N. J. B. Kraft, K. G. Smith, M. Vellend, and B. D. Inouye. 2011. Using null models to disentangle variation in community dissimilarity from variation in α-diversity. Ecosphere 2:art24.

Chase, J. M., B. J. McGill, D. J. McGlinn, F. May, S. A. Blowes, X. Xiao, T. M. Knight, O.

Purschke, and N. J. Gotelli. 2018. Embracing scale-dependence to achieve a deeper

understanding of biodiversity and its change across communities. Ecology Letters 21:1737–

1751.

Chase, J. M., and J. A. Myers. 2011. Disentangling the importance of ecological niches from

stochastic processes across scales. Philosophical Transactions of the Royal Society B:

Biological Sciences 366:2351–2363.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. Ecology Letters 8:1175–1182.

Durães, L., F. O. Roque, T. Siqueira, A. M. Santos, M. A. Borges, and R. S. Rezende. 2016.

Simulating the role of connectivity in shaping stream insect metacommunities under

colonization cycle dynamics. Ecological Modelling 334:19–26.

Flenner, I., and G. Sahln. 2008. Dragonfly community re-organisation in boreal forest lakes: rapid

species turnover driven by climate change? Insect Conservation and Diversity 1:169–179.

Germain, R. M., S. Y. Strauss, and B. Gilbert. 2017. Experimental dispersal reveals characteristic

scales of biodiversity in a natural landscape. Proceedings of the National Academy of

Sciences 114:4447–4452.

Gilbert, B., and J. M. Levine. 2017. Ecological drift and the distribution of species diversity.

Proceedings of the Royal Society B: Biological Sciences 284:20170507.

Hallmann, C. A., M. Sorg, E. Jongejans, H. Siepel, N. Hofland, H. Schwan, W. Stenmans, A.

Müller, H. Sumser, T. Hörren, D. Goulson, and H. de Kroon. 2017. More than 75 percent

decline over 27 years in total flying insect biomass in protected areas. PLOS ONE

12:e0185809.

Hawkins, C. P., H. Mykrä, J. Oksanen, and V. L. J. J. 2015. Environmental disturbance can increase

beta diversity of stream macroinvertebrate assemblages. Global Ecology and Biogeography

24:483–494.

Heino, J., A. S. Melo, and L. M. Bini. 2015a. Reconceptualising the beta diversity-environmental

heterogeneity relationship in running water systems. Freshwater Biology 60:223–235.

Heino, J., A. S. Melo, J. Jyrkänkallio-Mikkola, D. K. Petsch, V. S. Saito, K. T. Tolonen, L. M. Bini,

V. L. Landeiro, T. S. F. Silva, V. Pajunen, J. Soininen, and T. Siqueira. 2018. Subtropical

streams harbour higher genus richness and lower abundance of insects compared to boreal

streams, but scale matters. Journal of Biogeography 45:1983–1993.

Heino, J., A. S. Melo, T. Siqueira, J. Soininen, S. Valanko, and L. M. Bini. 2015b. Metacommunity

organisation, spatial extent and dispersal in aquatic systems: patterns, processes and

prospects. Freshwater Biology 60:845–869.

Hughes, J. M. 2007. Constraints on recovery: using molecular methods to study connectivity of

aquatic biota in rivers and streams. Freshwater Biology 52:616–631.

Hynes, H. B. N. 1975. The stream and its valley. Internationale Vereinigung Fur Theoretische und

Angewandte Limnologie, Verhandlungen 19:1–15.

Kraft, N. J. B., L. S. Comita, J. M. Chase, N. J. Sanders, N. G. Swenson, T. O. Crist, J. C. Stegen,

M. Vellend, B. Boyle, M. J. Anderson, H. V. Cornell, K. F. Davies, A. L. Freestone, B. D.

Inouye, S. P. Harrison, and J. A. Myers. 2011. Disentangling the Drivers of β Diversity

Along Latitudinal and Elevational Gradients. Science 333:1755–1758.

Lancaster, J., and B. J. Downes. 2017. A landscape-scale field experiment reveals the importance of

dispersal in a resource-limited metacommunity. Ecology 98:565–575.

Leibold, M. A., and J. M. Chase. 2018. Metacommunity Ecology. Princeton University Press,

Princeton, NJ.

Lister, B. C., and A. Garcia. 2018. Climate-driven declines in arthropod abundance restructure a

rainforest food web. Proceedings of the National Academy of Sciences 115:E10397–

E10406.

Macneale, K. H., B. L. Peckarsky, and G. E. Likens. 2005. Stable isotopes identify dispersal

patterns of stonefly populations living along stream corridors. Freshwater Biology 50:1117–

1130.

Mouquet, N., Y. Lagadeuc, V. Devictor, L. Doyen, A. Duputié, D. Eveillard, D. Faure, E. Garnier,

O. Gimenez, P. Huneman, F. Jabot, P. Jarne, D. Joly, R. Julliard, S. Kéfi, G. J. Kergoat, S.

Lavorel, L. Le Gall, L. Meslin, S. Morand, X. Morin, H. Morlon, G. Pinay, R. Pradel, F. M.

Schurr, W. Thuiller, and M. Loreau. 2015. REVIEW: Predictive ecology in a changing

world. Journal of Applied Ecology 52:1293–1310.

Mouquet, N., and M. Loreau. 2003. Community Patterns in Source-Sink Metacommunities. The

American Naturalist 162:544–557.

Muehlbauer, J. D., S. F. Collins, M. W. Doyle, and K. Tockner. 2014. How wide is a stream?

Spatial extent of the potential "stream signature" in terrestrial food webs using meta-

analysis. Ecology 95:44–55.

Myers, J. A., J. M. Chase, R. M. Crandall, and I. Jiménez. 2015. Disturbance alters beta-diversity

but not the relative importance of community assembly mechanisms. Journal of Ecology

103:1291–1299.

Myers, J. A., J. M. Chase, I. Jiménez, P. M. Jørgensen, A. Araujo-Murakami, N. Paniagua-

Zambrana, and R. Seidel. 2013. Beta-diversity in temperate and tropical forests reflects

dissimilar mechanisms of community assembly. Ecology Letters 16:151–157.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B.

O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2018.

vegan: Community Ecology Package.

Orrock, J. L., and R. J. Fletcher Jr. 2005. Changes in Community Size Affect the Outcome of

Competition. The American Naturalist 166:107–111.

Orrock, J. L., and J. I. Watling. 2010. Local community size mediates ecological drift and

competition in metacommunities. Proceedings of the Royal Society B: Biological Sciences

277:2185–2191.

Petsch, D. K., G. D. Pinha, J. D. Dias, and A. M. Takeda. 2015. Temporal nestedness in

Chironomidae and the importance of environmental and spatial factors in species rarity.

Hydrobiologia 745:181–193.

Roque, F. O., T. Siqueira, L. M. Bini, M. C. Ribeiro, L. R. Tambosi, G. Ciocheti, and S. Trivinho-

Strixino. 2010. Untangling associations between chironomid taxa in Neotropical streams

using local and landscape filters. Freshwater Biology 55:847–865.

Ruhi, A., X. Dong, C. H. McDaniel, D. P. Batzer, and J. L. Sabo. 2018. Detrimental effects of a

novel flow regime on the functional trajectory of an aquatic invertebrate metacommunity.

Global Change Biology 24:3749–3765.

Saito, V. S., M. V. Cianciaruso, T. Siqueira, A. A. Fonseca-Gessner, and S. Pavoine. 2016.

Phylogenies and traits provide distinct insights about the historical and contemporary

assembly of aquatic insect communities. Ecology and Evolution 6:2925–2937.

Saito, V. S., J. Soininen, A. A. Fonseca-Gessner, and T. Siqueira. 2015. Dispersal traits drive the

phylogenetic distance decay of similarity in Neotropical stream metacommunities. Journal

of Biogeography 42:2101–2111.

Siqueira, T., L. D. Durães, and F. de O. Roque. 2014. Predictive Modelling of Insect

Metacommunities in Biomonitoring of Aquatic Networks. Pages 109–126 *in* C. P. Ferreira

and W. A. C. Godoy, editors. Ecological Modelling Applied to Entomology. Springer

International Publishing.

Siqueira, T., C. G.-L. T. Lacerda, and V. S. Saito. 2015. How Does Landscape Modification Induce

Biological Homogenization in Tropical Stream Metacommunities? Biotropica 47:509–516.

Sundermann, A., S. Stoll, and P. Haase. 2011. River restoration success depends on the species pool

of the immediate surroundings. Ecological Applications 21:1962–1971.

Swan, C. M., and B. L. Brown. 2017. Metacommunity theory meets restoration: isolation may
mediate how ecological communities respond to stream restoration. Ecological Applications
27:2209–2219.

Tonkin, J. D., F. Altermatt, D. S. Finn, J. Heino, J. D. Olden, S. U. Pauls, and David. A. Lytle.
2018a. The role of dispersal in river network metacommunities: Patterns, processes, and
pathways. Freshwater Biology 63:141–163.

Tonkin, J. D., D. M. Merritt, J. D. Olden, L. V. Reynolds, and D. A. Lytle. 2018b. Flow regime
alteration degrades ecological networks in riparian ecosystems. Nature Ecology & Evolution
2:86.

Tonkin, J. D., S. Stoll, A. Sundermann, and P. Haase. 2014. Dispersal distance and the pool of taxa,
but not barriers, determine the colonisation of restored river reaches by benthic
invertebrates. Freshwater Biology 59:1843–1855.

Valente-Neto, F., L. Durães, T. Siqueira, and F. O. Roque. 2017. Metacommunity detectives:
Confronting models based on niche and stochastic assembly scenarios with empirical data
from a tropical stream network. Freshwater Biology 63:86–99.

Vásquez, D., R. W. Flowers, and M. Springer. 2009. Life history of five small minnow mayflies
(Ephemeroptera: Baetidae) in a small tropical stream on the Caribbean slope of Costa Rica.
Aquatic Insects 31:319–332.

Vellend, M. 2010. Conceptual synthesis in community ecology. The Quarterly review of biology
85:183–206.

Vellend, M. 2016. The Theory of Ecological Communities. Mpb Series: 57 ed. edition. Princeton
University Press, Princeton.

Vellend, M., D. S. Srivastava, K. M. Anderson, C. D. Brown, J. E. Jankowski, E. J. Kleynhans, N.
J. B. Kraft, A. D. Letaw, A. A. M. Macdonald, J. E. Maclean, I. H. Myers-Smith, A. R.
Norris, and X. Xue. 2014. Assessing the relative importance of neutral stochasticity in
ecological communities. Oikos 123:1420–1430.

633   Wallace, J. B., and N. H. Anderson. 1996. Habitat, life history, and behavioral adaptations of

634   aquatic insects. Pages 41–73 An Introduction to the Aquatic Insects of North America. Third

635   edition. Kendall/Hunt Publishing Company, Dubuque, Iowa.

636   Winegardner, A. K., B. K. Jones, I. S. Y. Ng, T. Siqueira, and K. Cottenie. 2012. The terminology

637   of metacommunity ecology. Trends in Ecology & Evolution 27:253–254.

Supplementary material

660   **Figure S1.** Graphical representation describing the general procedures for estimating beta diversity

661   (β), community size (CS), spatial extend (SE) and environmental heterogeneity (EH) within

662   watersheds (*n* = 20) in Brazil and Finland.

Basic model tested: β = *f* (CS,SE,EH)

664   * This was repeated for different dissimilaity coefficients, and for null model analyses

674 **Figure S2.** Geographical location and spatial distribution of streams in São Paulo State, Brazil

675 (circles in vermilion) and Finland (circles in blue).