Extensive splicing across the Saccharomyces cerevisiae genome

Stephen M. Douglass[1*], Calvin S. Leung[2], and Tracy L. Johnson[1,2*]

[1]Department of Molecular, Cell, and Developmental Biology, University of California, Los Angeles, Los

Angeles, CA 90095, USA

[2]Molecular Biology Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA

Correspondence: smdougla@ucla.edu, tljohnson@ucla.edu

Abstract

Pre-mRNA splicing is vital for the proper function and regulation of eukaryotic gene expression. *Saccharomyces cerevisiae* has been used as a model organism for studies of RNA splicing because of the striking conservation of the spliceosome components and its catalytic activity. Nonetheless, there are relatively few annotated alternative splice forms, particularly when compared to higher eukaryotes. Here, we describe a method to combine large scale RNA sequencing data to accurately discover novel splice isoforms in *Saccharomyces cerevisiae*. Using our method, we find extensive evidence for novel splicing of annotated intron-containing genes as well as genes without previously annotated introns and splicing of transcripts that are antisense to annotated genes. By incorporating several mutant strains at varied temperatures, we find conditions which lead to differences in alternative splice form usage. Despite this, every class and category of alternative splicing we find in our datasets is found, often at lower frequency, in wildtype cells under normal growth conditions. Together, these findings show that there is widespread splicing in *Saccharomyces cerevisiae*, thus expanding our view of the regulatory potential of RNA splicing in yeast.

Author Summary

Pre-mRNA splicing is a fundamental step in eukaryotic gene expression. *Saccharomyces cerevisiae*, also known as brewer's yeast, is a model organism for the study of pre-mRNA splicing in eukaryotes. Through the process of pre-mRNA splicing, a single gene is capable of encoding multiple mature mRNA products, but it is often difficult to identify the splice events that lead to these mRNA products. Here, we describe a method to accurately discover novel splice events in *Saccharomyces cerevisiae* and find evidence for extensive splicing in *Saccharomyces*. By utilizing a variety of strains and growth conditions, we are able to characterize many splice forms and correlate cellular conditions with prevalence of novel splice events.

Introduction

Eukaryotic genes are composed of coding sequences termed exons interrupted by non-coding sequences called introns. Introns are removed from RNA by the large macromolecular complex known as the spliceosome through the process of RNA splicing. By selectively including different combinations of exons, a single gene can produce multiple RNA products. Pre-mRNA splicing is crucial for the proper expression of eukaryotic genes, and spliceosome components are highly conserved from yeast to mammals at the sequence, structure, and functional levels [1, 2].

Despite the high conservation between the spliceosomal components in yeast and mammals, *Saccharomyces cerevisiae* has a more compact genome with approximately 300 annotated intron-containing genes. Even with the relatively streamlined splicing landscape in *S. cerevisiae*, the prevalence of alternative splicing in this organism has not been well characterized. Several studies have found instances of alternative splicing in individual genes in *S. cerevisiae* [3, 4, 5], and there have been high-throughput methods to find novel splicing [6,

7, 8, 9, 10]. Most of these methods focus primarily or entirely on intron retention or exon
skipping, while there has been little description of novel 5' or 3' splice site usage.

Here we describe a method for discovery of unannotated splice sites in *Saccharomyces*
cerevisiae by RNA-seq. We tested the method with wildtype strains as well as strains and
conditions that do not lead to direct changes in the splicing machinery itself, but that impact
broad cellular conditions, RNA turnover, chromatin remodeling, and histone composition. Many
of these strains and conditions help us observe changes in splicing that are not or are rarely
detected in wildtype cells under normal growth conditions. We also analyze several strains that
include *prp43-1*, a temperature sensitive mutation of the *prp43* gene, an RNA helicase directly
involved in spliceosome disassembly to determine if modulating disassembly might affect our
ability to detect aberrant splice site sampling. Interestingly, this mutation of *prp43* leads to a
decrease in splicing of both annotated introns as well as novel splicing.

We compare our results with those of earlier studies and find good agreement, however over
two thirds of the splice forms our method predicts, 676 total novel splice forms, have not been
previously described. Within known intron-containing genes (ICGs), we find that novel isoforms
generate longer introns in samples grown at higher temperature. We also find that deletion of
xrn1, an evolutionarily-conserved 5' to 3' exonuclease, leads to an accumulation of novel splice
isoforms that do not use either an annotated 5' nor an annotated 3' splice site, both within
known ICGs and unannotated ICGs. While most of our novel splice forms use a known 5' splice
site and a novel 3' splice site, we find that deletion of *ume6*, a component of a histone
deacetylase complex that regulates early meiosis and that has been recently shown to have a
regulatory role during mitosis [11], causes a dramatic increase in the use of novel 5' splice sites
within ICGs. We also find evidence for splicing of RNAs that are antisense to annotated
transcripts. Together, our results indicate that there is significantly more splicing than previously
thought. This suggests that the opportunity for splicing in the form of "latent" introns is a key
feature of the yeast genome. Furthermore, changes in the activity of the gene expression
machinery or the cells' environment can significantly alter this rich splicing landscape.

Results

Alternative splicing is widespread in *Saccharomyces cerevisiae*

In order to explore the extent of splicing across the yeast transcriptome in a high confidence
manner, we implemented a novel approach to allow us to leverage a large amount of RNA
sequence data while imposing stringent filters (figure 1). Briefly, a large number of RNA-seq
datasets are combined such that each novel splice form needs at least a single read that aligns
without mismatches to the novel junction to pass the initial filter. This read cannot also align to
known transcripts or the *Saccharomyces cerevisiae* genome, even with a large number of
mismatches. Once a splice junction has been identified in this way, all reads are realigned to
the newly discovered splice forms, annotated transcripts, and the genome to find the optimum
alignment. Each novel splice form is then scored based on how well its 5' splice site, 3' splice
site, and predicted branch point fit the consensus sequence for *Saccharomyces cerevisiae*
splice signals. These scores are then used to compute p-values that represent how likely a
splice product score of this strength is to occur by chance (figure S1).

To capture as many novel splice forms as possible, we incorporated datasets from a variety of
strains and experimental conditions that are known or suspected to have an impact of splicing,
some accessed from previous studies [12, 13] and some novel (Table S1). In addition to 4
wildtype samples grown at three different temperatures, these samples include 2 biological
replicates each of two deletions in genes involved in decay, including *xrn1*, a 5'-3' exonuclease,
and *upf1*, required for efficient nonsense-mediated decay. We also include *htz1Δ* which
encodes the histone variant H2A.Z and *swr1Δ* which is required to exchange H2A for H2A.Z in
chromatin as well as double mutants *swr1Δ xrn1Δ*, *swr1Δ upf1Δ*, *htz1Δ xrn1Δ*, and *htz1Δ upf1Δ*.
Cells that lack H2A.Z are found to have impaired splicing of intron-containing genes (ICGs),
particularly genes that have suboptimal splice sites [13, 14]. We also analyze RNA-seq data
from *snf2Δ,* which leads to an increase in use of non-canonical branch point and 5' splice site
sequences in annotated ICGs and *ume6Δ*, which derepresses genes implicated in meiosis in
Saccharomyces cerevisiae, and the double mutants *upf1Δ snf2Δ*, *xrn1Δ snf2Δ*, and *ume6Δ*
snf2Δ. Previous studies suggest that *snf2Δ* increases splicing by altering ribosomal protein
gene expression and *ume6Δ* allows expression of genes that are usually repressed. We also
include *set1Δ* and *set2Δ*, deletion mutations in histone methyltransferase genes. Finally, since
we previously showed that the DEAH protein Prp43 contributes to disassembly of suboptimal
spliceosomes using a *prp43* DAMP allele [13], we included a temperature sensitive mutant of
prp43, *prp43-1*, as well as *set1Δ prp43-1* and *set2Δ prp43-1*. All *set1Δ*, *set2Δ*, and *prp43-1*
strains contribute two samples to our workflow, one grown at 25° and one at 37°. Taken
together, these datasets represent 29 samples across a variety of *Saccharomyces cerevisiae*
strains and growth temperatures. By leveraging a large number of datasets, we are able to
discover more novel splice sites and determine cellular conditions that lead to changes in
alternative splice site usage.

Strain	Read counts	Aligned read counts
Wildtype	20395478	14160763
xrn1Δ	36132564	25787231
upf1Δ	32005528	22034981
swr1Δ	38830226	21964031
htz1Δ	35969862	18508451
swr1Δ xrn1Δ	86744422	50807249
swr1Δ upf1Δ	11369368	6729660
htz1Δ xrn1Δ	52891892	35039812
htz1Δ upf1Δ	49460562	25424884
Wildtype	35059908	27722481
snf2Δ	34016283	25657587
upf1Δ	21328241	14465110
upf1Δ snf2Δ	31564582	23487340
xrn1Δ	29688304	20282088

xrn1Δ snf2Δ	34289534	27043467
ume6Δ	40071056	30161996
ume6Δ snf2Δ	22534282	17424922
Wildtype 25° C	126355512	97015734
Wildtype 37° C	140962504	82278728
set1Δ 25° C	120413012	75538896
set1Δ 37° C	150376422	84044314
set2Δ 25° C	115930956	61396568
set2Δ 37° C	105592626	63182213
prp43-1 25° C	113138574	78671916
prp43-1 37° C	93850396	54354046
prp43-1 set1Δ 25° C	180103858	107502423
prp43-1 set1Δ 37° C	116668118	70474357
prp43-1 set2Δ 25° C	129498126	93547450
prp43-1 set2Δ 37° C	107964066	68127119

Using our method, we discover evidence for 944 novel splice events across 408 transcripts with
p < 0.05 (Table 1). Of these novel events, the majority are novel splice products found within
known ICGs, using either the annotated 5' splice site with a novel 3' splice site (Table S2,
n=537) or using the annotated 3' splice site with a novel 5' splice site (Table S2, n=129).
Additionally, we found several cases of novel splicing that do not use any annotated splice sites
within annotated ICGs (Table S2, n=22) and in unannotated ICGs (Table S2, n=198). We also
find evidence for splice forms that are antisense to known transcripts (Table S2, n=50).

	All		p<0.05		p<0.01		p<0.001	
	Events	Transcripts	Events	Transcripts	Events	Transcripts	Events	Transc
All	6187	1937	944	408	852	386	586	
Antisense	2504	1110	50	41	46	37	33	
Intronless	2709	1141	198	149	168	132	107	
New 3'	561	196	537	194	502	191	368	
New 5'	140	95	129	90	113	84	63	
New both	273	120	22	18	15	12	7	

Many of the events we discover using our method are low abundance. About half of the total
events are represented by fewer than five read counts across our datasets (figure S2A).
However, these low abundance events are high confidence due to our approach's stringent
discovery protocol and p-value cutoff. Furthermore, these high confidence, low abundance
novel splice events reveal splicing that is unlikely to be found in the high abundance data.
Different classes of novel splicing are more prevalent in our low abundance novel splicing data.
Specifically, novel splicing within annotated ICGs is more common in our high abundance data,
while novel splicing within unannotated ICGs and antisense RNAs are more common in our low
abundance data (figure S2C).

Validation of novel splicing

In order to best validate our method by RT-PCR, we chose candidates with a variety of read
counts and splice site scores representing each category of novel splicing we find, including
novel 5' splice site with annotated 3' splice site, novel 3' splice site with annotated 5' splice site,
both novel splice sites within an annotated ICG, novel splicing in an unannotated ICG, and
splicing of transcripts that are antisense to annotated genes. In addition, we selected some
genes which show only a single novel splice form and others with several. The results of our
validation are shown in Figure 2 using oligonucleotides listed in table S3. *BIG1*, an integral
membrane protein gene of the endoplasmic reticulum, shows a single novel splice event
utilizing a novel 3' splice site with an annotated 5' splice site in our data with 19 total read
counts spread roughly evenly across our 29 samples and a splice site score p-value of 0.0003.
The novel splice event in *BIG1* is in-frame with the annotated form, and would be predicted to
generate a protein product that has 6 additional amino acids. *SIM1*, a gene thought to
participate in DNA replication, shows a total of four novel splice forms, each utilizing an
annotated 3' splice site and novel 5' splice site with total read counts ranging from 2 to 35 and
splice site p-values ranging from 0.018 to 0.0004. The intron in *SIM1* is located in the 5' UTR,
therefore the novel splice events would not be expected to yield a new protein product, although
potential regulation in the 5' UTR could be altered. *MCR1*, a gene involved in ergosterol
biosynthesis in mitochondria, shows a single novel splice form utilizing both novel 5' and 3'
splice sites within an annotated ICG with 49 total read counts, 9 of which are found across our
snf2Δ samples, and a splice site p-value of 0.004. The annotated and novel introns for *MCR1*
are both found in the 5' UTR and are therefore unlikely to yield different protein products.
However, we previously showed that changes in *SNF2* expression can affect splicing of others
transcripts to alter mitochondrial function [15]. *SPF1*, which encodes an ion transporter of the
ER membrane, is an unannotated ICG that shows five novel splice forms in our data with read
counts ranging from 1 to 4 across all of our samples and p-value scores ranging from 0.004 to 6
x 10^{-6}. Of the five novel splice forms found in *SPF1*, only one is in frame, which would produce
a protein product that has 134 fewer amino acids. Finally, we find evidence for splicing in the
antisense direction to *LEU4*, with a total of six splice forms with counts ranging from 1 to 33
across all samples and splice site p-values ranging from 0.0001 to 2.8 x 10^{-6}. Not surprisingly,
of the 48 sequence reads that derive from spliced reads antisense to *LEU4*, 28 come from
samples which are deleted of *SET2*, a histone methyltransferase that has previously been
implicated in suppressing antisense transcription [16]. Together, we validate each category of
novel splicing we observe in our data and validate candidates ranging from a single read count
to 49 read counts and from splice site p-values ranging from 0.018 to 2.8 x 10^{-6}.

Primer	Sequence
BIG1 F	GTTGCTTATTATGTGTGGAAGCTTTTG
BIG1 R	CACCTGGTCTACGTTACAATACTCC
SPF1 F	CCCAGAGA GCCACAAGTT GATCTTG
SPF1 R	CATTTAGAGGCAATCTTGACCTGTTGC
SIM1 F	GTGCTA CCCAACTACT TACATTCCT
SIM1 R	CTTGGCTAAGGCAGCAGATGAAAC
MCR1 F	CCTTGATTGGTGTCTTGTCGAGAGAG
MCR1 R	ACTCACAGGAGTGTATGGTCTCACC
LEU4 antisense	GAGAGTATTATTGCTCTTGCTGAGC
LEU4 R	CTTGTTGGGAAGGGTCATCCTTAG

Comparison with other sequence-based approaches for alternative splicing

Even though our method filters (1) putative splice forms with strong similarity to the
Saccharomyces cerevisiae genome, (2) known transcripts, and (3) those found outside of
genes, the splice sites identified by our method include many reported in recent studies. Of the
522 novel splice sites described in Kawashima et al. [6], 420 are discoverable by our method,
236 are found in our raw data and 189 pass our $p < 0.05$ filter. Of those described in Schreiber
et al. [7], 248 out of 314 of the described splice events are discoverable by our method, 214 are
in our data and 185 pass our $p < 0.05$ filter. Qin et al. [10] describe a method of identification of
novel splice forms by lariat sequencing, a process which reveals 5' splice sites and branch
points but not 3' splice sites. Of the 45 novel 5' splice sites found in their work, 11 are present
in our raw data and 9 pass our $p < 0.05$ filter with at least one corresponding 3' splice site.
Gould et al. [9] combined lariat sequencing with RNA-seq to identify both 5' and 3' splice sites
along with novel branch point sequences. Of the 213 novel splice sites they report, 194 are
discoverable by our method, 133 are found in our raw data, and 114 pass our $p < 0.05$ filter.
The overlap between our work and these previous studies illustrates the power of our approach.
None of the splice sites described by Qin et al. are found in any of the three other studies, while
each of Kawashima et al., Gould et al., and Schreiber et al. have greater overlap with our work
than with one another (figure 3). Taken together, of the 944 splice junctions predicted with $p <$
0.05 by our method, 268 have been previously described and 676 are novel.

Saccharomyces cerevisiae contains antisense transcripts that undergo splicing

In addition to discovering novel splice sites within genes already known to contain introns and
unannotated ICGs, our method allows us to discover splicing that is antisense to annotated
transcripts, primarily in the degradation mutant strains *xrn1Δ* and *upf1Δ* and the histone
methyltransferase mutant *set2Δ*. Previous studies report that Set2 suppresses antisense
transcription [16]. We find evidence for 50 antisense novel splice events spread across 41
transcripts that pass our statistical criteria. Of these 41 transcripts, 37 show a single novel
splice event, two transcripts have two unique novel splice events, one transcript has 3 separate
events, and a single transcript, which is antisense to the *LEU4* gene, has 6 unique novel splice
events. While most of the antisense splice events in our data have low read counts, the *LEU4*
isoforms together account for 51 reads across 11 samples, including wildtype, *set2Δ*, and *set2Δ*

prp43-1 at both 25° and 37°, *set1Δ* and *set1Δ prp43-1* at 37° only, and *xrn1Δ*, *swr1Δ upf1Δ*, and
upf1Δ snf2Δ. Together, *LEU4* antisense splicing represents over 20% of our total antisense
read counts across all samples. These six isoforms arise from 3 different 5' splice sites and 4 3'
splice sites. 5 out of 6 of these isoforms generate similar mature mRNAs with an intron in the
size range from 115 nucleotides to 129 nucleotides and are therefore indistinguishable by RT-
PCR (figure 2E). The remaining form is generated from a unique 5' splice and 3' splice and
causes an intron of 464 nucleotides, and is low abundance in our RNA sequence data, with only
a single read count in a single sample, *set2Δ* at 37°.

In unannotated ICGs, we find 149 transcripts that show novel splicing in the sense direction and
37 that undergo novel splicing in the antisense direction. Interestingly, the number of transcripts
that show novel splicing in both the sense and antisense direction is just one, *DJP1*. If a set of
149 genes and a set of 37 genes are each randomly chosen from all *Saccharomyces cerevisiae*
transcripts, the expected value of the overlap is a single transcript, suggesting that within
unannotated ICGs, presence of novel splicing in the sense direction does not significantly
impact the chances of novel spicing of an antisense transcript or vice versa. Within annotated
ICGs, 220 transcripts undergo novel splicing in the sense direction and 4 have novel splicing in
the antisense direction, indicating no correlation between the splicing of annotated intron-
containing genes and their corresponding antisense transcripts.

Prp43 is required for efficient splicing of annotated and novel introns

Several of the strains in our analysis include *prp43-1*, a temperature sensitive mutation that is
viable at 25° C but not at 37° C. Prp43 is an RNA helicase that has a vital role in spliceosome
disassembly and is required for efficient mRNA splicing in *Saccharomyces cerevisiae*. *PRP43*
has also been implicated in ribosome biogenesis [17], and we previously showed that
decreasing levels of *PRP43* using a DAMP allele can suppress splice defects [13]. To
characterize the consequences of the *prp43-1* mutation in splicing, we compared the *prp43-1*
strain to a wildtype strain, a *set1Δ prp43-1* strain to a *set1Δ* strain, and a *set2Δ prp43-1* strain to
a *set2Δ* strain at both 25° and 37° C. As expected from its role in splicing, *prp43-1* shows a
decrease in the splicing of annotated introns in *Saccharomyces cerevisiae* (figure 4A).
Interestingly, despite earlier findings that reducing levels of wildtype Prp43 can suppress splice
defects and promote splicing of weak introns [13], we find that strains with *prp43-1* show less
novel splicing than their counterparts with wildtype *PRP43* (figure 4B).

Elevated temperature favors novel introns that are longer than their annotated introns

Across our datasets, we detect fewer novel splice events in high temperature samples than
would be expected by sequence depth alone. This is unsurprising given that many of our lower
temperature samples are mutants that lead to accumulation of normally degraded products,
such as *xrn1Δ* and *upf1Δ*. While the total number of novel splice events is underrepresented at
high temperature, the novel splice products generated differ in intron length. When cells are
grown at 37° C, novel splicing within annotated ICGs tends to favor intron sizes that are larger
than novel splice forms from cells that are grown at 25° or 30° C (figure 5A). This can be
explained by effects observed in our most common class of novel splice events, those that use

an annotated 5' splice site with a novel 3' splice site. At higher temperature, these novel 3'
splice sites tend to be further downstream than the annotated splice sites. We find that most of
the temperature-enriched splice events are more commonly found in our samples with wildtype
prp43, since *prp43-1* decreases both annotated and novel splicing (figure 5B). These results
are consistent with work that finds that intron structure can function to control alternative splicing
in yeast [18]. Nonetheless, our data reveal a number of previously unreported events.

As an example, *TMH18*, a mitochondrial membrane protein gene, has an annotated intron that
is 96 nucleotides long and two common novel splice isoforms discovered by our method, both of
which generate longer introns. The isoform that is highly enriched in high temperature samples
contains a 161 nucleotide intron, while the isoform that is not favored at high temperature
contains a 128 nucleotide intron. We speculate that the increase in temperature destabilizes
the secondary structure of some pre-mRNA molecules to allow access to normally inaccessible
splice sites. This may lead to an increase in use of distant novel splice sites. To confirm this,
we analyzed the predicted pre-mRNA secondary structure using MFOLD [19]. The predicted
secondary structure with the most favorable free energy shows that the annotated 5' splice site,
the annotated 3' splice site, and the novel 3' splice site that is not favored at high temperature
are all readily accessible, while the novel 3' splice site that is favored only at high temperature is
not (figure 5C).

Interestingly, these distant splice sites may be under evolutionary pressure to be unable to code
for protein, as 74/84 (88%) of these temperature-enriched splice sites contain premature
termination codons. This is similar to the fraction of PTC-generating events found in the totality
of our novel splicing events in annotated intron-containing genes, with 572/636 (90%) containing
premature termination codons. This agrees with results described by Kawashima et al. [6] that
find that stress conditions, including heat shock, cause an increase in non-productive novel
splice usage. Together these data suggest that alternative splicing may be a mechanism for
reconfiguring the transcriptome in response to stress.

Xrn1 deletion increases splice forms that do not use annotated splice sites

Novel splicing is found most commonly in strains in which *xrn1* has been deleted. These strains
account for approximately 9% of our total sequence depth but 19% of our total novel splice
counts. Interestingly, *xrn1* deletion mutants are particularly enriched in novel splice forms that
do not use any annotated splice sites. Roughly 18% of novel splice form counts that use either
an annotated 3' or 5' splice site derive from an *xrn1* deletion mutant, while 31% of those that
utilize two novel splices are found in an *xrn1* deletion strain. Two examples of unannotated
ICGs impacted strongly by *xrn1* deletion are *AGC1* and *MRM2*, both involved in mitochondrial
function. Interestingly, these genes' expression increases in *xrn1Δ*, and they each have their
highest RPKM values in the five samples in which *xrn1* is deleted [12, 13]. The 50 unannotated
ICGs that are most enriched in our *xrn1Δ* strains only have a single GO term in common,
"intracellular membrane-bounded organelle," further suggesting an impact on transcripts
important for mitochondrial function. Interestingly, a 2012 study found that Xrn1 is critical for the
translation of genes necessary for mitochondrial function in *Saccharomyces cerevisiae* [20].
Together, this suggests a role for Xrn1 in the regulation of alternative splice products of
mitochondrial transcripts.

Ume6Δ-derived increase in IC-RPG expression leads to increase in IC-RPG alternative splicing?

Our *ume6Δ* strain is highly enriched in novel splice events which utilize an annotated 3' splice site with a novel 5' splice site. Of the newly discovered splice sites which utilize a novel 5' splice site and a canonical 3' splice site, 59% are in intron-containing ribosomal protein genes (IC-RPGs). However, of the events that are found disproportionately more frequently in *ume6Δ*, 70% are in IC-RPGs. Our RNA sequence data suggests that IC-RPG expression increases slightly in *ume6Δ*. Novel splicing in IC-RPGs increases by 70%, and 5' novel IC-RPG splicing increases by 91% in *ume6Δ*-containing samples relative to samples with wildtype *ume6*.

Previous studies have shown that Ume6 is degraded under meiotic conditions [21]. This raises the interesting possibility that genetic manipulations that remove Ume6 may lead to changes in the unannotated splicing landscape that are related to what occurs under meiotic conditions. Ongoing experiments will test this model.

Discussion

In this study we show that there are many rarely utilized splice sites in *Saccharomyces cerevisiae*. Our methodology is capable of discovering many novel splice forms by utilizing a large number of RNA-seq datasets. We are able to do this in a robust, high-confidence manner by excluding reads that can be explained by sequencing error or genomic DNA contamination and filtering based on existing splice site census sequence data. Overall, we find a strong preference for novel splicing using a known 5' splice site and a novel 3' splice site within annotated ICGs, which represent over half of our statistically significant novel splice products. However, due to our high total sequencing depth and variety of strains and experimental conditions we are still able to find a relatively large number of novel splice forms that utilize known 3' splice sites with novel 5' splice sites, those that utilize novel 3' and 5' splice sites within annotated ICGs, those that utilize two novel splice sites within unannotated ICGs, and splicing of transcripts that are antisense to annotated genes. The large number of novel splice events that we discover allow us to correlate changes in splice site preferences with different mutant strains and experimental conditions. While it has been tempting to call splicing events that have not been previously annotated as "errors," we believe that these data actually reveal the remarkable substrate flexibility of an evolutionarily conserved enzyme. Moreover, it stands to reason that if splicing is to contribute to adaptation and, ultimately, evolution of multicellular organisms, then an array of sequences, not simply those that match a strong consensus, need to be recognized and spliced. The results described here provide a window into this sequence landscape.

In the adjoining manuscript [22], the authors find 229 "protointrons" – rapidly evolving, inefficiently spliced introns. Of these, we define 60 as novel introns with an additional 9 found in our RNA sequence data but filtered out due to low splice site scores. The limited overlap between the methods highlights that neither of our methods has reached saturation. Furthermore, the 10 strains in our analysis with the greatest normalized overlap with Talkish et

al. are the 10 strains that include either *xrn1Δ* or *upf1Δ,* suggesting that overlap between the
methods is more driven by stabilization of protointron-containing splice products than biological
similarity between "hungry spliceosome" conditions and the strains used here. Talkish et al. confirm
by RT-PCR several splice events that would be removed from our analysis due to poor splice
signals, indicating that our method's stringency likely eliminates many true splice products. The
approach used here scores putative novel splice sites by learning from annotated splice sites,
so as additional protointrons with unusual splice sites are discovered and validated, our
method's ability to discover these forms will increase.

The analysis of several mutant strains allows insights into splice site selection in
Saccharomyces cerevisiae. We find general trends in our data as well as specific effects for
particular strains or conditions. For example, elevated temperature leads to an increase in
novel splice form intron length, *xrn1Δ* leads to a large increase in novel splice forms that do not
use an annotated splice site, and *ume6Δ* leads to an increase in novel splicing of intron-
containing ribosomal protein genes.

Despite the fact that analyzing many mutant strains can increase our understanding of splice
site selection, every new category and class of splice form that we find is observed in our
wildtype data. While *xrn1Δ* dramatically increases the number of splice products that arise from
use of two novel splice sites, we see many examples of splice products that use two novel
splice sites in our wildtype samples. This holds true for elevated temperature, *ume6Δ*, and
set2Δ, as well as splicing of transcripts that are antisense to known transcripts, splicing that
uses a single annotated splice site, and splicing in unannotated ICGs. Together, this indicates
that while mutant strains are useful for correlating genetic changes with splicing changes, even
wildtype *Saccharomyces cerevisiae* in normal conditions are capable of producing a large
variety of splice products. Taken together, these findings illustrate the diversity and depth of
splicing in *Saccharomyces cerevisiae*, and also show the presence of latent introns that are
found across the genome.

Methods

Public datasets

376 *Saccharomyces cerevisiae* wildtype, *xrn1Δ*, *upf1Δ*, *swr1Δ*, *htz1Δ*, *swr1Δ xrn1Δ*, *swr1Δ upf1Δ*,
377 *htz1Δ xrn1Δ*, and *htz1Δ upf1Δ* strain RNA-seq data were download from GEO (accession
378 number GSE97416). Additional wildtype, *upf1Δ*, and *xrn1Δ* RNA-seq data as well as *snf2Δ*,
379 *ume6Δ*, *upf1Δ snf2Δ*, *xrn1Δ snf2Δ*, and *ume6Δ snf2Δ* strain RNA-seq data were also downloaded
380 from GEO (accession number GSE94404).

Yeast culture and Sequencing

Wildtype, *set1Δ*, *set2Δ*, *prp43-1*, *set1Δ prp43-1*, and *set2Δ prp43-1* strains were grown at 25°C
to OD 0.3. Then, cultures were equally split, half remaining at 25°C and half shifting to 37°C for
four hours. 10 mL of cells were pelleted from each sample and RNA extraction was performed.
Then, 20 μg total RNA per sample was treated with DNAse I (Roche) and depleted of rRNA with
the Ribo-Zero Gold rRNA Removal Kit (Illumina). RNA-seq libraries were prepared using the
Illumina TruSeq v2 RNA Kit. 50 base pair paired-end reads were generated on an Illumina
HiSeq 4000.

Sequence alignment

RNA sequence data were combined and aligned to the *Saccharomyces cerevisiae* SacCer3
genome reference and Ares Lab Yeast Intron Database Version 3 [23] in a single step using
STAR [24] allowing up to six mismatches and no unannotated gaps. Sequence reads that fail to
align in this step are then aligned to the SacCer3 genome again, this time allowing no
mismatches, a single gap, at most one alignment locus, and at least 15 nucleotide overhang on
each end of the alignment. Successful alignments in this step define putative novel introns. In
a final alignment step, all reads are aligned to the SacCer3 genome, the Ares Intron Database,
and newly defined putative novel introns in one step, allowing at most 1 alignment locus with up
to 2 mismatches. Counts for novel splice forms are based on those derived from this alignment
step.

Splice site scoring

5' splice site, 3' splice site, and branch point scores were generated based on how closely each
splicing signal matches the consensus sequence for that signal. For each position within a
splice site, the total number of adenine, cytosine, guanine, and thymine bases present at that
position in annotated splice signals was determined based on the Ares Lab Intron Database,
and then the proportion of each nucleotide at that position was found by dividing by the total
number of annotated splice products. The score for each splice site was calculated as follows:

$$Score = \sum_{i=1}^{k} N_i / X$$

Where i is the position within the splice signal, k is the number of positions in the splice signal,
N_i is the count of that specific nucleotide at that position, and X is the total counts for the splice
signal. To score each putative novel splice form, the 5' and 3' splice sites are determined from
RNA-seq data, while branch point sequences were chosen by selecting the highest scoring
possible branch point sequence within a maximum distance of 200 nucleotides from the 3'
splice site. The score for each putative novel splice site is simply the product of its 5' splice site
score, its 3' splice site score, and its branch point score. Putative novel splice sites were
considered antisense if the score in the antisense direction is higher than the score in the sense
direction.

Statistical analysis of splice sites

P-values for putative novel splice sites were generated by comparing the potential splice site to all possible combinations of 5' splice sites, 3' splice sites, and branch points. Based on a six nucleotide 5' splice site, a three nucleotide 3' splice site, and a seven nucleotide branch point, there are 4,294,967,296 possible combinations of splice signals that can be described. To convert our putative novel splice form scores to p-values we compared the score to the scores of all possible combinations of splice signals. The raw p-value is the fraction of these combinations that score equal to or better than the putative novel form, which also represents the chance of seeing a splice score as good or better than this by chance. We then correct the raw p-values for multiple hypothesis testing using the Bonferroni correction by multiplying each raw p-value by the number of tests conducted, which is the total number of putative novel splice sites times two, since each splice site score is calculated in both the sense and antisense direction. All adjusted p-values greater than one are then set to one.

Splicing Efficiency Calculation

To calculate the splicing efficiency of annotated splice sites, reads were aligned with STAR [24] allowing only a single alignment locus, only annotated splice sites, and at most two mismatches. Aligned reads within ICGs were categorized as exonic, spliced, or unspliced. We generated normalized spliced and unspliced read counts by dividing the raw counts in each category by the number of possible alignments that can fall into that category. This equates to read length minus one for spliced reads and the intron length plus the read length minus one for unspliced reads. Splicing efficiency is then computed as normalized spliced counts divided by the sum of normalized spliced and normalized unspliced counts.

RNA Folding

To find the optimum secondary structure for TMH18, we used the MFOLD web server with the pre-mRNA sequence accessed from the Saccharomyces Genome Database (https://www.yeastgenome.org/) and default MFOLD parameters [19; http://unafold.rna.albany.edu/?q=mfold]. The optimum secondary structure was visualized using MFOLD.

RNA isolation and RT-PCR

RNA was isolated from log phase yeast by hot phenol:chloroform:isoamyl alcohol (PCA) extraction with SDS. The RNA was then precipitated with ethanol. 20 μg of total RNA was DNase-treated with 30 U DNase 1 (Roche) for 1 hour at 25°C. 1 μg of DNase-treated RNA was used for cDNA synthesis. cDNA synthesis was performed using the Maxima First Strand cDNA Synthesis Kit (ThermoFisher). 1 μL of cDNA was used for the Taq PCR reaction using gene-specific primers to analyze splicing.

Data Visualization

Venn diagrams to view the overlap between this and previous works were generated using Venny 2.1.0 [27]. Box plots and bar graphs were generated using the MATLAB functions boxplot and bar, respectively.

Accession numbers

Data generated in this study is available under GEO accession number GSE120497.

Acknowledgements

We thank the Dr. Tracy Johnson lab (UCLA) for comments and suggestions to improve the manuscript and Manuel Ares, Jr. for generously sharing unpublished data.

References

1. Wahl MC, Will CL, Luhrmann R. The spliceosome: design principles of a dynamic RNP machine. Cell. 2009;136: 701–718.

2. Li X, Liu S, Jiang J, Zhang L, Espinosa S, Hill RC, et al. CryoEM structure of Saccharomyces cerevisiae U1 snRNP offers insight into alternative splicing. Nat Commun. 2017 Oct 19;8(1):1035.

3. Juneau K, Nislow C, Davis RW. Alternative splicing of PTC7 in Saccharomyces cerevisiae determines protein localization. Genetics. 2009 Sep;183(1): 185-94.

4. Hossain MA, Rodriguez CM, Johnson TL. Key features of the two-intron Saccharomyces cerevisiae gene SUS1 contribute to its alternative splicing. Nucleic Acids Res. 2011 Oct;39(19): 8612-27.

5. Hossain MA, Claggett JM, Edwards SR, Shi A, Pennebaker SL, Cheng MY, et al. Posttranscriptional Regulation of Gcr1 Expression and Activity Is Crucial for Metabolic Adjustment in Response to Glucose Availability. Mol Cell. 2016 May 5;62(3): 346-358.

6. Kawashima T, Douglass S, Gabunilas J, Pellegrini M, Chanfreau GF. Widespread use of non-productive
alternative splice sites in Saccharomyces cerevisiae. PLoS Genet. 2014 Apr 10;10(4): e1004249.

7. Schreiber K, Csaba G, Haslbeck M, Zimmer R. Alternative Splicing in Next Generation Sequencing Data
of Saccharomyces cerevisiae. PLoS One. 2015 Oct 15;10(10): e0140487.

8. Plass M, Codony-Servat C, Ferreira PG, Vilardell J, Eyras E. RNA secondary structure mediates
alternative 3'ss selection in Saccharomyces cerevisiae. RNA. 2012 Jun;18(6): 1103-15.

9. Gould GM, Paggi JM, Guo Y, Phizicky DV, Zinshteyn B, Wang ET, et al. Identification of new branch
points and unconventional introns in Saccharomyces cerevisiae. RNA. 2016 Oct;22(10): 1522-34.

10. Qin D, Huang L, Wlodaver A, Andrade J, Staley JP. Sequencing of lariat termini in S. cerevisiae reveals
5⊠ splice sites, branch points, and novel splicing events. RNA. 2016 Feb; 22(2): 237–253.

11. Lardenois A, Stuparevic I, Liu Y, Law MJ, Becker E, Smagulova F, et al. The conserved histone
deacetylase Rpd3 and its DNA binding subunit Ume6 control dynamic transcript architecture during
mitotic growth and meiotic development. Nucleic Acids Research. 2015 Jan; 43(1): 115-28.

12. Venkataramanan S, Douglass S, Galivanche AR, Johnson TL. The chromatin remodeling complex
Swi/Snf regulates splicing of meiotic transcripts in *Saccharomyces cerevisiae*. Nucleic Acids Research.
2017; 45(13): 7708-7721.

13. Neves LT, Douglass S, Spreafico R, Venkataramanan S, Kress TL, Johnson TL. The histone variant
H2A.Z promotes efficient cotranscriptional splicing in *S. cerevisiae*. Genes & Development. 2017; 31(7):
702-717.

14. Nissen KE, Homer CM, Ryan CJ, Shales M, Krogan NJ, Patrick KL, et al. The histone variant H2A.Z
promotes splicing of weak introns. Genes & Development. 2017; 31(7): 688-701.

15. Awad AM, Venkataramanan S, Nag A, Galivanche AR, Bradley MC, Neves LT, et al. Chromatin-
remodeling SWI/SNF complex regulates coenzyme Q6 synthesis and a metabolic shift to respiration in
yeast. Journal of Biological Chemistry. 2017;292(36): 14851-14866.

16. Venkatesh S, Li H, Gogol MM, Workman JL. Selective suppression of antisense transcription by Set2-
mediated H3K36 methylation. Nature Communications. 2016; 7: 13610.

17. Leeds NB, Small EC, Hiley SL, Hughes TR, Staley JP. The splicing factor Prp43p, a DEAH box ATPase,
functions in ribosome biogenesis. Molecular Cell Biology. 2006; 26(2): 513-22.

18. Meyer M, Plass M, Pérez-Valle J, Eyras E, Vilardell J. Deciphering 3'ss selection in the yeast genome
reveals an RNA thermosensor that mediates alternative splicing. Molecular Cell. 2011; 43(6): 1033-9.

19. Zucker M. On finding all suboptimal foldings of an RNA molecule. Science. 1989 Apr; 244(4900): 48-
52.

20. Sinturel F, Bréchemier-Baey D, Kiledjian M, Condon C, Bénard L. Activation of 5'-3' exoribonuclease
Xrn1 by cofactor Dcs1 is essential for mitochondrial function in yeast. PNAS. 2012; 109(21): 8264-9.

21. Mallory MJ, Cooper KF, Strich R. Meiosis-specific destruction of the Ume6p repressor by the Cdc20-
directed APC/C. Molecular Cell. 2007; 27(6): 951-61.

22. Talkish J, Igel AH, Perriman RJ, Shiue L, Katzman S, Munding EM, et al. Rapidly evolving protointrons
in *Saccharomyces* genomes revealed by a hungry spliceosome. bioRxiv. 2019.

23. Grate L., Ares M. Jr. Searching yeast intron data at Ares lab web site. Methods Enzymol. 2002; 350:
380–392.

24. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq
aligner. Bioinformatics. 2013; 29: 15–21.

25. Oliveros, JC. Venny: An interactive tool for comparing lists with Venn's diagrams. 2015.
http://bioinfogp.cnb.csic.es/tools/venny/index.html.

Figure 1. Workflow for discovery of novel splice forms. Multiple RNA sequence datasets are
consolidated then aligned to the *Saccharomyces cerevisiae* genome and annotated transcripts liberally,
allowing up to six mismatches but no gaps. Reads that fail to align are again aligned, but now allowing a
single gap corresponding to a putative intron and no mismatches. This step defines all putative novel
introns. Once again all reads are aligned to the *Saccharomyces cerevisiae* genome, known transcripts,
and putative novel introns allowing up to two mismatches, which determines each putative novel
intron's read counts. Consensus sequences are determined from known *Saccharomyces cerevisiae* 5'
splice site, 3' splice site, and branch point sequences, and each putative novel intron is scored based on
sequence similarity to the consensus sequences.

Figure 2. RT-PCR of representative novel splice isoforms. (A) *BIG1*; novel 3' splice site within
annotated ICG. (B) *SIM1*; 5' UTR intron, novel 5' splice site within annotated ICG. (C) *MCR1*; 5' UTR
intron, novel 5' and 3' splice sites within an annotated ICG. (D) *SPF1*; intron within an unannotated ICG.
(E) *LEU4* antisense; antisense transcript of the *LEU4* gene containing an intron.

Figure 3. Agreement with previous studies. Venn diagram showing the overlap between the novel
splice forms discovered here and those described previously.

Figure 4. *prp43-1* leads to a decrease in splicing. (A) Strains with the mutated *prp43-1* gene show a
decrease in splicing efficiency of annotated introns relative to strains with wildtype *PRP43*. (B) Evidence
of novel splicing is lower in strains with *prp43-1* versus wildtype *PRP43*.

Figure 5. Elevated temperature leads to longer introns. (A) Boxplot of temperature-enriched splice
forms versus temperature-independent splice forms shows higher temperature leads to an increase in
intron size. Temperature-enriched splice forms have much higher counts at higher temperature, but are
still present at lower temperature. (B) Shifting *prp43-1* to higher temperature causes a dramatic
decrease in these temperature-enriched splice forms. (C) Predicted structure of TMH18 with annotated
and novel splice sites labeled.

Figure S1. Putative novel splice site scores. Each putative novel splice site was scored by similarity to known *Saccharomyces cerevisiae* splice signals. The splice site score was plotted against putative intron length for novel splice products within annotated ICGs (blue) and unannotated ICGs (red).

Figure S2. Summary of novel splice products. (A) Over half of the data have fewer than 6 counts across all 29 of our datasets. (C) Novel splicing outside of annotated intron-containing genes and splicing of RNAs that are antisense to known transcripts is observed.