Title

Response relationships between CO_2, CH_4 and N_2O emissions and microbial functional groups in
wetland sediments after trace metal addition.

Authors

Georgios Giannopoulos*, Katherine Hartop, Bonnie Brown, Rima Franklin

Dept. Biology, Virginia Commonwealth University, Richmond, VA 23284, USA.

Current address: School of Chemistry, Chemical and Environmental Technology Lab. Aristotle University
of Thessaloniki, 546 40 Thessaloniki, Greece.

11
Abstract (176 words)

Biochemical nitrogen removal relies heavily on wetland microbes, prior to entering downstream
ecosystems. Microbes utilize oxy-reductases that contain metal-cofactors to reduce NO3 and NO2 to N2.
These transformations are linked also to other C cycling processes and may emit N2O, CH4 and CO2 as
well. In this incubation study we tested the effects of supplementing wetland sediments with trace-
metals (Cu, Mo, Fe) at µM levels on nitrogen kinetics, gaseous emissions (N2O, CO2 and CH4) and genes
abundances for the overall microbial community, related to NO2 reduction (denitrification and DNRA),
N2O reduction and methanogenesis. Trace metal availability and specifically Cu, enhances the reduction
of N2O to N2, increases the production of CH4 and has no effect on CO2 emissions from wetland
sediments. When sediments were supplied with Cu, higher gene abundances were observed for Cu-
reductases *nirK* and *nosZ*. Mo addition increased CH4 emissions from sediments and higher *mcrA* gene
abundances were observed. Emerging from this investigation, trace-metals bio-availability may directly
or indirectly regulate denitrification in the environment; an effect observed so far only in pure microbial
cultures.

27
28 **Keywords: trace metals, denitrification, greenhouse gases, sediments, microbial groups**
29
30

Introduction

The removal of many upstream anthropogenic pollutants (industrial, urban and agricultural) prior to entering downstream coastal and marine ecosystems relies on wetland microbes. Out of those pollutants, nitrogen (N) pollutants (exceeding 60 Tg N y^{-1}) are of major concern because they are linked to water quality degradation, eutrophication and increased greenhouse emissions (carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O)) (Anderson et al., 2010;Tian et al., 2015). Wetland microbes reduce the N load through various biochemical pathways i.e. assimilation, reduction and oxidation, but often this ecosystem service comes with the release of significant amounts of CO_2, N_2O and CH_4.

Microbial processes of nitrification, denitrification or nitrifier-denitrification, methanogenesis and methanotrophy are responsible for the production of N_2O and CH_4 in wetland sentiments (Soares et al., 2016). In particular N_2O is a very potent greenhouse gas, with a global warming potential ~ 300 and ~ 60 greater than CO_2 and CH_4, respectively, with important implications in ozone depletion. N_2O can be further reduced to inert N_2 by microbes that possess the only known enzyme to reduce N_2O; the nitrous oxide reductase (NOS)(Thomson et al., 2012). It is estimated that ~ 17 Tg N p.a. is returned to the atmosphere via denitrification in wetlands and terrestrial ecosystems globally, releasing ~ 48 Tg N p.a. to coastal and marine ecosystems (Schlesinger, 2009). Besides N cycling, wetlands are important ecosystems in C cycling as well. Despite their relative small coverage (~ 10 % globally), wetlands store large amounts of C and emit considerable amounts of CH_4 (144 Tg CH_4 p.a.)(IPPC, 2014). CH_4 emissions are the trade-off balance between the microbial processes of methanogenesis and methanotrophy.

The production and consumption of N_2O and CH_4 is catalyzed by oxy-reductases that utilize a metal co-factor (Mo, Cu, Fe and S) (Glass and Orphan, 2012). The reduction of NO_3^- is catalyzed by a periplasmic or membrane bound molybdenum (Mo) nitrate reductase (NAR) (Schwarz et al., 2009), NO_2^- is reduced to NO by either a non-metal cytochrome cd_1 (Einsle et al., 1999) or Cu-cofactor (Adman et al., 1995) nitrite reductase (NIR; expressed by *nirS* and *nirK* respectively). NO is reduced to N_2O by nitric oxide reductase (NOR) and contains heme-Fe co-factor; two types of NOR are distinguished, cNOR which is found in denitrifying bacteria and receives electrons from cytochrome *c* and qNOR which is typically found in pathogenic non-denitrifying bacteria as well as in denitrifiers and receives electrons directly from quinol (Shiro, 2012). Finally N_2O is reduced by the Cu-containing NOS (Rosenzweig, 2000). NO_2^-, besides being a denitrification intermediate, it is also a substrate for dissimilatory nitrite reduction to ammonium (DRNA) and this permutation has been linked to C/N (Kraft et al., 2014;Yoon et al., 2015;van den Berg et al., 2017). Likewise, denitrification may be linked or co-occur with other C cycling processes.

Interestingly, members of the NC10 phylum (e.g. *M. oxyfera*) are able to couple methane oxidation to

denitrification (NO_2^--dependent methane oxidation (N-DAMO)) utilizing a Cu-dependent particulate

methane monoxygenase (pMMO) and a Fe-rich cd_1 nitrite reductase (Deutzmann et al., 2014). While

extensive research has tested the relative contribution of key environmental factors (pH, NO_3^-/NO_2^-, O_2,

C/N) in regulating anaerobic microbial processes and specifically denitrification (Thomson et al., 2012),

we argue that the abundance and bio-availability of trace metals in the environment could considerably

control greenhouse gas emissions in the C and N biochemical cycle.

Metals in the environment have been traditionally seen as unwanted pollutants. This is because trace

metal accumulation is linked to toxicity and inhibition of ecosystem processes (Samanidou and

Papadoyannis, 1992). Contrariwise, low levels of trace metals may limit crop production, and impede

animal and human nutrition (Teklić et al., 2013). Concentrations exceeding $mg.L^{-1}$ range for Cu, Mo, Fe,

Zn and Pb severely inhibited denitrification in soils, sediments, water-bodies and waste waters (Labbé et

al., 2003;Magalhães et al., 2007;Liu et al., 2016). The recent advances in our understanding of

denitrification using model organisms demonstrated the complete reduction of NO_3^- to N_2 requires trace

metals at µM to pM levels. In fact, *in vitro* studies have shown that lack of Cu, Mo or Fe severely inhibits

denitrification or methane cycling, due to the formation of non-functional enzymes typically lacking the

respective metal co-factor. In soil bacteria *P. denitrificans* and *Pseudomonas stutzeri* Cu is required to

express to a functional Cu-containing NOS dimer (Granger and Ward, 2003;Felgate et al., 2012;Black et

al., 2016). In methanotrophs oxidizing CH_4 to methanol (CH_3OH) by MMO, the switch over between a Cu-

or Fe-dependent MMO is regulated by the environmental availability of Cu and Fe respectively (Murrell

et al., 2000;Bollinger Jr, 2010). All three classes of methanogens utilize a common cobalt dependent

Methyl-coenzyme M reductase (MCR; *mcrA*), whereas in the earlier steps of methanogenesis are

catalyzed by Fe-, Ni- and Zn- dependent ferrodoxins and dehydrogenases (Glass and Orphan, 2012).

Based on the above observations, few studies have optimized the trace metal concertation for

denitrification in a culture medium (Hahnke et al., 2014) and suggested the optimal levels of trace

metals in biological wastewater treatment plants (He et al., 2015;Wintsche et al., 2018).

Despite the great progress in metal ecotoxicity and metalloenzyme biochemistry, our understanding of

trace-metal availability in ecosystem C and N biochemical cycles is generally limited to aquatic

ecosystems primarily focusing on productivity, NH_4^+/NO_3^- and CO_2 assimilation rates (Twining et al.,

2007;Glass et al., 2012;Moore et al., 2013;Romero et al., 2013;Schoffman et al., 2016). Here, we

investigated the effects of trace metal (µM) addition on the microbial biochemistry of wetland

sediments focusing primarily on NO_2- reduction and greenhouse gas kinetics. We applied molecular

techniques (qPCR) to quantify changes in the microbial ecology of wetland sentiment in an effort to

better understand the intrinsic link between microbial ecology and process rates. We hypothesized that

(*i*) the bio-availability of metals (Mo, Fe and Cu) at μM levels will increase denitrification rates and

specifically Cu availability will reduce N_2O emissions. Also, considering C respiration, (*iia*) CO_2

emissions will be driven by denitrification and (*iib*) no effect on CH_4 emissions is expected.

Materials and Methods

Soil samples and experimental set-up

A combined (5 sites; 5-10 cm from surface) soil sample and soil water were collected from the tidal-

fluvial point bar deposit of the Pamunkey River (N 37.557451, W -76.972521) using a PVC pipe (Ø 10

cm). Soil sediments had 25 ±4 %DW organic matter (OM), 79 ±6 % moisture, 0.18 ±0.05 $g.cm^{-3}$ bulk

density, C/N ratio of 11 ±1.5 and pH 6.1 ±0.3. Soil water, at the collection time, contained 0.1 ±0.03 mM

NO_3^-, 0.1 ±0.02 mM NO_2^-, 0.16 ±0.04 mM SO_4^-, 0.4 ±0.1 mM Cl^-, and 0.3 ±0.1 mM NH_4^+, 0.11 ±0.01 mM

K^+, 0.26 ±0.1 mM Mg^{2+} and 0.29 ±0.1 mM Ca^{2+}. Samples were kept cool (ice-cooler) during transport and

stored at 4 ^{o}C overnight prior to processing. Any large stones and roots were removed, then 1000 g wet

soil was homogenized with an additional 1500 ml of filtered soil-water (Cellulose, 10 μm, Millipore,

USA). Soil slurry aliquots of 50 ml were transferred to 125 ml bottles (Wheaton, USA). Bottles were

crimped with rubber septa and were kept for 3 days in dark (25 ^{o}C) to let any residual O_2 to be

consumed by basal respiration. Then, all treatments received 250 μl aliquots of 2 M KNO_3 reaching a

final concentration of 10 mM. A 400 μl volume of ammonium molybdate ((NH_4)Mo_7O_{24}; 1 mM), iron

sulfate ($FeSO_4$; 18.5 mM) and copper sulfate ($CuSO_4$; 6.5 mM) stock solution were injected to the

bottles, adding 4, 74 and 26 μM respectively, according to the treatment plan (Table 1). Bottles were

vortexed briefly (30 s at 6000 rpm), flushed with N_2 (130 mins) and incubated at 25 ^{o}C in dark. Through-

out the incubation several gas and liquid samples were taken anaerobically (as indicated in the graphs).

At 96 hours, bottles were un-crimped in an O_2-free chamber, then 1 g of wet soil was taken

anaerobically and was frozen instantly for further molecular analyses.

Analytical techniques

Gas samples of 5 ml were stored in 3 ml exetainer vials (Labco, UK), that were previously flushed (N_2; 5

mins) and vacuumed. Gas samples were analyzed for N_2O, CO_2, CH_4 and O_2 concentration with gas

chromoatography (GC – 2014 equipped with FID, TCD and ECD; Shimadzu, USA) as in Morrissey and

Franklin (2015). For each gas, total production was determined as the sum of the gas accumulated in the

headspace and the gas dissolved in the slurry; the latter was estimated using the relationships described

by Heincke and Kaupenjohann (1999). From each bottle, 1.5 ml slurry samples were collected,

centrifuged (10000 x g for 5 mins) and the supernatant was diluted 10x, filtered (0.22 μm) and stored (-4

°C) prior to ion analysis. The concentration of NO_3^-, NO_2^-, SO_4^{2-} and NH_4^+ was determined by ion-

chromatography (ICS – 5000+, Dionex, USA) according to the manufacturer's instructions using columns

Dionex IonPac AS17 (2 × 250 mm) and IonPac CS12 (A – 5 μm, 3 × 150 mm) and a Dionex conductivity

sensor. Soil water content and OM were determined gravimetrically, soil pH was determined with a

glass electrode pH probe (Laquact; Horiba, USA) in soil-water extract (after centrifugation 4000 x g for

20 mins).

Molecular techniques

DNA was extracted from frozen soil samples (0.3 g) using the DNEasy Kit from Qiagen (USA) following

the manufacturer's instructions. The DNA yields were, on average, 42 ±2 ng.μl^{-1}. The retrieved DNA was

quantified using a Nanodrop spectrometer (Thermo, USA) and visualized on a 1.2 % agarose gel for

integrity. DNA samples were diluted to 3 ng.μl^{-1} and were further tested (see qPCR protocols on Table 2)

for the abundance of microbial groups typically found in wetlands: total eubacteria (*eub-518*), nitrite

reducers (denitrification: *nirK*, *nirS* and DNRA: *nrfA*), nitrous oxide reducers (*nosZ*) and methanogens

(*mcrA*). SensiFAST™ SYBR® No-ROX Kit (Bioline, UK) polymerase x2 mix was used according to the

manufacturer's instructions and real time data were collected using a Bio-rad CFX-384 (Bio-rad, USA)

real time PCR engine.

Statistical analyses

Kinetic and molecular datasets were tested for normality (RStudio [*base*]; Shapiro.test). Non-parametric

statistical tests were applied to assess any differences in the mean of ranks between the treatments

(RStudio [*agricolae*]; *Kruskal-Wallis* with *Bonferroni* correction). Basic statistics (mean and standard

error) were summarized with RStudio [*dyplr*] and plotted with RStudio [*ggplot2*]. For all statistical tests

$p<0.05$ was considered significant.

Results

The effects of trace metal addition on NO_3^-, NO_2^- and NH_4^+ kinetics

At the end of the incubation, all treatments had visible gas bubbles on the surface and within the slurry,

indicating microbial activity and gas production. We observed yellow and black patches within the slurry

possibly due to Fe and Mn reduction indicating anaerobic redusive conditions. Denitrfication

commenced with out any lag between the control and trace metal additions. Approximatelly, 12 mM

NO_3^- were consumed and 0.3 mM NO_2^- remained, while NH_4+ ranged from 200 to 750 µM at the end of

the incubation (**Figure 1**). At 48 h we observed a reduction in the remaining NO_2^- and a simultaneous

increase in the NH_4^+ concentration for all treatments. The remaining NO_3^- and NO_2^- at the end of the

incubation did not differ significantly when compared to the control. The treatments receiving additonal

Mo + Fe ($p \leq 0.01$) and Mo + Fe + Cu ($p \leq 0.01$) had significantly higher NH_4^+ than the control, and an

increasing trend in NH_4+ concentration for all treatments was observed towards the end of the

incubation, perhaps due to OM mineralization.

The effects of trace metal addition on CO_2, CH_4 and N_2O cumulative gas kinetics

Cumulative CO_2 production increased rapidly between 12 to 24 h, and continued to increase up to ~ 24

mM C-CO_2 at the end of the incubation. There were no significant differences between the control and

each metal treatment, except the Cu and the Mo + Fe treatment (Figure 2.a). A faster rate of CH_4

accumulation in the microcosms was observed between 6 to 12 h of incubation, earlier than CO_2. CH_4

continued to accumulate during the incubation and reached levels ranging from ~ 4 to 6.5 mM C-CH_4. At

the end of the incubation, we observed that the Cu treatment had significantly accumulated more CH_4

than the control treatment (Figure 2.b). Cumulative N_2O emissions rapidly increased between 24 to 48 h,

matching the increased NO_3^- consumption. Interestingly, the control, Mo, Fe and Mo + Fe treatment

accumulated significantly more N_2O than the remaining treatments that contained Cu (Cu, Fe + Cu, Mo +

Cu and Mo + Fe + Cu). These differences remained until the end of the incubation (Figure 2.c). Contrary

to CO_2 and CH_4 emissions, N_2O emissions reached a plateau between 48 to 96 h, indicating that N_2O

emissions were limited by the exhaustion of the available NO_3^- and not by C.

The effects of trace metal addition on targeted microbial groups

Average total DNA was ~ 52 ng.μl^{-1}.g^{-1} sediment with the Mo + Fe + Cu treatment having significantly

(p=0.03) greater DNA yield (62 ±2 SE) that the control (44 ±5 SE). The average bacterial abundance was

6.75E+8 *eub* copies g^{-1} sediment with the Fe, Fe + Cu and Mo + Cu treatments having significantly

(p=0.005) more *eub* gene copies than the control (**Figure 3.a**). The microbial group of methanogens

(*mcrA*) was relatively more abundant in the treatments containing additional Mo (Mo, Mo + Cu, Mo + Fe

and Mo + Fe + Cu; **Figure 3.c**) when compared to the control. Regarding the microbial groups involved in

N cycling we observed no effect of trace metals on DNRA NO_2^- reducers (*nrfA*; **Figure 3.b**). Interestingly,

trace metal addition affected the abundance of *nirK* ($p<0.001$; **Figure 3.d**), *nirS* ($p<0.001$; **Figure 3.e**) and

nosZ ($p=0.001$; **Figure 3.f**) denitrifying microbial groups. NO_2^- reducers utilizing Cu-NIR (*nirK*) were more

abundant in all treatments when compared to the control, except when Fe was added. Microbial groups

having a cytochrome cd_1-NIR (*nirS*) were relatively more abundant in the Mo and Fe + Cu treatment.

Lastly, the *nosZ* microbial group responsible for the reduction of N_2O to N_2 was more abundant in the

treatments Fe+Cu, Mo+Cu and Mo+Fe+Cu than the control. Although *nosZ* abundance was relatively

higher in all trace metal additions, the effects of single Cu addition on *nosZ* did not differ significantly.

Discussion

We observed that trace metal additions (Mo, Fe, Cu and their combinations) in wetland sediments

regulated denitrification kinetics and greenhouse gas emissions, and altered the abundance of targeted

microbial groups that are typically found in wetlands. The key interpretation of the incubation

experiments is that trace metal availability and specifically Cu, enhances the reduction of N_2O to N_2, the

production of CH_4 and has no effect on CO_2 emissions from wetland sentiments. The increase in the

denitrifiers' associated physiological activity to reduce N_2O when supplied with additional Cu

corresponded to the relative higher abundance of Cu-containing genes - nitrous oxide (*nosZ*) and

nititrite (*nirK*) reductase, but not to the abundance of nitrite reduces *nirS* and *nrfA,* suggesting a direct

relationship between Cu availiabilty and denitrification process in soils.

The absence of substantial amounts of NO_2 and N_2O in the bottles for the Cu-containing treatments (Cu,

Fe + Cu, Mo + Cu and Mo + Fe + Cu) and an increase in the respective gene abundance indicates that a

balanced denitrification process NO_3^- to N_2. Microbes with a complete denitrification strategy harvest

the energy from the last reduction step (N_2O to N_2) in addition to the rest of the denitrification steps,

which may be particularly advantageous when there are limited electron acceptors available in the

environment. We do not exclude other partial denitrification processes or other anaerobic processes

simultaneously co-occurring with denitrification as subjected by the residual N_2O in the treatments Cu-

containing treatments (≤2.3 mM $N-N_2O$). The increasing NH_4+ concentration (<1 mM) towards the end

of the incubation could indicate DNRA and /or N mineralization processes commonly co-occurring in

wetland sediments (White and Reddy, 2009). However, the lack of significant differences in DNRA gene

abundance indicates towards anaerobic N mineralization. Although we didn't measure NO in the

treatments, we assume that it was quickly utilized due to its potent cytotoxicity at µM levels (Chaudhari

et al., 2017). In the treatments accumulating notable amounts of N_2O (≥ 1mM) N_2O cytotoxicity may

occur and it can be releaved by the addition of vitamin B12, Co or methionine (Sullivan et al., 2013). The

continuous accumulation of CO_2 in all treatments, as proxy for microbial activity, firstly, excludes any

indirect acute cytotoxicity from NO due to NO_2^- reduction or N_2O accumulation and secondly idicates

that the availible electron donor (organic C) was in excess of the added NO_3^-. The magnitute of CO_2

accumulation was predictable; treatments with complete dentrification, thus having less residual N_2O

had accumulated more CO_2 by the end of the incubation. We assumed that the lack of additional Ni

would inhibit MCR, key enzyme in methanogenesis (see *Possible mechanisms and other regulators*).

Contrary to our hypothesis *iib,* trace metals had a significant effect on CH_4 emissions.

Other studies and trace metal levels

Several studies have looked into the effects of trace metal abundance or addition on C and N cycling in

sediments and peat soils. Typically those studies report a negative effect of trace metals on

biogeochemical processes, however the range of trace metal concentration far exceed the levels of the

current study. For instance, Keller and Wade amended peat slurries with ~ 100 µM trace element

solution and saw significant decrease in CH_4 emission, probably due to Cu induced toxicity. Similarly,

Magalhães et al. (2007) observed an 85% inhibition of denitrification accompanied with accumulation of

substantial N_2O and NO_2^- amounts at 79 Cu µg.g^{-1} sediment (equiv. 1.2 mM Cu) in estuaries. In a follow-

up study, lower denitrification rates due to Cu (~ 0.9 mM) were accompanied with a decline in *nirK*, *nirS*

and *nosZ* microbial group abundance and *β*-diversity (Magalhães et al., 2011). A more recent study that

tested the effects of Cu pollution (addition of 100 µg.L-1 ~ 1.6 µM) on urban wetland sediments, found

that Cu addition significantly reduced CH_4 emissions, but had no effect on N_2O emissions when

compared to the control (freshwater sediments; Doroski et al. (2019)). This divergent observation could

be due to low NO_3^- concentration (18 µM) in relation to available C or the presence of substantial

amounts of soil Cu (9.4 µg.g^{-1} sediment) at the start of the experiment in both control and Cu treatment.

Possible mechanisms

251 Denitrification is an anaerboic respiratory process that utilizes NO_3^- as a terminal electron acceptor

252 (Equation (2)) accompanied by energy transfer (Thomson et al., 2012). Thus, for a complete

253 denitrification reaction, two molecules of NO_3^- will be converted to one molecule of N_2 with the

254 transferring of 10 electrons to the electron recipient (NO_3^-, NO_2^-, NO and N_2O). Therefore, microbial

groups unable to utlize the complete denitrification pathway will harvest less energy, and/or will have to compensate the energy loss by adjusting their metabolic rates for example by enhancing the rate of NO_3^- reduction. Such bioenergetic adjustment was observed in chemostat cultures of *P. denitrificans*; the *nosZ* deficient strain was unable to complete the last step of denitrification and had lower biomass and higher NO_3^- reduction rates (Giannopoulos et al., 2017). At RNA level, Cu-decifient cells had more transcripts of *nosZ* and Cu-scavenging genes to compensate for the loss of N_2O reduction due to depleted Cu availiability (Felgate et al., 2012). And finally, strains deficient in Cu – transporters and chaperones were unable to reduce N_2O to N_2, although nosZ was expressed, forming a non-functional NOR (Sullivan et al., 2013). Therefore we postulate that, at community level, microbes utilizing the complete dentrification pathway will have a physiological advantage than those unbale to do so. This was supported by the increased abundances of NO_2^- reducers (*nirK*) and partly of the N_2O reducers (nosZ) and of the overal microbial abundance (eub); corroborating the fact that denitrifcation genes were up-regulated in single denitrifiers (*P.stutezi* at 0.05 Cu mg.L^{-1}; Black et al. (2016)).

Considering net CH_4 emissions, Mo clearly enhanced CH_4 accumulation (Mo) and abundance of *mcrA* genes in the Mo containing treatments. A Mo co-factor is typically required for the formylmethanofuran dehydrogenase (*fwd*) that catalyzes the reduction of CO_2 to formyl-methanofuran in the first step of methanogenesis by reduction of CO_2 with electrons from H_2 (Glass and Orphan, 2012). Mo availioability and the increasing partial pressure of CO_2 in the bottles during the incubation could have triggered a shift towards a Mo-based methanogenesis pathway. Another interesting aspect in CH_4 emissions and Cu, is the competition between methanotrophs and denitrifiers for the "Cu monopoly" (Chang et al., 2018), which should be further investigated in environmental samples. Eventhough the purpose of this study wasn't to quantify the relative contribution between methanotrophs (MMO) and methanogens (MCR) in the net CH_4 emissions, the higher abundance of methanogens (*mcrA*) and noticable accumulation of CH_4 in all metal treatments indicates the important role of trace metal availiability in CH_4 cycling in environmental samples.

*** Trace metal limitation in the environment and other regulators***

The abundance of bio-availiable metals in the environment, typically in the ion form, is often observed at several magnitute orders less compared to the routinely reported total metal content. Such trivial amounts could be unavailiable to microorganisms due to physiochemical interactions and plant uptake (Giller et al., 1998). At neutral and alkaline pH levels, metals tend to be effectively immobilized as inorganic compounds (metal – oxides, –hydroxides and – carbonates). Additionally, organic complexes

such humic ligands may bind metal cations making them unavailiable to microbes and plants. For

example, the particularly low concentrations of dissolved Cu, Fe and Mo (~ 5, 1 and 0.2 $\mu g.L^{-1}$,

respectively) and the copious dissolved organic C (~ 80 $mg.L^{-1}$) in siberian peatlands (Raudina et al.,

2017), could trigger significant emissions of CH_4 and N_2O as our climate is warming due to metal

limitations (Basiliko and Yavitt, 2001;Voigt et al., 2017). On the other hand, in temperate salt marshes

with reducing conditions, SO_4^- may suppress CH_4 emissions, but the abundant H_2S may reduce the

availiability of trace metals with important implications in N cycling and N_2O emissions (Gauci et al.,

2004;Butterbach-Bahl et al., 2013). In this study we added 26 μM and 74 μM of SO_4^- as $CuSO_4$ and $FeSO_4$

respectively that could potentially be reduced to H_2S, however the initial and final SO_4^- concentration

remained at comperable levels (Table 4).

Other well known environmental factors controlling denitrification is O_2, NH_4^+, NO_3^-/NO_2^- and C

(Thomson et al., 2012). Synthesizing the findings from other chemostat (Baumann et al., 1996;Felgate et

al., 2012;Giannopoulos et al., 2017;Hartop et al., 2017;Conthe et al., 2018) and incubation (Koike and

Hattori, 1975;Okereke, 1993;Morley et al., 2008) studies, it appears that the relative importance of the

aforementioned factors as control of denitrification is O_2> NH_4, NO_3/NO_2> C/N or C quality, with trace

metal bio-availability exerting an additional effect on each factor. In this study we did not indent to

quantify the relative contribution of each environmental factor on denitrification. Instead we

demonstrated the importance of trace metal availability on ecosystem processes contributing to the

release of N_2O, CH_4 and CO_2.

Future challenges

Trace metal limitition in denitrification biochemistry and in general the C and N cycle may also be

important in agricultural ecosystems besides wetlands. The continous fertilization with macronutrients

(NPK) and intentisive cropping has resulted in low concentrations of microelements (trace metals) in

soils and edible crops (hidden hunger; (Alloway, 2008);Teklić et al. (2013)). The additon of essential

micronutrients particularly Cu, Mo and Fe to agricultural lands has been previously suggested as a

possible strategy to mitigate N_2O emissions (Richardson et al., 2009), but few assessments have been

reported. Felgate et al. (2012) and Black et al. (2016) recomended a Cu concentration of 13 μM and 50

μM (equiv. 150–200 $\mu g.g^{-1}$ Cu in soils) based on studies with *P. denitrificans* and *Ps. Stutezi*

respectively. Building upon prior knowledge, we demonstrate that effects of metal addition (2.5 Cu, 26.7

Fe and 7.8 Mo $\mu g.g^{-1}$ sediment) on greenhouse gas emissions in environmental samples (sediments) are

due to subsequent changes in the microbial community structure. We are currently missing quantitative

data on optimal levels of soluble trace metals for biochemical processes in C and N cycling in a variety of

different soil types. Agricultural and flooded soils are particularly interesting because they tend to have
higher N_2O yields than marine and aquatic ecosystems (Beaulieu et al., 2011). Future studies should
investigate the effects of trace metal bio-fertilization in soils under commercial crops to assess any
beneficial effects in greenhouse gas mitigation and crop quality, as in Montoya et al. (2018). The type of
trace metal fertilizer is particularly interesting to investigate because chelators based fertilizers may
scavenge the metal co-factors and dissable key enzymes of the N cycle (urease, ammonia
monooxygenase and nitrite reductase) prior to N_2O reduction (Moffett et al., 2012;Montoya et al.,
2018).

Assessing the abundance of microbial groups using qPCR showed significant differences between the
control and the treatments. In this short term batch experiment, differences in microbial group
abundance could also illustrate changes in in the diversity of the microbial groups, for example between
the abundance of copiotrophic and / or fast growing (*r* strategists) microbes upon NO_3^- and trace metal
addtion, relative to slow growing (*k* strategists) microbes (Fierer et al., 2012). Further information on
microbial diversity and meta-transcriptomic profiles under comperative treatments is needed to
understand and evaluate the ecological importance of trace metal limitation. Such 'omic studies should
be accompanied by high-throughput metabolite or process rate analysis either in the lab or in the field.

Conlcusion

Emerging from this investigation, our results suggest that trace metal bio-availability may, directly or
indirectly, regulate or co-limit denitrification kinetics and induce a shift in the microbial community in
environmental samples, favoring a more active microbial community that is able to quickly acquire the
bio-available metals and incorporate them in functional oxy-reductases. This observation has important
implications in wetland C and N cycling. Considering also agro-ecosystems that are becoming limited in
trace metals trace metal amendments should be evaluated and considered as an alternative way to
mitigate N_2O emissions and improve the nutritional quality of crops.

Acknowledgements

346 I wish to thank Dr Ioannis Ipsilantis for his comments on an earlier version of this manuscript.

Figure 1 Average a) NO_3^-, b) NO_2^- and c) NH_4^+ concentration (mM; n=4, ±SE) per treatment throughout

350 the incubation (96 h). Levels of significance: *≤0.5, ** ≤0.01, *** ≤0.001 and *ns* denotes non-significant.

351 For Bonferroni test results, see SI.

354   **Figure 2** Cumulative gas concentration of CH_4, CO_2 and N_2O (n=4; ±SE). Levels of significance: *≤0.5, **

355   ≤0.01, *** ≤0.001 and ns denotes non-significant. For Bonferroni test results, see SI.

Figure 3 Average gene copies per g sediment per treatment at the end of the incubation (96 h; n=4; ±SE). Treatments with the same letter are not significantly different.

357
358

360   Table 1 Experimental treatments.

Treatment	Replicates	Nitrate (KNO$_3$; mM)	Molybdenum ((NH$_4$)Mo$_7$O$_{24}$; µM)	Iron (FeSO$_4$; µM)	Copper (CuSO$_4$; µM)
Control	4	10	--	--	--
Cu	4	10	--	--	26
Fe	4	10	--	74	--
Mo	4	10	4	--	--
Fe + Cu	4	10	--	74	26
Mo + Cu	4	10	4	--	26
Mo + Fe	4	10	4	74	--
Mo + Fe + Cu	4	10	4	74	26

 Table 2 Primers and reaction conditions for qPCR assays

Target	Primer	PCR Mix (15 µl)	Standard	Reaction Conditions	Reference
Eubacteria (*16S rRNA*)	Eub338 Eub518	1.2 ng template 0.1 µM each primer	*Desulfovibrio desulfuricans* ATCC 27774	95°C for 4 min, then 40 cycles of 30 s at 95°C, 30 s at 55.5°C, and 60 s at 72°C	(Fierer et al., 2005)
Denitrifiers (*nirS*)	cd3aF R3cd	10 ng DNA template 0.1 µM each primer	*Paracoccus dentrificans* ATCC 17741	95°C for 4 min, then 50 cycles of 30 s at 95°C, 30 s at 56°C and 60 s at 72°C	(Throback et al., 2004)
Denitrifiers (*nirK*)	nirKq-F nirK1040	1.5 ng DNA template 0.35 µM each primer	*Pseudomonas nitroreducens* ATCC 13867	15 min at 95°C, 9 touchdown cycles of 95°C for 15 s, 68°C for 60 s, and 81.5°C for 30 s (-1° per cycle for annealing); then 28 cycles of 95°C for 15 s, 60 °C for 60 s, and 81.5 °C for 30 s	(Smith et al., 2007)
Denitrifiers (*nosZ*)	nosZ1F nosZ2F	10 ng DNA template 1 µM each primer	*Pseudomonas fluorescens* C7R12	15 min at 95°C, 6 cycles of 95°C for 15 s, 67°C for 30 s with a touchdown of −1°C by cycle, 72°C for 30 s, and 80°C for 15 s (acquisition data step); 40 cycles of 95°C for 15 s and 62°C for 15 s, 72°C for 30 s, and 80°C for 15 s; and 1 cycle at 95°C for 15 s and 60°C for 15 s, to 95°C for 15 s	(Henry et al., 2006)
DNRA (*nrfA*)	nrfA6F nrfA6R	10 ng DNA template 0.3 µM conc. primers	*Escherichia coli* ATCC 11775	50°C for 2 min, 95°C for 8.5 min, and 50 cycles of 20 s at 94°C, 40 s at 54.5°C, and 10 s at 72°C	(Takeuchi, 2006)
Methanogens (*mcrA*)	mlas mcrA-rev	2 ng DNA template 0.6 µM conc. mlas 0.7 µM conc. mcrA-rev	*Methanococcus voltae* ATCC Strain A3	95°C for 5 min, then 50 cycles of 20 s at 95°C, 20 s at 59°C, and 45 s at 72°C	(Steinberg and Regan, 2009)

363 **Table 3** Average DNA yield per treatment and post-hoc analysis results of the probability of treatment
364 differences (n=4, ±SE).

Treatment	Total DNA		Means of the ranks*	
	$(ng.\mu l^{-1})$			
Control	44.3	±5	8.8	b
Cu	44.0	±2	6.0	b
Fe	50.6	±1	15.6	ab
Mo	57.0	±3	22.0	ab
Mo + Cu	52.8	±3	17.4	ab
Mo + Fe	53.3	±5	16.9	ab
Fe + Cu	53.3	±4	17.4	ab
Mo + Fe + Cu	62.3	±2	28.0	a

365 *Treatments with the same letter are not significantly different.*

368 **Table 4** Initial (0 h) and final (96 h) SO_4^- concentration in the treatments (n=4, ±SE).

Treatment	Initial $[SO_4^-]$ μM		Final $[SO_4^-]$ μM	
Control	4.9	±0.8	8.5	±3.5
Cu	23.4	±2.9	20.25	±2.0
Fe	68.9	±2.3	56.95	±8.4
Fe + Cu	99.2	±6.3	78.0	±33.3
Mo	15.0	±3.8	17.5	±2.7
Mo + Cu	21.4	±0.9	25.4	±2.7
Mo + Fe	68.1	±3.9	63.5	±5.0
Mo + Fe + Cu	133.0	±18.0	138.3	±21.0

Adman, E.T., Godden, J.W., and Turley, S. (1995). The Structure of Copper-nitrite Reductase from
*   Achromobacter cycloclastes at Five pH Values, with NO−2 Bound and with Type II Copper*
*   Depleted. Journal of Biological Chemistry 270, 27458-27474.*
Alloway, B.J. (2008). Micronutrient Deficiencies in Global Crop Production. Dordrecht, The Netherlands:
*   Springer.*
Anderson, B., Bartlett, K., Frolking, S., Hayhoe, K., Jenkins, J., and Salas, W. (2010). "Methane and Nitrous
*   Oxide Emissions from Natural Sources". (Washington DC, USA: United States Environmental*
*   Protection Agency).*
Basiliko, N., and Yavitt, J.B. (2001). Influence of Ni, Co, Fe, and Na additions on methane production in
*   Sphagnum-dominated Northern American peatlands. Biogeochemistry 52, 133-153.*
Baumann, B., Snozzi, M., Zehnder, A.J.B., and Van Meer, J.R.D. (1996). Dynamics of denitrification activity
*   of Paracoccus denitrificans in continuous culture during aerobic-anaerobic changes. Journal of*
*   Bacteriology 178, 4367-4374.*
Beaulieu, J.J., Tank, J.L., Hamilton, S.K., Wollheim, W.M., Hall, R.O., Mulholland, P.J., Peterson, B.J.,
*   Ashkenas, L.R., Cooper, L.W., Dahm, C.N., Dodds, W.K., Grimm, N.B., Johnson, S.L., Mcdowell,*
*   W.H., Poole, G.C., Valett, H.M., Arango, C.P., Bernot, M.J., Burgin, A.J., Crenshaw, C.L., Helton,*
*   A.M., Johnson, L.T., O'brien, J.M., Potter, J.D., Sheibley, R.W., Sobota, D.J., and Thomas, S.M.*
*   (2011). Nitrous oxide emission from denitrification in stream and river networks. Proceedings of*
*   the National Academy of Sciences 108, 214-219.*
Black, A., Hsu, P.C., Hamonts, K.E., Clough, T.J., and Condron, L.M. (2016). Influence of copper on
*   expression of nirS, norB and nosZ and the transcription and activity of NIR, NOR and N2 OR in the*
*   denitrifying soil bacteria Pseudomonas stutzeri. Microb Biotechnol 9, 381-388.*
Bollinger Jr, J.M. (2010). Biochemistry: Getting the metal right. Nature 465, 40-41.
Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S. (2013).
*   Nitrous oxide emissions from soils: how well do we understand the processes and their controls?*
*   Philosophical Transactions of the Royal Society B: Biological Sciences 368.*
Chang, J., Gu, W., Park, D., Semrau, J.D., Dispirito, A.A., and Yoon, S. (2018). Methanobactin from
*   Methylosinus trichosporium OB3b inhibits N2O reduction in denitrifiers. The ISME Journal.*
Chaudhari, S.S., Kim, M., Lei, S., Razvi, F., Alqarzaee, A.A., Hutfless, E.H., Powers, R., Zimmerman, M.C.,
*   Fey, P.D., and Thomas, V.C. (2017). Nitrite Derived from Endogenous Bacterial Nitric Oxide*
*   Synthase Activity Promotes Aerobic Respiration. mBio 8.*
Conthe, M., Wittorf, L., Kuenen, J.G., Kleerebezem, R., Van Loosdrecht, M.C.M., and Hallin, S. (2018). Life
*   on N2O: deciphering the ecophysiology of N2O respiring bacterial communities in a continuous*
*   culture. ISME J 12, 1142-1153.*
Deutzmann, J.S., Stief, P., Brandes, J., and Schink, B. (2014). Anaerobic methane oxidation coupled to
*   denitrification is the dominant methane sink in a deep lake. Proceedings of the National*
*   Academy of Sciences 111, 18273-18278.*
Doroski, A.A., Helton, A.M., and Vadas, T.M. (2019). Greenhouse gas fluxes from coastal wetlands at the
*   intersection of urban pollution and saltwater intrusion: A soil core experiment. Soil Biology and*
*   Biochemistry 131, 44-53.*
Einsle, O., Messerschmidt, A., Stach, P., Bourenkov, G.P., Bartunik, H.D., Huber, R., and Kroneck, P.M.H.
*   (1999). Structure of cytochrome c nitrite reductase. Nature 400, 476.*
Felgate, H., Giannopoulos, G., Sullivan, M.J., Gates, A.J., Clarke, T.A., Baggs, E., Rowley, G., and
*   Richardson, D.J. (2012). The impact of copper, nitrate and carbon status on the emission of*
*   nitrous oxide by two species of bacteria with biochemically distinct denitrification pathways.*
*   Environmental Microbiology 14, 1788-1800.*

Fierer, N., Jackson, J.A., Vilgalys, R., and Jackson, R.B. (2005). Assessment of Soil Microbial Community
Structure by Use of Taxon-Specific Quantitative PCR Assays. Applied and Environmental
Microbiology 71, 4117-4120.
Fierer, N., Lauber, C.L., Ramirez, K.S., Zaneveld, J., Bradford, M.A., and Knight, R. (2012). Comparative
metagenomic, phylogenetic and physiological analyses of soil microbial communities across
nitrogen gradients. ISME J 6, 1007-1017.
Gauci, V., Matthews, E., Dise, N., Walter, B., Koch, D., Granberg, G., and Vile, M. (2004). Sulfur pollution
suppression of the wetland methane source in the 20th and 21st centuries. Proceedings of the
National Academy of Sciences of the United States of America 101, 12583-12587.
Giannopoulos, G., Sullivan, M.J., Hartop, K.R., Rowley, G., Gates, A.J., Watmough, N.J., and Richardson,
D.J. (2017). Tuning the modular Paracoccus denitrificans respirome to adapt from aerobic
respiration to anaerobic denitrification. Environ Microbiol 19, 4953-4964.
Giller, K.E., Witter, E., and Mcgrath, S.P. (1998). Toxicity of heavy metals to microorganisms and
microbial processes in agricultural soils: a review. Soil Biology and Biochemistry 30, 1389-1414.
Glass, J., and Orphan, V.J. (2012). Trace metal requirements for microbial enzymes involved in the
production and consumption of methane and nitrous oxide. Frontiers in Microbiology 3, 1.
Glass, J.B., Axler, R.P., Chandra, S., and Goldman, C.R. (2012). Molybdenum limitation of microbial
nitrogen assimilation in aquatic ecosystems and pure cultures. Frontiers in Microbiology 3, 331.
Granger, J., and Ward, B.B. (2003). Accumulation of nitrogen oxides in copper-limited cultures of
denitrifying bacteria. Limnology and Oceanography 48, 313-318.
Hahnke, S.M., Moosmann, P., Erb, T.J., and Strous, M. (2014). An improved medium for the anaerobic
growth of Paracoccus denitrificans Pd1222. Front Microbiol 5, 18.
Hartop, K.R., Sullivan, M.J., Giannopoulos, G., Gates, A.J., Bond, P.L., Yuan, Z., Clarke, T.A., Rowley, G.,
and Richardson, D.J. (2017). The metabolic impact of extracellular nitrite on aerobic metabolism
of Paracoccus denitrificans. Water Research 113, 207-214.
He, Z., Geng, S., Pan, Y., Cai, C., Wang, J., Wang, L., Liu, S., Zheng, P., Xu, X., and Hu, B. (2015).
Improvement of the trace metal composition of medium for nitrite-dependent anaerobic
methane oxidation bacteria: Iron (II) and copper (II) make a difference. Water Res 85, 235-243.
Heincke, M., and Kaupenjohann, M. (1999). Effects of soil solution on the dynamics of N2O emissions: a
review. Nutrient Cycling in Agroecosystems 55, 133-157.
Henry, S., Bru, D., Stres, B., Hallet, S., and Philippot, L. (2006). Quantitative detection of the nosZ gene,
encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, narG, nirK,
and nosZ genes in soils. Applied and Environmental Microbiology 72, 5181-5189.
Ippc (2014). "Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the
Fifth Assessment Report of the Intergovernmental Panel on Climate Change ", (eds.) Core Writing
Team, R.K. Pachauri & L.A. Meyer. (Geneva, Switzerland: IPPC).
Koike, I., and Hattori, A. (1975). Growth Yield of a Denitrifying Bacterium, Pseudomonas denitrificans,
under Aerobic and Denitrifying Conditions. Microbiology 88, 1-10.
Kraft, B., Tegetmeyer, H.E., Sharma, R., Klotz, M.G., Ferdelman, T.G., Hettich, R.L., Geelhoed, J.S., and
Strous, M. (2014). Nitrogen cycling. The environmental controls that govern the end product of
bacterial nitrate respiration. Science 345, 676-679.
Labbé, N., Parent, S., and Villemur, R. (2003). Addition of trace metals increases denitrification rate in
closed marine systems. Water Research 37, 914-920.
Liu, Y., Liu, Y., Zhou, H., Li, L., Zheng, J., Zhang, X., Zheng, J., and Pan, G. (2016). Abundance, composition
and activity of denitrifier communities in metal polluted paddy soils. Sci Rep 6, 19086.
Magalhães, C., Costa, J., Teixeira, C., and Bordalo, A.A. (2007). Impact of trace metals on denitrification
in estuarine sediments of the Douro River estuary, Portugal. Marine Chemistry 107, 332-341.

Magalhães, C.M., Machado, A., Matos, P., and Bordalo, A.A. (2011). Impact of copper on the diversity, abundance and transcription of nitrite and nitrous oxide reductase genes in an urban European estuary. FEMS Microbiol Ecol 77, 274-284.

Moffett, J.W., Tuit, C.B., and Ward, B.B. (2012). Chelator-induced inhibition of copper metalloenzymes in denitrifying bacteria. Limnology and Oceanography 57, 272-280.

Montoya, M., Castellano-Hinojosa, A., Vallejo, A., Álvarez, J.M., Bedmar, E.J., Recio, J., and Guardia, G. (2018). Zinc fertilizers influence greenhouse gas emissions and nitrifying and denitrifying communities in a non-irrigated arable cropland. Geoderma 325, 208-217.

Moore, C.M., Mills, M.M., Arrigo, K.R., Berman-Frank, I., Bopp, L., Boyd, P.W., Galbraith, E.D., Geider, R.J., Guieu, C., Jaccard, S.L., Jickells, T.D., La Roche, J., Lenton, T.M., Mahowald, N.M., Marañón, E., Marinov, I., Moore, J.K., Nakatsuka, T., Oschlies, A., Saito, M.A., Thingstad, T.F., Tsuda, A., and Ulloa, O. (2013). Processes and patterns of oceanic nutrient limitation. Nature Geoscience 6, 701-710.

Morley, N., Baggs, E.M., Dorsch, P., and Bakken, L. (2008). Production of NO, N_2O and N_2 by extracted soil bacteria, regulation by NO_2^- and O_2 concentrations. FEMS Microbiology Ecology 65, 102-112.

Morrissey, E.M., and Franklin, R.B. (2015). Evolutionary history influences the salinity preference of bacterial taxa in wetland soils. Front Microbiol 6, 1013.

Murrell, J.C., Mcdonald, I.R., and Gilbert, B. (2000). Regulation of expression of methane monooxygenases by copper ions. Trends in Microbiology 8, 221-225.

Okereke, G.U. (1993). Growth yield of denitrifiers using nitrous oxide as a terminal electron acceptor. World Journal of Microbiology and Biotechnology 9, 59-62.

Raudina, T.V., Loiko, S.V., Lim, A.G., Krickov, I.V., Shirokova, L.S., Istigechev, G.I., Kuzmina, D.M., Kulizhsky, S.P., Vorobyev, S.N., and Pokrovsky, O.S. (2017). Dissolved organic carbon and major and trace elements in peat porewater of sporadic, discontinuous, and continuous permafrost zones of western Siberia. Biogeosciences 14, 3561-3584.

Richardson, D., Felgate, H., Watmough, N., Thomson, A., and Baggs, E. (2009). Mitigating release of the potent greenhouse gas N(2)O from the nitrogen cycle - could enzymic regulation hold the key? Trends Biotechnol 27, 388-397.

Romero, I.C., Klein, N.J., Sañudo-Wilhelmy, S.A., and Capone, D.G. (2013). Potential trace metal co-limitation controls on N_2 Fixation and NO_3- uptake in lakes with varying trophic status. Frontiers in Microbiology 4, 54.

Rosenzweig, A.C. (2000). Nitrous oxide reductase from Cu_A to Cu_Z. Nature Structural Biology 7, 169.

Samanidou, V., and Papadoyannis, I. (1992). Study of Heavy Metal Pollution in the Waters of Axios and Aliakmon Rivers in Northen Greece. J. Environ. Sci. Health A27, 587-601.

Schlesinger, W.H. (2009). On the fate of anthropogenic nitrogen. Proc Natl Acad Sci U S A 106, 203-208.

Schoffman, H., Lis, H., Shaked, Y., and Keren, N. (2016). Iron-Nutrient Interactions within Phytoplankton. Front Plant Sci 7, 1223.

Schwarz, G., Mendel, R.R., and Ribbe, M.W. (2009). Molybdenum cofactors, enzymes and pathways. Nature 460, 839-847.

Shiro, Y. (2012). Structure and function of bacterial nitric oxide reductases: nitric oxide reductase, anaerobic enzymes. Biochim Biophys Acta 1817, 1907-1913.

Smith, C.J., Nedwell, D.B., Dong, L.F., and Osborn, A.M. (2007). Diversity and abundance of nitrate reductase genes (narG and napA), nitrite reductase genes (nirS and nrfA), and their transcripts in estuarine sediments. Appl Environ Microbiol 73, 3612-3622.

Soares, J.R., Cassman, N.A., Kielak, A.M., Pijl, A., Carmo, J.B., Lourenço, K.S., Laanbroek, H.J., Cantarella, H., and Kuramae, E.E. (2016). Nitrous oxide emission related to ammonia-oxidizing bacteria and mitigation options from N fertilization in a tropical soil. Scientific Reports 6, 30349.

Steinberg, L.M., and Regan, J.M. (2009). mcrA-targeted real-time quantitative PCR method to examine
methanogen communities. Appl Environ Microbiol 75, 4435-4442.
Sullivan, M.J., Gates, A.J., Appia-Ayme, C., Rowley, G., and Richardson, D.J. (2013). Copper control of
bacterial nitrous oxide emission and its impact on vitamin B_{12}-dependent metabolism.
Proceedings of the National Academy of Sciences 110, 19926-19931.
Takeuchi, J. (2006). Habitat Segregation of a Functional Gene Encoding Nitrate Ammonification in
Estuarine Sediments. Geomicrobiology Journal 23, 75-87.
Teklić, T., Lončarić, Z., Kovačević, V., and Singh, B.R. (2013). Metallic trace elements in cereal grain - a
review: how much metal do we eat? Food and Energy Security 2, 81-95.
Thomson, A.J., Giannopoulos, G., Pretty, J., Baggs, E.M., and Richardson, D.J. (2012). Biological sources
and sinks of nitrous oxide and strategies to mitigate emissions. Philosophical Transactions of the
Royal Society B: Biological Sciences 367, 1157-1168.
Throback, I.N., Enwall, K., Jarvis, A., and Hallin, S. (2004). Reassessing PCR primers targeting nirS, nirK
and nosZ genes for community surveys of denitrifying bacteria with DGGE. FEMS Microbiol Ecol
49, 401-417.
Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., Banger, K., Tao, B., Pan, S., Liu, M., Zhang, C.,
Bruhwiler, L., and Wofsy, S. (2015). Global methane and nitrous oxide emissions from terrestrial
ecosystems due to multiple environmental changes. Ecosystem Health and Sustainability 1, 1-20.
Twining, B.S., Mylon, S.E., and Benoit, G. (2007). Potential role of copper availability in nitrous oxide
accumulation in a temperate lake. Limnology and Oceanography 52, 1354-1366.
Van Den Berg, E.M., Rombouts, J.L., Kuenen, J.G., Kleerebezem, R., and Van Loosdrecht, M.C.M. (2017).
Role of nitrite in the competition between denitrification and DNRA in a chemostat enrichment
culture. AMB Express 7, 91.
Voigt, C., Marushchak, M.E., Lamprecht, R.E., Jackowicz-Korczyński, M., Lindgren, A., Mastepanov, M.,
Granlund, L., Christensen, T.R., Tahvanainen, T., Martikainen, P.J., and Biasi, C. (2017). Increased
nitrous oxide emissions from Arctic peatlands after permafrost thaw. Proceedings of the
National Academy of Sciences 114, 6238-6243.
White, J.R., and Reddy, K.R. (2009). "Biogeochemical Dynamics I: Nitrogen Cycling in Wetlands," in The
Wetlands Handbook, eds. E. Maltby & T. Barker. (Oxford,UK: Blackwell Publishing Ltd), 800.
Wintsche, B., Jehmlich, N., Popp, D., Harms, H., and Kleinsteuber, S. (2018). Metabolic Adaptation of
Methanogens in Anaerobic Digesters Upon Trace Element Limitation. Front Microbiol 9, 405.
Yoon, S., Cruz-Garcia, C., Sanford, R., Ritalahti, K.M., and Loffler, F.E. (2015). Denitrification versus
respiratory ammonification: environmental controls of two competing dissimilatory NO_3^-/NO_2^-
reduction pathways in Shewanella loihica strain PV-4. ISME J 9, 1093-1104.