Title

Trace metal availability affects greenhouse gas emissions and microbial functional group abundance in
freshwater wetland sediments

Authors

Georgios Giannopoulos*, Katherine R. Hartop[§], Bonnie L. Brown[‡], Rima B. Franklin

Department of Biology, Virginia Commonwealth University, Richmond, VA 23284, USA.

Correspondence: Georgios Giannopoulos (Georgios.Z.Giannopoulos@gmail.com)

Current addresses:

*Environmental Technology Laboratory, School of Chemistry, Aristotle University of Thessaloniki, 546 40
Thessaloniki, Greece.

[‡]Department of Biological Sciences, University of New Hampshire, Durham, NH 03824, USA.

[§]Royal Society of Chemistry, Cambridge CB4 0WF, UK.
15

Abstract

We investigated the effects of trace metal additions on microbial nitrogen and carbon cycling using freshwater wetland sediment microcosms amended with μM concentrations of copper (Cu), molybdenum (Mo), iron (Fe), and all combinations. In addition to monitoring inorganic nitrogen transformations (NO_3^-, NO_2^-, N_2O, NH_4^+) and carbon mineralization (CO_2, CH_4), we tracked changes in functional gene abundance associated with denitrification (*nirS*, *nirK*, *nosZ*), DNRA (*nrfA*), and methanogenesis (*mcrA*). Greater availability of Cu led to more complete denitrification (i.e., less N_2O accumulation) and a higher abundance of the *nirK* and *nosZ* genes, which encode for Cu-dependent reductases. We found sparse evidence of DNRA activity and no consistent effect on CO_2 production. Contrary, net CH_4 production was stimulated by the trace metal amendments and the Mo additions, in particular, led to increased *mcrA* gene abundance. Taken together, these findings demonstrate that trace metal effects on microbial physiology, which have heretofore only been studied in pure culture, can impact community-level function. We observed direct and indirect effects on both nitrogen and carbon biogeochemistry that culminated in increased production of greenhouse gasses, and the shifts in functional group abundance that we documented suggest these responses may have been mediated through changes in microbial community composition. Overall, this work supports a more holistic consideration of metal effects on environmental microbial communities that recognizes the key role that metal limitation plays in microbial physiology.

Keywords: trace metals, denitrification, methane, carbon mineralization, greenhouse gases, wetlands, *nirK, nirS, nosZ, nrfA, mcrA*

Introduction

Wetland microbes are important for removing many anthropogenic pollutants from surface waters, effectively preventing the contaminants from entering downstream coastal and marine ecosystems. Nitrogen (N) pollutant release (> 60 Tg N y^{-1} globally) to the environment is of particular concern because it is linked to water quality degradation, eutrophication, and increased greenhouse emissions (carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O)) (Anderson et al., 2010;Tian et al., 2015). In freshwater wetland sediments, the microbial processes of nitrification, denitrification or coupled-denitrification, methanogenesis, and methanotrophy involve N_2O and CH_4 as metabolic byproducts, in addition to CO_2 (Soares et al., 2016). Nitrous oxide is an especially potent greenhouse gas, with a global warming potential (GWP) 298 and 25 times greater per 100 years than CO_2 and CH_4, respectively, and with important implications in ozone depletion (Ravishankara et al., 2009;IPCC, 2013). Some microbes can reduce N_2O to inert N_2 as part of denitrification, though there is only one enzyme known to facilitate this reaction: the nitrous oxide reductase (NOS)(Thomson et al., 2012). It is estimated that ~17 Tg N y^{-1} is returned to the atmosphere via denitrification in wetlands and terrestrial ecosystems globally, releasing ~48 Tg N y^{-1} to coastal and marine ecosystems (Schlesinger, 2009). Wetlands are equally important for their role in carbon (C) cycling. Despite their relatively small coverage (~10 % of global land area), these ecosystems store large amounts of C and emit considerable amounts of CH_4 (144 Tg CH_4 y^{-1})(IPCC, 2014). These net CH_4 emissions represent a balance between the microbial processes of methanogenesis and methanotrophy.

The microbial production and consumption of N_2O and CH_4 is catalyzed by oxy-reductases that utilize metal co-factors (Mo, Cu, Fe) (Glass and Orphan, 2012). For denitrification, the pathway includes the reduction of NO_3^- to NO_2^-, catalyzed by a periplasmic or membrane bound molybdenum (Mo) nitrate reductase (NAR) (Schwarz et al., 2009), and the further reduction of NO_2^- to NO by either a non-metal cytochrome cd_1 (Einsle et al., 1999) or a Cu co-factor (Adman et al., 1995) nitrite reductase (NIR), expressed by *nirS* and *nirK,* respectively. NO is further reduced to N_2O by nitric oxide reductase (NOR), which contains a heme-Fe co-factor (Shiro, 2012). Finally, N_2O is reduced by a Cu-containing NOS (Rosenzweig, 2000). Similar metal complexes are key in methane cycling. For example, Fe-, Ni-, and Zn- dependent ferrodoxins and dehydrogenases catalyze the early steps of methanogenesis, and all three classes of methanogens utilize a common methyl-coenzyme M reductase (MCR; *mcrA*) to catalyze the terminal steps (Glass and Orphan, 2012). Anaerobic methanotrophs such as members of the NC10 phylum (e.g., *M. oxyfera*) are able to couple CH_4 oxidation to denitrification (NO_2^--dependent methane oxidation (N-DAMO)) utilizing a Cu-dependent particulate methane monoxygenase (pMMO) and a Fe-rich cd_1 nitrite reductase (Deutzmann et al., 2014;Cheng et al., 2019). Through these sorts of interconnected pathways and processes, the abundance

and bioavailability of trace metals in the environment can impact C and N biogeochemistry and help regulate

associated greenhouse gas emissions.

Although extensive research has considered the relative contribution of key environmental factors (e.g., pH,

$NO_{3/2}^-$, O_2, and C/N) in regulating anaerobic microbial processes and specifically denitrification (Wallenstein

et al., 2006), our understanding of the impact of metal availability is more limited. Metals in the

environment have been traditionally seen as unwanted pollutants, and trace metal accumulation has been

linked to toxicity and inhibition of ecosystem processes (Samanidou and Papadoyannis, 1992). For example,

concentrations exceeding $mg \cdot L^{-1}$ range for Cu, Mo, Fe, Zn and Pb severely inhibited denitrification in soils,

sediments, surface water bodies, and waste waters (Labbé et al., 2003;Magalhães et al., 2007;Liu et al.,

2016). However, given the dependence of many microbial enzymes on metal co-factors, it is also possible for

the opposite problem to occur – wherein function is limited due to an inadequate supply of trace metals.

This scenario has received limited study in the environment, but is well documented in case studies with

model organisms. In fact, *in vitro* studies have shown that lack of Cu, Mo, or Fe severely inhibits

denitrification or methane cycling due to the formation of non-functional enzymes typically lacking the

respective metal co-factor. For example, in the soil bacteria *Paracoccus denitrificans* and *Pseudomonas*

stutzeri, Cu is required to express to a functional NOS dimer and reduce N_2O to N_2 (Granger and Ward,

2003;Felgate et al., 2012;Black et al., 2016). In methanotrophs oxidizing CH_4 to methanol (CH_3OH), the switch

between a Cu-dependent pMMO or Fe-dependent soluble MMO is regulated by the environmental

availability of each metal (Murrell et al., 2000;Bollinger Jr, 2010).

Despite the great progress in understanding metal ecotoxicity and metalloenzyme biochemistry, our

knowledge of how trace metal availability affects microbial activity in the environment is generally limited to

selected processes (e.g., NH_4^+/NO_3^- and CO_2 assimilation) in aquatic ecosystems (Twining et al., 2007;Glass et

al., 2012;Moore et al., 2013;Romero et al., 2013;Schoffman et al., 2016). A broader understanding of how

metal availability regulates microbial processes, especially in soils and sediments, is necessary if we are to

fully appreciate environmental controls on ecosystem C and N cycling. In this study, we investigated the

effects of trace metal additions on the microbial biogeochemistry of freshwater wetland sediments focusing

primarily on NO_3^-/NO_2^- reduction and greenhouse gas kinetics, and used quantitative (qPCR) to assess

changes in the abundance of key functional groups associated with these processes. We hypothesized that

greater bioavailability of Mo, Fe, and Cu in the sediments would increase denitrification and that greater Cu

availability would reduce N_2O emissions. We also predicted that C mineralization rates would be impacted by

metal additions, specifically higher CO_2 emissions driven by greater denitrification rates. Furthermore, we

expected net CH_4 emissions to be lower due to a combination of decreased CH_4 production, since denitrifiers

will outcompete methanogens for C substrates once metal limitations are removed, and increased CH_4

consumption, because of stimulation of methanotrophs through Fe and Cu additions.

Materials and Methods

Soil samples

Five soil samples (~350 g each, 5 – 15 cm from surface and ~2 m apart) were collected using a PVC pipe

(tube dia. 10 cm and length ~15 cm) from a tidal-fluvial point bar deposit in the Pamunkey River (N

37.557451, W -76.972521) during October 2016. Samples were kept in a cooler on ice during transport to

the lab and stored overnight (4^oC) until experimental setup (below) and characterization using standard

methods (SSSA, 1996). Sediments had 25 (± 4 S.E.) % organic matter (OM), 79 (± 6) % gravimetric moisture

content, 0.18 (± 0.05) $g.cm^{-3}$ bulk density, C/N ratio of 11.0 (± 1.5), and pH 6.1 (± 0.3). Porewater, collected

from aliquots of soil via centrifugation, contained 0.10 (± 0.03) mM NO_3^-, 0.07 (± 0.02) mM NO_2^-, 0.16 (±

0.04) mM SO_4^{2-}, 0.4 (± 0.1) mM Cl^-, 0.3 (± 0.1) mM NH_4^+, 0.11 (± 0.01) mM K^+, 0.26 (± 0.10) mM Mg^{2+}, and

0.29 (± 0.10) mM Ca^{2+}.

Experimental set-up

After large stones and roots were removed, the five soil samples were combined in approximately equal

parts to form a composite sample for subsequent experiments. An aliquot (1000 g) of this composite soil was

homogenized briefly with an additional 1500 ml of filtered porewater (Cellulose, 10-μm pore-size, Millipore,

Burlington, Massachusetts, USA) using a commercial blender (30 s at max speed). Soil slurry aliquots (50 ml)

were then transferred to sterile 125-ml glass bottles (Wheaton, Millville, New Jersey, USA). Bottles were

crimped with rubber septa and incubated for 3 days in the dark (25^oC) to allow residual O_2 to be consumed

by basal respiration. Then, each bottle received a 250-μl aliquot of 2 M KNO_3 for a final calculated NO_3^-

concentration of 10 mM. According to the treatment plan (**Table 1**), replicate (*n*=4) bottles also received

aliquots (200 μl) from aqueous stock solutions of ammonium heptamolybdate ((NH_4)$_6Mo_7O_{24}$; 1 mM), iron

sulfate ($FeSO_4$; 18.5 mM) and copper sulfate ($CuSO_4$; 6.5 mM), yielding final concentrations of 28, 74 and 26

μM of Mo, Fe, and Cu respectively. Bottles were vortexed briefly (30 s at 6000 rpm), flushed with N_2 (60

min), and incubated in the dark (25^oC). Gas (5 ml) and soil slurry (1.5 ml) samples were taken anaerobically

using a sterile syringe after 0, 6, 12, 24, 48, 72, and 96 h. Following the 96 h sampling, bottles were opened

in an O_2-free chamber and 1 g of wet soil was removed and immediately frozen (-20^oC) for future molecular

analyses.

Analytical techniques

Gas samples were stored in 3-ml Exetainer vials (Labco, UK) that were previously flushed (N_2; 5 min) and

vacuumed. Concentrations of N_2O, CO_2, and CH_4 were determined with gas chromatography (GC-14A

equipped with FID, TCD and ECD; Shimadzu, Columbia, Maryland, USA) as in Morrissey and Franklin (2015).

For each gas, total production was determined as the sum of the gas accumulated in the headspace and the

gas dissolved in the slurry; the latter was estimated using the relationships described by Heincke and

Kaupenjohann (1999). Liquid samples were obtained from each slurry sample by centrifugation (10,000 x g

for 5 min) and the supernatant was diluted 10-fold, filtered (0.22-μm pore size), and stored frozen (-20^{o}C)

until ion analysis. The concentrations of NO_3^-, NO_2^-, SO_4^{2-} and NH_4^+ were determined by ion-chromatography

(ICS – 5000+, Dionex, Sunnydale, California, USA) according to the manufacturer's instructions using columns

Dionex IonPac AS17 (2 × 250 mm) and IonPac CS12 (A – 5 μm, 3 × 150 mm) and a Dionex conductivity sensor.

Molecular techniques

DNA was extracted from frozen soil samples (~1 g wet weight, equivalent to ~0.3 g dry weight) using the

DNEasy Kit from Qiagen (Germantown, Maryland, USA) following the manufacturer's instructions. The

retrieved DNA was quantified using a Nanodrop spectrometer (Thermo, Wilmington, Delaware, USA) and

visualized on a 1.2 % agarose gel for integrity. The DNA yields were, on average, 42 (± 2) ng·μl^{-1}. DNA

samples were diluted to 3 ng·μl^{-1} prior to use in quantitative PCR (qPCR) to determine the abundance of

microbial groups typically found in wetland sediments: total eubacteria (*eub-518*), nitrite reducers

(denitrification: *nirK* and *nirS*; DNRA: *nrfA*), nitrous oxide reducers (*nosZ*), and methanogens (*mcrA*).

SensiFAST™ SYBR® No-ROX Kit Polymerase 2X mix (Bioline, UK) was used according to the manufacturer's

instructions. Primers were purchased from IDT (Integrative DNA Technologies, Skokie, Illinois, USA) and the

DNA for the standard curves was extracted from isolates obtained from ATCC (American Type Culture

Collection, Manassas, Virginia, USA). Data were collected using a BioRad CFX-384 Real Time System (BioRad,

Hercules, California, USA) and analyzed using CFX Manager software (Ver 3.1).

Estimation of the combined GHG emissions (CO_2 equivalents)

To compare overall GHG emissions across treatments, we estimated the combined "CO_2-equivalents" for

CO_2, N_2O, and CH_4 based on the 100 y GWP factors (IPCC, 2013). The combined CO_2-equivalents as g C per

bottle were calculated as follows:

$$CO_{2-eqv} = V * \{(12 * [CO_2]) + (298 * 14 * [N_2O]) + (25 * 12 * [CH_4])\}$$

171   where V is the volume, 298 and 25 is the global warming potential for 100 y per weight basis in CO_2-

172   equivalents for N_2O and CH_4, respectively; 12 the molecular weight of C per mole CO_2 and CH_4; 14 the

173   molecular weight of N per mole N_2O; [CO_2] and [CH_4] are the M concentrations for CO_2 and CH_4, respectively;

174   and, lastly, [N_2O] the M concentration of N_2O per mole N.

Statistical analyses

Kinetic and molecular datasets were tested for normality (RStudio [*base*]; Shapiro.test). Non-parametric *Kruskal-Wallis* with *Bonferroni* correction statistical tests were applied to assess differences in the mean of ranks between the treatments (RStudio [*agricolae*]). Basic statistics (mean and standard error) were summarized with RStudio [*dyplr*] (Boston, Massachusetts, USA) and plotted with SigmaPlot 14 (Systat Software Inc. San Jose, California, USA). For all statistical tests, $p \leq 0.05$ was considered significant.

Results

The effect of trace metal additions on NO_3^-, NO_2^-, and NH_4^+ kinetics

At the end of the incubation, all treatments had visible gas bubbles on the surface and within the slurry, indicating microbial activity and gas production. We observed yellow and black patches within the slurry possibly due to Fe and residual Mn reduction, indicating anaerobic reducing conditions. Nitrate reduction commenced at the same time for all the treatments (12-24 h, **Figure 1**), and ~12 mM NO_3^- was consumed by the end of the incubation. At that time, 0.3 mM NO_2^- remained across all treatments while NH_4^+ ranged from 0.20 to 0.75 mM (**Figure 1**). The treatments receiving additional Mo+Fe ($p \leq 0.01$) and Mo+Fe+Cu ($p \leq 0.01$) had significantly higher NH_4^+ than the control, and an increasing trend in NH_4^+ concentration for all treatments was observed towards the end of the incubation.

The effect of trace metal additions on N_2O, CH_4 and CO_2 cumulative gas kinetics

Matching the NO_3^- consumption in the slurry, N_2O emissions increased rapidly between 24 to 48 h (**Figure 2.a)**. Notably, the control, Mo, Fe, and Mo+Fe treatments accumulated significantly more N_2O than the remaining treatments (Cu, Fe+Cu, Mo+Cu and Mo+Fe+Cu) that all contained Cu. Total N_2O concentrations reached a plateau between 48 to 96 h, most likely due to exhaustion of the available NO_3^-. CH_4 accumulated quickly in the microcosms (6 to 12 h) and continued to accumulate up to ~6 mM. By the end of the incubation, all metal additions caused an increase in CH_4 production relative to the control, though the differences were only significant between the control and the Cu, Mo+Cu and Mo+Fe+Cu treatments (**Figure 2.b**).

Cumulative CO_2 production increased most rapidly between 12 to 24 h, and continued to increase up to ~20 mM CO_2 at the end of the incubation. There were no significant differences, except between the control and the Cu, Mo+Cu, Mo+Fe+Cu treatments (Figure 2.**c**).

The effects of trace metal addition on targeted microbial groups

The average bacterial abundance was 6.7×10^8 *eub* copies·g^{-1} sediment dry weight with the Fe, Fe+Cu and Mo+Cu treatments having significantly (p=0.005) more *eub* gene copies than the control (*eubacteria;* **Figure 3**). Trace metal addition affected the abundance of *nirK* (p<0.001), *nirS* (p<0.001), and *nosZ* (p=0.001) denitrifying microbial groups. Reducers of NO_2^- utilizing Cu-NIR (*nirK*) were more abundant in all treatments when compared to the control, though the increase due to Fe addition was not significant. Microbial groups having a cytochrome cd_1-NIR (*nirS*) were relatively more abundant in the Mo and Fe+Cu treatments. The *nosZ* microbial group responsible for the reduction of N_2O to N_2 was more abundant in the treatments that received Cu. We observed no effect of trace metals on DNRA NO_2^- reducers (*nrfA*; **Figure 3**) but found that methanogens were more abundant in the treatments containing additional Mo (Mo, Mo+Cu, Mo+Fe and Mo+Fe+Cu; *mcrA*; **Figure 3**) when compared to the control.

Discussion

We observed that trace metal (Mo, Fe, Cu, and their combinations) addition to wetland sediments regulated denitrification kinetics (**Figure 1**) and greenhouse gas emissions (**Figure 2**), and altered the abundance of microbial functional groups typically associated with N removal and C cycling (**Figure 3**). Previous studies have examined the effects of trace metal availability on these processes, often reporting negative effects (Giller et al., 1998;Magalhães et al., 2007;Magalhães et al., 2011;Deng et al., 2018;Keller and Wade, 2018). The range of metal concentrations previously tested far exceeds the levels applied in the current study. The key interpretation of this experiment is that trace metal availability, specifically Cu, enhances the reduction of N_2O to N_2, stimulates the production of CH_4, and has no effect on CO_2 emissions from wetland sediments. When N_2O, CH_4, and CO_2 were combined and expressed as CO_2-equivalents, all treatments receiving a Cu addition (Cu, Mo+Cu, Fe+Cu and Mo+Fe+Cu) had significantly lower CO_2-equivalent emissions (**Figure 4**), indicating the major role of Cu in anaerobic respiration (denitrification and methanogenesis) as a regulator of nutrient cycling and greenhouse gas emissions.

Denitrification

Denitrification is an anaerobic respiratory process that utilizes NO_3^- as a terminal electron acceptor with sequential reduction: $NO_3^- \rightarrow NO_2^- \rightarrow NO \rightarrow N_2O \rightarrow N_2$. In our experiments, none of the trace metal additions had an effect on NO_3^- removal (**Figure 1**), which is somewhat surprising since all bacterial nitrate reductases contain a Mo cofactor at their active sites (Moreno-Vivian et al., 1999). This may indicate that Mo was not

limiting to the denitrifiers in this set of soils. Following NO_2^- reduction, the production and consumption of

NO is an important denitrification intermediate. Although we didn't measure NO in our experiments, we

assume that it was quickly utilized or reduced to N_2O due to its potent cytotoxicity at µM levels (Chaudhari

et al., 2017;Hartop et al., 2017). In contrast, we saw a dramatic effect of Cu addition. Regardless of what

other metals were included, Cu amendments significantly enhanced the reduction of N_2O to N_2 (**Figure 2**)

and led to an increased abundance of nitrite (*nirK*) and nitrous oxide (*nosZ*) reductace genes (**Figure 3**),

which both code for Cu-containing enzymes. The especially strong effect of Cu is because the last step of

denitrification is catalyzed by a Cu-containing reductase (NOS) and, thus far, no other alternative pathway

has been suggested (Thomson et al., 2012). In laboratory experiments, Cu has similarly been found to be a

strong driver of denitrifying metabolism. For example, Cu-deficient cells can have more transcripts of *nosZ*

and Cu-scavenging genes to compensate for the loss of N_2O reduction due to depleted Cu availability

(Felgate et al., 2012). Also, strains deficient in Cu-transporters and chaperones may be unable to reduce N_2O

to N_2, even when *nosZ* is expressed and forming a non-functional NOR (Sullivan et al., 2013).

In the treatments accumulating notable amounts of N_2O (≥ 1 mM), N_2O cytotoxicity may occur and it can be

relieved by the addition of vitamin B12, Co, or methionine (Sullivan et al., 2013). We do not exclude other

partial denitrification processes or other anaerobic processes simultaneously co-occurring with

denitrification as subjected by the residual N_2O in the Cu-containing treatments (≤2.3 mM N_2O). The

continuous accumulation of CO_2 in all treatments firstly, as proxy for microbial activity, suggests against any

indirect acute cytotoxicity from NO due to NO_2^- reduction or N_2O accumulation. Secondly, cumulative CO_2

emissions indicate that the available electron donor (organic C) was in excess of the added NO_3^-. The

magnitude of CO_2 accumulation was predictable; treatments with less residual N_2O, and thus a more

complete denitrifation pathway, had accumulated more CO_2 by the end of the incubation.

DNRA

Another possible fate of the added NO_3^- is reduction to NH_4^+ via dissimilatory nitrite reduction to ammonium

(DNRA). Over the course of the study, we saw small accumulations of NH_4^+ (<1 mM; **Figure 1**), which suggests

this process could be happening to some degree, and observed variable effects depending on the type of

metal added. The accumulation of NH_4^+ was greatest in treatments that contained both Mo and Fe (i.e.,

Mo+Fe and Mo+Fe+Cu in **Figure 1**), which may indicate a synergistic response of the enzymes that reduce

NO_3^- (all nitrate reductases are Mo dependent) and the DNRA-specific nitrite reductase (*nrfA)*, which

includes a multi-heme complex. The addition of either Mo or Fe alone did not result in NH_4^+ accumulation.

The addition of Cu (added alone, or in combination with Mo or Fe) only led to a modest increase in NH_4^+.

Overall, the accumulated NH_4^+ (<1 mM) was relatively less to the amount of NO_3^- removed (>10 mM) and the

abundance of the DNRA marker gene *nrfA* was equally low (**Figure 3**), so it is unclear whether DNRA was occurring to an appreciable extent in our microcosms. Alternatively, one could postulate that anaerobic organic matter mineralization contributed to the accumulation of NH_4^+ and that observed differences due to metal addition were indirect responses. However, if this were the case, one would expect to see a correlation between final NH_4^+ concentration and total CO_2 production, a finding which is not evident. DNRA and N mineralization commonly co-occur in wetland sediments (White and Reddy, 2009), and additional analyses would be necessary to disentangle the two.

Effect on carbon mineralization

Wetland soils are large C sinks because anaerobic conditions constrain organic matter mineralization and oxidation. The lack of electron acceptors, such as: O_2, NO_3^-, and SO_4^{2-} in freshwater wetlands, may lead to fermentative and reductive conditions were organic C will be reduced to CH_4 (Kim et al., 2015;Cheng et al., 2019). In this study, the pattern of CO_2 emissions primarily reflected the magnitude of denitrification, as discussed earlier. Also, the addition of a more bio-energetically favourable substrate (NO_3^-) did not inhibit methanogenesis, nor did the addition of Fe and Cu increase CH_4 consumption as hypothesized, considering Fe- and Cu-methane monoxygenase pathways; rather, CH_4 accumulated throughout the incubations. The addition of 10 mM NO_3^-, 5 mM NO_2^-, and 1 mM N_2O in anoxic rice field slurries inhibited methanogenesis for 30, 25, and 20 d, respectively, and after that methanogenesis resumed (Klüber and Conrad, 1998). Perhaps, the addition of 10 mM NO_3^- was not sufficient to fully inhibit methanogenesis and/or the ample electron donors from organic matter alleviated any inhibitory effects.

In terms of trace metal amendment, Mo addition enhanced the abundance of *mcrA* genes (Mo, Mo+Cu, Mo+Fe and Mo+Fe+Cu; **Figure 3**). This was expected because a Mo co-factor is typically required for the formylmethanofuran dehydrogenase (*fwd*) that catalyzes the reduction of CO_2 to formyl-methanofuran in the first step of methanogenesis by reduction of CO_2 with electrons from H_2 (Glass and Orphan, 2012). The subsequent step in methanogenesis is catalyzed by MCR, which is a Ni-containing enzyme. Since we did not add Ni to any in any of our treatments, we assumed that MCR would not change (or would decrease due to compentation with denitrification). Mo addition and the increasing partial pressure of CO_2 in the bottles during the incubation could have triggered a shift towards a Mo-based methanogenesis pathway. In general, methanogens tend to grow more slowly than denitrifers. It could be that the *mcrA* response was not as strong due to insufficient time to increase in abundance (Enzmann et al., 2018). Another interesting aspect of Cu effects on CH_4 emissions is the competition between methanotrophs and denitrifiers for the "Cu monopoly" (Chang et al., 2018), which should be further investigated in environmental samples. Thus, the

higher abundance of methanogens (*mcrA*) and noticable accumulation of CH_4 in all metal treatments than
the control indicates an important role of trace metal availability in CH_4 cycling in environmental samples.

Potential environmental implications

Previous studies have examined the effects of trace metal abundance or addition on C- and N- cycling in
sediments, peat, and agricultural soils. Typically, those studies report a negative effect; however, the range
of trace metal concentrations in those studies far exceed the levels of the current study. For instance, Keller
and Wade (2018) amended peat slurries with ~200 µM trace element solution and saw significant decrease
in CH_4 emission, most likely due to Cu-induced toxicity. Similarly, Magalhães et al. (2007) observed 85%
inhibition of denitrification accompanied with accumulation of substantial N_2O and NO_2^- at 79 µg $Cu·g^{-1}$
sediment (equiv. 1200 µM Cu) in estuaries. In a follow-up study, lower denitrification rates due to Cu (~900
µM) were accompanied by a decline in the abundance and *β*-diversity associated with the *nirK*, *nirS*, and
nosZ microbial groups (Magalhães et al., 2011). A more recent study that tested the effects of Cu pollution
(addition of 100 $µg·L^{-1}$, ~1.6 µM) on urban freshwater wetland sediments found significantly reduced CH_4
emissions, but no change in N_2O emissions (Doroski et al. (2019). This divergent observation could be due to
low NO_3^- concentration (18 µM) in relation to available C or the presence of substantial amounts of soil Cu
(9.4 $µg·g^{-1}$ sediment) at the start of the experiment in both control and Cu treatment.

The abundance of bioavailable metals in the environment, typically in their ion forms, is often observed at
several magnitude orders less compared to the routinely reported total metal content. Such trivial amounts
could be unavailable to microorganisms due to physiochemical interactions and plant uptake (Giller et al.,
1998). At neutral and alkaline pH levels, metals tend to be effectively immobilized as inorganic compounds
(metal-oxides, -hydroxides and -carbonates). Additionally, organic complexes such humic ligands are known
to bind metal cations lessening their availablity to microbes and plants. For example, the particularly low
concentrations of dissolved Cu, Fe, and Mo (~5, 1 and 0.2 $µg·L^{-1}$, respectively) and the copious dissolved
organic C (~80 $mg·L^{-1}$) in Siberian peatlands (Raudina et al., 2017) could trigger significant emissions of CH_4
and N_2O with climate warming (Basiliko and Yavitt, 2001;Voigt et al., 2017). On the other hand, in temperate
salt marshes with reducing conditions, ample SO_4^{2-} may support abundant communities of sulfate-reducing
bacteria that outcompete methanogens and thus suppress CH_4 emissions, but the H_2S produced due to SO_4^{2-}
reduction may limit the availability of trace metals through the production of insoluble metal-sulfide
complexes with potential implications in N cycling and N_2O emissions (Gauci et al., 2004;Butterbach-Bahl et
al., 2013). In this study, we added 26 µM and 74 µM of SO_4^{2-} as $CuSO_4$ and $FeSO_4$, respectively, that could
potentially have been reduced to H_2S; however, the initial and final porewater SO_4^{2-} concentrations remained

at comparable levels (average change over time across treatments was only 2.8%, **Table 3**) and no sulfide
odor was detected when microcosms were opened.

Other well-known environmental factors controlling denitrification are O_2, NH_4^+, NO_3^-/NO_2^-, and C (Thomson
et al., 2012). Synthesizing the findings from other chemostat (Baumann et al., 1996;Felgate et al.,
2012;Giannopoulos et al., 2017;Hartop et al., 2017;Conthe et al., 2018) and incubation (Koike and Hattori,
1975;Okereke, 1993;Morley et al., 2008) studies, it appears that the relative importance of the
aforementioned factors in control of denitrification is $O_2 > NH_4^+$, $NO_{3/2}^- > C/N$ or C quality, with trace metal
bioavailability exerting an additional effect on each factor. The present results suggest that the effect Cu
exerts on N_2 and N_2O emissions from the environment may be more extensive than previously thought.
These findings are important not only for understanding fundamental controls on N_2O production, but also
for making predictions about potential future emissions given increasing nutrient and metal loads associated
with urban pollution. Interestingly, the effect of trace metal availability on denitrification enzymology is also
of considerable interest in the context of agriculture and food production, and it has even been suggested
that metal additions could be a possible strategy to mitigate high N_2O emissions associated with crop
production (Richardson et al., 2009).

Conclusions and future challenges

Our results demonstrate that the importance of trace metal availability in microbial physiology can affect
community-level function, including rates of anaerobic respiration, and manifest as differences in ecosystem
processes. We found that even short-term (96 hr) exposure to increased concentrations of trace metals led
to changes in community structure (functional group abundance) and greenhouse gas emissions (N_2O, CH_4,
and CO_2). In particular, our results suggest that trace metal bioavailability may, directly or indirectly, regulate
or co-limit denitrification kinetics and favor more active microbial communities that are able to quickly
acquire the bioavailable metals and incorporate them in functional oxy-reductases. Disentangling the
potential controls of various metals on denitrification, and co-varying effects on C cycling, requires more
experiments across a range of metal concentrations and biochemical processes in a variety of different soil
types and ecosystems. Future studies should investigate microbial diversity and meta-transcriptomic profiles
under comparative treatments to understand and evaluate the ecological importance of trace metal
limitation. Such 'omic studies should be accompanied by high-throughput metabolite or process rate analysis
either in the lab or in the field.

Acknowledgements

We thank Dr. Ioannis Ipsilantis for his comments on an earlier version of this manuscript. We also thank

Colin McKenny for assistance in field and lab, Gabriella Balasa for assistance with qPCR, Olivia De Meo for

setting up the GC and HPLC procedures, and Dr. Scott Neubauer for equipment access. This work was

supported by a grant from the National Science Foundation to Franklin, Brown, and Neubauer (DEB

1355059). This paper is contribution [# to be inserted at time of publication] from the VCU Rice Rivers

Center.

Figure 1 Average NO_3^- (a, b, c, d), NO_2^- (e, f, g, h), and NH_4^+ (i, j, k, l) concentrations (mM, n=4, mean ± SE) per treatment throughout the incubations.

Figure 2 Cumulative gas concentrations of N_2O (a, d), CH_4 (b, e) and CO_2 (c, f), through the incubation and at 96 h, respectively (*n*=4; mean ± SE). Within each bar graph (d, e, f), treatments with the same letter are not significantly different as determined via Kruskal-Wallis and Bonferroni *post hoc* testing.

Figure 3 Average gene copies per g sediment (dry weight) at the end of the incubation (96 h; *n*=4; mean ±
SE). Within each graph, treatments with the same letter are not significantly different as
determined via Kruskal-Wallis and Bonferroni *post hoc* testing.

Figure 4 Average cumulative CO_2-eqv. emissions (100 y^{-1}) per treatment at the end of the incubation (96 h; *n*=4; mean ± SE). Treatments with the same letter are not significantly different.

395   **Table 1** Experimental treatments (*n*= 4).

Treatment	Nitrate (KNO$_3$; mM)	Molybdenum ((NH$_4$)$_6$Mo$_7$O$_{24}$; μM)	Iron (FeSO$_4$; μM)	Copper (CuSO$_4$; μM)
Control	10	--	--	--
Cu	10	--	--	26
Fe	10	--	74	--
Fe+Cu	10	--	74	26
Mo	10	28	--	--
Mo+Cu	10	28	--	26
Mo+Fe	10	28	74	--
Mo+Fe+Cu	10	28	74	26

Table 2 Primers and reaction conditions for qPCR assays

Target	Primer	PCR Mix (15 µl)	Standard	Reaction Conditions	Reference
Eubacteria (*16S rRNA*)	*eub*338 *eub*518	1.2 ng template 0.1 µM each primer	*Desulfovibrio desulfuricans* ATCC 27774	95°C for 4 min, then 40 cycles of 30 s at 95°C, 30 s at 55.5°C, and 60 s at 72°C.	(Fierer et al., 2005)
Denitrifiers (*nirS*)	cd3aF R3cd	10 ng DNA template 0.1 µM each primer	*Paracoccus dentrificans* ATCC 17741	95°C for 4 min, then 50 cycles of 30 s at 95°C, 30 s at 56°C and 60 s at 72°C.	(Throback et al., 2004)
Denitrifiers (*nirK*)	*nirK*q-F *nirK*1040	1.5 ng DNA template 0.35 µM each primer	*Pseudomonas nitroreducens* ATCC 13867	15 min at 95°C, 9 touchdown cycles of 95°C for 15 s, 68 °C for 60 s, and 81.5°C for 30 s (-1° per cycle for annealing); then 28 cycles of 95°C for 15 s, 60°C for 60 s, and 81.5°C for 30 s.	(Smith et al., 2007)
Denitrifiers (*nosZ*)	*nosZ*1F *nosZ*2F	10 ng DNA template 1 µM each primer	*Pseudomonas fluorescens* C7R12	15 min at 95°C, 6 cycles of 95°C for 15 s, 67°C for 30 s with a touchdown of −1°C by cycle, 72°C for 30 s, and 80°C for 15 s (acquisition data step); 40 cycles of 95°C for 15 s and 62°C for 15 s, 72°C for 30 s, and 80°C for 15 s; and 1 cycle at 95°C for 15 s and 60°C for 15 s, to 95°C for 15 s.	(Henry et al., 2006)
DNRA (*nrfA*)	*nrfA*6F *nrfA*6R	10 ng DNA template 0.3 µM each primer	*Escherichia coli* ATCC 11775	50°C for 2 min, 95°C for 8.5 min, and 50 cycles of 20 s at 94°C, 40 s at 54.5°C, and 10 s at 72°C.	(Takeuchi, 2006)
Methanogens (*mcrA*)	mlas mcrA-rev	2 ng DNA template 0.6 µM *mlas* 0.7 µM *mcrA*-rev	*Methanococcus voltae* ATCC Strain A3	95°C for 5 min, then 50 cycles of 20 s at 95°C, 20 s at 59°C, and 45 s at 72°C.	(Steinberg and Regan, 2009)

Table 3 Initial (0 h) and final (96 h) SO_4^{2-} concentrations in the various treatments (n=4, mean ± SE).

Treatment	Initial [SO_4^{2-}] μM	Final [SO_4^{2-}] μM
Control	4.9 ± 0.8	8.5 ± 3.5
Cu	23.4 ± 2.9	20.25 ± 2.0
Fe	68.9 ± 2.3	56.95 ± 8.4
Fe+Cu	99.2 ± 6.3	78.0 ± 33.3
Mo	15.0 ± 3.8	17.5 ± 2.7
Mo+Cu	21.4 ± 0.9	25.4 ± 2.7
Mo+Fe	68.1 ± 3.9	63.5 ± 5.0
Mo+Fe+Cu	133.0 ± 18.0	138.3 ± 21.0

Adman, E.T., Godden, J.W., and Turley, S. (1995). The Structure of Copper-nitrite Reductase from Achromobacter cycloclastes at Five pH Values, with NO–2 Bound and with Type II Copper Depleted. Journal of Biological Chemistry 270, 27458-27474.

Anderson, B., Bartlett, K., Frolking, S., Hayhoe, K., Jenkins, J., and Salas, W. (2010). "Methane and Nitrous Oxide Emissions from Natural Sources". (Washington DC, USA: United States Environmental Protection Agency).

Basiliko, N., and Yavitt, J.B. (2001). Influence of Ni, Co, Fe, and Na additions on methane production in Sphagnum-dominated Northern American peatlands. Biogeochemistry 52, 133-153.

Baumann, B., Snozzi, M., Zehnder, A.J.B., and Van Meer, J.R.D. (1996). Dynamics of denitrification activity of Paracoccus denitrificans in continuous culture during aerobic-anaerobic changes. Journal of Bacteriology 178, 4367-4374.

Black, A., Hsu, P.C., Hamonts, K.E., Clough, T.J., and Condron, L.M. (2016). Influence of copper on expression of nirS, norB and nosZ and the transcription and activity of NIR, NOR and N2 OR in the denitrifying soil bacteria Pseudomonas stutzeri. Microb Biotechnol 9, 381-388.

Bollinger Jr, J.M. (2010). Biochemistry: Getting the metal right. Nature 465, 40-41.

Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S. (2013). Nitrous oxide emissions from soils: how well do we understand the processes and their controls? Philosophical Transactions of the Royal Society B: Biological Sciences 368.

Chang, J., Gu, W., Park, D., Semrau, J.D., Dispirito, A.A., and Yoon, S. (2018). Methanobactin from Methylosinus trichosporium OB3b inhibits N2O reduction in denitrifiers. The ISME Journal.

Chaudhari, S.S., Kim, M., Lei, S., Razvi, F., Alqarzaee, A.A., Hutfless, E.H., Powers, R., Zimmerman, M.C., Fey, P.D., and Thomas, V.C. (2017). Nitrite Derived from Endogenous Bacterial Nitric Oxide Synthase Activity Promotes Aerobic Respiration. mBio 8.

Cheng, C., Shen, X., Xie, H., Hu, Z., Pavlostathis, S.G., and Zhang, J. (2019). Coupled methane and nitrous oxide biotransformation in freshwater wetland sediment microcosms. Sci Total Environ 648, 916-922.

Conthe, M., Wittorf, L., Kuenen, J.G., Kleerebezem, R., Van Loosdrecht, M.C.M., and Hallin, S. (2018). Life on N2O: deciphering the ecophysiology of N2O respiring bacterial communities in a continuous culture. ISME J 12, 1142-1153.

Deng, D., Hu, M., Li, L., and Huang, Y. (2018). DenitrifyingMicrobial Communities in Heavy-Metal-Contaminated Paddy Soils near Electronic-Waste Processing Centers. Water, Air, & Soil Pollution 229.

Deutzmann, J.S., Stief, P., Brandes, J., and Schink, B. (2014). Anaerobic methane oxidation coupled to denitrification is the dominant methane sink in a deep lake. Proceedings of the National Academy of Sciences 111, 18273-18278.

Doroski, A.A., Helton, A.M., and Vadas, T.M. (2019). Greenhouse gas fluxes from coastal wetlands at the intersection of urban pollution and saltwater intrusion: A soil core experiment. Soil Biology and Biochemistry 131, 44-53.

Einsle, O., Messerschmidt, A., Stach, P., Bourenkov, G.P., Bartunik, H.D., Huber, R., and Kroneck, P.M.H. (1999). Structure of cytochrome c nitrite reductase. Nature 400, 476.

Enzmann, F., Mayer, F., Rother, M., and Holtmann, D. (2018). Methanogens: biochemical background and biotechnological applications. AMB Express 8, 1-1.

Felgate, H., Giannopoulos, G., Sullivan, M.J., Gates, A.J., Clarke, T.A., Baggs, E., Rowley, G., and Richardson, D.J. (2012). The impact of copper, nitrate and carbon status on the emission of nitrous oxide by two species of bacteria with biochemically distinct denitrification pathways. Environmental Microbiology 14, 1788-1800.

Fierer, N., Jackson, J.A., Vilgalys, R., and Jackson, R.B. (2005). Assessment of Soil Microbial Community Structure by Use of Taxon-Specific Quantitative PCR Assays. Applied and Environmental Microbiology 71, 4117-4120.

Gauci, V., Matthews, E., Dise, N., Walter, B., Koch, D., Granberg, G., and Vile, M. (2004). Sulfur pollution suppression of the wetland methane source in the 20th and 21st centuries. Proceedings of the National Academy of Sciences of the United States of America 101, 12583-12587.

Giannopoulos, G., Sullivan, M.J., Hartop, K.R., Rowley, G., Gates, A.J., Watmough, N.J., and Richardson, D.J. (2017). Tuning the modular Paracoccus denitrificans respirome to adapt from aerobic respiration to anaerobic denitrification. Environ Microbiol 19, 4953-4964.

Giller, K.E., Witter, E., and Mcgrath, S.P. (1998). Toxicity of heavy metals to microorganisms and microbial processes in agricultural soils: a review. Soil Biology and Biochemistry 30, 1389-1414.

Glass, J., and Orphan, V.J. (2012). Trace metal requirements for microbial enzymes involved in the production and consumption of methane and nitrous oxide. Frontiers in Microbiology 3, 1.

Glass, J.B., Axler, R.P., Chandra, S., and Goldman, C.R. (2012). Molybdenum limitation of microbial nitrogen assimilation in aquatic ecosystems and pure cultures. Frontiers in Microbiology 3, 331.

Granger, J., and Ward, B.B. (2003). Accumulation of nitrogen oxides in copper-limited cultures of denitrifying bacteria. Limnology and Oceanography 48, 313-318.

Hartop, K.R., Sullivan, M.J., Giannopoulos, G., Gates, A.J., Bond, P.L., Yuan, Z., Clarke, T.A., Rowley, G., and Richardson, D.J. (2017). The metabolic impact of extracellular nitrite on aerobic metabolism of Paracoccus denitrificans. Water Research 113, 207-214.

Heincke, M., and Kaupenjohann, M. (1999). Effects of soil solution on the dynamics of N2O emissions: a review. Nutrient Cycling in Agroecosystems 55, 133-157.

Henry, S., Bru, D., Stres, B., Hallet, S., and Philippot, L. (2006). Quantitative detection of the nosZ gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, narG, nirK, and nosZ genes in soils. Applied and Environmental Microbiology 72, 5181-5189.

Ipcc (2013). "Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment", (eds.) T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P.M. Midgley. (Cambridge, United Kingdom and New York, NY, USA Intergovernmental Panel on Climate Change).

Ipcc (2014). "Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change ", (eds.) Core Writing Team, R.K. Pachauri & L.A. Meyer. (Geneva, Switzerland: IPPC).

Keller, J.K., and Wade, J. (2018). No evidence for trace metal limitation on anaerobic carbon mineralization in three peatland soils. Geoderma 314, 95-101.

Kim, S.Y., Veraart, A.J., Meima-Franke, M., and Bodelier, P.L.E. (2015). Combined effects of carbon, nitrogen and phosphorus on CH4 production and denitrification in wetland sediments. Geoderma 259-260, 354-361.

Klüber, H.D., and Conrad, R. (1998). Effects of nitrate, nitrite, NO and N2O on methanogenesis and other redox processes in anoxic rice field soil. FEMS Microbiology Ecology 25, 301-318.

Koike, I., and Hattori, A. (1975). Growth Yield of a Denitrifying Bacterium, Pseudomonas denitrificans, under Aerobic and Denitrifying Conditions. Microbiology 88, 1-10.

Labbé, N., Parent, S., and Villemur, R. (2003). Addition of trace metals increases denitrification rate in closed marine systems. Water Research 37, 914-920.

Liu, Y., Liu, Y., Zhou, H., Li, L., Zheng, J., Zhang, X., Zheng, J., and Pan, G. (2016). Abundance, composition and activity of denitrifier communities in metal polluted paddy soils. Sci Rep 6, 19086.

Magalhães, C., Costa, J., Teixeira, C., and Bordalo, A.A. (2007). Impact of trace metals on denitrification in estuarine sediments of the Douro River estuary, Portugal. Marine Chemistry 107, 332-341.

Magalhães, C.M., Machado, A., Matos, P., and Bordalo, A.A. (2011). Impact of copper on the diversity, abundance and transcription of nitrite and nitrous oxide reductase genes in an urban European estuary. FEMS Microbiol Ecol 77, 274-284.

Moore, C.M., Mills, M.M., Arrigo, K.R., Berman-Frank, I., Bopp, L., Boyd, P.W., Galbraith, E.D., Geider, R.J., Guieu, C., Jaccard, S.L., Jickells, T.D., La Roche, J., Lenton, T.M., Mahowald, N.M., Marañón, E., Marinov, I., Moore, J.K., Nakatsuka, T., Oschlies, A., Saito, M.A., Thingstad, T.F., Tsuda, A., and Ulloa, O. (2013). Processes and patterns of oceanic nutrient limitation. Nature Geoscience 6, 701-710.

Moreno-Vivian, C., Cabello, P., Martinez-Luque, M., Blasco, R., and Castillo, F. (1999). Prokaryotic nitrate reduction: molecular properties and functional distinction among bacterial nitrate reductases. J Bacteriol 181, 6573-6584.

Morley, N., Baggs, E.M., Dorsch, P., and Bakken, L. (2008). Production of NO, N_2O and N_2 by extracted soil bacteria, regulation by NO_2^- and O_2 concentrations. FEMS Microbiology Ecology 65, 102-112.

Morrissey, E.M., and Franklin, R.B. (2015). Evolutionary history influences the salinity preference of bacterial taxa in wetland soils. Front Microbiol 6, 1013.

Murrell, J.C., Mcdonald, I.R., and Gilbert, B. (2000). Regulation of expression of methane monooxygenases by copper ions. Trends in Microbiology 8, 221-225.

Okereke, G.U. (1993). Growth yield of denitrifiers using nitrous oxide as a terminal electron acceptor. World Journal of Microbiology and Biotechnology 9, 59-62.

Raudina, T.V., Loiko, S.V., Lim, A.G., Krickov, I.V., Shirokova, L.S., Istigechev, G.I., Kuzmina, D.M., Kulizhsky, S.P., Vorobyev, S.N., and Pokrovsky, O.S. (2017). Dissolved organic carbon and major and trace elements in peat porewater of sporadic, discontinuous, and continuous permafrost zones of western Siberia. Biogeosciences 14, 3561-3584.

Ravishankara, A.R., Daniel, J.S., and Portmann, R.W. (2009). Nitrous Oxide (N2O): The Dominant Ozone-Depleting Substance Emitted in the 21st Century. Science 326, 123.

Richardson, D., Felgate, H., Watmough, N., Thomson, A., and Baggs, E. (2009). Mitigating release of the potent greenhouse gas N(2)O from the nitrogen cycle - could enzymic regulation hold the key? Trends Biotechnol 27, 388-397.

Romero, I.C., Klein, N.J., Sañudo-Wilhelmy, S.A., and Capone, D.G. (2013). Potential trace metal co-limitation controls on N_2 Fixation and NO_3- uptake in lakes with varying trophic status. Frontiers in Microbiology 4, 54.

Rosenzweig, A.C. (2000). Nitrous oxide reductase from Cu_A to Cu_Z. Nature Structural Biology 7, 169.

Samanidou, V., and Papadoyannis, I. (1992). Study of Heavy Metal Pollution in the Waters of Axios and Aliakmon Rivers in Northen Greece. J. Environ. Sci. Health A27, 587-601.

Schlesinger, W.H. (2009). On the fate of anthropogenic nitrogen. Proc Natl Acad Sci U S A 106, 203-208.

Schoffman, H., Lis, H., Shaked, Y., and Keren, N. (2016). Iron-Nutrient Interactions within Phytoplankton. Front Plant Sci 7, 1223.

Schwarz, G., Mendel, R.R., and Ribbe, M.W. (2009). Molybdenum cofactors, enzymes and pathways. Nature 460, 839-847.

Shiro, Y. (2012). Structure and function of bacterial nitric oxide reductases: nitric oxide reductase, anaerobic enzymes. Biochim Biophys Acta 1817, 1907-1913.

Smith, C.J., Nedwell, D.B., Dong, L.F., and Osborn, A.M. (2007). Diversity and abundance of nitrate reductase genes (narG and napA), nitrite reductase genes (nirS and nrfA), and their transcripts in estuarine sediments. Appl Environ Microbiol 73, 3612-3622.

Soares, J.R., Cassman, N.A., Kielak, A.M., Pijl, A., Carmo, J.B., Lourenço, K.S., Laanbroek, H.J., Cantarella, H., and Kuramae, E.E. (2016). Nitrous oxide emission related to ammonia-oxidizing bacteria and mitigation options from N fertilization in a tropical soil. Scientific Reports 6, 30349.

Sssa (1996). Methods of Soil Analysis Part 3—Chemical Methods. Madison, WI: Soil Science Society of America, American Society of Agronomy.

Steinberg, L.M., and Regan, J.M. (2009). mcrA-targeted real-time quantitative PCR method to examine methanogen communities. Appl Environ Microbiol 75, 4435-4442.

Sullivan, M.J., Gates, A.J., Appia-Ayme, C., Rowley, G., and Richardson, D.J. (2013). Copper control of bacterial nitrous oxide emission and its impact on vitamin B_{12}-dependent metabolism. Proceedings of the National Academy of Sciences 110, 19926-19931.

Takeuchi, J. (2006). Habitat Segregation of a Functional Gene Encoding Nitrate Ammonification in Estuarine Sediments. Geomicrobiology Journal 23, 75-87.

Thomson, A.J., Giannopoulos, G., Pretty, J., Baggs, E.M., and Richardson, D.J. (2012). Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. Philosophical Transactions of the Royal Society B: Biological Sciences 367, 1157-1168.

Throback, I.N., Enwall, K., Jarvis, A., and Hallin, S. (2004). Reassessing PCR primers targeting nirS, nirK and nosZ genes for community surveys of denitrifying bacteria with DGGE. FEMS Microbiol Ecol 49, 401-417.

Tian, H., Chen, G., Lu, C., Xu, X., Ren, W., Zhang, B., Banger, K., Tao, B., Pan, S., Liu, M., Zhang, C., Bruhwiler, L., and Wofsy, S. (2015). Global methane and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental changes. *Ecosystem Health and Sustainability* 1**,** 1-20.

Twining, B.S., Mylon, S.E., and Benoit, G. (2007). Potential role of copper availability in nitrous oxide accumulation in a temperate lake. *Limnology and Oceanography* 52**,** 1354-1366.

Voigt, C., Marushchak, M.E., Lamprecht, R.E., Jackowicz-Korczyński, M., Lindgren, A., Mastepanov, M., Granlund, L., Christensen, T.R., Tahvanainen, T., Martikainen, P.J., and Biasi, C. (2017). Increased nitrous oxide emissions from Arctic peatlands after permafrost thaw. *Proceedings of the National Academy of Sciences* 114**,** 6238-6243.

Wallenstein, M.D., Myrold, D.D., Firestone, M., and Voytek, M. (2006). Environmental controls on denitrifying communities and denitrification rates: insights from molecular methods. *Ecological Applications* 16**,** 2143-2152.

White, J.R., and Reddy, K.R. (2009). "Biogeochemical Dynamics I: Nitrogen Cycling in Wetlands," in *The Wetlands Handbook*, eds. E. Maltby & T. Barker. (Oxford,UK: Blackwell Publishing Ltd), 800.