Expanding the watch list for potential Ebola virus antibody escape mutations

Jagdish Suresh Patel[1,2*], Caleb J. Quates[1], Erin L. Johnson[1], F. Marty Ytreberg[1,3,4*]

[1]Center for Modeling Complex Interactions, University of Idaho, Moscow, Idaho, United States of America

[2]Department of Biological Sciences, University of Idaho, United States of America

[3]Department of Physics, University of Idaho, Moscow, Idaho, United States of America

[4]Institute for Bioinformatics and Evolutionary Biology, University of Idaho, Moscow, Idaho, United States of America

* Corresponding authors:

Email: thejagdishpatel@gmail.com (JSP)

Email: ytreberg@uidaho.edu (FMY)

23Abstract

24The 2014 outbreak of Ebola virus (EBOV) in Western Africa is the largest recorded filovirus

25disease outbreak and lead to the death of over 11,000 people. This deadly virus still poses a

26grave epidemic threat as evidenced by the current (since May 2018) EBOV outbreak in the

27Democratic Republic of the Congo which has already claimed the lives of over 250 people. One

28important strategy for combating EBOV epidemics is to anticipate how the evolution of EBOV

29might undermine treatment since the development of vaccines and antibody therapies are

30typically based on a single strain (often the 1976 Mayinga) of the EBOV envelope glycoprotein

31(GP). In a previous study we initiated a watch list of potential antibody escape mutations of

32EBOV by modeling interactions between EBOV GP and the monoclonal antibody KZ52. This

33watch list was generated using molecular modeling to estimate the effect of every possible

34mutation of GP. The final watch list containing 34 mutations were predicted to disrupt GP-KZ52

35binding but not to disrupt the ability of GP to fold and to form trimers. In this study, we expand

36our watch list by including three more monoclonal antibodies with distinct epitopes on GP,

37namely Antibody 100 (Ab100), Antibody 114 (Ab114) and 13F6-1-2. Our updated watch list

38contains 127 mutations, three of which have been seen in humans or are experimentally

39associated with reduced efficacy of antibody treatment. We believe mutations on this broad

40watch list require attention since they may be a signal of an evolutionary response from EBOV to

41treatment that could diminish the effectiveness of interventions.

Introduction

Ebola virus (EBOV) has five known species: Zaire EBOV, Sudan EBOV, Ta Forest EBOV, Bundibugyo EBOV, and Reston EBOV and can cause disease with up to 90% case fatality rates.[1, 2] The outbreak of Zaire EBOV infection in Western Africa between 2014 to 2016 is the largest recorded filovirus disease outbreak with death typically occurring 5-7 days after the appearance of symptoms.[3] This outbreak receded in 2016, but EBOV still constitutes an epidemic threat as evidenced by the current (since May 2018) Zaire EBOV outbreak in the Democratic Republic of the Congo which has already claimed lives of 271 people and more than 458 positive cases (WHO report, 4th December 2018).[4]

The 2014–2016 EBOV outbreak in West Africa led to clinical development of therapeutics and vaccines.[5, 6] The success of ZMapp, a cocktail of three chimeric monoclonal antibodies (mAbs) derived from immunized mice, in nonhuman primates (NHP) demonstrated the potential of mAb therapies against EBOV infection, and ZMapp is currently undergoing human trials.[7-9] Subsequently, several other mAbs were designed and demonstrated protection against EBOV infection and some of them are currently being used to manage the current EBOV outbreak. (https://www.nih.gov/news-events/news-releases/nih-begins-testing-ebola-treatment-early-stage-trial)[10, 11] To prepare for future outbreaks it is critical to anticipate and monitor EBOV evolution since it could lead to antibody escape mutants that could compromise treatment efforts. Sequencing studies have revealed that there is a significant genetic variation in EBOV. Most of the variation is found in the EBOV glycoprotein (GP) that is the target for the all mAbs currently under development and being used in the management of the current EBOV outbreak.[12-14]

67To address the potential threat of EBOV evolution outpacing antibody treatment efforts we

68previously initiated a watch list of potential antibody escape mutants for the EBOV GP.[15] We

69focused on the KZ52 mAb as it had a 3-D experimental structure bound to EBOV GP available.

70In this previous study, we combined use of FoldX software with molecular dynamics simulations

71to estimate folding and binding stabilities. We placed 34 mutations on the watch list by

72considering every possible EBOV GP mutation and choosing those that disrupt binding between

73GP and KZ52 but do not disrupt the ability of GP to fold and bind to form a complex. One of

74these 34 mutations (*N550K*) was seen in humans in a previous outbreak.

76The aim of this study is to expand our watch list of potential antibody escape mutants for EBOV

77GP. After we published our previous watch list, three more 3-D structures of mAbs interacting

78with different epitopes on EBOV GP (necessary for our molecular modeling approach) were

79published in the Protein Data Bank (PDB). In this study, we expand our watch list to include

80possible antibody escape mutations from three antibodies with distinct epitopes: Antibody 100

81(Ab100), Antibody 114 (Ab114) and 13F6-1-2. Our watch list now contains 127 mutations, three

82of which have been seen in humans or are experimentally associated with reduced efficacy of

83antibody treatment.

84Methods

85For a mutation to be placed on a watch list for EBOV it must: (1) disrupt binding to a protective

86antibody, and (2) leave the viral proteins functional. It is thus necessary to determine how amino

87acid mutations alter stabilities ($\Delta\Delta G$ values) for GP folding, forming a trimer and binding to the

88antibody. In our previous work, we have obtained $\Delta\Delta G$ values of GP folding, GP trimer

89formation and GP binding to KZ52 antibody using our molecular dynamics (MD) plus FoldX

90approach.[15] In this study we calculate $\Delta\Delta G$ values of binding for Ab100, Ab114 and 13F6-1-2

91antigen-antibody complexes using the same modeling approach.

92**Structure Preparation**

93**EBOV GP – Ab100 and Ab114 complexes**

94Structures of Ab100 and Ab114 bound to EBOV GP were obtained from the Protein Data Bank

95(PDB) accession number 5FHC.[16] The EBOV GP amino acid sequence was based on the 1976

96Mayinga strain, just like our previous study. The PDB file 5hfc.pdb contains coordinates of both

97Ab100 and Ab114 bound to EBOV GP (GP1 & GP2) monomer at different sites. This file was

98first modified to remove all but one copy each of GP1, GP2, antibody light chain and antibody

99heavy chain of Ab100 and Ab114 (one third of the GP-Ab100/Ab114 trimeric complex), and

100then was split into two files where one file had the GP – Ab100 complex and the other had the

101GP – Ab114 complex. The MODELLER software[17] was then used to build the missing

102residues. This included residues 190 – 213 that are predicted to be intrinsically disordered; the

103resulting complexes had no secondary structure content in this region. The full EBOV trimer

104protein complex was then created using the symexp command in PyMOL (see Figs 1A and 2A)

105and contained three copies each of GP1 (residues 33 - 278), GP2 (residues 502 – 599), heavy and

106light chains of the antibody.

108**Fig 1.** (A) Structure of EBOV GP trimer in complex with the Ab100 antibody. GP1 is gray, GP2

109is red and the heavy and the light chains of Ab100 are brown. (B) & (C) Mutations in the GP

110complex with $\Delta\Delta G_{Bind} > 2$ kcal/mol (i.e., above black dashed line) are considered disruptive and

111are highlighted using green stick representation. (D) $\Delta\Delta G_{Bind}$ values (gray circles) for all 19

112possible mutations at each site of GP1 (33 – 278) and GP2 (502 – 599). Different colors and

113counts in the legend indicate locations and number of mutations with $\Delta\Delta G_{\text{Bind}} > 2$ kcal/mol on

114the GP complex.

116**Fig 2.** (A) Structure of EBOV GP trimer in complex with the Ab114 antibody. GP1 is gray, GP2

117is yellow and the heavy and the light chains of Ab114 are orange. (B) Mutations in the GP

118complex with $\Delta\Delta G_{\text{Bind}} > 2$ kcal/mol (i.e., above black dashed line) are considered disruptive and

119are highlighted using green stick representation. (D) $\Delta\Delta G_{\text{Bind}}$ values (gray circles) for all 19

120possible mutations at each site of GP1 (33 – 278) and GP2 (502 – 599). Different colors and

121counts in the legend indicate locations and number of mutations with $\Delta\Delta G_{\text{Bind}} > 2$ kcal/mol on

122the GP complex.

123**EBOV mucin-like domain peptide – 13F6-1-2 antibody complex**
124Crystal structure of the 13F6-1-2 Fab fragment bound to its EBOV GP mucin-like domain (GP

125MLD) peptide epitope (11 amino acids, VEQHHRRTDND) was downloaded from the PDB

126using the 2QHR accession number.[18] Unlike other epitopes used in our previous and current

127study, this structure was based on the 1976 Eckron strain. However, the alignment of the 11-

128residue long peptide was 100% identical to the 1976 Mayinga strain. The PDB file 2qhr.pdb was

129edited to remove everything except GP MLD peptide (residues 404 – 414) and heavy and light

130chains of 13F6-1-2 antibody. (see Fig 3A) WHAT IF web server (https://swift.cmbi.umcn.nl)

131was then used to add the missing atoms to the complex structure.

133**Fig 3.** (A) Structure of EBOV GP MLD peptide bound to the 13F6-1-2 antibody. GP MLD

134peptide is in gray tube representation, and the heavy and the light chains of 13F6-1-2 are black.

135Mutations in the GP MLD peptide with $\Delta\Delta G_{\text{Bind}} > 2$ kcal/mol (i.e., above black dashed line) are

136considered disruptive and are highlighted using green stick representation. (B) $\Delta\Delta G_{\text{Bind}}$ values

137(gray circles) for all 19 possible mutations at each of the 11 sites of GP MLD peptide (404 –

138414). Different colors and counts in the legend indicate locations and number of mutations with

139$\Delta\Delta G_{\text{Bind}}$ >2 kcal/mol on the GP complex.

141**Molecular Dynamics Simulations**
142All three EBOV – antibody complexes were subjected to atomistic MD simulations using the

143protocol reported in our previous study. Briefly, the software package GROMACS 5.0.7[19] was

144used for all MD simulations with the Charmm22* forcefield.[20] GP MLD peptide – 13F6-1-2

145simulations were 100 ns, and the GP – Ab100 and Ab114 production simulations were run for a

146shorter 50 ns due to the large number of atoms in the final simulation box. During the production

147simulation snapshots were saved every 1 ns resulting in 50 snapshots for each of the GP – Ab100

148and GP – Ab114 systems, and 100 snapshots for the peptide – 13F6-1-2 complex.

149**FoldX**
150Snapshots of all three complexes were analyzed using FoldX software.[21, 22] As with our

151previous study, we initially minimize each snapshot six times in succession using the RepairPDB

152command to obtain convergence of the potential energy. BuildModel command was then used to

153generate all possible 19 single mutations in EBOV GP and GP MLD peptide at each amino acid

154site. Finally, the binding stability of the protein complex due to each mutation was estimated

155using AnalyseComplex command. For each mutation, we then estimated $\Delta\Delta G_{\text{Bind}}$ by averaging

156the FoldX results across all individual snapshot estimates.

158To calculate $\Delta\Delta G_{\text{Bind}}$ values for all possible 19 mutations at amino acid site of EBOV GP and GP

159MLD peptide, we carried out 980,400 FoldX calculations (344 GP residues × 19 possible

160mutations at each site × 50 MD snapshots × 3 copies of GP-Antibody in a complex) for each GP

161– Ab100 and GP – Ab114 complexes and 20,900 calculations (11-residue MLD peptide × 19

162possible mutations × 100 MD snapshots) for GP MLD peptide – 13F6-1-2 antibody complex.

163Averaging estimates across all individual snapshots ultimately resulted in 6,441 $\Delta\Delta G_{\text{Bind}}$ values

164each for Ab100 and Ab114 antibody complexes and 209 $\Delta\Delta G_{\text{Bind}}$ values for 13F6-1-2 antibody

165complex. (see SI Files)

166Results and discussion

167We have expanded the watch list generated by us in a previous study by including antibody

168escape mutations against three additional antibodies interacting with EBOV GP. These watch list

169mutations are those predicted to both disrupt GP - antibody binding and yet allow GP to fold and

170form trimers.

172The EBOV GP is a class I fusion protein consisting of disulfide-linked subunits, GP1 and GP2,

173that bind to form a chalice-shaped trimer. (see Figs 1A & 2A) Ab100 interacts at the base of the

174GP trimer (see Fig 1A). This interaction is similar to that of KZ52, the prototypic neutralizing

175antibody used in our previous study. However, Ab100 contacts GP1 and GP2 of a monomer and

176the disordered (residue 190 -213) loop of the neighboring monomer (see Fig 1A) in contrast to

177KZ52 which interacts with GP2 of a single monomer. The epitope for Ab114 spans both the

178glycan cap and the inner chalice of GP, (see Fig 2A) where it remains bound after proteolytic

179cleavage of the glycan cap and prevents interaction of cleaved GP to its host receptor.[16] 13F6-

1801-2 is a monoclonal antibody that binds to an 11-residue peptide located in the heavily

181glycosylated mucin-like domain (MLD) of the EBOV GP.[18] (see Fig 3A)

183Fig 4 is a graphical representation of our expanded watch list of EBOV mutations which includes

184KZ52, Ab100, Ab114 and 13F6-1-2 antibody complexes. The maximum functional stability

185change (i.e., folding and trimer formation) for all mutations is plotted against the corresponding

186change in the antibody binding affinity. The 127 mutations in the lower right quadrant are those

187that belong on the watch list since they are classified as both functional and disruptive to

188antibody binding. There were 21, 33, 39 and 34 mutations respectively in Ab100, Ab114, 13F6-

1891-2 and KZ52 antibody complexes. The specific mutations on the watch list are given in the

190Table 1 and shows that they are concentrated at just six residues in KZ52 and at five residues in

191each of the other three antibody complexes. (see Figs 1B & 1C, Fig 2B and Fig 3A) All of these

192mutations lie at the interface as one might intuitively expect from the structure, however, most

193mutations at the interface are not predicted to disrupt antibody binding.

195**Fig 4.** Maximum of folding stability, dimer binding stability (interaction of GP1 and GP2) or

196trimer binding stability (interaction of a GP1-GP2 dimer with other dimers), $\Delta\Delta G_{Max}$, as a

197function of $\Delta\Delta G_{Bind}$ for all antibody complexes. $\Delta\Delta G_{Max}$ values are considered to be zero for the

198intrinsically disordered 11-residue MLD peptide. Symbols in the inset legend indicate the

199corresponding antibody. Watch list mutations are shown as colored symbols and are predicted to

200disrupt binding to any one of the four antibodies (KZ52, Ab100, Ab114 and 13F6-1-2) but not to

201disrupt GP folding and trimer formation. Consistent with our previous study, mutations with

202$\Delta\Delta G_{Bind} > 2$ kcal/mol are considered disruptive to antibody binding and those with $-3 < \Delta\Delta G_{Max}$

203< 3 kcal/mol are considered functional. The number of watch list mutations associated with each

204antibody is shown in the legend.

206**Table 1. Watch list mutations**

Systems	Amino acid site	Predicted Escape Mutations
EBOV GP – KZ52	N506	W, Y
	P513	H, W
	N550	Q, K, P, F, H, I, E, R, W, V, Y, M
	P552	S, Q, K, T, F, A, H, G, R, W, V, Y
	G553	M
	G557	F, H, R, W, Y
EBOV GP – Ab100	P34	W, Y
	D47	R
	P198	F, W, Y, D, H
	E564	K, P, S, T, Y, A, R, G
	A568	F, W, Y, R, H
EBOV GP – Ab114	P116	F, W, Y, R, D, G, H
	G118	L, K, M, F, S, W, Y, R, N, C, Q, E, H
	T144	F, P, W, Y, D, E, H
	T223	F, W, Y, R, H
	L233	W
GP MLD peptide – 13F6-1-2	Q406	R, D, G, H, F, S, T, W, Y
	H407	P
	R409	A, N, D, C, Q, E, G, H, I, L, K, M, F, P, S, T, W, Y, V
	R410	G, W
	R411	R, Q, G, H, K, F, W, Y

208**Table 1.** Watch list mutations are shown for all four antibodies (KZ52, Ab100, Ab114 and 13F6-
2091-2) at a predicted amino acid sites on EBOV GP.

211If any of the mutations in Table 1 were to appear in a real population, it would indicate a possible

212increased risk of escaping the normal immune response or render treatments with these

213antibodies ineffective. Mutation *N550K* against KZ52 has already appeared in humans and

214thought to have been infected by gorillas in Central Africa between 2001 and 2003. This

215mutation is present in all sequenced isolates from that outbreak.

217Davidson et al.[23] conducted an alanine-scanning mutagenesis study on GP that can be

218qualitatively compared to our work. They individually mutated each residue of the GP protein to

219alanine and measured changes in GP-KZ52 binding affinities relative to the unmutated form.

220They identified five residues that are critical for KZ52 antibody binding: C511, N550, D552,

221G553, and C556. Three of these sites are found on our watch list in Table 1 (N550, D552, and

222G553) and 25 of the 34 (74%) watch list mutations are found at these three sites.

224In the study of MB-003,[24] a plant-derived monoclonal antibody cocktail composed of c13C6,

22513F6-1-2, and c6D8 used effectively in treatment of EBOV virus infection in non-human

226primates, was unable to protect two of six animals when initiated one- or two-days post-

227infection. Investigation of a mechanism of viral escape in one of the animals shown five

228nonsynonymous mutations in the monoclonal antibody target sites. Among these mutations

229*Q406R* and *R409C* were linked to a reduction in 13F6-1-2 antibody binding.[24] Both of these

230mutations were correctly identified by our modeling strategy and are present on the watch list.

232In spite of our efforts to expand the watch list, it remains incomplete and putative for several

233reasons. The list covers only four epitopes for which experimental structures of antibodies

234interacting with viral proteins are available. However, there are other epitopes known for EBOV

235GP.[10] With more experimental structures it would be possible to further expand the watch list.

236The size of the watch list depends on how we define the threshold for antibody binding (Fig 4).

237If the threshold is lowered from 2.0 kcal/mol, the watch list will naturally grow in size. The

238reasoning behind using a different cutoff for functional as compared to antibody binding is

239described in our previous study. The watch list only includes mutations that are individually

240predicted to disrupt antibody binding while remaining functional, however, it is possible that

241immune escape could arise by the cumulative effect of several changes on antibody binding.

242How multiple substitutions interact to produce cumulative effects on stability is not well

243understood and is an important consideration for future studies. The watch list has not been

244experimentally validated either in terms of mutational effects on GP folding and binding

245affinities, nor on the downstream immune system consequences. Our hope is that this work will

246encourage such research.

248In summary, we have expanded the watch list of potential antibody escape mutations of EBOV

249by including three more antibody complexes: EBOV GP – Ab100, GP – Ab114 and GP Mucin-

250like domain (GP MLD) peptide – 13F6-1-2. The watch list now contains 127 mutations in 21

251sites in EBOV GP. Mutations from the watch list that appear during an outbreak deserve

252attention since they may be a signal of an evolutionary response from the virus that could reduce

253the efficacy of treatment efforts. Ab114 has been recently approved for the first time to treat

254infected individuals during the current EBOV outbreak in Democratic Republic of Congo.

255(https://www.nih.gov/news-events/news-releases/nih-begins-testing-ebola-treatment-early-stage-

256trial) We hope our watch list will serve as a useful reference for the public health and emerging

257infectious disease monitoring agencies. The watch list can still be expanded if more experimental

258structures of EBOV – antibody complexes become available. In fact, as we were preparing this

259manuscript, the crystal structure of mAb CA45 bound to GP1 and GP2 interface of EBOV GP

260were published by Janus et al.[25] Lastly, we believe our in-silico approach could be applied to

261determine watch lists for other viruses provided experimental structures are available and for

262future design and optimization efforts of antibodies.

Acknowledgements

This research was supported by the Center for Modeling Complex Interactions sponsored by the NIGMS under award number P20 GM104420 and National Science Foundation EPSCoR Track-II under award number OIA1736253. Computer resources were provided in part by the Institute for Bioinformatics and Evolutionary Studies Computational Resources Core sponsored by the National Institutes of Health (P30 GM103324). This research also made use of the computational resources provided by the high-performance computing center at Idaho National Laboratory, which is supported by the Office of Nuclear Energy of the U.S. DOE and the Nuclear Science User Facilities under Contract No. DE-AC07-05ID14517. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1.	Holmes EC, Dudas G, Rambaut A, Andersen KG. The evolution of Ebola virus: Insights from the 2013–2016 epidemic. Nature. 2016;538:193.

2.	Reynolds P, Marzi A. Ebola and Marburg virus vaccines. Virus Genes. 2017;53(4):501-15.

3.	Wong G, He S, Leung A, Cao W, Bi Y, Zhang Z, et al. Naturally Occurring Single Mutations in Ebola Virus Observably Impact Infectivity. Journal of Virology. 2019;93(1):e01098-18.

4.	Maxmen A. Ebola detectives race to identify hidden sources of infection as outbreak spreads. Nature. 2018;564(7735):174-5. PubMed PMID: 30531881.

5.	Saphire EO, Schendel SL, Gunn BM, Milligan JC, Alter G. Antibody-mediated protection against Ebola virus. Nature Immunology. 2018;19(11):1169-78.

6.	Mirza MU, Vanmeert M, Ali A, Iman K, Froeyen M, Idrees M. Perspectives towards antiviral drug discovery against Ebola virus. Journal of Medical Virology. 0(ja).

7.	Borio L, Cox E, Lurie N. Combating Emerging Threats — Accelerating the Availability of Medical Therapies. New England Journal of Medicine. 2015;373(11):993-5. PubMed PMID: 26244879.

8.	Qiu X, Wong G, Audet J, Bello A, Fernando L, Alimonti JB, et al. Reversion of advanced Ebola virus disease in nonhuman primates with ZMapp. Nature. 2014;514:47.

9.	Mendoza EJ, Racine T, Kobinger GP. The ongoing evolution of antibody-based treatments for Ebola virus infection. Immunotherapy. 2017;9(5):435-50. PubMed PMID: 28357917.

10.	Saphire EO, Schendel SL, Fusco ML, Gangavarapu K, Gunn BM, Wec AZ, et al. Systematic Analysis of Monoclonal Antibodies against Ebola Virus GP Defines Features that Contribute to Protection. Cell. 2018;174(4):938-52.e13.

11.	Corti D, Misasi J, Mulangu S, Stanley DA, Kanekiyo M, Wollen S, et al. Protective monotherapy against lethal Ebola virus infection by a potently neutralizing antibody. Science. 2016:aad5224.

12.	Gire SK, Goba A, Andersen KG, Sealfon RSG, Park DJ, Kanneh L, et al. Genomic surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. Science. 2014;345(6202):1369-72.

13.	Tong Y-G, Shi W-F, Liu D, Qian J, Liang L, Bo X-C, et al. Genetic diversity and evolutionary dynamics of Ebola virus in Sierra Leone. Nature. 2015;524:93.

14. Park Daniel J, Dudas G, Wohl S, Goba A, Whitmer Shannon LM, Andersen Kristian G, et al. Ebola Virus Epidemiology, Transmission, and Evolution during Seven Months in Sierra Leone. Cell. 2015;161(7):1516-26.

15. Miller CR, Johnson EL, Burke AZ, Martin KP, Miura TA, Wichman HA, et al. Initiating a watch list for Ebola virus antibody escape mutations. PeerJ. 2016;4:e1674.

16. Misasi J, Gilman MSA, Kanekiyo M, Gui M, Cagigi A, Mulangu S, et al. Structural and molecular basis for Ebola virus neutralization by protective human antibodies. Science. 2016;351(6279):1343-6.

17. Webb B, Sali A. Comparative Protein Structure Modeling Using MODELLER. Current protocols in bioinformatics. 2016;54:5.6.1-5.6.37. PubMed PMID: 27322406; PubMed Central PMCID: PMCPMC5031415.

18. Lee JE, Kuehne A, Abelson DM, Fusco ML, Hart MK, Saphire EO. Complex of a Protective Antibody with Its Ebola Virus GP Peptide Epitope: Unusual Features of a Vλx Light Chain. Journal of Molecular Biology. 2008;375(1):202-16.

19. Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJ. GROMACS: fast, flexible, and free. Journal of computational chemistry. 2005;26(16):1701-18. PubMed PMID: 16211538.

20. Chen J, Im W, Brooks CL. Balancing Solvation and Intramolecular Interactions: Toward a Consistent Generalized Born Force Field. Journal of the American Chemical Society. 2006;128(11):3728-36.

21. Schymkowitz J, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. The FoldX web server: an online force field. Nucleic acids research. 2005;33(Web Server issue):W382-8. PubMed PMID: 15980494; PubMed Central PMCID: PMCPMC1160148.

22. Guerois R, Nielsen JE, Serrano L. Predicting Changes in the Stability of Proteins and Protein Complexes: A Study of More Than 1000 Mutations. Journal of Molecular Biology. 2002;320(2):369-87.

23. Davidson E, Bryan C, Fong RH, Barnes T, Pfaff JM, Mabila M, et al. Mechanism of Binding to Ebola Virus Glycoprotein by the ZMapp, ZMAb, and MB-003 Cocktail Antibodies. Journal of Virology. 2015;89(21):10982-92.

24. Kugelman Jeffrey R, Kugelman-Tonos J, Ladner Jason T, Pettit J, Keeton Carolyn M, Nagle Elyse R, et al. Emergence of Ebola Virus Escape Variants in Infected Nonhuman Primates Treated with the MB-003 Antibody Cocktail. Cell Reports. 2015;12(12):2111-20.

25. Janus BM, van Dyk N, Zhao X, Howell KA, Soto C, Aman MJ, et al. Structural basis for broad neutralization of ebolaviruses by an antibody targeting the glycoprotein fusion loop. Nature communications. 2018;9(1):3934.

Supporting information

S1 File. S1_Ab100_data.xlsx. Excel spreadsheet with estimated stability effects of all 6,441 mutations against Ab100.

S2 File. S2_Ab114_data.xlsx. Excel spreadsheet with estimated stability effects of all 6,441 mutations against Ab114.

S3 File. S3_13F6-1-2_data.xlsx. Excel spreadsheet with estimated stability effects of all 209 mutations against 13F6-1-2.

S4 File. S4_expanded_watchlist.xlsx. Excel spreadsheet with estimated binding stability effects of watch list mutations against all the four antibodies (KZ52, Ab100, Ab114, and 13F6-1-2).

Figure 1

Figure 2

Figure 3

Figure 4