Corresponding author: Gideon Nave

Assistant Professor of Marketing, Wharton School, University of Pennsylvania

Phone: (215) 898-8248 | Fax: (215) 898-2534 | Email: gnave@wharton.upenn.edu

https://marketing.wharton.upenn.edu/profile/gnave/

Authors: Amos Nadler[a], Colin F. Camerer[g], David T. Zava[c], Triana L. Ortiz[d], Neil V. Watson[e],
Justin M. Carré[f], & Gideon Nave[b,1]

Affiliations:

[a] Department of Economics, University of Toronto, 150 St. George Street, Toronto, ON, Canada, M5S
3G7; amos.nadler@utoronto.ca

[b] Marketing Department, The Wharton School of the University of Pennsylvania, 3730 Walnut St,
Philadelphia, PA, USA 19104; gnave@wharton.upenn.edu

[c] ZRT Laboratory, 8605 SW Creekside Place, Beaverton, OR, USA 97008; dzava@zrt.com

[d] Department of Psychology, Nipissing University, 100 College Drive, North Bay, ON, Canada, P1B 8L7;
trianao@nipissingu.ca

[e] Department of Psychology, Simon Fraser University, 8888 University Drive, Burnaby, BC, Canada,
V5A 1S6; nwatson@sfu.ca

[f] Department of Psychology, Nipissing University, 100 College Drive, North Bay, ON, Canada, P1B 8L7;
justinca@nipissingu.ca

[f] Department of the Humanities and Social Sciences, California Institute of Technology, 1200 E
California Blvd, MC 228-77, Pasadena, CA, USA 91125; camerer@hss.caltech.edu

[1]To whom correspondence should be addressed. Email: gnave@wharton.upenn.edu

Title: Does testosterone impair men's cognitive empathy? Evidence from two large-scale
randomized controlled trials

Short title/running head: Does testosterone impair cognitive empathy?

4
5
6

7
8

1 **Abstract**

The capacity to infer others' mental states (known as "mind reading" and "cognitive empathy")

is essential for social interactions across species, and its impairment characterizes

psychopathological conditions such as autism spectrum disorder and schizophrenia. Previous

studies reported that testosterone administration impaired cognitive empathy in healthy humans,

and that a putative biomarker of prenatal testosterone exposure (finger digit ratios) moderated the

effect. However, empirical support for the relationship has relied on small-sample studies with

mixed evidence. We investigate the reliability and generalizability of the relationship in two

large-scale double-blind placebo-controlled experiments in young men (N=243 and N=400),

using two different testosterone administration protocols. We find no evidence that cognitive

empathy is impaired by testosterone administration or associated with digit ratios. With an

unprecedented combined sample size, these results counter current theories and previous high-

profile reports, and demonstrate that previous investigations of this topic have been statistically

underpowered.

Key words: testosterone, cognitive empathy, mind reading, prenatal priming, steroid hormones,

pharmacology

Decades of research on neuroendocrinological influences on animal behaviour has provided a

reliable basis for exploring it in humans [1] and motivate a growing scientific focus on the

biological basis of social aptitudes and the causes of their deficits [2]. One important element of

social cognition is *"cognitive empathy"*, which constitutes the capacity to infer from observation

the emotions, beliefs, and goals of others.[1] This capacity exists across taxa [3] where its

impairment in humans characterizes a broad range of psychopathological conditions and is part

of the clinical diagnostic criteria for autism spectrum disorders (ASDs) [4].[2]

Testosterone-based biological theory of social cognition

A popular biopsychological model known as the Extreme Male Brain (EMB) hypothesis [5]

proposes that two distinct cognitive styles, *"systemizing"* and *"empathizing"*, typify males and

females respectively. The stereotypically male systemizing domain has no social dimension, and

in its extreme form social cognition is extinguished. Guided by observations that ASDs emerge

early in life and are substantially more prevalent among males[3], and that males typically score

lower than females in tests of cognitive empathy [6], the EMB hypothesis proposes that elevated

prenatal exposure to the sex steroid testosterone causes impairments in cognitive empathy,

through its masculinizing effect on the developing brain [7].

The EMB hypothesis found evidential support in a study that reported a correlation between

amniotic testosterone levels and ASD traits ([8], though see [9]), and has remained popular yet

controversial to date. Much of its research has relied on the assumption that the ratio between the

hand's second (index) to fourth (ring) digit (known as 2D:4D) is a developmental proxy for

---

[1] Cognitive empathy is the ability to interpret others' emotions and understand their behaviour vis-a-vis their emotional state; this is distinct from emotional empathy, which is the vicarious feeling of others' emotions along with them (Smith 2006).

[2] The DSM V criteria for ASDs include "Non-verbal communication problems, such as abnormal eye contact, posture, facial expressions, tone of voice and gestures, as well as an inability to understand these."

[3] ASD incidence rates vary widely by study, from 5.2 to 72.6 per 10,000 people and ratios range from 1.81 to 15.7 male:female.

prenatal testosterone exposure [10], which motivated examinations of correlations between

2D:4D and cognitive empathy and ASDs occurrences. While some studies provided supporting

evidence (e.g., [11]), several others failed to detect a relationship between digit ratio and

cognitive empathy [12,13]. Moreover, because it is not feasible to experimentally manipulate

prenatal testosterone exposure in humans (due to ethical considerations), findings along this line

of research have been correlational, which cannot establish causal relations [8].

Testing testosterone's causal effect on cognitive empathy

A handful of experiments attempted to address the above limitation by testing the effects of

testosterone administration on cognitive empathy in neurotypical adults, and investigating the

dependency of these effects on the 2D:4D biomarker [14–17]. This line of research critically

relies on an assumption originating in animal research, that *in utero* androgen exposure

moderates the activational effect of testosterone [18]. The seminal publication along this line of

research reported a placebo-controlled within-subject experiment of 16 healthy females, in which

exogenously administered testosterone strongly impaired cognitive empathy measured using the

"Reading the Mind in the Eyes Test" (RMET), a 36-item battery testing the ability to infer

others' emotional states and intentions from pictures of their eye regions [6] (see Fig. S2 in

Supplementary Material for example item). In addition to reporting a main effect of exogenous

testosterone reducing cognitive empathy, more than 50% of the individual differences in the

effect on the RMET were explained by the participants' variation in the right-hand 2D:4D,

implying involvement of prenatal testosterone exposure in the causal effect [17].

A similar experiment with roughly twice the sample size (*N*=33, all female sample) found a

much smaller[4] main effect (*P*=0.048, one-tailed), and no moderation by 2D:4D [16]. A third

experiment of 16 females found neither a main effect nor a moderation by 2D:4D [15]. Last, one

experiment investigated the effect of testosterone administration on the RMET in 30 healthy

males and found neither a main effect nor a moderation by the right-hand 2D:4D; however,

subsample analysis revealed that testosterone administration reduced cognitive empathy in

participants with relatively low (i.e., more masculine) left-hand 2D:4D, but no relationship for

high 2D:4D or either for the right hand [14].

Do the data support the hypothesis?

Despite these earlier findings the current literature on the effects of testosterone on cognitive

empathy is subjected to important limitations and results reveal weaknesses under scrutiny. First,

albeit there are a few parallel findings in terms of negative direction of effect of testosterone on

RMET performance, there is a lack of replicability across experiments, where only one of the

four studies observed a statistically significant (P<0.05) main effect of testosterone on the RMET

[17]. Moreover, this publication's report of a strong moderating role of the right-hand 2D:4D

was not replicated in any of the other studies (the only other report of an interaction between

testosterone administration and the 2D:4D was observed for the left hand [14]).[5]

A second concern is statistical power. Although the RMET is a noisy psychological instrument,[6]

and 2D:4D is, at best, a noisy proxy of prenatal testosterone exposure [19], all samples ranged

between 16 and 33 participants, which might have been too meager to credibly estimate a true

---

[4] The primary publication [17] had a statistical power of only 0.26 to detect the effect size found in the similar study with twice the sample size [16].

[5] One experiment (in males) reported a statistically significant moderating effect, but only for the left-hand 2D:4D; the two other experiments reported no moderation of the 2D:4D [15].

[6] The RMET has a test-retest reliability of 0.7 [6].

effect size. It is therefore impossible to know whether the inconsistencies in the literature are due

to an absence of a true association, or the result of false negative findings due to low statistical

power. Thus, these inconsistent results necessitate clarification through additional studies.

To this end, we conducted a powerful direct test of the activational and developmental effects of

testosterone on cognitive empathy by measuring the causal effect of exogenous testosterone and

the moderating role of putative prenatal androgenic biomarkers in two studies of healthy young

men. Our studies constitute the two largest behavioural testosterone administration experiments

conducted to date, with samples that were 15 and 25 times greater than the first study that

reported a statistically significant effect of testosterone on the RMET in females [17] and 7 and

12 times greater than the largest experiment in males [14]. In both studies we used a computer-

based version of the RMET to test the hypothesis that testosterone administration and its

purported developmental biomarkers affect cognitive empathy.

Methods

Experiment 1

Participants and experimental procedure. Two hundred forty-three males aged 18 to 55 (mean

age=23.63, SD=7.22) participated in the study and were mostly private Southern California

consortium students from diverse ethnic backgrounds (see Participants section and Table S1a in

Supplementary Material). The institutional review boards of Caltech and Claremont Graduate

University approved the study, all participants gave informed consent, no adverse events

occurred during any experimental session, and no participant or researcher was harmed. All data

and materials are available on the Open Science Framework (https://osf.io/hztfe/).

Participants registered by their preferred session dates and were added to cohorts of 13 to 16. They arrived at the lab at 9 a.m., signed informed consent forms, and had both of their hands scanned before being randomly assigned to private cubicles where they completed demographic and mood questionnaires (see Supplementary Material for all independent variables) and provided an initial saliva sample by passive drool. Next, participants proceeded to gel application (further details below), after which they were given printed material containing precautions and instructions prior to dismissal (experimental timeline shown in Fig. 1). All participants returned to the lab at 2 p.m., provided a second saliva sample, and began a battery of tasks that lasted approximately two hours. We did not randomize the order of the behavioral tasks, in similar fashion to previous studies [20], to standardize hormonal measurements (which have diurnal cycles) among participants. Following the experiment, participants completed an exit survey, where they indicated their beliefs about the treatment they had received using a five-point scale.

Treatment administration. Following initial intake, participants were escorted in groups of two to six to a semiprivate room. There they were provided *en masse* small plastic cups containing either 10 g of topical testosterone that is a widely prescribed transdermal testosterone gel with clearly mapped pharmacokinetics [21] (100 mg, Vogelxo^{TM}, N=123) or volume equivalent of inert placebo of similar viscosity and texture placebo (80% alcogel, 20% Versagel$^®$, N=118) under a double-blind protocol (see Fig. S1a in Supplementary Material for randomization protocol). Participants were instructed to apply the entirety of the gel container following manufacturer instructions.

Saliva samples. Each participant provided four passive drool saliva samples (upon arrival prior to treatment administration, shortly after returning for afternoon session, another closely following the RMET, and a final sample prior to exit survey) for subsequent assay (see

Supplemental Material for precise timing). To allow robust manipulation checks and obtain statistical control for hormonal markers of participants' biological states, we used liquid chromatography tandem mass spectrometry (LC-MS/MS, detection levels and precision are available in Supplementary Material Table S2) to measure the following salivary steroids: estrone, estradiol, estriol, testosterone, androstenedione, DHEA, 5-alpha DHT, progesterone, 17OH-progesterone, 11-deoxycortisol, cortisol, cortisone, and corticosterone (see Supplemental Material Table S7 for measurements).

Digit Ratio Measurements. Participants' hand scans acquired at intake were measured by two independent raters using and digital calipers to quantify 2D:4D; inter-rater correlation was 0.96 and their scores were averaged.

Behavioural task. We administered the adult version of the RMET developed by Baron-Cohen and colleagues [6] which shows the eye region of an actor's face, and a list of four words that describe emotional states and cognitive processes among which participants select the one that best described the person in the image (see Fig. S2 in the Supplemental Material for example item). The task was divided into two segments, baseline (morning) and post-treatment (afternoon), in a repeated measures design (see Fig. 1 below): each participant completed a half of the RMET in the morning (either part "A", items 1-18, or part "B", items 19-36) prior to receiving treatment, and the other half following treatment when, according to published pharmacokinetics, androgen levels are elevated and stable following exogenous application.

Psychological Questionnaires. Because there are various feasible channels through which testosterone could affect RMET performance (and affect being one of them), we measured mood using the PANAS-X scale [22], both pre- and post-treatment (see Table S1a in Supplementary

Material for aggregated responses).

Experiment 2

Participants. Experiment 2 included both students and participants from the general public for a

total sample of 400 participants (mean age=22.80, SD=4.68). The all-male sample was

composed predominantly of Caucasians and overall ethnic heterogeneity was representative of

the region (see Participants section and Table S1b in Supplementary Material). All accepted

participants completed the task and were included in the analysis (for pre-screening criteria see

Supplementary Material). The Nipissing University Research Ethics Board approved this study,

all participants gave informed consent, no adverse events occurred during any experimental

session, and no participant or researcher was harmed.

Experimental procedure. Participants arrived at one of three testing session times (10:00 a.m.,

12:30 p.m., or 2:30 p.m.) in cohorts of six and were brought individually into private testing

rooms to read and sign an informed consent form, receive a participant number, and complete

questionnaires (see Supplementary Material for all independent variables). Afterwards,

participants provided a 1-2 ml saliva sample before treatment administration, after which they

had their photos taken and hands scanned. Approximately two hours after arrival to the lab and

one-and-a-half hours after drug administration participants completed the RMET then provided

their final saliva sample. Upon session completion, participants received compensation and

completed an exit survey asking which treatment they believed they had received (see Figure

S1b in Supplementary Material).

Treatment administration. Following initial saliva sample collection, a researcher provided

two syringes pre-filled by a pharmacist following a double-blind protocol each containing 5.5 mg

of either placebo or testosterone gel (for a total of 11 mg). This is a newly approved nasal gel

used for the treatment of hypogonadism. Pharmacokinetic data indicate that serum testosterone

concentrations rise sharply within 15 minutes of testosterone gel application and remain elevated

(relative to placebo) up to 180 minutes post-application [23]. The gel was either Natesto® or the

volume equivalent of an inert placebo of similar viscosity and texture. Random assignment was

determined such that half the participants in every group received testosterone and half received

placebo such that total participants base was bisected with N=200 for both testosterone and

placebo groups (see Fig. S1b in Supplementary Material for randomization protocol).

Saliva samples. Each participant provided two saliva samples, with the first sample collection

time 30 minutes following arrival (and prior to gel application), and the second 120 minutes after

arrival. Participants provided passive drool into a 5 ml polystyrene tube while situated in their

individual testing rooms as instructed by a research assistant and samples were analyzed for pre-

and post-treatment testosterone and baseline cortisol using commercially available enzyme

immunoassay kits (DRG International) (see Table S2b in Supplementary Material for hormone

measures).

Digit Ratio Measurements. Participants' 2D:4D were measured by two independent raters

using hand scans and digital calipers with an inter-rater correlation of 0.86 and their scores were

averaged.

Behavioural task. Participants evaluated all 36 items of the RMET [6] as a single task.

Figure 1:

Comparison of experimental features to van Honk et al. (2011)

We note several differences between our study and the primary positive report of testosterone

administration on the RMET in 16 females [17].

Participant Sex. The EMB theory does not make any sex-specific predictions regarding the

developmental and activational effects of testosterone on cognitive empathy. However, van

Honk et al. [17] conducted their study in a female-only sample because the pharmacokinetics of

a single dose of testosterone had only been studied in females at the time [17]: *"We exclusively*

recruited women because the parameters ... for inducing neurophysiological effects ... are

known in women but not in men." (p. 3450). The recent pharmacokinetic mappings for short-

term single dose testosterone administrations [14,21] and availability of two unique

administration modalities of FDA-approved exogenous testosterone provided us with a reliable

foundation for testing the EMB hypothesis in men.

Drug dosage and delivery. van Honk et al. [17] used a sublingual testosterone administration

procedure, which causes a sharp increase in serum testosterone of 10-fold or more within 15

minutes, with a rapid decline to normal levels within 90 minutes in women [24]. It is important

to note that the pharmacokinetic data for sublingual administration (published in Fig. 1 of Tuiten

et al. [24]) show that at the time the task was performed—4 hours after sublingual

administration—participants' testosterone levels were the same across the testosterone and placebo groups. Moreover, the Tuiten et al. study that served as a justification for using a 4-hour delay had only 8 participants, and reported a statistically weak treatment effect (p=0.04, uncorrected for multiple comparisons).

In Experiment 1 we chose to administer testosterone using an FDA-approved transdermal gel for three reasons. First, transdermal gel had been extensively studied in the medical literature both prior and following its approval [25,26]. Second, one of our laboratories found reliable behavioural effects with robust manipulation check in serum using a single dose [27], and third, the pharmacokinetics of a single-dose of this testosterone administration method were mapped prior to the inception of our experiments by a study showing that plasma testosterone levels peaked 3 hours after single-dose exogenous transdermal administration, and stabilized at high levels between 4 and 7 hours following administration [21].[7] Therefore, we had all participants return to the lab 4.5 hours after receiving gel, when androgen levels were elevated and stable. We used a 100 mg transdermal dose, which quickly elevates then holds testosterone levels high and stable for approximately 24 hours [25] and was shown to generate effects on cognition, decision making, and other behaviours [27,29–31].

In Experiment 2 we used nasal delivery, following a recent study indicating that serum testosterone concentrations rise sharply within 15 minutes after Natesto® gel application and remain elevated for approximately three hours post application among hypogonadal males [23] (see Fig. S3 in Supplementary Material). This method conforms to our experimental paradigm's pharmacokinetic structure as serum testosterone approached its zenith in treated participants as they completed the RMET (see Supplementary Material Tables S2a and b for testosterone

---

[7] Subsequent studies measuring testosterone in serum in significantly larger sample sizes demonstrate an earlier hormonal peak at 60 minutes post administration with subsequent stabilization [14,25].

levels).

The doses in both experiments are commonly prescribed daily to men with low circulating

testosterone levels and serve as two distinct physical transport channels (transdermal and

intranasal, respectively) to reduce the probability that behavioural effects are transport channel

specific. Various studies show significant heterogeneity in change in testosterone levels

depending on delivery method, location of application in the body, and biofluid measured

[14,21,24,27,29,32]. However, all the exogenous delivery methods in this particular literature

cause a common hormonal trajectory characterized by a rapid initial rise, a peak above typical

circulating levels, and eventual return to baseline.

Experimental designs. van Honk et al. [17] used the same questions in pre- and post-treatment

testing. As testosterone treatment might affect participants' capacity to recall answers [33], this

design choice might have introduced memory confounds. In Experiment 1 we divided the RMET

into two portions, and administered each portion as either pre- or post-treatment measure in,

allowing us to capture baseline abilities while ruling out such confounds. In Experiment 2 we

conducted a between-subjects experiment that removes all effects of practice and recall from the

data.

Results

Manipulation check

In both experiments, pre-treatment (i.e., baseline) saliva testosterone levels were similar across

the two treatment groups, and post-treatment saliva testosterone levels were greater in the

testosterone group compared with the placebo group. In experiment 1, the mean logged pre-

treatment testosterone level was 5.58 pg/ml (*SD*=0.08) in the testosterone group and 5.77 pg/ml

(*SD*=0.09) in the placebo group (two-sided t-test: *P*=0.13, *t*(239)=1.50). Mean post-treatment

logged testosterone levels were 8.38 pg/ml (*SD*=0.15) in the testosterone group and 5.11 pg/ml

(*SD*=0.08) in the placebo group,[8] (*t*(239)=18.43, *P*<0.0001). Likewise in experiment 2, we find

similar mean baseline saliva testosterone concentrations between the groups, with l5.3 pg/ml

(SD=0.88) in the testosterone group and 5.4 pg/ml (*SD*=0.85) in the placebo group (two-sided t-

test: *P*=0.49, *t*(394)=0.69). The mean logged post-treatment saliva testosterone levels were (8.00

pg/ml, *SD*=1.27) in the testosterone group and 5.33 pg/ml (*SD*=0.73) in the placebo group (two-

sided t-test: *P*<0.001, *t*(396)=22.14)   (see Fig. 1 and Tables S2a and S2b in Supplementary

Material).

Consistent with previous reports, we found no treatment effects on mood and treatment

expectancy (e.g., [34]) (see Table S2a and S2b in Supplement), or levels of other hormones

unaffected by exogenous testosterone, as measured by LC-MS/MS in Experiment 1 or enzyme

immunoassay in Experiment 2 (see Tables S1a and S1b in Supplement).

Influence of Testosterone on RMET scores

Overall RMET scores in our samples were comparable with previous studies of similar

populations (see Fig. 2A below). Figure 2B shows baseline and post-treatment RMET scores in

Experiment 1, separated by treatment group and order. As expected, baseline (morning) RMET

performance were reliably correlated with afternoon scores (*r*(241)=0.40, *P*<0.001). In addition,

---

[8] Median testosterone levels (unlogged) of the testosterone group were 33.5 times that of the placebo group post-treatment.

participants' scores were, on average, slightly higher in the B portion of the test (A portion

average=13.54, *SD*=2.43; B portion average=13.95, *SD*=2.19, *t*(241)=2.53, *P*=0.01). Figure 2C

shows Experiment 2 scores by treatment groups.

Figure 2:

To test for the main effect of testosterone administration on cognitive empathy for Experiment 1,

we estimated linear regression models with the post-treatment (afternoon) RMET score as the

dependent variable, a binary treatment indicator (testosterone =1, placebo=0) as the independent

variable of primary interest, controlling for baseline performance, the order of the two portions

of the RMET (A and B) and additional control variables[9] (the results remain unchanged when

these control variables are excluded from the models; see Tables S3a and S4a in Supplementary

Material). Analogously, Experiment 2 data were analyzed using linear models with total RMET

score as the dependent variable, a binary treatment indicator as the chief independent variable of

interest, and control variables[10] (results remain unchanged with their exclusion from the models;

see Supplementary Material Tables S3b and S4b).

We found no reliable effect of testosterone administration on the RMET in Experiment 1

(beta=0.11, 95% CI=[−0.45, 0.68]; *t*(237)=0.37, *P*=0.71; Cohen's *d*=0.04, 95% CI=[−0.19,

0.28]). Thus, the effect's point estimate was positive and the 95% CI excluded the *d*=-0.49

reported in (van Honk et al. 2011) or any *negative* effects that are greater in magnitude than

d=0.19. A sample of at least 870 participants (in a between-subject design), or 435 subjects (in a

---

[9] These include RMET baseline scores, portion A or B, 2D:4D and treatment interactions, Cognitive Reflection Task (CRT) scores, math abilities, mood and affective measures, treatment expectancy, age, marital status, sexual preference, and all other measured hormones that were not influenced by testosterone treatment and may affect cognition and decision-making (e.g., cortisol [35,36]). The CRT control was added because performance is impaired by exogenous testosterone [29] and people with ASD outperform non-ASD age-matched controls [37].
[10] These include Cognitive Reflection Task (CRT) scores, factor 1 and 2 psychopathy measures, treatment expectancy, age, marital status, sexual preference, and all other measured hormones that were not influenced by testosterone treatment.

within-subject design), which is over 26 to 54 (within-subject 13 to 26) times greater than previous investigations, would be required to reliably detect even this "optimistic" negative effect size estimate with statistical power of 0.8. Regression analyses with comprehensive controls corroborate the absence of a main treatment effect, and the absence of moderation by 2D:4D (right hand, left hand, and their average), as implied by insignificant interaction coefficients (see Supplementary Material, Table S3a). Furthermore, in an analysis analogous to the previous positive report [17], we found no correlation between the treatment effect on the RMET and the right-hand 2D:4D in the testosterone group (r(123)=0.04, P=0.66, 95% CI=[−0.14, 0.22]) (see Fig. S4 in the Supplementary Material).

Experiment 2, which had 400 participants, could also not reject the null hypothesis (beta=0.27, 95% CI=[−0.49, 1.02]; t(398)=0.69, P=0.49; Cohen's d=0.04, 95% CI=[−0.15, 0.24]) and there was no significant treatment effect in any regression model (Supplementary Material Tables S3b and S4b). Similarly, the point estimate of the effect in Experiment 2 was positive, and the 95% CI did not include negative effects of testosterone administration on the RMET that were greater in magnitude than 0.15. Further analyses of each question in isolation using Chi-squared tests revealed no systematic differences between treatment conditions in any of the RMET items in both experimental datasets (see Supplementary Material Tables S5a and b).

Testing for effect of 2D:4D

Putative prenatal testosterone proxies (2D:4D, either right-hand, left-hand, or their average) did not correlate with RMET scores in both experiments or moderate the effect of testosterone administration, echoing other recent findings [14–16] (see Supplementary Material Tables S3a-S4b). These results are in line with previous reports showing no correlation between the 2D:4D

and RMET scores [12,38,39], and in contrast to the two papers reporting an interaction between

2D:4D and the exogenous testosterone's effect on the RMET [14,17].

Discussion

Our experiments used two notably large samples to test the effects of pharmacological

testosterone manipulation on cognitive empathy. Despite experimental differences between

them, their collected data exhibit the same results with robust statistical consistency, to

demonstrate a lack of effects of testosterone administration and 2D:4D on cognitive empathy.

These findings, and the literature as a whole, cast serious doubts on the proposal that testosterone

causally impairs cognitive empathy, for several reasons.

First, the low statistical power of previous investigations undermines their reliability in capturing

true effects. Even if we assume that a purported size of testosterone's negative effect on

cognitive empathy is the overly "optimistic" negative bound of our confidence interval for $d=$-

0.19 in Experiment 1, we find that all previous investigations of the topic were statistically

underpowered (<0.3 power). Second, the results of the previous small sample studies are

discrepant. Our large samples draw on drastically more data than all previous investigations

combined, and generalize across geographically, economically, and culturally distinct

populations (see Participants section of Supplementary Material). Our use of two different

experimental designs and testosterone administration protocols across these populations further

mitigates the concern that the outcomes were due to a particular experimental factor. Of note,

there are some design differences between our studies and previous investigations (see Table 1;

differences from [17] discussed above). However, even if those design differences led to a

complete abolishment of a "real" effect of testosterone on cognitive empathy, our results

demonstrate beyond a reasonable doubt that such an effect is not generalizable to both males and

females. Future work with females could employ a similar approach as ours characterized by

large samples from different geographies, distinct administration methods, and other design

features that strongly inform whether a relationship (or its absence) generalizes across sexes.

Table 1. Summary of literature linking testosterone administration and the RMET

	Study design				Results					
	Sex	**N**	**Design**	**Dose**	**Main effect**	**Effect Size**	**SE of Effect Size**	**95% CI for Effect Size**		**2D:4D moderation**
van Honk et al. (2011)	F	16	Within subject; repeated task	0.5 mg sublingual	One-tailed Wilcoxon P = 0.01	-0.49	0.25	-1.19	0.21	Low right hand ratio, negative relationship
Olsson et al. (2016)	F	33	Within subject; repeated task	50 mg transdermal	One-tailed P = 0.048	-0.33	0.13	-0.15	0.82	No
Bos et al. (2016)	F	16	Within subject; no repetition	0.5 mg sublingual	No effect P = 0.78	-0.10	0.10	-0.60	0.79	No
Carré et al. (2015)	M	30	Within subject	150 mg transdermal	No effect P = 0.25	0.20	0.35	-0.70	0.31	Low left hand ratio, negative relationship
Nadler et al. Experiment 1	M	241	Between subject; no repetition	100 mg transdermal	No effect P = 0.83	0.03	0.36	-0.19	0.28	No
Nadler et al. Experiment 2	M	400	Within subject; no repetition	11 mg intranasal	No effect P = 0.66	0.04	0.26	-0.15	0.24	No

In "repeated task" paradigms, participants completed the RMET twice: prior to any treatment in Experiment 1 in this study, and after receiving a placebo or testosterone in van Honk et al. (2011).
Effect size is calculated using Hedge's bias-correct effect size due to small sample size and unequal variances between treatment and control groups.

2
3 A third reason concerns the validity of the 2D:4D biomarker. The initial findings that prenatal

4 testosterone exposure correlates with 2D:4D are supported in non-clinical and clinical human

5 populations [11], as well as in preliminary causal evidence in relative phalanx/tibia lengths in

6 mice [18]. However, recent work highlights concerns regarding the reliability of 2D:4D as a

biomarker [41,42]. For example, the 2D:4D of complete androgen insensitivity syndrome

patients were found to be only somewhat feminized, and had the same variance as in healthy

controls, demonstrating that the preponderance of individual differences in the measure is not

attributable to the influence of testosterone exposure [19]. There is also longitudinal evidence

that 2D:4D systematically changes during childhood [43,44], which is unconformable with the

preposition that it accurately quantifies prenatal influences. Moreover, while many studies report

2D:4D sexual dimorphism [45,46], other studies suggest lack of ethnic universality of

dimorphism [47,48]. Finally, there is also a debate on whether sexual dimorphism is the product

of allometric shift in shape rather than hormonal influences [49,50].

Furthermore, many reports of correlations between 2D:4D and behavioural traits hold only for

subsets of the population (e.g., particular sex or race). Correlation sometimes holds only for the

right hand 2D:4D but in other times only for the left hand, or for the average of both hands [51].

Overall, significant results are seldom replicated, and few survive correction for multiple

comparisons or meta-analytic aggregations (e.g., [52]). These concerns belie the validity of the

measure as a biomarker and its capacity to detect reliable correlations with noisy psychological

constructs in studies of small samples.

Despite our dissenting results, the absence of evidence is not necessarily evidence of absence.

Specifically, the lack of an association between 2D:4D and cognitive empathy could be

attributable to the failure of the measure to serve as a reliable androgenic biomarker. We

therefore agree with Baron-Cohen et al. [53] (p. 6) that it is worthwhile to study the occurrence

of impaired cognitive empathy and other ASD traits in developmentally unique populations. One

such study reported mixed evidence of higher scores in some Autism Quotient self-rating

subscales among women with congenital adrenal hyperlexia (CAH) and lower scores in other

subscales, compared with their unaffected relatives [54], with no significant results in men.

However, results along this line of research, too, are far from being conclusive. For example, [9]

found that young females with and without CAH did not differ in autistic traits, and that amniotic

testosterone levels were not associated with scores from either sex individually or the entire

sample. Other longitudinal studies also found no association between various measures of

prenatal androgens measured in umbilical cord blood and amniotic fluid and autistic traits

[55,56]. Thus, further investigations, preferably using larger samples, are required for resolving

the inconsistencies in this literature.

To conclude, we tested testosterone's causal role in cognitive empathy across distinct

administration methods using notably large samples from two distinct populations, and found no

evidence of an effect of testosterone administration on RMET in young adult neurotypical

human males. While our results do not exclude all possible relationships between testosterone

and interpreting others' emotions and states of mind, our large-scale study and evaluation of

previous literature exhibit robust evidence of no causal relationship between activational and

purported developmental testosterone exposure and cognitive empathy.

Author contributions

A.N.: experimental design, manuscript, data analysis; G.N.: experimental design, manuscript,

data analysis; C.C.: manuscript; D.Z.: manuscript, hormonal assay; T.L.O.: experimental design,

data collection; hormonal assay; N.V.W.: manuscript; J.M.C.: experimental design, manuscript.

Acknowledgements. Funding for this work was generously provided by Caltech, Ivey Business

School, IFREE, Russell Sage Foundation, University of Southern California, INSEAD, Stockholm School of Economics, Wharton Neuroscience Initiative, the Natural Sciences and Engineering Council of Canada, and the Northern Ontario Heritage Fund Corporation. Special thank you to Jorge Barraza, Austin Henderson, Garrett Thoelen, Dylan Mandred, Kimberly Gilbert, Caelan Mathers, Emily Jeanneault, Nicole Marley, Kendra Maracle, Victoria Bass-Parcher, Nadia Desrosiers, Charlotte Miller, Brittney Robinson, Dalton Rogers, Megan Phillips, Brandon Reimer, Camille Gray, Christine Jessamine, and Brandon Reimer who assisted this study, and David Kimball for LC-MS/MS assay testing.

References

1. McCall C, Singer T. 2012 The animal and human neuroendocrinology of social cognition, motivation and behavior. *Nat. Neurosci.* **15**, 681–688.

2. Ferguson JN, Young LJ, Insel TR. 2002 The neuroendocrine basis of social recognition. *Front. Neuroendocrinol.* **23**, 200–224.

3. Decety J, Bartal IB-A, Uzefovsky F, Knafo-Noam A. 2016 Empathy as a driver of prosocial behaviour: highly conserved neurobehavioural mechanisms across species. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **371**, 20150077.

4. American Psychiatric Association. 2013 *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub.

5. Baron-Cohen S. 2002 The extreme male brain theory of autism. *Trends Cogn. Sci.* **6**, 248–254.

6. Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I. 2001 The 'Reading the Mind in the Eyes' test revised version: A study with normal adults, and adults with Asperger syndrome or high-functioning autism. *J. Child Psychol. Psychiatry* **42**, 241–251.

7. Zuloaga DG, Puts DA, Jordan CL, Breedlove SM. 2008 The role of androgen receptors in the masculinization of brain and behavior: what we've learned from the testicular feminization mutation. *Horm. Behav.* **53**, 613–626.

8. Auyeung B, Baron-Cohen S, Ashwin E, Knickmeyer R, Taylor K, Hackett G. 2009 Fetal testosterone and autistic traits. *Br. J. Psychol.* **100**, 1–22.

9. Kung KTF *et al.* 2016 No relationship between prenatal androgen exposure and autistic traits: convergent evidence from studies of children with congenital adrenal hyperplasia and of amniotic testosterone concentrations in typically developing children. *J. Child Psychol. Psychiatry* **57**, 1455–1462.

10. Manning JT. 2002 *Digit Ratio: A Pointer to Fertility, Behavior, and Health*. Rutgers University Press.

11. Lutchmaya S, Baron-Cohen S, Raggatt P, Knickmeyer R, Manning JT. 2004 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Hum. Dev.* **77**, 23–28.

12. Hönekopp J. 2012 Digit ratio 2D:4D in relation to autism spectrum disorders, empathizing, and systemizing: a quantitative review. *Autism Res.* **5**, 221–230.

13. Guyatt AL, Heron J, Le Cornu KB, Golding J, Rai D. 2015 Digit ratio and autism spectrum disorders in the Avon Longitudinal Study of Parents & Children: a birth cohort study. *BMJ Open*

14. Carré JM, Ortiz TL, Labine B, Moreau BJP, Viding E, Neumann CS, Goldfarb B. 2015 Digit ratio (2D:4D) and psychopathic traits moderate the effect of exogenous testosterone on socio-cognitive processes in men. *Psychoneuroendocrinology* **62**, 319–326.

15. Bos PA, Hofman D, Hermans EJ, Montoya ER, Baron-Cohen S, van Honk J. 2016 Testosterone reduces functional connectivity during the 'Reading the Mind in the Eyes' Test. *Psychoneuroendocrinology* **68**, 194–201.

16. Olsson A, Kopsida E, Sorjonen K, Savic I. 2016 Testosterone and estrogen impact social evaluations and vicarious emotions: A double-blind placebo-controlled study. *Emotion* **16**, 515–523.

17. van Honk J, Schutter DJ, Bos PA, Kruijt A-W, Lentjes EG, Baron-Cohen S. 2011 Testosterone administration impairs cognitive empathy in women depending on second-to-fourth digit ratio. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 3448–3452.

18. Zheng Z, Cohn MJ. 2011 Developmental basis of sexually dimorphic digit ratios. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 16289–16294.

19. Berenbaum SA, Bryk KK, Nowak N, Quigley CA, Moffat S. 2009 Fingers as a marker of prenatal androgen exposure. *Endocrinology* **150**, 5119–5124.

20. Zethraeus N, Kocoska-Maras L, Ellingsen T, von Schoultz B, Hirschberg AL, Johannesson M. 2009 A randomized trial of the effect of estrogen and testosterone on economic behavior. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 6535–6538.

21. Eisenegger C, von Eckardstein A, Fehr E, von Eckardstein S. 2013 Pharmacokinetics of testosterone and estradiol gel preparations in healthy young men. *Psychoneuroendocrinology* **38**, 171–178.

22. Watson D, Clark LA. 1999 The PANAS-X: Manual for the positive and negative affect schedule-expanded form.

23. Geniole SN *et al.* In press. Using a psychopharmacogenetic approach to identify the pathways through which and people for whom testosterone promotes aggression. *Psychol. Sci.*

24. Tuiten A, Van Honk J, Koppeschaar H, Bernaards C, Thijssen J, Verbaten R. 2000 Time course of effects of testosterone administration on sexual arousal in women. *Arch. Gen. Psychiatry* **57**, 149–53; discussion 155–6.

25. Swerdloff RS *et al.* 2000 Long-term pharmacokinetics of transdermal testosterone gel in hypogonadal men. *J. Clin. Endocrinol. Metab.* **85**, 4500–4510.

26. Resnick SM *et al.* 2017 Testosterone Treatment and Cognitive Function in Older Men With Low Testosterone and Age-Associated Memory Impairment. *JAMA* **317**, 717–727.

27. Nadler A, Jiao P, Alexander V, Johnson C, Zak P. 2017 The Bull of Wall Street: Experimental Analysis of Testosterone and Asset Trading. *Manage. Sci.* (doi:10.1287/mnsc.2017.2836)

28. Carré JM, Geniole SN, Ortiz TL, Bird BM, Videto A, Bonin PL. 2017 Exogenous Testosterone Rapidly Increases Aggressive Behavior in Dominant and Impulsive Men. *Biol. Psychiatry* **82**, 249–256.

29. Nave G, Nadler A, Zava D, Camerer C. 2017 Single-Dose Testosterone Administration Impairs Cognitive Reflection in Men. *Psychol. Sci.* **28**, 1398–1407.

30. Nave G, Nadler A, Dubois D, Zava D, Camerer C, Plassmann H. 2018 Single-dose testosterone administration increases men's preference for status goods. *Nat. Commun.* **9**, 2433.

31. Henderson A, Thoelen G, Nadler A, Barraza J, Nave G. 2018 Testing the influence of testosterone administration on men's honesty in a large laboratory experiment. *Sci. Rep.* **8**, 11556.

32. Geniole SN *et al.* 2019 Using a Psychopharmacogenetic Approach To Identify the Pathways Through Which—and the People for Whom—Testosterone Promotes Aggression. *Psychological Science*. **30**, 481–494. (doi:10.1177/0956797619826970)

33. Cherrier MM, Craft S, Matsumoto AH. 2003 Cognitive changes associated with supplementation of testosterone or dihydrotestosterone in mildly hypogonadal men: a preliminary report. *J. Androl.* **24**, 568–576.

34. Eisenegger C, Naef M, Snozzi R, Heinrichs M, Fehr E. 2010 Prejudice and truth about the effect of testosterone on human bargaining behaviour. *Nature* **463**, 356–359.

35. Margittai Z, Nave G, Van Wingerden M, Schnitzler A, Schwabe L, Kalenscher T. 2018 Combined Effects of Glucocorticoid and Noradrenergic Activity on Loss Aversion. *Neuropsychopharmacology* **43**, 334–341.

36. Margittai Z, Nave G, Strombach T, van Wingerden M, Schwabe L, Kalenscher T. 2016 Exogenous cortisol causes a shift from deliberative to intuitive thinking. *Psychoneuroendocrinology* **64**, 131–135.

37. Brosnan M, Lewton M, Ashwin C. 2016 Reasoning on the Autism Spectrum: A Dual Process Theory Account. *J. Autism Dev. Disord.* **46**, 2115–2125.

38. Masuya Y, Okamoto Y, Inohara K, Matsumura Y, Fujioka T, Wada Y, Kosaka H. 2015 Sex-different abnormalities in the right second to fourth digit ratio in Japanese individuals with autism spectrum disorders. *Mol. Autism* **6**, 34.

39. Voracek M, Dressler SG. 2006 Lack of correlation between digit ratio (2D: 4D) and Baron-Cohen's 'Reading the Mind in the Eyes' test, empathy, systemising, and autism-spectrum quotients in a general population sample. *Pers. Individ. Dif.* **41**, 1481–1491.

40. Huber SE, Lenz B, Kornhuber J, Müller CP. 2017 Prenatal androgen-receptor activity has organizational morphological effects in mice. *PLoS One* **12**, e0188752.

41. Valla JM, Ceci SJ. 2011 Can Sex Differences in Science Be Tied to the Long Reach of Prenatal Hormones?: Brain Organization Theory, Digit Ratio (2D/4D), and Sex Differences in Preferences and Cognition. *Perspect. Psychol. Sci.* **6**, 134–146.

42. Richards G. 2017 What is the evidence for a link between digit ratio (2D:4D) and direct measures of prenatal sex hormones? *Early Hum. Dev.* **113**, 71–72.

43. McIntyre MH, Ellison PT, Lieberman DE, Demerath E, Towne B. 2005 The development of sex differences in digital formula from infancy in the Fels Longitudinal Study. *Proc. Biol. Sci.* **272**, 1473–1479.

44. Trivers R, Manning J, Jacobson A. 2006 A longitudinal study of digit ratio (2D:4D) and other finger ratios in Jamaican children. *Horm. Behav.* **49**, 150–156.

45. Galis F, Ten Broek CMA, Van Dongen S, Wijnaendts LCD. 2010 Sexual dimorphism in the prenatal digit ratio (2D:4D). *Arch. Sex. Behav.* **39**, 57–62.

46. Putz DA, Gaulin SJC, Sporter RJ, McBurney DH. 2004 Sex hormones and finger length. *Evol. Hum. Behav.* **25**, 182–199.

47. Apicella CL, Tobolsky VA, Marlowe FW, Miller KW. 2016 Hadza hunter-gatherer men do not have more masculine digit ratios (2D:4D). *Am. J. Phys. Anthropol.* **159**, 223–232.

48. Galizzi MM, Nieboer J. 2015 Digit ratio (2D:4D) and altruism: evidence from a large, multi-ethnic sample. *Front. Behav. Neurosci.* **9**, 41.

49. Forstmeier W. In press. Avoiding misinterpretation of regression lines in allometry: is sexual dimorphism in digit ratio spurious? (doi:10.1101/298786)

50. Kratochvíl L, Flegr J. 2009 Differences in the 2nd to 4th digit length ratio in humans reflect shifts along the common allometric line. *Biology Letters*. **5**, 643–646. (doi:10.1098/rsbl.2009.0346)

51. Teatero ML, Netley C. 2013 A Critical Review of the Research on the Extreme Male Brain Theory and Digit Ratio (2D:4D). *J. Autism Dev. Disord.* **43**, 2664–2676.

52. Turanovic JJ, Pratt TC, Piquero AR. 2017 Exposure to fetal testosterone, aggression, and violent behavior: A meta-analysis of the 2D:4D digit ratio. *Aggress. Violent Behav.* **33**, 51–61.

53. Baron-Cohen S, Lombardo MV, Auyeung B, Ashwin E, Chakrabarti B, Knickmeyer R. 2011 Why are autism spectrum conditions more prevalent in males? *PLoS Biol.* **9**, e1001081.

54. Knickmeyer R, Baron-Cohen S, Fane BA, Wheelwright S, Mathews GA, Conway GS, Brook CGD, Hines M. 2006 Androgens and autistic traits: A study of individuals with congenital adrenal hyperplasia. *Horm. Behav.* **50**, 148–153.

55. Jamnadass ESL, Keelan JA, Hollier LP, Hickey M, Maybery MT, Whitehouse AJO. 2015 The perinatal androgen to estrogen ratio and autistic-like traits in the general population: a longitudinal pregnancy cohort study. *J. Neurodev. Disord.* **7**, 17.

56. Whitehouse AJ, Mattes E, Maybery MT, Dissanayake C, Sawyer M, Jones RM, Pennell CE, Keelan JA, Hickey M. 2012 Perinatal testosterone exposure and autistic-like traits in the general population: a longitudinal pregnancy-cohort study. *J. Neurodev. Disord.* **4**, 25.

Figure 1. Experimental Timelines

a. Experiment 1

b. Experiment 2

Note: horizontal axis not to scale

Figure 2. RMET distributions and scores across treatment groups

1 Figure legends
2
3 Figure 1

Time-coded experimental timeline shows that, following intake (morning), participants completed half of the RMET (portion A or B), provided a saliva sample, and received gel prior to being dismissed. Upon their return to the lab (afternoon), another saliva sample was collected prior to taking the second portion of the RMET (B or A) (Standard errors shown).	Timeline generalizes experimental sequence for all three start times for Experiment 2. Following arrival and completion of consent and self-report questionnaires, participants provided a saliva sample approximately 30 minutes prior to drug administration. The second sample was collected at the end of a two-hour protocol approximately one-and-a-half hours after drug administration and fifteen minutes after the RMET.

4
5
Figure 2
Fig. 2. (*A*) Cumulative distributions of scores from our experiments juxtaposed with the original paper [6]; scores for Experiment
1 portions A and B (both of which are unaffected by treatment) are combined into a total score: RMET scores in Experiment 1
was 27.5 (*SD*=3.9), and 25.6 (*SD*=3.9) in Experiment 2, which are similar to male students in [6] showing average score of 27.3
(*SD*=3.7). *(B)* Experiment 1 pre- and post-treatment RMET scores. No pre- or post-treatment differences were found between the
two groups, regardless of the order in which portions of the tests were taken (standard errors shown). (*C*) Experiment 2 RMET
scores. No differences were found between treatment groups.
13